SYNAPTIC RELEASE POTENTIATION
AT AGING AUDITORY RIBBON SYNAPSES

Running title: Synaptopathy at aging auditory ribbon synapses

Thibault PEINEAU[1,2#], Séverin BELLEUDY[1#], Susanna PIETROPAOLO[3], Yohan BOULEAU[1,2] and Didier DULON[1,2*]

[1]Neurophysiologie de la Synapse Auditive, INSERM UMRS 1120, Bordeaux Neurocampus, Université de Bordeaux, 33076 Bordeaux, France

[2] Institut de l'Audition, Centre Institut Pasteur/Inserm, 63 rue de Charenton, 75012 Paris

[3] INCIA, UMR 5287, CNRS, University of Bordeaux, Bat B2, Allée Geoffroy St. Hilaire, CS 50023, 33615, Pessac Cedex, France.

[*] Corresponding author: Didier DULON
didier.dulon@inserm.fr

# First co-authors, contributed equally to the study

Keywords: Auditory hair cells, Synaptic ribbons, Synaptopathy, Hyperacusis, Ca^{2+} channels, Exocytosis

Conflict of Interest
The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions
DD designed research; SB, TP, YB and DD performed research and analyzed the data. SP performed the Acoustic Startle Experiments. DD wrote the paper.

Funding & acknowledgments
This work was supported by a grant from the Fondation Pour l'Audition to DD (IDA CLINICAL TRANSLA - 010703E), and from the French Society of Audioprosthesis professionals, Entendre SAS (Saint Cyr l'Ecole 78210 France). Part of the content of the manuscript has previously appeared as a preprint online at bioRxiv doi: https://doi.org/10.1101/2020.05.15.097550.

ABSTRACT

Age-related hidden hearing loss is often described as a cochlear synaptopathy that results from a progressive degeneration of the inner hair cell (IHC) ribbon synapses. The functional changes occurring at these synapses during aging are not fully understood. Here, we characterized this aging process in IHCs of C57BL/6J mice, a strain which is known to carry a cadherin23 mutation and experiences early hearing loss with age. These mice, while displaying a large increase in auditory brainstem thresholds due to 50 % loss of IHC synaptic ribbons at middle age (postnatal day 365), paradoxically showed enhanced acoustic startle reflex suggesting a hyperacusis-like response. The auditory defect was associated with a large shrinkage of the IHCs' cell body and a drastic enlargement of their remaining presynaptic ribbons which were facing enlarged postsynaptic AMPAR clusters. Presynaptic Ca^{2+} microdomains and the capacity of IHCs to sustain high rates of exocytosis were largely increased, while on the contrary the expression of the fast repolarizing BK channels, known to negatively control transmitter release, was decreased. This age-related synaptic plasticity in IHCs suggested a functional potentiation of synaptic transmission at the surviving synapses, a process that could partially compensate the decrease in synapse number and underlie hyperacusis.

INTRODUCTION

Age-related hearing loss (ARHL or presbycusis) is the most prevalent form of sensory disability in human populations and adversely affects the quality of life of many elderly individuals. Simultaneously with ARHL, chronic tinnitus and hyperacusis often develop due to abnormal neural activity in the central auditory system whose origin may arise from cochlear pathologies (for review see Knipper et al., 2013). The development of ARHL has been generally considered to be associated with primary degeneration of cochlear hair cells, with secondary deafferentation associating degeneration of spiral ganglion neurons and ribbon synapses. However, recent studies have resulted in a paradigm shift in the understanding of the early cause of ARHL (Kujawa and Liberman, 2015; 2019). During aging, hearing thresholds in the elderly often remain normal but speech intelligibility become hampered, especially in noisy environment. This pathology in the elderly is referred as hidden hearing loss and has been proposed to be caused in mice by an early loss of ribbon synapses contacting the inner hair cells (IHCs), well before any major loss of hair cells and spiral ganglion neurons (Stamataki et al., 2006). While glutamate excitotoxicity, resulting in nerve terminal swelling, has been proposed to play a role in the setting of noise-induced hearing loss (Pujol and Puel, 1999; Hu et al., 2020), the molecular mechanisms underlying the degeneration of the cochlear ribbon synapses during aging remain to be elucidated. Oxidative damage due to an increase in ROS levels seems to plays a crucial role in ARHL (Fu et al., 2018) but how the ribbon synapses are functionally affected remain unknown.

The C57BL/6J inbred strain of mice has been extensively used as model for ARHL (Johnson et al., 1997; Spongr et al., 1997; Hequembourg and Liberman, 2001; Bartolomé et al., 2002; Henry, 2002). This mouse strain carries a cadherin23 mutation (*Cdh23* [753A], also known as *Ahl*), which affects inner ear structures, characterized by progressive degeneration of ribbon synapses and spiral ganglion neurons, and results in age-related hearing loss. *Ahl* is a recessive single nucleotide mutation at 753 (G=>A) on the *Cdh23* gene on mouse chromosome 10 (Noben-Trauth et al., 2003). *Cdh23* encodes cadherin 23, a protein necessary for inner ear development and maintenance of the sensory hair cell structures, in particular the tip and lateral links of the stereocilia at the hair bundle (Siemens et al., 2004; Söllner et al., 2004; Kazmierczak et al., 2007; Michel et al., 2005; Lagziel et al., 2005), but growing anatomical evidence also suggests that synaptic rearrangements on sensory hair cells also contribute to auditory functional decline in C57BL/6J mice (Stamataki et al., 2006; Zachary and Fuchs, 2015; Jiang et al., 2015; Jeng et al., 2021). Surprisingly, aging C57BL/6J mice, although showing cochlear ribbon synapse degeneration and hearing loss, display increased acoustic startle reflex amplitudes (Ouagazzal et al., 2006; Ison and Allen, 2003; Ison et al., 2007) and a time increase in recovery from ABR short-term adaptation (Walton et al., 1995). Whether this exaggerated startle reflex can be explained by functional changes of cochlear hair cell ribbon synapses remain to be explored. Our present study will show that aged C57BL/6J mice, while losing nearly up to 50 % of

their IHC ribbon synapses, have IHCs with reduced BK channel clusters, larger remaining presynaptic

ribbons (with increased Ca^{2+} microdomains and larger sustained exocytotic responses) facing

postsynaptic afferent boutons with larger AMPAR clusters. All these features indicated that synaptic

release potentiation occur at auditory IHC ribbon synapses during aging. This age-related synaptic

plasticity could be viewed as compensatory mechanisms of the decrease IHC synapse number and may

explain the hyperacusis-like startle reflex observed in aged C57BL/6J mice.

RESULTS

Increased ABR thresholds and hyperacusis-like effect on ASR

As previously described (Zheng et al., 1999; Zhou et al., 2006), CBA/J mice, used here as a reference,

showed no sign of rapid degradation of their hearing function with aging, as attested by constant click

ABR thresholds near 15 dB SPL up to one year of age (Fig.1A). On the contrary, C57BL/6J mice,

carrying the $Cdh23^{753A}$ mutation, displayed an early progressive increase in ABR thresholds with aging

(Fig.1A). Hearing thresholds started to increase significantly at postnatal day P180 to reach a mean 50

dB increase at P365 (click ABR thresholds above 65 dB SPL) as compared to young mature P30 mice.

Distortion product otoacoustic emissions (DPOAEs) in P365 C57BL/6J mice were still present but

significantly reduced, indicating that the electromechanical activity of the outer hair cells (OHCs) was

also affected (Fig.1D). However, counting hair cells on surface preparations of organs of Corti from

these P365 mice, in the mid-cochlea region (8-16 kHz), indicated only a moderate loss of OHCs and

IHCs (statistically not significant; Fig.2E), in good agreement with the study of Zachary and Fuchs

(2015). This moderate loss of OHCs suggested that the profound hearing loss occurring at P365 in

C57BL/6J mice arose from additional factors than a simple alteration of the cochlear amplifier. We

then compared the amplitude on the ABR wave-I as a function of sound intensity in young and old

mice. The wave-I is known to result from the synchronous activity of the auditory nerve fibers and can

provide an objective measure of ribbon synapses loss when measured at high sound intensity above 70

dB SPL. In P365 old mice, ABR wave-I amplitudes were largely reduced as compared to young

mature P30 mice, even at high sound intensity when IHCs are directly stimulated and bypass the

amplification process of the OHCs (Fig.1B). The large decrease in wave-I amplitude indicated a

decreased number of auditory nerve fibers activated by sound and/or a decrease in their firing rate or

synchrony. The latencies of the ABR wave-I were also significantly increased suggesting again that

the IHC ribbon synapses are affected (Fig.1C).

Surprisingly, at lower sound intensities tested (between 20-35 dB above background noise level), the

strength of the acoustic startle reflex (ASR) was found larger in old P365 C57BL/6J mice as compared

to young P60, indicating hyperacusis reactivity (Fig 1.E). The origin of this acoustic hyperactivity at

low sound intensities could arise from central neural reorganization (Willott 1996) and be consecutive

to functional defect of the IHC ribbon synapses in aging C57BL/6J mice that we are going to

characterize in this study. At loud sound intensities, 45 dB above background noise, the amplitude of the ASR responses were however diminished in old mice, likely reflecting the increase in ABR thresholds due to a drastic decrease in the number of ribbon synapses per IHCs.

Figure 1. Sensorineural hearing loss with Aging in C57BL/6J mice. (**A**) Click-ABR thresholds as a function of age in CBA/J mice (blue; n = 5 to 16) and C57BL/6J mice (black, n = 6 to 15). For each strain, asterisks indicate significant difference with p<0.05 as compared to P30 mice (two-way ANOVA). Inset show comparative mean ABR waves at 70 dB SPL in P30 and P365 C57BL/6J mice. (**B**) Wave-I amplitude (P1-N1) as a function of sound intensity in young P30 (black) and old P365 (pink) C57BL/6J mice. Asterisks indicate significant difference with p<0.05 (two-way ANOVA; p = 0.001). (**C**) Latency to reach peak amplitude of wave-I as a function of sound intensity. Significantly different with p=0.003 between 80 to 50 dB SPL (two-way ANOVA; n is indicated in graph B). (**D**) Distortion products otoacoustic emissions (DPOAEs; 2f1-f2 with f1 = 12.73 kHz and f2 = 15,26 kHz) as a function of sound intensity in young P30 (black) and old P360 (pink) C57BL/6J mice. Asterisk indicated significant difference with p<0.05 (two-way ANOVA, p = 0.01). (**E**) Startle reactivity to acoustic stimulation. The intensity of startle reaction (in mV) was ln-transformed and obtained through two conditions of white-noise bursts of 20 and 40 ms. Note that below 35 dB above background the startle reaction is increased at P360 whereas at 45 and 55 dB above background the startle reaction is decreased compared to P30 mice. Asteriks indicated significant difference with p<0.05 (two-way ANOVA, p = 0.007 between 20-35 dB and p = 0.005 between 45-55 dB).

Drastic loss of IHC synaptic ribbons with aging

Confocal immunofluorescence imaging of the synaptic ribbons in IHCs from C57BL/6J mice, in the cochlear region encoding between 8-16 kHz, showed a progressive and large decrease

in the number of synaptic ribbons with aging, from 16.3 ± 0.3 ribbons/IHC at P15 to 8.7 ± 0.5 ribbons/IHC at P365 (Fig.2A-C). This nearly 50% loss of synaptic ribbons in IHCs of C57BL/6J mice at P365 was in good agreement with previous studies (Zachary and Fuchs, 2015; Jeng et al., 2020). Counting the spiral ganglion neurons in the mid-frequency region of the cochlea (8-16 kHz) showed a mean 20 % loss of neuron in P365 mice as compared to P30 mice (Fig.2D-F). Remarkably, the age-related loss of neurons, averaging 20 %, was much less pronounced than the loss of IHC synaptic ribbons (50 %), suggesting that there was a large number of neurons surviving without dendritic/synaptic projection toward the organ of Corti as previously shown by Stamataki et al. (2006).

Figure 2. Synaptic ribbon loss in aging IHCs and comparative density of spiral ganglion neurons. **(A-C)** Confocal immunofluorescence imaging of IHCs from P30 and P360 C57BL/6J mice. Image resulted from stack reconstructions of 22 slices of 0.3 μm thickness, showing F-actin (green, labeling the hair bundles), otoferlin (blue; a synaptic Ca^{2+} sensor labeling the entire IHC body) and synaptic ribbons (Ribeye/CtBP2, red dots). The white bars indicated 8 μm. **(B)** Plot of synaptic ribbon counts per IHC as a function of age in C57BL/6J mice (black) in the mid-frequency cochlear region (8-16 kHz). Asterisks indicate significant difference with p<0.05 (comparison with P15 C57BL/6J mice, one-way ANOVA post-hoc Tukey test). 203, 306, 306, 306, 295 ribbons were respectively analyzed at P15, P30, P60, P180, P365 in 3 to 5 mice at each age. **(D-F)** Confocal immunofluorescence imaging of cochlear transversal sections from P30 and P360 mice (8-16 kHz area) where neurons and their afferent dendrites are labeled with dextran amines 3000. **(E)** Analyzed in the mid-cochlea (8-16kHz), comparative histogram of hair cells counts, SGNs and synaptic ribbons (3 mice at each age relative to young mature P30 C57BL/6J mice. The asterisks indicated statistical difference with p < 0.05 (from left to right p = 0.37, p = 0.14, p = 0.020 for SGNs and p = 1.01E-14 for ribbons, using unpaired t-test). Note the reduction in SGNs and the drastic decrease of the fluorescent labeling of the afferent nerve fibers projecting to the organ of Corti.

Enlarged presynaptic ribbons associated with larger postsynaptic AMPAR clusters

When analyzing the 3-D imaging reconstruction of the IHC synaptic ribbons from P365 C57BL/6J IHCs from high resolution confocal STED immunofluorescent sections, we found that these structures displayed a progressive increase in their mean size (volume) as compared to young P15 mice (Fig.3A). The mean volume of the synaptic ribbons increased nearly three-fold between P15 and P365 (Fig.3B). Interestingly, the increase in size of the ribbons started very early after P30, an age at which their number also started to decrease (Fig.2B) while hearing thresholds were still normal (Fig.1A).

Figure 3. IHC synaptic ribbons progressively increased in size with aging. (**A**) Example of high magnification confocal STED images (stack reconstruction of five 0.25 μm slices). The yellow line delineates the IHC contour. Ribbons were less numerous and larger in size in IHCs from P365 C57BL/6J mice. (**B**) The mean volume of the IHC synaptic ribbons was calculated in the mid-cochlear region (encoding 8-16 kHz), using the plugin 3D-Object counter of ImageJ, and was plotted as a function of age: P15 (203 ribbons - 40 IHCs - 3 mice), P30 (306 ribbons - 98 IHCs - 5 mice), P60 (306 ribbons - 95 IHCs - 3 mice), P180 (306 ribbons - 67 IHCs - 3 mice), P365 (295 ribbons - 43 IHCs - 3 mice). As indicated by the asterisks, when comparing with P15, the mean volume of the ribbons started significantly to increase significantly from P30 (p<0.05; one-way ANOVA).

We next investigated the changes occurring with aging in the spatial distribution of the synaptic ribbons at the basolateral membrane of the IHCs. It is worth recalling that larger presynaptic ribbons associated to low-spontaneous-rate afferent fibers are usually found at the modiolar side of the IHCs (i.e. the baso-lateral curved side of the IHCs directed toward the spriral ganglion neuron) while smaller ribbons associated with high-spontaneous-rate afferent

fibers predominates at the pillar side (Liberman, 1982; Liberman et al., 2011). We found that,

while a larger proportion of synaptic ribbons distributed at the modiolar side in young P30

IHCs (~60 %), the surviving ribbons in old P365 IHCs were evenly distributed at the pillar

and modiolar side (Fig.4A-B). The loss of synaptic ribbons with aging was more pronounced

at the modiolar side, ie at the side where most of the low-spontaneous (high threshold) fibers

make contact to IHCs (Fig 4C). An increase in volume was noted both in pillar and modiolar

ribbons (Fig 4D). Unlike young P30 IHCs, old P365 IHCs had modiolar and pillar ribbons

with similar volume (0.7 ± 0.1 μm^3 and 0.57 ± 0.07 μm^3, respectively; not significantly

different, unpaired t-test p = 0.28, Fig.4D). Similar preferential loss of synaptic modiolar

ribbons associated with an increase in ribbon volume was also reported after noise-induced

hearing loss (Liberman and Kujawa, 2017; Hickman et al., 2020).

224
225

Figure 4. Modiolar vs pillar distribution of the synaptic ribbons with aging. (**A-B**) The distribution of IHC synaptic ribbons was splitted in two cell areas: P (pillar side) and M (Modiolar side). The two regions were determined using the K-means clustering method for each cell (see Methods). The yellow lines in the inset confocal images was drawn to arbitrarily delineate the frontier between pillar and modiolar ribbons (green labeling F-actin, blue Otoferlin and red the synaptic ribbons). Note that in young P30 IHCs (n=16), synaptic ribbons are preferentially localized in the modiolar region (62.3 ± 3.5 % vs 37.7 ± 3.5; unpaired t-test p = 2.07 E-5). In old P365 IHCs (n=11), the remaining ribbons equally distributed in the modiolar and pillar side (50.7 ± 4.0 % vs 49.3 ± 4.0 %, unpaired t-test; p = 0.81). (**C**) Synaptic ribbons were significantly loss both at the modiolar side and pillar side. However, the loss was much more pronounced in the IHCs' modiolar side (9.56 ± 0.5 ribbons vs 3.27 ± 0.24 ribbons; unpaired t-test p =5.57 E-10) and pillar (5.9 ± 0.6 ribbons vs 3.6 ± 0.6 ribbons, unpaired t-test p = 0.016). (**D**) The increase in volume affected both modiolar (P30=0.24 ± 0.02 µm³ vs P365=0.7 ± 0.1 µm³ unpaired t-test, p = 3.7 E-11) and pillar (P30= 0.12 ± 0.01 µm³ vs P365= 0.57 ± 0.07 µm³, unpaired t-test p = 7.5 E-14) synaptic ribbons.
This comparative analysis was performed in IHCs of the mid-cochlear region (encoding 8-16 kHz).

Postsynaptic afferent endings to ribbon IHCs express AMPA-type ionotropic glutamate [glutamate receptor A2 (GluA2)]-containing AMPA receptors (Matsubara et al., 1996; Hu et al., 2020). To determine whether the organization of postsynaptic AMPARs facing the IHCs ribbon was modified in old P365 mice we performed GluA2 immunostaining and confocal fluorescence imaging (Fig.5). As compared to young ribbon synapses, old synapses had larger ribbons associated with larger AMPAR clusters, suggesting that the changes occurring with aging are both at the pre and postsynaptic side in good agreement with the EM morphological analysis of Stamataki et al. (2006).

Figure 5. Larger postsynaptic patches of GluA2 receptors in old IHCs. (**A to D**) Immuno-confocal imaging of IHCs (stack reconstructions of 22 slices of 0.3µm thickness) showing otoferlin (blue), the synaptic ribbons (Ribeye/CtBP2, red dots)

and the post-synaptic AMPA receptors (GluA2, green dots). The white bars indicate 5 µm in A and B, then 0.5µm in C
and D. (C) Comparative distribution of the volume of postsynaptic patches of AMPA GluA2 receptors. Note that the
GluA2 receptor volume significantly increased with aging in C57Bl/6J mice ($p < 0.001$; unpaired-t test); (n = 86
synapses; from 3 mice at P30 (16 IHCs) and P365 (29 IHCs) ; mid-cochlear region encoding 8-16 kHz). (D) Comparative
volume distribution of postsynaptic GluA2 and corresponding postsynaptic ribbon in P30 and P365 IHCs. Note a good
linear correlation between GluA2 and ribbon volumes (Pearson's coefficient r = 0.93 and 0.86, respectively for P30 and
P360).

Drastic IHC cell size reduction wih aging

To determine whether the loss of synaptic ribbons was associated with a cell size reduction in

old P365 IHCs, we labeled the cytosol of IHCs with an antibody against myo-7a, an

unconventional myosin essential for the hair bundle functional integrity and found in the

apical sterocilia as well as the cytoplasm of the hair cells (Hasson et al., 1995). We found a

drastic reduction of the basal infra-nuclear synaptic zone of the IHCs from an average of 15.7

$\pm$ 0.1 µm (n=34) in P21 IHCs to 12.4 $\pm$ 0.3 µm (n=50) in old P365 IHCs (unpaired-t test, p =

1.1 E-13), Fig. 6A-C). The cell size reduction of old IHCs was confirmed by another method

using whole-cell patch clamp measurement of the IHC resting membrane capacitance in

organs of Corti preparation ex-vivo (Fig.6D-E). P365 IHCs displayed a significant smaller

resting capacitance, measured at -70 mV, of 7.9 $\pm$ 0.4 pF (n=22) as compared to P21 IHCs

(10. 8 $\pm$ 0.1 pF, n=44, unpaired t-test, p = 3.7 E-10).

Figure 6. Cell size reduction of IHCs with aging. (A-B) Immuno-confocal imaging of IHCs labeled for myo-7a (blue) and CtBP2 (labeling the ribbons, red) in P21 (A) et P365 (B) mice (reconstructed stack images of 22 slices of 0.3 µm thick). The length of the infra-nuclear synaptic zone was measured from the center of the nucleus to the bottom of the cell as indicated by the yellow line. The scale bar indicated 5 µm. (C) Comparative histogram of the length of the infra-nuclear

280 synaptic zone. Asterisks indicated statistical significance with p < 0.0001, unpaired t-test.) Measurements were made
281 from IHCs of the mid-cochlear region (8-16 kHz). (D) Photomicrograph showing a row of IHCs in an ex-vivo organ of
282 Corti explant and a patch-clamp recording electrode allowing measurements of capacitive and ionic currents. (E)
283 Comparative mean IHC resting membrane capacitance, measured at -70 mV, in whole-cell patch-clamp configuration.
284 Asterisks indicated values with significant difference with p < 0.0001, unpaired-t test.
285

287 Higher Ca^{2+} current density and exocytosis in old IHCs

Although the loss of IHC ribbon synapses in C57BL/6J mice with aging is well documented

in the literature (see Wan and Corfas, 2015), the functional changes of the remaining "old"

IHC ribbon synapses has not yet been explored in detail at the cellular level. We investigated

here the changes in the calcium-evoked exocytotic responses of aging IHCs from ex-vivo

explants of the organ of Corti from aging C57BL/6J mice. We recall that Ca^{2+} is a key

messenger for normal hearing since it tunes the transduction of the analogic sound signal into

nerve impulses. This transduction occurs at the IHC synaptic active zone through the voltage-

activation of $Ca_v1.3$ Ca^{2+} channels (Brandt et al., 2003; Vincent et al., 2017) and the action of

the Ca^{2+} sensor otoferlin (Roux et al., 2006; Beurg et al., 2010; Michalski et al., 2017).

Increase in the density of L-type Ca^{2+} channels has long been recognized as a characteristic of

aging central neurons (Thibault and Landfield, 1996; Campbell et al., 1996) but possible

altered function in Ca^{2+} signaling has not been investigated in aging IHCs.

Figure 7. Ca^{2+} currents and exocytosis are exacerbated in IHCs from aged C57BL6/J mice. (**A**) Ca^{2+} current density of old P365 IHCs (pink) was larger as compared to young mature P21 IHCs (black). IHCs were subjected to a voltage ramp stimulation from -90 mV to +30 mV in 120 ms, giving a slope change in voltage of 1mV/ms. Comparative I_{Ca} maximum density measured at -10 mV indicated a significant increase in P360 IHCs (unpaired two-sample t-test, p=0.03) while the half-max voltage activation ($V_{1/2}$) of the Ca^{2+} currents remained unchanged (unpaired t-test, p = 0.06). (**B**) Kinetics of exocytosis were evoked by a voltage-steps from -80 mV to -10 mV with increasing duration from 5 ms to 80 ms. Note that brief stimulations below 25 ms, a time-frame addressing the release of the Readily Releasable Pool of vesicles, the exocytotic response (normalized to cell size) was similar in P30 and P360 IHCs (p = 0.2; two-way ANOVA). Longer stimulations, above 30 ms, which probed the activation of the secondarily releasable pool of vesicles, showed a significant increase in exocytosis (two-way ANOVA; p = 2.3 E-8), indicating that old IHCs were more likely to sustain high rates of vesicular release. (**C-D**) Representative examples of ΔC_m and I_{Ca}^{2+} recordings from P30 and P365 IHCs during a 20 ms and a 100 ms voltage-step stimulation from -80 mV to -10 mV. The number of IHCs, indicated in graph, were obtained from 5 mice at P365 and 6 mice at P30.

In the present study, whole cell patch-clamp recordings from old P365 IHCs showed a significantly increase in Ca^{2+} current density (Ca^{2+} current amplitude normalized to cell size) as compared to young mature P30 IHCs, rising from a mean peak value of 15.8 ± 1.75 pA/pF to 20.1 ± 1.9 pA/pF at -10 mV (Fig. 7A; unpaired t-test, p <0.05). The voltage-activation

curve of the Ca^{2+} currents in old P365 IHCs remained similar to the young mature P30 IHCs,

with a mean half-voltage activation ($V_{1/2}$) of -29.3 ± 1.1 and -26.7 ± 2.8 mV, respectively

(unpaired t-test, p = 0.06).

Patch-clamp measurement of membrane capacitance upon depolarizing voltage-steps from -

80 to -10 mV with increasing time duration allowed to compare the kinetics of vesicle

exocytosis in young and old IHCs. For brief stimulations below 25 ms, a time-frame

addressing the release of the Readily Releasable Pool (RRP) of vesicles, the exocytosis

response (normalized to cell size) was similar in P30 and P365 IHCs (Fig. 7B-C). However,

longer depolarization above 25 ms revealed a more sustained exocytotic response in old IHCs,

suggesting the mobilization of a more efficient secondarily releasable pool (SRP), i.e. a more

efficient vesicle recruitment in these cells. To investigate whether this increased SRP

exocytosis was due to higher Ca^{2+} microdomains at the ribbon active zones, we used the high

affinity Ca^{2+}-indicator Rhod-2 (K_d = 570 nM) which is known to allow a good monitoring of

low Ca^{2+} concentrations ranging from 20 nM to 20 µM (Del Nido et al., 1998). We choose to

use Rhod-2 rather than a low affinity Ca^{2+} dye such as OGB-5N with a Kd at 20 µM (Vincent

et al., 2014) because we wanted here to primarily probe the amplitude of the Ca^{2+}

microdomains, where the changes in Ca^{2+} concentrations are due to Ca^{2+} diffusion and are

lower than at the nanodomain range near the Ca^{2+} channels. Ca^{2+} microdomains allow vesicle

replenishment, a process occurring at the µm scale away from the ribbon and their associated

Ca^{2+} channels (Spassova et al., 2004; Johnson et al., 2005; Graydon et al., 2011). We found

that, upon 5 consecutive long 100 ms voltage-step stimulations (from -80 mV to -10 mV),

IHCs displayed brighter transient Ca^{2+} hot spots (measured in 1µm^2 areas) at their basal

synaptic area in P365 mice as compared to young P30 IHCs (Fig. 8A-B). This indicated the

expression of synaptic Ca^{2+} microdomains with larger amplitude in old P365 IHCs, a response

likely due a larger Ca^{2+} channel density at their ribbon active zones (Fig. 7A).

Simultaneously, in the same cells, the application of the 5 consecutive 100 ms stimulations,

generated a facilitated vesicular recruitment in old P365 IHCs (Fig.8D). Also as a

consequence of increased Ca^{2+} microdomains in old IHCs, the simultaneous recording of

Ca^{2+} currents showed an enhanced time-inactivation, indicative of an increased Ca^{2+}-

dependent inactivation due to higher intracellular Ca^{2+} concentration (Fig. 8C).

Figure 8. Larger synaptic Ca^{2+} signaling and increased sustained exocytosis in aged IHCs. (**A**) Example of confocal Ca^{2+} imaging, using the intracellular fluorescent Ca^{2+} indicator Rhod2, in an IHC. The cell was voltage-clamped at -80 mV and stimulated by 5 (100 ms) consecutive voltage-step to -10 mV. The photograph show two examples of square zones (1µm^2) below the cell nucleus where voltage-induced calcium increase was measured. (**B**) Comparative mean amplitude of transient fluorescent hot spots showed higher Ca2+ responses at the presynaptic active zones of old P365 IHCs (pink) as compared to young mature P30 IHCs (black). Hair cells were depolarized by 5 five consecutive 100 ms depolarizing steps from -80 mV to -10 mV. The values n in the graph indicated the number of active zones analyzed for young (6 IHCs from 6 mice) and old mice (5 IHCs from 5 mice), respectively. The peak of the 5 calcium responses ($\triangle F=F-F_0$ with F_0 as the fluorescent base line value averaged 0.5 s before stimulation) were significantly larger in old IHCs (two-way ANOVA (p = 8.8 E-5). The histogram compared the overall fluorescent response $\triangle F/F_0$ occurring during the 5 stimulations (0.5 to 2.5 s). The integral of the Ca^{2+} response (Area Under the Curve, AUC) was significantly larger in old P365 IHCs (unpaired t-test, p = 0.03). (**C**) Comparative mean of calcium current recorded at the same time during the 5 stimulations shown in B. Note that the inactivating part of the calcium current is increased in P365 IHCs. Asterisk indicate statistical analysis with unpaired t-test. (**D**) The concomitant exocytosis was also measured during the 5 stimulations shown in B Exocytosis measured as the variation of membrane capacitance was normalized for each IHC to their ribbon number or cell size; both normalization giving similar results). The exocytotic responses were larger in old P365 IHCs. The asterisks indicated statistical significant difference with p<0.05 (two-way ANOVA; p = 0.0015).

Aging promoted autophagy in IHCs through the recruitment of LC3B

The size reduction of the old IHCs (Fig.6) suggested that the degenerative process of the ribbon synapses was associated with a concomitant reduction of the cell surface membrane of IHCs during aging, suggesting a largely augmented uptake of plasma membrane through endocytosis. Like other type of cells, such as neurons or retinal cells, auditory hair cells utilize autophagic pathways to maintain cell homeostasis during excessive ROS production (Defourny et al., 2019; Fu et al., 2018). Autophagy is a degradation process where cytoplasmic and plasma membrane components are engulfed by vesicles called autophagosomes. To determine whether this process could explain the IHC size reduction with aging, we investigated the recruitment of LC3B, a structural protein of autophagosomal membranes. We assessed and compared the autophagosome numbers and surface area in young P30 and old P365 IHCs by quantifying LC3B puncta numbers/cell by using confocal immunofluoresence. We found that the number and overall surface area of LC3-positive structures in IHCs largely increased with aging (Fig.9), indicating an increased autophagosome formation. Similar investigations in P365 CBA/J mice, a strain which does not display early age-related hearing loss (Fig. 1), showed a weak expression of LC3B in IHCs (Fig. 9).

Figure 9. LC3-positive structures are prominent in aging IHCs. Representative confocal immunofluorescence images showing LC3B staining in IHCs from young P30 C57BL/6J (left), old P365 CBA/J (middle) and old P365 C57BL/6J (right) mice.The histograms below show the quantification of the LC3B puncta average number and average mean surface area per IHC. Images were obtained from processing confocal stack images (22 of 0.3 µm thickness) using ImageJ quantification tool (3D-object counter). Quantification was performed from 25 to 30 IHCs for each condition in the mid-cochlea (encoding region 8-16 kHz; 3 mice). Asterisks indicate statistical significance p < 0.05 (unpaired t test).

Clustering and voltage-dependent activation of BK channels are affected aging in IHCs

There is increasing evidence that oxydative damage due to an increase in ROS levels plays a crucial role in age-related degeneration of the synaptic ribbon synapses (Fu et al., 2018; Rousset et al., 2020). BK ($K_{Ca1.1}$) channels are known to be highly sensitive to oxidative stress and their activity is a good indicator of intracellular ROS production (Sahoo et al.,

2014; Hermann et al., 2015). Interestingly, a decrease in BK channel expression in IHCs has

been associated to noise-related ROS and peroxysome pathology (Delmaghani et al., 2015)

but never yet described in IHCs during age-relate hearing loss. It is worth recalling that BK

channels allows fast repolarization of IHCs and are essential for phase-locking of the auditory

afferent neurons (Kros et al., 1998; Skinner et al., 2003; Oliver et al., 2006; Kurt et al., 2012).

We here investigated, using immunofluorescent confocal microscopy, the comparative level

expression of BK channels in IHCs from young (P30) and old P365 C57BL/6J mice. In young

mature IHCs, BK channels are known to be organized in a dozen of clusters at the neck of the

IHCs below the cuticular plate (Pyott et al., 2004; Hafidi et al., 2005). We used 3-D

reconstructions from stacks of confocal images of immunostained organs of Corti (mid-

cochlear frequency region 8-16 kHz) to determine the mean total surface area of the BK

clusters per IHCs (Fig.10A-C). The results indicated that the membrane surface covered by

BK channel clusters was reduced almost threef

old in P365 IHCs as compared to P30 IHCs.

To further characterize the changes in BK channel function with aging, we performed whole

cell patch-clamp recordings in IHCs from P30 and P365 mice. Since BK currents in IHCs

have the peculiar property to be extremely fast activating currents (also termed IK_f; Kros et

al., 1998), these currents were recorded during brief 20 ms depolarizing steps and measured

2.5 ms after their onset (Fig.10D-F). BK currents in IHCs from young mature have also the

particularity to activate at extremely negative membrane potential (near -60 mV) (Lingle et

al., 2019). While the whole-cell conductance of BK channels from P365 IHCs was not

significantly different from P30 IHCs (0.26 .± 0.04 nS and 0.23 ±.0.03 nS, respectively), their

voltage-dependant activation showed a large positive shift of 7 mV (Fig.10F). Interestingly,

the decrease in size of the BK patches and the positive voltage-activation shift in old IHCs

somewhat resembled the phenotype observed in mice KO for LRRC52, a γ BK regulatory

subunits (Lang et al., 2019; Lingle et al., 2019), suggesting that the BK complex formed with

this γ subunit could be one the target of the degenerative oxydative process in aging

C57BL56/J mice IHCs.

Figure 10. Immunofluorescent confocal imaging of BK channels in aging IHCs. (**A**) Numerous large BK channel clusters (red) are localized extrasynaptically below the cuticular plate of IHCs in young mature P30 C57BL6/J mice. (B) Note the strong reduction in size of the BK channels clusters in IHCs of old P365 C57BL6/J mice.
(C) Histogram comparing the total surface area covered by the BK channel clusters per IHC in the mid-turn cochlea (region encoding 8-16 kHz). The total number of IHCs analyzed was 58 and 25 from P30 (n=3) and P365 (n=3) mice, respectively. Asterisks indicate statistical significance with p<0.05 (p = 1.12 E-7; unpaired t test).
(**D**) Comparative fast outward potassium currents (IKf), carried by BK channels, recorded from P30 (black, n=12) and P360 (pink, n=8) IHCs when using a brief depolarization step of 20 ms from -80 to -10 mV.
(**E**) Comparative BK current-voltage curves (n = 12 and 8 IHCs respectively in P30 and P360 mice, from 3 mice at each age). Significantly different two-way ANOVA, p = 2.7 E-8. IKf was measured 2.5 ms after the onset of the voltage-step stimulation. (**F**) Comparative BK conductance fitted with a Boltzmann curve indicated a V1/2 of -20.5 ± 5.7mV in P30 IHCs and of -13.7 ± 1.4 mV in P365 IHCs, respectively (unpaired t-test, p = 0.009). Note that voltage error across series resistance (Rs = 5 ± 2 MΩ) and leak current were compensated for each IHC. All values are expressed as means ± SEM.

DISCUSSION

Larger synaptic ribbons associated with increased Ca^{2+} signals

Ribbons are presynaptic organelles tethering glutamate-filled vesicles at the hair cell active zone (AZ). These intracellular structures, composed of the protein RIBEYE, are anchored by

the scaffold protein bassoon to the AZ plasma membrane, adjacent to Ca^{2+} $Ca_V1.3$ channels

(Brandt et al, 2003; Graydon et al., 2011; Jing et al., 2013; Vincent et al., 2017). Ribbons are

essential for high rate of transmitter release and for high temporal precision of neuronal

encoding of sound at the auditory afferent nerve fibers (Jean et al., 2018; Becker et al., 2018).

Each auditory IHC have 10 to 20 synaptic ribbons depending of their cochlear frequency

location (Meyer et al., 2009). Ribbons are heterogeneous in size and each ribbon makes a

single synaptic contact with one afferent nerve fiber. Remarkably, fibers associated to large

ribbons display low spontaneous rate and high-threshold have the tendency to be distributed

at the modiolar side of IHCs (Liberman et al., 2011). Nerve fibers associated with smaller

ribbons display high spontaneous rate and low-threshold activation, are mainly located toward

the pillar side of IHCs. Interestingly, we found here that old C57BL6/J IHCs preferentially

loose modiolar ribbons (known to be associated with high threshold low spontaneous fibers),

therefore mimicking what is happening during noise-induced hearing loss (Fernandez et al.,

2015; Liberman and Kujawa 2017; Hickman et al., 2020). Heterogeneity in the voltage-

dependence of Ca^{2+} channels and release site coupling has been recently proposed to

contribute to the firing fiber specificity, with modiolar ribbons being activated at more

negative potentials (Özçete and Moser 2021). In our study, we did not find evidence for a

voltage-shift in the voltage-dependence of Ca^{2+} channels with aging but we did observe an

increase in the density of Ca^{2+} currents. The increase in Ca^{2+} entry was further confirmed by

the stronger time-inactivation of the Ca^{2+} currents (Grant and Fuchs, 2008; Vincent et al.,

2017).

One of the most striking observations was that the IHC synaptic ribbons underwent a nearly

threefold increase in size with aging, confirming a certain degree of plasticity of these

synaptic structures. The enlargement of the presynaptic ribbons was in good agreement with a

previous ultrastructural study in aging C57BL/6J mice by Stamataki et al. (2006) and the

recent study of Jeng et al., (2020). Interestingly, the size of the young mature IHC synaptic

ribbons is known to correlate positively with the AZ Ca^{2+} microdomain amplitude, as well as

the number and gating of the $Ca_v1.3$ Ca^{2+} channels (Frank et al., 2009; Neef et al., 2018).

Also, hair cells of transgenic zebrafish with enlarged ribbons display larger ribbon-localized

Ca^{2+} currents (Sheets et al., 2017). In agreement with these previous studies, we found here,

in IHCs from aging C57BL/6J mice, that the remaining larger ribbons were associated with

larger Ca^{2+} microdomain amplitude, and larger density of calcium currents, reinforcing the

idea that Ca^{2+} ions regulate the size of the synaptic ribbons. In this context, It is worth

recalling that mitochondrial-Ca^{2+} uptake has been proposed to lower cellular NAD^+/NADH redox and regulates ribbon size, possibly by acting through the NAD^+ binding site of the B domain of RIBEYE (Wong et al., 2019). In vitro work has shown that NAD^+ and NADH can promote interactions between ribeye domains (Magupalli et al., 2008) and the size of the ribbons would be then sensitive to the subcellular synaptic NAD^+/NADH redox level (Wong et al., 2019), suggesting that the increase in size of the IHCs ribbons from aged C57BL/6J mice also results from NAD^+/NADH homeostasis disruption.

Figure 11. Synaptic IHC degeneration was associated with a strong potentiation of the remaining ribbon synapses in aging C57BL/6J mice. (**A**) Age-related hearing loss in P360 mice was associated with IHC shrinkage (cell size reduction), an important loss of afferent ribbon synapses (>50%), a drastic enlargement of the surviving synaptic ribbons (both at pillar and modiolar side), an increase in Ca^{2+} signaling (larger Ca^{2+} microdomain amplitude as indicated by the larger pink domains around the ribbons) and a large decrease in the fast repolarizing BK channel clusters (blue bars). (**B**) Hypothetical view of a young and old IHC ribbon synapse.

Increased capacity for sustained IHC synaptic exocytosis: Implication for the discharge pattern of the auditory nerve firing

As a consequence of larger Ca^{2+} microdomains, we found that IHCs from old C57BL/6J mice had larger and more sustained exocytotic responses. These larger release responses could also well be explained by a higher synaptic vesicle density at the active zone of the IHC ribbon synapses as suggested by the EM study of Stamataki et al. (2006). Surprisingly, the recent study of Jeng et al., (2020), which did not perform Ca^{2+} imaging, concluded that the size and kinetics of Ca^{2+}-dependent exocytosis in IHCs were unaffected in aging mice but this study did not normalize the IHC exocytotic responses to their cell size or to their ribbon synapse number, and therefore may have overlooked the differential phenotype.

What would be the consequences on the discharge pattern of the remaining afferent fibers of a more sustained release capacity in old ribbon synapses ? Fast adaptation in the discharge rate of the afferent auditory nerve fibers is thought to be one of their essential property for emphasizing the timing information of sound stimuli. This adaption takes the form of a fast increase in spike rate at stimulus onset that rapidly declines in a double exponential decay to a lower steady-state rate after a few tens of milliseconds (Taberner and Liberman 2005). The origin of this fast adaptation has been proposed to arise first from H^+ release block of Ca^{2+} channels and second to rapid depletion of synaptic vesicle release (Vincent et al., 2017). We predict that the increased capacity to sustain SRP exocytosis in old IHCs with larger ribbons would have the tendency to decrease the level of firing adaptation of the auditory fibers and therefore explain the altered recovery from short-term adaptation in old C57BL/6J mice (Walton et al., 1995).

Furthermore, we confirmed here the hyperacusis-like effect of the startle responses in old C57BL/6J mice (Ison et al., 2007). This increased response at low sound intensity could be explained by a release potentiation at the IHC ribbon synapses due to the presynaptic higher Ca^{2+} channel density associated with a larger ribbon coupled with a larger vesicular SRP, and increased expression of post-synaptic AMPARs. All these features would make the auditory ribbon synapses hypersensitive and underlie the hyperacusis-like reflex in old C57BL/6J mice.

IHC shrinkage and disruption of BK channel clusters

Another novel observation of our study was the drastic reduction in cell size of IHCs from aging C57BL/6J mice. Interestingly, tuberous sclerosis rapamycin-sensitive complex 1

(mTORC1) has been shown to be overactivated and promote oxidative damages to synapse

ribbon loss in aging C57BL/6 mice (Fu et al., 2018). We found that aged P365 IHCs had an

increase expression of the autophagy marker LC3B. The activation of the mTOR pathway

with aging could therefore trigger the cell size reduction of IHCs, by activating the autophagy

pathway, as proposed in many other cell types (Fingar et al., 2002; Fumarola et al., 2005). In

addition, our study described another novel remarkable feature of aging IHCs, a drastic

disruption of the BK channel clusters, suggesting that these potassium channels are also a

good indicator of the redox state and autophagy in IHCs. The expression and distribution of

BK channels could be regulated through the rapamycin insensitive mTOR complex 2

(mTORC2) as shown in the kidney glomerulus podocytes (Wang et al., 2019). Interestingly,

the disruption of the BK channel clusters in aging IHCs was associated with a large positive

shift in their voltage-activation curve, suggesting that the fast repolarizing property mediated

by BK channels which likely contributes to decrease transmitter release (Skinner et al., 2003),

would be largely hampered during sound stimuli in aging mice, an additional factor that may

also contribute to hyperacusis.

CONCLUSIONS

In conclusion, our study unraveled several novel functional and structural features of aging in

IHC ribbon synapses from C57BL/6J mice (Fig.10): 1) a large reduction in IHC cell size, 2) a

drastic enlargement of the presynaptic ribbons and postsynaptic AMPAR clusters 3) an

increased presynaptic Ca^{2+} signaling associated with a stronger sustained exocytotic response

4) a disruption of BK channel clusters and a drastic shift in their voltage-dependence

decreasing their potential negative feedback on neurotransmission. Overall, these results

suggested that aging IHCs ribbon synapses can undergo important structural and functional

plasticity that lead to synaptic potentiation that could explain the paradoxical hyperacusis-like

exaggeration of the acoustic startlle reflex observed in C57BL/6 mice.

MATERIALS and METHODS

Mice

Inbred C57BL/6J mice, which are homozygous for the recessive cadherin23 mutation

$Cdh23^{753A}$ (also known as Ah1) were used in the present study. Hearing thresholds and IHC

ribbon counts in mice of either sex were performed at postnatal ages: P15, P30, P60, P180

and P365. CBA/J mice (from Charles River) were also tested as controls at P50, P250 and P365. Animal care and all procedures were approved by the institutional care and use committee of the University of Bordeaux and the French Ministry of Agriculture (agreement C33-063-075).

Assessment of hearing thresholds (ABRs)

To record ABRs (Auditory Brainstem Responses, which represent the sound-evoked synchronous firing of the auditory cochlear nerve fibers and the activation of the subsequent central auditory relays), mice were anesthetized with intraperitoneal injection of xylazine (6 mg/ml, Rompun Bayer) and ketamine (80 mg/ml, Virbac) mixture diluted in physiological saline. The mouse was placed in a closed acoustic chamber and its body temperature was kept constant at 37°C during ABRs recording. For sound stimulus generation and data acquisition, we used a TDT RZ6/BioSigRZ system (Tucker-Davis Technologies). Click-based ABR signals were averaged after the presentation of a series of 512 stimulations. Thresholds were defined as the lowest stimulus for recognizable wave-I and II. The amplitude of ABR wave-I was estimated by measuring the voltage difference between the positive (P1) and negative (N1) peak of wave-I. Sound intensities of 10 to 90 dB SPL in 10 dB step, were tested.

Distortion product otoacoustic emissions (DPOAEs), which originate from the electromechanical activity of the outer hair cells (OHCs), were tested by using two simultaneous continuous pure tones with frequency ratio of 1.2 (f1 = 12.73 kHz and f2 = 15,26 kHz). DPOAEs were collected with the TDT RZ6/BioSigRZ system designed to measure the level of the "cubic difference tone" 2f1-f2.

Acoustic Startle Reflex (ASRs) measurements

Testing was conducted in four acoustic startle chambers for mice (SR-LAB, San Diego Instruments, San Diego, CA). Each chamber consisted of a nonrestrictive cylindrical enclosure attached horizontally on a mobile platform, which was in turn resting on a solid base inside a sound-attenuated isolation cubicle. A high-frequency loudspeaker mounted directly above the animal enclosure inside each cubicle produced a continuous background noise of 65 dBA and various acoustic stimuli in the form of white noise. Mechanical vibrations caused by the startle response of the mouse were converted into analog signals by a piezoelectric accelerometer attached to the platform. A total of 130 readings were taken at 0.5-ms intervals (i.e., spanning across 65 ms), starting at the onset of the startle stimulus. The

average amplitude over the 65 ms was used to determine the stimulus reactivity. The sensitivity of the stabilimeter was routinely calibrated to ensure consistency between chambers and across sessions.

Acoustic startle reflex was assessed during a session lasting for approximately 30 minutes, in which the mice were presented with a series of discrete pulse-alone trials of different intensities and durations. Ten pulse intensities were used: 69, 73, 77, 81, 85, 90, 95, 100, 110, and 120 dBA lasting for either 20 or 40 ms (background noise level: 65 dBA). A session began when the animals were placed into the Plexiglas enclosure. The mice were acclimatized to the apparatus for 2 minutes before the first trial began. The first six trials consisted of six pulse-alone trials of 120 dBA, comprising three trials of each duration. These trials served to stabilize the animals' startle response, and were analyzed separately. Subsequently, the animals were presented with five blocks of discrete test trials. Each block consisted of 20 pulse alone trials, one for each intensity and duration. All trials were presented in a pseudorandom order with an inter-trials interval of 14 seconds.

Tissue preparation and imunocytochemistry

Mice were deeply anesthetized with isofluorane (Vetflurane, Virbac). The ramps of the cochlear apparatus were dissected and prepared as previously described by Vincent et al. (2014). Inner ears (cochleae) were fixed by incubation in 4% paraformaldehyde in phosphate-buffered saline (PBS), pH 7.4, at 4°C overnight and washed with cold phosphate buffered saline (PBS). They were then incubated for 2 days in PBS solution containing 10% EDTA pH 7.4 at 4°C. The middle part of the organ of Corti (area encoding between 8 and 16 kHz) was then dissected and the tectorial membrane removed. The tissue was first incubated with PBS containing 30% normal horse serum and triton X100 0.5 % for 1 hour at room temperature. Synaptic ribbons were labeled with anti-CtBP2 (1/200 Goat polyclonal, Santa Cruz, USA ; cat # SC-5966 or 1/200 Mouse IgG1 monoclonal antibody BD Biosciences, cat # 612044). Otoferlin was labeled with a mouse monoclonal antibody (1/200 AbCam, cat # ab53233). Autophagy was detected by labeling LC3B protein with anti Rabbit monoclonal antibody (1/200 LC3B(D11) XP, Cell Signaling Technology, cat # 3868). BK channels were labeled with a Rabbit polyclonal antibody (1/200, Alomone, Israel, cat #APC-021. Explants of Organ of Corti were incubated overnight at 4°C with primary antibodies. The following fluorescent secondary antibodies 1/500 were then used: anti-Goat Donkey polyclonal Fluoprobes 547H (Interchim, cat# FP-SB2110), anti-Mouse Donkey polyclonal Fluoprobes 647H (Interchim,

cat# FP-SC4110), anti-Mouse Donkey polyclonal Alexa Fluor 488 (AbCam, cat# ab150109),

anti-Mouse Goat polyclonal Alexa Fluor 546 (Molecular Probes, cat# A-11003), anti-Mouse

Goat polyclonal Alexa Fluor 488 (Molecular Probes, cat# A-10680), anti-Rabbit Donkey

polyclonal Fluoprobes 594 (Interchim, cat# FP-SD5110). Actin-F was also used to visualize

hair cells (1/100, Phalloidin Fluoprobe 405, Interchim, Montlucon, France; cat # FP-

CA9870). For Image Acquisition, organ of Corti samples were analyzed using a confocal

laser scanning microscope Leica SP8 with a 63X oil immersion objective (NA = 1.4) and

white light laser (470 to 670 nm) (Bordeaux Imaging Center). Phalloidin was imaged by using

a diode laser at 405 nm also mounted on the microscope. For complete 3D-stack

reconstruction of IHCs, 25 to 30 images (0.3 µm thickness) were acquired as previously

described (Vincent et al, 2014). Ribbon volumes were calculated using the 3D-Objects

Counter Plugin of ImageJ. Image resolution was 0.038 µm/pixel. As previously described by

Hickman et al. (2020) and Jeng et al., (2020), the x,y,z coordinates of ribbons in each z-stack

were transformed into a coordinate system based on the modiolar-pillar polarity of the IHCs.

We determined the pillar versus modiolar distribution of the synaptic ribbons by using a *k*-

means clustering method in the z-stack position of each ribbon, with arbitrarily choosing the

number of cluster as 2 (OriginPro 9.1software, OriginLab, Northampton, USA). From these 2

clusters, we determine arbitrarily the border between the modiolar and pillar side in each IHC

by taking the half-distance between the two nearest ribbon of each cluster.

High resolution imaging of synaptic ribbons with confocal STED microscopy

For high-resolution imaging of IHC ribbons, we used a Leica DMI6000 TCS SP8 STED

microscope with 93X glycerol immersion objective (NA 1.3) (Bordeaux Imaging Center).

Mouse sensory organs were fixed and incubated with primary anti CtBP2 (1/100) antibodies

in conditions similar to those described above for confocal microscopy. Secondary antibodies

Goat anti- IgG1 Mouse polyclonal Alexa Fluor 647 (Jackson, cat# 115-605-205) were used to

label CtBP2 primary antibodies. The pulsed STED depletion laser (pulsed laser IR, Spectra

Physics Mai Tai) was set at 775 nm. Stack images were acquired with the following

parameters: scan rate 200 Hz, 775 nm depletion laser power between 8 and 15%, scans frames

accumulated per *X-Y* section 8 times, z- step size 0.25 µm, number of plans between 8 and

10, pixel size 20 nm giving an *X-Y* image size of 21 × 5 μm (1024 × 256 pixels).

Counting spiral ganglion neurons

The afferent nerve fibers of the auditory nerve were fluorescently labeled using a retrograde

track tracing technique with dextran amines (Fritzsch et al., 2016). Cochleae were extracted

from temporal bone of freshly euthanised mice and rapidly immerged intact into ice-cold

artificial perilymph (NaCl 135; KCl 5.8; $CaCl_2$ 1.3; $MgCl_2$ 0.9; NaH_2PO_4 0.7; Glucose 5.6;

Na pyruvate 2; HEPES 10, pH 7.4, 305 mOsm). A small crystal of tetramethylrhodamine

dextran amines 3000 (Life technologies, NY) was placed for 15-20 min at the edge of the

auditory nerve coming out at the internal auditory canal of the intact cochlea. The

intracochlear sensory tissues were then fixed by gentle perfusion of a 4% PFA solution for 30

min and rinse with artificial perilymph. They were then decalcified for 2 days in PBS solution

containing 10% EDTA pH 7.4 at 4°C. The cochleae were then sliced with a razor blade along

the longitudinal cochlear axes and the slices were then mounted on a glass-slide for

fluorescent confocal microscopy observation. The quantification of the spiral ganglion

neurons per surface area of the neural ganglion were made in the middle part of the cochlea in

a region encoding 8-16 kHz, by using the 3D object counter application of ImageJ software.

Ca^{2+} currents and whole-cell membrane capacitance measurement in IHCs

Measurements of resting membrane capacitance of IHCs were performed at P30 and P365

mice in the 20-40% normalized distance from the apex, an area coding for frequencies

ranging from 8 to 16 kHz, by using an EPC10 amplifier controlled by Patchmaster pulse

software (HEKA Elektronik, Germany), as previously described in detail (Vincent et al.,

2014, 2015). The organ of Corti was incubated in an extracellular perilymph-like solution

containing NaCl 135; KCl 5.8; $CaCl_2$ 5; $MgCl_2$ 0.9; NaH_2PO_4 0.7; Glucose 5.6; Na pyruvate

2; HEPES 10, pH 7.4, 305 mOsm. This extracellular solution was complemented with 0.25

µM of apamin (Latoxan; cat # L8407) and 1 µM of XE-991 (Tocris Bioscience; cat # 2000) to

block SK channels and KCNQ4 channels, respectively. The external Ca^{2+} concentration was

increased from 1.3 to 5 mM to enhance the amplitude of Ca^{2+} currents to levels nearby body

temperature. All experiments were performed at room temperature (22°C–24°C).

Patch pipettes were pulled with a micropipette Puller P-97 Flaming/Brown (Sutter Instrument,

Novato, CA, USA) and fire-polished with a Micro forge MF-830, (Narishige, Japan) to obtain

a resistance range from 3 to 5 MΩ. Patch pipettes were filled with a cesium-based

intracellular solution containing (in mM): CsCl 145; $MgCl_2$ 1; HEPES 5; EGTA 1; TEA 20,

ATP 2, GTP 0.3, pH 7.2, 300 mOsm. Measurements of the resting membrane capacitance

(cell size) of IHCs were obtained in whole-cell voltage-clamp configuration at -70 mV and

after 2 min equilibrium of the internal patch-pipette recording solution with the IHC cytosol.

BK currents

Whole-cell recordings of BK currents were recorded in P30 and P365 IHCs from ex-vivo

whole-mount preparation (Skinner et al., 2003) as described above for Ca^{2+} currents.

Recording pipettes were filled with a KCl-based intracellular solution containing in mM: 158

KCl, 2 $MgCl_2$, 1.1 EGTA, 5 HEPES, and 3.05 KOH, pH 7.20.

Intracellular Ca^{2+} imaging

Fluorescent Ca^{2+} signals were measured with a C2 confocal system and NIS-element imaging

software (Nikon) coupled to the FN1 upright microscope as previously described (Vincent et

al., 2014 ; 2017). For Ca^{2+} imaging in IHCs, in which membrane capacitance and Ca^{2+}

currents were simultaneously recorded, patch pipettes were filled with the same cesium-base

solution described above supplemented with 50 µM of the red fluorescent indicator Rhod2

(R14220, Invitrogen[TM], Thermofisher Scientific). The Ca^{2+} dye was excited with a 543 nm

Helium Neon Laser system (Melles Griot 05-LGR-193-381) coupled to a Nikon C2-confocal

FN1- upright microscope and emission fluorescence at 552-617 nm was recorded at 18

images/s (resolution 0.4 µm/pixel). Using the Nikon imaging system NIS-Elements, IHC

active zones were first individually identified by their fluorescent transient responses,

following a 100 ms stimulation (from -80 to -10 mV) and then the focal plane (with pinhole

1.1 µm) was adjusted in the z dimension to give the brightest value for each active zone.

Then, after a 1 min rest, each active zone was stimulated by a train of five consecutive 100 ms

stimulation, each separated by 250 ms. Fluorescence emission in each active zone was

subsequently measured with ImageJ software by drawing a region of interest of 1 μm^2 at the

synaptic zone (Fig.9B). Emission fluorescent signals were analyzed and normalized by the

ratio $\Delta F/F_0$ ratio, where F_0 was the fluorescence level before stimulation.

Statistical analysis

Results were analyzed with OriginPro 9.1software (OriginLab, Northampton, USA). Results

were analyzed with OriginPro 9.1software (OriginLab, Northampton, USA). Data were tested

for normal distribution using the Shapiro-Wilk normality test, and parametric or

nonparametric tests were applied accordingly. For statistical analyses with two data sets, two-

tailed unpaired t-tests or two-sided Mann-Whitney tests were used. For comparisons of more than two data sets, one-way ANOVA or two-way ANOVA followed by Tukey multiple comparison test were used when data was normally distributed. If a data set was not normally distributed Kruskal-Wallis tests followed by Dunn multiple comparison tests, respectively, was used. When more than two data sets were compared and significant differences were found, reported p values correspond to the post-hoc multiple comparison test. Values of $p < 0.05$ were considered significant. Results are expressed as mean ± SEM.

REFERENCES

Bartolomé MV, del CE, López LM, Carricondo F, Poch-Broto J, Gil-Loyzaga P. 2002. Effects of aging on C57BL/6J mice: an electrophysiological and morphological study. *Advances in oto-rhino-laryngology* **59**:106-11. doi: 10.1159/000059247

Becker L, Schnee ME, Niwa M, Sun W, Maxeiner S, Talaei S, Kachar B, Rutherford MA, Ricci AJ. 2018. The presynaptic ribbon maintains vesicle populations at the hair cell afferent fiber synapse. *eLife* **7**:e30241. doi: 10.7554/eLife.30241.

Beurg M, Michalski N, Safieddine S, Bouleau Y, Schneggenburger R, Chapman ER, Petit C, Dulon D. 2010. Control of exocytosis by synaptotagmins and otoferlin in auditory hair cells. *The Journal of neuroscience* **30**(40):13281-90.

Brandt A, Striessnig J, Moser T. 2003. Ca$_V$1.3 channels are essential for development and presynaptic activity of cochlear inner hair cells. *The Journal of neuroscience* **23**(34): 10832-10840.

Campbell LW, Hao SY, Thibault O, Blalock EM, and. Landfield PW. 1996. Aging Changes in Voltage-Gated Calcium Currents in Hippocampal CA1 Neurons. *The Journal of neuroscience* **16**(19):6286–6295.

Defourny J, Aghaie A, Perfettini I, Avan P, Delmaghani S, Petit C. 2019. Pejvakin-mediated pexophagy protects auditory hair cells against noise-induced damage. *PNAS* **116**(16):8010-8017.

Delmaghani S, Defourny J, Aghaie A, Beurg M, Dulon D, Thelen N, Perfettini I, Zelles T, Aller M, Meyer A, Emptoz A, Giraudet F, Leibovici M, Dartevelle S, Soubigou G, Thiry M, Vizi ES, Safieddine S, Hardelin JP, Avan P, Petit C. 2015. Hypervulnerability to Sound Exposure through Impaired Adaptive Proliferation of Peroxisomes. *Cell* **163**(4):894-906. doi: 10.1016/j.cell.2015.10.023

Del Nido PJ, Glynn P, Buenaventura P, Salama G, Koretsky AP. 1998. Fluorescence measurement of calcium transients in perfused rabbit heart using rhod 2. *The American journal of physiology* **274**(2):H728-41.

Fernandez KA, Jeffers PW, Lall K, Liberman MC, Kujawa SG**.** 2015. Aging after noise exposure: acceleration of cochlear synaptopathy in "recovered" ears. *The Journal of neuroscience* **35**(19):7509-20. doi: 10.1523/JNEUROSCI.5138-14.2015.

Fingar DC, Salama S, Tsou C, Harlow E, Blenis J. 2002. Mammalian cell size is controlled by mTOR and its downstream targets S6K1 and 4EBP1/eIF4E. *Genes & development* **16**(12):1472-87.

Frank T, Khimich D, Neef A, Moser T. 2009. Mechanisms contributing to synaptic Ca2+ signals and their heterogeneity in hair cells. *PNAS* **106**(11):4483-8.

Fritzsch B, Duncan JS, Kersigo J, Gray B, Elliott KL. 2016. Neuroanatomical Tracing Techniques in the Ear: History, State of the Art, and Future Developments. *Methods in molecular biology* **1427**:243-62. doi: 10.1007/978-1-4939-3615-1_14.

Fu X, Sun X, Zhang L, Jin Y, Chai R, Yang L, Zhang A, Liu X, Bai X, Li J, Wang H, Gao J. 2018. Tuberous sclerosis complex-mediated mTORC1 overactivation promotes age-related hearing loss. *Journal of Clinical Investigation* **128**(11):4938-4955.

Fumarola C, La Monica S, Alfieri RR, Borra E, Guidotti GG. 2005. Cell size reduction induced by inhibition of the mTOR/S6K-signaling pathway protects Jurkat cells from apoptosis. *Cell death and differentiation* **12**(10):1344-57.

Grant L and Fuchs P. 2008. Calcium- and calmodulin-dependent inactivation of calcium channels in inner hair cells of the rat cochlea. *Journal of neurophysiology* **99**(5):2183-93. doi: 10.1152/jn.01174.2007.

Graydon CW, Cho S, Li GL, Kachar B, von Gersdorff H. 2011. Sharp Ca^{2+} nanodomains beneath the ribbon promote highly synchronous multivesicular release at hair cell synapses. *The Journal of neuroscience* **31**(46):16637-50.

Hafidi A, Beurg M, Dulon D. 2005. Localization and developmental expression of BK channels in mammalian cochlear hair cells. *Neuroscience* **130**(2):475-84

Hasson T, Heintzelman MB, Santos-Sacchi J, Corey DP, and Mooseker MS. 1995. Expression in cochlea and retina of myosin VIIa, the gene product defective in Usher syndrome type 1B. *PNAS* **92** (21) 9815-9819

Hequembourg S and Liberman MC. 2001. Spiral ligament pathology: a major aspect of age-related cochlear degeneration in C57BL/6 mice. *JARO* **2**(2):118-29.

Henry KR. 2002. Sex- and age-related elevation of cochlear nerve envelope response (CNER) and auditory brainstem response (ABR) thresholds in C57BL/6 mice. *Hearing research* **170**(1-2):107-15.

Hermann A, Sitdikova GF, Weiger TM. 2015. Oxidative Stress and Maxi Calcium-Activated Potassium (BK) Channels. *Biomolecules* **5**(3):1870-911.

Hu N, Rutherford MA, Green SH. 2020. Protection of cochlear synapses from noise-induced
excitotoxic trauma by blockade of Ca2+-permeable AMPA receptors. *PNAS* **117**(7):3828-
3838.

Hickman TT, Hashimoto K, Liberman LD, Liberman MC. 2020. Synaptic migration and
reorganization after noise exposure suggests regeneration in a mature mammalian cochlea.
Scientific reports **10**(1):19945.

Ison JR and Allen PD. 2003. Low-frequency tone pips elicit exaggerated startle reflexes in
C57BL/6J mice with hearing loss. *JARO* **4**(4):495-504.

Ison JR, Allen PD, O'Neill WE. 2007 Age-related hearing loss in C57BL/6J mice has both
frequency-specific and non-frequency-specific components that produce a hyperacusis-like
exaggeration of the acoustic startle reflex. *JARO* **8**(4):539-50.

Jean P, Lopez de la Morena D, Michanski S, Jaime Tobón LM, Chakrabarti R, Picher MM,
Neef J, Jung S, Gültas M, Maxeiner S, Neef A, Wichmann C, Strenzke N, Grabner C, Moser
T. 2018. The synaptic ribbon is critical for sound encoding at high rates and with temporal
precision. *eLife* **7**:e29275.
824
825 **Jeng J**, Johnson SL, Carlton AJ, De Tomasi L, Goodyear RJ, De Faveri F, Furness DN, Wells
826 S, Brown SDM, Holley MC, Richardson GP, Mustapha M, Bowl MR, Marcotti W. 2020.
827 Age-related changes in the biophysical and morphological characteristics of mouse cochlear
828 outer hair cells. *The Journal of physiology* **598**(18):3891-3910.
829
830 **Jeng J**, Carlton AJ, Johnson SL, Brown SDM, Holley MC, Bowl MR, Marcotti W. 2021.
831 Biophysical and morphological changes in inner hair cells and their efferent innervation in the
832 ageing mouse cochlea. *The Journal of physiology* **599**(1):269-287.
833
Jiang XW, Li XR, Zhang YP. 2015. Changes of ribbon synapses number of cochlear hair
cells in C57BL/6Jmice with age. *International journal of clinical and experimental medicine*
8(10):19058-64. eCollection 2015.

Jing Z, Rutherford MA, Takago H, Frank T, Fejtova A, Khimich D, Moser T, Strenzke N.
2013. Disruption of the presynaptic cytomatrix protein bassoon degrades ribbon anchorage,
multiquantal release, and sound encoding at the hair cell afferent synapse. *The Journal of*
neuroscience **33**(10):4456-67. doi: 10.1523/JNEUROSCI.3491-12.2013

Johnson KR, Erway LC, Cook SA, Willott JF, Zheng QY. 1997. A major gene affecting age-
related hearing loss in C57BL/6J mice. *Hearing research* **114**(1-2):83-92

Johnson SL, Marcotti W, Kros CJ. 2005. Increase in efficiency and reduction in Ca2+
dependence of exocytosis during development of mouse inner hair cells. *The Journal of*
Physiology **563**(Pt 1):177-91.

Kazmierczak P, Sakaguchi H, Tokita J, Wilson-Kubalek EM, Milligan RA, Müller U,
Kachar B. 2007. Cadherin 23 and protocadherin 15 interact to form tip-link filaments in
sensory hair cells. *Nature* **449**(7158):87-91.

Kim KX, Payne S, Yang-Hood A, Li SZ, Davis B, Carlquist J, V-Ghaffari B, Gantz JA, Kallogjeri D, Fitzpatrick JAJ, Ohlemiller KK, Hirose K, Rutherford MA. 2019. Vesicular Glutamatergic Transmission in Noise-induced Loss and Repair of Cochlear Ribbon Synapses. *The Journal of neuroscience* **39** (23) 4434-4447; doi: 10.1523/JNEUROSCI.2228-18.2019.

Knipper M, Van Dijk P, Nunes I, Rüttiger L, Zimmermann U. 2013. Advances in the neurobiology of hearing disorders: recent developments regarding the basis of tinnitus and hyperacusis. *Progress in neurobiology* **111**:17-33. doi: 10.1016/j.pneurobio.2013.08.002.

Kros CJ, Ruppersberg JP, Rüsch A. 1998. Expression of a potassium current in inner hair cells during development of hearing in mice. *Nature* **394**(6690):281-4

Kujawa SG and Liberman MC. 2015. Synaptopathy in the noise-exposed and aging cochlea: Primary neural degeneration in acquired sensorineural hearing loss. *Hearing research* **330**(Pt B):191-9.

Kujawa SG and Liberman MC. 2019. Translating animal models to human therapeutics in noise-induced and age-related hearing loss. *Hearing research* **77**:44-52.

Kurt S, Sausbier M, Rüttiger L, Brandt N, Moeller CK, Kindler J, Sausbier U, Zimmermann U, van Straaten H, Neuhuber W, Engel J, Knipper M, Ruth P, Schulze H. 2012. Critical role for cochlear hair cell BK channels for coding the temporal structure and dynamic range of auditory information for central auditory processing. *FASEB journal* **26**(9):3834-43.

Lagziel A, Ahmed ZM, Schultz JM, Morell RJ, Belyantseva IA, Friedman TB. 2005. Spatiotemporal pattern and isoforms of cadherin 23 in wild type and waltzer mice during inner ear hair cell development. *Developmental Biology* **280**(2):295-306.

Lang I, Jung M, Niemeyer BA, Ruth P, Engel J. 2019. Expression of the LRRC52 γ subunit (γ2) may provide Ca 2+-independent activation of BK currents in mouse inner hair cells. *FASEB journal* **33**(11):11721-11734.

Liberman LD, Wang H, Liberman MC. 2011. Opposing gradients of ribbon size and AMPA receptor expression underlie sensitivity differences among cochlear-nerve/hair-cell synapses. *The Journal of neuroscience* **19**;31(3):801-8.

Liberman MC and Kujawa SG. 2017. Cochlear synaptopathy in acquired sensorineural hearing loss: Manifestations and mechanisms. *Hearing research* **349**:138-147.

Liberman MC. 1982. Single-neuron labeling in the cat auditory nerve. *Science* **216**(4551):1239-41.

Lingle CJ, Martinez-Espinosa PL, Yang-Hood A, Boero LE, Payne S, Persic D, V-Ghaffari B, Xiao M, Zhou Y, Xia X, Pyott SJ, Rutherford MA. 2019. LRRC52 regulates BK channel function and localization in mouse cochlear inner hair cells. *PNAS* **116**(37):18397-18403.

Magupalli VG, Schwarz K, Alpadi K, Natarajan S, Seigel GM, Schmitz F. 2008. Multiple RIBEYE-RIBEYE interactions create a dynamic scaffold for the formation of synaptic ribbons. *The Journal of neuroscience* **28**:7954– 7967.

Matsubara A, Laake JH, Davanger S, Usami S, Ottersen OP. 1996. Organization of AMPA receptor subunits at a glutamate synapse: a quantitative immunogold analysis of hair cell synapses in the rat organ of Corti. *The Journal of neuroscience* **16**(14):4457-67.

Meyer AC, Frank T, Khimich D, Hoch G, Riedel D, Chapochnikov NM, Yarin YM, Harke B, Hell SW, Egner A, Moser T. 2009. Tuning of synapse number, structure and function in the cochlea. *Nature Neuroscience* **12**(4):444-53. doi: 10.1038/nn.2293. Epub 2009 Mar 8.

Michalski N, Goutman JD, Auclair SM, Boutet de Monvel J, Tertrais M, Emptoz A, Parrin A, Nouaille S, Guillon M, Sachse M, Ciric D, Bahloul A, Hardelin JP, Sutton RB, Avan P, Krishnakumar SS, Rothman JE, Dulon D, Safieddine S, Petit C. 2017. Otoferlin acts as a Ca^{2+} sensor for vesicle fusion and vesicle pool replenishment at auditory hair cell ribbon synapses. *eLife* **7**;6. pii: e31013.

Michel V, Goodyear RJ, Weil D, Marcotti W, Perfettini I, Wolfrum U, Kros CJ, Richardson GP, Petit C. 2005. Cadherin 23 is a component of the transient lateral links in the developing hair bundles of cochlear sensory cells. *Developmental Biology* **280**(2):281-94

Neef J, Urban NT, Ohn TL, Frank T, Jean P, Hell SW, Willig KI, Moser T. 2018. Quantitative optical nanophysiology of Ca_{2+} signaling at inner hair cell active zones. *Nature Communication* **9**(1):290. doi: 10.1038/s41467-017-02612-y.

Noben-Trauth K, Zheng QY, Johnson KR. 2003. Association of cadherin 23 with polygenic inheritance and genetic modification of sensorineural hearing loss. *Nature Genetics* **35**(1):21-3. Epub 2003 Aug 10.

Oliver D, Taberner AM, Thurm H, Sausbier M, Arntz C, Ruth P, Fakler B, Liberman MC. 2006. The role of BKCa channels in electrical signal encoding in the mammalian auditory periphery. *The Journal of neuroscience* **26**(23):6181-9.

Ouagazzal AM, Reiss D, Romand R. 2006. Effects of age-related hearing loss on startle reflex and prepulse inhibition in mice on pure and mixed C57BL and 129 genetic background. *Behavior Brain Research*. **172**(2):307-15. doi: 10.1016/j.bbr.2006.05.018.

Özçete ÖD, Moser T. 2021. A sensory cell diversifies its output by varying Ca 2+ influx-release coupling among active zones. *EMBO Journal* **1**;40(5):e106010.

Pujol R and Puel JL. 1999. Excitotoxicity, synaptic repair, and functional recovery in the mammalian cochlea: a review of recent findings. *Annals of the New York Academy of Sciences* **884**:249-54.

Pyott SJ, Glowatzki E, Trimmer JS, Aldrich RW. 2004. Extrasynaptic localization of inactivating calcium-activated potassium channels in mouse inner hair cells. *The Journal of neuroscience* **24**(43):9469-74.

Rousset F, Nacher-Soler G, Coelho M, Ilmjarv S, Kokje VBC, Marteyn A, Cambet Y, Perny
M, Roccio M, Jaquet V, Senn P, Krause KH. 2020. Redox activation of excitatory pathways
in auditory neurons as mechanism of age-related hearing loss. Redox Biology **30**:101434.

Roux I, Safieddine S, Nouvian R, Grati M, Simmler MC, Bahloul A, Perfettini I, Le Gall M,
Rostaing P, Hamard G, Triller A, Avan P, Moser T, Petit C. 2006. Otoferlin, defective in a
human deafness form, is essential for exocytosis at the auditory ribbon synapse. *Cell*
127(2):277-89.

Sahoo N, Hoshi T, Heinemann SH. 2014. Oxidative modulation of voltage-gated potassium
channels. *Antioxid Redox Signal* **21**(6):933-52. doi: 10.1089/ars.2013.5614. Epub 2013 Oct
26. Review

Sheets L, He XJ, Olt J, Schreck M, Petralia RS, Wang YX, Zhang Q, Beirl A, Nicolson T,
Marcotti W, Trapani JG, Kindt KS. 2017. Enlargement of Ribbons in Zebrafish Hair Cells
Increases Calcium Currents But Disrupts Afferent Spontaneous Activity and Timing of
Stimulus Onset. *The Journal of neuroscience* **37**(26):6299-6313. doi:
10.1523/JNEUROSCI.2878-16.2017.
949
950 **Siemens J**, Lillo C, Dumont RA, Reynolds A, Williams DS, Gillespie PG, Müller U. 2004.
951 Cadherin 23 is a component of the tip link in hair-cell stereocilia. *Nature* **428**(6986):950-5.
952 Epub 2004 Mar 31.
953
954 **Skinner LJ**, Enée V, Beurg M, Jung HH, Ryan AF, Hafidi A, Aran JM, Dulon D. 2003.
955 Contribution of BK Ca2+-activated K+ channels to auditory neurotransmission in the Guinea
956 pig cochlea. *Journal of Neurophysiology* **90**(1):320-32.
957
958 **Söllner C**, Rauch GJ, Siemens J, Geisler R, Schuster SC, Müller U, Nicolson T; Tübingen
959 2000 Screen Consortium. 2004. Mutations in cadherin 23 affect tip links in zebrafish sensory
960 hair cells. *Nature* **428**(6986):955-9.
961
962 **Spassova MA**, Soboloff J, He L, Hewavitharana T, Xu W, Venkatachalam K, van Rossum
963 DB, Patterson RL, Gill DL. 2004. Calcium entry mediated by SOCs and TRP channels:
964 variations and enigma. Review *Biochimica et biophysica acta* **1742**(1-3):9-20.
965
Spongr VP, Flood DG, Frisina RD, Salvi RJ. 1997. Quantitative measures of hair cell loss in
CBA and C57BL/6 mice throughout their life spans. *The Journal of the Acoustical Society of*
America **101**(6):3546-53.

Stamataki S, Francis HW, Lehar M, May BJ, Ryugo DK. 2006. Synaptic alterations at inner
hair cells precede spiral ganglion cell loss in aging C57BL/6J mice. *Hearing research* **221**(1-
2):104-18. Epub 2006 Sep 26
972
973 **Taberner AM and Liberman MC**. 2005. Response properties of single auditory nerve fibers
974 in the mouse. *Journal of neurophysiology* **93**(1):557-69.

975 **Thibault O and. Landfield PW**. 1996. Increase in Single L-Type Calcium Channels in
976 Hippocampal Neurons During Aging. *Science* **272** : 1017-1020.

Vincent PF, Bouleau Y, Safieddine S, Petit C, Dulon D. 2014. Exocytotic machineries of vestibular type I and cochlear ribbon synapses display similar intrinsic otoferlin-dependent Ca2+ sensitivity but a different coupling to Ca2+ channels. *The Journal of neuroscience* **34**(33):10853-69.

Vincent PF, Bouleau Y, Petit C, Dulon D. 2015. A synaptic F-actin network controls otoferlin-dependent exocytosis in auditory inner hair cells. *eLife* **4**.

Vincent PF, Bouleau Y, Charpentier G, Emptoz A, Safieddine S, Petit C, Dulon D. 2017. Different Ca$_V$1.3 Channel Isoforms Control Distinct Components of the Synaptic Vesicle Cycle in Auditory Inner Hair Cells. *The Journal of neuroscience* **37**(11):2960-2975.

Walton JP, Frisina RD, Meierhans LR. 1995. Sensorineural hearing loss alters recovery from short-term adaptation in the C57BL/6 mouse. *Hearing research* **88**(1-2):19-26.

Wan G and Corfas G. 2015. No longer falling on deaf ears: mechanisms of degeneration and regeneration of cochlear ribbon synapses. *Hearing research* **329**:1-10.

Wang Y, Tao J, Wang M, Yang L, Ning F, Xin H, Xu X, Cai H, Zhang W, Yu K, Zhang X. 2019. Mechanism of Regulation of Big-Conductance Ca^{2+}-Activated K$^+$ Channels by mTOR Complex 2 in Podocytes. *Frontiers in physiology* **10**:167. doi: 10.3389/fphys.2019.00167. eCollection 2019.

Willott JF. 1996. Anatomic and physiologic aging: a behavioral neuroscience perspective. *Journal of the American Academy of Audiology* **7**(3):141-51.

Wong HC, Zhang Q, Beirl AJ, Petralia RS, Wang YX, Kindt K. 2019. Synaptic mitochondria regulate hair-cell synapse size and function. *eLife* **8**. pii: e48914. doi: 10.7554/eLife.48914.

Zachary SP and Fuchs PA. 2015. Re-Emergent Inhibition of Cochlear Inner Hair Cells in a Mouse Model of Hearing Loss. *The Journal of neuroscience* **35**(26):9701-6.

Zheng QY, Johnson KR, and Erway LC. 1999. Assessment of hearing in 80 inbred strains of mice by ABR threshold analyses. *Hearing research* **130**(1-2): 94–107.

Zhou X, Jen PH-S, Seburn KL, Frankel WN, and Zheng QY. 2006. Auditory brainstem responses in 10 inbred strains of mice. *Brain Research* **1091,** 16–26.