# Dopamine influences attentional rate modulation in Macaque posterior parietal cortex

Jochem van Kempen[1*], Christian Brandt[2], Claudia Distler[3], Mark A. Bellgrove[4], Alexander Thiele[1]

[1] Biosciences Institute, Newcastle University, NE1 7RU, Newcastle upon Tyne, United Kingdom.

[2] Institute of Clinical Research, University of Southern Denmark, 5230 Odense M, Denmark.

[3] Allgemeine Zoologie und Neurobiologie, Ruhr-Universität Bochum, 44801 Germany

[4] Turner Institute for Brain and Mental Health, School of Psychological Sciences, Monash University, Melbourne, Victoria 3800, Australia.

* Jochem.van-kempen@newcastle.ac.uk

## Conflict of interest statement

There are no conflicts of interest.

## Acknowledgements

Funded by Wellcome trust 093104 (JvK, AT), MRC MR/P013031/1 (JvK, AT). MAB is supported by a Senior Research Fellowship from the Australian National Health and Medical Research Council (NHMRC). Funded by strategic research partnership between Newcastle University and Monash University.

## Abstract

Selective attention facilitates the prioritization of task-relevant sensory inputs over those which are irrelevant. Although cognitive neuroscience has made great strides in understanding the neural substrates of attention, our understanding of its neuropharmacology is incomplete. Cholinergic and glutamatergic contributions have been demonstrated, but emerging evidence also suggests an important influence of dopamine (DA). DA has historically been investigated in the context of frontal/prefrontal function arguing that dopaminergic receptor density in the posterior/parietal cortex is sparse. However, this notion was derived from rodent data, whereas in primates DA innervation in parietal cortex matches that of many prefrontal areas. We recorded single- and multi-unit activity whilst iontophoretically administering dopaminergic agonists and antagonists to posterior parietal cortex of rhesus macaques engaged in a spatial attention task. Out of 88 neurons, 50 showed modulation of activity induced by drug administration. Dopamine inhibited firing rates across the population according to an inverted-U shaped dose-response curve. D1 receptor antagonists diminished firing rates in broad-spiking units according to a monotonically increasing function. Additionally, dopamine modulated attentional signals in broad, but not narrow-spiking cells. Finally, both drugs modulated the pupil light reflex. These data show that dopamine plays an important role in shaping neuronal responses and modulates attentional processing in macaque parietal cortex.

## Significance statement

Dopamine is critically involved in high-level cognitive functions, and dopaminergic dysfunctions pertain to ageing and neurological and psychiatric disorders. Most previous studies focused on dopaminergic effects on prefrontal activity or its role in basal ganglia

45  circuitry. The effects of dopamine in other brain areas such as parietal cortex, despite its well-

46  established role in cognition and cognitive dysfunction, have largely been overlooked. This

47  study is the first to show dopaminergic modulation of parietal activity in general, and specific

48  to spatial attention in the non-human primate, revealing cell-type specific effects of dopamine

49  on attentional modulation.

## Introduction

Selective attention refers to prioritization of behaviorally relevant, over irrelevant, sensory

inputs. Convergent evidence from human neuropsychological, brain imaging and non-human

primate studies shows that fronto-parietal networks are crucial for selective attention (Posner,

1990; Desimone and Duncan, 1995; Corbetta and Shulman, 2011). Neuromodulation of

attention-related activity in these networks occurs at least in part via glutamatergic (Herrero

et al., 2013) and cholinergic inputs (Warburton and Rusted, 1993; Levin and Simon, 1998;

Nelson et al., 2005; Sarter et al., 2005; Parikh et al., 2007; Furey et al., 2008; Herrero et al.,

2008; Dasilva et al., 2019). Multiple lines of evidence, however, also suggest dopaminergic

modulation (Bellgrove and Mattingley, 2008; Noudoost and Moore, 2011a; Soltani et al.,

2013; Thiele and Bellgrove, 2018). Here we sought to understand how dopamine applied to

macaque posterior parietal cortex (PPC) modulates attention-related activity.

The functional significance of dopamine is well-established for a number of brain areas,

including the frontal cortex (executive control) and basal ganglia (motor control). For these

regions, substantial across-species similarities allowed the development of mechanistic

models with clinical translational value for various disorders (e.g., Parkinson's disease,

schizophrenia or attention deficit hyperactivity disorder (ADHD)) (Arnsten et al., 2012;

Thiele and Bellgrove, 2018). Species differences with respect to dopaminergic innervation do

however exist for posterior cortical areas, including the PPC. Although sparse in rodents,

dopaminergic innervation of parietal areas in non-human primates is comparable in strength

and laminar distribution to prefrontal regions (Berger et al., 1991). Moreover, macaque PPC

has high densities of dopamine transporter (DAT) immunoreactive axons (Lewis et al., 2001).

These observations align with dense dopaminergic receptor expression in human PPC

(Caspers et al., 2013) and imaging studies of clinical disorders where medications targeting

dopamine receptors or transporters modulate parietal activity (Mehta et al., 2000). Given

these data and the clinical significance of PPC function, greater understanding of

dopaminergic effects in this region is warranted.

Selective attention relies heavily on PPC integrity and multiple lines of evidence suggest that

dopamine modulates attentional processes related to parietal function. First, dopamine

agonists reduce spatial inattention in neurological (Gorgoraptis et al., 2012) and psychiatric

patients with disorders such as schizophrenia (Maruff et al., 1995) and ADHD (Bellgrove et

al., 2008; Silk et al., 2014). Second, psychopharmacological studies in healthy volunteers

suggest that dopamine antagonists modulate parameters of spatial cueing paradigms (e.g.

validity effect), often associated with parietal function (Clark et al., 1989). Third, DNA

variation in a polymorphism of the dopamine transporter gene (DAT1) is associated with

individual differences in measures of spatial selective attention (Bellgrove et al., 2007, 2009;

Newman et al., 2014). Fourth, non-human primate studies revealed dopaminergic

contributions to working memory signals in dorsolateral prefrontal cortex (dlPFC) (Williams

and Goldman-Rakic, 1995), and modulation of dopaminergic signaling in frontal eye fields

(FEF) affects V4 neurons in a manner similar to attention and biases behavioral choices

(Noudoost and Moore, 2011a; Soltani et al., 2013). Dopamine thus contributes to working

memory, target selection and probably also spatial attention in dlPFC and FEF (Williams and

Goldman-Rakic, 1995; Noudoost and Moore, 2011a, 2011b; Clark and Noudoost, 2014).

Both areas are critical nodes of fronto-parietal attention networks. Thus, while dopaminergic

influences on frontal circuits are comparatively well understood, their effect on attention-

related activity in PPC is yet to be established.

Here we sought to address this knowledge gap by locally infusing dopamine or the selective

D1 receptor (D1R) antagonist SCH23390 into the PPC of two macaque monkeys during a

selective attention task. We show that single and multi-unit (SU, MU) activity is inhibited by

iontophoresis of dopaminergic drugs into intraparietal sulcus (IPS) gray matter. The effects of

the non-selective agonist dopamine (DA) followed an inverted U-shaped dose-response

curve, whereas the D1-selective antagonist SCH23390 dose-response followed a monotonic

function. Additionally, we found cell-type specific effects on attentional modulation whereby

DA affected attention-related activity in broad but not narrow-spiking units. Finally, both

drugs reduced the pupillary light reflex.

## Materials & Methods

### Procedures

All animal procedures were performed in accordance with the European Communities

Council Directive RL 2010/63/EC, the National Institute of Health's Guidelines for the Care

and Use of Animals for Experimental Procedures, and the UK Animals Scientific Procedures

Act. Animals were motivated to engage in the task through fluid control at levels that do not

affect animal physiology and have minimal impact on psychological wellbeing (Gray et al.,

2016).

### Surgical preparation

The monkeys were implanted with a head post and recording chambers over the lateral intraparietal sulcus under sterile conditions and under general anesthesia. Surgery and postoperative care conditions have been described in detail previously (Thiele et al., 2006).

### Behavioral paradigms

Stimulus presentation and behavioral control was regulated by Remote Cortex 5.95 (Laboratory of Neuropsychology, National Institute for Mental Health, Bethesda, MD). Stimuli were presented on a cathode ray tube (CRT) monitor at 120 Hz, 1280 × 1024 pixels, at a distance of 54 cm.

The location of the saccade field (SF) was mapped using a visually guided saccade task. Here, monkeys fixated centrally for 400 ms after which a saccade target was presented in one of nine possible locations (8-10° from fixation, equally spaced between). After a random delay (800-1400 ms, uniformly distributed) the fixation point was extinguished, which indicated to the monkey to perform a saccade towards the target. Online analysis of visual, sustained and saccade related activity determined an approximate SF location which guided our subsequent receptive field (RF) mapping. The location and size of RFs were measured as described previously (Gieselmann and Thiele, 2008), using a reverse correlation method. Briefly, during fixation, a series of black squares (1-3° size, 100% contrast) were presented for 100 ms at pseudorandom locations on a 9 × 12 grid (5-25 repetitions for each location) on a bright background. RF eccentricity ranged from 2.5° to 17° and were largely confined to the contralateral visual field.

The main task and stimuli have been described previously (Thiele et al., 2016; van Kempen et al., 2020). In brief, stimuli were presented on a cathode ray tube (CRT) monitor (120 Hz,

1280 × 1024 pixels, 55 cm from the animal). The monkey initiated a trial by holding a lever

and fixating a white fixation spot (0.1°) displayed on a grey background (1.41 cd/m$^2$). After

425/674 ms [monkey 1/monkey 2] three colored square wave gratings (2° - 6°, dependent on

RF size and distance from fixation) appeared equidistant from the fixation spot, one of which

was centered on the RF of the recorded neuron. Red, green and blue gratings (see Table 1 for

color values) were presented with an orientation at a random angle to the vertical meridian

(the same orientation for the three gratings in any given session). The locations of the colors,

as well as the orientation, were pseudorandomly assigned between recording sessions and

held constant for a given recording session. Gratings moved perpendicular to the orientation,

whereby the direction of motion was pseudorandomly assigned for every trial. After a

random delay (570-830/620-940 ms [monkey 1/monkey 2], uniformly distributed in 1 ms

steps) a central cue appeared that matched the color of the grating that would be relevant on

the current trial. After 980-1780/1160-1780 ms [monkey 1/monkey 2] (uniformly distributed

in 1 ms steps), one pseudorandomly selected grating changed luminance (dimmed). If the

cued grating dimmed, the monkey had to release the lever to obtain a reward. If a non-cued

grating dimmed, the monkey had to ignore this and wait for the cued grating to dim. This

could happen when the second or third grating changed luminance (each after 750-1130/800-

1130 ms [monkey 1/monkey 2], uniformly distributed in 1 ms steps). Drugs were

administered in blocks of 36 trials. The first block was always a control block. Thereafter,

drug blocks and recovery blocks were alternated until the animal stopped working (number of

block reversals, median ± interquartile range = 12 ± 6).

## Identification of recording sites

The location of the IPS was initially guided by means of postoperative structural magnetic resonance imaging (MRI), displaying the recording chamber. During each recording, neuronal response properties where determined using SF and RF mapping tasks. During the SF mapping task, we targeted cells that showed spatially selective persistent activity and preparatory activity before the execution of a saccadic eye movement.

## Electrode-pipette manufacturing

We recorded from the lateral (and in a few occasions medial) bank of the IPS using custom-made electrode-pipettes that allowed for simultaneous iontophoretic drug application and extracellular recording of spiking activity (Thiele et al., 2006). The location of the recording sites in one of the monkeys was verified in histological sections stained for cyto- and myeloarchitecture (Distler and Hoffmann, 2001) .

The manufacture of the electrodes was similar to the procedures described by Thiele et al., (2006), with minor changes to the design in order to reach areas deeper into the IPS, such as the ventral part of the lateral intraparietal area (LIPv). We sharpened tungsten wires (125 μm diameter, 75 mm length, Advent Research Materials Ltd., UK) by electrolytic etching off the tip (10-12 mm) in a solution of $NaNO_2$ (172.5 g), KOH (85 g) and distilled water (375 ml). We used borosilicate glass capillaries with three barrels (custom ordered, Hilgenberg GmBH, www.hilgenberg-gmbh.de), with the same dimensions as those described previously (Thiele et al., 2006). The sharpened tungsten wire was placed in the central capillary and secured in place by bending the non-sharpened end (approximately 10 mm) of the wire over the end of the barrel. After marking the location of the tip of the tungsten wire, shrink tubing was placed around the top and bottom of the glass. The glass was pulled around the tungsten wire using a

PE-21 Narishige microelectrode puller with a heating coil made from Kanthal wire (1 mm

diameter, 13 loops, inner loop diameter 3 mm) and the main (sub) magnet set to 30 (0) and

the heater at 100. The electrode-pipette was placed such that the tip of the tungsten wire

protruded 11 mm from the bottom of the heating coil. After pulling, we filled the central

barrel (with the tungsten electrode inside) with superglue using a syringe and fine flexible

injection cannula (MicroFil 28 AWG, MF28G67-5, World Precision Instruments, Ltd.). We

found that if we did not fill (most of) the central barrel with superglue after pulling, the

recorded signal was often very noisy, possibly due to small movements of the animal (such as

drinking), which caused the free tungsten wire to resonate inside the glass. Using a micro

grinder (Narishige EG-400), we removed excess glass, sharpened the tip of the electrode and

opened the flanking barrels of the pipette. This pulling procedure resulted in a pulled

electrode part of approximately 2.5 cm length, with gradually increasing diameter, from ~10

µm to ~200 µm, over the first 12 mm of the electrode-pipette.

### Electrode-pipette filling and iontophoresis

Electrode-pipettes were back-filled with the same drug in both pipettes using a syringe, filter

units (Millex® GV, 22 µm pore diameter, Millipore Corporation) and fine flexible injection

cannula (MicroFil 34 AWG, MF34G-5, World Precision Instruments, Ltd.). The pipettes

were connected to the iontophoresis unit (Neurophore-BH- 2, Medical systems USA) with

tungsten wires (125 µm diameter) inserted into the flanking barrels. Because of the

exploratory nature of these recordings (it is unknown whether DA influences parietal neurons

during spatial attention tasks and what modulation can be expected with different amounts of

drug applied), we used a variety of iontophoretic ejection currents (20 - 90 nA). The details

regarding concentration and pH of the drugs were: Dopamine (0.1M in water for injections,

pH 4-5) and SCH23390 (0.005-0.1M in water for injections, pH 4-5).

Data acquisition

Stimulus presentation, behavioral control and drug administration was regulated by Remote

Cortex 5.95 (Laboratory of Neuropsychology, National Institute for Mental Health, Bethesda,

MD). Raw data were collected using Remote Cortex 5.95 (1-kHz sampling rate) and by

Cheetah data acquisition (32.7-kHz sampling rate, 24-bit sampling resolution) interlinked

with Remote Cortex 5.95. Data were replayed offline, sampled with 16-bit resolution and

band-pass filtered (0.6-9 kHz). Spikes were sorted manually using SpikeSort3D (Neuralynx).

Eye position and pupil diameter were recorded using a ViewPoint eyetracker (Arrington

research) at 220 Hz. Pupil diameter was recorded in 40 out of 54 recording sessions.

Pupillometry

Pupil diameter was low pass filtered (10 Hz) using a second order Butterworth filter. Baseline

activity was estimated as the average activity before stimulus onset (-300 to -50 ms), which

was used to normalize the pupil diameter time course. Stimulus evoked pupil constriction

was baseline corrected (on a trial by trial basis) and averaged in a 250 ms time window

centered on 500 ms after stimulus onset. Cue-evoked (250 ms window centered on 500 ms

after cue onset) and pre-dimming (-300 to -50 ms) pupil diameter, was baseline subtracted

with a pre-cue baseline (-300 to -50 ms). For visualization, pupil diameter in each epoch was

scaled to a range from zero to one, before averaging across trials.

### Analysis of cell type.

We distinguished between different cell types based on the duration of the extracellular spike waveform as described in Thiele et al. (2016). Specifically, we classified cells based on the peak-to-trough ratio, i.e. the duration between the peak and the trough of the interpolated (cubic spline) spike waveform. To test whether the distribution of peak-to-trough distance of the spike waveforms was unimodal (null hypothesis) or bimodal, indicating that our distribution contained different cell types, a modified Hartigan's dip test was used (Ardid et al., 2015; Thiele et al., 2016). We used a cut-off of 250 µs to classify cells as narrow or broad spiking, as this was where our distribution revealed the main 'dip' (Figure 3A-B).

### Fano factor

The variability of neural responses was quantified using Fano factors ($FF$), computed as the ratio between the variance ($\sigma^2$) and the mean ($\mu$) spike counts within the time window of interest, defined as:

$$FF = \frac{\sigma^2}{\mu}$$

### Drug modulation

The strength of the effect of drug application on neural activity (firing rates) was determined via a drug modulation index ($drugMI$), defined as:

$$drugMI = \frac{drug_{on} - drug_{off}}{drug_{on} + drug_{off}}$$

with $drug_{on}$ as the neural activity when drug was applied, and $drug_{off}$ the activity when the drug was not applied. This index ranges from -1 to 1, with zero indicating no modulation due

to drug application and with positive values indicating higher activity when the drug was

applied and conversely, negative values indicating lower activity.

Quantification of attentional rate modulation.

To quantify the difference between neural responses when attention was directed towards the

RF versus away from the RF, we computed the area under the receiver operating

characteristic (AUROC) curve. Stemming from signal detection theory (Green and Swets,

1966), this measure represents the difference between two distributions as a single scalar

value, taking into account both the average difference in magnitude as well as the variability

of each distribution. This value indicates how well an ideal observer would be able to

distinguish between two distributions, for example the neural response when attention is

directed towards versus away from its RF. It is computed by iteratively increasing the

threshold and computing the proportion (from the first sample to the threshold) of hits and

false alarms (FA), i.e. the correct and false classification as samples belonging to one of the

activity distributions. The ROC curve is generated by plotting the proportions of hits against

the proportion of FAs, and AUROC is taken as the area under the ROC curve. An AUROC of

0.5 indicates that the two distributions were indistinguishable, whereas an AUROC of 0 or 1

indicates that the two distributions were perfectly separable.

Experimental design and statistical analysis

We recorded single (SU, n=40) and multi-unit (MU, n=48) activity (total 88 units; 64 from

monkey 1, 24 from monkey 2) from 2 male rhesus macaque monkeys (*Macaca mulatta,* age

9-11 years, weight 8-12.9 kg).

To determine whether DA significantly affected neural activity across the population of units,

we used two-sided paired-sample Wilcoxon signed rank tests. For comparisons within one

recording, e.g. spike rates across trials for different conditions, we used analysis of variance

(ANOVA) with three factors: attention (towards or away from the RF), drug (on or off) and

stimulus direction. To test whether drug application affected behavioral performance, we

used sequential linear mixed effects models with attention and drug as fixed effects and with

the recording number as a random effect, to account for the repeated measurements in the

data.

To test for significant linear or quadratic trends in the drug dose-response curve, we used

sequential linear mixed effects models and likelihood ratio tests. Specifically, we tested

whether a first order (linear) polynomial fit was better than a constant (intercept-only) fit and

subsequently whether a second order (non-monotonic) polynomial fit was better than a linear

fit. The modulation due to drug application of the neural response $y$ was modelled as a linear

combination of polynomial basis functions of the iontophoretic ejection current ($X$):

$$y \sim \beta_0 + \beta_1 X + \beta_2 X^2$$

, with $\beta$ as the polynomial coefficients. When a significant quadratic relationship was found,

we used the two-lines approach to determine whether this relationship was significantly U-

shaped (Simonsohn, 2017). Error bars in all figures indicate the standard error of the mean

(SEM), unless stated otherwise.

We selected which cells to include in each of the analyses based on the output of the 3-factor

ANOVA described above. For example, if we wanted to investigate whether drug application

affected attentional modulation of firing rates, we only included cells that revealed a main or

interaction effect for both attention and drug application.

## Code

Data analyses were performed using custom written scripts in Matlab (the Mathworks) and RStudio (RStudio Team (2016). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA URL http://www.rstudio.com). Data and analysis scripts necessary to reproduce these results are available upon reasonable request.

## Results

We recorded activity from 88 single and multi-units from intraparietal sulcus (IPS) in two awake, behaving Macaque monkeys performing a selective attention task (Figure 1A). Of these cells, 86 (97.7%) showed a visual response to stimulus onset and 74 (84.1%) were modulated by attention (Figure 1B). During recording, we used an electrode-pipette combination to iontophoretically administer dopaminergic (DA) drugs in the vicinity of the recorded cells (Thiele et al., 2006). Across the two monkeys, we recorded from 59 units whilst administering the unselective agonist dopamine and from 29 units during which we administered the selective D1R antagonist SCH23390. Firing rates in 36 (61%) and 14 (48.3%) units were modulated by application of the unselective agonist dopamine and SCH23390, respectively (Figure 1B). Thus, around half the units were modulated both by attention and drug application (Figure 1B), which is comparable to cholinergic modulation of attention induced activity in macaque V1 and FEF (Herrero et al., 2008; Dasilva et al., 2019).

Figure 2A illustrates the population activity (from all units) aligned to stimulus onset, cue onset and the first-dimming event, for both the no-drug and the drug conditions. For a given drug condition, neural activity between attention conditions did not differ when aligned to stimulus onset but started to diverge approximately 200 ms after cue onset, indicating which of the three gratings was behaviorally relevant on that trial, and diverged further leading up to

the first dimming event. Across the population, dopamine strongly reduced firing rates

throughout the duration of the trial, including during baseline periods as well as stimulus and

cue presentation. The effects of SCH23390 were of the same sign but weaker. Although drug

induced changes to attentional modulation of neural activity appear relatively small at the

population level (using visual inspection, compare the difference between the dark and light

blue lines and the difference between the red and orange lines), a subset of neurons revealed

an interaction between attention and drug application (Figure 1B), as illustrated for an

example neuron in Figure 2B, and these effects depended on the cell types affected (see

below). Next, we specifically examined units that were modulated by attention and/or drug

application and investigated whether activity modulation due to attention and drug

application mapped onto different cell types.

Cells were classified as narrow or broad-spiking cells according to the median duration of the

peak-to-trough time of the spike waveforms (Figure 3A & B). These cell types have

previously been found to respond differently to dopaminergic drug application (Jacob et al.,

2013, 2016). Although narrow and broad-spiking cells have been argued to respectively

constitute inhibitory interneurons and excitatory pyramidal cells (Mitchell et al., 2007), a

more recent study found that output cells in primary motor cortex (unequivocal pyramidal

cells) had a narrow action potential waveform (Vigneswaran et al., 2011), and most

pyramidal cells in macaque PFC express the Kv3.1b potassium channel, associated with the

generation of narrow spikes (Soares et al., 2017). Therefore, the narrow-broad categorization

solely allows to distinguish between 2 different cell-type categories, without mapping this

classification specifically onto interneurons or pyramidal cells, let alone a more fine-grained

distinction.

The application of DA reduced firing rates across the population of both broad and narrow

spiking units, and for both the attend towards and away from RF conditions (Figure 3C).

Fano factors were unaffected by dopamine application (Figure 3D). The application of

SCH23390 elicited a small but significant reduction of the average firing rates of broad-

spiking units during both attention conditions (Figure 3E) without affecting FFs (Figure 3F).

Dopaminergic drug application thus mainly inhibited cellular activity, without affecting the

rate variability, as quantified by FFs.

To investigate whether dopamine affected attention-specific activity, we tested whether

attention AUROC values were modulated by drug application. Attention AUROC values

indicate how well an ideal observer can distinguish between neural activity during attend RF

or attend away trials. A value of 0.5 indicates that the distributions are indistinguishable,

whereas values of 0 or 1 indicate perfectly distinguishable distributions. The application of

the non-specific agonist dopamine reduced AUROC values for broad-spiking cells, whereas

narrow-spiking cells were unaffected (Figure 4A). SCH23390 application did not modulate

AUROC values for either cell type (Figure 4B). Dopamine thus had a cell-type specific effect

on attentional rate modulation.

We applied dopaminergic drugs with a variety of iontophoretic ejection currents (20-90 nA).

Since dopamine has previously been shown to modulate neural activity according to an

inverted U-shaped dose-response curve (Vijayraghavan et al., 2007), with maximal

modulation at intermediate dopamine levels, we tested whether the ejection current was

predictive of the firing rate modulation associated with drug application, estimated by a drug

modulation index (drugMI). Specifically, we used sequential linear mixed effects model

analyses and likelihood ratio tests to test for linear and quadratic trends. U-shaped trends

were verified using the two-lines approach (Materials & Methods). The non-specific agonist

dopamine displayed a non-monotonic relationship with drugMI ($\chi^2_{(1)} = 7.18$, $p = 0.007$) and

revealed an inverted U-shaped curve ($p < 0.05$) in which intermediate ejection currents

elicited the most negative drugMI, i.e. the largest inhibition of activity (Figure 4C). For

SCH23390, on the other hand, we found a monotonic dose-response relationship ($\chi^2_{(1)}$ = 4.21,

p = 0.040), with more inhibition of firing rates with higher drug ejection currents (Figure

4D). To investigate whether drug dosage was also predictive of attentional rate modulation,

we performed the same analysis on the difference score (drug – no drug) of attention

AUROC values. Neither dopamine ($\chi^2_{(1)}$ = 0.95, p = 0.330), nor SCH23390 ($\chi^2_{(1)}$ = 0.33, p =

0.568) dosage were predictive of attention AUROC (data not shown). Comparable results

were obtained when these analyses were limited to only broad spiking cells, or cells that

showed both an attention and drug effect.

As drug application strongly inhibited firing rates across the population and we found cell-

type specific effects on attention-specific activity, we next investigated whether drug

application affected behavioral performance (Figure 5). To this end, we used sequential

multilevel model analyses to test for fixed effects of attention and drug application, as well as

their interaction, on RT. Neither attention (Dopamine: $\beta$ = -13.49±8.88, p = 0.132;

SCH23390: $\beta$ = 2.86±11.34, p = 0.802), nor drug application (Dopamine: $\beta$ = -3.47±8.88, p =

0.697; SCH23390: $\beta$ = 10.38±11.34, p = 0.363) nor their interaction (Dopamine: $\beta$ =

2.87±5.62, p = 0.611; SCH23390: $\beta$ = -4.33±7.17, p = 0.548) were predictive of RT for either

drug. Given the focal nature of micro-iontophoretic drug application (Herz et al., 1969), the

absence of an effect of drug application on behavioral performance is not surprising and in-

line with comparable work on DA in PFC (Vijayraghavan et al., 2007; Jacob et al., 2013,

2016).

Interestingly, however, we found that the application of both DA and SCH23390 influenced

pupil diameter. We conducted a sliding-window Wilcoxon signed rank analysis for each 200

ms window, in 10 ms increments, comparing baseline-normalized pupil diameter on drug

compared to no-drug trials (Figure 6A). This analysis revealed a significant difference in

pupil diameter that started after stimulus onset and lasted until after cue onset. Specifically,

we found a small, but significant, modulation of the pupillary light reflex (Figure 6). The

magnitude of the constriction of the pupil upon stimulus onset was reduced during

dopaminergic drug application compared to control trials, but neither drug influenced pupil

diameter during any other time window (Figure 6B-E). Another sliding window analysis

using a two factor (drug by attention) repeated measures ANOVA revealed no effect of

attention (main or interaction) on pupil diameter (data not shown). Thus, locally applied

dopaminergic drugs in parietal cortex modulated the pupillary light reflex.

## Discussion

We tested the effects of dopaminergic drugs on PPC activity during spatial selective

attention. The non-specific agonist dopamine inhibited activity according to an inverted U-

shaped dose-response curve, whereas the D1R antagonist decreased firing rates for broad-

spiking units following a monotonic dose-response curve. Dopamine additionally reduced

attention-related firing rate modulations in broad-spiking units. Finally, local drug application

in parietal cortex decreased the pupillary light reflex. This is the first study (to the best of our

knowledge) revealing the role of dopaminergic modulation on attention-related activity in

parietal cortex.

### General and cell-type specific dopaminergic modulation in parietal cortex

We distinguished between broad and narrow-spiking units. Even though, as discussed above,

this classification does not reflect a one-to-one mapping onto interneurons and pyramidal

cells, this categorization may explain some of our results (Jacob et al., 2013, 2016).

Dopamine has a well-established role in modulating prefrontal signaling, supporting

cognitive functions such as working memory and attention (Williams and Goldman-Rakic,

1995; Watanabe et al., 1997; Vijayraghavan et al., 2007; Noudoost and Moore, 2011b; Clark

and Noudoost, 2014; Thiele and Bellgrove, 2018; Ott and Nieder, 2019). D1R and D2R are

expressed broadly throughout the cortex and fulfil complementary roles in prefrontal

cognitive control (Ott and Nieder, 2019). Although D2Rs have been implicated in rule coding

(Ott et al., 2014), modulation of working memory is mostly associated with D1R stimulation

or blockade (Sawaguchi et al., 1990; Sawaguchi and Goldman-Rakic, 1991, 1994; Williams

and Goldman-Rakic, 1995). Moreover, while manipulation of either receptor subtype in FEF

can modulate behavioral choices (Soltani et al., 2013), only D1R blockade in FEF elicited

activity resembling attentional effects in extrastriate visual areas (Noudoost and Moore,

2011a). Interestingly, D1R expression is higher in FEF pyramidal cells compared to

interneurons (Mueller et al., 2018, 2019). Here, dopaminergic drugs affected broad-spiking

more than narrow-spiking units. Although it is unknown whether dopamine receptor

expression differs across cell types in PPC, if expression is similar to the FEF, modulation of

parietal attentional signals might rely on higher expression of D1R compared to D2R in

broad-spiking putative pyramidal cells.

It is remarkable that the majority of the recorded neurons were inhibited by dopamine and

SCH23390 application, as previous studies (in prefrontal cortex) found mixed responses to

unselective dopamine (Jacob et al., 2013) or D1R stimulation (Williams and Goldman-Rakic,

1995; Vijayraghavan et al., 2007). These effects could theoretically be due to our

recording/iontophoresis setup. As both agonists and antagonists elicited responses of the

same sign, effects unrelated to specific drugs could have been ruled out by control recordings

using saline or by compensating ejection currents. Similar control experiments from our lab

(and other labs) have, however, never resulted in systematic (condition specific) effects

(Herrero et al., 2008, 2013, 2017; Thiele et al., 2012; Jacob et al., 2013; Ott et al., 2014; Ott

and Nieder, 2016; Dasilva et al., 2019). Further, the cell-type specific effects and U-shaped

dose-response curve argue against our results being an iontophoresis artefact.

These effects may alternatively be explained by drug dosages. Although Jacob et al. (2013)

used a variety of ejection currents (25-100 nA) and the proportion of inhibited or excited cells

did not differ by dosage, activity increases have been found for low, and decreases for high

D1R-agonist and antagonist dosages (Williams and Goldman-Rakic, 1995; Vijayraghavan et

al., 2007). Indeed, while our sample size using lower dosages was small, lower ejection

currents predicted positive and less negative modulation. At the dosages used in this study,

dopamine could have mostly inhibitory effects. Vijayraghavan et al. (2007) found that low

doses (10-20 nA) of D1-agonists reduced overall firing rates, but increased spatial specificity

of prefrontal neurons, whereas high dosages (20-100 nA) further reduced activity and

abolished spatially selective information. Given that our study was unrelated to spatial

specificity (i.e. saccade field tuning), we were unable to assess this particular feature, but

dopaminergic influences may still enhance spatial tuning of PPC despite an overall reduction

in activity.

Another factor that could explain the low number of dopamine-excited units is the short

block duration used in our task. Cells excited by dopamine respond slower to drug

application than inhibited cells, with an average modulation up-ramp time constant of 221.9 s

(Jacob et al., 2013). In our task, with a median trial duration of approximately 8 s, a block (36

trials) lasted approximately 288 s. Dopamine-excited neurons could have only started to show

modulation towards the end of the block, resulting in a population of largely inhibited units.

In sum, dopaminergic effects on (task-related) activity are complex (Seamans and Yang,

2004) and depend on various factors not controlled for in this study, such as endogenous

levels of dopamine. Within prefrontal cortex, coding can be enhanced by D1R agonists, and

diminished by antagonists (Vijayraghavan et al., 2007; Ott et al., 2014), or vice-versa

(Williams and Goldman-Rakic, 1995; Noudoost and Moore, 2011a). Indeed, dopaminergic

effects show regional variability across different brain areas, even within PFC (Arnsten et al.,

2012). Thus, the mechanisms discussed above might not apply to PPC. Future studies are

needed to further elucidate cell-type and receptor-subtype specific effects of dopamine in

parietal cortex during task performance.

### Dopaminergic dose-response curve

Dopamine receptor stimulation follows an inverted-U shaped dose-response curve whereby

too little or too much stimulation leads to suboptimal behavioral performance (Arnsten et al.,

1994; Zahrt et al., 1997) or neural coding (Vijayraghavan et al., 2007). Whereas optimal

levels of dopamine receptor stimulation can stabilize and tune neural activity, suboptimal

levels decrease neural coding and behavioral performance.

Here we found an inverted-U shaped dose-response curve for the unselective agonist

dopamine, and a monotonic function for the D1R antagonist SCH23390. Rather than

predicting neural coding for attention, however, ejection currents were merely predictive of

drug modulation indices, without any relationship to attention AUROC values. However, our

sample size, especially for SCH23390, might have been too small to reliably determine the

shape of the dose-response curve. Additionally, the dopaminergic effects might partly be

driven by receptor subtypes (e.g. D2R) not usually associated with modulation of delay

period activity. While this study provides evidence for a role of dopamine in parietal cortex

during cognitive tasks, further research is required to elucidate the exact underlying

mechanisms.

## Dopaminergic modulation of the pupil light reflex

The pupil light reflex (PLR) transiently constricts the pupil after exposure to increases in illumination or presentation of bright stimuli (Loewenfeld, 1993; McDougal and Gamlin, 2014). Recent studies have shown that covert attention can modulate this behavioral reflex (Naber et al., 2013; Binda and Murray, 2015a, 2015b). Subthreshold FEF microstimulation respectively enhances or reduces the PLR when a light stimulus is presented inside or outside the saccade field (Ebitz and Moore, 2017). The PLR thus depends both on luminance changes and the location of spatial attention. We found that dopaminergic drug application in parietal cortex reduced the PLR. These results are in agreement with the electrophysiological results, as drug administration also reduced attentional rate modulation. Two (non-exclusive) mechanisms have been proposed by which FEF can modulate the PLR (Binda and Gamlin, 2017). By direct or indirect projections to the olivary pretectal nucleus, or via indirect projections to constrictor neurons in the Edinger-Westphal nucleus. For the latter, these projections are hypothesized to pass through extrastriate visual cortex and/or the superior colliculus (SC). Subthreshold microstimulation of the intermediate (SCi), but not superficial (SCs), layers of the SC elicits a short latency pupillary dilation (Wang et al., 2012; Joshi et al., 2016). Whereas the SCs receives input from early visual areas, including the retina, the SCi receives input from higher-order association cortices. Along with preparing and executing eye movements, the SCi is involved in directing covert attention (Kustov and Lee Robinson, 1996; Ignashchenkova et al., 2004; Muller et al., 2005; Lovejoy and Krauzlis, 2010), and provides an essential contribution to the selection of stimuli amongst competing distractors (McPeek and Keller, 2002, 2004; reviewed in Mysore and Knudsen, 2011). Moreover, the SC receives dense projections from parietal cortex (Kuypers and Lawrence, 1967; Becker, 1989), and has been hypothesized to play an important role in pupil diameter modulation (Wang and Munoz, 2015). It is currently unclear whether dopaminergic

modulation of frontal (or parietal) cortex modulates SC activity, but this pathway seems a

strong candidate for the modulation of the PLR (Wang and Munoz, 2015) that we

encountered in this study through DA application.

Dopamine is an important modulator of high-level cognitive functions, both in the healthy

and ageing brain as well as for various clinical disorders (Robbins and Arnsten, 2009;

Arnsten et al., 2012; Thiele and Bellgrove, 2018). Although dopaminergic effects within PFC

have been elucidated in some detail, the effects of dopamine in other brain areas such as

parietal cortex, despite its well-established role in cognition and cognitive dysfunction, has

largely been overlooked. This study is the first to show dopaminergic modulation of parietal

activity in general, and activity specific to spatial attention in the non-human primate. Our

work encourages future studies of dopaminergic involvement in parietal cortex, thereby

gaining a broader understanding of neuromodulation in different networks for cognition.

## References

References

Ardid S, Vinck M, Kaping D, Marquez S, Everling S, Womelsdorf T (2015) Mapping of

functionally characterized cell classes onto canonical circuit operations in primate

prefrontal cortex. J Neurosci.

Arnsten AFT, Cai JX, Murphy BL, Goldman-Rakic PS (1994) Dopamine D1 receptor

mechanisms in the cognitive performance of young adult and aged monkeys.

Psychopharmacology (Berl).

Arnsten AFT, Wang MJ, Paspalas CD (2012) Neuromodulation of Thought: Flexibilities and

Vulnerabilities in Prefrontal Cortical Network Synapses. Neuron 76:223–239.

Becker W (1989) The neurobiology of saccadic eye movements. Metrics. Rev Oculomot Res

3:13–67 Available at: http://www.ncbi.nlm.nih.gov/pubmed/2486323.

Bellgrove MA, Barry E, Johnson KA, Cox M, Dáibhis A, Daly M, Hawi Z, Lambert D,

Fitzgerald M, McNicholas F, Robertson IH, Gill M, Kirley A (2008) Spatial attentional

bias as a marker of genetic risk, symptom severity, and stimulant response in ADHD.

Neuropsychopharmacology 33:2536–2545.

Bellgrove MA, Chambers CD, Johnson KA, Daibhis A, Daly M, Hawi Z, Lambert D, Gill M,

Robertson IH (2007) Dopaminergic genotype biases spatial attention in healthy children.

Mol Psychiatry 12:786–792.

Bellgrove MA, Johnson KA, Barry E, Mulligan A, Hawi Z, Gill M, Robertson I, Chambers

CD (2009) Dopaminergic Haplotype as a Predictor of Spatial Inattention in Children

With Attention-Deficit/Hyperactivity Disorder. Arch Gen Psychiatry 66:1135 Available

at:

http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archgenpsychiatry.2009.120

.

Bellgrove MA, Mattingley JB (2008) Molecular genetics of attention. In: Annals of the New

York Academy of Sciences, pp 200–212.

Berger B, Gaspar P, Verney C (1991) Dopaminergic innervation of the cerebral cortex:

Unexpected differences between rodents and primates. Trends Neurosci 14:21–27

Available at: http://linkinghub.elsevier.com/retrieve/pii/016622369190179X [Accessed

January 14, 2015].

Binda P, Gamlin PD (2017) Renewed Attention on the Pupil Light Reflex. Trends Neurosci

40:455–457 Available at: http://www.cell.com/trends/neurosciences/pdf/S0166-

2236(17)30122-4.pdf [Accessed July 26, 2017].

Binda P, Murray SO (2015a) Spatial attention increases the pupillary response to light

changes. J Vis 15:1–1.

Binda P, Murray SO (2015b) Keeping a large-pupilled eye on high-level visual processing.

Trends Cogn Sci 19:1–3 Available at: http://dx.doi.org/10.1016/j.tics.2014.11.002.

Caspers S, Schleicher A, Bacha-Trams M, Palomero-Gallagher N, Amunts K, Zilles K (2013)

Organization of the human inferior parietal lobule based on receptor architectonics.

Cereb Cortex 23:615–628.

Clark CR, Geffen GM, Geffen LB (1989) Catecholamines and the covert orientation of

attention in humans. Neuropsychologia 27:131–139.

Clark KL, Noudoost B (2014) The role of prefrontal catecholamines in attention and working

memory. Front Neural Circuits 8:33 Available at:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3986539&tool=pmcentrez&

rendertype=abstract [Accessed November 5, 2014].

Corbetta M, Shulman GL (2011) Spatial Neglect and Attention Networks. Annu Rev

Neurosci 34:569–599 Available at:

http://www.annualreviews.org/doi/abs/10.1146/annurev-neuro-061010-113731.

Dasilva M, Brandt C, Gotthardt S, Gieselmann MA, Distler C, Thiele A (2019) Cell class-

specific modulation of attentional signals by acetylcholine in macaque frontal eye field.

Proc Natl Acad Sci:201905413 Available at:

http://www.pnas.org/lookup/doi/10.1073/pnas.1905413116.

Desimone R, Duncan J (1995) Neural Mechanisms of Selective Visual Attention. Annu Rev

Neurosci 18:193–222 Available at:

http://www.annualreviews.org/doi/abs/10.1146/annurev.ne.18.030195.001205.

Distler C, Hoffmann K-P (2001) Cortical Input to the Nucleus of the Optic Tract and Dorsal

Terminal Nucleus (NOT-DTN) in Macaques: a Retrograde Tracing Study. Cereb Cortex

11:572–580 Available at: https://academic.oup.com/cercor/article-

lookup/doi/10.1093/cercor/11.6.572.

Ebitz RB, Moore T (2017) Selective Modulation of the Pupil Light Reflex by

Microstimulation of Prefrontal Cortex. J Neurosci 37:5008–5018 Available at:

http://www.jneurosci.org/content/jneuro/early/2017/04/21/JNEUROSCI.2433-

16.2017.full.pdf [Accessed April 24, 2017].

Furey ML, Pietrini P, Haxby J V., Drevets WC (2008) Selective effects of cholinergic

modulation on task performance during selective attention. Neuropsychopharmacology.

Gieselmann MA, Thiele A (2008) Comparison of spatial integration and surround

suppression characteristics in spiking activity and the local field potential in macaque

V1. Eur J Neurosci 28:447–459.

Gorgoraptis N, Mah YH, MacHner B, Singh-Curry V, Malhotra P, Hadji-Michael M, Cohen

D, Simister R, Nair A, Kulinskaya E, Ward N, Greenwood R, Husain M (2012) The

effects of the dopamine agonist rotigotine on hemispatial neglect following stroke. Brain

135:2478–2491.

Gray H, Bertrand H, Mindus C, Flecknell P, Rowe C, Thiele A (2016) Physiological,

Behavioral, and Scientific Impact of Different Fluid Control Protocols in the Rhesus

Macaque (Macaca mulatta). Eneuro 3:1–15.

Green DM, Swets JA (1966) Signal Detection Theory and Psychophysics. New York: Wiley.

Herrero JL, Gieselmann M a, Sanayei M, Thiele A (2013) Attention-induced variance and

noise correlation reduction in macaque v1 is mediated by NMDA receptors. Neuron

78:729–739 Available at:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3748348&tool=pmcentrez&

rendertype=abstract [Accessed January 6, 2015].

Herrero JL, Gieselmann MA, Thiele A (2017) Muscarinic and Nicotinic Contribution to

Contrast Sensitivity of Macaque Area V1 Neurons. Front Neural Circuits 11:1–15.

Herrero JL, Roberts MJ, Delicato LS, Gieselmann MA, Dayan P, Thiele A (2008)

Acetylcholine contributes through muscarinic receptors to attentional modulation in V1.

Nature 454:1110–1114 Available at:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2666819&tool=pmcentrez&

rendertype=abstract [Accessed October 7, 2014].

Herz A, Zieglgänsberger W, Färber G (1969) Microelectrophoretic studies concerning the

spread of glutamic acid and GABA in brain tissue. Exp Brain Res 9 Available at:

http://link.springer.com/10.1007/BF00234456.

Ignashchenkova A, Dicke PW, Haarmeier T, Thier P (2004) Neuron-specific contribution of

the superior colliculus to overt and covert shifts of attention. Nat Neurosci 7:56–64

Available at: http://www.nature.com/articles/nn1169.

Jacob SN, Ott T, Nieder A (2013) Dopamine regulates two classes of primate prefrontal

neurons that represent sensory signals. J Neurosci 33:13724–13734 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/23966694 [Accessed November 24, 2014].

Jacob SN, Stalter M, Nieder A (2016) Cell-type-specific modulation of targets and distractors

by dopamine D1 receptors in primate prefrontal cortex. Nat Commun.

Joshi S, Li Y, Kalwani RM, Gold JI (2016) Relationships between Pupil Diameter and

Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. Neuron

89:221–234 Available at: http://dx.doi.org/10.1016/j.neuron.2015.11.028.

Kustov AA, Lee Robinson D (1996) Shared neural control of attentional shifts and eye

movements. Nature 384:74–77 Available at: http://www.nature.com/articles/384074a0.

Kuypers HGJM, Lawrence DG (1967) Cortical projections to the red nucleus and the brain

stem in the rhesus monkey. Brain Res 4:151–188 Available at:

http://dx.doi.org/10.1016/j.brainres.2016.01.006.

Levin ED, Simon BB (1998) Nicotinic acetylcholine involvement in cognitive function in

animals. Psychopharmacology (Berl) 138:217–230 Available at:

http://link.springer.com/10.1007/s002130050667.

Lewis DA, Melchitzky DS, Sesack SR, Whitehead RE, Auh S, Sampson A (2001) Dopamine

transporter immunoreactivity in monkey cerebral cortex: Regional, laminar, and

ultrastructural localization. J Comp Neurol 432:119–136.

Loewenfeld IE (1993) The pupil: anatomy, physiology, and clinical applications. Detroit:

Wayne State University.

Lovejoy LP, Krauzlis RJ (2010) Inactivation of primate superior colliculus impairs covert

651           selection of signals for perceptual judgments. Nat Neurosci 13:261–266 Available at:

652           http://www.nature.com/articles/nn.2470.

653       Maruff P, Hay D, Malone V, Currie J (1995) Asymmetries in the covert orienting of visual

654           spatial attention in schizophrenia. Neuropsychologia 33:1205–1223.

655       McDougal DH, Gamlin PD (2014) Autonomic Control of the Eye. In: Comprehensive

656           Physiology, pp 439–473. Hoboken, NJ, USA: John Wiley & Sons, Inc. Available at:

657           http://doi.wiley.com/10.1002/cphy.c140014 [Accessed July 26, 2017].

658       McPeek RM, Keller EL (2002) Saccade target selection in the superior colliculus during a

659           visual search task. J Neurophysiol 88:2019–2034 Available at:

660           http://jn.physiology.org/content/88/4/2019%5Cnhttp://jn.physiology.org/content/88/4/20

661           19.short%5Cnhttp://jn.physiology.org/content/jn/88/4/2019.full.pdf%5Cnhttp://www.nc

662           bi.nlm.nih.gov/pubmed/12364525.

663       McPeek RM, Keller EL (2004) Deficits in saccade target selection after inactivation of

664           superior colliculus. Nat Neurosci 7:757–763 Available at:

665           http://www.nature.com/articles/nn1269.

666       Mehta MA, Owen AM, Sahakian BJ, Mavaddat N, Pickard JD, Robbins TW (2000)

667           Methylphenidate Enhances Working Memory by Modulating Discrete Frontal and

668           Parietal Lobe Regions in the Human Brain. J Neurosci 20:RC65–RC65 Available at:

669           http://www.ncbi.nlm.nih.gov/pubmed/10704519.

670       Mitchell JF, Sundberg K a, Reynolds JH (2007) Differential Attention-Dependent Response

671           Modulation across Cell Classes in Macaque Visual Area V4. Neuron 55:131–141

Available at: http://www.ncbi.nlm.nih.gov/pubmed/17610822 [Accessed November 25,

2014].

Mueller A, Krock RM, Shepard S, Moore T (2019) Dopamine Receptor Expression Among

Local and Visual Cortex-Projecting Frontal Eye Field Neurons. Cereb Cortex:1–17

Available at: https://academic.oup.com/cercor/advance-

article/doi/10.1093/cercor/bhz078/5481679.

Mueller A, Shepard SB, Moore T (2018) Differential Expression of Dopamine D5 Receptors

across Neuronal Subtypes in Macaque Frontal Eye Field. Front Neural Circuits.

Muller JR, Philiastides MG, Newsome WT (2005) Microstimulation of the superior

colliculus focuses attention without moving the eyes. Proc Natl Acad Sci 102:524–529

Available at: http://www.pnas.org/cgi/doi/10.1073/pnas.0408311101.

Mysore SP, Knudsen EI (2011) The role of a midbrain network in competitive stimulus

selection. Curr Opin Neurobiol 21:653–660 Available at:

http://linkinghub.elsevier.com/retrieve/pii/S0959438811000924.

Naber M, Alvarez GA, Nakayama K (2013) Tracking the allocation of attention using human

pupillary oscillations. Front Psychol.

Nelson CL, Sarter M, Bruno JP (2005) Prefrontal cortical modulation of acetylcholine release

in posterior parietal cortex. Neuroscience 132:347–359 Available at:

https://linkinghub.elsevier.com/retrieve/pii/S0306452205000229.

Newman DP, Cummins TDR, Tong JHS, Johnson BP, Pickering H, Fanning P, Wagner J,

Goodrich JTT, Hawi Z, Chambers CD, Bellgrove M a. (2014) Dopamine Transporter

Genotype Is Associated with a Lateralized Resistance to Distraction during Attention

Selection. J Neurosci 34:15743–15750 Available at:

http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.2327-14.2014 [Accessed

November 19, 2014].

Noudoost B, Moore T (2011a) Control of visual cortical signals by prefrontal dopamine.

Nature 474:372–375 Available at:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3117113&tool=pmcentrez&

rendertype=abstract [Accessed October 10, 2014].

Noudoost B, Moore T (2011b) The role of neuromodulators in selective attention. Trends

Cogn Sci 15:585–591 Available at:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3351278&tool=pmcentrez&

rendertype=abstract [Accessed October 15, 2014].

Ott T, Jacob SN, Nieder A (2014) Dopamine Receptors Differentially Enhance Rule Coding

in Primate Prefrontal Cortex Neurons. Neuron 84:1317–1328 Available at:

http://linkinghub.elsevier.com/retrieve/pii/S0896627314010125 [Accessed December 4,

2014].

Ott T, Nieder A (2016) Dopamine D2 Receptors Enhance Population Dynamics in Primate

Prefrontal Working Memory Circuits. Cereb Cortex:1–13 Available at:

http://cercor.oxfordjournals.org/cgi/doi/10.1093/cercor/bhw244.

Ott T, Nieder A (2019) Dopamine and Cognitive Control in Prefrontal Cortex. Trends Cogn

Sci 23:213–234 Available at: https://doi.org/10.1016/j.tics.2018.12.006.

Parikh V, Kozak R, Martinez V, Sarter M (2007) Prefrontal Acetylcholine Release Controls

Cue Detection on Multiple Timescales. Neuron 56:141–154 Available at: http://ac.els-

cdn.com/S0896627307006745/1-s2.0-S0896627307006745-main.pdf?_tid=c3c68606-

4baa-11e7-b903-

00000aab0f6b&acdnat=1496858425_98631c2399f8a1cfe667358952b60577 [Accessed

June 7, 2017].

Posner M (1990) The Attention System Of The Human Brain. Annu Rev Neurosci 13:25–42.

Robbins TW, Arnsten AFT (2009) The neuropsychopharmacology of fronto-executive
function: monoaminergic modulation. Annu Rev Neurosci 32:267–287 Available at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2863127&tool=pmcentrez&
rendertype=abstract [Accessed July 20, 2014].

Sarter M, Hasselmo ME, Bruno JP, Givens B (2005) Unraveling the attentional functions of
cortical cholinergic inputs: Interactions between signal-driven and cognitive modulation
of signal detection. Brain Res Rev 48:98–111.

Sawaguchi T, Goldman-Rakic P (1991) D1 dopamine receptors in prefrontal cortex:
involvement in working memory. Science (80- ) 251:947–950 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/1825731.

Sawaguchi T, Goldman-Rakic PS (1994) The role of D1-dopamine receptor in working
memory: local injections of dopamine antagonists into the prefrontal cortex of rhesus
monkeys performing an oculomotor delayed-response task. J Neurophysiol 71:515–528
Available at: http://www.physiology.org/doi/10.1152/jn.1994.71.2.515.

Sawaguchi T, Matsumura M, Kubota K (1990) Effects of dopamine antagonists on neuronal
activity related to a delayed response task in monkey prefrontal cortex. J Neurophysiol
63:1401–1412 Available at: http://www.physiology.org/doi/10.1152/jn.1990.63.6.1401.

Seamans JK, Yang CR (2004) The principal features and mechanisms of dopamine
modulation in the prefrontal cortex. Prog Neurobiol 74:1–58 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/15381316 [Accessed July 15, 2014].

Silk TJ, Newman DP, Eramudugolla R, Vance A, Bellgrove MA (2014) Influence of

methylphenidate on spatial attention asymmetry in adolescents with attention deficit

hyperactivity disorder (ADHD): Preliminary findings. Neuropsychologia 56:178–183.

Simonsohn U (2017) Two-Lines: A Valid Alternative to the Invalid Testing of U-Shaped

Relationships with Quadratic Regressions. Ssrn.

Soares D, Goldrick I, Lemon RN, Kraskov A, Greensmith L, Kalmar B (2017) Expression of

Kv3.1b potassium channel is widespread in macaque motor cortex pyramidal cells: A

histological comparison between rat and macaque. J Comp Neurol 525:2164–2174.

Soltani A, Noudoost B, Moore T (2013) Dissociable dopaminergic control of saccadic target

selection and its implications for reward modulation. Proc Natl Acad Sci U S A

110:3579–3584 Available at:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3587234&tool=pmcentrez&

rendertype=abstract [Accessed December 29, 2014].

Thiele A, Bellgrove MA (2018) Neuromodulation of Attention. Neuron 97:769–785

Available at: https://doi.org/10.1016/j.neuron.2018.01.008 [Accessed February 27,

2018].

Thiele A, Brandt C, Dasilva M, Gotthardt S, Chicharro D, Panzeri S, Distler C (2016)

Attention Induced Gain Stabilization in Broad and Narrow-Spiking Cells in the Frontal

Eye-Field of Macaque Monkeys. J Neurosci 36:7601–7612 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/27445139.

Thiele A, Delicato LS, Roberts MJ, Gieselmann MA (2006) A novel electrode-pipette design

for simultaneous recording of extracellular spikes and iontophoretic drug application in

awake behaving monkeys. J Neurosci Methods 158:207–211 Available at:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2666830&tool=pmcentrez&

rendertype=abstract [Accessed November 18, 2014].

Thiele A, Herrero JL, Distler C, Hoffmann K-P (2012) Contribution of cholinergic and

GABAergic mechanisms to direction tuning, discriminability, response reliability, and

neuronal rate correlations in macaque middle temporal area. J Neurosci 32:16602–16615

Available at: http://www.ncbi.nlm.nih.gov/pubmed/23175816.

van Kempen J, Gieselmann MA, Boyd M, Steinmetz NA, Moore T, Engel T, Thiele A (2020)

Top-down coordination of local cortical state during selective attention.

Vigneswaran G, Kraskov A, Lemon RN (2011) Large Identified Pyramidal Cells in Macaque

Motor and Premotor Cortex Exhibit "Thin Spikes": Implications for Cell Type

Classification. J Neurosci 31:14235–14242.

Vijayraghavan S, Wang M, Birnbaum SG, Williams G V, Arnsten AFT (2007) Inverted-U

dopamine D1 receptor actions on prefrontal neurons engaged in working memory. Nat

Neurosci 10:376–384 Available at: http://www.ncbi.nlm.nih.gov/pubmed/17277774

[Accessed November 24, 2014].

Wang C-A, Boehnke SE, White BJ, Munoz DP (2012) Microstimulation of the Monkey

Superior Colliculus Induces Pupil Dilation Without Evoking Saccades. J Neurosci

32:3629–3636 Available at:

http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.5512-11.2012.

Wang C-A, Munoz DP (2015) A circuit for pupil orienting responses: implications for

cognitive modulation of pupil size. Curr Opin Neurobiol 33:134–140 Available at:

http://dx.doi.org/10.1016/j.conb.2015.03.018.

Warburton DM, Rusted JM (1993) Cholinergic control of cognitive resources.

Neuropsychobiology 28:43–46 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/8255409.

Watanabe M, Kodama T, Hikosaka K (1997) Increase of Extracellular Dopamine in Primate

Prefrontal Cortex During a Working Memory Task. J Neurophysiol 78:2795–2798

Available at: http://www.physiology.org/doi/10.1152/jn.1997.78.5.2795.

Williams G V, Goldman-Rakic PS (1995) Modulation of memory fields by dopamine D1

receptors in prefrontal cortex. Nature 376:572–575.

Zahrt J, Taylor JR, Mathew RG, Arnsten AFT (1997) Supranormal Stimulation of D 1

Dopamine Receptors in the Rodent Prefrontal Cortex Impairs Spatial Working Memory

Performance. J Neurosci 17:8528–8535 Available at:

http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.17-21-08528.1997.

# Figures

Figure 1. Experimental paradigm and unit selectivity. (**A**) Behavioral paradigm. The monkey held a

lever and fixated on a central fixation spot to initiate the trial. One of 3 colored gratings was presented

inside the receptive field (RF) of the neurons under study. After a variable delay a cue matching one

of the grating colors surrounded the fixation spot, indicating which grating was behaviorally relevant

(target). In pseudorandom order the stimuli decreased in luminance (dimmed). Upon dimming of the

target, the monkey had to release the lever. (**B**) Proportion of units that are visually responsive,

modulated by attention or drug application. Inset shows a Venn diagram of unit selectivity. Note that

1 unit was not selective for any of the experimental factors.

Figure 2. Population activity and example unit. (**A**) Population histograms for all cells recorded

during dopaminergic drug application that were selective for attention and/or drug application.

Population activity aligned to stimulus onset (left), cue onset (middle) and the first dimming event

(right), for the non-specific agonist dopamine (top) and the D1 antagonist SCH23390 (bottom). Error

bars denote ±1 SEM. (**B**) Activity from a representative cell recorded during application of the non-

specific agonist dopamine.This cell's activity, aligned to the first dimming event, was significantly

modulation by attention, drug application and showed a significant interaction between these factors.

The grey bar indicates the time window used for statistical analyses. Statistics: two-factor ANVOVA.

Figure 3. Dopaminergic modulation of firing rates across broad and narrow spiking units. (**A**)

Average spike waveforms for the population of units. (**B**) Distribution of peak-to-trough ratios.

Statistics: calibrated Hartigan's dip test (Ardid et al., 2015). (**C**) Average firing rates between no drug

and drug conditions for the non-specific agonist dopamine for attend RF (left) and attend away (right)

conditions. (**D**) Fano factors between no drug and drug conditions for the non-specific agonist

dopamine. (**E-F** Same conventions as (**C-D**) but for the D1 antagonist SCH23390. Only units that

revealed a main or interaction effect for the factor drug were included in this analysis. Statistics: two-

sided Wilcoxon signed rank test.

Figure 4. Dopaminergic modulation of AUROC values and dose-response curves. (**A-B**) Area under

the receiver operating characteristic (AUROC) curve between no drug and drug conditions for the

non-specific agonist dopamine (**A**) and the D1-receptor antagonist SCH23390 (**B**). Only cells that

revealed a main or interaction effect for the factors drug and attention were included in this analysis.

Statistics: Wilcoxon signed rank tests. (**C-D**) Drug modulation index plotted against ejection current

for the non-specific agonist dopamine (**C**) and the D1 antagonist SCH23390 (**D**). Note the reversed y-

axis. Solid and dotted lines represent significant model fits (applied to all cells simultaneously) and

their 95% confidence intervals, respectively. A monotonic relationship is shown if a first-order fit was

better than a constant fit, and a non-monotonic relationship is shown if a second-order fit was better

than a linear fit. $U^+$ indicates a significant U-shaped relationship. Only cells that revealed a main or

interaction effect for the factor drug were included in this analysis. Statistics: linear mixed-effects

model analysis.

Figure 5. Behavioral performance is unaffected by iontophoretic application of dopaminergic

drugs.Average RT on attend RF and attend away trials for the non-specific agonist dopamine (**A**) and

the D1 antagonist SCH23390 (**B**). Dots represent average RT during a single recording session.

Statistics: linear mixed-effects model analysis. Error bars denote ±1 SEM.

Figure 6. Modulation of pupil diameter by dopamine in Parietal cortex. (**A**) Baseline normalized pupil
time course aligned to stimulus onset (left), cue onset (middle) and the first dimming event (right).
The grey bar indicates the times where drug application brought about a significant difference in pupil
diameter. (**B**-**E**) Average normalized pupil diameter during pre-stimulus baseline period (**B**), after
stimulus onset, baseline corrected (**C**), after cue onset, corrected for pupil diameter before cue onset
(**D**), and before the first dimming event, corrected for pupil diameter before cue onset (**E**). Shaded
regions denote ±1 SEM. Statistics: Wilcoxon signed rank test. FDR correction was applied for the
analysis in panel **A**.

858    Tables

859    Table 1. Color values used for the 3 colored gratings across recording sessions and subjects, indicated

860    as [RGB] – luminance (cd/m$^2$). a = Undimmed values, b = dimmed values.

|  | *Red* | *Green* | *Blue* |
|---|---|---|---|
| *Monkey 1*<br><br>*Early recordings (n=29)* | a. [255 0 0] - 14.5<br><br>b. [100 0 0] - 1.4 | a. [0 128 0] – 9.1<br><br>b. [0 70  0] – 1.9 | a. [60 60 255] - 11.5<br><br>b. [10 10 140] – 2.2 |
| *Monkey 2*<br><br>*Early recordings (n=5)* | a. [220 0 0] – 12.8<br><br>b. [180 0 0] – 7.7 | a. [0 135 0] – 12.9<br><br>b. [0 110  0] – 7.3 | a. [60 60 255] – 12.2<br><br>b. [35 35 220] – 7.4 |
| *Monkey 1/2 (n=12/8)*<br><br>*Late recordings* | a. [220 0 0] – 12.8<br><br>b. [140 0 0] – 4.2 | a. [0 135 0] – 12.9<br><br>b. [0 90  0] – 4.6 | a. [60 60 255] – 12.2<br><br>b. [30 30 180] – 4.6 |

861