# HerpesDRG: a comprehensive resource for human herpesvirus antiviral drug resistance genotyping

**Authors:**

Charles OJ[1], Venturini C[1], Goldstein RA[2], Breuer J [1,3]

[1]Department of Infection, Immunity and Inflammation, University College London, Great Ormond Street Institute of Child Health, London WC1N 1EH, United Kingdom

[2]Division of Infection and Immunity, University College London, London WC1E 6BT, United Kingdom

[3]Great Ormond Street Hospital for Children National Health Service Foundation Trust, London WC1N 1LE, United Kingdom

Corresponding author:  Charles OJ - oscar.charles.18@ucl.ac.uk

**Abstract**

The prevention and treatment of many herpesvirus associated diseases is based on the utilization of antiviral therapies, however therapeutic success is limited by the development of drug resistance. A comprehensive point of truth of resistance conferring mutations has been missing but would be important to aid the development of antiviral drugs and in management of infections. We therefore developed HerpesDRG, a drug resistance mutation database for all the known important genes and current treatment options, built from a systematic review of available genotype to phenotype literature. The database is released along with an R package to provide a low barrier of entry to variant resistance annotation and clinical implication analysis from common sanger and NGS sequence data. This represents the first openly available and community maintainable knowledgebase of drug resistance mutations for the human herpes viruses (HHV), developed for the community of researchers and clinicians tackling HHV drug resistance.

**Availability**

27  The HerpesDRG database is available at github.com/ojcharles/herpesdrg-db. The R package for

28  resistance genotyping data is available at github.com/ojcharles/herpesdrg. A user-friendly

29  webserver is available at cmv-resistance.ucl.ac.uk/herpesdrg

**Introduction**

Herpesviruses and their associated diseases are a major health problem worldwide for

immunocompromised patients. The prevention and treatment of human cytomegalovirus

(HCMV) and herpes simplex virus (HSV) disease for example are essential in management of solid

organ transplant (SOT) & hematopoietic stem cell transplant (HSCT) recipients (Ariza-Heredia et

al., 2014; Emery, 2001; Lee et al., 2019; Limaye, 2002). Although a handful of treatment options

are available such as Acyclovir, Ganciclovir and Letermovir, emergence of resistance causing

mutations are now found for all approved drugs and pose a significant threat due to aggressive

disease course and a greater mortality risk (Bacon et al., 2003; Kakiuchi et al., 2017; Komatsu et

al., 2014; Lurain and Chou, 2010; Patel et al., 2014).

Drug resistance in a clinical setting is often diagnosed by sequencing followed by variant

annotation (genotyping), rather than isolating and phenotypically characterising (phenotyping)

due to cost and time benefits (Gu et al., 2019; Russell, 2018). Genotyping does however require

that any variants present have previously been phenotyped (Lurain and Chou, 2010). In these

experiments a single novel mutation is transferred into a reference virus, or a clinical isolate is

found with similar characteristics. The impact of this mutation on drug sensitivity can then be

determined by a plaque reduction assay (PRA), Bacterial Artificial Chromosome (BAC) reporter

assay or similar, returning an $IC_{50}$ against a given drug. It is typical to report the mutations impact

on antiviral sensitivity as a fold change relative to the control strain $IC_{50}$, where HCMV is currently

the only human herpes virus (HHV) with consensus guidelines relating fold change to a "resistant"

or "susceptible" phenotype (Kotton et al., 2018; Danve et al., 2002; Chou et al., 2005; Sauerbrei et

al., 2011).

An issue is that those previously phenotyped data are published in a disparate collection of

reports. This has meant that researchers often must perform a review of the scientific literature

for identified mutations in a viral isolate, which is time consuming, especially when there are

several reports for a given mutation (Paolucci et al., 2021; van der Beek et al., 2013, p. 20).

Therefore there have been published efforts to collate this data previously such as in reviews,

however the data contained is not updateable and requires reformatting for informatics (Lurain

and Chou, 2010; Sauerbrei et al., 2016). Advancing our toolbox such as by developing a

comprehensive mutation phenotype map that is community maintainable and in an open format

would solve these data requirement problems for genotypic methods. Further, it could play an

important role in developing improved antiviral therapies (Chou, 2020; Razonable et al., 2020).

User-friendly tools are also important to enable researchers to make use of such databases

(Chevillotte et al., 2010; Hayer et al., 2013). Any such tool built to analyse HHV sequence data

should support both sanger consensus sequence outputs, as this is still widely adopted, as well as

modern Next Generation Sequencing (NGS) variant outputs (Streck et al., 2023). NGS has become

a popular option and provides advantages such sensitivity to low-level variants that helps in early

decision making (Chin et al., 2013; Garrigue et al., 2016; Guermouche et al., 2020) and evidence

that accumulation of low-level variants may be associated with poor clinical outcome (Houldcroft

et al., 2016; Venturini et al., 2022). Additionally there is interest in sequencing wider than just

DNA polymerase (HSV1/2:UL30, HCMV:UL54, varicella-zoster virus (VZV):ORF28) and kinase

(HSV1/2: UL30, HCMV:UL97, VZV:ORF36) genes as the number of antiviral targets has grown

(Kawashima et al., 2017; Lurain and Chou, 2010; Serris et al., 2022). For example HCMV mutants

in UL27, UL51, UL56 and UL89 are not sequenced in many diagnostic labs (Kotton et al., 2018).

Here we present an open-source comprehensive database that links HHV mutations to impact on

drug sensitivity phenotypes that is interrogatable and maintainable by the community. The

database is released along with analytical tools suitable for resistance annotation of modern

sequence data from HCMV, HSV1, HSV2, VZV and HHV6.To the best of our knowledge published

comparable resources are comparatively affected by issues such as not covering all genes affecting

current clinically relevant drugs, inability to update data after publication and closed-source

databases that cannot be community verified (Chevillotte et al., 2010; López-Aladid et al., 2019;

Lurain and Chou, 2010; Sauerbrei et al., 2016).

**Database Construction**

Literature regarding mutation resistance phenotype were identified by a comprehensive PubMed

("Database resources of the National Center for Biotechnology Information," 2018) search with

key terms "Cytomegalovirus [Title] AND resistance [Title/Abstract]", likewise for the other HHVs.

Papers were inspected with regular expressions to detect the occurrence of mutations, from

where literature article data was manually extracted. Each entry in the database represents the

relationship between a mutation, study, control species, assay method, metadata, and fold change,

where fold change values are stored for 11 currently clinically important drugs. Multiple entries

may be present for the same variant where either a single publication provides different test

methodology to produce independent sensitivity results, or where the same variant is tested

across multiple publications. This Fold-change values come in the form of a numeric decimal value

if fold changes are possible to extract, or as a string "Resistance" or "Polymorphism" if only a

phenotype is described. Study reference data comes in the form of a title, a HTTP link and a DOI.

Assay information columns record the method of strain generation, i.e., "marker transfer", or

"isolated strain", the test method such as "PRA" or "dye uptake assay" and the control strain used

to generate a fold change. For some well-studied mutations, co-occurring mutations have been

observed to amplify resistance and are recorded where relevant (Chou, 2015; Chou et al., 2016).

Finally, any metadata such as when the entry was created, any notes and a status flag. Only rows

with the status "A" for active are returned in the R package and webserver.

It is well understood that a single key mutation is almost always the causative agent that results

in drug resistance for the HHV's, while any other mutations are often neutral polymorphisms. In

this database there are entries from isolates with multiple co-occurring mutations, often clinical

isolates. We use the following rules when including these entries: a) If the strain is sensitive in

vitro, any uncharacterised mutations are accepted as polymorphisms. B) Where a strain has a

single uncharacterised mutation, and the others are known polymorphisms, the single unknown

mutation is entered with the strains fold change. C) If there are multiple uncharacterised

mutations the entry is recorded along with co-mutations and set the status "O" for obsolete, it will

not feature in processes. d) If at a later date with new data only a single uncharacterised mutation

now is present for an entry as in (c), the entry is re-looked at and treated as in (b).

**Drug resistance genotyping**

To facilitate the resistance annotation of virus sequence data we developed Herpes Drug

Resistance Genotyping "HerpesDRG", a R package that enables the assessment of resistance

mutations present from variant VCF or FASTA sequence data (Figure 1). FASTA format sequences

are aligned to the selected viruses reference genome with MAFFT (Katoh and Standley, 2013)

using the "—add –keeplength" parameters, such that both whole genome sequences or genetic

fragments (e.g. UL23, UL54) can be inputted. SNP-sites is used to extract SNP's and downstream

functions identify indels (Page et al., 2016). For detection of low frequency mutations Variant Call

Format (VCF >= version4.0) and default Varscan2 output file are accepted, where the data has

been mapped to the relevant NCBI reference strain (Koboldt et al., 2012). All input formats are

converted to a variant which is processed using the VariantAnnotation package to predict their

effect on coding regions (Obenchain et al., 2014; R Core Team, 2014). The HerpesDRG

"call_resistance" function produces the key output, an annotated variant table including

resistance data present in the database. In the case of a mutation present having multiple entries

in the database, all relevant entries are returned.

Figure1: Graphical representation of the HerpesDRG annotation pf FASTA sequence input or VCF

inputs (dashed lines). Q – Query sequence, R – NCBI Reference sequence.

A concise clinical output can then be generated from this table using the "make_clin_table"

function (Figure 2). Here for each drug HerpesDRG identifies the mutation of maximal fold change

found in the sample present at greater than 10% frequency, then allocates a category to that drug

accordingly: High level (maximum fold change above 15), Moderate level (maximal fold change

between 5-15), Low level (the maximum fold change between 2-5), Polymorphism (less than 2,

or recorded as such), Resistant (only anecdotal resistant data was returned), NA (no mutations

returned) and evidence strength as: "Good, in vitro", or NA (no evidence). These fold change cut-

off values are in line with recommendations from 'The Third International Consensus Guidelines

on the Management of Cytomegalovirus in Solid-organ Transplantation' (Kotton et al., 2018)

which we use consistently across all viruses.

Figure2: Clinical overview table produced by the "make_clin_table" function for the example data

A10.vcf. Example code on GitHub.

**Usage**

Install HerpesDRG by running the R command: devtools::install_github("ojcharles/herpesdrg").

This will take care of installing other dependencies and the database. Inputs for the

"call_resistance" function takes as input any of the aforementioned file types, along with which

virus to genotype against and flags for whether to return all synonymous / non synonymous

mutations (default is only resistance mutants). The above examples were all based on HCMV for

consistency but work equivalently for the other herpesviruses.

**User interface**

The HerpesDRG toolset is accessible through a user-friendly web interface included in the R

package and available at cmv-resistance.ucl.ac.uk/herpesdrg/. Here users upload sequence data

as described above and select the virus of interest from the drop-down list. There the full

resistance mutations table, the simplified clinical overview and additional graphics are generated

automatically. Specific terms of use for the hosted instance are included with the shiny

application. The application was developed using the shiny framework (Chang et al., n.d.). We

include example files for whole genome analysis (VCF, varscan tab, FASTA), and specific gene

regions FASTA.

**Conclusion**

We have developed a database that comprehensively captures known drug resistance mutations

and their impact on antiviral sensitivity for the antiviral treated Humen Herpes Viruses. The

database is released in an open-source format such that it can be community maintained acting

as a point of truth for future research and informatic tools. We release this database with an

associated tool that allows a low barrier to analysing presence of resistance in sequencing data.

**Funding**

O.J.C is supported by the UKRI Medical Research Council grant "MR/N013867/1". C.V. is funded

by Wellcome Trust Grant "204870/Z/16/Z". J.B. receives funding from the NIHR UCL/UCLH

Biomedical Research Centre.

**Conflicts of interest**

None declared.

**Acknowledgements**

We would like to thank Steven A Kemp and Salvatore Camiolo for their testing of HerpesDRG.

**References**

Ariza-Heredia, E.J., Nesher, L., Chemaly, R.F., 2014. Cytomegalovirus diseases after hematopoietic
stem cell transplantation: A mini-review. Cancer Lett. 342, 1–8.
https://doi.org/10.1016/j.canlet.2013.09.004

Bacon, T.H., Levin, M.J., Leary, J.J., Sarisky, R.T., Sutton, D., 2003. Herpes Simplex Virus Resistance
to Acyclovir and Penciclovir after Two Decades of Antiviral Therapy. Clin. Microbiol. Rev.
16, 114–128. https://doi.org/10.1128/CMR.16.1.114-128.2003

Chang, W., Cheng, J., Yihui, X., Jonathan, M., n.d. shiny: Web Application Framework for R. R
package version 1.3.2.

Chevillotte, M., von Einem, J., Meier, B.M., Lin, F.-M., Kestler, H.A., Mertens, T., 2010. A new tool
linking human cytomegalovirus drug resistance mutations to resistance phenotypes. Antiviral
Res. 85, 318–327. https://doi.org/10.1016/j.antiviral.2009.10.004

Chin, E.L., da Silva, C., Hegde, M., 2013. Assessment of clinical analytical sensitivity and specificity
of next-generation sequencing for detection of simple and complex mutations. BMC Genet.
14, 6. https://doi.org/10.1186/1471-2156-14-6

Chou, S., 2020. Advances in the genotypic diagnosis of cytomegalovirus antiviral drug resistance.
Antiviral Res. 176, 104711. https://doi.org/10.1016/j.antiviral.2020.104711

Chou, S., 2015. Rapid *In Vitro* Evolution of Human Cytomegalovirus UL56 Mutations That Confer
Letermovir Resistance. Antimicrob. Agents Chemother. 59, 6588–6593.
https://doi.org/10.1128/AAC.01623-15

Chou, S., Ercolani, R.J., Lanier, E.R., 2016. Novel Cytomegalovirus UL54 DNA Polymerase Gene
Mutations Selected In Vitro That Confer Brincidofovir Resistance. Antimicrob. Agents
Chemother. 60, 3845–3848. https://doi.org/10.1128/AAC.00214-16

Chou, S., Van Wechel, L.C., Lichy, H.M., Marousek, G.I., 2005. Phenotyping of Cytomegalovirus
Drug Resistance Mutations by Using Recombinant Viruses Incorporating a Reporter Gene.
Antimicrob. Agents Chemother. 49, 2710–2715. https://doi.org/10.1128/AAC.49.7.2710-
2715.2005

Danve, C., Morfin, F., Thouvenot, D., Aymard, M., 2002. A screening dye-uptake assay to evaluate in
vitro susceptibility of herpes simplex virus isolates to acyclovir. J. Virol. Methods 105, 207–
217. https://doi.org/10.1016/s0166-0934(02)00103-9

Database resources of the National Center for Biotechnology Information, 2018. . Nucleic Acids Res.
46, D8–D13. https://doi.org/10.1093/nar/gkx1095

Emery, V., 2001. Investigation of CMV disease in immunocompromised patients. J. Clin. Pathol. 54,
84–88. https://doi.org/10.1136/jcp.54.2.84

Garrigue, I., Moulinas, R., Recordon-Pinson, P., Delacour, M.-L., Essig, M., Kaminski, H., Rerolle, J.-
P., Merville, P., Fleury, H., Alain, S., 2016. Contribution of next generation sequencing to
early detection of cytomegalovirus UL97 emerging mutants and viral subpopulations analysis
in kidney transplant recipients. J. Clin. Virol. 80, 74–81.
https://doi.org/10.1016/j.jcv.2016.04.017

Gu, W., Miller, S., Chiu, C.Y., 2019. Clinical Metagenomic Next-Generation Sequencing for Pathogen
Detection. Annu. Rev. Pathol. 14, 319–338. https://doi.org/10.1146/annurev-pathmechdis-
012418-012751

Guermouche, H., Burrel, S., Mercier-Darty, M., Kofman, T., Rogier, O., Pawlotsky, J.-M., Boutolleau,
D., Rodriguez, C., 2020. Characterization of the dynamics of human cytomegalovirus
resistance to antiviral drugs by ultra-deep sequencing. Antiviral Res. 173, 104647.
https://doi.org/10.1016/j.antiviral.2019.104647

Hayer, J., Jadeau, F., Deléage, G., Kay, A., Zoulim, F., Combet, C., 2013. HBVdb: a knowledge
database for Hepatitis B Virus. Nucleic Acids Res. 41, D566–D570.
https://doi.org/10.1093/nar/gks1022

Houldcroft, C.J., Bryant, J.M., Depledge, D.P., Margetts, B.K., Simmonds, J., Nicolaou, S., Tutill,
H.J., Williams, R., Worth, A.J.J., Marks, S.D., Veys, P., Whittaker, E., Breuer, J., 2016.
Detection of Low Frequency Multi-Drug Resistance and Novel Putative Maribavir Resistance
in Immunocompromised Pediatric Patients with Cytomegalovirus. Front. Microbiol. 7.
https://doi.org/10.3389/fmicb.2016.01317

Kakiuchi, S., Tsuji, M., Nishimura, H., Yoshikawa, T., Wang, L., Takayama-Ito, M., Kinoshita, H.,
Lim, C.-K., Fujii, H., Yamada, S., Harada, S., Oka, A., Mizuguchi, M., Taniguchi, S., Saijo,
M., 2017. Association of the Emergence of Acyclovir-Resistant Herpes Simplex Virus Type 1
With Prognosis in Hematopoietic Stem Cell Transplantation Patients. J. Infect. Dis. 215, 865–
873. https://doi.org/10.1093/infdis/jix042

Katoh, K., Standley, D.M., 2013. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Mol. Biol. Evol. 30, 772–780.
https://doi.org/10.1093/molbev/mst010

Kawashima, M., Nemoto, O., Honda, M., Watanabe, D., Nakayama, J., Imafuku, S., Kato, T.,
Katsuramaki, T., 2017. Amenamevir, a novel helicase–primase inhibitor, for treatment of
herpes zoster: A randomized, double-blind, valaciclovir-controlled phase 3 study. J. Dermatol.
44, 1219–1227. https://doi.org/10.1111/1346-8138.13948

Koboldt, D.C., Zhang, Q., Larson, D.E., Shen, D., McLellan, M.D., Lin, L., Miller, C.A., Mardis,
E.R., Ding, L., Wilson, R.K., 2012. VarScan 2: Somatic mutation and copy number alteration
discovery in cancer by exome sequencing. Genome Res. 22, 568–576.
https://doi.org/10.1101/gr.129684.111

Komatsu, T.E., Pikis, A., Naeger, L.K., Harrington, P.R., 2014. Resistance of human cytomegalovirus
to ganciclovir/valganciclovir: A comprehensive review of putative resistance pathways.
Antiviral Res. 101, 12–25. https://doi.org/10.1016/j.antiviral.2013.10.011

Kotton, C.N., Kumar, D., Caliendo, A.M., Huprikar, S., Chou, S., Danziger-Isakov, L., Humar, A.,
Group,  on behalf of the T.T.S.I.C.C., 2018. The Third International Consensus Guidelines on
the Management of Cytomegalovirus in Solid-organ Transplantation. Transplantation 102,
900–931. https://doi.org/10.1097/TP.0000000000002191

Lee, D.H., Zuckerman, R.A., AST Infectious Diseases Community of Practice, 2019. Herpes simplex
virus infections in solid organ transplantation: Guidelines from the American Society of
Transplantation Infectious Diseases Community of Practice. Clin. Transplant. 33, e13526.
https://doi.org/10.1111/ctr.13526

Limaye, A.P., 2002. Ganciclovir-Resistant Cytomegalovirus in Organ Transplant Recipients. Clin.
Infect. Dis. 35, 866–872. https://doi.org/10.1086/342385

López-Aladid, R., Guiu, A., Mosquera, M.M., López-Medrano, F., Cofán, F., Linares, L., Torre-
Cisneros, J., Vidal, E., Moreno, A., Aguado, J.M., Cordero, E., Martin-Gandul, C., Carratalá,
J., Sabé, N., Niubó, J., Cervera, C., Capón, A., Cervilla, A., Santos, M., Bodro, M., Muñoz, P.,
Fariñas, M.C., Antón, A., Aranzamendi, M., Montejo, M., Pérez-Romero, P., Len, O., Marcos,
M.Á., 2019. Improvement in detecting cytomegalovirus drug resistance mutations in solid
organ transplant recipients with suspected resistance using next generation sequencing. PLOS
ONE 14, e0219701. https://doi.org/10.1371/journal.pone.0219701

Lurain, N.S., Chou, S., 2010. Antiviral Drug Resistance of Human Cytomegalovirus. Clin. Microbiol.
Rev. 23, 689–712. https://doi.org/10.1128/CMR.00009-10

Obenchain, V., Lawrence, M., Carey, V., Gogarten, S., Shannon, P., Morgan, M., 2014.
VariantAnnotation : a Bioconductor package for exploration and annotation of genetic
variants. Bioinformatics 30, 2076–2078. https://doi.org/10.1093/bioinformatics/btu168

Page, A.J., Taylor, B., Delaney, A.J., Soares, J., Seemann, T., Keane, J.A., Harris, S.R., 2016. SNP-
sites: rapid efficient extraction of SNPs from multi-FASTA alignments. Microb. Genomics 2.
https://doi.org/10.1099/mgen.0.000056

Paolucci, S., Campanini, G., Cassaniti, I., Tebaldi, A., Novazzi, F., Fratini, A., Meini, A., Girelli, F.,
Palumbo, L., Plebani, A., Baldanti, F., 2021. Emergence of Letermovir-resistant HCMV UL56
mutant during rescue treatment in a liver transplant recipient with ganciclovir-resistant
infection HCMV: a case report. BMC Infect. Dis. 21, 994. https://doi.org/10.1186/s12879-
021-06694-4

Patel, S.J., Kuten, S.A., Knight, R.J., Hong, D.M., Gaber, A.O., 2014. Resolution of Mild Ganciclovir-
Resistant Cytomegalovirus Disease with Reduced-Dose Cidofovir and CMV-Hyperimmune
Globulin. J. Transplant. 2014, 342319. https://doi.org/10.1155/2014/342319

R Core Team, 2014. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Razonable, R.R., Inoue, N., Pinninti, S.G., Boppana, S.B., Lazzarotto, T., Gabrielli, L., Simonazzi, G.,
Pellett, P.E., Schmid, D.S., 2020. Clinical Diagnostic Testing for Human Cytomegalovirus
Infections. J. Infect. Dis. 221, S74–S85. https://doi.org/10.1093/infdis/jiz601

Russell, D.A., 2018. Sequencing, Assembling, and Finishing Complete Bacteriophage Genomes, in:
Clokie, M.R.J., Kropinski, A.M., Lavigne, R. (Eds.), Bacteriophages: Methods and Protocols,
Volume 3, Methods in Molecular Biology. Springer, New York, NY, pp. 109–125.
https://doi.org/10.1007/978-1-4939-7343-9_9

Sauerbrei, A., Bohn-Wippert, K., Kaspar, M., Krumbholz, A., Karrasch, M., Zell, R., 2016. Database
on natural polymorphisms and resistance-related non-synonymous mutations in thymidine
kinase and DNA polymerase genes of herpes simplex virus types 1 and 2. J. Antimicrob.
Chemother. 71, 6–16. https://doi.org/10.1093/jac/dkv285

Sauerbrei, A., Taut, J., Zell, R., Wutzler, P., 2011. Resistance testing of clinical varicella-zoster virus
strains. Antiviral Res. 90, 242–247. https://doi.org/10.1016/j.antiviral.2011.04.005

Serris, A., Pouvaret, A., Loiseau, C., Abid, H., Burrel, S., Fourgeaud, J., Rouzaud, C., Lanternier, F.,
Boutolleau, D., Frange, P., 2022. Pritelivir for recurrent aciclovir-resistant herpes simplex
virus 2 infections in immunocompromised patients. J. Antimicrob. Chemother. dkac165.
https://doi.org/10.1093/jac/dkac165

Streck, N.T., Espy, M.J., Ferber, M.J., Klee, E.W., Razonable, R.R., Gonzalez, D., Sayada, C., Heaton,
P.R., Chou, S., Binnicker, M.J., 2023. Use of next-generation sequencing to detect mutations
associated with antiviral drug resistance in cytomegalovirus. J. Clin. Microbiol. e0042923.
https://doi.org/10.1128/jcm.00429-23

van der Beek, M.T., Vermont, C.L., Bredius, R.G.M., Marijt, E.W.A., van der Blij-de Brouwer, C.S.,
Kroes, A.C.M., Claas, E.C.J., Vossen, A.C.T.M., 2013. Persistence and Antiviral Resistance of
Varicella Zoster Virus in Hematological Patients. Clin. Infect. Dis. 56, 335–343.
https://doi.org/10.1093/cid/cis879

Venturini, C., Colston, J.M., Charles, O., Best, T., Atkinson, C., Forrest, C., Williams, C., Rao, K.,
Worth, A., Thorburn, D., Harber, M., Griffiths, P., Breuer, J., 2022. Persistent low-level
variants in a subset of HCMV genes are highly predictive of poor outcome in
immunocompromised patients with cytomegalovirus infection.
https://doi.org/10.1101/2022.09.12.22279849

311