β-catenin perturbations control differentiation programs in mouse embryonic
stem cells

Elisa Pedone[1,2,+], Mario Failli[3,4], Gennaro Gambardella[3,4], Rossella De Cegli[3], Diego
di Bernardo[3,4] and Lucia Marucci[1,2,5,+].

[1]Department of Engineering Mathematics, University of Bristol, Bristol BS8 1UB, UK. [2]School
of Cellular and Molecular Medicine, University of Bristol, Bristol BS8 1TD, UK. [3]Telethon
Institute of Genetic and Medicine Via Campi Flegrei 34, 80078 Pozzuoli, Italy.[4]Department of
Chemical, Materials and Industrial Production Engineering, University of Naples Federico II,
80125 Naples, Italy. [5]BrisSynBio, Bristol BS8 1TQ, UK.

[+] corresponding authors; elisa.pedone@bristol.ac.uk; lucia.marucci@bristol.ac.uk

Abstract

The Wnt/β-catenin pathway is involved in development, cancer and embryonic stem
cell (ESC) maintenance; its dual role in stem cell self-renewal and differentiation is still
controversial. Here, we applied an elegant *in vitro* system enabling conditional β-
catenin control in β-catenin null mouse ESCs. We report that moderate induction of
exogenous β-catenin enhances epiblast stem cell (EpiSC) derivation *in vitro*. Using a
different genetic model and β-catenin chemical modulation, we derived a new protocol
for ESCs to EpiSCs differentiation, based on NDiff227 and the GSK3α/β inhibitor
Chiron, that is more efficient than standard ActivinA/Fgf2-based protocols. Finally, we
report that moderate β-catenin levels favour early stem cell commitment towards
mesoderm if the protein is overexpressed only in the 'ground state' of pluripotency
conditions, or endoderm if the overexpression is maintained during the differentiation,
unravelling the controversial role of this signalling pathway at the exit from
pluripotency.

Introduction

Pluripotent Cells (PCs) are characterized by indefinite proliferative and differentiation
potential and their identity is determined by the balance between signals promoting
self-renewal and differentiation. The first step for stem cell differentiation is the exit
from the pluripotent state, tightly controlled by underlying gene regulatory network
dynamics which can drive specific lineage commitment. During murine development
in vivo, embryonic stem cells (ESCs), that represent the naïve pluripotent state of the
early epiblast[1-3], convert into the late epiblast and finally in terminally differentiated

somatic cells. ESCs can be derived from the pre-implantation epiblast and used to

study cell pluripotency and differentiation, providing an excellent *in vitro* system for

understanding signalling pathway interplay in cell fate decision making.

In serum-based cultures, mouse ESCs (hereafter called ESCs) are heterogeneous for

the expression of pluripotency genes[1,2,4-10], while, when cultured in serum-free media

supplemented with inhibitors of MEK1/2 (PD) and GSK3α/β (Chiron) and in presence

or not of the Leukaemia Inhibitory Factor-LIF (2i or 2i+LIF)[3], a uniform self-renewal

condition known as 'ground state' of pluripotency is established; it is characterized by

homogenous gene expression[11-14], genome demethylation[15-17] and stable naïve

pluripotency[18,19]. Mouse epiblast stem cells (hereafter called EpiSCs) also represent

a relevant *in vitro* model because of their similarities with embryonic stem cells of

human origin[20]. EpiSCs, derived from the post-implantation epiblast, are also capable

of differentiating in all the germ-layers[20,21]; however, they differ from ESCs in

morphology, clonogenicity, gene expression, epigenome status and, most importantly,

ability to contribute to chimaeras[14,20-22]. EpiSCs require ActivinA and the fibroblast

growth factor 2 (FGF2)[20,21] for *in vitro* expansion; of note, FGF signalling pathway

activation, while promoting EpiSC self-renewal, induces ESC differentiation[23,24].

FGF2 treatment, in combination or not with ActivinA and LIF/STAT3 pathway

inhibitors, has been used for ESCs differentiation into EpiSCs both in serum-based

and serum-free culture conditions[25-28], although with low efficiency. Self-renewing

EpiSCs have been recently established by simultaneous activation and inhibition of

the Wnt/β-catenin pathway[29]; however, the effect of these perturbations on ESCs-

EpiSCs direct transition has not been fully explored.

The Wnt/β-catenin is a highly conserved signalling pathway involved in ESCs self-

renewal[30] and cell-cycle progression[31]. β-catenin levels are tightly controlled by the

active transcription of negative regulators working at different levels of the signalling

cascade[32]: Axin2[33-35] is part of the disruption complex whereas DKK1[36] binds the Wnt

receptor complex attenuating cellular response upon activation of the pathway. These

negative feedback loops contribute to the emergence of nonlinear dynamics in the

Wnt/β-catenin pathway, proved to be important in different biological and

developmental aspects (as reviewed in[37]), ESCs pluripotency[10] and somatic cell

reprogramming[10,38-41]. The role of the canonical Wnt pathway in early *in vivo*

developmental stages and the requirement of its activation for ESC self-renewal have

been a matter of intense research, often generating contradictory results[3,42-47].

Pluripotency incompetence has been reported in two independent studies using β-

catenin[-/-] ECSs[42,44]; this phenotype was, however, contradicted in later studies with

newly generated β-catenin[-/-] cell lines, which normally self-renew in both serum and

2i+LIF (hereafter called 2i/L), but present some differentiation defects when LIF-

deprived[45-47]. Such knock-out models provide an excellent *in vitro* system to study β-

catenin function on ESC decision making.

Here, we take advantage of the β-catenin[-/-] ESC line generated by Aulicino and

colleagues[45], where the entire β-catenin coding sequence was removed to avoid

possible compensatory mechanisms from aberrant truncated isoforms, to study the

effect of β-catenin perturbations on the exit from pluripotency and differentiation.

Different β-catenin doses have been indirectly achieved in the past by mutating the

adenomatous polyposis coli gene (APC)[48]; teratomas from the mutants with the

highest β-catenin transcriptional activity showed major differentiation defects in the

neuroectoderm, dorsal mesoderm and endoderm lineages. Of note, results in[48]

suggest that active β-catenin nuclear translocation (different across mutants) might

also be involved in the observed differentiation impairment. Cellular models enabling

direct modulation of β-catenin are necessary to systematically associate protein

perturbations to pluripotency and differentiation phenotypes.

We tuned β-catenin levels in β-catenin[-/-] ESCs applying an improved inducible

system[49] and measured both the global gene expression following 2i/L withdrawal,

and the efficiency of ESC-EpiSC transition *in vitro*. Comparing the response to

differentiation stimuli of ESCs expanded in serum/LIF or 2i/L, we demonstrated that

moderate β-catenin overexpression in β-catenin[-/-] ESCs enhances the differentiation

efficiency into EpiSCs. Moreover, short-term expansion in 2i/L predisposes ESCs to

such transition, challenging the hypothesis of a synergistic effect of the ERK, Wnt/β-

catenin and STAT pathways on EpiSCs derivation *in vitro* that would require further

investigation. These results were recapitulated by exposing wild-type ESCs to low

doses of the GSK3α/β inhibitor Chiron, further confirming our findings and providing

an improved protocol for fast and efficient *in vitro* derivation of EpiSCs. Finally, the

transcriptome of ESCs expressing different β-catenin levels confirmed what we and

others reported about β-catenin dispensable requirement for pluripotency

establishment[45-47,49], while suggesting that specific β-catenin perturbations cause a
bias towards the endoderm lineage.

Overall, our study highlights that synergistic effects of β-catenin doses and culture
conditions control *in vitro* ESC fate decision making.

Results

Wnt/β-catenin pathway perturbations control *in vitro* generation of EpiSC

To study the role of the Wnt/β-catenin pathway in EpiSC derivation *in vitro*, we used
the C1-EF1a-rtTA_TRE3G-DDmCherryβ-catenin^{S33Y} (hereafter called C1) ESC line
we previously generated[49]. Briefly, β-catenin$^{-/-}$ ESCs[45] were modified to stably
express a doxycycline-inducible fusion protein comprising the conditional destabilising
domain (DD), the mCherry fluorescent protein and the constitutive active β-
catenin^{S33Y}(Figure 1A)[50]. The inducer molecule doxycycline (Doxy) enables
transcriptional initiation, while trimethoprim (TMP) allows protein stabilisation by
inactivating the DD (Figure 1A)[49]. The use of a constitutively active and conditional β-
catenin^{S33Y} form, uncoupled from upstream endogenous regulations and in a knock-
out background, would help in dissecting β-catenin functions in the ESC-EpiSC
transition, avoiding compensatory mechanisms and possible off-target effects of
stable over-expression and chemical compounds.

We confirmed, in C1 cells, the correct induction (Figure 1A, inset and[49]), intracellular
distribution and functionality of the exogenous protein upon input administration[49]. We
previously confirmed the dispensable role of β-catenin in pluripotent culture conditions
and showed, using Alkaline Phosphatase Staining, that moderate β-catenin
overexpression (i.e., TMP10µM_Doxy10ng/mL) can protect cells from exiting
pluripotency in the absence of both serum and LIF[49]. Following these results, we
measured the efficiency of EpiSCs derivation when different doses of exogenous β-
catenin are overexpressed in pluripotent conditions and/or during differentiation,
adapting an existing differentiation protocol[29,51] (see Methods for details). To
appreciate cellular response changes depending on the culture condition, C1 ESCs
were expanded either in serum/LIF (hereafter called FBS/L) or in 2i/L. ESCs in the
'ground state' media (i.e. 2i/L)[3] have been extensively characterised for their
transcriptional and epigenetic homogeneity[12,14] and resemblance of the pre-
implantation epiblast[11,15-18]. However, prolonged culture in 2i/L results in epigenetic

changes impairing normal differentiation *in vitro* and development *in vivo*[52]. Therefore, we opted for a short-term culture in 2i/L (3 passages), sufficient to obtain cell homogeneity while avoiding possible aberrations.

ESCs from FBS/L or 2i/L (Figure 1B, C) were cultured for 48 hrs either in DMSO (Figure 1B) or in presence of maximum TMP (10µM) combined with low (10ng/mL) or saturating (100ng/mL) Doxy (Figure 1C). The concentrations of Doxy were extrapolated from flow cytometry measurements of the mCherry signal to provide two doses (moderate and high) of the exogenous protein (Figure 1A, inset). We next seeded 1.5×10^4 cells/cm^2 on fibronectin-coated plates in NDiff227[26] supplemented with ActivinA, FGF2[29] and different combinations of DMSO, TMP and Doxy (Figure 1B, C and Methods). Cells kept under these conditions for 4 days, with media refreshed after the first 2 culture days, were subsequently analysed for the expression of the Epiblast marker Fgf5 by qPCR (Figure 1B, C).

TMP/Doxy pre-treatment before differentiation (Figure 1C) did not alter the basal Fgf5 expression of FBS/L ESCs (T0 samples in Figure 1D), whereas that of TMP10µM_Doxy10ng/mL treated ESCs from 2i/L culture was higher compared to the TMP10µM_Doxy100ng/mL and the TMP10µM treated samples (T0 samples in Figure 1E).

Upon 4 days of differentiation, in all Doxy/TMP treated conditions, Fgf5 levels were higher when C1 ESCs were pre-cultured in 2i/L, suggesting that the latter predisposes cells to more efficiently differentiate into EpiSCs (Figure 1E). In both scenarios (i.e., from FBS/L or 2i/L), β-catenin overexpression in pluripotent conditions only (i.e., "Before" in Figure 1D, E), or both before and during differentiation (i.e., "Always" in Figure 1D, E), did not increase Fgf5 levels as compared to the control (i.e. TMP10µM-treated C1 ESCs, Figure 1D, E). Interestingly, FBS/L pre-cultured C1 ESCs induced with a low amount of Doxy (TMP10µM_Doxy10ng/mL "During Differentiation" sample, Figure 1D) converted into EpiSCs more efficiently as compared to all other treatments from FBS/L (Figure 1D).

Altogether, these results indicate that both the cell culture media and the doses of exogenous β-catenin strongly influence how cells respond to the ActivinA/FGF2 differentiation stimulus, with the 2i/L pre-culture enabling more homogeneous differentiation towards Epiblast, and moderate β-catenin overexpression in FBS/L-derived ESCs during the differentiation protocol only improving EpiSCs establishment *in vitro*.

Figure 1. Dual-input control of β-catenin doses in EpiSC derivation *in vitro*.

A Dual-input regulation system consisting of the doxycycline responsive element and the conditionally destabilised mCherryβ-catenin^{S33Y} module. Doxycycline (Doxy) and trimethoprim (TMP) allow mCherryβ-catenin^{S33Y} transcription initiation and protein stabilisation, respectively. **(A, inset)** Flow cytometry profile of C1 ESCs treated for 24 hrs with TMP10μM and the indicated concentrations of Doxy. **B, C** Experimental scheme ESC to EpiSC differentiation. FBS/L and 2i/L C1 ESCs were pre-treated either with DMSO **(B)** or TMP10μM and Doxy10-100ng/mL **(C)**. Following 48 hrs of treatment, cells were seeded on fibronectin in NDiff227 and exposed to ActvinA/FGF2 and different combination of DMSO, Doxy and TMP for 4 days before being collected for RNA extraction. After 2 days, the media was changed, and the drugs were refreshed. **D, E** Fgf5 mRNA levels measured in FBS/L **(D)** and 2i/L **(E)** C1 ESCs, after 4 days of differentiation in NDiff227+ActivinA/FGF2 and different combination of DMSO, Doxy and TMP. Data are means±SEM (n=7, D (T0, Doxy10-100ng/mL "During Differentiation", Doxy10ng/mL T0, Doxy100ng/mL T0, Doxy100ng/mL "Before" ad Doxy100ng/mL "Always"); n=6, D (Doxy10ng/mL "Before"); n=5, D (TMP10μM "During

Differentiation"); n=2, E). p-values from two-tailed unpaired t test computed by comparing each sample with C1 TMP10µM "During Differentiation" are shown, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. Dots represent individual data with similar values overlapping.

Chemical modulation of the canonical Wnt pathway enhances Epiblast derivation from ESCs

The above results (Figure 1) and the need to define a protocol for efficient derivation of EpiSCs *in vitro* motivated us to explore the differentiation potential of wild-type ESCs when deprived of pluripotency factors and exposed to different chemical perturbations of the Wnt/β-catenin pathway (Figure 2A). We took advantage of the *miR-290-mCherry/miR-302-eGFP[53]* ESC line (hereafter called dual reporter ESCs), which allows fluorescent tracking of the exit from pluripotency. Specifically, naïve dual reporter ESCs express the mCherry reporter only, and progressively start expressing also the GFP reporter when allowed to differentiate into EpiSCs[53].

We measured the transition from ESCs to EpiSCs using flow cytometry in at least 2 independent 4-day time-courses (Figure 2A). Before differentiation, dual reporter ESCs were cultured in FBS/L and treated for 48 hrs with DMSO (Figure 2B) or with Chiron (1-3µM, Figure 2C, D respectively) to pre-activate the canonical Wnt pathway, or were maintained in 2i/L for 3 passages (i.e., 1 week; Figure 2E). At Day 0, 1.5×10^4 cells/cm^2 were seeded on fibronectin-coated plates and exposed to different combination of drugs added to the NDiff227[26]: ActivinA+FGF2+DMSO (AFD); ActivinA+FGF2+Ch1µM (AFC1); ActivinA+FGF2+Ch3µM (AFC3); Ch1µM (C1); Ch1µM+XAV2µM (C1X2); Ch3µM (C3); Ch3µM+XAV2µM[29] (C3X2) (Figure 2A). The GFP signal was analyzed using flow cytometry every day for 4 days and the media changed after the first 2 culture days (Figure 2A).

Dual reporter ESCs pre-cultured in FBS/L in absence of β-catenin activation showed, already at Day 1, a GFP+ cell percentage increase in all but AFD treatment; AFD-treated cells efficiently transited towards the Epi state later in the experiment, but they were unable to rich other treatment GFP+ percentage levels (Figure 2B).

FBS/L Ch1µM pre-cultured dual reporter ESCs showed an overall increased percentage of GFP+ cells compared to DMSO-treated ESCs (compare Figure 2B, C). However, the fold-increase at Day 4 with respect to Day 0 of the best performing conditions (i.e., C1 and C1X2; Figure 2C) was lower in Ch1µM pre-treated cells (13- and 12-fold increase, respectively; Figure 2C) in comparison with the DMSO (23- and

22-fold increase, respectively; Figure 2B). This suggests, as in the experiments
performed in a β-catenin null background (Figure 1), that the ESC to EpiSC conversion
is favoured by β-catenin activation during the differentiation only, while its prolonged
stimulation might reduce the differentiation efficiency. This was confirmed in Ch3μM
pre-treated cells, where the best performing conditions (i.e., C3 and C3X2; Figure 2D)
also showed an overall lower percentage of GFP+ cells as compared to DMSO-treated
ESCs. Moreover, the simultaneous activation/inhibition of the pathway (i.e., C3X2
condition) had opposite effects given cells pre-culture conditions: the percentage of
GFP+ cells increased over-time in DMSO pre-treated cells (20-fold increase at Day 4
with respect to Day 0; Figure 2B), while GFP expression was quite low at the end of
the time-course in Ch3μM pre-treated ESCs (3.6-fold increase at Day 4 with respect
to Day 0; Figure 2D). Interestingly, Chiron pre-treatment had no positive effect on the
differentiation efficiency using the standard differentiation protocol (AFD), as the GFP+
percentage increase in AFD-treated cells at Day 4, as compared to Day 0, was almost
the same in Ch1μM pre-cultured cells (7.4-fold increase; Figure 2C) and lower in
Ch3μM (1.9-fold increase; Figure 2D) and 2i/L (1.1-fold increase; Figure 2E) pre-
cultured cells as compared to the DMSO condition (7.4-fold increase; Figure 2B). In
ESCs pre-cultured in 2i/L, there was not much difference across the differentiation
protocols, as can be seen by the overall higher p-values computed through a one-way
ANOVA test (compare Figure 2B-E). Finally, the combined treatment with ActivinA,
FGF2 and Chiron (AFC1 and AFC3; Figure 2B-E) did not change EpiSCs derivation
efficiency, as compared to the standard ActivinA/FGF2 treatment (Figure 2B-E).
The ESC-EpiSC transition was proven to be a highly heterogeneous process involving
massive cell death[54]. Although we did not observe significant changes in cell viability
across experiments, we noticed strong variability in the efficiency of *in vitro* EpiSCs
derivation: in Supplementary Figure 2, we show the average of at least 2 experiments
where the overall percentage of GFP+ cells was almost halved across conditions, as
compared to data in Figure 2. Still, results in Supplementary Figure 2 confirm that the
addition of Chiron during differentiation, in combination or not with XAV2μM (C1 and
C1X2 or C3 and C3X2; Supplementary Figure 2A-D), enables a much more efficient
conversion towards EpiSCs as compared to the other protocols. Also, we confirmed
that, if pre-cultured in FBS/L, no pre-activation of the Wnt/β-catenin pathway and
differentiation in media enriched with Ch1μM or Ch1μM±XAV2μM enables efficient

EpiSCs derivation (C1 and C1X2; Supplementary Figure 2A), in agreement with β-

catenin genetic perturbation results in Figure 1.

Figure 2. Chemical perturbation of the Wnt/β-catenin in EpiSC derivation *in vitro*.

255 **A** Experimental scheme of Epiblast differentiation. Dual reporter ESCs cultured in FBS/L and

256 pre-treated for 48 hrs with DMSO and Chiron (1-3 μM), or in 2i/L for 3 passages, were seeded

on fibronectin in NDiff227 supplemented with different combination of drugs (ActivinA+FGF2+DMSO (AFD); ActivinA+FGF2+Ch1μM (AFC1); ActivinA+FGF2+Ch3μM (AFC3); Ch1μM (C1); Ch1μM+XAV2μM (C1X2); Ch3μM (C3); Ch3μM+XAV2μM (C3X2)). The GFP signal from the EpiSC marker was measured by flow cytometry every 24 hrs for 4 consecutive days. After the first 2 days, the media was changed, and the drugs were refreshed. **B-E** Percentage of GFP+ cells calculated over the total amount of living cells in DMSO **(B)**, Ch1μM **(C)**, Ch3μM **(D)** and 2i/L **(E)** pre-cultured dual reporter ESCs. Histograms from Day 0 and Day 4 of each condition are shown as insets. Data are means±SEM (n=2, B (Day1, Day3 AFD, Day4 AFD), C (Day 1), D (Day1, Day3 AFC3), E (Day0, Day1 AFC3-C3X2); n=3, B (Day0, Day2, Day3 AFC1-AFC3-C1-C1X2-C3-C3X2, Day4 AFC1-AFC3-C1-C1X2-C3-C3X2), C (Day0, Day2 AFD-AFC1-C1X2, Day3, Day4), D (Day0, Day2 AFD-AFC3-C3X2, Day3 AFD-C3-C3X2, Day4), E (Day1 AFD, Day2 AFD-AFC3-C3X2, Day3, Day4); n=6, C (Day2 C1), D (Day2 C3), E (Day1 C3, Day2 C3). p-values from one-way ANOVA are shown across samples for each day. Dots represent individual data with similar values overlapping. Colour-blind safe combinations were selected using colorbrewer2 (https://colorbrewer2.org/#type=sequential&scheme=BuGn&n=3).

To confirm these results at single cell level, we monitored single-cell ESC-EpiSC transition in 71 hrs time-lapse microscopy experiments (Supplementary Movies 1-3; see Methods for details). FBS/L pre-cultured dual reporter ESCs were imaged every 60 minutes while constantly stimulated with AFD (Supplementary Movie 1) or Chiron 1-3μM (Supplementary Movies 2 and 3, respectively). In the AFD condition, a few GFP+ cells started to appear after 57 hrs (Supplementary Movie 1), whereas in the C1 and C3 culture conditions the transition started already after 10-14 hrs, with a peak around 40-43 hrs (Supplementary Movies 2 and 3). The GFP signal from cells differentiated in the presence of Chiron3μM was weaker as compared to the Chiron1μM treatment (compare Supplementary Movies 2 and 3).

These results confirm that strength and time of β-catenin stabilization are important to tip the balance between cell fates. In agreement with our previous observations in FBS/L pre-cultured knock-out cells (see TMP10μM_Doxy10ng/ml "During Differentiation" sample in Figure 1D), data collected using the dual reporter cell line confirm that high levels of β-catenin can be detrimental to efficiently derive EpiSCs *in vitro*. Indeed, a moderate activation or the simultaneous treatment with both an activator (i.e., Chiron) and an inhibitor (i.e., XAV) of the Wnt/β-catenin pathway

improve ESC direct differentiation into EpiSC, effectively replacing ActivinA and FGF2
requirement (Figure 2B, Supplementary Figure 2A).

Transcriptome and WGCNA analysis of ESC exit from pluripotency with varying
β-catenin doses

Next, we studied C1 ESCs exit from the 'ground state' of pluripotency, i.e., upon 2i/L
withdrawal, by RNA-sequencing; such monolayer differentiation protocol does not
induce a specific cell fate and is ideal to observe any β-catenin dependent
differentiation bias.

C1 ESCs cultured in 2i/L for 3 passages were treated for 48 hrs with saturating
concentrations of Doxy and TMP (100ng/mL and 10μM, respectively) to induce the
expression of the exogenous fusion protein (Figure 3A). Taking advantage of the
mCherry-tag for exogenous β-catenin overexpression, Doxy/TMP treated C1 ESCs
were sorted into two different subpopulations: C1 with Middle and High β-catenin
levels (hereafter called C1M and C1H samples, respectively; Figure 3A and
Supplementary Figure 3A). Cells were then seeded at 1.5×10^4 cells/cm^2 on gelatin-
coated plates and cultured in NDiff227 media±inducers for 4 days before being
transcriptionally profiled (see Methods for details; Figure 3A). Sequencing provided
snapshots of the C1, C1M and C1H samples transcriptome in pluripotent condition,
and of the differentiated counterparts (Day 4 samples) cultured in NDiff227 and DMSO
(i.e., upon Doxy/TMP withdrawal; hereafter called C1DMV and C1DHV), or in NDiff227
and TMP/Doxy (hereafter called C1DMDT and C1DHDT) during differentiation. The
C1 and C1D samples refer to cells treated only with TMP10μM in the pluripotent and
differentiated states, respectively.

Principal Components Analysis (PCA, Figure 3B) showed three main clusters: one
group included C1, C1M and C1H in the 'ground state' of pluripotency, another group
included C1D ESCs and the final group contained all perturbed samples (C1DMV,
C1DHV, C1DMDT and C1DHDT) after 4 days of differentiation.

We next performed the Weighted Gene Correlation Network Analysis (WGCNA) to
pinpoint the processes associated with β-catenin perturbations. The WGCNA[70] is a
computational approach that allows the identification of gene modules based on their
correlation and network-derived topological properties. We applied the WGCNA to our

sequencing data and identified 13 different modules (Supplementary Figure 3B; see
Methods for details). We then defined the principal component of each module (i.e.,
eigenmodule, ME) and calculated the Pearson Correlation Coefficient (r) of each
eigenmodule with time (Supplementary Figure 3C) or β-catenin doses (Figure 3C).
Next, for each module, we performed a functional annotation analysis on the genes
with a module membership |kME| ≥ 0.8 (Methods) to identify the biological processes
and pathways enriched (Supplementary Table 1), focusing the attention on the
modules having $r > 0.5$ and p-value < 0.05 with the examined traits (Figure 3C,
Supplementary Figure 3C, Methods).
We found four modules significantly correlating (i.e., Green r = 0.92/p-value = 9e-14,
Blue r = 0.82/p-value = 1e-08 , Black r = 0.67/p-value = 3e-05 and Brown r =0.57/p-
value = 7e-04; Supplementary Figure 3C, Supplementary Table 1) and three anti-
correlating (i.e., Turquoise r = -0.99/p-value = 2e-24, yellow r = -0.75/p-value = 1e-06
and Pink r = -0.56/p-value = 8e-04; Supplementary Figure 3C, Supplementary Table
1) with time, meaning that these modules were regulated during the differentiation.
The biological processes (list of genes with a kME > 0.8; Supplementary Table 1)
related to these modules with a FDR < 0.05 showed an enrichment of genes involved
in translation, rRNA processing, ribosomal biogenesis and stem cell division
(Supplementary Figure 3D, F Supplementary Table 1, Green and Brown), protein
transport and processes associated with cellular respiration (Supplementary Figure
3E, Supplementary Table 1, Blue), positive regulation of growth, ncRNA processing
and neuronal tube formation and development (Supplementary Figure 3F,
Supplementary Table 1, Brown), regulation of tissue remodelling, embryonic and
forebrain development, stem cell population maintenance and cell homeostasis
(Supplementary Figure 3G, Supplementary Table 1, Turquoise), nuclear division and
meiosis and organelle fission (Supplementary Figure 3H, Supplementary Table 1,
Pink). For the pathway analysis, we found an enrichment in ribosome (Supplementary
Table 1, Green and Blue), Hedgehog signalling pathway (Supplementary Table 1,
Green), oxidative phosphorylation and diseases associated with oxidative stress
(Supplementary Table 1, Blue and Turquoise), cancer (Supplementary Table 1, Brown
and Turquoise) and regulation of the actin cytoskeleton (Supplementary Table 1,
Yellow). Of note, there were no significantly enriched BPs and/or pathways for the
genes from the Black, Yellow and Pink modules (Supplementary Table 1, Black,
Yellow and Pink).

To gain more information about the effects of β-catenin at the exit from pluripotency,

we correlated the eigenmodule of each module with β-catenin doses (Figure 3C, see

Methods for details) and found three modules correlating with β-catenin doses (i.e.,

Tan r = 0.81/p-value = 2e-8, Purple r = 0.71/p-value = 7e-6 and Yellow/Green r =

0.69/p-value =1 e-5; Figure 3C, Supplementary Table 1). The biological processes (list

of genes with a kME > 0.8; Supplementary Table 1) corresponding to these modules

with a FDR < 0.05 showed enrichment in cell division (Figure 3D, Supplementary

Table 1, Tan), metabolism and negative regulation of neuronal death (Figure 3E,

Supplementary Table 1, Purple). Finally, the pathways analysis showed a significant

enrichment only in Lysosome and N-Glycan biosynthesis from the Purple module,

whereas pentose phosphate pathways and autophagy from the Green/Yellow

(Supplementary Table 1, Purple and GreenYellow).

Overall, the WGCNA analysis confirmed the expected major transcriptional and

metabolic changes associated with the exit from the pluripotent status, and confirmed

previously reported β-catenin functions on cell survival and proliferation[31].

A

B

C

D

E

Figure 3. Transcriptome analysis of monolayer differentiation experiments upon β-catenin perturbations and WGCNA of the genes correlating with β-catenin doses.

A Experimental scheme of the monolayer differentiation protocol. 2i/L C1 ESCs were pre-treated with TMP10μM and Doxy100ng/mL for 48 hrs; living cells were then sorted from the Dapi negative fraction of TMP-treated cells (C1), whereas β-catenin overexpressing cells from Doxy/TMP-treated samples were FACS-sorted from the mCherry fraction and divided into Middle (C1M) and High (C1H) subpopulations. 1.5×10^4 cells/cm^2 from each individual population were then seeded on gelatin in NDiff227 supplemented with DMSO or TMP±Doxy100ng/mL. After 4 days of differentiation in NDiff227, cells were collected and processed for the RNA sequencing. During the protocol, the media was changed, and the drugs refreshed after 2 culture days. **B** Principal Component Analysis (PCA) of all samples; the average of biological replica is shown. **C** Eigenmodules correlating with β-catenin doses; the Pearson Correlation Coefficient (*r*) and relative p-value are shown. **D, E** Bar-chart of the top-ten enriched biological processes (BP) with FDR < 0.05 from genes belonging to the Tan **(D)** and Purple **(E)** WGCNA modules.

Next, to look at comparisons between specific pairs of samples, we performed a Gene Ontology (GO) and Functional Annotation analysis on the differentially expressed

genes filtered for the false discovery rate (FDR < 0.05) and log fold change (logFC <

-2 or > 2; see Methods for details). To identify gene ontologies specific for each

experimental perturbation, we focused on the Biological Processes (BPs) enriched

only in one comparison (red bars in Figure 4A-D, Supplementary Figure 4A, B).

Starting from the pluripotent condition, when comparing C1M and C1H with C1 we

found that the first 10 biological processes with a FDR < 0.05 were mainly enriched in

cell cycles and metabolism (Supplementary Figure 4A, B, Supplementary Tables 2,

3). Interestingly, the genes exclusively upregulated in C1H compared to C1 were

related to tissue differentiation (i.e., eye morphogenesis and urogenital system

development) and DNA methylation involved in gamete generation (Supplementary

Figure 4B, Supplementary Table 3). Only a few signalling pathways were enriched in

C1M ESCs compared to the control cell line C1 (Supplementary Tables 2, 3). These

results, together with the PCA in Figure 3B, confirm previous observations about the

dispensable function β-catenin has in pluripotent culture conditions[45-47,49] and suggest

a bias towards differentiation in C1H ESCs (Supplementary Figure 4B, Supplementary

Table 3).

We then analysed the genes differentially expressed at Day 4 upon β-catenin

perturbation as compared to C1D ESCs. Upregulated genes mostly contributed to the

BPs significantly enriched (Figure 4A-D, Supplementary Tables 4-7), while

downregulated genes only contributed to enrich a few processes, namely general

metabolic processes (e.g., regulation of transporter and cation channel activity) and

mesenchymal to epithelial transition (Figure 4A-D, Supplementary Tables 4-7). Genes

exclusively upregulated in the C1DMV *vs* C1D comparison belonged to the mesoderm

lineage (i.e., cardiovascular system development; Figure 4A and Supplementary

Table 4), while, in the C1DMDT *vs* C1D comparison, upregulated genes were enriched

for the endoderm lineage (i.e., urogenital system; Figure 4B and Supplementary Table

5). Nevertheless, mesoderm and endoderm lineages were represented in both

comparisons. GO performed on the C1DHV and C1DHDT comparisons with C1D

showed only a few differences in the enriched BPs, that indeed did not define a bias

toward a specific lineage (Figure 4C, D, Supplementary Tables 6, 7).

These results suggest that the major changes in the differentiation program initiated

upon 2i/L withdrawal are induced by moderate β-catenin doses and are influenced by

the timing of protein overexpression. The pathway enrichment analysis showed the

upregulation of protein metabolism in C1DMV and C1DHV (Supplementary Tables 4,

6), MAPK signalling pathway (Supplementary Table 4) in C1DMV, and ECM-receptor interaction and PI3K-AKT signalling pathway in C1DHV (Supplementary Table 6).

To gauge insights into specific differentiation programs, we selected sets of markers for naïve and general pluripotency, early post-implantation epiblast, ectoderm, mesoderm, endoderm, germ cell and trophectoderm[51], and clustered our samples according to the expression of these gene sets.

Naïve pluripotency genes were downregulated during differentiation in all samples, indicating the successful exit of cells from pluripotency (Figure 4E). Pluripotent C1M and C1H samples clustered together (Figure 4E) although close to C1 ESCs, in agreement with previous observations (Figure 3B). ESCs differentiated in presence of DMSO (i.e., C1DMV and C1DHV; Figure 4E) clustered together, similarly to samples differentiated in presence of Doxy and TMP (i.e., C1DMDT and C1DHDT, Figure 4E); still, a large number of genes (e.g., Klf5, Tcl1, Klf2 and Nr0b1) showed a different pattern among C1D, C1DMV, and C1DHV, discriminating ESCs with different β-catenin doses (Figure 4E). These results support the hypothesis of a β-catenin-dependent effect on transcriptional changes.

The clustering across samples was similar for general pluripotency markers (Figure 4F). In accordance with previous reports about the transition to primed pluripotency[55-58], in the majority of differentiated samples Sox2 was downregulated and Utf1, Zfp281 and Lin28 were upregulated (Figure 4F). Moreover, almost all the genes were lower in C1DMV, C1DMDT, C1DHV and C1DHDT as compared to C1D (exceptions were Lin28a, higher in C1DHDT and Sox2, higher in C1DHV; Figure 4F), and showed different behaviours in DMSO- (i.e., C1DMV and C1DHV; Figure 4F) *vs* Doxy/TMP-treated samples (i.e., C1DMDT, C1DHDT; Figure 4F), confirming that the duration of β-catenin overexpression has an influence on cell identity. Zfp281, Zic2 and Utf1 showed a similar pattern in β-catenin overexpressing cells as compared to C1D ESCs, all being downregulated (Figure 4F). Zfp281 is a Zinc finger transcription factor implicated in regulating stem cell pluripotency[59,60], and recently reported as a bidirectional regulator of the ESC-EpiSC transition in cooperation with Zic2, another zinc finger protein[61]. The undifferentiated embryonic cell transcription factor 1 (Utf1) is expressed in ESCs and is involved in many aspects of gene expression control (e.g., transcription factor[62] and epigenome organization[63,64]), playing an important regulatory role during the exit of ESCs from pluripotency[63,64]. The concomitant

reduction of Zfp281, Zic2 and Urf1 suggests a global change in the chromatin
organization of β-catenin overexpressing ESCs en route to differentiation (Figure 4F).
Early post-implantation epiblast genes were mostly upregulated in primed ESCs
compared to the pluripotent condition with no evident differences across treatments in
naïve ESCs (Figure 4G). The exception was Foxd3, downregulated in both naïve and
primed β-catenin overexpressing cells as compared to the controls C1 and C1D ESCs
(Figure 4G). Interestingly, Dnmt3a levels, although similar at T0, were very different at
Day 4, with C1DMV/C1DHV and C1DMDT/C1DHDT showing 80% and 70% gene
expression reduction as compared to the control C1D, respectively (Figure 4G); also,
samples constantly exposed to Doxy/TMP (i.e., C1DMDT and C1DHDT) showed
higher Dnmt3a expression than DMSO-treated ESCs (i.e., C1DMV and C1DHV;
Figure 4G). Similar observations hold for Dnmt3b, where the reduction compared to
C1D was of about 60% for C1DMV/C1DHV and 10% for C1DMDT/C1DHDT (Figure
4G). Dnmt3a, b and Foxd3 are DNA and chromatin remodelling factors, respectively;
Dnmt enzymes methylate genomic regions, whereas Foxd3 reduces active and
enhances inactive histone marks by recruiting the Lysine-specific demethylase 1
(Lsd1)[65]. The reduced expression of those genes in β-catenin overexpressing cells,
including the pluripotent markers Utf1 discussed above, supports the hypothesis of
differentially methylated DNA.
We then screened for a large panel of lineage-priming factors. Clustering based on
the ectoderm lineage genes showed that pluripotent C1 and C1M are more similar
than C1H (Figure 4H); following 2i/L withdrawal, the clustering resembled those of
previous sets (Figure 4E-G), with C1DMV/C1DHV and C1DMDT/C1DHDT grouping
together (Figure 4H). Genes from this lineage had different and sometimes opposite
expression across samples, making difficult to identify a clear pattern associated with
β-catenin perturbations. Nevertheless, we distinguished two groups of genes: one
group (i.e., Dlx3, Pou3f2, Otx1 and Pou3f3) was mainly downregulated during
differentiation, whereas the other group (i.e., Nes, Ascl1, Cdh2 and Pax6) had the
opposite trend (Figure 4H).
When looking at mesoderm markers (Figure 4I), differentiated samples clustered
similarly to the previous data set; however, C1DMV/C1DHV were more similar to C1D
than in the previous comparisons (Figure 4E-H) and genes showed different
expression patterns across conditions. Lhx1, Lefty1/2, Meox1, Hoxb1 and Bmp4 were
mainly upregulated upon differentiation and no major differences were observed

across samples in pluripotent conditions. The exception in this group was Lhx1 that was higher in β-catenin overexpressing cells depending on the timing of overexpression (compare C1DMV/C1DHV with C1DMDT/C1DHDT in Figure 4I). In contrast, genes such as Nodal, Kdr, Mixl1, Gsc, Foxf1 and Zic1 got downregulated when exiting pluripotency and the levels across conditions were not significantly changing (Figure 4I). Of note, although the behaviour of individual genes is hard to interpret, C1D ESCs were very different from all differentiated samples with perturbations of β-catenin, stressing the relevance of the latter for mesoderm specification[46].

The endoderm lineage was the one most influenced by β-catenin perturbations: C1D cells were unable to induce the expression of endoderm-related genes (compare C1 and C1D in Figure 4J), whereas in all perturbed ESCs their expression increased over time. As previously observed (Figure 4E-I), samples clustered together based on the duration of β-catenin overexpression rather than on its dose (i.e., C1DMV/C1DHV and C1DMDT/C1DHDT, Figure 4J). Moreover, because of their impaired differentiation, C1D clustered together with pluripotent samples (Figure 4J). The C1DHDT showed the highest expression of the 50% of the endoderm-associated genes (namely, Cxcr4, Gata4 and Sox7) as compared to all other differentiated samples (i.e., C1D, C1DMV, C1DMDT and C1DHV). Slightly different was the behaviour of Cxcr4, that decreased over time; however, its expression is indicative of definitive endoderm differentiation[66] that most probably is not happening in the short protocol we applied. These observations, together with the incapability of C1D cells to embark differentiation toward this lineage, strongly support previous knowledge about the β-catenin requirement for endoderm organization[46,67].

In the analysis of the germ cell lineage markers, all genes showed a rather heterogeneous expression pattern across samples (Figure 4K). Pluripotent C1M and C1H clustered together and close to C1, and differentiated samples clustered based on the duration of β-catenin perturbation (i.e., C1DMV/C1DHV and C1DMDT/C1DHDT). Dazl, Prdm1 and Ddx4 showed a clearer pattern in pluripotent conditions, being upregulated in C1M and C1H compared to C1 ESCs (Figure 4K).

Finally, when looking at trophectoderm markers, clustering showed similarity of C1 and C1M, as for the ectoderm and endoderm lineages (Figure 4H, J, respectively). C1DMDT and C1DHDT were similar for the expression of trophectoderm genes, whereas C1DMV was part of a different branch clustering with C1D; C1DHV was more

closely related to C1DMDT and C1DHDT (Figure 4L). Moreover, 90% of trophectoderm genes got downregulated during differentiation (namely, Eomes, Elf3 and Cdx1) in all the conditions; the only exception was Cdx2 that was upregulated in C1D, C1MDV and C1DMDT in comparison with the corresponding T0 (Figure 4L). Eomes was recently reported to control the exit from pluripotency by acting on the chromatin status[68]; its behaviour in naïve C1M and C1H ESCs support the theory of a different chromatin conformation in pluripotent β-catenin overexpressing cells (Figure 4L).

Accounting for the fact that ectoderm is a default linage of the monolayer differentiation protocol we applied[69], sequencing results suggest that β-catenin overexpression in a knock-out background favours rescuing defects in differentiation towards endoderm more than mesoderm. Indeed, mesodermal genes were mostly downregulated when β-catenin was overexpressed, whereas endodermal genes were all upregulated as compared to the control (Figure 4I, J). Moreover, we observed that lineage differentiation was influenced by the duration of protein overexpression rather than the dose. According to that, there was a transition from mesoderm to endoderm following moderate but constant β-catenin overexpression (compare C1DMV and C1DMDT in Figure 4A, B). Nevertheless, endoderm was an enriched gene ontology in all considered comparisons (Figure 4A-D). Finally, the observed expression of pluripotency markers Zfp281, Zic2 and Utf1, the early post-implantation markers Dnmt3a-b and Foxd3 and the trophectoderm marker Eomes suggests a reorganization of the epigenome in naïve C1M and C1H ESCs and upon monolayer differentiation of C1DMV, C1DMDT, C1DHV and C1DHDT ESCs.

Figure 4. Gene ontology and clustergram of the differential expressed genes in control and perturbed ESCs.

A-D Bar-chart of the top-ten enriched biological processes (BP) with FDR < 0.05 from differentially expressed genes in C1DMV **(A),** C1DMDT **(B),** C1DHV **(C)** and C1DHDT **(D)** compared to C1D ESCs. Black and grey bars represent upregulated and downregulated BPs,

respectively. In red bars, the BPs exclusively enriched in the indicated condition. **E-L** Clustergram over heatmaps of Naïve **(E)** and general pluripotency **(H)**, early post-implantation **(G)**, ectoderm **(H)**, mesoderm **(I)**, endoderm **(J)**, germ cell **(K)** and trophectoderm **(L)** lineages from pluripotent and differentiated ESCs expressing different β-catenin amount. Each column is the average of 4 biological replicates.

Discussion

The role of the Wnt/β-catenin pathway as a pluripotency gatekeeper has been matter of many studies and debates[30,42,48,71-74]; while modulation of the canonical Wnt pathway has been extensively proved to be important for EpiSC *in vivo* derivation[75,76], self-renewal[77] and *in vitro* lineage differentiation[78-80], its relevance for the exit from pluripotency and for ESC-EpiSC direct transition have not been explored thoroughly. In this work, we proved that genetic β-catenin manipulation or chemical perturbation of the canonical Wnt pathway controls ESC fate at the exit from pluripotency.

Using two different cellular models, we found that moderate β-catenin doses in β-catenin$^{-/-}$ ESCs or its stabilization in wild-type ESCs increases the differentiation efficiency of ESCs into EpiSCs, similarly to what was reported for EpiSCs self-renewal[29]: EpiSCs can be maintained in Chiron3μM/XAV2μM cultures, with self-renewal regulated by both Axin2 and β-catenin[29]; here we found that, in absence of pre-activation of the canonical Wnt pathway, the Chiron3μM/XAV2μM treatment has a better effect than Chiron3μM alone on ESC-EpiSC transition at the end of the differentiation experiments (Day 3 and 4 in Figure 2B and Supplementary 2A), while the opposite was observed in the early phases of the protocol (Day 1 and 2 in Figure 2B and Supplementary 2A). This could be explained by the fact that Axin2 expression is β-catenin-dependent, and β-catenin stabilization gets induced by Chiron treatment; therefore, the effect of XAV might be relevant only upon Axin2 stabilisation. Alternatively, it might also suggest a switch in the effect of Chiron3μM/XAV2μM stimulation in a specific moment during the differentiation process, probably when ESCs start to acquire a more stable EpiSC identity (as in[29]). Although both hypotheses are in agreement with what was reported by Kim and colleagues[29], the first one seems to be more plausible looking at the response in Ffg5 (Figure 1) and GFP expression (Figure 2B, Supplementary Figure 2A), i.e., when a moderate activation of the Wnt/β-catenin pathway was induced. Indeed, both β-catenin$^{-/-}$ exposed to a low concentration of the input Doxy (i.e., TMP10 μM _Doxy10ng/mL, "During Differentiation" in Figure

1D) and the dual reporter ESCs treated with Chiron1µM (i.e., C1 in Figure 2B) showed

the highest differentiation efficiency, suggesting that Chiron1µM treatment alone is

enough to keep β-catenin doses within a range that would instead require XAV

treatment if using higher doses of Chiron (i.e., 3µM), and that Chiron1µM is mimicking

the effect obtained by β-catenin genetic manipulation. Furthermore, XAV treatment in

combination with Chiron1µM reduced the percentage of differentiated cells at Day 1

(Figure 2B and Supplementary 2A); this could be due to the stoichiometry between β-

catenin, Ainx2 and the two inhibitors. We reason that, in Chiron1µM/XAV2µM, XAV is

in excess with respect to the amount of transcribed Axin2, therefore all the translated

protein gest stabilised, and the final effect is a strong inhibition of the signalling

pathway. Nevertheless, this effect might get attenuated with time, probably when a

balance between the drug and the target is achieved. This theory seems to be

supported by the behaviour of cells treated with Chiron1µM or Chiron3µM; the first

showed a GFP peak within the first 2 days followed by GFP stabilisation, while the

second showed the opposite trend (i.e., GFP reduction over time), suggesting that

prolonged β-catenin stabilization in FBS/L can impair ESCs differentiation towards

EpiSCs. These results pose the question of what would have happened if cells were

stimulated for longer with Chiron1µM. There are two possible scenarios; in the first,

EpiSCs, derived from ESCs, would self-renew in Chiron1µM as well as in

Chiron3µM/XAV2µM. In the second scenario, the transition from Chiron1µM to Chiron

3µM would be preferred for the maintenance of established EpiSCs. However, the

latter scenario would be in contrast with the results showed in[29], where it was proved

immediate differentiation or death of EpiSCs expanded in Chiron3µM±PD1µM. Of

note, the basal medium used in[29] for EpiSC self-renewal is based on minimal essential

medium (GMEM) supplemented with 10% FBS[29], therefore differences might be

expected.   Finally, we found that 2i/L pre-cultured ESCs better differentiate into

EpiSCs, regardless of further modulation of the pathway during differentiation. It has

been shown that, during the differentiation process, ESCs downregulate Oct4

expression and undergo a temporary cell cycle arrest[54]; when differentiation has been

accomplished, Oct4 levels and cell cycle progression are restored to support the clonal

expansion of the EpiSC population. However, rare Oct4-/Cdx2+ cells might appear

during the initial phase of the differentiation process; this cellular fraction does not

respond to the cell cycle arrest, and therefore has a proliferative advantage[54]. The

simultaneous elimination of differentiating incompetent cells (i.e., Oct4-/Cdx2+) and

the re-established proliferation of differentiated cells (i.e., Oct4+/Cdx2-) ensure a homogenous population of EpiSCs[54].

We and others reported a reduced proliferation rate of ESCs stimulated with Chiron both in FBS/L[31] and 2i/L[13]. In agreement with these observations, we hypothesise that Chiron treatment might reduce the proliferation of Oct4-/Cdx2+ cells, establishing a balance between differentiating competent (i.e., Oct4-/Cdx2-) and incompetent (i.e., Oct4-/Cdx2+) cells. The 2i/L pre-culture might additionally homogenise the expression of various genes, including Oct4[81] and reduce the appearance of Cdx2+ cells.

Overall, these results confirm the effect β-catenin has on preparing cells to appropriately respond to the differentiation stimuli and suggest that both the duration and the dose of β-catenin overexpression control cell differentiation *in vitro*. Moreover, they enabled us to define an improved protocol for EpiSCs derivation *in vitro*, based on NDiff227 and Chiron1µM.

RNA sequencing performed in ESCs during the exit from the naïve 'ground state' of pluripotency[51] showed that, in β-catenin overexpressing cells (in particular in C1DMV), Dnmt3a and b had an expression pattern similar to the one observed in Rex1-high ESCs differentiated using a similar protocol[51], indicating that moderate β-catenin overexpression in naïve ESCs influences DNA methylation associated with the exit from pluripotency. Interestingly, we observed a significant upregulation of endodermal genes in β-catenin overexpressing cells, indicating a requirement of β-catenin for this specific fate. This phenotype was previously reported in the β-catenin null cell line generated by Lyashenko and colleagues[46], where the defect in endoderm lineage differentiation was rescued by overexpressing both wild-type or transcriptional incompetent β-catenin[46]; in contrast, mesoderm and ectoderm induction seemed to not require β-catenin[46]. With our approach that enables dose- and time-controlled β-catenin overexpression, we were able to define the amount of protein and the optimal window of overexpression to facilitate mesoderm (i.e., C1DMV; Figure 4A, Supplementary Table 4) or endoderm (i.e., C1DMDT; Figure 4B, Supplementary Table 5) differentiation, and confirmed that the ectoderm lineage is not affected by β-catenin loss and therefore is not influenced by its overexpression.

In the future, it will be of great interest to use our inducible system to interrogate the effect of more complex β-catenin dynamics on stem-cell identity and to further investigate the role of the β-catenin transcriptional activity in pluripotent and differentiated cells of both murine and human origin.

Authors Contributions

E.P. designed and performed experiments; M.F. and G.G. performed the WGNCA analysis; M.F. performed the GO; R.D.C. performed the Differential Expression analysis; E.P., and L.M. analysed data; E.P., M.F., R.D.C. and L.M. wrote the paper; D.d.B. supervised the bioinformatics analysis; L.M. supervised the entire project.

Acknowledgements

We thank Dr Robert Blelloch for the *miR-290-mCherry/miR-302-eGFP* ESC line; Dr Andre Hermann and Dr Lorena Sueiro Ballesteros (Flow Cytometry Facility, University of Bristol), Dr Mark Jepson and Alan Leard (Wolfson Imaging Facility, University of Bristol) and the Next Generation Sequencing Core (TIGEM, Naples) for their support. This work was funded by Medical Research Council (grant MR/N021444/1) to L.M., by the Engineering and Physical Sciences Research Council (grants EP/R041695/1 and EP/S01876X/1 to L.M.), EC funding H2020 (FET OPEN 766840-COSY-BIO) to L.M., BrisSynBio, a BBSRC/EPSRC Synthetic Biology Research Centre (BB/L01386X/1) to L.M, STAR-University of Naples Federico II grant to G.G. and Fondazione Telethon grant to D.d.B.

Methods

Cell line derivation

C1 cell lines were previously derived in[49] by a double lentiviral infection of β-catenin[-/-] ESCs[45] with the EF1a-rtTA (Neomycin) plasmid followed by the pLVX_TrE3G-DDmCherryβ-catenin[S33Y](Puromycin). Cells were selected with Neomycin after the first round and with Puromycin after the last infection. The dual reporter ESCs were a gift from Dr Blelloch[53].

ESCs were cultured on gelatin-coated dishes in Dulbecco's modified Eagle's medium (DMEM) supplemented with 15% fetal bovine serum (FBS, Sigma), 1 x nonessential amino acids, 1 x GlutaMax, 1 x2-mercaptoethanol and 1000 U/mL LIF (Peprotech). To note, that for the 2i/L culture, cells were kept for 3 passages (around 1week) in serum-free NDiff227-based media supplemented with 1000 U/mL LIF, 3μM of the GSK-3α/β inhibitor Chiron-99021 (Selleck, S1263) and 1μM of the MEK inhibitor PD0325901 (Selleck, S1036).

Epiblast Differentiation

For EpiSC derivation *in vitro*, FBS/L and 2i/L 1.5×10^4 cells/cm^2 were seeded on 10µg/mL Fibronectin-coated 12-well plates in NDiff227 (Takara, Y40002) and, according to the experiment, stimulated with DMSO, TMP10µM (Sigma, T7883), Doxy 10-100ng/mL (Sigma, D9891), human ActivinA 10ng/mL (Peptrotech, 120-14E), human FGF2 10ng/mL (Peptrotech, 100-18B) Chiron1-3µM and the XAV939 2µM (Sigma, 575545) for 4 days with the media and drugs refreshed after the first 2 culture days (Figures 1B, C and 2A).

Monolayer differentiation

Sorted C1, C1M and C1H ESCs were plated at 1.5×10^4 cells/cm^2 on gelatin-coated 12-well plates in plain NDiff227 and stimulated with DMSO or TMP10µM±Doxy10-100ng/mL for 4 days with the media and drugs refreshed after the first 2 culture days (Figure 3A).

Drugs pre-treatment

Some experimental conditions required pre-treatment of cells. For β-catenin overexpression in Figure 1C, C1 ESCs were stimulated for 48 hrs with TMP10µM and Doxy10-100ng/mL before the EpiSC differentiation, whereas for pre-activation of the canonical Wnt pathway in Figure 2A, C-E and Supplementary Figure 2B-D, dual reported ESCs were exposed to Chiron1-3µM (Figure 2C, Supplementary Figure 2B and Figure 2D, Supplementary Figure 2C, respectively) for 48 hrs or cultured for 3 passages in 2i/L (Figure 2E, Supplementary Figure 2D), before the differentiation.

Flow cytometry analysis

ESCs from a 12-well plate were washed with sterile Phosphate-Buffered Saline (PBS, Gibco), incubated with 80 µL of trypsin for 2–3′ at room temperature and collected with 120 µL of PBS 2% FBS containing DAPI as cell viability marker. Cell suspension was analysed using the BD LSR Fortessa and 10,000 living cells were recorded for each sample. The % of GFP positive cells was calculated over living cells, gated as DAPI negative, using the FlowJo V10 software.

Flow activated cell sorting (FACS)

ESCs were washed with sterile phosphate-buffered saline (PBS, Gibco), trypsinised for 2-3′ at room temperature and centrifuged at 1000 × g for 5′. Pelleted cells were resuspended in 500 µL of plain NDiff227 media supplemented with DAPI. The mCherry positive fraction was sorted from DAPI negative using the BD Influx high-speed 16-parameter fluorescence activated cell sorter.

qPCR

For quantitative PCR, the total RNA, extracted from cells using the RNeasy kit (Qiagen), was retrotranscribed (Thermo Fischer, RevertAid Reverse Transcriptase EP0441) and the cDNA used as template for each qPCR reaction in a 15 µL reaction volume. iTaq Universal SYBR Green Supermix (1725120, Bio-Rad) was used with the Qiagen Rotor-Gene System. To eliminate the contamination from genomic DNA, the RNeasy Plus Mini Kit (Qiagen, 74134) was used to purify the total RNA used for the RNA Sequencing. The primers used were: Actin-Fwd: ACGTTGACATCCGTAAAGACCT, Actin-Rev: GCAGTAATCTCCTTCTGCATCC; Fgf5-Fwd: AAAACCTGGTGCACCCTAGA, Fgf5-Rev: CATCACATTCCCGAATTAAGC).

Time-lapse experiments

1.5×10^4 cells were seeded on 10µg/mL fibronectin-coated µ-Slide 8 Well Glass Bottom (Ibidi, 80827) and imaged with a Leica DMI6000 inverted epifluorescence microscope equipped with the Photometrics Prime 95B sCMOS camera (1200x1200 11µm pixels, 8 ,12 bit or 16 bit, 70 fps full frame) and an environmental control chamber (Solent) for long- term temperature control and CO2 enrichment. The Adaptive Focus Control (AFC) ensures focus is maintained during multiple acquisition cycles. Images were acquired from two channels (phase contrast and green fluorescence) with a 20X objective every 60min for 71 hrs. ImageJ version: 2.0.0-rc-69/1.52p was used to improve the GFP signal, apply a Gaussian blur filter (Sigma radius 2) and combine channels.

QuantSeq 3' RNA sequencing library preparation.

Preparation of libraries was performed with a total of 100ng of RNA from each sample using QuantSeq 3'mRNA-Seq Library prep kit (Lexogen, Vienna, Austria) according to manufacturer's instructions. Total RNA was quantified using the Qubit 2.0

fluorimetric Assay (Thermo Fisher Scientific). Libraries were prepared from 100ng of

total RNA using the QuantSeq 3' mRNA-Seq Library Prep Kit FWD for Illumina

(Lexogen GmbH). Quality of libraries was assessed by using screen tape High

sensitivity DNA D1000 (Agilent Technologies). Libraries were sequenced on a

NovaSeq 6000 sequencing system using an S1, 100 cycles flow cell (Illumina Inc.).

Amplified fragmented cDNA of 300 bp in size were sequenced in single-end mode with

a read length of 100 bp.

Illumina novaSeq base call (BCL) files are converted in fastq file through bcl2fastq

[http://emea.support.illumina.com/content/dam/illuminasupport/documents/document

ation/software_documentation/bcl2fastq/bcl2fastq2-v2-20-software-guide-15051736-

03.pdf] (version v2.20.0.422).

QuantSeq 3' RNA sequencing data processing and analysis.

For analysis, sequence reads were trimmed using bbduk software

(https://jgi.doe.gov/data-and-tools/bbtools/bb-tools-user-guide/usage-guide/) (bbmap

suite 37.31) to remove adapter sequences, poly-A tails and low-quality end bases

(regions with average quality below 6). Alignment was performed with STAR 2.6.0a3[82]

on mm10 reference assembly obtained from cellRanger website

(https://support.10xgenomics.com/single-cell-gene-expression/software/release-

notes/build#mm10_3.0.0; Ensembl assembly release 93). Expression levels of genes

were determined with htseq-count[83] using Gencode/Ensembl gene model. We have

filtered out all genes having < 1 cpm in less than n_min samples and Perc MM reads

> 20% simultaneously. Differential expression analysis was performed using edgeR[84],

a statistical package based on generalized linear models, suitable for multifactorial

experiments. The threshold for statistical significance chosen was False Discovery

Rate (FDR) < 0.05 (GSE148879). The lists of differentially expressed genes (DEGs),

for each comparison, with a threshold of logFC > 2 for the induced and logFC < -2 for

the inhibited transcripts (Supplementary Tables 2-7) were used for the Functional

Annotation analysis.

The data were deposited in GEO with the accession number GSE148879.

Weighted Gene Correlation Network Analysis (WGCNA)

Quant-seq 3' mRNA data of 32 samples was used to construct a gene co-expression

network by applying Weighted Gene Correlation Network Analysis (WGCNA)[70] from

the WGCNA package in the R statistical environment version 3.6. Briefly, we first

computed the Pearson correlation coefficient among all pairs of expressed genes and

then an appropriate value of the soft-thresholding power (β=6) giving a scale-free

topology fitting index (R^2) ≥ 0.85 was selected to build the weighted adjacency matrix.

The weighted adjacency matrix was further transformed into a topological overlap

matrix (TOM)[85] and the resulting dissimilarity matrix used for hierarchical clustering.

Gene modules were finally identified by cutting the hierarchical dendrogram with the

dynamic tree cut algorithm from dynamicTreeCut package in R[86] statistical

environment with standard parameters, except for cutHeight we set equal to 0.25 and

deepSplit we set equal to 1. The value of deepSplit parameter was selected after

performing a cluster stability analysis. Briefly, for each possible value of deepSplit

parameter (i.e., 0, 1, 2, 3 or 4), modules were identified for both the full dataset and

50 resampled datasets. Then, the clustering solution obtained for the full dataset was

compared with each resampled solution by mean of Adjusted Rand Index (ARI)[87]. The

solution giving the highest average ARI was used for the clustering analysis as

described above. Finally, to identify which clusters were correlated with β-catenin

expression doses or differentiation time we correlated the first principal component of

each gene module (i.e., the eigenmodule) with the traits of interest. The eigenmodule

can be considered as a "signature" of the module gene expression. Modules correlated

with the traits with a p-value < 0.01 were considered statistically significant and used

for further analyses.

Functional Annotation Analysis

Differentially expressed genes (either logFC > 2 or logFC < -2) and module "hubs"

having high module membership (also known as |KME| > 0.8) within the module were

analysed for the enrichment in GO Biological Processes[88] and KEGG Pathways[89] via

the clusterProfiler package in R statistical environment[90]. The threshold for statistical

significance was FDR < 0.05, the top-ten BPs were represented as -log10 (FDR;

Figure 3D-E, Supplementary Figure 3D-H, and Figure 4A-D, Supplementary Figure

4A, B).

Statistical analysis

Differences between samples were analysed by two-tailed unpaired t tstest and one-way ANOVA using Matlab (MathworksMatlab R2019a, update 9.6.0.1307630). A p-value lower than 0.05 was considered statistically significant.

Clustergram over heatmaps were generated using the clustergram function in Matlab (MathworksMatlab R2019a, update 9.6.0.1307630) that applies the Euclidean distance metric and average linkage. The data have been standardized across all samples for each gene and have 0 as mean and 1 as standard deviation.

References

Evans, M. J. & Kaufman, M. H. Establishment in culture of pluripotential cells from mouse embryos. *Nature* **292**, 154-156, doi:10.1038/292154a0 (1981).

Martin, G. R. Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *Proc Natl Acad Sci U S A* **78**, 7634-7638, doi:10.1073/pnas.78.12.7634 (1981).

Ying, Q. L., Wray, J., Nichols, J., Batlle-Morera, L., Doble, B., Woodgett, J., Cohen, P. & Smith, A. The ground state of embryonic stem cell self-renewal. *Nature* **453**, 519-523, doi:10.1038/nature06968 (2008).

Brook, F. A. & Gardner, R. L. The origin and efficient derivation of embryonic stem cells in the mouse. *Proc Natl Acad Sci U S A* **94**, 5709-5712, doi:10.1073/pnas.94.11.5709 (1997).

Williams, R. L., Hilton, D. J., Pease, S., Willson, T. A., Stewart, C. L., Gearing, D. P., Wagner, E. F., Metcalf, D., Nicola, N. A. & Gough, N. M. Myeloid leukaemia inhibitory factor maintains the developmental potential of embryonic stem cells. *Nature* **336**, 684-687, doi:10.1038/336684a0 (1988).

Smith, A. G., Heath, J. K., Donaldson, D. D., Wong, G. G., Moreau, J., Stahl, M. & Rogers, D. Inhibition of pluripotential embryonic stem cell differentiation by purified polypeptides. *Nature* **336**, 688-690, doi:10.1038/336688a0 (1988).

Ying, Q. L., Nichols, J., Chambers, I. & Smith, A. BMP induction of Id proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3. *Cell* **115**, 281-292, doi:10.1016/s0092-8674(03)00847-x (2003).

8   Niwa, H., Burdon, T., Chambers, I. & Smith, A. Self-renewal of pluripotent
embryonic stem cells is mediated via activation of STAT3. *Genes Dev* **12**, 2048-
2060, doi:10.1101/gad.12.13.2048 (1998).

9   Matsuda, T., Nakamura, T., Nakao, K., Arai, T., Katsuki, M., Heike, T. & Yokota,
T. STAT3 activation is sufficient to maintain an undifferentiated state of mouse
embryonic stem cells. *EMBO J* **18**, 4261-4269, doi:10.1093/emboj/18.15.4261
(1999).

10   Marucci, L., Pedone, E., Di Vicino, U., Sanuy-Escribano, B., Isalan, M. &
Cosma, M. P. beta-catenin fluctuates in mouse ESCs and is essential for
Nanog-mediated reprogramming of somatic cells to pluripotency. *Cell Rep* **8**,
1686-1696, doi:10.1016/j.celrep.2014.08.011 (2014).

11   Boroviak, T., Loos, R., Lombard, P., Okahara, J., Behr, R., Sasaki, E., Nichols,
J., Smith, A. & Bertone, P. Lineage-Specific Profiling Delineates the Emergence
and Progression of Naive Pluripotency in Mammalian Embryogenesis. *Dev Cell*
35, 366-382, doi:10.1016/j.devcel.2015.10.011 (2015).

12   Marks, H., Kalkan, T., Menafra, R., Denissov, S., Jones, K., Hofemeister, H.,
Nichols, J., Kranz, A., Stewart, A. F., Smith, A. & Stunnenberg, H. G. The
transcriptional and epigenomic foundations of ground state pluripotency. *Cell*
149, 590-604, doi:10.1016/j.cell.2012.03.026 (2012).

13   Godwin, S., Ward, D., Pedone, E., Homer, M., Fletcher, A. G. & Marucci, L. An
extended model for culture-dependent heterogenous gene expression and
proliferation dynamics in mouse embryonic stem cells. *NPJ Syst Biol Appl* **3**,
19, doi:10.1038/s41540-017-0020-5 (2017).

14   Ghimire, S., Van der Jeught, M., Neupane, J., Roost, M. S., Anckaert, J.,
Popovic, M., Van Nieuwerburgh, F., Mestdagh, P., Vandesompele, J., Deforce,
D., Menten, B., Chuva de Sousa Lopes, S., De Sutter, P. & Heindryckx, B.
Comparative analysis of naive, primed and ground state pluripotency in mouse
embryonic stem cells originating from the same genetic background. *Sci Rep*
8, 5884, doi:10.1038/s41598-018-24051-5 (2018).

15   Ficz, G., Hore, T. A., Santos, F., Lee, H. J., Dean, W., Arand, J., Krueger, F.,
Oxley, D., Paul, Y. L., Walter, J., Cook, S. J., Andrews, S., Branco, M. R. &
Reik, W. FGF signaling inhibition in ESCs drives rapid genome-wide
demethylation to the epigenetic ground state of pluripotency. *Cell Stem Cell* **13**,
351-359, doi:10.1016/j.stem.2013.06.004 (2013).

16 Habibi, E., Brinkman, A. B., Arand, J., Kroeze, L. I., Kerstens, H. H., Matarese,
F., Lepikhov, K., Gut, M., Brun-Heath, I., Hubner, N. C., Benedetti, R., Altucci,
L., Jansen, J. H., Walter, J., Gut, I. G., Marks, H. & Stunnenberg, H. G. Whole-
genome bisulfite sequencing of two distinct interconvertible DNA methylomes
of mouse embryonic stem cells. *Cell Stem Cell* **13**, 360-369,
doi:10.1016/j.stem.2013.06.002 (2013).

17 Leitch, H. G., McEwen, K. R., Turp, A., Encheva, V., Carroll, T., Grabole, N.,
Mansfield, W., Nashun, B., Knezovich, J. G., Smith, A., Surani, M. A. & Hajkova,
P. Naive pluripotency is associated with global DNA hypomethylation. *Nat*
Struct Mol Biol **20**, 311-316, doi:10.1038/nsmb.2510 (2013).

18 Nichols, J. & Smith, A. Pluripotency in the embryo and in culture. *Cold Spring*
Harb Perspect Biol **4**, a008128, doi:10.1101/cshperspect.a008128 (2012).

19 Alexandrova, S., Kalkan, T., Humphreys, P., Riddell, A., Scognamiglio, R.,
Trumpp, A. & Nichols, J. Selection and dynamics of embryonic stem cell
integration into early mouse embryos. *Development* **143**, 24-34,
doi:10.1242/dev.124602 (2016).

20 Tesar, P. J., Chenoweth, J. G., Brook, F. A., Davies, T. J., Evans, E. P., Mack,
D. L., Gardner, R. L. & McKay, R. D. New cell lines from mouse epiblast share
defining features with human embryonic stem cells. *Nature* **448**, 196-199,
doi:10.1038/nature05972 (2007).

21 Brons, I. G., Smithers, L. E., Trotter, M. W., Rugg-Gunn, P., Sun, B., Chuva de
Sousa Lopes, S. M., Howlett, S. K., Clarkson, A., Ahrlund-Richter, L.,
Pedersen, R. A. & Vallier, L. Derivation of pluripotent epiblast stem cells from
mammalian embryos. *Nature* **448**, 191-195, doi:10.1038/nature05950 (2007).

22 Han, D. W., Tapia, N., Joo, J. Y., Greber, B., Arauzo-Bravo, M. J., Bernemann,
C., Ko, K., Wu, G., Stehling, M., Do, J. T. & Scholer, H. R. Epiblast stem cell
subpopulations represent mouse embryos of distinct pregastrulation stages.
Cell **143**, 617-627, doi:10.1016/j.cell.2010.10.015 (2010).

23 Stavridis, M. P., Lunn, J. S., Collins, B. J. & Storey, K. G. A discrete period of
FGF-induced Erk1/2 signalling is required for vertebrate neural specification.
Development **134**, 2889-2894, doi:10.1242/dev.02858 (2007).

24 Kunath, T., Saba-El-Leil, M. K., Almousailleakh, M., Wray, J., Meloche, S. &
Smith, A. FGF stimulation of the Erk1/2 signalling cascade triggers transition of

pluripotent embryonic stem cells from self-renewal to lineage commitment.
Development **134**, 2895-2902, doi:10.1242/dev.02880 (2007).

25   Greber, B., Wu, G., Bernemann, C., Joo, J. Y., Han, D. W., Ko, K., Tapia, N.,
Sabour, D., Sterneckert, J., Tesar, P. & Scholer, H. R. Conserved and divergent
roles of FGF signaling in mouse epiblast stem cells and human embryonic stem
cells. *Cell Stem Cell* **6**, 215-226, doi:10.1016/j.stem.2010.01.003 (2010).

26   Guo, G., Yang, J., Nichols, J., Hall, J. S., Eyres, I., Mansfield, W. & Smith, A.
Klf4 reverts developmentally programmed restriction of ground state
pluripotency. *Development* **136**, 1063-1069, doi:10.1242/dev.030957 (2009).

27   Johansson, B. M. & Wiles, M. V. Evidence for involvement of activin A and bone
morphogenetic protein 4 in mammalian mesoderm and hematopoietic
development. *Mol Cell Biol* **15**, 141-151, doi:10.1128/mcb.15.1.141 (1995).

28   Yao, S., Chen, S., Clark, J., Hao, E., Beattie, G. M., Hayek, A. & Ding, S. Long-
term self-renewal and directed differentiation of human embryonic stem cells in
chemically defined conditions. *Proc Natl Acad Sci U S A* **103**, 6907-6912,
doi:10.1073/pnas.0602280103 (2006).

29   Kim, H., Wu, J., Ye, S., Tai, C. I., Zhou, X., Yan, H., Li, P., Pera, M. & Ying, Q.
L. Modulation of beta-catenin function maintains mouse epiblast stem cell and
human embryonic stem cell self-renewal. *Nat Commun* **4**, 2403,
doi:10.1038/ncomms3403 (2013).

30   Sato, N., Meijer, L., Skaltsounis, L., Greengard, P. & Brivanlou, A. H.
Maintenance of pluripotency in human and mouse embryonic stem cells
through activation of Wnt signaling by a pharmacological GSK-3-specific
inhibitor. *Nat Med* **10**, 55-63, doi:10.1038/nm979 (2004).

31   De Jaime-Soguero, A., Aulicino, F., Ertaylan, G., Griego, A., Cerrato, A.,
Tallam, A., Del Sol, A., Cosma, M. P. & Lluis, F. Wnt/Tcf1 pathway restricts
embryonic stem cell cycle through activation of the Ink4/Arf locus. *PLoS Genet*
13, e1006682, doi:10.1371/journal.pgen.1006682 (2017).

32   Stamos, J. L. & Weis, W. I. The beta-catenin destruction complex. *Cold Spring*
Harb Perspect Biol **5**, a007898, doi:10.1101/cshperspect.a007898 (2013).

33   Jho, E. H., Zhang, T., Domon, C., Joo, C. K., Freund, J. N. & Costantini, F.
Wnt/beta-catenin/Tcf signaling induces the transcription of Axin2, a negative
regulator of the signaling pathway. *Mol Cell Biol* **22**, 1172-1183,
doi:10.1128/mcb.22.4.1172-1183.2002 (2002).

34   Chia, I. V. & Costantini, F. Mouse axin and axin2/conductin proteins are
functionally equivalent in vivo. *Mol Cell Biol* **25**, 4371-4376,
doi:10.1128/MCB.25.11.4371-4376.2005 (2005).

35   Leung, J. Y., Kolligs, F. T., Wu, R., Zhai, Y., Kuick, R., Hanash, S., Cho, K. R.
& Fearon, E. R. Activation of AXIN2 expression by beta-catenin-T cell factor. A
feedback repressor pathway regulating Wnt signaling. *J Biol Chem* **277**, 21657-
21665, doi:10.1074/jbc.M200139200 (2002).

36   Glinka, A., Wu, W., Delius, H., Monaghan, A. P., Blumenstock, C. & Niehrs, C.
Dickkopf-1 is a member of a new family of secreted proteins and functions in
head induction. *Nature* **391**, 357-362, doi:10.1038/34848 (1998).

37   Pedone, E. & Marucci, L. Role of beta-Catenin Activation Levels and
Fluctuations in Controlling Cell Fate. *Genes (Basel)* **10**,
doi:10.3390/genes10020176 (2019).

38   Aulicino, F., Theka, I., Ombrato, L., Lluis, F. & Cosma, M. P. Temporal
perturbation of the Wnt signaling pathway in the control of cell reprogramming
is modulated by TCF1. *Stem Cell Reports* **2**, 707-720,
doi:10.1016/j.stemcr.2014.04.001 (2014).

39   Ho, R., Papp, B., Hoffman, J. A., Merrill, B. J. & Plath, K. Stage-specific
regulation of reprogramming to induced pluripotent stem cells by Wnt signaling
and T cell factor proteins. *Cell Rep* **3**, 2113-2126,
doi:10.1016/j.celrep.2013.05.015 (2013).

40   Kimura, M., Nakajima-Koyama, M., Lee, J. & Nishida, E. Transient Expression
of WNT2 Promotes Somatic Cell Reprogramming by Inducing beta-Catenin
Nuclear Accumulation. *Stem Cell Reports* **6**, 834-843,
doi:10.1016/j.stemcr.2016.04.012 (2016).

41   Lluis, F., Pedone, E., Pepe, S. & Cosma, M. P. Periodic activation of Wnt/beta-
catenin signaling enhances somatic cell reprogramming mediated by cell
fusion. *Cell Stem Cell* **3**, 493-507, doi:10.1016/j.stem.2008.08.017 (2008).

42   Anton, R., Kestler, H. A. & Kuhl, M. Beta-catenin signaling contributes to
stemness and regulates early differentiation in murine embryonic stem cells.
FEBS Lett **581**, 5247-5254, doi:10.1016/j.febslet.2007.10.012 (2007).

43   Soncin, F., Mohamet, L., Eckardt, D., Ritson, S., Eastham, A. M., Bobola, N.,
Russell, A., Davies, S., Kemler, R., Merry, C. L. & Ward, C. M. Abrogation of E-
cadherin-mediated cell-cell contact in mouse embryonic stem cells results in

reversible LIF-independent self-renewal. *Stem Cells* **27**, 2069-2080,
doi:10.1002/stem.134 (2009).

44 Wagner, R. T., Xu, X., Yi, F., Merrill, B. J. & Cooney, A. J. Canonical Wnt/beta-
catenin regulation of liver receptor homolog-1 mediates pluripotency gene
expression. *Stem Cells* **28**, 1794-1804, doi:10.1002/stem.502 (2010).

45 Aulicino, F., Sottile, F., Pedone, E., Lluis, F., Marucci, L. & Cosma, M. P.
Canonical Wnt pathway controls mESCs self-renewal through inhibition of
spontaneous differentiation via β-catenin/TCF/LEF functions. *bioRxiv*,
doi:10.1101/661777 (2019).

46 Lyashenko, N., Winter, M., Migliorini, D., Biechele, T., Moon, R. T. & Hartmann,
C. Differential requirement for the dual functions of beta-catenin in embryonic
stem cell self-renewal and germ layer formation. *Nat Cell Biol* **13**, 753-761,
doi:10.1038/ncb2260 (2011).

47 Wray, J., Kalkan, T., Gomez-Lopez, S., Eckardt, D., Cook, A., Kemler, R. &
Smith, A. Inhibition of glycogen synthase kinase-3 alleviates Tcf3 repression of
the pluripotency network and increases embryonic stem cell resistance to
differentiation. *Nat Cell Biol* **13**, 838-845, doi:10.1038/ncb2267 (2011).

48 Kielman, M. F., Rindapaa, M., Gaspar, C., van Poppel, N., Breukel, C., van
Leeuwen, S., Taketo, M. M., Roberts, S., Smits, R. & Fodde, R. Apc modulates
embryonic stem-cell differentiation by controlling the dosage of beta-catenin
signaling. *Nat Genet* **32**, 594-605, doi:10.1038/ng1045 (2002).

49 Pedone, E., Postiglione, L., Aulicino, F., Rocca, D. L., Montes-Olivas, S.,
Khazim, M., di Bernardo, D., Pia Cosma, M. & Marucci, L. A tunable dual-input
system for on-demand dynamic gene expression regulation. *Nat Commun* **10**,
4481, doi:10.1038/s41467-019-12329-9 (2019).

50 Sadot, E., Conacci-Sorrell, M., Zhurinsky, J., Shnizer, D., Lando, Z., Zharhary,
D., Kam, Z., Ben-Ze'ev, A. & Geiger, B. Regulation of S33/S37 phosphorylated
beta-catenin in normal and transformed cells. *J Cell Sci* **115**, 2771-2780 (2002).

51 Kalkan, T., Olova, N., Roode, M., Mulas, C., Lee, H. J., Nett, I., Marks, H.,
Walker, R., Stunnenberg, H. G., Lilley, K. S., Nichols, J., Reik, W., Bertone, P.
& Smith, A. Tracking the embryonic stem cell transition from ground state
pluripotency. *Development* **144**, 1221-1234, doi:10.1242/dev.142711 (2017).

52 Choi, J., Huebner, A. J., Clement, K., Walsh, R. M., Savol, A., Lin, K., Gu, H.,
Di Stefano, B., Brumbaugh, J., Kim, S. Y., Sharif, J., Rose, C. M., Mohammad,

A., Odajima, J., Charron, J., Shioda, T., Gnirke, A., Gygi, S., Koseki, H.,

Sadreyev, R. I., Xiao, A., Meissner, A. & Hochedlinger, K. Prolonged Mek1/2

suppression impairs the developmental potential of embryonic stem cells.

Nature **548**, 219-223, doi:10.1038/nature23274 (2017).

53 Parchem, R. J., Ye, J., Judson, R. L., LaRussa, M. F., Krishnakumar, R.,

Blelloch, A., Oldham, M. C. & Blelloch, R. Two miRNA clusters reveal

alternative paths in late-stage reprogramming. *Cell Stem Cell* **14**, 617-631,

doi:10.1016/j.stem.2014.01.021 (2014).

54 Turco, M. Y., Furia, L., Dietze, A., Fernandez Diaz, L., Ronzoni, S., Sciullo, A.,

Simeone, A., Constam, D., Faretta, M. & Lanfrancone, L. Cellular heterogeneity

during embryonic stem cell differentiation to epiblast stem cells is revealed by

the ShcD/RaLP adaptor protein. *Stem Cells* **30**, 2423-2436,

doi:10.1002/stem.1217 (2012).

55 Zhang, J., Ratanasirintrawoot, S., Chandrasekaran, S., Wu, Z., Ficarro, S. B.,

Yu, C., Ross, C. A., Cacchiarelli, D., Xia, Q., Seligson, M., Shinoda, G., Xie,

W., Cahan, P., Wang, L., Ng, S. C., Tintara, S., Trapnell, C., Onder, T., Loh, Y.

H., Mikkelsen, T., Sliz, P., Teitell, M. A., Asara, J. M., Marto, J. A., Li, H., Collins,

J. J. & Daley, G. Q. LIN28 Regulates Stem Cell Metabolism and Conversion to

Primed Pluripotency. *Cell Stem Cell* **19**, 66-80, doi:10.1016/j.stem.2016.05.009

(2016).

56 Fidalgo, M., Huang, X., Guallar, D., Sanchez-Priego, C., Valdes, V. J.,

Saunders, A., Ding, J., Wu, W. S., Clavel, C. & Wang, J. Zfp281 Coordinates

Opposing Functions of Tet1 and Tet2 in Pluripotent States. *Cell Stem Cell* **19**,

355-369, doi:10.1016/j.stem.2016.05.025 (2016).

57 Luo, Z., Gao, X., Lin, C., Smith, E. R., Marshall, S. A., Swanson, S. K., Florens,

L., Washburn, M. P. & Shilatifard, A. Zic2 is an enhancer-binding factor required

for embryonic stem cell specification. *Mol Cell* **57**, 685-694,

doi:10.1016/j.molcel.2015.01.007 (2015).

58 Betschinger, J., Nichols, J., Dietmann, S., Corrin, P. D., Paddison, P. J. &

Smith, A. Exit from pluripotency is gated by intracellular redistribution of the

bHLH transcription factor Tfe3. *Cell* **153**, 335-347,

doi:10.1016/j.cell.2013.03.012 (2013).

59 Brandenberger, R., Wei, H., Zhang, S., Lei, S., Murage, J., Fisk, G. J., Li, Y.,

Xu, C., Fang, R., Guegler, K., Rao, M. S., Mandalam, R., Lebkowski, J. &

Stanton, L. W. Transcriptome characterization elucidates signaling networks
that control human ES cell growth and differentiation. *Nat Biotechnol* **22**, 707-
716, doi:10.1038/nbt971 (2004).

60   Wang, Z. X., Teh, C. H., Chan, C. M., Chu, C., Rossbach, M., Kunarso, G.,
Allapitchay, T. B., Wong, K. Y. & Stanton, L. W. The transcription factor Zfp281
controls embryonic stem cell pluripotency by direct activation and repression of
target genes. *Stem Cells* **26**, 2791-2799, doi:10.1634/stemcells.2008-0443
(2008).

61   Mayer, D., Stadler, M. B., Rittirsch, M., Hess, D., Lukonin, I., Winzi, M., Smith,
A., Buchholz, F. & Betschinger, J. Zfp281 orchestrates interconversion of
pluripotent states by engaging Ehmt1 and Zic2. *EMBO J* **39**, e102591,
doi:10.15252/embj.2019102591 (2020).

62   Okuda, A., Fukushima, A., Nishimoto, M., Orimo, A., Yamagishi, T.,
Nabeshima, Y., Kuro-o, M., Nabeshima, Y., Boon, K., Keaveney, M.,
Stunnenberg, H. G. & Muramatsu, M. UTF1, a novel transcriptional coactivator
expressed in pluripotent embryonic stem cells and extra-embryonic cells.
EMBO J **17**, 2019-2032, doi:10.1093/emboj/17.7.2019 (1998).

63   Kooistra, S. M., Thummer, R. P. & Eggen, B. J. Characterization of human
UTF1, a chromatin-associated protein with repressor activity expressed in
pluripotent cells. *Stem Cell Res* **2**, 211-218, doi:10.1016/j.scr.2009.02.001
(2009).

64   Jia, J., Zheng, X., Hu, G., Cui, K., Zhang, J., Zhang, A., Jiang, H., Lu, B., Yates,
J., 3rd, Liu, C., Zhao, K. & Zheng, Y. Regulation of pluripotency and self-
renewal of ESCs through epigenetic-threshold modulation and mRNA pruning.
Cell **151**, 576-589, doi:10.1016/j.cell.2012.09.023 (2012).

65   Respuela, P., Nikolic, M., Tan, M., Frommolt, P., Zhao, Y., Wysocka, J. & Rada-
Iglesias, A. Foxd3 Promotes Exit from Naive Pluripotency through Enhancer
Decommissioning and Inhibits Germline Specification. *Cell Stem Cell* **18**, 118-
133, doi:10.1016/j.stem.2015.09.010 (2016).

66   Yasunaga, M., Tada, S., Torikai-Nishikawa, S., Nakano, Y., Okada, M., Jakt, L.
M., Nishikawa, S., Chiba, T., Era, T. & Nishikawa, S. Induction and monitoring
of definitive and visceral endoderm differentiation of mouse ES cells. *Nat*
Biotechnol **23**, 1542-1550, doi:10.1038/nbt1167 (2005).

67   Engert, S., Burtscher, I., Liao, W. P., Dulev, S., Schotta, G. & Lickert, H.
Wnt/beta-catenin signalling regulates Sox17 expression and is essential for
organizer and endoderm formation in the mouse. *Development* **140**, 3128-
3138, doi:10.1242/dev.088765 (2013).

68   Tosic, J., Kim, G. J., Pavlovic, M., Schroder, C. M., Mersiowsky, S. L., Barg,
M., Hofherr, A., Probst, S., Kottgen, M., Hein, L. & Arnold, S. J. Eomes and
Brachyury control pluripotency exit and germ-layer segregation by changing the
chromatin state. *Nat Cell Biol* **21**, 1518-1531, doi:10.1038/s41556-019-0423-1
(2019).

69   Ying, Q. L., Stavridis, M., Griffiths, D., Li, M. & Smith, A. Conversion of
embryonic stem cells into neuroectodermal precursors in adherent
monoculture. *Nat Biotechnol* **21**, 183-186, doi:10.1038/nbt780 (2003).

70   Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation
network analysis. *BMC Bioinformatics* **9**, 559, doi:10.1186/1471-2105-9-559
(2008).

71   Ogawa, K., Nishinakamura, R., Iwamatsu, Y., Shimosato, D. & Niwa, H.
Synergistic action of Wnt and LIF in maintaining pluripotency of mouse ES cells.
Biochem Biophys Res Commun **343**, 159-166, doi:10.1016/j.bbrc.2006.02.127
(2006).

72   Hao, J., Li, T. G., Qi, X., Zhao, D. F. & Zhao, G. Q. WNT/beta-catenin pathway
up-regulates Stat3 and converges on LIF to prevent differentiation of mouse
embryonic stem cells. *Dev Biol* **290**, 81-91, doi:10.1016/j.ydbio.2005.11.011
(2006).

73   Singla, D. K., Schneider, D. J., LeWinter, M. M. & Sobel, B. E. wnt3a but not
wnt11 supports self-renewal of embryonic stem cells. *Biochem Biophys Res*
Commun **345**, 789-795, doi:10.1016/j.bbrc.2006.04.125 (2006).

74   Takao, Y., Yokota, T. & Koide, H. Beta-catenin up-regulates Nanog expression
through interaction with Oct-3/4 in embryonic stem cells. *Biochem Biophys Res*
Commun **353**, 699-705, doi:10.1016/j.bbrc.2006.12.072 (2007).

75   Tsukiyama, T. & Ohinata, Y. A modified EpiSC culture condition containing a
GSK3 inhibitor can support germline-competent pluripotency in mice. *PLoS*
One **9**, e95329, doi:10.1371/journal.pone.0095329 (2014).

76   Sugimoto, M., Kondo, M., Koga, Y., Shiura, H., Ikeda, R., Hirose, M., Ogura,
A., Murakami, A., Yoshiki, A., Chuva de Sousa Lopes, S. M. & Abe, K. A simple

and robust method for establishing homogeneous mouse epiblast stem cell
lines by wnt inhibition. *Stem Cell Reports* **4**, 744-757,
doi:10.1016/j.stemcr.2015.02.014 (2015).

77 Sumi, T., Oki, S., Kitajima, K. & Meno, C. Epiblast ground state is controlled by
canonical Wnt/beta-catenin signaling in the postimplantation mouse embryo
and epiblast stem cells. *PLoS One* **8**, e63378,
doi:10.1371/journal.pone.0063378 (2013).

78 Liu, K., Sun, Y., Liu, D. & Ye, S. Inhibition of Wnt/beta-catenin signaling by
IWR1 induces expression of Foxd3 to promote mouse epiblast stem cell self-
renewal. *Biochem Biophys Res Commun* **490**, 616-622,
doi:10.1016/j.bbrc.2017.06.086 (2017).

79 Osteil, P., Studdert, J. B., Goh, H. N., Wilkie, E. E., Fan, X., Khoo, P. L., Peng,
G., Salehin, N., Knowles, H., Han, J. J., Jing, N., Fossat, N. & Tam, P. P. L.
Dynamics of Wnt activity on the acquisition of ectoderm potency in epiblast
stem cells. *Development* **146**, doi:10.1242/dev.172858 (2019).

80 Kurek, D., Neagu, A., Tastemel, M., Tuysuz, N., Lehmann, J., van de Werken,
H. J. G., Philipsen, S., van der Linden, R., Maas, A., van, I. W. F. J., Drukker,
M. & Ten Berge, D. Endogenous WNT signals mediate BMP-induced and
spontaneous differentiation of epiblast stem cells and human embryonic stem
cells. *Stem Cell Reports* **4**, 114-128, doi:10.1016/j.stemcr.2014.11.007 (2015).

81 Munoz Descalzo, S., Rue, P., Garcia-Ojalvo, J. & Martinez Arias, A.
Correlations between the levels of Oct4 and Nanog as a signature for naive
pluripotency in mouse embryonic stem cells. *Stem Cells* **30**, 2683-2691,
doi:10.1002/stem.1230 (2012).

82 Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut,
P., Chaisson, M. & Gingeras, T. R. STAR: ultrafast universal RNA-seq aligner.
Bioinformatics **29**, 15-21, doi:10.1093/bioinformatics/bts635 (2013).

83 Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with
high-throughput sequencing data. *Bioinformatics* **31**, 166-169,
doi:10.1093/bioinformatics/btu638 (2015).

84 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics **26**, 139-140, doi:10.1093/bioinformatics/btp616 (2010).

85 Yip, A. M. & Horvath, S. Gene network interconnectedness and the generalized
topological overlap measure. *BMC Bioinformatics* **8**, 22, doi:10.1186/1471-
2105-8-22 (2007).

86 Langfelder, P., Zhang, B. & Horvath, S. Defining clusters from a hierarchical
cluster tree: the Dynamic Tree Cut package for R. *Bioinformatics* **24**, 719-720,
doi:10.1093/bioinformatics/btm563 (2008).

87 Hubert, L., Arabie, P. . Comparing partitions. *Journal of Classification* **2**, 193-
218, doi:10.1007/BF01908075 (1985).

88 Ashburner, M., Ball, C. A., Blake, J. A., Botstein, D., Butler, H., Cherry, J. M.,
Davis, A. P., Dolinski, K., Dwight, S. S., Eppig, J. T., Harris, M. A., Hill, D. P.,
Issel-Tarver, L., Kasarskis, A., Lewis, S., Matese, J. C., Richardson, J. E.,
Ringwald, M., Rubin, G. M. & Sherlock, G. Gene ontology: tool for the
unification of biology. The Gene Ontology Consortium. *Nat Genet* **25**, 25-29,
doi:10.1038/75556 (2000).

89 Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes.
Nucleic Acids Res **28**, 27-30, doi:10.1093/nar/28.1.27 (2000).

90 Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for
comparing biological themes among gene clusters. *OMICS* **16**, 284-287,
doi:10.1089/omi.2011.0118 (2012).

1188

Supplementary Figure 2. Replica of *in vitro* ESC-EpiSC differentiation experiments upon chemical β-catenin perturbations

A-D Percentage of GFP+ cells calculated over the total amount of leaving cells in DMSO **(B)**, Ch1µM **(C)**, Ch3µM **(D)** and 2i/L **(E)** pre-cultured dual reporter ESCs. Histograms from Day 0 and Day 4 of each condition are shown as inset. Data are means±SEM (n=2, A (Day1 C1, Day3 AFC1, Day4 AFD-AFC3), B (Day 1 AFC1), D (Day4 AFD); n=3, A (Day1 AFD-AFC1-AFC3-C1X2-C3-C3X2, Day2, Day3 AFD-

AFC3-C1-C1X2-C3-C3X2, Day4 AFC1-C1-C1X2-C3-C3X2), B (Day1 AFD-C1-C1X2,

Day2, Day3, Day4), C (Day1, Day2, Day3, Day4), D (Day0, Day1, Day2, Day3, Day4);

n=6, A (Day0), B (Day0), C (Day0). p-values from one-way ANOVA are shown across

samples for each day. Dots represent individual data with similar values overlapping.

Colour-blind   safe   combinations   were   selected   using   colorbrewer2

(https://colorbrewer2.org/#type=sequential&scheme=BuGn&n=3).

1202
1203
1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

A

B

C

1217

1218

1219

1220

Supplementary Figure 3. FACS gating strategy and WGCNA of the genes correlating with time.

A Gating strategy used to sort C1M and C1H ESCs, following 48 hrs treatment with TMP10µM and Doxy100ng/mL. C1 ESCs treated with TMP10µM were used as negative control. **B** Clustering dendrogram of genes, with dissimilarity based on

1226 topological overlap, together with the assigned module colours; grey genes are
1227 unassigned to any module. **C** Eigenmodules correlating with time; the Pearson
1228 correlation coefficient (*r*) and relative p-value are shown. **D-H** Bar-chart of the top-ten
1229 enriched biological processes (BP) with FDR < 0.05 from genes belonging to the
1230 Green (**D**), Blue **(E**), Brown **(F)**, turquoise **(G)** and Pink **(H)** WGCNA modules.

Supplementary Figure 4. Gene ontology of the differential expressed genes in pluripotent ESCs.

A, B Bar-chart of the top-ten enriched biological processes (BP) with FDR < 0.05 from differentially expressed genes in in C1M **(A)** and C1H **(B)** compared to C1 ESCs. Black and grey bars represent upregulated and downregulated BPs, respectively. In red bars, the BPs exclusively enriched in the indicated condition.

Supplementary Table 1

List of identified module genes correlating with time (Supplementary Figure 3) or β-catenin doses (Figure 3) before and after the filtering for the |kME| ≥ 0.8; GO and pathway analysis.

Supplementary Table 2

List of differentially expressed genes before and after the filtering for the FDR (< 0.05) and log fold change (-2 > log > 2); GO and pathway analysis in pluripotent C1M vs C1 ESCs.

Supplementary Table 3

1277 List of differentially expressed genes before and after the filtering for the FDR (< 0.05)
1278 and log fold change (-2 > log > 2); GO and pathway analysis in pluripotent C1H vs C1
1279 ESCs.
1280

1281 **Supplementary Table 4**
1282 List of differentially expressed genes before and after the filtering for the FDR (< 0.05)
1283 and log fold change (-2 > log > 2); GO and pathway analysis in differentiated C1DMV
1284 vs C1D ESCs.
1285

1286 **Supplementary Table 5**
1287 List of differentially expressed genes before and after the filtering for the FDR (< 0.05)
1288 and log fold change (-2 > log > 2); GO and pathway analysis in differentiated C1DMDT
1289 vs C1D ESCs.
1290

1291 **Supplementary Table 6**
1292 List of differentially expressed genes before and after the filtering for the FDR (< 0.05)
1293 and log fold change (-2 > log > 2); GO and pathway analysis in differentiated C1HV vs
1294 C1D ESCs.
1295

1296 **Supplementary Table 7**
1297 List of differentially expressed genes before and after the filtering for the FDR (< 0.05)
1298 and log fold change (-2 > log > 2); GO and pathway analysis in differentiated C1DHDT
1299 vs C1D ESCs.
1300

1301 **Supplementary Movie 1**
1302 71 hrs time-lapse of dual reporter ESCs without pre-activation of the canonical Wnt
1303 pathway, differentiated in NDiff227 supplemented with ActivinA+Fgf2+DMSO.
1304

1305 **Supplementary Movie 2**
1306 71 hrs time-lapse of dual reporter ESCs without pre-activation of the canonical Wnt
1307 pathway, differentiated in NDiff227 supplemented with Chiron 1µM.
1308

1309 **Supplementary Movie 3**

1310   71 hrs time-lapse of dual reporter ESCs without pre-activation of the canonical Wnt

1311   pathway, differentiated in NDiff227 supplemented with Chiron 3μM.

1312