Optimal coordination and reorganization of photosynthetic properties in C₄ grasses

Haoran Zhou[1,2,3], Erol Akçay[1], Brent Helliker[1]

[1]Department of Biology, University of Pennsylvania, Philadelphia, PA 19104, USA

[2]School of the Environmental, Yale University, New Haven, CT 06511, USA

[3]University Corporation for Atmospheric Research, Boulder, CO 80301, USA

Correspondence: Haoran Zhou

Phone: 1-215-808-7042

Email: zhouhaoran06@gmail.com

ORCID iD: 0000-0002-5852-9535

Erol Akçay: eakcay@sas.upenn.edu

Brent Helliker: helliker@sas.upenn.edu

Running title: Optimal coordination in C₄ photosynthesis

Conflict of interest: None declared.

Paper type: Original Article

Word count: 5368

Number of figures: 7

Number of tables: 1

Supplementary material:

Supplementary Material I: supplementary dataset

Number of supplementary figures: 4

Number of supplementary tables: 11

Funding:

HZ and this research is supported by the NOAA Climate and Global Change Postdoctoral

Fellowship Program, administered by UCAR's Cooperative Programs for the Advancement

of Earth System Science (CPAESS) under award #NA18NWS4620043B and is also

supported by the Dissertation Completion Fellowship provided by the Graduate Division of

School of Arts and Sciences, University of Pennsylvania. BH is supported by NSF-IOS

award 1856587.

Abstract

C_3 and C_4 are major functional types in terrestrial biosphere models, with photosynthesis

traits as important input parameters. The evolution of C_4 required reorganizations of Calvin-

Benson-cycle and coordination of C_4-cycle enzymes, resulting in divergences of

physiological traits between C_3 and C_4. In addition, photosynthesis further optimized after the

evolution of C_4 causing diversification within C_4 lineages due to different evolutionary

histories. We combined optimality modeling, physiological measurements and phylogenetic

analysis to examine how various aspects of C_4 photosynthetic machinery were reorganized

and coordinated within C_4 lineages and as compared to closely-related C_3 in grasses.

Optimality models and measurements indicated a higher maximal electron transport to

maximal Rubisco carboxylation ratio (J_{max}/V_{cmax}) in C_4 than C_3, consistent with the optimal

prediction to maximize photosynthesis. The coordination between Calvin-Benson and C_4

cycles (V_{pmax}/V_{cmax}), however, is in line with the optimal modeling results under 200 ppm, as

opposed to current CO_2. Such inconsistencies can be explained by a slowly declining

assimilation rate beyond optimal V_{pmax}/V_{cmax}. Although rapid coordination occurred early in

C_4 evolution, C_4 is still under optimizing processes and photosynthetic measures have

continued to increase across time. Lastly, better understandings of J_{max}/V_{cmax}, V_{pmax}/V_{cmax} and

fluorescence-based-electron-transport proffer enhanced approaches to parameterize terrestrial

biosphere models.

Key words C_4 photosynthesis, electron transport, grass, optimality, J_{max}/V_{cmax}, V_{pmax}/V_{cmax},

evolution

Introduction

C_3 and C_4 are major photosynthesis pathways and important functional types identified in the

terrestrial biosphere models, e.g., Earth System Models (ESMs) and Land Surface Models

(LSMs) (Booth et al., 2012; Schaefer et al., 2012; Croft et al., 2017; Rogers et al., 2017).

Photosynthetic traits and parameters of C_3 and C_4 photosynthesis, e.g., maximal Rubisco

carboxylation rate (V_{cmax}) and maximal electron transport (J_{max}), are crucial inputs in these

models. Such parameters, as well as the J_{max}/V_{cmax}, are well-documented for C_3

photosynthesis, however, the C_4 photosynthesis are still underrepresented currently due to

lack of empirical reports and estimation (Bellasio *et al.*, 2016). The evolution of C_4 occurred

over an extended period that encompassed many different climate change scenarios; therefore

C_4 could respond to the future climate change differently than C_3 (Ward et al., 1999; Leakey

et al., 2009; Wittmer et al., 2010; Morgan et al., 2011; Reich et al., 2018). Additionally, it has

recently been proposed that by taking a lineage-based, or evolutionary, approach to ESM and

LSM parameterization that a more realistic approach to functional diversity can be captured

by these large-scale models (Griffith et al. 2020). Thus, it is crucial to identify the trait

divergences of C_4 photosynthesis from C_3, especially with regard to the photosynthetic

parameters.

C_4 photosynthesis evolved as a response to inefficiencies of C_3 photosynthesis that are

exacerbated under certain environmental conditions: low CO_2, drought, high temperature and

high light (Ehleringer & Monson, 1993; Ehleringer *et al.*, 1997; Edwards and Smith, 2010;

Zhou et al., 2018). Rubisco, the CO_2 carboxylating enzyme of the Calvin-Benson (CB) cycle

also assimilates O_2 as the first reaction of photorespiration and consequently reduces CB

cycle efficiency up to 30% in C_3 species (Ehleringer *et al.*, 1991; Bauwe *et al.*, 2010; Raines

2011). The C_4 pathway concentrates CO_2 around Rubisco and dramatically reduces

photorespiration by segregating atmospheric CO_2 uptake by Phosphoenolpyruvate

carboxylase (PEPc) and CB-cycle into two compartments within the leaf. The operation of C_4

cycle does have additional ATP costs for which C_3 plants do not remunerate (Hatch, 1987).

In sum, the assembly of C_4 photosynthesis broke the balance between CO_2 carboxylation and

electron transport of the CB cycle, which existed in C_3 ancestors: C_4 photosynthesis elevates

the efficiency of CO_2 carboxylation at the expense of using more energy from electron

transport (Sage, 2001, 2004, 2016; Christin & Osborne, 2014; Lundgren *et al.*, 2019). Thus, a

further reorganization of resource allocation between Rubisco carboxylation and electron

transport and the coordination between CB and C_4 cycle should occur along with, or as a

consequence of, C_4 formation. Furthermore, selection optimized the function of the C_4

through intermediate steps through adjusting physiological traits (Christin & Osborne, 2014;

Sage, 2016) and different C_4 lineages evolved at different time points and endured different

evolutionary histories. We therefore expect diversification in photosynthetic traits and

parameters among C_4 lineages. If C_4 photosynthesis is continuously under extended

optimization after its formation (Edwards, 2019; Heyduk et al., 2019), such diversification

would also represent evolutionary trends between photosynthetic parameters and

evolutionary time/ages. In the current study, we examine how various aspects of C_4

photosynthetic machinery— nitrogen allocation between Rubisco carboxylation and electron

transport, and CB and C_4 cycle— were reorganized and coordinated within C_4 lineage, as

well as compared to closely-related C_3 species, and whether the coordination of C_4

machineries are on a walk toward or have already reached the optimal state.

The relative ratio among V_{cmax}, J_{max} and maximal PEP carboxylation rate (V_{pmax}) could

represent the coordination within CB cycle and between CB and C_4 cycles. The resource

allocation between the Rubisco carboxylation and electron transport is represented by

J_{max}/V_{cmax}. Although J_{max}/V_{cmax} has been empirically measured in numerous C_3 species

(Wullschleger, 1993) and examined with optimal modeling results of J_{max}/V_{cmax} for C_3

(Walker *et al.*, 2014; Kromdijk & Long, 2016; Quebbeman & Ramirez, 2016), there have

been far fewer measurements or predictions in C_4. The coordination among CB cycle and the

C_4 cycle could be represented by the relationships among V_{cmax}, J_{max} and V_{pmax}. As a

relationship between J_{max} and V_{cmax} has been established above, V_{pmax}/V_{cmax} could be taken

into consideration to depict a complete picture of the functional coordination (Yin *et al.*,

2016).

In C_3 plants J_{max}/V_{cmax} displays acclimation with changing environmental factors

(temperature, CO_2, light, and water availability; Onoda *et al.*, 2005; Rodriguez-Calcerrada *et*

al., 2008; Kromdijk & Long, 2016; Yin *et al.*, 2018), and it is expected that J_{max}/V_{cmax} and

V_{pmax}/V_{cmax} might also vary in C_4. Sage and McKown (2006) proposed that C_4 may show less

plasticity and acclimation in phenotypical traits in response to global climate change, due to

their complex anatomical and biochemical features, e.g., the structural and physiological

integration of the mesophyll-bundle sheath complex. Combining theoretical predictions and

empirical examination of how J_{max}/V_{cmax} and V_{pmax}/V_{cmax} vary with environment could

elucidate the acclimation capability C_4 and further show if acclimation occurs in an optimal

manner. Such an understanding of C_4 responses to changing climate would reduce the

uncertainty of global carbon flux prediction (Beerling & Quick, 1995; Quebbeman &

Ramirez, 2016; Croft et al., 2017; Rogers et al., 2017).

In addition to the coordination between CB and C_4 cycle detailed above, we expect a

coordination between cyclic electron transport and total electron transport. Cyclic-electron

transport, which produces ATP only, has been proposed to be enhanced C_4 plants as

compared to C_3 plants to fulfill the extra ATP requirements of C_4 cycle (Takabayashi *et al.*,

2005; Nakamura et al., 2013; Munekage, 2016; Munekage & Taniguchi, 2016; Yin & Struik,

2018). Chlorophyll fluorescence has been widely used for C_3 species to estimate the J_{max} (Yin

et al., 2009; Bellasio et al., 2016); however, chlorophyll fluorescences, representing the

electron transport associated with linear electron transport, differs between C_3 and C_4 species.

Lack of details in the proportion of linear electron transport in total electron transport in C_4

species, therefore, hindered the use of fluorescence method to estimate electron transport in

C_4 species. We expected such a ratio will be lower in C_4 than that in C_3. Establishing this

ratio will provide for an additional tool to estimate J_{max} using fluorescence measurements, as

well as V_{cmax} and V_{pmax} for C_4 species.

In the current study, we first used physiological models coupling photosynthesis, hydraulics

and nitrogen stoichiometry to predict the optimal J_{max}/V_{cmax} and V_{pmax}/V_{cmax}. Then, we

performed in vivo experiments to estimate these parameters empirically on grass lineages

including C_3 and C_4 species selected from the PACMAD clade (Grass Phylogeny Working

Group II, 2012; Spriggs *et al.*, 2014) and compared these to published in vitro measurements.

By sampling multiple independent origins of C_4 species within a phylogenetic context

(Edwards *et al.*, 2007; Cavender-Bares *et al.*, 2009), we were able to use phylogenetic

comparative methods to examine the divergence of traits between C_3 and C_4 species and, to

detect whether there are continuous evolutionary trends. In sum, we used optimality

modeling, physiological measurements and evolutionary comparative methods to examine

evolutionary trends, the approach to optimality, and to gain a better formal understanding of

J_{max}/V_{cmax} and V_{pmax}/V_{cmax} in C_4 photosynthesis to reduce uncertainty in modeling the

carbon/nitrogen cycle, vegetation dynamics and productivity. Specifically, we aimed to test

and examine the following hypotheses and questions. (1) Reorganization in resource

allocation between Rubisco carboxylation and electron transport and coordination between

CB cycle and C_4 cycle in C_4 resulted in higher J_{max}/V_{cmax} and lower flr-ETR/J_{max} than C_3. (2)

C$_4$ values for J_{max}/V_{cmax} and V_{pmax}/V_{cmax} are optimized the for current environment

conditions. (3) Although evolving millions of years ago, selection has continued to optimize

C$_4$ after the evolution of the CCM, and this will yield a positive evolutionary trend between

photosynthesis parameters and evolutionary age. (4) To examine the acclimation capability of

C$_4$ species using predictions and measurements of how J_{max}/V_{cmax} and V_{pmax}/V_{cmax} vary with

environmental change.

Materials and Methods

Plant material

We cultivated 35 closely related species, including nine C$_3$ and 26 C$_4$. The species belong to

nine independent origins of closely-related C$_3$ and C$_4$ lineages. Detailed cultivation

information is the same as Zhou *et al.* (2020).We, then, extracted the dated phylogenetic tree

for the species from the dated phylogenetic trees (Spriggs *et al.*, 2014) (Fig. S1).

Physiological Modeling

Optimal J_{max}/V_{cmax} and V_{pmax}/V_{cmax} Based on the C$_3$ and C$_4$ models constructed in Zhou *et*

al. (2018), which incorporates the soil-plant-air water continuum into traditional C$_3$ and C$_4$

photosynthesis models (Farquhar et al., 1980; von Caemmerer, 2000), we added

stochiometric correlations between photosynthesis parameters and nitrogen. Using such a

framework, we can model the optimal J_{max}/V_{cmax} and V_{pmax}/V_{cmax} simultaneously considering

the following nitrogen stoichiometry:

The total nitrogen is the sum of different components (Evans, 1989):

$N_{org} = N_P + N_E + N_R + N_S + N_O,$ (1)

in which N_P represents the nitrogen in pigment proteins, N_E represents the nitrogen for the

electron transport system, N_R represents the nitrogen of Rubisco, N_S represents nitrogen in

soluble proteins except for Rubisco, and N_O represents additional organic leaf nitrogen not

invested in photosynthetic functions.

In order to model the optimal J_{max}/V_{cmax} and V_{pmax}/V_{cmax}, we need to consider the nitrogen

stoichiometry among J_{max}, V_{cmax} and V_{pmax}. We used empirical relationships found in previous

studies (Evans & Poorter, 2001; Niinemets & Tenhunen, 1997; Quebbeman & Ramirez,

2016):

$N_P + N_E = 0.079 J_{max} + 0.0331 \chi,$ (2)

$N_S = \nu J_{max},$ (3)

$N_R = V_{cmax}/(6.25 \times V_{cr} \times \xi),$ (4)

$N_{PEP} = V_{pmax}/(6.72 \times V_{pr} \times \xi),$ (5)

χ is the concentration of chlorophyll per unit area (µmol Chl m^{-2}), 0.079 is in mmol N s

(µmol)$^{-1}$, and 0.0331 is in mmol N (µmol Chl)$^{-1}$, $\nu \approx 0.3$ (mmol N s (µ mol)$^{-1}$. V_{cr} is the

specific activity of Rubisco (the maximum rate of RuBP carboxylation per unit Rubisco; $\approx$

20.5 µmol CO_2 (g Rubisco)$^{-1}$s^{-1}) and 6.25 is grams RuBisCO per gram nitrogen in RuBisCO.

V_{pr} is the specific activity of PEPc, that is, the maximum rate of RuBP carboxylation per unit

PEPc ($\approx$ 181.7 µmol CO_2 (g PEPC)$^{-1}$s^{-1}), 6.72 is grams PEPc per gram nitrogen in PEPC

(calculated from the amino acids composition of Fujita et al., 1984), and ξ is the mass in

grams of one millimole of nitrogen equal to 0.014 g N (mmol N)$^{-1}$.

Further, we simplify the equation (2) by assuming there is a coordination of resource

allocation between chlorophyll and electron transport for saturated light intensity, which

determines the J_{max}. We make this assumption for the light saturated condition and use the

empirical equation of Croft et al. (2017) to equation (2)

$\chi = \frac{1000 J_{max}}{2.49} / \eta,$ (6)

where η is the average molar mass for chlorophyll (900 g/mol). Thus,

$N_{org} - N_O = 0.079 J_{max} + 0.0331 \chi + \nu J_{max} + \frac{V_{cmax}}{6.25 \times V_{cr} \times \xi} + V_{pmax}/(6.72 \times V_{pr} \times \xi),$ (7)

When the light intensity varies, the following function is used to adjust the electron transport

rate (Ögren & Evans, 1993):

$J = \frac{I_2 + J_{max} - \sqrt{(I_2 + J_{max})^2 - 4\Theta I_2 J_{max}}}{2\Theta}.$ (8)

Also, all the photosynthetic parameters are temperature-sensitive (Zhou *et al.*, 2018).

In the optimal modeling processes, we set N_{org}- N_O as constant of 129 mmol N m^{-2} (which

yield a V_{cmax}= 39 μmol m^{-2} s^{-1}, J_{max}= 195 μmol m^{-2} s^{-1} and V_{pmax}= 78 μmol m^{-2} s^{-1}, if assuming

J_{max}/V_{cmax}=5 and V_{pmax}/V_{cmax}=2). Using these models, we modeled the assimilation rates with

different J_{max}/V_{cmax} from 1 to 8 of 0.01 interval and different V_{pmax}/V_{cmax} from 0.5 to 5 of 0.01

to find the globally optimal assimilation rate with respect to both J_{max}/V_{cmax} and V_{pmax}/V_{cmax}.

The corresponding J_{max}/V_{cmax} or J_{max}/V_{cmax} under the highest assimilation rates represent the

optimal ratios. Then, we also model the locally optimal J_{max}/V_{cmax} and V_{pmax}/V_{cmax} when

constraining the corresponding V_{pmax}/V_{cmax} and J_{max}/V_{cmax} with the average measured values

respectively.

Using the model described above, we were able to model the optimal J_{max}/V_{cmax} and

V_{pmax}/V_{cmax} under different environmental gradients: CO_2 of 200, 300, 400, 500 and 600 ppm;

VPD and ψ_S of (0 MPa, 0.15) (0.625, -0.5), (1.25, -1), (1.875, -1.5), and (2.5, -2); light

intensity of 2000, 1600, 1200, 800 and 400 μmolm^{-2}s^{-1}; temperature of 15, 20, 25, 30 and 35

^{o}C. Since there is potential uncertainty for stochiometric relationships, we performed

sensitivity analysis for the total nitrogen (from 100% to 50% with 10% interval of the regular

nitrogen) for optimal J_{max}/V_{cmax} and V_{pmax}/V_{cmax} results and we also performed sensitivity

analysis for stoichiometry of PEPC by varying the $1/(6.72 \times V_{pr} \times \xi)$ term in Eq. 5 from 50%

to 800%. For the C_3 pathway, all the modeling process are similar with the C_4 and a same

value of N_{org}- N_O is used, except that a simplified version of equation (7) is used as below

(Quebbeman & Ramirez, 2016):

$$N_{org} - N_O = 0.079 J_{max} + 0.0331\chi + \nu J_{max} + \frac{V_{cmax}}{6.25 \times V_{cr} \times \xi},$$ (9)

Effects of decreasing V_{cmax} Using the model, we also simulated the effect of decreasing V_{cmax}

on the assimilation rate of both C_3 and C_4 pathway. In this modeling process, we hold J_{max}

and other photosynthetic parameters constant as the initial modeling condition as above, but

varying the V_{cmax} to 100%, 90%, 80%, 70%, 60% and 50% of the original values of C_3.

Gas exchange and fluorescence measurements

We measured A/C_i curves using a LI-6400XT (LI-COR Inc., Lincoln, NE, USA) for all the

species by setting the CO_2 concentrations as 400, 200, 50, 75, 100, 125, 150, 175, 200, 225,

250, 275, 300, 325, 350, 400, 500, 600, 700, 800, 1000, 1200, 1400 ppm under light intensity

of 2000 $\mu mol m^{-2}s^{-1}$. Data were recorded when the intercellular CO_2 concentration

equilibrated 2-5 minutes. Fluorescence was measured along with A/C_i curves using a 2 cm^2

fluorescence chamber head. After the change of CO_2 concentration, the quantum yield was

measured by multiphase flash when A reached a steady state (Bellasio *et al.*, 2014). The leaf

temperatures were controlled at 25°C, VPD varied at 1-1.7kPa and the flow rate of 500 μmol

s^{-1} for all the measurements. The cuvette was covered by Fun-Tak to lessen leakiness. We

used the estimation method in Zhou *et al.* (2019) for A/C_i curves to estimate in vivo V_{cmax},

J_{max}, and V_{pmax} with one slight methodological change. Since it is thought that the region of

V_{cmax} limit A/C_i is very narrow in C_4 species, we assigned the C_i regions limited by carbonic

anhydrase and V_{cmax} with a very low criteria of 5 Pa or below. We let the data points with C_i

ranging from 5 to 60 Pa CO_2 to be freely determined by which of the four potential

limitation states to minimize the estimation error. Using this method, we avoided the

potential bias of including optimal perspectives to the estimation method which could occur

when directly assigning the cross points co-limited by V_{cmax}, V_{pmax} and J_{max}. Fluorescence

results were used to calculate the flr-ETR (Genty et al., 1989).

Furthermore, in vivo V_{cmax} estimated from A/C_i could be a source of uncertainty in C_4

species. In order to provide additional support to the measured J_{max}/V_{cmax} and V_{pmax}/V_{cmax}

value, we collected in vitro measured values for V_{cmax} and V_{pmax} from previous research,

which includes eleven studies with 87 averaged results reported under current and varying

environmental conditions (Supplementary Material I). Since it is impossible to obtain in vitro

J_{max} and the estimation of J_{max} from A/C_i curves is considered reliable, we also obtain the

corresponding A/C_i curves from these studies if they were reported, to obtain the J_{max}. The

combination of in vivo and in vitro measurements should yield a good representation of

current J_{max}/V_{cmax} and V_{pmax}/V_{cmax} states in the C_4 plants.

Chlorophyll measurements and leaf nitrogen

Chlorophyll were measured using the spectrophotometer method (Porra et al., 1989). We cut

the fresh leaves of species into pieces of 0.5 mm long (total leaf area was measured) and

submerged the fragments into DMF. After all the Chlorophyll was extracted and the leave

turned white, the supernatant was used to measure the absorption under 663.8 nm and 646

nm. Total Chlorophyll concentrations were calculated using the equation of Porra et al.

(1989). We measure leaf nitrogen contents for each sample using the CHNOS analyzer

(ECS4010, Costech Analytical Technologies, Inc., Valencia, CA).

Phylogenetic analysis

We fitted each of the photosynthetic parameters (V_{cmax}, J_{max}, J_{max}/V_{cmax}, Total Chl, flr-

ETR/J_{max}, flr-ETR, V_{pmax} and V_{pmax}/V_{cmax}) to six different evolutionary models falling into

Brownian Motion model and Ornstein-Uhlenbeck Model using the R package of

"mvMORPH" (Table S1). The small-sample-size corrected version of Akaike information

criterion (AICc, the lower AICc, the better fit) and Akaike weights (AICw, the higher AICw,

the better fit) were used as criteria to figure out the best-fitted model. We used likelihood-

ratio test (LRT) method to test whether one model variant performs significantly better than

others and to determine whether there are significant differences between C_3 and C_4 species.

We also extract the evolutionary ages for each C_4 species from the dated phylogeny (Spriggs

et al., 2014). We regressed the above photosynthetic traits with evolutionary ages to detect

potential evolutionary trends.

Results

In vivo and in vitro J_{max}/V_{cmax} follow the global optima in C_4, but V_{pmax}/V_{cmax} does not.

In vivo and in vitro J_{max}/V_{cmax} are consistent with the optimal predictions under current CO_2

conditions, but V_{pmax}/V_{cmax} fell in to the optimal range under CO_2 of 200 ppm. The global

optima modelling results indicated maximal photosynthesis at the J_{max}/V_{cmax} of 5-6.2, which

is relatively constant across different CO_2 concentrations, while the optimal range for

V_{pmax}/V_{cmax} for C_4 species is 1.4-2.2 at CO_2 of 200 ppm, but decreases to 1-1.6 when CO_2

reaches 400 and 600 ppm (Fig. 1b, d, f). The averaged in vitro and in vivo J_{max}/V_{cmax} are

consistent with the global optimal predictions under CO_2 of 400 ppm (Fig. 1, Fig. 2a,

Supplementary Material I), as well as the locally optimal predictions controlling V_{pmax}/V_{cmax}

at the in vivo and in vitro level (Fig. 2). However, the averages of in vitro and in vivo

V_{pmax}/V_{cmax} are beyond the optimal predictions of global optima at CO_2 of 400 ppm, but the

measurement results are consistent with the optimal condition at CO_2 of 200 ppm (Fig. 1, 3,

4, S2, Supplementary Material I). The 3D images and the contour plots also illustrate that

when J_{max}/V_{cmax} is at the optimal range, beyond the optimal range of V_{pmax}/V_{cmax}, the

assimilation surface is quite flat: photosynthesis declines, but quite slowly; however, when

V_{pmax}/V_{cmax} drops outside of the optimal ranges, there are sharp decreases of photosynthesis

(Fig. 1). In vitro measurements indicated large variations of J_{max}/V_{cmax} and V_{pmax}/V_{cmax} at the

species level, and in vivo results, within a small scale of variation, fell completely into the

range of in vitro results (Fig. 2a and 3a).

C_4 species have higher J_{max}/V_{cmax} and lower flr-ETR/J_{max} than C_3 species.

Phylogenetic analysis shows the J_{max}/V_{cmax} follows the Ornstein-Uhlenbeck model with a

higher stable state for C_4 species and a lower stable state for C_3 (Table 1; Fig. 1a). Such an

empirical relationship is consistent with the optimal modeling predictions for C_3 and C_4. We

looked further into how such a higher J_{max}/V_{cmax} in C_4 species is reached by comparing

individual empirical parameters. C_4 species has significantly higher stable states of J_{max}, but

significantly lower stable states of V_{cmax} and nitrogen content than their closely-related C_3

(Table 1). In order to examine the potential effects of decreasing V_{cmax} on assimilation rate,

we held the J_{max} as constant and changed the V_{cmax} from 100% to 50% of the original C_3

parameter values in the C_3 and C_4 models. A decrease in V_{cmax} will significantly decrease the

assimilation rates of C_3 species from 10 °C to 35°C under different CO_2 concentrations, while

decreasing V_{cmax} has little effects on the assimilation rates of C_4 species (Fig. 5).

The flr-ETR/J_{max} was significantly higher in C_3 species than that in their closely related C_4

species, with the stable states of 1.08 and 0.64 for C_3 and C_4 respectively (Table 1).

Phylogenetic analysis indicated C_3 and C_4 species had the similar stable states of ETR, but C_4

has a higher total electron transport rate, J_{max} (Table 1).

The positive evolutionary trends of photosynthetic parameters

Plotting the photosynthetic parameters with evolutionary ages, extracted from a dated

phylogeny for the multiple lineages, allows us to look for further evolutionary trends in C_4

and their closely related C_3 species. Regressions of evolutionary age versus photosynthetic

traits provide signals for long-term directional trends in photosynthetic machinery following

the establishment of C_4 photosynthesis (Fig. 6). A_{max}, J_{max}, total chlorophyll, V_{cmax} and V_{pmax}

showed significant positive correlations with evolutionary age in C_4 species, but not C_3

species, while nitrogen, J_{max}/V_{cmax}, V_{pmax}/V_{cmax} and flr-ETR$/J_{max}$ did not show significant

correlation with the evolutionary age.

Optimal variation of J_{max}/V_{cmax} and V_{pmax}/V_{cmax} with environmental conditions.

To understand how the J_{max}/V_{cmax} and V_{pmax}/V_{cmax} varied theoretically in response to

environmental changes (Fig. 7a, b, Fig. S3a, b), we calculated their optimal value for varying

CO_2 concentrations, water limitations, temperatures, and light intensities. The optimal

J_{max}/V_{cmax} is predicted to increase linearly in C_3 and, but increase a little bit and maintain

relatively constant in C_4 species with increasing CO_2 concentration (Fig. 7a). The optimal

J_{max}/V_{cmax} decreases similarly in both C_3 and C_4 species along with increasing water limitation

and increases similarly with decreasing light intensity and increasing temperatures (Fig. 7).

The changes of J_{max}/V_{cmax} with water limitation, light intensity, and temperature are non-

linear, with the rate-of-change increasing greatly after a threshold (water limitation of ψ_S=-1,

VPD=1.25, the light intensity of 800 μmol m^{-2} s^{-1} and temperature of 30 ^{o}C). The optimal

V_{pmax}/V_{cmax} decreases along with the increase of the CO_2 concentration, especially when CO_2

increases from 200 ppm to 300 ppm, but the change is little when CO_2 is above 400 ppm

(Fig. 7c, S3c). However, V_{pmax}/V_{cmax} is relatively constant with the varying of water limitation

conditions and light intensity (Fig. 7c,d, S3c,d). V_{pmax}/V_{cmax} decreases with the rise in

temperature from 15 to 35 °C (Fig. 7d, S3d).

Sensitivity analysis for optimal J_{max}/V_{cmax} and V_{pmax}/V_{cmax}

Since there is a great variation of total nitrogen content in plants and there is uncertainty in

the stoichiometry of PEPC, we performed sensitivity analysis and found the optimal

modeling of J_{max}/V_{cmax} and V_{pmax}/V_{cmax} are robust. For C_3 photosynthesis, optimal J_{max}/V_{cmax}

increases slightly with decreasing total nitrogen (7.7% increase of J_{max}/V_{cmax} with 50%

reduction in total nitrogen), while for C_4, optimal J_{max}/V_{cmax} decreases with total nitrogen

(16.2% increase of J_{max}/V_{cmax} with 50% reduction in total nitrogen) (Fig. S4a). The optimal

V_{pmax}/V_{cmax} increases somewhat with decreasing total nitrogen (8.7% increase of V_{pmax}/V_{cmax}

with 50% reduction in total nitrogen) (Fig. S4a). The optimal J_{max}/V_{cmax} is relatively constant

with the change of stoichiometry of PEPC from 50% to 200%, but increases as the

stoichiometry increases from 200% to 800% (Fig. S3b). The optimal V_{pmax}/V_{cmax} is relatively

robust with the change of stoichiometry of PEPC.

Discussion

Our modeling efforts provide an explanation for the observed variation in J_{max}/V_{cmax} and

V_{pmax}/V_{cmax}, and why V_{pmax}/V_{cmax} appears to be optimized for the lower bounds of atmospheric

CO_2 of the Pleistocene. Our reported value of V_{pmax}/V_{cmax} are comparable with previous

studies (Kubien *et al.*, 2003; Pengelly *et al.*, 2010; Yin *et al.*,2011, 2016; Pignon and Long,

2020), and two recent papers also indicated that the coordination between CB and C_4 cycles

might represent a legacy of ancient low CO_2 conditions (Sundermann et al., 2018; Pignon and

Long, 2020). All extant C_4 species have gone through a low CO_2 bottleneck over the last 5

million years (Edwards et al., 2010). This bottleneck may have resulted in a strong selection

to increase V_{pmax}/V_{cmax} to maintain a high assimilation rate under the low CO_2 of glacial

maxima (~200 ppm). As CO_2 has risen, first with the beginning of the Holocene interglacial,

and then again with the continual burning of fossil fuels, V_{pmax}/V_{cmax} remained constant and

consequently exceeded the optimal V_{pmax}/V_{cmax} at higher CO_2. The effects of a too-high

V_{pmax}/V_{cmax} on assimilation rate are, however, minimal, thus the selection against increasing

V_{pmax}/V_{cmax} was likely weak. The explanation directly rests on the topology of assimilation

surface: when J_{max}/V_{cmax} and V_{pmax}/V_{cmax} are lower than the optimal states, assimilation rate

declines greatly; but when J_{max}/V_{cmax} and V_{pmax}/V_{cmax} exceed the optimal states, the decrease

of assimilation rate is minimal. Similarly, if a high J_{max}/V_{cmax} was selected for a given species

under, for example, growth at relatively low light conditions, the high J_{max}/V_{cmax} could well

be maintained for a long time even when moving to a high light environment. Such small

changes in assimilation rate may not change fitness enough to drive evolution toward optimal

states in natural plants; artificial selection and manipulation to change the J_{max}/V_{cmax} and

V_{pmax}/V_{cmax} toward the optimal states, however, might show potential in regard to increasing

total assimilation rate and productivity (Walker et al., 2018; Pignon and Long, 2020).

It has been hypothesized that after the evolution of the full C_4 CCM that selection across

various habitats would select for further physiologically optimal adjustments (Williams et al.,

2013; Stata et al., 2019); we find unequal support for such adjustments through time as

several physiological measures were positively related to evolutionary age, but there were no

trends with photosynthetic coordination. From this, we conclude that initially there was very

strong selection for coordination between the CB cycle, light reactions and the C_4 CCM, and

that changes in secondary or tertiary traits led to an increase in, for example, maximum CO_2

assimilation rate through time. Zhou et al. (2020) illustrated that hydraulic traits could be just

such an example of the secondary traits: the formation of C_4 pathway allowed for lower

stomatal conductance which triggered the decline of leaf hydraulic conductance and

capacitance over evolutionary time to further maximize the assimilation rate. Additionally,

anatomical organization within the leaf, like the ratio of mesophyll cells to bundle-sheath

cells, the 3D arrangement of cells and shifts in intercellular airspace could also be selected

upon through time towards a more optimal C_4 photosynthetic machine (Wang et al., 2017;

Alonso-Cantabrana et al., 2018; Edwards, 2019).

Higher J_{max}/V_{cmax} in C_4 than that in C_3 indicated a change in resource allocation, namely

nitrogen, between the light reactions and the CB cycle, and as a crucial evolutionary step for

elevating C_4 efficiency, it is important to look at how the reallocation may have occurred and

the larger-scale consequences. The modeling results indicate that a decrease of Rubisco

content is favored in C_4, because overall nitrogen requirements decrease and such a reduction

has minimal effects on net assimilation rate. Significantly lower V_{cmax} in all of our C_4 species

and lower Rubisco in previous studies confirmed the assertion (Brown, 1978; Ku *et al.,* 1979;

Sage and Pearcy, 1987; Sharwood *et al.*, 2016). Any surplus nitrogen not invested in Rubisco

could be distributed among three broad categories: 1) reallocated to the light reactions or 2)

stored or used to construct new tissues or 3) simply not taken up from the growth

environment, thus reducing total plant nitrogen requirements. Tissue *et al.* (1995) and

Ghannoum *et al.* (2010) detected lower Rubisco content and higher chlorophyll and thylakoid

content, supporting resource reallocation from RuBP carboxylation to electron transport

within the leaf. Our measurements provide evidence that the coordination of J_{max}/V_{cmax}

resulted from a mix of hypotheses 1) and 3). The significantly higher J_{max} and lower V_{cmax} in

C_4 than their closely-related C_3 species supports a reallocation of hypothesis 1). In addition,

hypothesis 3) likely occurred together with hypothesis 1) because C_4 species have

significantly lower nitrogen content. Hypothesis 2), not exclusive to hypotheses 1) and 3),

could be supported by evidence that C_4 plants maintain larger leaf areas (Ripley *et al.*, 2008).

These hypotheses are connected to potential ecological ramifications. Firstly, in a nitrogen-

depleted habitat, C_4 could have a competitive advantage as confirmed by Ripley *et al.* (2008;

although Sage & Pearcy, 1987, found no evidence for this). In habitats where nitrogen is not

limiting, the excess nitrogen could be used to construct more leaf area (Sage & Pearcy, 1987;

Anten *et al.*, 1995; Ripley *et al.*, 2008), and greater leaf area in the early stages of growth was

indeed seen by Atkinson *et al.* (2016). On the other hand, the lack of nitrogen reallocation

from the CB cycle to the light reactions may indicate physiological constraints in fertile

habitats. For example, photorespiration in C_3 plants is proposed to enhance the nitrate

metabolism (Oaks, 1994; Rachmilevitch *et al.*, 2004; Bauwe *et al.*, 2010; Bloom, 2015),

therefore the formation of CCM, which inhibits photorespiration, may reduce overall plant-

available nitrogen for C_4. In addition, the increase of J_{max} in C_4 is due to an enhanced cyclic

electron transport, while maintaining the linear electron transport at the same level of C_3.

Elevating cyclic electron transport is therefore a potentially important step in engineering C_4

photosynthesis into C_3 crops.

Empirical evidence indicates adaptive plasticity (or acclimation) of C_4 photosynthetic

coordination occurs under different environmental conditions, consistent with optimal

predictions. Contrary to what Sage and McKown (2006) proposed, C_4 exhibited significant

acclimation capability with varying CO_2 (Pinto et al. 2014; Pinto et al. 2016), water

availability (Sharwood et al. 2014), light intensity (Sharwood et al. 2014; Pengelly et al.

2010; Sonawane, 2016) and temperature (Sonawane, 2016; Pittermann and Sage 2001;

Kubien and Sage 2004; Serrano-Romero and Cousins, 2020) in both J_{max}/V_{cmax} and

V_{pmax}/V_{cmax} (Supplementary Material I). The varying trends in J_{max}/V_{cmax} are generally

consistent between the empirical measurements and optimal modeling predictions in the

current study: higher resource allocation to electron transport under shade, elevated CO_2,

high water availability and higher temperature. However, C_3 and C_4 species might respond

differently to environmental variations, especially in response to CO_2 variation shown in the

optimal modeling results. The V_{pmax}/V_{cmax} exceededs the optimal prediction when acclimated

to the current environmental conditiona, perhaps because V_{pmax}/V_{cmax} is constraint by a legacy

of acclimation to historical CO_2. These acclimation responses should be considered and

included in ESMs and LSMs to predict responses of C_3 and C_4 ecosystems to future climate

change (Rogers et al., 2017; Smith and Keenan, 2020).

Our improved estimates of J_{max}/V_{cmax}, V_{pmax}/V_{cmax} and flr- ETR/J_{max} for C_4 plants could

directly benefit terrestrial biosphere models. Although J_{max}, V_{cmax} and V_{pmax} are key input

parameters in global-scale models (Beerling & Quick, 1995; Zaehle et al., 2005; Bonan et al.,

2011; Walker et al., 2014), it is difficult and perhaps not feasible to measure all parameters

for numerous sites. If one the ratioed parameters described here, either J_{max}/V_{cmax}, V_{pmax}/V_{cmax}

and/or flr- ETR/J_{max}, is obtained then other parameters could be estimated. Using J_{max}/V_{cmax},

V_{pmax}/V_{cmax} and flr- ETR/J_{max} is especially crucial in C_4 species because in vivo estimation of

V_{cmax} and V_{pmax} are more difficult and less reliable, and in vitro measurements are not easily

performed over broad taxonomic or spatial scales. Our results suggest a value of around 5.5

for J_{max}/V_{cmax} and a value of 2 for V_{pmax}/V_{cmax} supported by optimality modeling and

empirical measurements. With an empirical value of around 60% found in the current study,

the measurement of flr- ETR could provide another tool to further estimate J_{max}, V_{cmax} and

V_{pmax}. Previous theoretical analysis also inferred values of flr- ETR/J_{max} of 60% and 50% in

C_4 (Yin et al., 2011; Yin & Struik, 2012; Yin & Struik, 2018).

The evolution of C_4 photosynthesis required the reorganization and coordination of the CB-

cycle, the light reactions and the PEPc-based CCM. Strong divergence in J_{max}/V_{cmax} between

C_4 and C_3 species indicates the resource allocation changes between light reactions and the

CB cycle that were necessary to support the enhanced ATP requirement of the C_4 CCM

(Osborn & Sack, 2012; Zhou et al., 2018). Observed J_{max}/V_{cmax} were within the predicted

optimal zone suggesting that the resource reallocation between Rubisco carboxylation and

electron transport are operating near optimality under current environmental conditions;

however, the long tail exceeding the optimal J_{max}/V_{cmax} in empirical measurements indicates

multiple species have over-allocated to electron transport, perhaps a legacy of native

ecological conditions. The coordination between CB and C_4 cycles was in line with the

optimal conditions under 200 ppm representing an over-allocation of resources for current

environmental conditions, but there is little costs to assimilation rate due to this lack of

optimality. Rapid coordination occurred early in C_4 evolution, but it appears that C_4

photosynthesis is still under selection for further optimization. The enhanced understanding

of the evolution-based photosynthetic reorganization and coordination in C_4 photosynthesis,

along with our ratio-based approach to obtain photosynthetic parameters can lead to a better

parameterization of terrestrial biosphere models for C_4.

Acknowledgements

We sincerely thank Dr. Matteo Detto (Princeton University) for his help in building the

optimal models for J_{max}/V_{cmax}.

Reference

Alonso-Cantabrana H., Cousins A.B., Danila F., Ryan T., Sharwood R.E., von Caemmerer

S., Furbank R.T. (2018). Diffusion of CO_2 across the mesophyll-bundle sheath cell

interface in a C_4 plant with genetically reduced PEP carboxylase activity. *Plant*

Physiology, 178, 72–81.

Anten N.P., Schieving F., Medina E., Werger M.J.A., Schuffelen P. (1995). Optimal leaf area

indices in C_3 and C_4 mono-and dicotyledonous species at low and high nitrogen

availability. *Physiologia plantarum*, 95, 541-550.

Atkinson R.R., Mockford E.J., Bennett C., Christin P.A., Spriggs E.L., Freckleton R.P.,

Thompson K., Rees M., Osborne C.P. (2016). C_4 photosynthesis boosts growth by

altering physiology, allocation and size. *Nature Plants*, 2, 16038.

Bauwe H., Hagemann M., Fernie A.R. (2010). Photorespiration: players, partners and

origin. *Trends in plant science*, 15, 330-336.

Beerling D.J., Quick W.P. (1995). A new technique for estimating rates of carboxylation and

electron transport in leaves of C_3 plants for use in dynamic global vegetation

models. *Global Change Biology*, 1, 289-294.

Bellasio C., Beerling D.J., Griffiths H. (2016). An Excel tool for deriving key photosynthetic

parameters from combined gas exchange and chlorophyll fluorescence: theory and

practice. *Plant, cell & environment*, 39, 1180-1197.

Bellasio C., Burgess S.J., Griffiths H., Hibberd J.M. (2014). A high throughput gas exchange

screen for determining rates of photorespiration or regulation of C_4 activity. *Journal of*

experimental botany, 65, 3769-3779.

Bloom A.J. (2015). Photorespiration and nitrate assimilation: a major intersection between

plant carbon and nitrogen. *Photosynthesis Research*, 123, 117-128.

Bonan G.B., Lawrence P.J., Oleson K.W., Levis S., Jung M., Reichstein M., Lawrence D.M.,

Swenson S.C. (2011). Improving canopy processes in the Community Land Model

version 4 (CLM4) using global flux fields empirically inferred from FLUXNET

data. *Journal of Geophysical Research: Biogeosciences*, 116(G2).

Booth B.B., Jones C.D., Collins M., Totterdell I.J., Cox P.M., Sitch S., Huntingford C., Betts

R.A., Harris G.R. and Lloyd J. (2012). High sensitivity of future global warming to land

carbon cycle processes. *Environmental Research Letters*, 7, p.024002.

Brown R.H. (1978). A difference in N use efficiency in C_3 and C_4 plants and its implications

in adaptation and evolution 1. *Crop Science*, 18, 93-98.

von Caemmerer S. (2000). Biochemical models of photosynthesis. In Techniques in Plant

Sciences. CSIRO Publishing, Colingwood, Australia, pp 91–122.

Cavender-Bares J., Kozak K.H., Fine P.V., Kembel S.W. (2009). The merging of community
ecology and phylogenetic biology. *Ecology letters*, 12, 693-715.

Christin P.A., Osborne C.P. (2014). The evolutionary ecology of C$_4$ plants. *New*
Phytologist, 204, 765-781.

Croft H., Chen J.M., Luo X., Bartlett P., Chen B., Staebler R.M. (2017). Leaf chlorophyll
content as a proxy for leaf photosynthetic capacity. *Global change biology*, 23, 3513-
3524.

Edwards E.J. (2019). Evolutionary trajectories, accessibility and other metaphors: the case of
C4 and CAM photosynthesis. *New Phytologist*, 223, 1742-1755.

Edwards E.J., Osborne C.P., Strömberg C.A.E., Smith S.A., C$_4$ Grasses Consortium. (2010).
The origins of C$_4$ grasslands: integrating evolutionary and ecosystem science. *Science*,
328, 587–591.

Edwards E.J., Still C.J., Donoghue M.J. (2007). The relevance of phylogeny to studies of
global change. *Trends in Ecology & Evolution*, 22, 243-249.

Ehleringer J.R., Cerling T.E., Helliker B.R. (1997). C$_4$ photosynthesis, atmospheric CO_2, and
climate. *Oecologia*, 112, 285-299.

Ehleringer J.R., Monson R.K. (1993). Evolutionary and ecological aspects of photosynthesis
pathway variation. *Annual Review of Ecology and Systematics*, 24, 411-439.

Ehleringer J.R., Sage R.F., Flanagan L.B., Pearcy R.W. (1991). Climate change and the
evolution of C$_4$ photosynthesis. *Trends in ecology & evolution*, 6, 95-99.

Evans J.R. (1989). Partitioning of nitrogen between and within leaves grown under different
irradiances. *Functional Plant Biology*, 16, 533–548, doi:10.1071/PP9890533.

Evans J., Poorter H. (2001). Photosynthetic acclimation of plants to growth irradiance: the
relative importance of specific leaf area and nitrogen partitioning in maximizing carbon
gain. *Plant, Cell & Environment*, 24, 755-767.

Farquhar G.D., Von Caemmerer S., Berry J.A. (1980). A biochemical model of
photosynthetic carbon dioxide assimilation in leaves of 3-carbon pathway species.
Planta, 149, 78–90.

Fujita N., Miwa T., Ishijima S., Izui K., Katsuki H. (1984). The primary structure of
phosphoenolpyruvate carboxylase of Escherichia coli. Nucleotide sequence of the ppc
gene and deduced amino acid sequence. *The Journal of Biochemistry*, 95, 909-916.

Genty B., Briantais J.M., Baker N.R. (1989). The relationship between the quantum yield of
photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochimica*
et Biophysica Acta (BBA)-General Subjects, 990, 87-92.

Ghannoum O., Evans J.R., von Caemmerer S. (2010). Nitrogen and water use efficiency of
C_4 plants. In: C_4 photosynthesis and related CO_2 concentrating mechanisms (eds.
Raghavendra, A.S. & Sage, R.F.). Springer Science, The Netherlands, Dordrecht, pp.
129–146.

Grass Phylogeny Working Group II. 2012. New grass phylogeny resolves deep evolutionary
relationships and discovers C_4 origins. *New Phytologist*, 193, 304-312.

Griffith D.M., Osborne C.P., Edwards E.J., Bachle S., Beerling D.J., Bond W.J., Gallaher
T.J., Helliker B.R., Lehmann C.E., Leatherman L. and Nippert J.B. (2020). Lineage-
based functional types: characterising functional diversity to enhance the representation
of ecological behaviour in Land Surface Models. *New Phytologist*, 228, 15-23.

Hatch M.D. (1987). C_4 photosynthesis: a unique blend of modified biochemistry, anatomy
and ultrastructure. *Biochimica et Biophysica Acta (BBA)-Reviews on Bioenergetics*, 895,
81-106.

Heyduk K., Moreno-Villena J.J., Gilman I.S., Christin P.A., and Edwards E.J. 2019. The
genetics of convergent evolution: insights from plant photosynthesis. *Nature Review*
Genetics, 20, 485-493.

Kromdijk J., Long S.P. (2016). One crop breeding cycle from starvation? How engineering
crop photosynthesis for rising CO_2 and temperature could be one important route to
alleviation. *Proceedings of the Royal Society B: Biological Sciences*, 283, p.20152578.

Ku M.S.B., Schmitt M.R., Edwards G.E. (1979). Quantitative determination of RuBP
carboxylase-orygenase protein in leaves of several C_3 and C_4 plants. *Journal of*
experimental botany, 114, 89–98.

Kubien D.S., von Caemmerer S., Furbank R.T., Sage R.F. (2003). C_4 photosynthesis at low
temperature. A study using transgenic plants with reduced amounts of Rubisco. *Plant*
Physiology, 132, 1577-1585.

Kubien D.S., Sage R.F. (2004). Low-temperature photosynthetic performance of a C_4 grass
and a co-occurring C_3 grass native to high latitudes. *Plant, Cell & Environment*, 27, 907-
916.

Leakey A.D. (2009). Rising atmospheric carbon dioxide concentration and the future of C_4
crops for food and fuel. *Proceedings of the Royal Society B: Biological Sciences*, 276,
2333-2343. doi: 10.1098/rspb.2008.1517

Lundgren M.R., Dunning L.T., Olofsson J.K., MorenoVillena J.J., Bouvier J.W., Sage T.L.,
Khoshravesh R., Sultmanis S., Stata M., Ripley B.S., Vorontsova M.S. (2019). C_4
anatomy can evolve via a single developmental change. *Ecology letters*, 22, 302-312.

Morgan J.A., LeCain D.R., Pendall E., Blumenthal D.M., Kimball B.A., Carrillo Y.,
Williams D.G., Heisler-White J., Dijkstra F.A., and West M. (2011). C_4 grasses prosper
as carbon dioxide eliminates desiccation in warmed semi-arid grassland. *Nature,* 476,
202-206. doi: 10.1038/nature10274.

Munekage Y.N. (2016). Light harvesting and chloroplast electron transport in NADP-malic
enzyme type C_4 plants. *Current opinion in plant biology*, 31, 9-15.

Munekage Y.N., Taniguchi Y.Y. (2016). Promotion of cyclic electron transport around
photosystem I with the development of C_4 photosynthesis. *Plant and Cell*
Physiology, 57, 897-903.

Nakamura N., Iwano M., Havaux M., Yokota A., Munekage Y.N. (2013). Promotion of
cyclic electron transport around photosystem I during the evolution of NADP–malic
enzyme-type C_4 photosynthesis in the genus *Flaveria*. *New Phytologist*, 199, 832-842.

Niinemets Ü., Tenhunen J.D. (1997). A model separating leaf structural and physiological
effects on carbon gain along light gradients for the shade-tolerant species *Acer*
saccharum. *Plant Cell & Environment*, 20, 845–866.

Oaks A. (1994). Efficiency of nitrogen utilization in C_3 and C_4 cereals. *Plant Physiology*,
106: p.407.

Ögren E., Evans J.R. (1993). Photosynthetic light-response curves. *Planta*, 189, 182-190.

Onoda Y., Hikosaka K., Hirose T. (2005). The balance between RuBP carboxylation and
RuBP regeneration: a mechanism underlying the interspecific variation in acclimation of
photosynthesis to seasonal change in temperature. *Functional Plant Biology*, 32, 903-
910.

Osborne C.P., Sack L. (2012). Evolution of C_4 plants: a new hypothesis for an interaction of
CO_2 and water relations mediated by plant hydraulics. *Philosophical Transactions of the*
Royal Society B: Biological Sciences, 367, 583-600.

Pengelly J.J., Sirault X.R., Tazoe Y., Evans J.R., Furbank R.T., von Caemmerer S. (2010).
Growth of the C_4 dicot *Flaveria bidentis*: photosynthetic acclimation to low light through
shifts in leaf anatomy and biochemistry. *Journal of experimental botany*, 61, 4109-4122.

Pignon C.P., Long S.P. (2020). Retrospective analysis of biochemical limitations to
photosynthesis in 49 species: C_4 crops appear still adapted to pre-industrial atmospheric
[CO2]. *Plant, Cell & Environment*, 43, 2606-2622.

Pinto H., Powell J.R., Sharwood R.E., Tissue D.T., Ghannoum O. (2016). Variations in
nitrogen use efficiency reflect the biochemical subtype while variations in water use

efficiency reflect the evolutionary lineage of C_4 grasses at inter-glacial CO_2. *Plant, cell*

& environment, 39, 514-526.

Pinto H., Sharwood R.E., Tissue D.T., Ghannoum O. (2014). Photosynthesis of C_3, C_3–C_4,

and C_4 grasses at glacial CO_2. *Journal of Experimental Botany*, 65, 3669-3681.

Pittermann J., Sage R.F. (2001). The response of the high altitude C_4 grass *Muhlenbergia*

montana (Nutt.) AS Hitchc. to long-and short-term chilling. *Journal of Experimental*

Botany, 52, 829-838.

Porra R.J., Thompson W.A., Kriedemann P.E. (1989). Determination of accurate extinction

coefficients and simultaneous equations for assaying chlorophylls a and b extracted with

four different solvents: verification of the concentration of chlorophyll standards by

atomic absorption spectroscopy. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*,

975, 384-394.

Quebbeman J.A., Ramirez J.A. (2016). Optimal allocation of leaf-level nitrogen: Implications

for covariation of V_{cmax} and J_{max} and photosynthetic downregulation. *Journal of*

Geophysical Research: Biogeosciences,121, 2464-2475.

Rachmilevitch S., Cousins A.B., Bloom A.J. (2004). Nitrate assimilation in plant shoots

depends on photorespiration. *Proceedings of the National Academy of Sciences*, 101,

11506-11510.

Raines C.A. (2011). Increasing photosynthetic carbon assimilation in C_3 plants to improve

crop yield: current and future strategies. *Plant physiology*, 155, 36-42.

Reich P.B., Hobbie S.E., Lee T.D., Pastore M.A. (2018). Unexpected reversal of C_3 versus C_4

grass response to elevated CO_2 during a 20-year field experiment. *Science*, 360, 317-320.

doi: 10.1126/science.aas9313

Ripley B.S., Abraham T.I., Osborne C.P. (2008). Consequences of C_4 photosynthesis for the

partitioning of growth: a test using C_3 and C_4 subspecies of *Alloteropsis semialata* under

nitrogen-limitation. *Journal of experimental botany*, 59, 1705-1714.

Rogers A., Medlyn B.E., Dukes J.S., Bonan G., Von Caemmerer S., Dietze M.C., Kattge J.,

Leakey A.D., Mercado L.M., Niinemets Ü., Prentice I.C. (2017). A roadmap for

improving the representation of photosynthesis in Earth system models. *New Phytologist*,

213, 22-42. doi: 10.1111/nph.14283

Rodriguez-Calcerrada J., Reich P.B., Rosenqvist E., Pardos J.A., Cano F.J., Aranda I. (2008).

Leaf physiological versus morphological acclimation to high-light exposure at different

stages of foliar development in oak. *Tree physiology*, 28, 761-771.

Sage R.F. 2001. Environmental and evolutionary preconditions for the origin and
diversification of the C_4 photosynthetic syndrome. *Plant Biology*, 3, 202–213.

Sage R.F. (2004). The evolution of C_4 photosynthesis. *New Phytologist*, 161, 341–370.

Sage R.F. (2016). Tracking the evolutionary rise of C_4 metabolism. *Journal of experimental*
botany, 67, 2919-2922.

Sage R.F., McKown A.D. (2006). Is C_4 photosynthesis less phenotypically plastic than C_3
photosynthesis? *Journal of experimental botany*, 57, 303-317.

Sage R.F., Pearcy R.W. (1987). The nitrogen use efficiency of C_3 and C_4 plants: I. Leaf
nitrogen, growth, and biomass partitioning in Chenopodium album (L.) and *Amaranthus*
retroflexus (L.). *Plant Physiology*, 84, 954-958.

Schaefer K., Schwalm C.R., Williams C., Arain M.A., Barr A., Chen J.M., Davis K.J.,
Dimitrov D., Hilton T.W., Hollinger D.Y., Humphreys E. (2012). A model-data
comparison of gross primary productivity: Results from the North American Carbon
Program site synthesis. *Journal of Geophysical Research: Biogeosciences*, 117(G3).

Serrano-Romero, E.A., Cousins, A.B. (2020). Cold acclimation of mesophyll conductance,
bundle-sheath conductance and leakiness in *Miscanthus× giganteus. New*
Phytologist 226, 1594-1606.

Sharwood R.E., Sonawane, B.V., Ghannoum O. (2014). Photosynthetic flexibility in maize
exposed to salinity and shade. *Journal of Experimental Botany*, 65, 3715-3724.

Sharwood R.E., Ghannoum O., Whitney S.M. (2016). Prospects for improving CO_2 fixation
in C_3-crops through understanding C_4-Rubisco biogenesis and catalytic
diversity. *Current Opinion in Plant Biology*, 31, 135-142.

Smith N.G., Keenan T.F. (2020). Mechanisms underlying leaf photosynthetic acclimation to
warming and elevated CO_2 as inferred from least-cost optimality theory. *Global Change*
Biology, 26, 5202-5216.

Sonawane, B.V. (2016). Environmental regulation of CO_2 concentrating mechanisms in C_4
grasses with different biochemical subtypes (Doctoral dissertation, Western Sydney
University (Australia)).

Spriggs, E.L., Christin, P.A. and Edwards, E.J. (2014). C_4 photosynthesis promoted species
diversification during the Miocene grassland expansion. *Plos one*, 9, p.e97722.

Stata M., Sage T.L., Sage R.F. (2019). Mind the gap: the evolutionary engagement of the C_4
metabolic cycle in support of net carbon assimilation. *Current Opinion in Plant Biology*,
49, 27-34. doi: 10.1016/j.pbi.2019.04.008

Sundermann E., Lercher M. and Heckmann D. (2018). Modeling Cellular Resource

Allocation Reveals Low Phenotypic Plasticity of C_4 Plants and Infers Environments of

C_4 Photosynthesis Evolution. bioRxiv (Preprint). Available at

doi: https://doi.org/10.1101/371096 (Accessed April 15, 2020).

Takabayashi A., Kishine M., Asada K., Endo T., Sato F. (2005). Differential use of two

cyclic electron flows around photosystem I for driving CO_2-concentration mechanism in

C_4 photosynthesis. *Proceedings of the National Academy of Sciences*, 102, 16898-16903.

Tissue D.T., Griffin K.L., Thomas R.B., Strain B.R. (1995). Effects of low and elevated CO_2

on C_3 and C_4 annuals. II. Photosynthesis and leaf biochemistry. *Oecologia*, 101, 21-28.

Walker A.P., Beckerman A.P., Gu L., Kattge J., Cernusak L.A., Domingues T.F., Scales J.C.,

Wohlfahrt G., Wullschleger S.D., Woodward F.I. (2014). The relationship of leaf

photosynthetic traits–V_{cmax} and J_{max}–to leaf nitrogen, leaf phosphorus, and specific leaf

area: a meta-analysis and modeling study. *Ecology and evolution*, 4, 3218-3235.

Walker B.J., Drewry D.T., Slattery R.A, VanLoocke A., Cho Y.B., Ort D.R. (2018).

Chlorophyll can be reduced in crop canopies with little penalty to photosynthesis. *Plant*

Physiology, 176, 1215-1232.

Wang S., Tholen D., and Zhu X.G. (2017). C_4 photosynthesis in C_3 rice: a theoretical analysis

of biochemical and anatomical factors. *Plant, cell & environment*, 40, 80-94.

Ward J.K., Tissue D.T., Thomas R.B., Strain B.R. (1999). Comparative responses of model

C_3 and C_4 plants to drought in low and elevated CO_2. *Global Change Biology* 5: 857-

867.

Williams B.P., Johnston I.G., Covshoff S., Hibberd J.M. (2013). Phenotypic landscape

inference reveals multiple evolutionary paths to C_4 photosynthesis. *Elife*, 2, p.e00961.

doi: 10.7554/eLife.00961

Wittmer M.H., Auerswald K., Bai Y., Schaeufele R., Schnyder H. (2010). Changes in the

abundance of C_3/C_4 species of Inner Mongolia grassland: evidence from isotopic

composition of soil and vegetation. *Global Change Biology*, 16, 605-616. doi:

10.1111/j.1365-2486.2009.02033.x

Wullschleger S.D. (1993). Biochemical limitations to carbon assimilation in C_3 plants - A

retrospective analysis of the A/C_i curves from 109 species. *Journal of Experimental*

Botany, 44, 907-920.

Yin X., Van Der Putten P.E., Driever S.M., Struik P.C. (2016). Temperature response of

bundle-sheath conductance in maize leaves. *Journal of experimental botany*, 67, 2699-

2714.

Yin X., Schapendonk A.H., Struik P.C. (2018). Exploring the optimum nitrogen partitioning

to predict the acclimation of C_3 leaf photosynthesis to varying growth

conditions. *Journal of experimental botany*, 70, 2435-2447.

Yin X., Struik P.C. (2012). Mathematical review of the energy transduction stoichiometries

of C_4 leaf photosynthesis under limiting light. *Plant, cell & environment*, 35, 1299-1312.

Yin X., Struik P.C. (2018). The energy budget in C_4 photosynthesis: insights from a cell-

type-specific electron transport model. *New Phytologist*, 218, 986-998.

Yin X., Struik P.C., Romero P., Harbinson J., Evers J.B., Van Der Putten P.E., Vos J.A.N.

(2009). Using combined measurements of gas exchange and chlorophyll fluorescence to

estimate parameters of a biochemical C_3 photosynthesis model: a critical appraisal and a

new integrated approach applied to leaves in a wheat (*Triticum aestivum*) canopy. *Plant,*

Cell & Environment, 32, 448-464.

Yin X., Sun Z., Struik P.C., Van der Putten P.E., Van Ieperen W.I.M., Harbinson J. (2011).

Using a biochemical C_4 photosynthesis model and combined gas exchange and

chlorophyll fluorescence measurements to estimate bundle-sheath conductance of maize

leaves differing in age and nitrogen content. *Plant, cell & environment*, 34, 2183-2199.

Zaehle S., Sitch S., Smith B., Hatterman F. (2005). Effects of parameter uncertainties on the

modeling of terrestrial biosphere dynamics. *Global Biogeochemical Cycles* 19: GB3020.

Zhou H, Akçay E, Helliker BR. 2019. Estimating C_4 photosynthesis parameters by fitting

intensive A/C_i curves. *Photosynthesis research*, 141, 181-194.

Zhou H., Helliker B.R., Huber M., Dicks A., Akçay E. (2018). C_4 photosynthesis and climate

through the lens of optimality. *Proceedings of the National Academy of Sciences*, 115,

12057-12062.

Zhou H., Akçay E., Edwards E., and Helliker B.R. (2020). The legacy of C_4 evolution in the

hydraulics of C_3 and C4 grasses. bioRxiv.

doi: https://doi.org/10.1101/2020.05.14.097030.

Table 1 Phylogenetic results of the best-fitted models and their parameters for

photosynthesis parameters (summarizing Table S2-S10).

Property	Model	Model type	AICw	Root C$_3$	Root C$_4$
J_{max}/V_{cmax}	Model 6*	OU	1.000	1.59	5.90
V_{cmax}	Model 6*	OU	0.769	59.52	23.03
J_{max}	Model 6*	OU	0.583	88.94	132.16
Total Chl	Model 1	BM	0.293	0.36	
flr-ETR/J_{max}	Model 5*	OU	1.000	1.08	0.64
Chl a/b	Model 5	OU	0.995	3.40	4.61
flr-ETR	Model 1	BM	**0.397**	88.94	
V_{pmax}	Model 4	OU	0.854	41.52	
V_{pmax}/V_{cmax}	Model 4	OU	0.875	1.97	
N	Model 5*	OU	0.562	3.62	2.53

767 If the values for C$_3$ and C$_4$ were different, it meant there were significant different values for

768 C$_3$ and C$_4$ species (the evolutionary model with two different values of the root fit

769 significantly better than the evolutionary model with the similar root).

Figures

Fig. 1 Modeling results of assimilation rate with respect to maximal electron transport to maximal Rubisco carboxylation (J_{max}/V_{cmax}) and maximal PEP carboxylation to maximal Rubisco carboxylation (V_{pmax}/V_{cmax}) under CO_2 concentration of 200, 400 and 600 ppm. Other environmental conditions are soil water potential (ψ_s)=-0.5 MPa, VPD=0.625, temperature of 25 °C and light intensity of 2000 μmol m^{-2} s^{-1}. Left: 3D plot; right: corresponding contour plot.

Fig. 2 Empirical measurements (a) and optimal modeling results (b) of J_{max}/V_{cmax} for C3 and

C4 under ψ_S=-0.5 MPa, VPD=0.625, temperature of 25 °C and light intensity of 2000 μmol

m^{-2} s^{-1}, the cultivating environmental condition. In (b), the black line represents C3, solid red

line represents C4 modeling results with controlling V_{pmax}/V_{cmax} at the in vivo measurement

level, dashed red line represents C4 modeling results with controlling V_{pmax}/V_{cmax} at the in

vitro measurement level.

Fig. 3 Empirical measurements (a) and optimal modeling results (b) of V_{pmax}/V_{cmax} for C$_4$

under ψ_S=-0.5 MPa, VPD=0.625, temperature of 25 ^{o}C and light intensity of 2000 μmol m^{-2} s^{-1}

^{1}, the cultivating environmental condition. In (b), solid red line represents C$_4$ modeling results

with controlling J_{max}/V_{cmax} at the in vivo measurement level, dashed red line represents C$_4$

modeling results with controlling J_{max}/V_{cmax} at the in vitro measurement level.

Fig. 4 Modeling results of assimilation rate with varying J_{max}/V_{cmax} and V_{pmax}/V_{cmax} for C$_4$

under different CO_2 concentration and ψ_S=-1MPa, VPD=1.25, temperature of 25 $^{\circ}$C and light

intensity of 2000 μmol m^{-2} s^{-1}, the common grassland growth condition. Modeling results

were obtained by controlling the other parameter at the in vivo measurement level.

Fig. 5 Modeling results of changing V_{cmax} on assimilation rates for C_3 (acd) and for C_4 (bdf)

under different CO_2. Solid black line: the initial modeling condition of V_{cmax} (a typical C_3

value of 69 μmol m^{-2} s^{-1}); dashed black line: 90% of the initial V_{cmax}; dotted black line: 80%

of the initial V_{cmax}; solid grey line: 70% of the initial V_{cmax}; dashed grey line: 60% of the

initial V_{cmax}; dotted grey line: 50% of the initial V_{cmax}. C_3 and C_4 parameters shared similar

parameters except for the carbon concentration mechanism and different hydraulic

conductance to mimic the initial origin of C_4 (see Zhou et al (2020) for other parameters).

815 **Fig. 6** The regression for maximal assimilation rate (A_{max}), J_{max}, total cholorophyll (Total

816 chl), V_{cmax}, V_{pmax}, nitrogen concentration, J_{max}/V_{cmax}, V_{pmax}/V_{cmax} and fluorescence-estimated

817 electron transport to J_{max} (ETR/ J_{max}) vs. the evolutionary age for the nine origins to show the

818 evolutionary trend within C_4 (red) and within their closely-related C_3 species (black).

819
820
821

Fig. 7 Modeling results of optimal J_{max}/V_{cmax} and V_{pmax}/V_{cmax} for C_3 (black lines) and C_4 (red

lines) under different environmental conditions. (a) Solid line: different CO_2; dashed line:

different water limitation conditions (1: saturated water; 2: ψ_S=-0.5 MPa, VPD=0.625; 3:

ψ_S=-1 MPa, VPD=1.25 MPa; 4: ψ_S=-1.5 MPa, VPD=1.875; 5: ψ_S=-2 MPa, VPD=2.5). (b)

Solid line: different light intensities; dashed line: different temperature. Modeling results

were obtained by controlling the other parameter at the in vivo measurement level.