Selective B cell depletion upon infusion of replication-incompetent anti-CD19 CAR lentivirus

Craig M. Rive[a,1], Eric Yung[a,1], Lisa Dreolini[a], Daniel J. Woodsworth[a], Robert A. Holt [a,b,c,2]

[a] Canada's Michael Smith Genome Sciences Centre, BC Cancer, Vancouver, BC, V5Z 1L3, Canada.
[b] Department of Medical Genetics, University of British Columbia, Vancouver, BC, V6T 1Z4, Canada.
[c] Department of Molecular Biology & Biochemistry, Simon Fraser University, Burnaby, BC, V5A 1S6, Canada.
[1] Co-first author
[2] Correspondence: rholt@bcgsc.ca, Canada's Michael Smith Genome Sciences Centre, BC Cancer, Vancouver, British Columbia, Canada

Abstract

Anti-CD19 CAR-T therapy for B cell malignancies has shown clinical success, but a major limitation is the logistical complexity and high cost of manufacturing autologous cell products. Direct infusion of viral gene transfer vectors to initiate *in vivo* CAR-T transduction, expansion and anti-tumor activity could provide an alternative, universal approach for CAR-T and related immune effector cell therapies that circumvents *ex vivo* cell manufacturing. To explore the potential of this approach we first evaluated human and mouse $CD8^+$ T cells transduced with VSV-G pseudotyped lentivectors carrying an anti-CD19CAR-2A-GFP transgene comprising either an 1D3 or FMC63 anti-CD19 binding domain. To evaluate CD19 antigen-driven CAR-T proliferation *in vitro* we co-cultured transduced murine T cells with an excess of irradiated splenocytes. We observed markedly greater expansion over a 9 week period for CAR-transduced T cells compared to control T cells transduced with a GFP-only transgene cultured under the same conditions (mean fold expansion +/- SD above control: ID3-CD19CAR-GFP modified T cells, 12.2 +/- 0.09 (p < 0.002); FMC63-CD19CAR-GFP modified T cells 8.8 +/- 0.03 (p < 0.004)). CAR-T cells isolated at the end-point of these co-cultures showed potent B cell directed cytolytic activity *in vitro*. Next, we administered approximately 20 million replication-incompetent lentiviral particles carrying either ID3-CD19CAR-GFP, FMC63-CD19CAR-GFP, or GFP-only transgene to wild-type C57BL/6 mice by tail vein infusion and monitored the dynamics of immune cell subsets isolated from peripheral blood at weekly intervals. We saw emergence of a persistent CAR-transduced $CD3^+$ T cell population beginning at week 3-4 that reached a maximum of 13.5 +/- 0.58 % (mean +/- SD) and 7.8 +/- 0.76% of the peripheral blood $CD3^+$ T cell population in mice infused with ID3-CD19CAR-GFP lentivector or FMC63-CD19CAR-GFP lentivector, respectively. This was followed by a rapid decline in each group of the B cell content of peripheral blood. Complete B cell aplasia was apparent by week 5 and was sustained until the end of the protocol (week 8). None of these changes were observed in mice infused with GFP-only control lentivector, and significant CAR positive populations were not observed within other immune cell subsets, including macrophage, natural killer, or B cells. Modest weight loss of 5.5 +/- 2.97 % (mean +/- SD) was observed in some animals receiving an anti-CD19CAR-GFP transgene during the protocol. These results indicate that direct IV injection of lentiviral particles carrying an anti-CD19 CAR transgene can transduce T cells that then fully ablate endogenous B cells in wild type mice. Based on these results it may be useful to further explore, using existing vectors, the feasibility of systemic gene therapy as a modality for CAR-T and related genetically-engineered Immune Effector Cell therapies.

Keywords: CAR-T, Universal CAR, Immune Effector Cells, Gene transfer, Gene therapy, Leukemia, Lymphoma, Cancer

Introduction

Ordinarily, a cancer patient's immune system can recognize cancer cells as being altered, and mount a natural immune response. T cells are known to be the main mediators of the anti-cancer immune response but sometimes the natural T cell response is ineffective; T cells may fail to recognize tumor cells, fail to activate, or fail to sustain a response long enough to have an impact. It is possible for these limitations to be overcome by genetic engineering strategies that involve isolating a sample of a patient's T lymphocytes, genetically modifying and activating the cells *ex vivo*, and then re-administering them to the same patient. The genetic modification step involves introducing into the T cells an extra gene that carries instructions for a new antigen receptor that may be a recombinant alpha-beta T cell receptor (αβTCR) or a Chimeric Antigen Receptor (CAR).

Currently, most genetically engineered Immune Effector Cell (IEC) therapies require several steps including i) manufacturing plasmid encoding the synthetic transgene and ancillary plasmids, ii) delivery of these plasmids into producer cells to generate virus, iii) collection of patient autologous T cells and

integration of the transgene into the genomes of these cells by viral transduction, iv) expansion of the transduced T cells *ex vivo* to produce the therapeutic cells, v) extensive release testing of the manufactured product, vi) preconditioning lymphodepletion of the patient, vii) infusion of the modified autologous cells, and viii) intensive clinical and immunological monitoring. Thus, there is considerable infrastructure and expertise required to deliver IEC treatments safely and successfully. The complexity of manufacturing IECs is a critical limitation to the cell therapy field, and a major contributor to the high cost of reimbursement of new cell-based interventions coming to market.

Much of the growing interest in genetically engineered IECs as a new class of therapeutic is driven by the impressive results CAR-T therapies have shown in patients with relapsed/refractory B-cell Acute Lymphoblastic Leukemia or Non-Hodgkin Lymphoma. These patients, for whom standard therapies have failed, have historically had dismal outcomes with survival measured in weeks or months. Remarkably, CAR-T therapy targeting CD19 has led to meaningful and durable remissions for large proportion of these patients in many different studies[1–7].

Ideally, to reduce the logistical burden of IEC therapies, a master allogeneic cell bank would be manufactured centrally and distributed to clinical sites for infusion as a "universal" cell product. For this approach it would be necessary to source large numbers of primary, non-immortalized cells from allogeneic donors. Many donors would be needed because there is a limit to the number of cells that can be obtained per donor, resulting in non-homogeneous products susceptible to batch-to-batch variation. Of course, allogeneic therapies also face the fundamental limitations of susceptibility to graft rejection (caused by recognition of allo-antigens on donor cells by the recipient's endogenous T cells) and graft-versus-host-disease (caused by systemic reactivity of donor cells to the recipient's allo-antigens). Despite these major hurdles to universal cell therapies, the manufacturing constraints associated with personalized autologous products are so severe that "AlloCAR-T" therapy has become an area of intense interest. To engineer universal IECs one approach has been to knockout the endogenous TCR in the T cell product to attenuate anti-host activity, and to delete receptors in the cell product for immunosuppressant drugs such as alemtuzumab (anti-CD52) to confer resistance, such that those drugs can then be administered to the recipient to selectively inhibit host immunity and prevent graft rejection, but not inhibit the function of the grafted allogeneic cells. Indeed some successes have been demonstrated in pre-clinical work[8] and in case studies[9,10] with gene-edited allo-CAR-Ts. Likewise, the immortalized IEC cell line NK-92 has shown encouraging safety and efficacy in a recent phase 1 trial, but the requirement of irradiating the cells pre-infusion to reduce oncogenic risk will likely remain limiting for this approach.[11] The generation of a bank of allogeneic CAR-T cells by *in vitro* differentiation and maturation of hematopoietic stem and progenitor cell precursors is an alternative and highly compelling strategy. Demonstrated in principle, this approach awaits further optimization.[12]

The best approach to universal IEC therapy may be to sidestep cell manufacturing entirely and infuse gene transfer vector directly in order to transduce T cells *in vivo*, redefining these treatments as gene therapies rather than cell therapies. The ability to safely administer gene transfer vectors directly, as is done for other modes of gene therapy[13–16] would be transformative for cancer immunotherapy.

In vivo gene delivery has previously been attempted, preclinically, for CAR-T therapy. Nanoparticles have been demonstrated to be able to carry a CAR transgene payload and transduce CD3[+] cells, but this required flanking piggyBAC sequences and expression of a transposase to enable persistent CAR expression [17] and resulted in modification of non-T cells as well as T cells. In another recent study lentivirus modified to specifically target CD8[+] T cells was engineered and tested for transfer of CAR constructs *in vivo* [18]. Transduction was successful, but with room for improvement in efficiency and specificity. Here, we describe the initial experiments we performed to test *in vivo* transfer of a CAR transgene to murine T cells using a conventional, clinically compatible lentiviral vector.

Results

During our early pre-clinical development of an anti-CD19 CAR intervention for B cell malignancies we explored how CAR-T cells behave *in vitro* in the context of antigen driven expansion. Using a standard 2nd generation anti-CD19 CAR design[19] (FMC63 scFv CD19 binding domain, CD8α hinge and transmembrane domain, 4-1BB co-stimulatory domain, CD3z signaling domain and 2A-GFP tag) and standard VSV-G pseudotyped lentivector, we transduced healthy donor human Peripheral Blood Mononuclear Cells (PBMC) *in vitro* and cultured these for nine weeks. We observed that CAR-modified T-cells expand rapidly and selectively relative to GFP transduced T cells when stimulated with irradiated allogeneic PBMC as a source of CD19 antigen[**Figure 1A**]. Expanded CAR-T cells showed robust CD19-dependent target cell killing *in vitro* with 66.0 +/- 0.14%; (mean +/- SD) target cell death at 1:1 effector:target ratio, 90.2 +/- 2.62% target cell death at 4:1 effector:target ratio, and 94.4 +/- 0.35% target cell death at 10:1 effector:target ratio [**Figure 1B**].

These early observations showing that B cells are able to drive robust expansion of functional CD19 CAR-T cells *in vitro*, together with the recent reports on T cell modification *in situ* noted above[17,18] were encouraging and prompted us to explore the potential for *in vivo* transduction and expansion of CAR-T cells using a murine model system. We constructed a CAR where we changed the scFv binding domain from FMC63[20] (a monoclonal antibody raised against human CD19) to 1D3[21] (a monoclonal antibody raised against mouse CD19). 2.5×10^5 splenocytes isolated from wildtype C57BL/6 mice were transduced with lentivirus encoding either the 1D3-CD19CAR-GFP (mouse), FMC63-CD19CAR-GFP (human) or GFP-only transgene then stimulated with a further excess of irradiated autologous murine splenocytes as a source of murine CD19 antigen. We observed that under these conditions murine T-cells modified with either of the CARs expanded rapidly and selectively. For murine T cells modified with the human CAR expansion was 8.8+/- 0.03 (mean +/- SD) fold higher at day 63 than GFP transduced T cells (p < 0.004 paired, two-tail T test), and for murine T cells modified with the murine CAR expansion was 12.9 +/- 0.09 fold higher at day

63, relative to GFP transduced T cells (p < 0.002 paired, two-tail T test). The murine control T cells transduced with GFP-only lentivirus showed limited but observable expansion, possibly been driven by the original CD3/CD28 activation prior to transduction [**Figures 2A and 2B**]. This is contrast to human GFP-only control T cells (Figure 1A) which were not CD3/CD28 activated prior to transduction, and did not expand in subsequent co-culture (Materials and Methods). Expanded murine CAR-T cells showed robust and selective killing of murine B cells *in vitro*. The 1D3-CD19CAR-GFP modified murine T cells killed 24.8 +/- 1.69% (mean +/- SD) of murine B cells in the 5 hour co-culture and the FMC63-CD19CAR-GFP modified murine T cells killed 18.5 +/- 1.08% of murine B cells in the 5 hour co-culture [**Figure 2C**].

Interestingly, this result indicates that the FMC63 scFv is able to bind mouse as well as human CD19, to the degree necessary for CAR-T activation in this experimental system. To our knowledge this has not previously been reported.

We then administered lentivirus encoding each CAR or GFP-only lentivirus to C57BL/6 wild type mice (age 4-6 weeks) by tail vein injection, and drew peripheral blood at regular intervals. We observed that GFP-expressing T cells appear in the blood of mice 3 weeks after the initial injection of $1.8\text{-}2 \times 10^7$ infectious units (IU) of either the 1D3- or FMC63-CD19CAR-GFP lentivirus, but not after injection of the same dose of GFP-only lentivirus [**Figures 3A and 3B**]. Specifically, we saw emergence of a persistent 1D3-CD19CAR-GFP transduced CD3+ T cell population beginning at week 3 that reaching a maximum of 13.5 +/- 0.58% (mean +/- SD) of the peripheral blood CD3+ T cell population by week 5 [**Figures 3A and 4A**]. This coincided with the loss of B cells, seen 5 weeks after the initial injection [**Figures 3A and 4B**]. In experiments where we infused murine T cells modified with the FMC63-CD19CAR-GFP construct we also saw the emergence of a persistent FMC63-CD19CAR-GFP transduced CD3+ T cell population beginning at week 5 that reached a maximum of 7.8 +/-0.76% of the peripheral blood CD3+ T cell population by week 5 [**Figures 3A and 4B**]. Again, this coincided with a loss of B cells, seen 5 weeks after the initial injection. This further confirms functional recognition of mouse

CD19 by FMC63 **[Figures 3A and 4B]** . With either CAR, B cell aplasia was sustained until the end of the protocol (week 8). Mice treated with 1.8-2 x10^7 IU of the GFP control lentivirus had no GFP expressing T cells and did not show the same B cell loss.

To assess the ability of lentiviral injections to transduce PBMCs other than T cells, we looked for GFP expression in different immune cell subsets including the CD20/CD45R/B220 B cell population, CD11b$^+$ macrophage, and CD335$^+$ natural killer (NK) cells. We found that only T cells expressed the reporter GFP gene, indicative of CAR expression, and our interpretation is that interaction between the *in vivo* transduced CAR-T cells and endogenous B cells drives CAR-T cell expansion and CAR-T mediated B cell depletion, resulting eventually in B cell aplasia [**Figure 4**]. The only overt change observed in mice treated with either 1D3- or FMC63-CD19CAR lentivirus was a modest decrease in weight of 5.5 +/- 2.97% (mean +/- SD) and 3.6 +/- 1.78%, respectively over the first week after treatment [**Figure 5**].

Discussion

Here we report the observation that upon intravenous infusion, replication-incompetent lentiviral particles carrying an anti-CD19 CAR transgene can effectively transduce T cells, leading to CAR-T proliferation and B cell aplasia. This supports the notion that CAR-T cells generated in this manner could also have anti-tumor activity in the setting of B cell malignancies, with limited toxicity. If further testing indicates this is the case, then the potential of this protocol for "universal CAR-T therapy" warrants careful consideration. From our perspective there are many issues to explore when considering systemic gene therapy as an alternative to *ex-vivo* manufacturing of CAR-T cells, and these factors are inter-related:

Efficiency and target cell specificity. For clinically relevant anti-tumor efficacy, a sufficient number of target T cells needs to be encountered by virus, successfully transduced, and reliably and persistently express the receptor transgene at physiologically relevant levels. Most lentiviral gene transfer vectors used clinically, including the vector we use here, are pseudotyped using vesicular stomatitis virus G

glycoprotein (VSV-G). VSV-G binds the LDL receptor expressed on the surface of most cell types including T cells. Upon intravenous (IV) injection of virus, T cells plus all other hematopoietic cells and vascular endothelial cells are susceptible to transduction. While successfully transduced T cells expressing a CAR may undergo antigen driven expansion *in vivo*, bystander cells can be problematic if the act as viral sinks. Thus, it will be necessary to define empirically the optimal viral dosing regimen in this setting. Further implications to the transduction of bystander cells are unique to the setting of anti-CD19 CAR-T therapy. For example, expression of the CAR transgene in malignant B cells by inadvertent transduction of tumor blasts is problematic, because the CAR can bind CD19 antigen on the same cell and block engagement by cytolytic CAR-T cells. This is a known but rare mechanism of relapse in conventional *ex vivo* protocols [22] and could be exacerbated by *in vivo* transduction. Another consideration is that while receptor transgene expression in CD8$^+$ cytolytic T cells and CD4$^+$ helper T cells is desirable,[23] expression in regulatory T cells may be lead to inhibition rather than activation of immunoreactivity.

Interestingly, murine T cells have previously been demonstrated to have lower transduction efficiencies, compared to human T cells and compared to other murine cell types[24–27], when transduced with HIV-1 based lentiviral vectors. The viral vector system we describe here avoids some of the known barriers of murine T cells. The entry block is avoided by the use of VSV-G pseudotyped virus. Reduced Tat activation is avoided entirely by our use of a Tat-independent transfer vector with a chimeric Rous Sarcoma virus LTR. Reduced LTR promoter activity is compensated by the use of an internal EF1α promoter driving the transgene. Some reduction in lentiviral integration can still be expected in murine T cells due to inefficient nuclear import of the pre-integration complex.[26] Empirically, however, we find that while our lentiviral vectors transduce murine T cells less efficiently than they transduce human T cells **[Figures S1]**, the level of transduction of murine T cells *in vitro* and *in vivo* is sufficient.

Immunogenicity. A well recognized issue that can limit the utility of systemic gene therapy is the inherent immunogenicity of viral gene transfer vectors. For example, pre-existing immunity to adenoviral vectors is common in the population due the high incidence of exposure to natural adenoviruses. Other viruses, such as Maloney Murine Leukemia Virus (MMLV) and adeno-associated viruses have limited pre-existing immunity but are immunogenic when used as gene transfer vectors. Pre-existing immunoreactivity to lentivirus is generally absent in the population, and acute immunoreactivity to infused CAR-bearing virus is mitigated by CAR-T mediated depletion of the B cell compartment including the potentially anti-lentiviral antibody producing B cells. These dynamics will be important to evaluate but we do not expect immunogenicity will be limiting given that IEC therapies are typically single dose administrations. It is already well established that patients who respond to CAR-T therapy do so after a single infusion, and that re-infusing a non-responsive patient does not provide significant benefit.[28]

Safety. Insertional mutagenesis of integrating viral vectors led to highly regrettable negative outcomes in early gene therapy trials and severe setbacks to the gene therapy field.[29–33] However, these interventions were developed to treat rare genetic immunodeficiencies that required modification of $CD34^+$ hematopoietic stem cells, and oncogenesis was due to insertional mutagenesis specifically within the hematopoietic stem cell compartment. Conversely, no detrimental effects of insertional mutagenesis have been reported from any CAR-T clinical trial using retroviral vectors to date. Some but not all CAR-T protocols include a step to purify $CD4^+/CD8^+$ T cells before *ex vivo* transduction, which minimizes risk. For *in vivo* transduction, where T cells cannot be segregated, a theoretical risk of insertional mutagenesis remains and further highlights the importance of target cell selectivity.

The use of HIV-based lentivirus for gene transfer also raises concern about the theoretical risk of transmission of replication competent lentivirus (RCL). We use a modified 2nd generation lentiviral vector system with the minimal viral components included in the 3rd generation system split across three plasmids instead of four. These alterations abolish the potential for recombination events that can lead to formation of RCL in target cells. Testing of hundreds of cell therapy products generated with third generation vectors as well screening clinical trial participants after infusion of cell product failed to detect RCL, [34,35] demonstrating the safety of lentiviral gene therapy in this regard.

Patient conditioning. CAR-T protocols and other clinical IEC protocols typically include lymphodepleting chemotherapy with cyclophosphamide and/or fludarabine as pre-conditioning. While it is not yet known exactly how this facilitates response, it is thought to improve engraftment by depleting endogenous T effector cells that compete for resources, and deplete regulatory T cells that may otherwise suppress the infused CAR-T population. While it would not be productive to perform complete lymphoablation prior to a viral infusion protocol (because endogenous T cells must be present for virus to transduce), mild preconditioning may be beneficial, such that depletion leaves enough endogenous T cells in circulation to support viral transduction and conversion to CAR-Ts, while at the same time providing space for robust CAR-T expansion and engraftment. This could be further tested, but may be unnecessary given our results show full ablation of endogenous B cells after lentiviral infusion **[Figure 2]** without any pre-conditioning lymphodepletion.

In summary, anti-CD19 CAR-T therapy for B cell malignancies has shown clinical success, but a major limitation is the logistical complexity and high cost of manufacturing autologous cell products. This has led to increasing interest in industry and academia in universal (non-personalized) strategies. Here we have shown that direct IV injection of lentiviral particles carrying the anti-CD19 CAR transgene can transduce T cells that then fully ablate endogenous B cells in wild type mice, indicating that direct infusion of viral gene transfer vectors to initiate *in vivo* T cell transduction, expansion and anti-tumor activity is a potential universal approach.

Materials and Methods

Media. RPMI-1640 supplemented media (sRPMI-1640) consisted of 2 mM GlutaMAX (Invitrogen, 35050-061), 1 mM MEM non-essential amino acid (Invitrogen, 11140-050), 1 mM sodium pyruvate (Invitrogen, 11360-070), 10 mM HEPES (Invitrogen, 15630-808), 100 U/mL penicillin- streptomycin (Invitrogen, 15140-122), MycoZap prophylactic (Lonza, VZA-2031), 0.05mM 2-Beta Mercaptoethanol (Sigma, 60-42-2, 2-βME added to media for mouse cell cultures only) and 10% heat-inactivated fetal bovine serum (Invitrogen, 12484-028). For expansion cultures IL-2 (StemCell, 78036.3) was added to the sRPMI-1640 media. Cells were cultured in sRPMI-1640 with a final IL-2 concentration of 300IU/ml, unless otherwise stated. FACS media consisted of 1x D-PBS (Sigma, 59331C) with 2% HI-FCS. In house MACS separation buffer consisted of D-PBS with 2mM EDTA (ThermoFisher, 15575020) and 2% HI-FBS.

Production of lentiviral gene transfer vectors. Our lentivector system is a modified 2^{nd} generation system that uses three separate plasmids to split the lentiviral genome as previously described.[36,37] The system is comprised of a packaging plasmid encoding gag, pol and rev, an envelope expressing plasmid and a transfer plasmid encoding the transgene. Biosafety of the 2^{nd} generation design was improved by removal of tat from the packaging plasmid with the concurrent replacement of the transfer vector 5' Long terminal repeat (LTR) with chimeric sequence for Tat-independent expression and a self-inactivating modified 3' LTR.[38] These modifications serve to prevent the formation of replication competent virus.

Synthetic transgene sequences encoding 1) GFP, 2) FMC63-CD19CAR-2A-GFP and 3) 1D3-CD19CAR-2A-GFP were manufactured (ATUM), subcloned into the transfer plasmid and verified by Sanger sequencing (Genewiz). To manufacture virus particles, envelope, packaging, and transfer plasmid were co-transfected into the packaging cell line HEK-293T (clone 17; ATCC) using TransIT-LT1 (Mirus, MIR2305). Media containing the lentivirus was collected after transfection, filtered and then ultracentrifuged at 25,000 rpm or 106,750 xg, for 90 minutes at 4˚C (Optima XE-90, Beckman-Coulter with SW 32 Ti swinging bucket rotor) to pellet the virus. Viral pellets were resuspended in 1x D-PBS and aliquoted for long term storage at -80C. Functional viral titers were established by infection of K562 cells (ATCC) and EL4 cells (ATCC) determined by transducing $5x10^4$ EL4 or K562 cells (ATCC) with 1, 2, 4, 8, 16, or 32 µL of concentrated viral supernatant. Performed in duplicate, cells were suspended in a final volume of 500uL of sRPMI-1640 media, in the 24-well format. Cells were then incubated for 72 hours, before being washed, centrifuged, and resuspended in FACS media. Acquisition was performed on the BD LSRFortessa cell analyser, and analysed using FlowJo and GraphPad Prism software version 8.0.0 for windows (GraphPad Software, California, USA) [**Figure S1** and **Figure S2**].

Sample acquisition and storage. Human PBMCs were isolated by Ficoll gradient purification as per manufactures protocol (Ficoll, GE-17-1440, GE Healthcare) from human peripheral blood leukapheresis pack (StemCell). 100-150µl of C57BL/6 blood was collected in lithium heparin capillary tubes via tail vein blood sampling pre-treatment and then every week post-treatment for 8 weeks. The blood samples were diluted with 1 - 1.5 ml of cold RPMI, which was overlayed with 4ml of Ficoll and C57BL/6 PBMCs were isolated following the manufacturer's protocol. C57BL/6 PBMCs isolated from mice were analysed within 24 hours of isolation and were resuspended in sRPMI-1640 at a concentration of $1x10^6$ cells per ml, placed into a T25 culture flask (156367, ThermoFisher), and rested overnight in the cell incubator at 37˚C with 5% CO_2 airflow before being used in any analysis. Human PBMCs and C57BL/6 PBMCs not immediately used in any experiments were stored down by resuspending cells in freezing media (HI-FBS with 10% DMSO, Fisher Scientific, BP231-100) in aliquots of $5-10x10^6$ cells per ml and frozen down to -80˚C at a speed of 1˚C/min prior to storage in liquid nitrogen.

The C57BL/6 splenocytes used in the *in vitro* assays were acquired from naive or untreated C57BL/6 mice, aged 4-8 weeks old. All C57BL/6 were however processed the same way by first manually dissociating

the spleens and filtering them through a 70 µm cell strainer (Corning, CLS431751) into a 50mL falcon tube, centrifuging 400xg for 10 min (Eppendorf Centrifuge 5810R, rotor A-4-62) at room temperature to pellet the cells. The splenocyte samples were resuspended in 5mL of ACK (Ammonium-Chloride-Potassium) Lysing Buffer (Thermofisher), before incubating the cells on ice for no longer than 4 minutes. 10 mL of sRPMI-1640 media was added to the cells to neutralize the ACK buffer and wash the splenocytes. The splenocytes were then centrifuged at 400xg for 10 min (Eppendorf Centrifuge 5810R, rotor A-4-62) at room temperature, the supernatant was removed, and the splenocytes washed again before being resuspended in sRPMI-1640 media and cell numbers obtained using the Countess automated cell counter (Thermofisher). Splenocytes were stored down by resuspending cells in freezing media (HI-FBS with 10% DMSO, Fisher Scientific, BP231-100) in aliquots of $5-10x10^6$ cells per ml and frozen down to -80˚C at a speed of 1˚C/min prior to storage in liquid nitrogen.

In vitro FMC63-CD19CAR-GFP modified T cell expansion and testing using human PBMC. $5x10^5$ Human PBMC were rested in RPMI-1640 supplemented media at $1x10^6$ cells/mL with 300IU/ml of IL-2 for 24 hours and then transduced at a MOI of 5 (calculated based on titration data) with either FMC63- or GFP-lentivirus. Transduced cells were then incubated for 48 hours before cells were washed with RPMI-1640 supplemented media, centrifuged at 300 x g for 5 mins at room temperature (Eppendorf centrifuge 5810R) and resuspended in RPMI-1640 supplemented media with 300IU/ml of IL-2 and $5x10^7$ irradiated allogeneic PBMC (50Gy). Cells were cultured with 50% of the media being removed and replaced with fresh media every third day, or split to maintain cells at $1x10^6$ cells/mL. Cells were counted using a Countess automated cell counter (Thermofisher), and aggregate counts were determined on day 24 and 63[**Figure 1A**].

To assess the cytolytic activity of FMC63-CD19CAR-GFP, $1x10^5$ cells expanded from transduced human PBMCs (effectors) were labelled with cell proliferation dye eFluor 450 (eBiosciences) and co-cultured with sRPMI-1640 in 5mL Falcon round-bottom polystyrene tubes (Fisher) for 4 hours

with CD19[+] 721.221 (ATCC) or CD19[-] K562 (ATCC) cells (targets) using $1x10^5$ (1:1), $4x10^5$ (4:1) or $1x10^6$ (10:1) target cells [**Figure 1B**]. Prior to the co-culture, the target cells (721.211 and K562 cells) were stained with the cell proliferation dye eFluor 670 (eBiosciences). At the end of the co-cultures, the cells were stained with DAPI (Thermofisher). Flow cytometry data acquisition was performed using the BD LSRFortessa cell analyser with BD FACSDiva software and the analysis was performed using FlowJo software, version 10 for Windows GraphPad Prism software version 8.0.0 for windows (GraphPad Software, California, USA) [**Figure S3**].

In vitro 1D3- and FMC63-CD19CAR-GFP modified T cell expansion and testing using C57BL/6 splenocytes. C57BL/6 splenocytes were removed from liquid nitrogen storage, thawed, and resuspended in sRPMI-1640 with 300 IU/ml of IL-2, and 30ng/ml of CD3 (clone 145-2C11, 100314, Biolegend) and 30ng/ml CD28 (clone 37.51, 102112, Biolegend) antibodies, at a concentration of $1x10^6$ cells/ml. The splenocytes were then treated with 1D3- CD19CAR-GFP, FMC63-CD19CAR-GFP, or GFP-only viral supernatant at a MOI of 5 (calculated based on titration data) and incubated for 72 hours. Following transduction, cells were washed by resuspending the cells in sRPMI-1640, centrifuging them at 400xg for 10 mins at room temperature (Eppendorf Centrifuge 5810R, rotor A-4-62), removing the supernatant, and repeating the process a further 2 times. The cells were resuspended in sRPMI-1640 media containing 300IU/ml of IL-2 at a concentration of $2.5x10^5$ cells per ml, with 1 ml being aliquoted per well, in the 6-well culture plate format. Irradiated autologous murine splenocytes (50Gy) were resuspended in sRPMI-1640 media containing 300IU/ml of IL-2 at a concentration of $2.5x10^7$/ml and 1 ml was added to the transduced splenocytes giving a final ratio of 1:100 transduced splenocytes to irradiated splenocytes. Every third day, 1ml of the culture media was removed and 1ml of fresh sRPMI-1640 media with 600IU/ml of IL-2 was added (giving a final concentration of 300IU/ml IL-2). On days 0, 14, 28, and 63, the cells were counted using the Countess automated cell counter (Thermofisher) and an aliquot of $3x10^4$ cells were stained with fluorescent

labeled antibodies and analyzed using flow cytometry [**Figure 2A and Figure 2B**].

In a 96-well plate format with a final volume of 200µl (sRPMI-1640), 1×10^5 1D3-, FMC63-CD19CAR, or GFP-transduced T cells from day 63 of the expansion were co-cultured for 5 hours with 1×10^5 B cells isolated from another sample of C57BL/6 splenocytes, using the Miltenyi MACS mouse CD19 B-cell isolation beads, following the manufacturer's protocol and using a MACS separation buffer made in-house. At the end of the co-culture, the cells were stained with fluorescent labeled antibodies and analyzed using flow cytometry [**Figure 2C**]. Cells were washed with 500µl of D-PBS and centrifuged at 400xg for 10 minutes at room temperature (Eppendorf Centrifuge 5810R, rotor A-4-62). The supernatant was removed and the cell pellet resuspended in 50µl of D-PBS before being stained with CD20-PE (clone A1SB12, 12-0201-82, eBioscience), CD3-efluor 450 (clone 17A2, 48-0032-82, eBioscience), and the live/dead marker, FVS780 (565388, BD Biosciences). Cells and antibodies were incubated at 4ºC for 30 min, then washed by adding 2mL of FACS media to each tube and centrifuging for 10 minutes at 400xg at 4˚C followed by removal of the supernatant before resuspension in 500µl of cold FACS media. Flow cytometry acquisition was performed using the BD LSRFortessa cell analyser with BD FACSDiva software. The analysis was performed using FlowJo software, version 10 for windows and GraphPad Prism software version 8.0.0 for windows (GraphPad Software, California, USA) [**Figure S4**].

In vivo testing. All animal experiments were assessed and approved by the University of British Columbia's Animal Care Committee under ethics certificate #A17-0107. For the *in vivo* testing, male C57BL/6 mice, aged 4 to 6 weeks old, were obtained from the BC Cancer Research Centre's Animal Resource Centre. The experimental mice received one tail vein infusion on day 1 which involved the injection of 200µl of D-PBS containing $1.8-2 \times 10^7$ total infectious units of either 1D3-, FMC63-CD19CAR-GFP, or GFP-only lentiviruses. Mice were monitored closely for the first 72 hours, and then once a day until the experimental endpoint. The mice were also weighed periodically and

blood collected in lithium heparin capillary tubes via tail vein sampling as previously described. At the experimental endpoint, the mice were humanly euthanized and spleen and blood tissue collected and processed as previously described. The cells were stained with fluorescent labeled antibodies and analyzed using flow cytometry [**Figure 3 and Figure 4**]

For flow cytometry acquisition and analysis, 1×10^5 cells were stained with CD3-efluor 450 (clone 17A2, 48-0032-82, eBioscience), CD90.2-AF700 (clone 30-H12, 105319, Biolegend), CD20-PE (clone A1SB12, 12-0201-82, eBioscience), CD45R/B220-APC-CY7 (clone RA3-6B2, 103223, Biolegend), CD335-APC (clone 29A1.4, 137608, Biolegend), and CD11B-BV605 (clone M1/79, 101237, Biolegend) conjugated flow antibodies. Cells and antibodies were incubated at 4ºC for 30 min, then washed by adding 2mL of FACS media to each tube and centrifuging for 10 minutes at 400xg at 4˚C, discarding the supernatant before resuspension in 500µl of cold FACS media. Acquisition was performed on the BD LSRFortessa cell analyser, and analysed using FlowJo and GraphPad Prism software version 8.0.0 for windows (GraphPad Software, California, USA) [**Figure S5**].

For all flow analyses, we used the CD20-PE antibody as a B cell marker in place of a CD19 antibody (**Figure 2, Figure 3, Figure 4**) to avoid any possible interference between anti-CD19 antibody and anti-CD19 CARs in these mixed cell populations.

Author Contributions and Notes

R.A.H, C.M.R and E.Y. designed the research, C.M.R, E.Y, L.D., and D.J.W. performed the research, C.M.R analyzed the data; all authors interpreted the results and wrote the manuscript.
The authors declare no conflict of interest.

Acknowledgments

This research was supported by the British Columbia Cancer Foundation, the Leon Judah Blackmore Foundation and the BioCanRx network.

References

1.   Neelapu, S. S. *et al.* Axicabtagene ciloleucel CAR T-cell

therapy in refractory large B-Cell lymphoma. *N. Engl. J. Med.* **377**, 2531–2544 (2017).

2. Maude, S. L. *et al.* Tisagenlecleucel in children and young adults with B-cell lymphoblastic leukemia. *N. Engl. J. Med.* **378**, 439–448 (2018).

3. Park, J. H. *et al.* Long-term follow-up of CD19 CAR therapy in acute lymphoblastic leukemia. *N. Engl. J. Med.* **378**, 449–459 (2018).

4. Hay, K. A. *et al.* Factors associated with durable EFS in adult B-cell ALL patients achieving MRD-negative CR after CD19 CAR T-cell therapy. *Blood* **133**, 1652–1663 (2019).

5. Hirayama, A. V *et al.* High rate of durable complete remission in follicular lymphoma after CD19 CAR-T cell immunotherapy. *Blood* **134**, 636–640 (2019).

6. Schuster, S. J. *et al.* Chimeric antigen receptor T Cells in refractory B-Cell lymphomas. *N. Engl. J. Med.* **377**, 2545–2554 (2017).

7. Grigor, E. J. M. *et al.* Risks and Benefits of Chimeric Antigen Receptor T-Cell (CAR-T) Therapy in Cancer: A Systematic Review and Meta-Analysis. *Transfusion Medicine Reviews* **33**, 98–110 (2019).

8. Sommer, C. *et al.* Preclinical Evaluation of Allogeneic CAR T Cells Targeting BCMA for the Treatment of Multiple Myeloma. *Mol. Ther.* **27**, 1126–1138 (2019).

9. Qasim, W. *et al.* Molecular remission of infant B-ALL after infusion of universal TALEN gene-edited CAR T cells. *Sci. Transl. Med.* **9**, (2017).

10. Brudno, J. N. *et al.* Allogeneic T Cells That Express an Anti-CD19 Chimeric Antigen Receptor Induce Remissions of B-Cell Malignancies That Progress After Allogeneic Hematopoietic Stem-Cell Transplantation Without Causing Graft-Versus-Host Disease. *J. Clin. Oncol.* **34**, 1112–21 (2016).

11. Williams, B. A. *et al.* A phase I trial of NK-92 cells for refractory hematological malignancies relapsing after autologous hematopoietic cell transplantation shows safety and evidence of efficacy. *Oncotarget* **8**, 89256–89268 (2017).

12. Shukla, S. *et al.* Progenitor T-cell differentiation from hematopoietic stem cells using Delta-like-4 and VCAM-1. *Nat. Methods* **14**, 531–538 (2017).

13. Dunbar, C. E. *et al.* Gene therapy comes of age. *Science* **359**, eaan4672 (2018).

14. Rangarajan, S. *et al.* AAV5-Factor VIII Gene Transfer in Severe Hemophilia A. *N. Engl. J. Med.* **377**, 2519–2530 (2017).

15. Russell, S. *et al.* Efficacy and safety of voretigene neparvovec (AAV2-hRPE65v2) in patients with RPE65-mediated inherited retinal dystrophy: a randomised, controlled, open-label, phase 3 trial. *Lancet* **390**, 849–860 (2017).

16. Christine, C. W. *et al.* Safety and tolerability of putaminal AADC gene therapy for Parkinson disease. *Neurology* **73**, 1662–1669 (2009).

17. Smith, T. T. *et al.* In situ programming of leukaemia-specific t cells using synthetic DNA nanocarriers. *Nat. Nanotechnol.* **12**, 813–822 (2017).

18. Pfeiffer, A. *et al.* In vivo generation of human CD 19- CAR T cells results in B-cell depletion and signs of cytokine release syndrome . *EMBO Mol. Med.* **10**, (2018).

19. Imai, C. *et al.* Chimeric receptors with 4-1BB signaling capacity provoke potent cytotoxicity against acute lymphoblastic leukemia. *Leukemia* **18**, 676–684 (2004).

20. Nicholson, I. C. *et al.* Construction and characterisation of a functional CD19 specific single chain Fv fragment for immunotherapy of B lineage leukaemia and lymphoma. *Mol. Immunol.* **34**, 1157–1165 (1997).

21. Kochenderfer, J. N., Yu, Z., Frasheri, D., Restifo, N. P. & Rosenberg, S. A. Adoptive transfer of syngeneic T cells transduced with a chimeric antigen receptor that recognizes murine CD19 can eradicate lymphoma and normal B cells. *Blood* **116**, 3875–86 (2010).

22. Ruella, M. *et al.* Induction of resistance to chimeric antigen receptor T cell therapy by transduction of a single leukemic B cell. *Nat. Med.* **24**, 1499–1503 (2018).

23. Turtle, C. J. *et al.* CD19 CAR-T cells of defined CD4+:CD8+ composition in adult B cell ALL patients. *J. Clin. Invest.* **126**, 2123–2138 (2016).

24. Baumann, J. G. *et al.* Murine T Cells Potently Restrict Human Immunodeficiency Virus Infection. *J. Virol.* **78**, 12537–12547 (2004).

25. Kerkar, S. P. *et al.* GeneticKerkar, S. P., Sanchez-Perez, L., Yang, S., Borman, Z. a, Muranski, P., Ji, Y., … Restifo, N. P. (2011). Genetic engineering of murine CD8+ and CD4+ T cells for preclinical adoptive immunotherapy studies. Journal of Immunotherapy (Hagerstown, Md. : *J. Immunother.* **34**, 343–52 (2011).

26. Tsurutani, N. *et al.* Nuclear Import of the Preintegration Complex Is Blocked upon Infection by Human Immunodeficiency Virus Type 1 in Mouse Cells. *J. Virol.* **81**, 677–688 (2007).

27. Delville, M. *et al.* A Nontoxic Transduction Enhancer Enables Highly Efficient Lentiviral Transduction of Primary Murine T Cells and Hematopoietic Stem Cells. *Mol. Ther. - Methods Clin. Dev.* **10**, 341–347 (2018).

28. Bezerra, E., Gauthier, J. & Hirayama, A. V. Factors associated with response, CAR-T cell *in vivo* expansion, and progression-free survival after repeat infusions of CD19 CAR-T cells. *Present. 61st Am. Soc. Hematol. Annu. Meet. Expo. Orlando, FL. December 7-10, 2019* (2019).

29. Blaese, R. M. *et al.* T lymphocyte-directed gene therapy for ADA-SCID: Initial trial results after 4 years. *Science (80-.).* **270**, 475–480 (1995).

30. Cavazzana-Calvo, M. *et al.* Gene therapy of human severe combined immunodeficiency (SCID)-X1 disease. *Science (80-.).* **288**, 669–672 (2000).

31. Anderson, W. F., Blaese, R. M. & Culver, K. The ADA human gene therapy clinical protocol: Points to Consider response with clinical protocol, July 6, 1990. *Hum. Gene Ther.* **1**, 331–362 (1990).

32. Bordignon, C. *et al.* Transfer of the ADA gene into bone marrow cells and peripheral blood lymphocytes for the treatment of patients affected by ADA-deficient SCID. *Hum. Gene Ther.* **4**, 513–20 (1993).

33. Fischer, A., Hacein-Bey, S. & Cavazzana-Calvo, M. Gene therapy of severe combined immunodeficiencies. *Nature Reviews Immunology* **2**, 615–621 (2002).

34. Cornetta, K. *et al.* Absence of Replication-Competent Lentivirus in the Clinic: Analysis of Infused T Cell Products. *Mol. Ther.* **26**, 280–288 (2018).

35. Marcucci, K. T. *et al.* Retroviral and Lentiviral Safety Analysis of Gene-Modified T Cell Products and Infused HIV and Oncology Patients. *Mol. Ther.* **26**, 269–279 (2018).

36. Zufferey, R., Nagy, D., Mandel, R. J., Naldini, L. & Trono, D. Multiply attenuated lentiviral vector achieves efficient gene delivery *in vivo*. *Nat. Biotechnol.* **15**, 871–875 (1997).

37. Naldini, L. *et al.* In vivo gene delivery and stable transduction of nondividing cells by a lentiviral vector. *Science (80-.).* **272**,

263–267 (1996).
38.   Dull, T. *et al.* A Third-Generation Lentivirus Vector with a Conditional Packaging System. *J. Virol.* **72**, 8463 (1998).

Figures

Figure 1: Proliferation and cytolytic properties of T cells from human PBMC transduced with FMC63-CD19CAR-GFP or GFP-only Lentivirus. **Figure 1A**: $5x10^5$ Human healthy donor PBMCs were transduced with the FMC63-CD19CAR-GFP or GFP-only lentivirus at a MOI of 5. Transduced PBMC were then co-cultured with an excess of irradiated allogeneic PBMCs which provided a source of B cells expressing the CD19 antigen. The FMC63-CD19CAR-GFP human T cells showed robust expansion (orange, square), compared to GFP transduced cell (green, triangle) which did not expand. **Figure 1B**: $1x10^5$ FMC63-CD19CAR-GFP human T cells from day 63 of the expansion assays were co-cultured with CD19+ (721.221) or CD19- (K562) target cells for 4h at effector to target (E:T) ratios of 1:1, 4:1, and 10:1 Target cell death was measured through the shift in the FVS780 signal, measured by flow cytometry [**Figure S3**]. Expanded human CAR-T cells showed robust and selective killing of the CD19$^+$ 721.221 cells. At a ratio of 1:1 (E:T) the human CAR-T cells killed 66.0% +/- 0.14% (mean +/- SD) of B cells, at a ratio of 4:1 (E:T) the human CAR-T cells killed 90.2% +/- 2.62% of B cells, and at a ratio of 10:1, the human CAR-T cells killed 94.4% +/- 0.35%) of B cells.Two-way ANOVA, Sidak's multiple comparison ***$p<0.001$. Inset: DAPI signal intensity in target cell populations, showing the shift to higher DAPI signal and corresponding target cell death in the CD19$^+$ (orange) population compared to the lower DAPI signal and lack of cytolysis in the CD19- (cyan) population.

Figure 2: Proliferation and cytolytic properties of CD3+ T cells from C57BL/6 splenocytestransduced with 1D3-CD19CAR-GFP, FMC63-CD19CAR-GFP, or GFP only transgenes. **Figure 2A**: 2.5x10⁵ C57BL/6 splenocytes were transduced with 1D3-CD19CAR-GFP or GFP-only lentivirus at an MOI of 5. The transduced splenocytes were then co-cultured with an excess of irradiated C57BL/6 splenocytes which provided a source of B cells expressing the CD19 antigen. The 1D3-CD19CAR transduced splenocytes showed a 12.2+/- 0.09 (mean +/- SD) fold greater expansion (red, square), compared to GFP transduced (green, triangle) (p < 0.002 paired, two-tail T test). **Figure 2B**: 2.5x10⁵ C57BL/6 splenocytes were transduced with the FMC63-CD19CAR-GFP or GFP-only lentivirus at an MOI of 5. Splenocytes were then co-cultured with irradiated C57BL/6 splenocytes as in Figure 1A. The FMC63-CD19CAR transduced splenocytes showed 8.8 +/- 0.03 fold greater expansion (orange, square), than GFP transduced (cyan, triangle), (p < 0.004 paired, two-tail T test). **Figure 2C**: 1x10⁵ FMC63-CD19CAR-GFP or 1D3-CD19CAR murine T cells from day 63 of the expansion assays were co-cultured for 5 hours with B cells isolated from another sample of C57BL/6 splenocytes at a E:T ratio of 1:1. Target cell death was measured through the shift in the FVS780 signal, measured by flow cytometry [**Figure S3**]. Expanded murine CAR-T cells showed robust and selective killing of murine B cells, with the 1D3-CD19CAR T cells killing 24.8 +/- 1.69% of B cells and the FMC63-CD19CAR T cells killing 18.5 +/- 1.08 % of B cells in 5 hour co-cultures. Two-way ANOVA, Sidak's multiple comparison ***p<0.001.

Figure 3: Analysis of Total CD20[+], CD45R/B220[+] B cells and GFP[+], CD3[+], CD90.2[+] T cells isolated from C57BL/6 whole blood pre- and post-treatment with 1D3-CD19CAR-GFP CAR, FMC63-CD19 CAR-GFP or control (GFP-only) lentiviruses. **Figure 3A**: C57BL/6 wildtype mice were treated with a single IV injection lentivirus (1.8-2 x10[7] IU in 200µl PBS), to deliver either the 1D3-CD19CAR-GFP CAR, FMC63-CD19 CAR, or control (GFP-only). Mice treated with the ID3-CD19CAR lentivirus had a maximum 1D3-CD19CART cell population of 42.5 +/- 1.07% (mean +/- SD) within their peripheral blood by week 5 (white bars) which was accompanied by B cell aplasia (grey bars) by week 5 which was maintained until the end of the experiment (week 8). Mice treated with the FMC63-CD19CAR lentivirus had a maximum FMC63-CD19CAR T cell population of 26.1 +/- 1.01% within their peripheral blood by week 5 (white bars) which was accompanied by B cell aplasia (grey bars) by week 5 which was maintained until the end of the experiment (week 8). **Figure 3B**: Representative flowgrams from **Figure 3A**

Figure 4: Analysis of C57BL/6 whole blood pre- and post-treatment with 1D3-CD19CAR-GFP CAR, FMC63-CD19 CAR-GFP or control (GFP-only) lentiviruses. **Figure 4A***:* Total (grey bars) and GFP+ (white bars) CD3+, CD90.2+ T cells, **Figure 4B**: Total (grey bars) and GFP+ (white bars) CD20+, CD45R/B220+ B cells, **Figure 4C**: Total (grey bars) and GFP+ (white bars) CD11b+ macrophage and other non-T cells, and **Figure 4D**: Total (grey bars) and GFP+ (white bars) CD335+ NK and NKT cells.

Figure 5: Mouse weights (grams) pre- and post treatment. **Figure 5A**: 1D3-CD19CAR-GFP lentivirus, **Figure 5B**: FMC63-CD19CAR-GFP lentivirus, and **Figure 5C**: GFP-only lentivirus or PBS. In **Figure 5A** and **5B**, the mice that were treated with the 1D3-CD19CAR-GFP lentivirus or the FMC63-CD19CAR-GFP showed a 5.5 +/- 2.97% (mean +/- SD) and 3.6 +/- 1.78% reduction in weight, respectively, within 5-7 days after receiving treatment. Once the mice weights recovered, they exhibited no other visible adverse effects.

Supplementary Materials

Figure S1: Viral supernatant titration flow gating protocol. After debris was eliminated by FSC-A vs. SSC-A gating, cell doublets and clumps were then eliminated by FSC-H vs. FSC-A gating followed by SSC-A vs. SSC-H gating. The live cells were determined by gating on the FVS780 negative cells which was on the GFP positive cells to determine the percentage of GFP expressing cells in order to measure transduction efficiency. The flow collected data was analysed using Flow Jo v10, for windows and GraphPad Prism software version 8.0.0 for windows (GraphPad Software, California, USA).

Figure S2: 1D3-CD19CAR-GFP, FMC63-CD19CAR-GFP, and GFP lentivirus transduction efficiency. **Figure S4A**: Titration of 1D3- CD19CAR-GFP and FMC63-CD19CAR-GFP viral supernatant titration over EL4 (green and purple, respectively) and K562 (red and blue, respectively) cell lines. While the transduction efficiency appears significantly better in the human K562s ($p < 0.001$ ANOVA), the transduction of a mouse cell line, EL4, indicates this lentivirus is able to transduce mouse cells, albeit at a lower frequency. **Figure S4B**: 2.5×10^5 splenocytes were transduced with 1D3-CD19CAR-GFP, FMC63-CD19CAR-GFP, and GFP lentivirus at an MOI of 5 (calculated based on **Figure S2A**). The percentage (%) of GFP+, CD3+ T cells within the whole splenocyte sample was determined by assessing 3.0×10^4 cells via flow cytometry 72 hours after transduction with virus. 1D3-CD19CAR-GFP had a transduction efficiency of 17.1% (+/-1.23%), FMC63-CD19CAR-GFP 16.3% (+/- 2.6%), and GFP 12.2% (+/- 3.85%).

Figure S3: Flow cytometry gating protocol for the *in vitro* FMC63-CD19CAR-GFP T cell expansion and cytolytic activity using human PBMCs. Debris, cell doublets, and clumps were eliminated by FSC-A vs. FSC-H gating followed by SSC-A vs. SSC-H gating. The target cells, stained with CP eFluor 670 were selected for and gated on DAPI to obtain the percentage of live vs dead target cells. The flow collected data was analysed using Flow Jo v10, for windows and GraphPad Prism software version 8.0.0 for windows (GraphPad Software, California, USA) [**Figure 1B**].

Figure S4: *In vitro* 1D3- or FMC63-CD19CAR-GFP T cell cytolytic activity using C57BL/6 splenocytes. Debris, cell doublets, and clumps were eliminated by FCS-A vs SSC-A gating followed by FSC-A vs. FSC-H and SSC-A vs. SSC-H gating. T and B cells were then identified by CD3 vs. CD20 gating. CD20 positive cells were selected for and gated on FVS-780 to obtain the percentage of live vs. dead target cells. The flow collected data was analysed using Flow Jo v10, for windows and GraphPad Prism software version 8.0.0 for windows (GraphPad Software, California, USA) [**Figure 2**]

Figure S5: Flow gating protocol for the analysis of mouse blood pre- and post-treatment with lentiviruses. Debris was eliminated by FSC-A vs. SSC-A gating, cell doublets and clumps were then eliminated by FSC-H vs. FSC-A gating followed by SSC-A vs. SSC-H gating. T and B cells were then identified based on expression of CD3 or CD20 and NK cells and monocytes based on the expression of CD335 or CD11b, respectively. The CD3 positive cells were then gated for expression of CD90.2, and these CD3 and CD90.2 positive cells were gated for GFP expression(the CD19CAR reference gene marker). CD20 positive cells were gated for CD45R/B220 expression, and the CD20 and CD45R/B220 positive cells were then gated for GFP expression. NK cells (CD335) or monocytes (CD11b) were individually gated for GFP expression. The flow collected data was analysed using Flow Jo v10, for windows and GraphPad Prism software version 8.0.0 for windows (GraphPad Software, California, USA) [**Figure 3 and Figure 4**]