Accurate prediction of *cis*-regulatory modules reveals a prevalent regulatory

genome of humans

Pengyu Ni[1]

Zhengchang Su [1,*]

1 Department of Bioinformatics and Genomics, the University of North Carolina at Charlotte,

9201 University City Boulevard, Charlotte, NC, 28223, USA

Contact Information:

Zhengchang Su

Email:zcsu@uncc.edu

Tel: +01-704-687-7996

Fax: +01-704-687-8667

Running title: Accurate prediction of *cis*-regulatory modules in human genomes

Abstract

Annotating all *cis*-regulatory modules (CRMs) and transcription factor (TF) binding sites(TFBSs) in genomes remains challenging. We tackled the task by integrating putative TFBSs motifs found in available 6,092 datasets covering 77.47% of the human genome. This approach enabled us to partition the covered genome regions into a CRM candidate (CRMC) set (56.84%) and a non-CRMC set (43.16%). Intriguingly, like known enhancers, the predicted 1,404,973 CRMCs are under strong evolutionary constraints, suggesting that they might be *cis*-regulator. In contrast, the non-CRMCs are largely selectively neutral, suggesting that they might not be cis-regulatory. Our method substantially outperforms three state-of-the-art methods (GeneHancers, EnhancerAtlas and ENCODE phase 3) for recalling VISTA enhancers and ClinVar variants, as well as by measurements of evolutionary constraints. We estimated that the human genome might encode about 1.46 million CRMs and 67 million TFBSs, comprising about 55% and 22% of the genome, respectively; for both of which, we predicted 80%. Therefore, the *cis*-regulatory genome appears to be more prevalent than originally thought.

Introduction

cis-regulatory sequences, also known as *cis*-regulatory modules (CRMs) (i.e., promoters, enhancers, silencers and insulators), are made of clusters of short DNA sequences that are recognized and bound by specific transcription factors (TFs)[1]. CRMs display different functional states in different cell types in multicellular eukaryotes during development and physiological homeostasis, and are responsible for specific transcriptomes of cell types[2]. A growing body of evidence indicates that CRMs are at least as important as coding sequences (CDSs) to account for inter-species divergence[3, 4] and intra-species diversity[5], in complex traits. Recent genome-wide association studies (GWAS) found that most complex trait-associated single nucleotide polymorphisms (SNPs) do not reside in CDSs, but rather lie in non-coding sequences (NCSs)[6, 7], and often overlap or are in linkage disequilibrium (LD) with TF

binding sites (TFBSs) in CRMs[8]. It has been shown that complex trait-associated variants systematically

disrupt TFBSs of TFs related to the traits [8], and that variation in TFBSs affects DNA binding, chromatin

modification, transcription[9-11], and susceptibility to complex diseases[12, 13] including cancer[14-17].

In principle, variation in a CRM may result in changes in the affinity and interactions between TFs and

their binding sites, leading to alterations in histone modifications and target gene expressions in

relevant cells[18, 19]. These alterations in molecular phenotypes can lead to changes in cellular and

organ-related phenotypes among individuals of a species[20, 21]. However, it has been difficult to link

non-coding variants to complex traits[18, 22], largely because of our lack of a good understanding of all

CRMs, their constituent TFBSs and target genes in genomes[23].

Fortunately, the recent development of ChIP-seq techniques for locating histone marks[24] and

TF bindings in genomes in specific cell/tissue types[25] has led to the generation of enormous amount of

data by large consortia such as ENCODE[26], Roadmap Epigenomics[27] and Genotype-Tissue Expression

(GTEx)[28], as well as individual labs worldwide[29]. These increasing amounts of ChIP-seq data for

relevant histone marks and various TFs in a wide spectrum of cell/tissue types provide an

unprecedented opportunity to predict a map of CRMs and constituent TFBSs in the human genome.

Many computational methods have been developed to explore these data individually or jointly[30]. For

instance, as the large number of binding-peaks in a typical TF ChIP-seq dataset dwarfs earlier motif-

finding tools (e.g., MEME[31] and BioProspector[32]) to find TFBSs of the ChIP-ed TF, new tools (e.g.,

DREME[33], MEME-ChIP[34], XXmotif [35] and Homer[36]) have been developed. However, some of

these tools (e.g. MEME-ChIP) were designed to find primary motifs of the ChIP-ed TF in short sequences

(~200bp) around the binding-peak summits in a small number of selected binding peaks in a dataset due

to their slow speed. Some faster tools (e.g. Homer, DREME, and XXmotif) are based on the

discriminative motif-finding schema[37] by finding overrepresented k-mers in a ChIP-seq dataset, but

they often fail to identify TFBSs with subtle degeneracy. As TFBSs form CRMs for combinatory regulation

in higher eukaryotes [1, 38], tools such as SpaMo [39], CPModule [40] and CCAT [41] have been

developed to identify multiple closely located motifs as CRMs in a single ChIP-seq dataset. However,

these tools cannot predict CRMs containing novel TFBSs, because they all depend on a library of known

motifs (e.g., TRANSFAC [42] or JASPAR [43]) to scan for cooperative TFBSs in binding peaks. Due

probably to the difficulty to find TFBS motifs in a mammalian TF ChIP-seq dataset that may contain tens

of thousands of binding peaks, few efforts have been made to explore entire sets of an increasing

number of TF ChIP-seq datasets to simultaneously predict CRMs and constituent TFBSs [44-47].

On the other hand, as a single histone mark is not a reliable CRM predictor, a great deal of

efforts have been made to predict CRMs based on multiple histone marks and chromatin accessibility

(CA) data from the same cell/tissue types using various machine-learning methods, including hidden

Markov models[48], dynamic Bayesian networks[49], time-delay neural networks[50], random

forest[51], and support vector machines (SVMs)[52]. Many enhancer databases have also been created

either by combining results of multiple such methods[53-55], or by identifying overlapping regions of CA

and histone mark tracks in the same cell/tissue types[56-60]. In particular, the ENCODE phase 3

consortium[26] recently identified 926,535 candidate *cis*-regulatory elements (cCREs) based on overlaps

between millions of DNase I hypersensitivity sites (DHSs)[61] and transposase accessible sites (TASs)[62],

active promoter histone mark H3K4me3[63] peaks, active enhancer mark H3K27ac[64] peaks, and

insulator mark CTCT[65] peaks in a large number of cell/tissue types. Although CRMs predicted by these

methods are often cell/tissue type-specific, their applications are limited to cell/tissue types for which

the required datasets are available[26, 48, 49, 66]. The resolution of these methods is also low[48, 49,

66] and often lacks TFBSs information[26, 48, 49, 66], particularly for novel motifs, although some

predictions provide TFBSs locations by finding matches to known motifs[54, 55, 59]. Moreover, results

of these methods are often inconsistent[67-70], e.g., even the best-performing tools (DEEP and CSI-

ANN) have only 49.8% and 45.2%, respectively, of their predicted CRMs overlap with the DHSs in Hela

cells[52]; and only 26% of predicted ENCODE enhancers in K562 cells can be experimentally verified[67].

The low accuracy of these methods might be due to the fact that CA and histone marks alone are not

reliable predictors of active CRMs [52, 67, 68, 70].

It has been shown that TF binding data are more reliable for predicting CRMs than CA and

histone mark data, particularly, when multiple closely located binding sites for key TFs were used [52,

67, 68, 70]. Moreover, although primary binding sites of a ChIP-ed TF tended to be enriched around the

summits of binding peaks, TFBSs of cooperators of the ChIP-ed TFs tend to appear at the two ends of

binding peaks[71, 72]. With this recognition, instead of predicting cell/tissue type specific CRMs using CA

and histone marks data, we proposed to first predict a largely cell-type agnostic or static map of CRMs

and constituent TFBBs in the genome by integrating all available TF ChIP-seq datasets for different TFs in

various cell/tissue types[46, 47], just as has been done for identifying all genes encoded in the genome

using gene expression data from all cell/tissue types[73]. We also proposed to appropriately extend

short binding peaks to the typical length of enhancers, so that more TFBSs for cooperators of the ChIP-

ed TF could be included [71, 72], and thus, full-length CRMs could be identified[46, 47]. Once a map of

CRMs and constituent TFBSs is available, the specificity of CRMs in any cell/tissue type can be

determined using one or few epigenetic mark datasets collected in the cell/tissue type[26], because

when anchored by correctly predicted CRMs, the accuracy of epigenetic marks for predicting active

CRMs could be largely improved [68]. Although our earlier implementation of this strategy, dePCRM,

resulted in promising results using even insufficient datasets available then[46, 47], we were limited by

three technical hurdles. First, although existing motif-finders such as DREME used in dePCRM worked

well for relatively small ChIP-seq datasets from organisms with smaller genomes such as the fly [47],

they are unable to work on very large entire datasets from mammalian cells/tissues, so we had to split a

large dataset into smaller ones for motif finding in the entire dataset[46], which may compromise the

accuracy of motif finding and complicate subsequent data integration. Second, although the distances

and interactions between TFBSs in a CRM are critical, both were not considered in our earlier scoring

functions [46, 47], potentially limiting the accuracy of predicted CRMs. Third, the earlier "branch-and-

bound" approach to integrate motifs found in different datasets is not efficient enough to handle a

much larger number of motifs found in an ever increasing number of large ChIP-seq datasets from

human cells/tissues[46, 47]. To overcome these hurdles, we developed dePCRM2 based on an ultrafast,

accurate motif-finder ProSampler[72], a novel effective combinatory motif pattern discovery method,

and scoring functions that model essentials of both the enhanceosome and billboard models of

CRMs[74-76]. Using available 6,092 ChIP-seq datasets covering 77.47% of the human genome after

extending the binding peaks, dePCRM2 was able to partition the covered genome regions into a CRM

candidate (CRMC) set and a non-CRMC set, and predict 201 unique TF binding motif families in the

CRMCs. Both evolutionary and independent experimental data indicate that at least the vast majority of

the predicted 1,404,973 CRMCs might be functional, while at least the vast majority of the predicted

non-CRMCs might not be functional.

Results

The dePCRM2 pipeline

TFs in higher eukaryotes tend to cooperatively bind to their TFBSs in CRMs[1]. Different CRMs of the

same gene are structurally similar and closely located[77]. For example, in the locus control region

(LCR) of the hemoglobin genes in the mouse genome, multiple enhancers with similar combinations of

TFBSs regulate the expression of different hemoglobin genes in different tissues and developmental

stages [78]. Moreover, functionally related genes are often regulated by the same sets of TFs in different

cell types during development and in maintaining physiological homeostasis[1]. Due to the clustering

nature of TFBSs of cooperative TFs in a CRM, if we extend the called short binding peaks of a TF ChIP-seq

dataset from the two ends and reach the typical size of a CRM (500~3,000bp)[79], the extended peaks

would have a great chance to contain TFBSs of cooperative TFs[46, 47, 72]. For instance, if two different

TFs cooperatively regulate the same regulons in several cell types, then at least some of the extended

peaks of datasets for the two TFs from these cell types should contain the TFBSs of both TFs, and even

have some overlaps if the CRMs are reused in different cell types. Therefore, if we have a sufficient

number of ChIP-seq datasets for different TFs from the same and different cell types, we are likely to

include datasets for some cooperative TFs, and their TFBSs may co-occur in some extended peaks. Based

on these observations, we designed dePCRM[46, 47] and dePCRM2 to predict CRMs and constituent

TFBSs by identifying overrepresented co-occurring patterns of motifs found by a motif-finder in a large

number of TF ChIP-seq datasets. dePCRM2 overcomes the aforementioned shortcomings of dePCRM as

follows. First, using an ultrafast and accurate motif-finder ProSampler[72], we can find significant motifs

in available ChIP-seq datasets of any size (Figures 1A and 1B) without the need to split large datasets

into small ones[46]. Second, after identifying highly co-occurring motifs pairs (CPs) in the extended

binding peaks in each dataset (Figure 1C), we cluster highly similar motifs in the CPs and find a unique

motif (UM) in each resulting cluster (Figure 1D). Third, we model distances and interactions among

cognate TFs of the binding sites in a CRM by constructing interaction networks of the UMs based on the

distance between the binding sites and the extent to which biding sites in the UMs cooccur to improve

prediction accuracy (Figure 1E). Fourth, we identify as CRMCs closely located clusters of binding sites of

the UMs along the genome (Figure 1F), thereby partitioning genome regions covered by the extended

binding peaks into a CRMCs set and a non-CRMCs set. Fifth, we evaluate each CRMC using a novel score

that considers not only the number of TFBSs in a CRM, but also the distances between the TFBSs, their

quality scores and all pair-wise cooccurring frequencies between their motifs (Figure 1G). Lastly, we

compute a p-value for each S_{CRM} score, so that CRMs and constituent TFBSs can be predicted at

different significant levels using different S_{CRM} score or p-value cutoffs. Clearly, as the number of UMs

is a small constant number constrained by the number of TF families encoded in the genome, the

downstream computation based on the set of UMs runs in a constant time, thus dePCRM2 is highly

scalable. The source code of dePCRM2 is available at http://github.com/zhengchangsulab/pcrm2

Figure 1. Schematic of the dePCRM2 pipeline. A. Extend each binding peak in each dataset to its two ends to reach
a preset length, i.e., 1,000bp. B. Find motifs in each dataset using ProSampler. C. Find CPs in each dataset. For
clarity, only the indicated CPs are shown, while those formed between motifs in pairs P_1 and P_2 in dataset d_1, and
so on, are omitted. D. Construct the motif similarity graph, cluster similar motifs and find UMs in the resulting
motif clusters. Each node in the graph represents a motif, while weights on the edges are omitted for clarity.
Clusters are connected by edges of the same color and line type. E. Construct UM interaction networks. Each node
in the networks represents a UM, while weights on the edges are omitted for clarity. F. Project binding sites in the
UMs back to the genome and link adjacent TFBSs along the genome, thereby identifying CRMs and non-CRMs.
G. Evaluate each CRMC by computing its S_{CRM} score and the associated p-value.
176
177
Unique motifs recall most known TF motifs families and have distinct patterns of interactions.

ProSampler identified at least one motif in 5,991 (98.70%) of the 6092 ChIP-seq datasets

(Supplementary Note) but failed to find any motifs in the remaining 101 (1.66%) datasets that all contain

less than 310 binding peaks (Table S1), indicating that they are likely of low quality. As shown in Figure

2A, the number of motifs found in a dataset generally increases with the increase in the number of

binding peaks in the dataset, but enters a saturation phase and stabilizes around 250 motifs when the

number of binding peaks is beyond 40,000. In total, ProSampler identified 856,793 motifs in the 5,991

datasets. dePCRM2 found co-occurring motif pairs (CPs) in each dataset (Figure 1C) by computing a

cooccurring score S_c for each pair of motifs in the dataset (formula 2). As shown in Figure 2B, S_c scores

show a trimodal distribution. dePCRM2 selected as CPs the motif pairs that accounted for the mode

with the highest S_c scores, and discarded those that accounted for the other two modes with lower S_c

scores, because these low-scoring motif pairs were likely to co-occur by chance. In total, dePCRM2

identified 4,455,838 CPs containing 226,355 (26.4%) motifs from 5,578 (93.11%) of the 5,991 datasets.

Therefore, we further filtered out 413 (6.89%) of the 5,991 datasets because each had a low S_c score

compared with other datasets. Clearly, more and less biased datasets are needed to rescue their use in

the future for more complete predictions. Clustering the 226,355 motifs in the CPs resulted in 245

clusters consisting of 2~72,849 motifs, most of which form a complete similarity graph or clique,

indicating that member motifs in a cluster are highly similar to each other (Figure S1A). dePCRM2 found

a UM in 201 (82.04%) of the 245 clusters (Figure S1B and Table S1) but failed to do so in 44 clusters due

to the low similarity between some member motifs (Figure S1A). Binding sites of the 201 UMs were

found in 39.87~100% of the sequences in the corresponding clusters, and in only 1.49% of the clusters

binding sites were not found in more than 50% of the sequences due to the low quality of member

motifs (Figure S2). Thus, this step retained most of putative binding sites in most clusters. The UMs

contain highly varying numbers of binding sites ranging from 64 to 13,672,868 with a mean of 905,288

(Figure 2C and Table S1), reminiscent of highly varying number of binding peaks in the datasets

(Supplementary Note). The lengths of the UMs range from 10 to 21bp with a mean of 11bp (Figure 2D),

which are in the range of the lengths of known TF binding motifs, although they are biased to 10bp due

to the limitation of the motif-finder to find longer motifs. As expected, a UM is highly similar to its

member motifs that are highly similar to each other (Figure S1A). For example, UM44 contains 250

highly similar member motifs (Figure 2E). Of the 201 UMs, 117 (58.2%) match at least one of the 856

annotated motifs in the HOCOMOCO [80] and JASPAR[81] databases, and 92 (78.63%) match at least

two (Table S2), suggesting that most UMs might consist of motifs of different TFs of the same TF

family/superfamily that recognize highly similar motifs, a well-known phenomenon[82, 83]. Thus, a UM

might represent a motif family/superfamily for the cognate TF family/superfamily. For instance, UM44

matches known motifs of nine TFs of the "ETS" family ETV4~7, ERG, ELF3, ELF5, ETS2 and FLI1, a known

motif of NFAT5 of the "NFAT-related factor" family, and a known motif of ZNF41 of the "more than 3

adjacent zinc finger factors" family (Figure 2F and Table S2). The high similarity of these motifs suggest

that they might form a superfamily. The remaining 84 (43.28%) of the 201 UMs might be novel motifs

recognized by unknown TFs (Figure S1B and Table S1). On the other hand, 64 (71.91%) of the 89

annotated motif TF families match one of the 201 UMs (Table S3), thus, our predicted UMs include most

of the known TF motif families.

To model interactions between cognate TFs of the UMs, we computed an interaction score

S_{INTER} based on distances and cooccurrence levels between binding sites of two UMs (formula 3), which

largely improves our earlier score (data not shown) that only considers cooccurring frequencies of

binding sites in two motifs [46, 47]. As shown in Figure 2G, there are clear interaction patterns between

putative cognate TFs of many UMs, many of which are supported by experimental evidence. For

example, in a cluster formed by 10 UMs (Figure 2H), seven of them (UM126, UM146, UM79, UM223,

UM170, UM103 and UM159) match known motifs of MESP1/ZEB1, TAL1::TCF3, ZNF740,

MEIS1/TGIF1/MEIS2/MEIS3, TCF4/ZEB1/CTCFL/ZIC1/ZIC4/SNAI1, GLI2/GLI3 and KLF8, respectively. At

least a few of them are known collaborators in transcriptional regulation. For example, GLI2 cooperates

with ZEB1 to repress the expression of *CDH*1 in human melanoma cells via directly binding to two close

binding sites at the *CDH*1 promoter[84]; ZIC and GLI cooperatively regulate neural and skeletal

development through physical interactions between their zinc finger domains [85]; and ZEB1 and TCF4

reciprocally modulate their transcriptional activities to regulate the expression of *WNT*[86], to name a

few.

Figure 2. Prediction of UMs. A. Relationship between the number of predicted motifs in a dataset and the size of the
dataset (number of binding peaks in the dataset). The datasets are sorted in ascending order of their sizes. B.
Distribution of cooccurrence scores (Sc) of motif pairs found in a dataset. The dotted vertical line indicates the
cutoff value (0.7) of Sc for predicting cooccurring pairs (CPs). C. Number of putative binding sites in each of the

UMs sorted in ascending order. D. Distribution of the lengths of the UMs and known motifs in the HOCOMOCO
and JASPAR databases. E. The logo and similarity graph of the 250 member motifs of UM44. In the graph, each
node in blue represents a member motif, and two member motifs are connected by an edge in green if their similarity
is greater than 0.8 (SPIC score). Four examples of member motifs are shown in the right panel. F. UM44 matches
known motifs of nine TFs of the "ETS", "NFAT-related factor", and "more than 3 adjacent zinc finger factors"
families. G. Heatmap of the interaction networks of the 201 UMs, while names of the UMs are omitted for clarity.
H. A blowup view of the indicated cluster in G, formed by 10 UMs, of which UM126, UM146, UM79, UM223,
UM170, UM103 and UM159 match known motifs of MESP1|ZEB1, TAL1::TCF3, ZNF740,
MEIS1|TGIF1|MEIS2|MEIS3, TCF4|ZEB1|CTCFL|ZIC1|ZIC4|SNAI1, GLI2|GLI3 and KLF8, respectively. Some
of these TFs are known collaborators in transcriptional regulation.
248
249
An appropriate extension of original binding peaks greatly increases the power of datasets

By concatenating closely located binding sites of the UMs along the genome, dePCRM2 partitioned the

77.47% of the genome that are covered by the extended binding peaks (Supplementary Note) in two

exclusive sets (Figure 1F), i.e., the CRMC set containing 1,404,973 CRMCs with a total length of bp

(56.84%) covering 44.03% of the genome, and the non-CRMC set containing 1,957,936 sequence

segments with a total length of 1,032,664,424bp (43.16%) covering 33.44% of the genome. Interestingly,

only 57.88% (776,999,862bp) of genome positions of the CRMCs overlap those of the original binding

peaks. Hence, dePCRM2 only retained 61.40% of genome positions covered by the original peaks, and

abandoned the remaining 38.60% of nucleotide position. These abandoned positions covered by

originally called binding peaks might not enrich for TFBSs, which is in agreement with earlier findings

about the noisy nature of TF ChIP-seq data [87-89]. On the other hand, the remaining 42.12%

(565,448,583bp) genome positions of the CRMCs overlap those of the extended parts of the original

peaks, indicating that TFBSs of cooperative TFs are indeed enriched in the extended parts as has been

shown earlier[46, 47, 71, 72], and dePCRM2 is able predict CRMs that are not covered by any binding

peaks. Thus, by appropriately extending original binding peaks, we could greatly increase the power of

datasets. Based on the overlap between a CRMC and original binding peaks in a cell/tissue type

(Materials and Methods), dePCRM2 predicted functional states of the 57.88% of the CRMCs in at least

one of the cell/tissue types from which binding peaks were called. However, dePCRM2 was not able to

predict the functional states of the remaining 42.12% of the CRMCs that do not overlap any original

binding peaks in the datasets. The predicted CRMCs and constituent TFBSs are available at https://cci-

bioinfo.uncc.edu/

The CRMCs are unlikely predicted by chance

To further evaluate the predicted CRMCs, we computed a S_{CRM} score for each CRMC (formula

4). As shown in Figure 3A, the distribution of the S_{CRM} scores of the CRMCs is strongly right-skewed

relative to that of the Null CRMCs (Materials and Methods), indicating that the CRMCs generally score

much higher than the Null CRMCs, and thus are unlikely produced by chance. Based on the distribution

of the S_{CRM} scores of the Null CRMCs, dePCRM2 computed a p-value for each CRMC (Figure 3A). With

the increase in the S_{CRM} cutoff α ($S_{CRM} \geq$ α), the associated p-value cutoff drops rapidly, while both the

number of predicted CRMs and the proportion of the genome covered by the predicted CRMs decrease

slowly (Figure 3B), indicating that most CRMCs have low p-values. For instance, with α increasing from

56 to 922, p-value drops precipitously from 0.05 to 1.00×10^{-6} (5×10^5 fold), while the number of

predicted CRMs decreases from 1,155,151 to 327,396 (3.53 fold), and the proportion of the genome

covered by the predicted CRMs decreases from 43.47% to 27.82% (1.56 fold) (Figure 3B). Predicted

CRMs contain from 20,835,542 (p-value $\leq 1 \times 10^{-6}$) to 31,811,310 (p-value $\leq$ 0.05) non-overlapping

putative TFBSs that consist of from 11.47% (p-value$\leq 1 \times 10^{-6}$) to 16.54% (p-value $\leq$ 0.05) of the genome

(Figure 3C). In other words, dependent on p-value cutoffs ($1 \times 10^{-6} \sim$ 0.05), 38.05~41.23% of nucleotide

positions of the predicted CRMs are made of putative TFBSs (Figure 3C), and most of predicted CRMs

(93.99~95.46%) and constituent TFBSs (93.20~94.67%) are located in non-exonic sequences (NESs)

(Figure 3C), comprising 26.66~42.47% and 10.94~16.03% of NESs, respectively (Figure 3D). Surprisingly,

dependent on p-value cutoffs ($1 \times 10^{-6} \sim$ 0.05), the remaining 4.54~6.01% and 5.33~6.80% of the

predicted CRMs and constituent TFBSs, respectively, are in exonic sequences (ESs, including CDSs, 5'-

and 3'-untranslated regions), respectively (Figure 3C), in agreement with an earlier report[90].

Figure 3. Prediction of CRMs using different S_{CRM} cutoffs. A. Distribution of S_{CRM} scores of the CRMCs and Null CRMCs. The inset is a blowup view of the indicated region. The vertical dashed lines indicate the associated p-values of the S_{CRM} cutoffs mentioned in the main text. B. Number of the predicted CRMs, proportion of the genome predicted to be CRMs and the associated p-value as functions of the S_{CRM} cutoff α. C. Percentage of the genome that are predicted to be CRMs and TFBSs in ESs and NESs using various S_{CRM} cutoffs and associated p-values. D. Percentage of NESs that are predicted to be CRMs and TFBSs using various S_{CRM} cutoffs and associated p-values. E. Distribution of the lengths of CRMs predicted using different S_{CRM} cutoffs and associated p-values.

The S_{CRM} score captures the length feature of enhancers

As shown in Figure 3E, the CRMCs with a mean length of 981bp are generally shorter than VISTA

enhancers with a mean length of 2,049bp. Specifically, 621,842 (44.26%) of the 1,404,973 CRMCs are

shorter than the shortest VISTA enhancer (428bp), suggesting that they might be short CRMs (such as

promoters or short enhancers) or components of long CRMs. However, these shorter CRMCs (< 428bp)

comprise only 7.42% of the total length of the CRMCs. The remaining 733,132 (55.74%) CRMCs

comprising 92.58% of the total length of the CRMCs are longer than the shortest VISTA enhancer

(428bp), thus most of them are likely full-length CRMs. Therefore, predicted CRMC positions in the

genome are mainly covered by full-length or longer CRMCs. As expected, with the increase in α

(decrease in p-value cutoff), the distribution of the lengths of the predicted CRMs shifts to right and

even surpass that for VISTA enhancers (Figure 3E), indicating shorter CRMCs can be effectively filtered

out by a higher S_{CRM} cutoff α (a smaller p-value). The remaining CRMCs might be different type of CRMs

with different length features. For instance, at a rather stringent S_{CRM} cutoff α =676 (p=5X10^{-6}), 976,345

(69.49%) shorter CRMCs with a mean length of 387bp were filtered out (Figure 3E), the remaining

428,628 (30.51%) CRMCs have similar length distribution (mean length of 2292bp) to that of VISTA

enhancers (mean length of 2049bp) (Figure 3E), which are mainly involved in development-related

functions and are generally longer than other types of enhancers [91]. However, it is worth noting that

VISTA enhancers may not necessarily all be in their full-length forms, because even a portion of an

enhancer could be still partially functional[1], and it is still technically difficult to validate very long

enhancers in transgene animal models in a large scale. Therefore, it is not surprising that with even

more stringent S_{CRM} cutoffs, the predicted CRMs could be longer than VISTA enhancers (Figure 3C), and

they are likely super-enhancers for cell differentiation of development[92]. Taken together, these

results suggest that the S_{CRM} score captures the length feature of enhancers.

The CRMCs and non-CRMCs show dramatically distinct evolutionary behaviors

To see how effectively dePCRM2 partitions the covered genome regions into the CRMC set and the non-

CRMC set, we compared their evolutionary behaviors with those of the entire set of VISTA enhancers

using the GERP[93] and phyloP[94] scores of their nucleotide positions in the genome. Both the GERP

and the phyloP scores quantify conservation levels of genome positions based on nucleotide

substitutions in alignments of multiple vertebrate genomes. The larger a positive GERP or phyloP score

of a position, the more likely it is under negative/purifying selection; and a GERP or phyloP score around

zero means that the position is selectively neutral or nearly so[93, 94]. However, a negative GERP or

phyloP score is cautiously related to positive selection[93, 94]. For convenience of discussion, we

consider a position with a GERP or phyloP score within an interval centering on 0 [-δ,+ δ] (δ >0) to be

selectively neutral or nearly so, and a position with a score greater than δ to be under negative

selection. We define proportion of neutrality of a set of positions to be the size of the area under the

density curve of the distribution of the scores of the positions within the window [-δ,+ δ]. Because ESs

evolve quite differently from NESs, we focused on the CRMCs and constituent TFBSs in NESs, and left

those that overlap ESs in another analysis (Jing Chen, Pengyu Ni, Jun-tao Guo and Zhengchang Su). The

choice of δ = 0.1, 0.2, 0.3, 0.4 ,0.5, 1 and 2 gave similar results (data not shown), so we choose δ =1 in

the subsequent analyses. As shown in Figure 4A, GERP scores of VISTA enhancers show a trimodal

distribution with a small peak around score -5, a blunt peak around score 0, a sharp peak around score

3.5, and a small proportion of neutrality of 0.23, indicating that most nucleotide positions of VISTA

enhancers are under strong evolutionary selection, particularly, negative selection. This result is

consistent with the fact that VISTAT enhancers are mostly ultra-conserved[95], development-related

enhancers[96, 97]. The 0.23 proportion of neutrality of the VISTA enhancer positions indicates that this

proportion of positions might simply serve as non-functional spacers between adjacent TFBSs.

Interestingly, there are 942 genome regions in the VISTA database, which failed to be validated as active

enhancers in transgenic assays, and we found that they had similar GERP and phyloP score distributions

as VISTA enhancers, although the former set is slightly less conserved than the latter set (Figure S3),

suggesting that most of these "validated negative regions (VNRs) might actually have *cis*-regulatory

functions under conditions that might have not be tested. In contrast, the distribution of the GERP scores

of the non-CRMCs (1,034,985,426 bp) in NESs displays a sharp peak around score 0, with low right and

left shoulders, and a high proportion of neutrality of 0.71 (Figure 4A), suggesting that most positions of

the non-CRMCs are selectively neutral or nearly so, and thus are likely to be nonfunctional. The

remaining 0.29 portion of positions of the non-CRMCs seem to be under varying levels of selection

(Figure 4A), so they might have other functions than cis-regulation. Intriguingly, the distribution of the

GERP scores of the 1,292,356 CRMCs (1,298,719,954bp) in NESs has a blunt peak around score 0, with

high right and left shoulders, and a small proportion of neutrality of 0.31 (Figure 4A). Thus, like VISTA

enhancers, most positions of the CRMCs are also under strong evolutionary selections, and thus, are

likely to be functional, while the small proportion (0.31) of neutrality indicates that this proportion of

positions in the CRMCs might simply serve as non-functional spacers, instead of TFBSs. Notably, the

distribution of GERP scores of the CRMCs lack obvious peaks around scores -5 and 3.5 (Figure 4A),

therefore, the average selection strength on the CRMCs is weaker than that on VISTA enhancers (but

see the section "The higher the S_{CRM} score of a CRMC, the stronger evolutionary constraint it is under"

). Nonetheless, this is expected considering the ultra-conversation nature of the small set of

development-related VASTA enhancers[95-97]. In any rate, the dramatic differences between the

evolutionary behaviors of the non-CRMCs and those of the CRMCs strongly suggests that dePCRM2

largely partitions the covered genome regions into a cis-regulatory CRMC set and a non-cis-regulatory

non-CRMC set. Similar results were obtained using the phyloP scores, although they display quite

different distributions than the GERP scores (Figure S4A).

To see why dePCRM2 abandoned the 38.60% nucleotide positions covered by the original

binding peaks in predicting the CRMCs, we plotted the distribution of their conservation scores. As

shown in Figure 4A, these abandoned positions have a GERP score distribution almost identical to those

in the non-CRMCs, indicating that, like the non-CRMCs, they are largely selectively neutral, and thus,

unlikely to be cis-regulatory, strengthening our earlier argument that they might not contain TFBSs.

Therefore, dePCRM2 is able to accurately distinguish cis-regulatory and non-cis-regulatory parts in both

the original binding peaks and their extended parts. As shown in Supplementary Note, the 10 CRM

function-related elements datasets (Tables S4~S8) that we collected for validating the predicted CRMCs

are strongly biased to the covered genome regions relative to the uncovered regions. To see why this is

possible, we plotted the distributions of conservation scores of the positions of the covered and

uncovered regions in NESs. Interestingly, the uncovered regions have a GERP score distribution and a

proportion of neutrality (0.59) in between those of the covered regions (0.49) and those of the non-

CRMCs (0.71) (Figure 4A), indicating that the uncovered regions are more evolutionarily selected than

the non-CRMCs as expected, but less evolutionary selected than the covered regions. This implies that

the uncovered regions contain functional elements such as CRMs, but their density could be lower than

that of the covered regions. Assuming that the total length of CRMs in a region is proportional to the

total length of evolutionarily constrained parts in the region, the proportion of uncovered regions that

might be CRMs could be estimated to be (1-0.59)/(1-0.49)=80.04% of that in the covered regions.

Therefore, it appears that existing studies are strongly biased to more evolutionary constrained regions

due probably to their large effect sizes and more critical functions. Similar results were obtained using

the phyloP scores (Figure S4A).

Figure 4. CRMCs and non-CRMCs in NESs show different evolutionary behaviors measured by GERP scores.
A. Distributions of the GERP scores of nucleotide positions of VISTA enhancers, CRMCs, non-CRMCs, abandoned genome regions covered by original binding peaks, genome regions covered by extended binding peaks and genome regions uncovered by extended binding peaks. The area under the density curves in the score interval [-1, 1] is defined as the proportion of neutrality of the sequences. B. Proportion of neutrality of CRMCs with a S_{CRM} score in different intervals in comparison with that of the non-CRMCs (a). The inset shows the distributions of the GERP scores of the non-CRMCs and CRMCs with S_{CRM} scores in the intervals indicted by color and letters. C. Proportion of neutrality of CRMs predicted using different S_{CRM} score cutoffs and associated p-values in comparison with those of the non-CRMCs (a) and CRMCs (b). The inset shows the distributions of the GERP scores of the non-CRMCs, CRMCs and the predicted CRMs using the S_{CRM} score cutoffs and p-values indicated by color and letters. The dashed lines in B and C indicate the saturation levels.

409   **The higher the S$_{CRM}$ score of a CRMC, the stronger evolutionary constraint it is under**

410   To see whether the S_{CRM} score of a CRMC captures the strength of evolutionary selection that it is

411   under, we plotted the distributions of the conservation scores of subsets of the CRMCs with a S_{CRM}

412   score in different non-overlapping intervals. Remarkably, even the subset with S_{CRM} scores in the lowest

413   interval [0, 1) has a smaller proportion of neutrality (0.56) than the non-CRMCs (0.71) (Figure 4B),

414   indicating that even these low-scoring CRMCs with short lengths (Figure 3E) are more likely to be under

415   strong evolutionary constraints than the non-CRMCs, and thus might be more likely cis-regulatory. With

416   the increase in the lower bound of S_{CRM} intervals, the proportion of neutrality of the corresponding

417   subsets of CRMCS drops rapidly, followed by a slow linear decrease around the interval [1000,1400)

418   (Figure 4B). Therefore, the higher the S_{CRM} score of a CRMC, the more likely it is under strong

419   evolutionary constraint, suggesting that the S_{CRM} score indeed captures the evolutionary behavior of a

420   CRM as a functional element, in addition to its length feature (Figure 3E). The same conclusion can be

421   drawn from the phyloP scores (Figure S4B).

422

423   We next examined the relationship between the conservation scores of the predicted CRMs and

424 S_{CRM} score cutoffs α (or p-value cutoffs) used for their predictions. As shown in Figure 4C, even the

425   CRMs predicted at a low α have a much smaller proportion of neutrality (e.g., 0.31 for the smallest α=0,

426   i.e., the entire CRMC set) than the non-CRMCs (0.71), suggesting that most of the predicted CRMs might

427   be authentic although some short ones may not be in full-length, while the non-CRMCs might contain

428   few false negative CRMCs. With the increase in α (decrease in p-value cutoff), the proportion of

429   neutrality of the predicted CRMs decreases but slowly, entering a saturation phase (Figure 4C).

430   Interestingly, at very high S_{CRM} score cutoffs, the predicted CRMs evolve like VISTA enhancers, with a

431   trimodal GERP score distribution, and thus might be involved in development[98, 99]. For instance, at α=

432   13,750, the distribution of GERP scores of the predicted CRMs displays a peak around score -5 and a

peak around score 3.5, with a small proportion of neutrality of 0.24 (Figure 4C) (it is 0.23 for VISTA

enhancers, Figure 4A). Thus, the higher α (i.e., the smaller the p-value cutoff), the more likely the

predicted CRMs are under strong evolutionary constraints. The infinitesimal decrease in the proportion

of neutrality of predicted CRMs with the increase in S_{CRM} cutoffs (Figure 4C) strongly suggests that the

predicted CRMs, particularly those at a low p-value cutoff, are under similarly strong evolutionary

constraints, close to the possibly highest saturation level to which ultra-conserved VISTA enhancers are

subject. Therefore, it is highly likely that at low p-value cutoffs, specificity of the predicted CRMs might

approach the possibly highest level that the VISTA enhancers achieve. However, without the availability

of a gold standard negative CRM set in the genome[23], we could not explicitly calculate the specificity

of the predicted CRMs at different p-value cutoffs. Similar results are observed using the phyloP scores

(Figure S4C).

444
dePCRM2 achieves high sensitivity and likely high specificity for recalling functionally validated CRMs

and non-coding SNPs

To further evaluate the accuracy of dePCRM2, we calculated the sensitivity (recall rate or true positive

rate (TPR)) of CRMs predicted at different S_{CRM} cutoffs α and associated p-values for recalling a variety

of CRM function-related elements located in the covered genome regions in the 10 experimentally

determined datasets in various cell/tissue types (Tables S4~S8, Materials and Methods). Here, if a

predicted CRM and an element overlap each other by at least 50% of the length of the shorter one, we

say that the CRM recalls the element. As shown in Figure 5A, with the increase in the p-value cutoff, the

sensitivity for recalling the elements in all the 10 datasets increases rapidly and becomes saturated well

before p-value increases to 0.05 (α ≥ 56). Figures S5A~S5J show examples of the predicted CRMs

overlapping and recalling the elements in the 10 datasets. Particularly, at p-value cutoff 5x10^{-5} (α=412),

the predicted 593,731 CRMs covering 36.63% of the genome (Figure 3C) recall 100% of VISTA

enhancers[79] and 91.61% of ClinVar SNPs[79] (Figure 5A). The rapid saturation of sensitivity for

recalling these two types of validated functional elements at such a low p-value cutoff once again

strongly suggests that dePCRM2 also achieves very high specificity, although we could not explicitly

compute it for the aforementioned reason. On the other hand, even at the higher p-value cutoff 0.05

(α=56), the predicted 1,155,151 CRMs covering 43.47% of the genome (Figure 3C) only achieve varying

intermediate levels of sensitivity for recalling FANTOM5 promoters (FPs)(88.77%)[100], FANTOM5

enhancers (FEs) (81.90%)[101], DHSs (74.68%)[61], TASs (84.32%)[29], H3K27ac (82.96%)[29], H3K4me1

(76.77%)[29], H3K4me3 (86.96%)[29] and GWAS SNPs (64.50%)[102], although all are significantly higher

than that (15%) of randomly selected sequences with matched lengths from the covered genome

regions (Figure 5A).

To find out the reasons for such varying sensitivity of dePCRM2 for recalling different types

elements in the 10 datasets, we plotted the distribution of GERP scores of the recalled and uncalled

elements in each dataset by our predicted CRMs at p-value <0.05. Since we have already plotted the

distribution of the entire set of VISTA enhancers (Figure 4A), to avoid redundancy, we instead plotted

the distribution for the CRMs (VISTA-CRMs) that overlap and recall the 785 VISTA enhancers in the

covered regions. As shown in Figure 5B, like the predicted CRMs, the recalled elements in all the

datasets are under strong evolutionary selections (at p-value <0.05), thus are likely functional. However,

VISTA-CRMs, recalled ClinVar SNPs and recalled FPs evolve more like VISTA enhancers with a trimodal

GERP score distribution (Figure 4A), suggesting that they are under stronger evolution constraints than

the other recalled element types. These results are not surprising, as we mentioned earlier VISTA

enhancers are mostly ultra-conserved, development related enhancers[95-97], while ClinVar SNPs were

identified for their conserved critical functions[103], and promoters are well-known to be more

conserved than are enhancers[104]. In stark contrast, like the non-CRMCs, all unrecalled elements in the

10 datasets are largely selectively neutral, and thus, are unlikely to be functional, with the exception

that the 10,350 (2.57%) unrecalled ClinVar SNPs display a trimodal distribution and there are no

unrecalled VISTA enhancers (Figure 5B). Notably, proportions of neutrality of unrecalled PEs (0.59) and

PFs(0.63) are smaller than that of the non-CRMCs (0.71) (Figure 5B), suggesting we might miss a small

portion of authentic PEs and PFs (see below for false negative rate (FNR) estimations of our CRMs).

Nevertheless, assuming that at least most of unrecalled elements in the datasets except the VISTA and

ClinVar datasets, are non-*cis*-regulatory, we estimated that the false discovery rate (FDR) of the

remaining eight datasets might be up to from 11.23% (1-0.8877) for FPs to 35.50% (1-0.6450) for GWAS

SNPs. Such high FDRs for CA (DHSs and TASs) and histone marks are consistent with an earlier study[68].

Interestingly, the trimodal distribution of GERP scores of the 2.57% of unrecalled ClinVar SNPs displays a

large peak around score 0 and two small peaks around -5 and 3.5, with a proportion of neutrality 0.40

(Figure 5B), indicating that about 40% of the relevant SNPs might be selectively neutral, and thus non-

functional. We therefore estimated the FDR of the ClinVar SNP dataset to be about 0.40*2.57%=1.03%.

Hence, like VISTA enhancers, ClinVar SNPs are a reliable set for evaluating CRM predictions. The peak of

the unrecalled ClinVar SNPs around score 3.5 (Figure 5B), indicates that the relevant SNPs are under

strong purifying selection, and thus might be functional, but were missed by dePCRM2. We therefore

estimate our predictions (at p-value <0.05) might have a FNR < 2.57%-1.03%=1.54%. In other words, the

real sensitivity (=1-FNR) for dePCRM2 to recall authentic ClinVar SNPs might be higher than the

calculated 97.54% (Figure 5A). These estimates are supported by the zero FNR and 100% sensitivity for

our predicted CRMs to recall VISTA enhancers (Figure 5A) and a simulation to be described later.

The zero, very low (<1.03%) and low (11.23%) FDRs of VISTA enhancers, ClinVar SNPs and FPs

datasets, respectively, are clearly related to the high reliability of the experimental methods used to

characterize them. However, the low FDRs might also be related to the highly conserved nature of these

elements (Figure 5B), as their critical functions and large effect sizes may facilitate their correct

characterization. In this regard, we note that the intermediately high FDRs of the FEs(18.10%),

DHSs(25.32), TASs (15.68%), H3K4m3 (13.04%), H3K4m1 (23.23%) and H3K27ac (17.04%) datasets might

be due to the facts that bidirectional transcription[105], CA[68, 70, 106] and histone marks[68, 70] are

not unique to active enhancers. The very high FDR of GWAS SNPs (35.5%) might be due to the fact that a

lead SNP associated with a trait may not necessarily be located in a CRM and causal; rather, some

variants in a CRM, which are in LD with the lead SNP, are the culprits[102, 107]. Example of GWAS SNPs

in LD with positions in a CRM are shown in Figures S5K and S5L. Interestingly, many recalled ClinVar

SNPs (42.59%) and GWAS SNPs (38.18%) are located in critical positions in predicted binding sites of the

UMs (e.g., Figures S5D and S5F).

In addition, we found that 722 (76.65%) of the 942 VNRs in the VISTA database fall in the

covered 77.47% genome regions. At a p-value cutoff of 0.05, the predicted CRMs recall 711 (98.48%) of

the 722 VNRs. Interestingly, recalled VNR positions evolve similarly to the VISTA enhancer positions,

while unrecalled VNR positions evolve similarly to the non-CRMC positions (Figure S3). These results

strongly suggest that recalled VNRs might be true enhancers that function in conditions yet to be tested,

as acknowledged by the VISTA team [108]. On the other hand, most unrecalled VNRs might not be *cis*-

regulatory.

Figure 5. Validation of the predicted CRMs by 10 experimentally determined sequence elements datasets. A. Sensitivity (recall rate or TPR) of the predicted CRMs and control sequences as a function of p-value cutoff for recalling the sequence elements in the datasets. The dashed vertical lines indicate the p-value ≤0.05 cutoff. B. Distributions of GERP scores of the recalled and unrecalled elements in each dataset in comparison with those of the predicted CRMs at p≤0.05 and non-CRMCs. Note that there are no unrecalled VISTA enhancers, and the distribution of the recalled 785 VISTA enhancers in the covered genome regions (not shown) is almost identical to the entire set of 976 VISTA enhancers (Figure 4A). The curve labeled by VISTA-CRMs is the distribution of CRMs that overlap and recall the 785 VISTA enhancers.

dePCRM2 outperforms state-of-the-art methods for predicting CRMs

We compared our predicted CRMs at p-value ≤ 0.05 ($S_{CRM} < 56$) with three most comprehensive sets of

predicted enhancers/promoters, i.e., GeneHancer 4.14[55], EnhancerAtals2.0[59] and cCREs[26]. For

convivence of discussion, we call these three sets enhancers or cCREs. GeneHancer 4.14 is the most

updated version containing 394,086 non-overlapping enhancers covering 18.99% (586,582,674bp) of the

genome (Figure 6A). These enhancers were predicted by integrating multiple sources of both predicted

and experimentally determined CRMs, including VISTA enhancers[79], ENCODE phase 2 enhancer-like

regions[109], ENSEMBL regulatory build[53], dbSUPER[110], EPDnew promoters[111], UCNEbase[112],

CraniofacialAtlas[113], FPs[100] and FEs [101]. Enhancers from ENCODE phase 2 and ESEMBL were

predicted based on multiple tracks of epigenetic marks using the well-regarded tools ChromHMM[48]

and Segway[114]. Of the GeneHancer enhancers, 388,407 (98.56%) have at least one nucleotide located

in the covered genome regions, covering 18.89% of the genome (Figure 6A). EnhancerAtlas 2.0 contains

7,433,367 overlapping cell/tissue-specific enhancers in 277 cell/tissue types, which were predicted by

integrating 4,159 TF ChIP-seq, 1,580 histone mark, 1,113 DHS-seq, and 1,153 other enhancer function-

related datasets, such as FEs[115]. After removing redundancy (identical enhancers in difference

cell/tissues), we ended up with 3,452,739 EnhancerAtlas enhancers that may still have overlaps,

covering 58.99% (1,821,795,020bp) of the genome (Figure 6A), and 3,417,629 (98.98%) of which have at

least one nucleotide located in the covered genome regions, covering 58.78% (1,815,133,195bp) of the

genome (Figure 6A). cCREs represents the most recent CRM prediction by the ENCODE phase 3

consortium[26], containing 926,535 non-overlapping cell type agnostic enhancers and promoters

covering 8.20% (253,321,371bp) of the genome. The cCREs were predicted based on overlaps among

703 DHS, 46 TAS and 2,091 histone mark datasets in various cell/tissue types produced by ENCODE

phases 2 and 3 as well as the Roadmap Epigenomics projects[26]. Of these cCREs, 917,618 (99.04%)

have at least one nucleotide located in the covered genome regions, covering 8.13% (251,078,466bp) of

the genome (Figure 6A). Thus, due probably to the aforenoted reasons, these three sets of predicted

enhancers and cCREs also are strongly biased to the covered regions relative to the uncovered regions.

Both the number (1,155,151) and genome coverage (43.47%) of our predicted CRMs (p-value<0.05) are

larger than those of GeneHancer enhancers (388,407 and 18.89%) and of cCREs (917,618 and 8.12%),

but smaller than those of EnhancerAtlas enhancers (3,417,629 and 58.78%), in the covered regions.

To make the comparisons fair, we first computed the sensitivity of these three sets of enhancers

and cCREs for recalling VISTA enhancers, ClinVar SNPs and GWAS SNPs in the covered regions. We

omitted FPs, FEs, DHSs, TASs and the three histone marks for the valuation as they were used in

predicting CRMs by GeneHancer 4.14, EnhancerAtlas 2.0 or ENCODE phase 3 consortium. We also

excluded VISTA enhancers for evaluating GeneHancer enhancers as the former were compiled in the

latter [55]. Remarkably, our predicted CRMs outperform EnhancerAtlas enhancers for recalling VISTA

enhancers (100.00% vs 94.01%) and ClinVar SNPs (97.43% vs 7.03%) (Figure 6B), even though our CRMs

cover a smaller proportion of the genome (43.47% vs 58.78%, or 35.22% more) (Figure 6A), indicating

that dePCRM2 has both higher sensitivity and specificity than the method behind EnhancerAtlas 2.0[59].

However, our CRMs underperform EnhancerAtlas enhancers for recalling GWAS SNPS (64.50% vs

69.36%, or 7.54% more) (Figure 6B). As we indicated earlier, the lower sensitivity of dePCRM2 for

recalling GWAS SNPs might be due to the fact that an associated SNP may not necessarily be causal

(Figures S5K and S5L). The higher sensitivity of EnhancerAtlas enhancers for recalling GWAS SNPs might

be simply thanks to their 35.22% more coverage of the genome (58.78%) than that of our predicted

CRMs (43.47%) (Figure 6A). Our predicted CRMs outperform cCREs for recalling VISTA enhancers (100%

vs 85.99%), ClinVar SNPs (97.43% vs 18.28%) and GWAS SNPs (64.50% vs 15.74%) (Figure 6B). Our

predicted CRMs also outperform GeneHancer enhancers for recalling ClinVar SNPs (97.43% vs 33.16%)

and GWAS SNPs (64.50% vs 34.11%) (Figure 6B). However, no conclusion can be drawn from these

results about the specificity of our predicted CRMs compared with GeneHancer enhancers and cCREs,

because our predicted CRMs cover a higher proportion of the genome than both of them (43.47% vs

18.89% and 8.20%). On the other hand, both GeneHancer 4.14 enhancers and cCREs outperform

EnhancerAtlas enhancers for recalling ClinVar SNPs (33.16% and 18.28% vs 7.03%)(Figure 6B), even

though they have a much smaller genome coverage than EnhancerAtlas enhancers (18.89% and 8.20%

vs 58.78%) (Figure 6A), indicating that they have higher specificity than EnhancerAtlas enhancers.

As shown in Figure 6C, the intersections/overlaps between the four predicted

CRMs/enhancer/cCREs sets are quite low. For instance, EnhancerAtlas enhancers, GeneHancer

enhancers and cCREs share 926,396,395bp (50.85%), 414,806,711bp (70.72%), and 194,709,825bp

(76.86%) of their nucleotide positions with our predicted CRMs, corresponding to 69.01%, 30.90% and

14.51% of the positions of our CRMs (Figure 6C), respectively. There are only 105,606,214bp shared by

all the four sets, corresponding to 5.80%, 18.00%, 41.69% and 7.87% of nucleotide positions covered by

EnhancerAtlas enhancers, GeneHancer enhancers, cCREs and our CRMs, respectively. As expected, the

50.85%, 70.72% and 76.86% of their nucleotide positions that EnhancerAtlas enhancers, GeneHancer

enhancers and cCREs share with our CRMs, respectively, evolve similarly to our predicted CRMs,

although those of GeneHancer enhancers and cCREs are under slightly higher evolutionary constraints

than our CRMs (Figure 6D). However, at a higher S_{CRM} cutoff, e.g. α=3,000 (p<2.2x10^{-302}), our predicted

CRMs are even under stronger evolutionary constraints than the shared GeneHancer enhancers and

cCREs positions (Figure 6D). Therefore, the shared GeneHancer enhancers and cCREs positions just

evolve like subsets of our predicted CRMs with higher S_{CRM} scores. By stark contrast, like the non-CRMCs,

the remaining 49.14%, 29.28% and 23.13% of their nucleotide positions that EnhancerAtlas enhancers,

GeneHancer enhancers and cCREs do not share with our CRMs, respectively, are largely selectively

neutral, although they all have slightly smaller proportion of neutrality than that of the non-CRMCs

(0.66, 0.63 and 0.61 vs. 0.71, respectively) (Figure 6D), due probably to the small FNR (<1.54%) of our

predicted CRMs. Nonetheless, these results strongly suggest that the vast majority of the unshared

positions of the three sets of predicted enhancers/eCREs are selectively neutral, and thus might be

nonfunctional. It appears that the predicted enhancers/cCREs in the three sets that overlap our CRMs

are likely to be authentic, while most of those that do not might be false positives. Hence, we estimated

the FDR of EnhancerAtlas enhancers, GeneHancer enhancers and cCREs to be around 49.14%, 29.28%

and 23.13%, respectively. Therefore, it is highly likely that GeneHancer 4.14 and cCREs might largely

under-predict enhancers as evidenced the fact that they are targeted at evolutionarily more constrained

elements (Figure 6D), even though they have rather high FDRs around 29.28% and 23.12%, respectively

(Figure 6D), while EnhancerAtlas 2.0 might largely over-predict enhancers with a very high FPR around

49.14% (Figure 6D).

Finally, we compared the lengths of the four sets of predicted CRMs/enhancers/cCREs with

those of VISTA enhancers . As shown in Figure 6E, the distribution of the lengths of cCREs has a narrow

high peak at 345bp with a mean length of 273bp and a maximal length of 350bp. It is highly likely that

the vast majority of authentic cCREs are just components of long CRMs, because even the longest cCREs

(350bp) is shorter and the shortest VISTA enhancer (428bp). The highly uniform lengths of the predicted

cCREs also indicate the limitation of the underlying prediction pipeline[26]. The distribution of

GeneHancer enhancers oscillates with a period of 166bp (Figure 6E), which might be an artifact of the

underlying algorithm for combining results from multiple sources [55]. Moreover, with a mean length of

1,488bp, GeneHancer enhancers are shorter than the VISTA enhancers (with a mean length 2,049bp)

(Figure 6E). EnhancerAtlas enhancers also have a shorter mean length (680bp) than VISTA enhancers

(3049bp) (Figure 6E). Our predicted CRMs at p-value <0.05 have a mean length of 1,162bp, thus also are

shorter than that of VISTA enhancers (Figure 6E). However, as we indicated earlier, with a more

stringent p-value cutoff 5×10^{-6}, the resulting 428,628 predicted CRMs have almost an identical length

distribution as the VISTA enhancers (Figure 3E). Taken together, these results unequivocally indicate

that our predicted CRMs are much more accurate than the three state-of-the-art predicted

enhancer/cCRE sets for both the nucleotide positions and lengths of CRMs/enhancers/cCREs.

Figure 6. Comparison of the performance of dePCRM2 and three state-of-the-art methods. A. Proportion of the
genome that are covered by enhancers/CRMs predicted by the four methods (All), and proportion of genome
regions covered by predicted enhancers/CRMs that at least partially overlap the covered genome regions (With
overlap). B. Sensitivity for recalling VISTA enhancers, ClinVar SNPs and GWAS SNPs, by the predicted
enhancers/CRMs that at least partially overlap the covered genome regions. C. Upset plot showing numbers of
nucleotide positions shared among the predicted CRMs, GeneHancer enhancers, EnhancerAtlas enhancers and
cCREs. D. Distributions of GERP scores of nucleotide positions of the CRMs predicted at p-value $\leq$ 0.05 and p-value
$\leq$ 2.2X10^{-308}, and the non-CRMCs, as well as of nucleotide positions that GeneHancer enhancers, EnhancerAtlas
enhancers and cCREs share and do not share with the predicted CRMs at p-value $\leq$ 0.05. E. Distributions of lengths
of the four sets of predicted enhancers/CRMs in comparison to that of VISTA enhancers. The inset is a blow-up
view of the axes defined region.
647
648　**At least half of the human genome might code for CRMs**
649
What is the proportion of the human genome coding for CRMs and TFBSs? The high accuracy of our

predicted CRMs and constituent TFBSs might well position us to more accurately address this interesting

and important, yet unanswered question[116, 117]. To this end, we took a semi-theoretic approach.

Specifically, we calculated the expected number of true positives and false positives in the CRMCs in

each non-overlapping S_{CRM} score interval based on the predicted number of CRMCs and the density of

S_{CRM} scores of Null CRMCs in the interval (Figure 7A), yielding 1,383,152 (98.45%) expected true

positives and 21,821 (1.55%) expected false positives in the CRMCs (Figure 7B). The vast majority of the

21,821 expected false positive CRMCs have a low S_{CRM} score < 4 (inset in Figure 7A) with a mean length

of 28 bp, comprising 0.02% (21,821x28/3,088,269,832) of the genome and 0.05% (0.0002/0.4403) of the

total length of the CRMCs, i.e., a FDR of 0.05% for nucleotide positions (Figure 7C). On the other hand, as

the CRMCs miss 2.49% of ClinVar SNPs in the covered genome regions (Figure 5A), the FNR of

partitioning the genome in CRMCs and non-CRMCs would be < 2.49%(1-0.40)=1.49%, given the

proportion of neutrality of 0.4 for the unrecalled ClinVar SNPs (Figure 5B). False negative CRMCs would

make up 0.67% of the genome and 1.99% of the total length of the non-CRMCs, meaning a false

omission rate (FOR) of 1.99% for nucleotide positions (Figure 7C). Hence, true CRM positions in the

covered regions would make up 44.68% (44.03%-0.02%+0.67%) of the genome (Figure 7C). In addition,

as we argued earlier, the CRMC density in the uncovered 22.53% genome regions is about 80.04% of

that in the covered regions, thus, CRMCs in the uncovered regions would be about 10.40% (0.2253 x

0.4468x0.8004/0.7747) of the genome (Figure 7C). Taken together, we estimated about 55.08%

(44.68%+10.40%) of the genome to code for CRMs, for which we have predicted 79.90% [(44.03-

0.02)/55.08]. Moreover, as we predicted that about 40% of CRCs are made up of TFBSs (Figure 3C), we

estimated that about 22.03% of the genome might encode TFBSs. Furthermore, assuming a mean length

1,162bp for CRMs (the mean length of our predicted CRMs at p-value <0.05), and a mean length of

10bp for TFBSs (Figure 2D), we estimated that the human genome would encode about 1,463,872 CRMs

(3,088,269,832x0.5508/1,162) and 67,034,584 TFBSs (3,088,269,832x0.2203/10).

Figure 7. Estimation of the portion of the human genome encoding CRMs. A. Expected number of true positive and false positive CRMCs in the predicted CRMCs in each one-unit interval of the S_{CRM} score. The inset is a blow-up view of the axes defined region. B. Expected cumulative number of true positives and false positives with the increase in S_{CRM} score cutoff for predicting CRMs. The inset is a blow-up view of the axes defined region. C. Proportions of the genome that are covered and uncovered by the extended binding peaks and estimated proportions of CRMCs in the regions. Numbers in the braces are the estimated proportions of the genome being the indicated sequence types, and numbers in the boxes are proportions of the indicated sequence types in the covered regions or the uncovered regions, so they are summed to 1.

Discussion

Identification of all functional elements, in particular, CRMs in genomes has been the central task in the postgenomic era, and enormous CRM function-related data have been produced to achieve the goal[23, 118]. Although great progresses have been made to predict CRMs in the genomes [26, 53, 55, 59, 119] using these data, most existing methods attempt to predict cell/tissue specific CRMs using CA and multiple tracks of histone marks collected in the same cell /tissue types[26, 48, 55, 59, 114]. These methods are limited by the scope of applications[26, 48, 114], low resolution of predicted CRMs[26, 59], lack of constituent TFBS information[26, 59], and high FDRs[68](Figure 6D). To overcome

these limitations, we proposed a different approach to first predict a cell type agnostic or static map of

CRMs and constituent TFBBs in the genome[46, 47] by identifying repeatedly cooccurring patterns of

motifs found in appropriately expanded binding peaks in a large number of TF ChIP-seq datasets for

different TFs in various cell/tissue types. Since it is mainly TFBSs in a CRM that define its structure and

function, it not surprising that TF ChIP-seq data are a more accurate predictor of CRMs than CA and

histone mark data[52, 68, 70]. Therefore, our approach might hold promise for more accurate

predictions of CRMs and constituent TFBSs, notwithstanding computational challenges. Once a map of

CRMs and constituent TFBBs in the gnome is available, functions of CRMs and constituent TFBSs in

cell/tissue types could be studied in a more focused and cost-effective ways. Another advantage of our

approach is that we do not need to exhaust all TFs and all cell/tissue types of the organism in order to

predict most, if not all, of CRMs and constituent TFBBSs in the genome as we demonstrated earlier[46,

47], because CRMs are often repeatedly used in different cell/tissue types, developmental stages and

physiological homeostasis[1]. Moreover, by appropriately extending the binding peaks in each dataset,

we could largely increase the chance to identify cooperative motifs and full-length CRMs, thereby

increasing the power of existing data, thereby further reducing the number of datasets needed as we

have demonstrated in this and previous studies [46, 47]. We might only need a large but limited

number of datasets to predict most, if not all, CRMs and TFBSs in the genome, as predicted UMs and

CRMs enters a saturation phase when more than few hundreds of datasets were used for the

predictions as we showed earlier [46]. Our earlier application of the approach resulted in very promising

results in the fly[47] and human[46] genomes even using a relatively small number of strongly biased

datasets available then. However, the earlier implementations were limited by computational

inadequacies of underlying algorithms to find and integrate motifs in ever increasing number of large TF

ChIP-seq datasets in mammalian cell/tissues[46, 47]. In this study, we circumvent the limitations by

developing the new pipeline dePCRM2 based on an ultrafast and accurate motif finder ProSampler, an

efficient motif pattern integration method, and a novel CRM scoring function that captures essential

features of full-length CRMs.

Remarkably, dePCRM2 enables us to partition the 77.47% genome regions covered by the

extended binding peaks in 6,079 TF ChIP-seq datasets into two exclusive sets, i.e., the CRMCs and non-

CRMCs. Multiple pieces of evidence strongly suggest that the partition might be highly accurate. First,

the vast majority of the CRMCs are unlikely predicted by chance as suggested by their small p-values

(Figure 3B). Second, even the subset of the CRMCs with the lowest S_{CRM} scores ((0,1]) are under

stronger evolutionary constraints than the non-CRMCs (Figures 4B and S4B), indicating that even these

low-scoring CRMCs are more likely to be functional than non-CRMCs, not to mention CRMCs with higher

S_{CRM} scores that are under stronger evolutionary constraints (Figures 4C, 4D, S4C and S4D). Third, the

vast majority of the CRMCs are under similarly strong evolutionary constraints, and a subset of the

CRMCs with high S_{CRM} scores evolve like the ultra-conserved, development-related VISTA enhancers

with trimodal GERP score distributions (Figures 4A and S4A). Fourth, all experimentally validated VISTA

enhancers and almost all (97.51%) of well-documented ClinVar SNPs in the covered genome regions are

recalled by the CRMCs (Figure 5A), indicating that the CRMCs have a very low FNR. Finally, our

simulation studies indicate that the CRMCs have a very low FDR of 0.05%, and the non-CRMCs have a

low FOR of 1.99% (Figure 7C), strongly suggesting that both sensitivity and specificity of our predicted

CRMs are very high. To the best of our knowledge, we are the first to accurately partition large regions

(77.47**%)** of the genome into a set (CRMCs) that are highly likely to be *cis*-regulatory, and a set (non-

CRMCs) that are highly unlikely to be *cis*-regulatory.

Accurate prediction of the length of CRMs is also critical, but it appeared to be a difficult problem

as evidenced by the peculiar distributions of the lengths of GeneHancer enhancers and cCREs (Figure 6E).

Although 44.26% (621,841) of our predicted 1,404,973 CRMCs are shorter than the shortest (428bp)

VISTA enhancer, and thus are likely CRM components, they comprise only 7.42% of the total length of the

CRMCs. The remaining 55.74% (783,132) of the CRMCs comprising 92.58% of the total length of the

CRMCs, are longer than the shortest (428bp) VISTA enhancer, and thus are likely full-length CRMs.

Therefore, the vast majority of the predicted CRMC positions in the genome might be covered by full-

length CRMs. Very short CRMCs tend to have small S_{CRM} scores and be under weak evolutionary

constraints, and thus can be effectively filtered out using more stringent S_{CRM} cutoffs (Figures 3E, 4C and

S4C). It has been shown that an enhancer's length and evolutionary behavior are determined by its

regulatory tasks [91], and conserved enhancers are active in development [98, 99], while fragile enhancers

are associated with evolutionary adaptation [98]. CRMCs with different S_{CRM} cutoffs might belong to

different functional types as indicated by their different evolutionary behaviors (4A, 4C, S4A and S4C) and

length distributions (Figures 3E). For example, like VISTA enhancers, CRMs predicted at high S_{CRM} cutoffs

tend to be longer (Figure 3E) and under stronger evolutionary constrains (Figures 4C and S4C), thus might

be mainly involved in development, whereas CRMs predicted at lower S_{CRM} cutoffs tend to be shorter

(Figure 3E) and under weaker evolutionary constrains (Figures 4C and S4C), thus might be mainly involved

in non-development related functions. On the other hand, the failure to predict full-length CRMs of short

CRM components might be due to insufficient data coverage on the relevant loci in the genome. This is

reminiscent of our earlier predicted, even shorter CRMCs (mean length = 182bp) using a much smaller

number and less diverse 670 datasets[46]. As we argued earlier[46] and confirmed here by the much

longer CRMCs (mean length = 981bp) predicted using the much larger and more diverse datasets albeit

still strongly biased to a few TFs and cell/tissue types (Supplementary Note). We anticipate that full-length

CRMs of these short CRM components can be predicted using even larger and more diverse TF ChIP-seq

data. Thus, efforts should be made in the future to increase the genome coverage and reduce data biases

by including more untested TFs and untested cell types in the TF ChIP-seq data generation.

Interestingly, our predicted CRMs (at p-value < 0.05) achieve perfect (100.00%) and very high

(97.43) sensitivity for recalling VISTA enhancers [79] and ClinVAR SNPs [103], respectively, but varying

intermediate sensitivity ranging from 64.50% (for GWAS SNPs) to 88.77% (for FPs) for recalling other

CRM function-related elements in the remaining eight datasets (Figure 5A). It appears that such varying

sensitivity is due to varying FDRs ranging from 0% (for VISTA enhancers) to 35.5% (for GWAS SNPs) of

the methods used to characterize the elements (Figure 5B). Our finding that DHSs, TASs, and histone

mark (H3K4m1, H3K4m3 and H3K27ac) peaks have high FDRs for predicting CRMs is consistent with an

earlier study showing that histone marks or CA were less accurate predictor of active enhancers than TF

binding data[68]. Thus, it is not surprising that our predicted CRMs substantially outperforms the three

state-of-the-art sets of predicted enhancers/cCREs, i.e., GeneHancer 4.14 [55], EnhancerAtals2.0 [59]

and cCREs[26], both for recalling VISTA enhancers (we excluded GeneHancer enhancers for this

evaluation since VISTA enhancers were a part of it) and ClinVar SNPs (Figure 6B) and for predicting the

lengths of CRMs (Figure 6E), because these three sets were mainly predicted based on overlaps between

multiple tracks of CA and histone marks in various cell/tissue type. Although great efforts have been

made to improve the accuracy of EnhancerAtlas 2.0 enhancers[59], GeneHancer 4.14 enhancers[55]

and cCREs[26], they still suffer quite high FDRs (49.14%, 29.28% and 23.12%, respectively).

Although dePCRM2 can predict functional states of CRMCs in a cell/tissue type that have original

binding peaks overlapping the CRMs, it cannot predict the functional states of CRMs in the extended

parts of the original binding peaks in a cell/tissue if the CRMs do not overlap any available binding peaks

of all TFs tested in the cell/tissue type. However, the functional state of each CRM in the map in any

cell/tissue type could be predicted based on overlap between the CRM and a single or few epigenetic

mark datasets such as CA, H3K27ac and/or H3K4m3 data collected from the very cell/tissue type.

Anchored by correctly predicted CRMs, these epigenetic marks could accurately predict the functional

states of the CRMs[68]. Thus, our approach might be more cost-effective for predicting both a static

map of CRMs as well as constituent TFBSs in the genome and their functional states in various cell/tissue

types.

Remarkably, although originally called binding peaks is the strongly biased to few cell types and

TFs (Supplementary Note), and the 6,092 TF ChIP-seq datasets cover only 40.96% of the genome, after

moderately extending the binding peaks, we increased the genome coverage to 77.47%, an 89.14 %

increase. Nucleotide positions of the extended parts of the peaks contribute 42.12% positions of the

predicted CRMCs. Therefore, appropriate extension of called binding peaks in the datasets can

substantially increase the power of available data. On the other hand, we abandoned 38.60% of

positions covered by the original binding peaks, which might be nonfunctional as they evolve like the

non-CRMCs (Figures 4A and S4A). Therefore, originally called binding peaks cannot be equivalent to

CRMs or parts of CRMs as has also been shown earlier[87-89], and integration of multiple TF ChIP-seq

datasets as demonstrated in this study is necessary for accurate genome-wide predictions of CRMs.

The proportion of the human genome that is functional is a topic under hot debate [109, 120]

and a wide range from 5% to 80% of the genome has been suggested to be functional based on

difference sources of evidence [23, 61, 116, 120, 121]. The major disagreement is for the proportion of

functional NCSs in the genome, mainly CRMs, which have been coarsely estimated to comprise from 8%

to 40% of the genome [109, 120]. Moreover, a wide range of CRM numbers from 400,000 [109] to more

than a few million [23, 59] has been suggested to be encoded in the human genome. However, to our

best knowledge, no estimate has been made on substantial evidence. Our predicted CRMCs cover

44.03% of the genome, which is lower than EnhancerAtlas enhancers (58.99%)[59] do. The much higher

accuracy of our predicted CRMs suggests that cCREs (7.9%)[26] and GeneHancer enhancer might

underpredict, whereas EnhancerAtlas 2.0 might overpredict CRMs. Based on the estimated FDR and FNR

of the predicted CRMCs and non-CRMCs as well as the estimated density of CRMs in the uncovered

regions relative to the covered regions (Figure 7C), we estimated that about 55.08% and 22.03% of the

genome might code for CRMs and TFBSs, respectively, which encode about 1.46 million CRMs and 67

million TFBSs. Therefore, the number of our predicted CRMs is almost four times more than an earlier

estimate of 400,000 [109], and they are encoded by a higher (55.08%) proportion of the genome than

earlier thought 40%[109, 120]. We estimated that our true positive CRMs cover 44.01% (44.03-0.02) of

the genome, therefore, we might have predicted 79.90 % (44.01/55.08) CRM positions encoded in the

genome. In summary, it appears that the *cis*-regulatory genome is more prevalent than originally

thought.

Conclusions

We have developed a new highly accurate and scalable pipeline dePCRM2 for predicting CRMs

and constituent TFBSs in large genomes by integrating a large number of TF ChIP-seq datasets for

various TFs in a variety of cell/tissue types of the organisms. Applying dePCRM2 to all available ~6,000

TF ChIP-seq datasets, we predicted an unprecedentedly complete, high resolution map of CRMs and

constituent TFBSs in 77.47% of the human genome covered by extended binding peaks of the datasets.

Evolutionary and experimental data suggest that dePCRM2 achieves very high prediction sensitivity and

specificity. Based on the predictions, we estimated that about 55% and 22% of the genome might code

for CRMs and TFBSs, encoding about 1.46 million CRMs and 67 million TFBSs, respectively; for both of

which we predicted about 80%. Therefore, the *cis*-regulatory genome is more prevalent than originally

thought. With the availability in the future of more diverse and balanced data covering more regions of

the genome, it is possible to predict a more complete map of CRMs and constituent TFBSs in the

genome.

Materials and Methods

Datasets

We downloaded 6,092 TF ChIP-seq datasets from the Cistrome database[29]. The binding peaks in each

dataset were called using a pipeline for uniform processing[29]. We filtered out binding peaks with a

read depth score less than 20. For each binding peak in each dataset, we extracted a 1,000 bp genome

sequence centering on the middle of the summit of the binding peak. We downloaded 976

experimentally verified enhancers and 942 negatively validated regions (NVRs) from the VISTA Enhancer

database[79], 424,622 ClinVar SNPs from the ClinVar database[103], 32,689 enhancers[101] and

184,424 promoters[100] from the FANTOM5 project website, 91,369 GWAS SNPs from GWAS

Catalog[102], and 122,468,173 DHSs in 1,353 datasets (Table S4), 29,520,736 transposase-accessible

sites (TASs) in 1,059 datasets (Table S5), 99,974,447 H3K27ac peaks in 2,539 datasets (Table S6),

77,500,232 H3K4me1 peaks in 1,210 datasets (Table S7), and 70,591,888 H3K4me3 peaks in 2,317

datasets (Table S8) from the Cistrome database[29].

Measurement of the overlap between two different datasets

To evaluate the extent to which the binding peaks in two datasets overlap with each other, we calculate

an overlap score $S_0(d_i, d_j)$ between each pair of datasets d_i and d_j, defined as,

$$S_0(d_i, d_j) = \frac{1}{2} \times \left(\frac{o(d_i, d_j)}{|d_i|} + \frac{o(d_i, d_j)}{|d_j|} \right), \qquad (1)$$

where $o(d_i, d_j)$ is the number of binding peaks in d_i and d_j that overlap each other by at least one bp.

Parameters for accuracy evaluation

We use the following definitions to evaluate the accuracy of datasets and predictions. $Sensitivity =$

recall rate $= TPR$ (true positive rate) $= \frac{TP}{TP+FN}$, FNR (false negative rate) $= \frac{FN}{TP+FN}$, Specificity $=$

$\frac{TN}{FP+TN}$, FPR(false positive rate) $\frac{FP}{FP+TN}$, FDR (false discorery rate) $= \frac{FP}{TP+FP}$, and

OR (false ommision rate) $= \frac{FN}{FN+TN}$, where TP is true positives; FN, false negatives; FP, false positives;

and TN, true negatives.

The dePCRM2 pipeline

Step 1: Find motifs in each dataset using ProSampler[72](Figures 1A and 1B).

Step 2. Compute pairwise motif co-occurring scores and find co-occurring motif pairs (CPs): As True

motifs are more likely to co-occur in the same sequence than spurious ones, to filter out false positive

motifs, we find overrepresented CPs in each dataset (Figure 1C). Specifically, for each pair of motifs

$M_d(i)$ and $M_d(j)$ in each data set d, we compute their co-occurring scores S_c defined as,

$$S_c\left(M_i(i), M_j(j)\right) = \frac{o\left(M_d(i), M_d(j)\right)}{max\{|M_d(i)|,|M_d(i)|\}}, \tag{2}$$

where $|M_d(i)|$ and $|M_d(j)|$ are the number of binding peaks containing TFBSs of motifs $M_d(i)$ and

$M_d(j)$, respectively; and $o\left(M_d(i), M_d(j)\right)$ the number of binding peaks containing TFBSs of both the

motifs in d. We identify CPs with an $S_c \geq \beta$. We choose β such that the component with the highest

scores in the trimodal distribution S_c is kept (Figures 1C and 2B) (by default $\beta = 0.7$).

Step 3. Construct a motif similarity graph and find unique motifs (UMs): We combine highly similar

motifs in the CPs from different datasets to form a UM presumably recognized by a TF or highly similar

TFs of the same family/superfamily[122]. Specifically, for each pair of motifs $M_a(i)$ and $M_b(j)$ from

different datasets a and b, respectively, we compute their similarity score S_s using our SPIC[123] metric.

We then build a motif similarity graph using motifs in the CPs as nodes and connecting two motifs with

their S_s being the weight on the edge, if and only if (iff) $S_s>\beta$ (by default, β=0.8, Figure 1D). We apply

the Markov cluster (MCL) algorithm [124] to the graph to identify dense subgraphs as clusters. For each

cluster, we merge overlapping sequences, extend each sequence to a length of 30bp by padding the

same number of nucleotides from the genome to the two ends, and then realign the sequences to form

a UM using ProSampler[72](Figure 1D).

Step 4. Construct the interaction networks of the UMs/TFs: TFs tend to repetitively cooperate with each

other to regulate genes in different contexts by binding to their cognate TFBSs in CRMs. The relative

distances between TFBSs in a CRM often do not matter (billboard model), but sometimes they are

constrained by the interactions between cognate TFs (enhanceosome model) [74-76]. To model

essential features of both scenarios, we compute an interaction score between each pair of UMs, U_i and

U_j, defined as,

890
$$S_{INTER}(U_i, U_j) = \frac{1}{|D(U_i,U_j)|}\sum_{d\in D(U_i,U_j)}\left(\frac{1}{|d(U_i)|} + \frac{1}{|d(U_j)|}\right)\sum_{s\in S(d(U_i),(d(U_j))}\frac{150}{r(s)}, \qquad (3)$$

where $D(U_i, U_j)$ is the datasets in which TFBSs of both U_i and U_j occur, $d(U_k)$ the subset of dataset d,

containing at least one TFBS of U_k, $S(d(U_i), (d(U_j))$ the subset of d containing TFBSs of both U_i and U_j,

and $r(s)$ the shortest distance between any TFBS of U_i and any TFBS of U_j in a sequence s. We

construct UM/TF interaction networks using the UMs as nodes and connecting two nodes with their

S_{INTER} being the weight on the edge (Figure 1E). Therefore, the S_{INTER} score allows flexible adjacency and

orientation of TFBSs in a CRM (billboard model) and at the same time, it rewards motifs with binding

sites co-occurring frequently in a shorter distance in a CRM (enhanceosome model), particularly within a

nucleosome with a length of about 150bp[74, 75, 125]**.**

Step 5. Partition the covered genome regions into a CRM candidate (CRMC) set and a non-CRMC set: We

project TFBSs of each UM back to the genome, and link two adjacent TFBSs if their distance $d \leq 300bp$

(roughly the length of two nucleosomes). The resulting linked DNA segments are CRMCs, while DNA

segments in the covered regions that cannot be linked are non-CRMCs (Figure 1F).

Step 6. Evaluate each CRMC: We compute a CRM score for a CRMC containing n TFBSs ($b_1, b_2, ..., b_n$),

defined as,

$$S_{CRM}(b_1, b_2 \cdots, b_n) = \frac{2}{n-1} \times \sum_{i=1}^{n} \sum_{j>i} S_{INTER}[U(b_i), U(b_j)] \times [S(b_i) + S(b_j)], \qquad (4)$$

where $U(b_k)$ is the UM of TFBS b_k, $S_{INTER}[U(b_i), U(b_j)]$ the weight on the edge between $U(b_i)$ and

$U(b_j)$, in the interaction networks, and $S(b_k)$ the score of b_k based on the position weight matrix

(PWM) of $U(b_k)$. Only TFBSs with a positive score are considered. Thus, S_{CRM} considers the number of

TFBSs in a CRMC, as well as their quality and strength of all pairwise interactions.

Step 7. Predict CRMs: We create the Null interaction networks by randomly reconnecting the nodes with

the edges in the interaction networks constructed in Step 4. For each CRMC, we generate a Null CRMC

that has the same length and nucleotide compositions as the CRMC using a third order Markov chain

model[72]. We compute a S$_{CRM}$ score for each Null CRMC using the Null interaction networks, and the

binding site positions and PWMs of the UMs in the corresponding CRMC. Based on the distribution of

the S$_{CRM}$ scores of the Null CRMCs, we compute an empirical p-value for each CRMC, and predict those

with a p-value smaller than a preset cutoff as CRMs in the genome (Figure 1G).

Step 8. Prediction of the functional states of CRMs in a given cell type: For each predicted CRM at p-

value <0.05, we predict it to be active in a cell/tissue type, if its constituent binding sites of the UMs

whose cognate TFs were tested in the cell/tissue type overlap original binding peaks of the TFs;

otherwise, we predict the CRM to be inactive in the cell/tissue type. If the CRM does not overlap any

binding peaks of the TFs tested in the cell/tissue type, we assign its functional state in the cell/tissue

type "TBD" (to be determined).

Generation of control sequences for validation

To create a set of matched control sequences for validating the predicted CRMs using experimentally

determined elements used in Figure 5A, for each predicted CRMC, we produced a control sequence by

randomly selecting a sequence segment with the same length as the CRMC from the genome regions

covered by the extended binding peaks. To calculate the S$_{CRM}$ score of a control sequence, we assigned it

the TFBS positions and their UMs according to those in the counterpart CRMC. Thus, the control set

contains the same number and length of sequences as in the CRMCs, but with arbitrarily assigned TFBSs

and UMs.

Authors' contributions

ZS conceived the project. ZS and PN developed the algorithms and PN carried out all computational

experiments and analysis. ZS and PN wrote the manuscripts. All authors read and approved the final

manuscript.

Funding

The work was supported by US National Science Foundation (DBI- 1661332). The funding bodies played

no role in the design of the study and collection, analysis, and interpretation of data and in writing the

manuscript.

Ethics approval and consent to participate

Not applicable.

Competing interests

The authors declare no competing financial interests.

REFERENCES

1.   Davidson EH. The Regulatory Genome: Gene Regulatory Networks In Development And
Evolution: Academic Press; 2006.
2.   Wilczynski B, Furlong EE. Dynamic CRM occupancy reflects a temporal map of developmental
progression. Molecular systems biology. 2010;6:383. Epub 2010/06/24. doi: msb201035 [pii]

10.1038/msb.2010.35 [doi]. PubMed PMID: 20571532; PubMed Central PMCID: PMC2913398.
3.   King M, Wilson A. Evolution at two levels in humans and chimpanzees. Science. 1975;188:107-
16.
4.   Rubinstein M, de Souza FS. Evolution of transcriptional enhancers and animal diversity. Philos
Trans R Soc Lond B Biol Sci. 2013;368(1632):20130017. Epub 2013/11/13. doi: rstb.2013.0017 [pii]

10.1098/rstb.2013.0017 [doi]. PubMed PMID: 24218630; PubMed Central PMCID: PMC3826491.
5.   Siepel A, Arbiza L. Cis-regulatory elements and human evolution. Curr Opin Genet Dev.
2014;29:81-9. Epub 2014/09/15. doi: 10.1016/j.gde.2014.08.011. PubMed PMID: 25218861; PubMed
Central PMCID: PMCPMC4258466.

6.   Hindorff LA, Sethupathy P, Junkins HA, Ramos EM, Mehta JP, Collins FS, et al. Potential etiologic
and functional implications of genome-wide association loci for human diseases and traits. Proc Natl
Acad Sci U S A. 2009;106(23):9362-7. Epub 2009/05/29. doi: 0903103106 [pii]

10.1073/pnas.0903103106 [doi]. PubMed PMID: 19474294; PubMed Central PMCID: PMC2687147.
7.   Ramos EM, Hoffman D, Junkins HA, Maglott D, Phan L, Sherry ST, et al. Phenotype-Genotype
Integrator (PheGenI): synthesizing genome-wide association study (GWAS) data with existing genomic
resources. Eur J Hum Genet. 2014;22(1):144-7. Epub 2013/05/23. doi: ejhg201396 [pii]

10.1038/ejhg.2013.96 [doi]. PubMed PMID: 23695286; PubMed Central PMCID: PMC3865418.
8.   Maurano MT, Humbert R, Rynes E, Thurman RE, Haugen E, Wang H, et al. Systematic
Localization of Common Disease-Associated Variation in Regulatory DNA. Science. 2012;337(6099):1190-
5. doi: 10.1126/science.1222794.
9.   Kasowski M, Kyriazopoulou-Panagiotopoulou S, Grubert F, Zaugg JB, Kundaje A, Liu Y, et al.
Extensive variation in chromatin states across humans. Science. 2013;342(6159):750-2. Epub
2013/10/19. doi: 10.1126/science.1242510. PubMed PMID: 24136358; PubMed Central PMCID:
PMCPMC4075767.
10.   Kilpinen H, Waszak SM, Gschwind AR, Raghav SK, Witwicki RM, Orioli A, et al. Coordinated
effects of sequence variation on DNA binding, chromatin structure, and transcription. Science.
2013;342(6159):744-7. Epub 2013/10/19. doi: science.1242463 [pii]

10.1126/science.1242463 [doi]. PubMed PMID: 24136355.
11.   McVicker G, van de Geijn B, Degner JF, Cain CE, Banovich NE, Raj A, et al. Identification of
genetic variants that affect histone modifications in human cells. Science. 2013;342(6159):747-9. Epub
2013/10/19. doi: science.1242429 [pii]

10.1126/science.1242429 [doi]. PubMed PMID: 24136359.
12.   Smith E, Shilatifard A. Enhancer biology and enhanceropathies. Nat Struct Mol Biol.
2014;21(3):210-9. doi: 10.1038/nsmb.2784.
13.   Mathelier A, Shi W, Wasserman WW. Identification of altered cis-regulatory elements in human
disease. Trends Genet. 2015;31(2):67-76. Epub 2015/02/01. doi: 10.1016/j.tig.2014.12.003. PubMed
PMID: 25637093.
14.   Herz HM, Hu D, Shilatifard A. Enhancer malfunction in cancer. Mol Cell. 2014;53(6):859-66. Epub
2014/03/25. doi: 10.1016/j.molcel.2014.02.033. PubMed PMID: 24656127; PubMed Central PMCID:
PMCPmc4049186.
15.   Ongen H, Andersen CL, Bramsen JB, Oster B, Rasmussen MH, Ferreira PG, et al. Putative cis-
regulatory drivers in colorectal cancer. Nature. 2014;512(7512):87-90. doi: 10.1038/nature13602. Epub
2014 Jul 23.
16.   Khurana E, Fu Y, Chakravarty D, Demichelis F, Rubin MA, Gerstein M. Role of non-coding
sequence variants in cancer. Nat Rev Genet. 2016;17(2):93-108. Epub 2016/01/20. doi:
10.1038/nrg.2015.17. PubMed PMID: 26781813.
17.   Zhou S, Treloar AE, Lupien M. Emergence of the Noncoding Cancer Genome: A Target of Genetic
and Epigenetic Alterations. Cancer discovery. 2016;6(11):1215-29. Epub 2016/11/04. doi: 10.1158/2159-
8290.cd-16-0745. PubMed PMID: 27807102; PubMed Central PMCID: PMCPMC5117653.
18.   Whitaker JW, Chen Z, Wang W. Predicting the human epigenome from DNA motifs. Nat
Methods. 2015;2015 Mar;12(3):265-72.
19.   Wang M, Zhang K, Ngo V, Liu C, Fan S, Whitaker JW, et al. Identification of DNA motifs that
regulate DNA methylation. Nucleic Acids Res. 2019;47(13):6753-68. Epub 2019/07/25. doi:
10.1093/nar/gkz483. PubMed PMID: 31334813; PubMed Central PMCID: PMCPMC6649826.

20.   Ward LD, Kellis M. Interpreting noncoding genetic variation in complex traits and human
disease. Nat Biotechnol. 2012;30(11):1095-106. Epub 2012/11/10. doi: nbt.2422 [pii]

10.1038/nbt.2422 [doi]. PubMed PMID: 23138309; PubMed Central PMCID: PMC3703467.
21.   Pai AA, Pritchard JK, Gilad Y. The genetic and mechanistic basis for variation in gene regulation.
PLoS Genet. 2015;11(1):e1004857. doi: 10.1371/journal.pgen.. eCollection 2015 Jan.
22.   Albert FW, Kruglyak L. The role of regulatory variation in complex traits and disease. Nat Rev
Genet. 2015;16(4):197-212. Epub 2015/02/25. doi: 10.1038/nrg3891. PubMed PMID: 25707927.
23.   Gasperini M, Tome JM, Shendure J. Towards a comprehensive catalogue of validated and target-
linked human enhancers. Nat Rev Genet. 2020;21(5):292-310. Epub 2020/01/29. doi: 10.1038/s41576-
019-0209-0. PubMed PMID: 31988385.
24.   Barski A, Cuddapah S, Cui K, Roh TY, Schones DE, Wang Z, et al. High-resolution profiling of
histone methylations in the human genome. Cell. 2007;129(4):823-37. Epub 2007/05/22. doi: S0092-
8674(07)00600-9 [pii]

10.1016/j.cell.2007.05.009 [doi]. PubMed PMID: 17512414.
25.   Johnson DS, Mortazavi A, Myers RM, Wold B. Genome-wide mapping of in vivo protein-DNA
interactions. Science. 2007;316(5830):1497-502. PubMed PMID: 17540862.
26.   Moore JE, Purcaro MJ, Pratt HE, Epstein CB, Shoresh N, Adrian J, et al. Expanded encyclopaedias
of DNA elements in the human and mouse genomes. Nature. 2020;583(7818):699-710. Epub
2020/07/29. doi: 10.1038/s41586-020-2493-4. PubMed PMID: 32728249; PubMed Central PMCID:
PMCPMC7410828.
27.   Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, Heravi-Moussavi A, et al. Integrative analysis
of 111 reference human epigenomes. Nature. 2015;518(7539):317-30. doi: 10.1038/nature14248.
28.   Consortium G. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis:
multitissue gene regulation in humans. Science. 2015;348(6235):648-60. Epub 2015/05/09. doi:
10.1126/science.1262110. PubMed PMID: 25954001; PubMed Central PMCID: PMCPMC4547484.
29.   Mei S, Qin Q, Wu Q, Sun H, Zheng R, Zang C, et al. Cistrome Data Browser: a data portal for ChIP-
Seq and chromatin accessibility data in human and mouse. Nucleic Acids Res. 2017;45(D1):D658-d62.
Epub 2016/10/30. doi: 10.1093/nar/gkw983. PubMed PMID: 27789702; PubMed Central PMCID:
PMCPMC5210658.
30.   Kleftogiannis D, Kalnis P, Bajic VB. Progress and challenges in bioinformatics approaches for
enhancer identification. Brief Bioinform. 2016;17(6):967-79. Epub 2015/12/05. doi:
10.1093/bib/bbv101. PubMed PMID: 26634919; PubMed Central PMCID: PMCPMC5142011.
31.   Bailey TL, Elkan C. Fitting a mixture model by expectation maximization to discover motifs in
biopolymers. Proc Int Conf Intell Syst Mol Biol. 1994;2:28-36. PubMed PMID: 7584402.
32.   Liu X, Brutlag DL, Liu JS. BioProspector: discovering conserved DNA motifs in upstream
regulatory regions of co-expressed genes. Pac Symp Biocomput. 2001:127-38. PubMed PMID: 11262934.
33.   Bailey TL. DREME: motif discovery in transcription factor ChIP-seq data. Bioinformatics.
2011;27(12):1653-9. Epub 2011/05/06. doi: btr261 [pii]

10.1093/bioinformatics/btr261 [doi]. PubMed PMID: 21543442; PubMed Central PMCID:
PMC3106199.
34.   Machanick P, Bailey TL. MEME-ChIP: motif analysis of large DNA datasets. Bioinformatics.
2011;27(12):1696-7. Epub 2011/04/14. doi: btr189 [pii]

10.1093/bioinformatics/btr189 [doi]. PubMed PMID: 21486936; PubMed Central PMCID:
PMC3106185.

35.   Hartmann H, Guthohrlein EW, Siebert M, Luehr S, Soding J. P-value-based regulatory motif
discovery using positional weight matrices. Genome Res. 2013;23(1):181-94. Epub 2012/09/20. doi:
gr.139881.112 [pii]

10.1101/gr.139881.112 [doi]. PubMed PMID: 22990209; PubMed Central PMCID: PMC3530678.
36.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple combinations of lineage-
determining transcription factors prime cis-regulatory elements required for macrophage and B cell
identities. Mol Cell. 2010;38(4):576-89. Epub 2010/06/02. doi: 10.1016/j.molcel.2010.05.004. PubMed
PMID: 20513432; PubMed Central PMCID: PMCPMC2898526.
37.   Sinha S. Discriminative motifs. Journal of computational biology : a journal of computational
molecular cell biology. 2003;10(3-4):599-615. Epub 2003/08/26. doi: 10.1089/10665270360688219
[doi]. PubMed PMID: 12935347.
38.   Bailey TL, Machanick P. Inferring direct DNA binding from ChIP-seq. Nucleic Acids Res. 2012.
Epub 2012/05/23. doi: gks433 [pii] 10.1093/nar/gks433 [doi]. PubMed PMID: 22610855.
39.   Whitington T, Frith MC, Johnson J, Bailey TL. Inferring transcription factor complexes from ChIP-
seq data. Nucleic Acids Res. 2011;39(15):e98. Epub 2011/05/24. doi: gkr341 [pii]

10.1093/nar/gkr341 [doi]. PubMed PMID: 21602262; PubMed Central PMCID: PMC3159476.
40.   Sun H, Guns T, Fierro AC, Thorrez L, Nijssen S, Marchal K. Unveiling combinatorial regulation
through the combination of ChIP information and in silico cis-regulatory module detection. Nucleic Acids
Res. 2012;40(12):e90. Epub 2012/03/17. doi: gks237 [pii]

10.1093/nar/gks237 [doi]. PubMed PMID: 22422841; PubMed Central PMCID: PMC3384348.
41.   Jiang P, Singh M. CCAT: Combinatorial Code Analysis Tool for transcriptional regulation. Nucleic
Acids Res. 2014;42(5):2833-47. Epub 2013/12/25. doi: 10.1093/nar/gkt1302. PubMed PMID: 24366875;
PubMed Central PMCID: PMCPMC3950699.
42.   Wingender E, Chen X, Fricke E, Geffers R, Hehl R, Liebich I, et al. The TRANSFAC system on gene
expression regulation. Nucleic Acids Res. 2001;29:281-3.
43.   Vlieghe D, Sandelin A, De Bleser PJ, Vleminckx K, Wasserman WW, van Roy F, et al. A new
generation of JASPAR, the open-access repository for transcription factor binding site profiles. Nucleic
Acids Res. 2006;34(Database issue):D95-7. PubMed PMID: 16381983.
44.   Yip KY, Cheng C, Bhardwaj N, Brown JB, Leng J, Kundaje A, et al. Classification of human genomic
regions based on experimentally determined binding sites of more than 100 transcription-related
factors. Genome biology. 2012;13(9):R48. Epub 2012/09/07. doi: gb-2012-13-9-r48 [pii]

10.1186/gb-2012-13-9-r48 [doi]. PubMed PMID: 22950945; PubMed Central PMCID: PMC3491392.
45.   Kheradpour P, Kellis M. Systematic discovery and characterization of regulatory motifs in
ENCODE TF binding experiments. Nucleic Acids Res. 2014;42(5):2976-87. Epub 2013/12/18. doi:
10.1093/nar/gkt1249. PubMed PMID: 24335146; PubMed Central PMCID: PMCPMC3950668.
46.   Niu M, Tabari E, Ni P, Su Z. Towards a map of cis-regulatory sequences in the human genome.
Nucleic Acids Res. 2018;46(11):5395-409. Epub 2018/05/08. doi: 10.1093/nar/gky338. PubMed PMID:
29733395; PubMed Central PMCID: PMCPMC6009671.
47.   Niu M, Tabari ES, Su Z. De novo prediction of cis-regulatory elements and modules through
integrative analysis of a large number of ChIP datasets. BMC Genomics. 2014;15(1):1047. doi:
10.186/471-2164-15-1047.
48.   Ernst J, Kellis M. ChromHMM: automating chromatin-state discovery and characterization.
Nature methods. 2012;9(3):215-6. Epub 2012/03/01. doi: 10.1038/nmeth.1906. PubMed PMID:
22373907; PubMed Central PMCID: PMCPMC3577932.

49.   Hoffman MM, Ernst J, Wilder SP, Kundaje A, Harris RS, Libbrecht M, et al. Integrative annotation
of chromatin elements from ENCODE data. Nucleic Acids Res. 2013;41(2):827-41. Epub 2012/12/12. doi:
gks1284 [pii]

10.1093/nar/gks1284 [doi]. PubMed PMID: 23221638; PubMed Central PMCID: PMC3553955.
50.   Firpi HA, Ucar D, Tan K. Discover regulatory DNA elements using chromatin signatures and
artificial neural network. Bioinformatics. 2010;26(13):1579-86. Epub 2010/05/11. doi: btq248 [pii]

10.1093/bioinformatics/btq248 [doi]. PubMed PMID: 20453004; PubMed Central PMCID:
PMC2887052.
51.   Rajagopal N, Xie W, Li Y, Wagner U, Wang W, Stamatoyannopoulos J, et al. RFECS: a random-
forest based algorithm for enhancer identification from chromatin state. PLoS Comput Biol.
2013;9(3):e1002968. Epub 2013/03/26. doi: 10.1371/journal.pcbi.1002968 [doi]

PCOMPBIOL-D-12-01314 [pii]. PubMed PMID: 23526891; PubMed Central PMCID: PMC3597546.
52.   Kleftogiannis D, Kalnis P, Bajic VB. DEEP: a general computational framework for predicting
enhancers. Nucleic Acids Res. 2015;43(1):e6. Epub 2014/11/08. doi: 10.1093/nar/gku1058. PubMed
PMID: 25378307; PubMed Central PMCID: PMCPMC4288148.
53.   Zerbino DR, Wilder SP, Johnson N, Juettemann T, Flicek PR. The ensembl regulatory build.
Genome Biol. 2015;16:56.(doi):10.1186/s13059-015-0621-5.
54.   Ashoor H, Kleftogiannis D, Radovanovic A, Bajic VB. DENdb: database of integrated human
enhancers. Database : the journal of biological databases and curation. 2015;2015. Epub 2015/09/08.
doi: 10.1093/database/bav085. PubMed PMID: 26342387; PubMed Central PMCID: PMCPmc4560934.
55.   Fishilevich S, Nudel R, Rappaport N, Hadar R, Plaschkes I, Iny Stein T, et al. GeneHancer:
genome-wide integration of enhancers and target genes in GeneCards. Database (Oxford). 2017;2017.
Epub 2017/06/13. doi: 10.1093/database/bax028. PubMed PMID: 28605766; PubMed Central PMCID:
PMCPMC5467550.
56.   Chen C, Zhou D, Gu Y, Wang C, Zhang M, Lin X, et al. SEA version 3.0: a comprehensive extension
and update of the Super-Enhancer archive. Nucleic Acids Res. 2020;48(D1):D198-d203. Epub
2019/11/02. doi: 10.1093/nar/gkz1028. PubMed PMID: 31667506; PubMed Central PMCID:
PMCPMC7145603.
57.   Kang R, Zhang Y, Huang Q, Meng J, Ding R, Chang Y, et al. EnhancerDB: a resource of
transcriptional regulation in the context of enhancers. Database (Oxford). 2019;2019. Epub 2019/01/29.
doi: 10.1093/database/bay141. PubMed PMID: 30689845; PubMed Central PMCID: PMCPMC6344666.
58.   Zhang G, Shi J, Zhu S, Lan Y, Xu L, Yuan H, et al. DiseaseEnhancer: a resource of human disease-
associated enhancer catalog. Nucleic Acids Res. 2018;46(D1):D78-d84. Epub 2017/10/24. doi:
10.1093/nar/gkx920. PubMed PMID: 29059320; PubMed Central PMCID: PMCPMC5753380.
59.   Gao T, Qian J. EnhancerAtlas 2.0: an updated resource with enhancer annotation in 586
tissue/cell types across nine species. Nucleic Acids Res. 2020;48(D1):D58-d64. Epub 2019/11/20. doi:
10.1093/nar/gkz980. PubMed PMID: 31740966; PubMed Central PMCID: PMCPMC7145677.
60.   Cheneby J, Gheorghe M, Artufel M, Mathelier A, Ballester B. ReMap 2018: an updated atlas of
regulatory regions from an integrative analysis of DNA-binding ChIP-seq experiments. Nucleic Acids Res.
2018;46(D1):D267-d75. Epub 2017/11/11. doi: 10.1093/nar/gkx1092. PubMed PMID: 29126285;
PubMed Central PMCID: PMCPMC5753247.
61.   Thurman RE, Rynes E, Humbert R, Vierstra J, Maurano MT, Haugen E, et al. The accessible
chromatin landscape of the human genome. Nature. 2012;489(7414):75-82. Epub 2012/09/08. doi:
nature11232 [pii]

10.1038/nature11232 [doi]. PubMed PMID: 22955617.

62.   Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition of native chromatin
for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome
position. Nature methods. 2013;10(12):1213-8. Epub 2013/10/08. doi: 10.1038/nmeth.2688. PubMed
PMID: 24097267; PubMed Central PMCID: PMCPmc3959825.
63.   Aday AW, Zhu LJ, Lakshmanan A, Wang J, Lawson ND. Identification of cis regulatory features in
the embryonic zebrafish genome through large-scale profiling of H3K4me1 and H3K4me3 binding sites.
Developmental biology. 2011;357(2):450-62. Epub 2011/03/26. doi: 10.1016/j.ydbio.2011.03.007.
PubMed PMID: 21435340; PubMed Central PMCID: PMCPMC3273848.
64.   Creyghton MP, Cheng AW, Welstead GG, Kooistra T, Carey BW, Steine EJ, et al. Histone H3K27ac
separates active from poised enhancers and predicts developmental state. Proc Natl Acad Sci U S A.
2010;107(50):21931-6. Epub 2010/11/26. doi: 1016071107 [pii]

10.1073/pnas.1016071107 [doi]. PubMed PMID: 21106759; PubMed Central PMCID: PMC3003124.
65.   Kim TH, Abdullaev ZK, Smith AD, Ching KA, Loukinov DI, Green RD, et al. Analysis of the
vertebrate insulator protein CTCF-binding sites in the human genome. Cell. 2007;128(6):1231-45. Epub
2007/03/27. doi: S0092-8674(07)00205-X [pii]

10.1016/j.cell.2006.12.048 [doi]. PubMed PMID: 17382889; PubMed Central PMCID: PMC2572726.
66.   Won KJ, Chepelev I, Ren B, Wang W. Prediction of regulatory elements in mammalian genomes
using chromatin signatures. BMC Bioinformatics. 2008;9:547. Epub 2008/12/20. doi: 1471-2105-9-547
[pii]

10.1186/1471-2105-9-547 [doi]. PubMed PMID: 19094206; PubMed Central PMCID: PMC2657164.
67.   Kwasnieski JC, Fiore C, Chaudhari HG, Cohen BA. High-throughput functional testing of ENCODE
segmentation predictions. Genome Res. 2014;24(10):1595-602. Epub 2014/07/19. doi:
10.1101/gr.173518.114. PubMed PMID: 25035418; PubMed Central PMCID: PMCPMC4199366.
68.   Dogan N, Wu W, Morrissey CS, Chen KB, Stonestrom A, Long M, et al. Occupancy by key
transcription factors is a more accurate predictor of enhancer activity than histone modifications or
chromatin accessibility. Epigenetics Chromatin. 2015;8:16. Epub 2015/05/20. doi: 10.1186/s13072-015-
0009-5. PubMed PMID: 25984238; PubMed Central PMCID: PMCPMC4432502.
69.   Catarino RR, Stark A. Assessing sufficiency and necessity of enhancer activities for gene
expression and the mechanisms of transcription activation. Genes Dev. 2018;32(3-4):202-23. Epub
2018/03/02. doi: 10.1101/gad.310367.117. PubMed PMID: 29491135; PubMed Central PMCID:
PMCPMC5859963.
70.   Arbel H, Basu S, Fisher WW, Hammonds AS, Wan KH, Park S, et al. Exploiting regulatory
heterogeneity to systematically identify enhancers with high accuracy. Proc Natl Acad Sci U S A.
2019;116(3):900-8. Epub 2019/01/02. doi: 10.1073/pnas.1808833115. PubMed PMID: 30598455;
PubMed Central PMCID: PMCPMC6338827.
71.   Goi C, Little P, Xie C. Cell-type and transcription factor specific enrichment of transcriptional
cofactor motifs in ENCODE ChIP-seq data. BMC Genomics. 2013;14 Suppl 5:S2. Epub 2014/02/26. doi:
1471-2164-14-S5-S2 [pii]

10.1186/1471-2164-14-S5-S2 [doi]. PubMed PMID: 24564528; PubMed Central PMCID:
PMC3852067.
72.   Li Y, Ni P, Zhang S, Li G, Su Z. ProSampler: an ultra-fast and accurate motif finder in large ChIP-
seq datasets for combinatory motif discovery. Bioinformatics. 2019. Epub 2019/05/10. doi:
10.1093/bioinformatics/btz290. PubMed PMID: 31070745.
73.   Allen JE, Pertea M, Salzberg SL. Computational gene prediction using multiple sources of
evidence. Genome Res. 2004;14(1):142-8. Epub 2004/01/07. doi: 10.1101/gr.1562804 [doi]

14/1/142 [pii]. PubMed PMID: 14707176; PubMed Central PMCID: PMC314291.

74.   Arnosti DN, Kulkarni MM. Transcriptional enhancers: Intelligent enhanceosomes or flexible
billboards? J Cell Biochem. 2005;94(5):890-8.

75.   Yanez-Cuna JO, Kvon EZ, Stark A. Deciphering the transcriptional cis-regulatory code. Trends
Genet. 2013;29(1):11-22. doi: 10.1016/j.tig.2012.09.007. Epub Oct 23.

76.   Vockley CM, Barrera A, Reddy TE. Decoding the role of regulatory element polymorphisms in
complex disease. Curr Opin Genet Dev. 2017;43:38-45. Epub 2016/12/17. doi:
10.1016/j.gde.2016.10.007. PubMed PMID: 27984826.

77.   Snetkova V, Skok JA. Enhancer talk. Epigenomics. 2018;10(4):483-98. Epub 2018/03/28. doi:
10.2217/epi-2017-0157. PubMed PMID: 29583027; PubMed Central PMCID: PMCPMC5925435.

78.   Li Q, Peterson KR, Fang X, Stamatoyannopoulos G. Locus control regions. Blood.
2002;100(9):3077-86. Epub 2002/10/18. doi: 10.1182/blood-2002-04-1104. PubMed PMID: 12384402;
PubMed Central PMCID: PMCPMC2811695.

79.   Visel A, Minovitsky S, Dubchak I, Pennacchio LA. VISTA Enhancer Browser--a database of tissue-
specific human enhancers. Nucleic Acids Res. 2007;35(Database issue):D88-92. Epub 2006/11/30. doi:
10.1093/nar/gkl822. PubMed PMID: 17130149; PubMed Central PMCID: PMCPMC1716724.

80.   Kulakovskiy IV, Vorontsov IE, Yevshin IS, Sharipov RN, Fedorova AD, Rumynskiy EI, et al.
HOCOMOCO: towards a complete collection of transcription factor binding models for human and
mouse via large-scale ChIP-Seq analysis. Nucleic Acids Res. 2018;46(D1):D252-d9. Epub 2017/11/16. doi:
10.1093/nar/gkx1106. PubMed PMID: 29140464; PubMed Central PMCID: PMCPMC5753240.

81.   Mathelier A, Fornes O, Arenillas DJ, Chen CY, Denay G, Lee J, et al. JASPAR 2016: a major
expansion and update of the open-access database of transcription factor binding profiles. Nucleic Acids
Res. 2016;44(D1):D110-5. Epub 2015/11/05. doi: 10.1093/nar/gkv1176. PubMed PMID: 26531826;
PubMed Central PMCID: PMCPMC4702842.

82.   Lambert SA, Jolma A, Campitelli LF, Das PK, Yin Y, Albu M, et al. The Human Transcription
Factors. Cell. 2018;175(2):598-9. Epub 2018/10/06. doi: 10.1016/j.cell.2018.09.045. PubMed PMID:
30290144.

83.   Ambrosini G, Vorontsov I, Penzar D, Groux R, Fornes O, Nikolaeva DD, et al. Insights gained from
a comprehensive all-against-all transcription factor binding motif benchmarking study. Genome Biol.
2020;21(1):114. Epub 2020/05/13. doi: 10.1186/s13059-020-01996-3. PubMed PMID: 32393327.

84.   Perrot CY, Gilbert C, Marsaud V, Postigo A, Javelaud D, Mauviel A. GLI2 cooperates with ZEB1 for
transcriptional repression of CDH1 expression in human melanoma cells. Pigment Cell Melanoma Res.
2013;26(6):861-73. Epub 2013/07/31. doi: 10.1111/pcmr.12149. PubMed PMID: 23890107.

85.   Koyabu Y, Nakata K, Mizugishi K, Aruga J, Mikoshiba K. Physical and functional interactions
between Zic and Gli proteins. J Biol Chem. 2001;276(10):6889-92. Epub 2001/01/12. doi:
10.1074/jbc.C000773200. PubMed PMID: 11238441.

86.   Sánchez-Tilló E, de Barrios O, Valls E, Darling DS, Castells A, Postigo A. ZEB1 and TCF4
reciprocally modulate their transcriptional activities to regulate Wnt target gene expression. Oncogene.
2015;34(46):5760-70. Epub 2015/09/21. doi: 10.1038/onc.2015.352. PubMed PMID: 26387539.

87.   Mendoza-Parra MA, Van Gool W, Mohamed Saleem MA, Ceschin DG, Gronemeyer H. A quality
control system for profiles obtained by ChIP sequencing. Nucleic Acids Res. 2013;41(21):e196. Epub
2013/09/17. doi: 10.1093/nar/gkt829. PubMed PMID: 24038469; PubMed Central PMCID:
PMCPMC3834836.

88.   Marinov GK, Kundaje A, Park PJ, Wold BJ. Large-scale quality analysis of published ChIP-seq data.
G3 (Bethesda). 2014;4(2):209-23. doi: 10.1534/g3.113.008680.

89.   Devailly G, Mantsoki A, Michoel T, Joshi A. Variable reproducibility in genome-scale public data:
A case study using ENCODE ChIP sequencing resource. FEBS Lett. 2015;589(24 Pt B):3866-70. Epub

2015/12/02. doi: 10.1016/j.febslet.2015.11.027. PubMed PMID: 26619763; PubMed Central PMCID:
PMCPMC4686001.
90. Stergachis AB, Neph S, Reynolds A, Humbert R, Miller B, Paige SL, et al. Developmental fate and
cellular maturity encoded in human regulatory DNA landscapes. Cell. 2013;154(4):888-903. Epub
2013/08/21. doi: S0092-8674(13)00891-X [pii]

10.1016/j.cell.2013.07.020 [doi]. PubMed PMID: 23953118.
91. Li L, Wunderlich Z. An Enhancer's Length and Composition Are Shaped by Its Regulatory Task.
Front Genet. 2017;8:63. Epub 2017/06/08. doi: 10.3389/fgene.2017.00063. PubMed PMID: 28588608;
PubMed Central PMCID: PMCPMC5440464.
92. Pott S, Lieb JD. What are super-enhancers? Nat Genet. 2014;47(1):8-12. doi: 0.1038/ng.3167.
93. Cooper GM, Goode DL, Ng SB, Sidow A, Bamshad MJ, Shendure J, et al. Single-nucleotide
evolutionary constraint scores highlight disease-causing mutations. Nat Methods. 2010;7(4):250-1. doi:
10.1038/nmeth0410-250.
94. Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. Detection of nonneutral substitution rates on
mammalian phylogenies. Genome Res. 2010;20(1):110-21. Epub 2009/10/28. doi:
10.1101/gr.097857.109. PubMed PMID: 19858363; PubMed Central PMCID: PMCPMC2798823.
95. Visel A, Prabhakar S, Akiyama JA, Shoukry M, Lewis KD, Holt A, et al. Ultraconservation identifies
a small subset of extremely constrained developmental enhancers. Nat Genet. 2008;40(2):158-60.
PubMed PMID: 18176564.
96. Bejerano G, Pheasant M, Makunin I, Stephen S, Kent WJ, Mattick JS, et al. Ultraconserved
elements in the human genome. Science. 2004;304(5675):1321-5. Epub 2004/05/08. doi:
10.1126/science.1098119. PubMed PMID: 15131266.
97. Katzman S, Kern AD, Bejerano G, Fewell G, Fulton L, Wilson RK, et al. Human genome
ultraconserved elements are ultraselected. Science. 2007;317(5840):915. Epub 2007/08/19. doi:
10.1126/science.1142430. PubMed PMID: 17702936.
98. Li S, Kvon EZ, Visel A, Pennacchio LA, Ovcharenko I. Stable enhancers are active in development,
and fragile enhancers are associated with evolutionary adaptation. Genome Biol. 2019;20(1):140. Epub
2019/07/17. doi: 10.1186/s13059-019-1750-z. PubMed PMID: 31307522; PubMed Central PMCID:
PMCPMC6631995.
99. Woolfe A, Goodson M, Goode DK, Snell P, McEwen GK, Vavouri T, et al. Highly conserved non-
coding sequences are associated with vertebrate development. PLoS Biol. 2005;3(1):e7. Epub
2005/01/05. doi: 10.1371/journal.pbio.0030007 [doi]. PubMed PMID: 15630479; PubMed Central
PMCID: PMC526512.
100. Forrest AR, Kawaji H, Rehli M, Baillie JK, de Hoon MJ, Haberle V, et al. A promoter-level
mammalian expression atlas. Nature. 2014;507(7493):462-70. Epub 2014/03/29. doi:
10.1038/nature13182. PubMed PMID: 24670764; PubMed Central PMCID: PMCPMC4529748.
101. Andersson R, Gebhard C, Miguel-Escalada I, Hoof I, Bornholdt J, Boyd M, et al. An atlas of active
enhancers across human cell types and tissues. Nature. 2014;507(7493):455-61. Epub 2014/03/29. doi:
nature12787 [pii]

10.1038/nature12787 [doi]. PubMed PMID: 24670763.
102. Buniello A, MacArthur JAL, Cerezo M, Harris LW, Hayhurst J, Malangone C, et al. The NHGRI-EBI
GWAS Catalog of published genome-wide association studies, targeted arrays and summary statistics
2019. Nucleic Acids Res. 2019;47(D1):D1005-D12. Epub 2018/11/18. doi: 10.1093/nar/gky1120. PubMed
PMID: 30445434; PubMed Central PMCID: PMCPMC6323933.
103. Landrum MJ, Lee JM, Benson M, Brown GR, Chao C, Chitipiralla S, et al. ClinVar: improving
access to variant interpretations and supporting evidence. Nucleic Acids Res. 2018;46(D1):D1062-d7.

Epub 2017/11/23. doi: 10.1093/nar/gkx1153. PubMed PMID: 29165669; PubMed Central PMCID:
PMCPMC5753237.
104. Villar D, Berthelot C, Aldridge S, Rayner TF, Lukk M, Pignatelli M, et al. Enhancer evolution across
20 mammalian species. Cell. 2015;160(3):554-66. Epub 2015/01/31. doi: 10.1016/j.cell.2015.01.006.
PubMed PMID: 25635462; PubMed Central PMCID: PMCPMC4313353.
105. Young RS, Kumar Y, Bickmore WA, Taylor MS. Bidirectional transcription initiation marks
accessible chromatin and is not specific to enhancers. Genome Biol. 2017;18(1):242. Epub 2017/12/30.
doi: 10.1186/s13059-017-1379-8. PubMed PMID: 29284524; PubMed Central PMCID: PMCPMC5747114.
106. Chereji RV, Eriksson PR, Ocampo J, Prajapati HK, Clark DJ. Accessibility of promoter DNA is not
the primary determinant of chromatin-mediated gene regulation. Genome Res. 2019;29(12):1985-95.
Epub 2019/09/13. doi: 10.1101/gr.249326.119. PubMed PMID: 31511305; PubMed Central PMCID:
PMCPMC6886500.
107. MacArthur J, Bowler E, Cerezo M, Gil L, Hall P, Hastings E, et al. The new NHGRI-EBI Catalog of
published genome-wide association studies (GWAS Catalog). Nucleic Acids Res. 2017;45(D1):D896-D901.
Epub 2016/11/29. doi: 10.1093/nar/gkw1133. PubMed PMID: 27899670; PubMed Central PMCID:
PMCPMC5210590.
108. Gorkin DU, Barozzi I, Zhao Y, Zhang Y, Huang H, Lee AY, et al. An atlas of dynamic chromatin
landscapes in mouse fetal development. Nature. 2020;583(7818):744-51. Epub 2020/07/31. doi:
10.1038/s41586-020-2093-3. PubMed PMID: 32728240; PubMed Central PMCID: PMCPMC7398618.
109. Bernstein BE, Birney E, Dunham I, Green ED, Gunter C, Snyder M. An integrated encyclopedia of
DNA elements in the human genome. Nature. 2012;489(7414):57-74. Epub 2012/09/08. doi:
nature11247 [pii]

10.1038/nature11247 [doi]. PubMed PMID: 22955616; PubMed Central PMCID: PMC3439153.
110. Khan A, Zhang X. dbSUPER: a database of super-enhancers in mouse and human genome.
Nucleic Acids Res. 2015. Epub 2015/10/07. doi: 10.1093/nar/gkv1002. PubMed PMID: 26438538.
111. Dreos R, Ambrosini G, Cavin Perier R, Bucher P. EPD and EPDnew, high-quality promoter
resources in the next-generation sequencing era. Nucleic Acids Res. 2013;41(Database issue):D157-64.
Epub 2012/11/30. doi: 10.1093/nar/gks1233. PubMed PMID: 23193273; PubMed Central PMCID:
PMCPMC3531148.
112. Dimitrieva S, Bucher P. UCNEbase--a database of ultraconserved non-coding elements and
genomic regulatory blocks. Nucleic Acids Res. 2013;41(Database issue):D101-9. Epub 2012/11/30. doi:
10.1093/nar/gks1092. PubMed PMID: 23193254; PubMed Central PMCID: PMCPMC3531063.
113. Wilderman A, VanOudenhove J, Kron J, Noonan JP, Cotney J. High-Resolution Epigenomic Atlas
of Human Embryonic Craniofacial Development. Cell Rep. 2018;23(5):1581-97. Epub 2018/05/03. doi:
10.1016/j.celrep.2018.03.129. PubMed PMID: 29719267; PubMed Central PMCID: PMCPMC5965702.
114. Hoffman MM, Buske OJ, Wang J, Weng Z, Bilmes JA, Noble WS. Unsupervised pattern discovery
in human chromatin structure through genomic segmentation. Nat Methods. 2012;9(5):473-6. Epub
2012/03/20. doi: nmeth.1937 [pii]

10.1038/nmeth.1937 [doi]. PubMed PMID: 22426492; PubMed Central PMCID: PMC3340533.
115. Gao T, He B, Liu S, Zhu H, Tan K, Qian J. EnhancerAtlas: a resource for enhancer annotation and
analysis in 105 human cell/tissue types. Bioinformatics. 2016;32(23):3543-51. Epub 2016/08/16. doi:
10.1093/bioinformatics/btw495. PubMed PMID: 27515742; PubMed Central PMCID: PMCPMC5181530.
116. Pennacchio LA, Bickmore W, Dean A, Nobrega MA, Bejerano G. Enhancers: five essential
questions. Nat Rev Genet. 2013;14(4):288-95. Epub 2013/03/19. doi: nrg3458 [pii]

10.1038/nrg3458 [doi]. PubMed PMID: 23503198.

117. Kellis M, Wold B, Snyder MP, Bernstein BE, Kundaje A, Marinov GK, et al. Defining functional
DNA elements in the human genome. Proc Natl Acad Sci U S A. 2014;111(17):6131-8. doi:
10.1073/pnas.1318948111. Epub 2014 Apr 21.
118. Snyder MP, Gingeras TR, Moore JE, Weng Z, Gerstein MB, Ren B, et al. Perspectives on ENCODE.
Nature. 2020;583(7818):693-8. Epub 2020/07/29. doi: 10.1038/s41586-020-2449-8. PubMed PMID:
32728248; PubMed Central PMCID: PMCPMC7410827.
119. Wang X, He L, Goggin SM, Saadat A, Wang L, Sinnott-Armstrong N, et al. High-resolution
genome-wide functional dissection of transcriptional regulatory regions and nucleotides in human. Nat
Commun. 2018;9(1):5380. Epub 2018/12/21. doi: 10.1038/s41467-018-07746-1. PubMed PMID:
30568279; PubMed Central PMCID: PMCPMC6300699.
120. Stamatoyannopoulos JA. What does our genome encode? Genome Res. 2012;22(9):1602-11.
Epub 2012/09/08. doi: 22/9/1602 [pii]

10.1101/gr.146506.112 [doi]. PubMed PMID: 22955972; PubMed Central PMCID: PMC3431477.
121. King DC, Taylor J, Elnitski L, Chiaromonte F, Miller W, Hardison RC. Evaluation of regulatory
potential and conservation scores for detecting cis-regulatory modules in aligned mammalian genome
sequences. Genome Res. 2005;15(8):1051-60. Epub 2005/07/15. doi: 10.1101/gr.3642605. PubMed
PMID: 16024817; PubMed Central PMCID: PMCPMC1182217.
122. Weirauch MT, Yang A, Albu M, Cote AG, Montenegro-Montero A, Drewe P, et al. Determination
and inference of eukaryotic transcription factor sequence specificity. Cell. 2014;158(6):1431-43. doi:
10.016/j.cell.2014.08.009.
123. Zhang S, Jiang L, Du C, Su Z. SPIC: A novel information contents based similarity metric for
comparing transcription factor binding site motifs. BMC Syst Biol. 2013;7(Suppl 2):S14. Epub S14. doi:
doi:10.1186/1752-0509-7-S2-S14.
124. van Dongen S, Abreu-Goodger C. Using MCL to extract clusters from networks. Methods in
molecular biology (Clifton, NJ). 2012;804:281-95. Epub 2011/12/07. doi: 10.1007/978-1-61779-361-5_15
[doi]. PubMed PMID: 22144159.
125. Vockley CM, McDowell IC, D'Ippolito AM, Reddy TE. A long-range flexible billboard model of
gene activation. Transcription. 2017;8(4):261-7. Epub 2017/06/10. doi:
10.1080/21541264.2017.1317694. PubMed PMID: 28598247; PubMed Central PMCID:
PMCPMC5574526.

1348