Back to feedback: aberrant sensorimotor control in music performance under

pressure

Shinichi Furuya [1][3] [*], Reiko Ishimaru [2] [*], Takanori Oku [1], and Noriko Nagata [2]

1) Sony Computer Science Laboratories Inc. (Sony CSL), Japan

2) School of Science and Technology, Kwansei Gakuin University, Japan

3) Sophia University, Japan

*: These authors are equally contributing co-first authors.

Keywords: choking under pressure; feedback control; robustness of sensorimotor

control; psychological stress; music performance anxiety

Running Title: Sensorimotor instability by state anxiety

Number of Figures: 2 (2 color figures included)

Correspondence to: Shinichi Furuya, PhD

Sony Computer Science Laboratories Inc. (Sony CSL), Japan

E-mail: furuya@csl.sony.co.jp

Conflicts of interest: All authors (SF, RI, TO, NN) have no conflicts of financial and/or

non-financial interest to declare.

21
22

ABSTRACT

Precisely timed production of dexterous actions is often destabilized in anxiogenic situations. Previous studies demonstrated that cognitive functions such as attention and working memory as well as autonomic nervous functions are susceptible to induced anxiety in skillful performance while playing sports or musical instruments. However, it is not known whether the degradation of motor functions, sensory perception, or sensorimotor control underlies such a compromise of skillful performance due to psychophysiological distress. Here, we addressed this issue through a series of behavioral experiments, which demonstrated no detrimental effects of the stress on the perceptual accuracy and precision of the finger movements in pianists. By contrast, after transiently delaying the timing of tone production while playing the piano, the local tempo was abnormally disrupted only under pressure. The results suggest that psychological stress degraded the temporal stability of movement control due to an abnormal increase in sensory feedback gain but not temporal perception or motor precision. A learning experiment further demonstrated that the temporal instability of auditory-motor control under pressure was alleviated after practicing piano while ignoring delayed auditory feedback but not after practicing while compensating for the delayed feedback. Together, these findings suggest an abnormal transition from feedforward to feedback control in expert piano performance in anxiogenic situations, which can be mitigated through specialized sensorimotor training that involves piano practice while volitionally ignoring the delayed provision of auditory feedback.

INTRODUCTION

Skillful behaviors are readily spoiled in anxiogenic situations such as the Olympic games and music competitions. Suboptimal performance of skillful actions due to psychological stress and fear under pressure is a common problem across trained individuals, such as athletes, musicians, and surgeons, and sometimes affects the future career prospects of these individuals. State anxiety typically compromises spatiotemporal control of dexterous movements [1-6]. For example, stuttering in speech is exaggerated under pressure [7], which indicates the detrimental effects of psychological stress on precisely timed execution of motor sequences. Similarly, psychological stress triggers rhythmic distortion of sequential finger movements in musical performance [8-10]. These deleterious effects of psychological stress on motor precision have been extensively argued in terms of their relation to abnormal attention under pressure [1,5,11-14]. For example, skilled baseball players were more aware of the direction of bat motion in a higher pressure condition, and it was speculated that the exaggerated attention to the well-learned processes of skill execution led to abnormal modulation of skill execution and thereby increased the movement variability under pressure [5]. In contrast to cognitive malfunctions such as abnormal attention, it has not been elucidated what malfunctions in sensory-motor processes underlie the degradation of precision of skillful motor actions due to psychological stress. Understanding this is essential to shed light on neuropsychological mechanisms subserving the robustness and flexibility of sensorimotor skills in various environments.

At least three malfunctions of the sensorimotor system are candidates that may compromise timing control of motor sequences due to psychophysiological distress. The

first candidate is abnormal temporal perception due to psychological stress. It has been reported that fear and anxiety distort time perception, such as interval timing and simultaneity [15-17]. For example, a recent study showed that experimentally inducing anxiety resulted in underestimating the duration of temporal intervals [17]. The second candidate is abnormal motor functions under pressure. It has been reported that induced stress elevated finger force fluctuation during a precision grip [18,19], which suggests the degradation of motor functions in stressful situations. However, most of these previous studies targeted individuals without any history of extensive training, which questions the time perception and motor functions of experts under pressure.

The third candidate is sensory feedback control of movements. A key element of skillful behaviors is the robustness of skills against external perturbation. The balance between the robustness and flexibility of sensorimotor skills dynamically shifts over a course of learning [20-22]. At the early stage of skill acquisition, a large gain of sensory feedback enables fast skill improvements with training. Through extensive training, such a skill reaches its plateau and is stabilized. The stabilized skill is minimally susceptible to sensory perturbations due to feedforward control, which has been demonstrated in trained individuals [23]. For example, singers but not nonmusicians can keep singing accurately even by artificially shifting the pitch of the voice [24] or by blocking somatosensory afferent information via vocal-fold anesthesia [25]. Similarly, during piano playing, delayed tone production perturbs the subsequent keystroke motions to a lesser extent in expert pianists than in nonmusicians [26]. Accordingly, it is possible that psychological pressure destabilizes experts' skills, which mediates abnormal increases in the timing fluctuation of skilled motor actions under pressure [9].

Here, we addressed the effects of psychological stress on time perception, motor functions, and sensorimotor control of expert pianists. We focused on musicians because many musicians suffer from state anxiety when performing on the stage, so-called performance anxiety [27,28], which occurs in up to 90% of musicians. The stability of sensorimotor control was assessed by providing transient sensory perturbation during piano playing [29]. By using sensory perturbation, we also tested whether sensorimotor training can alleviate the detrimental effects of psychological stress on sensory-motor skills.

MATERIALS & METHODS

Participants

In the present study, 11 (7 females, 25.4 ± 4.9 yrs old) and 30 (23 females, 24.1 ± 6.0 yrs old) right-handed pianists with no history of neurological disorders participated in experiments 1 and 2, respectively. All of the pianists majored in piano performance in a musical conservatory and/or had extensive and continuous private piano training under the supervision of a professional pianist/piano professor. The age at which the participants had started to play the piano was before 8 years old. In accordance with the Declaration of Helsinki, the experimental procedures were explained to all participants. Informed consent was obtained from all participants prior to participating in the experiment, and the experimental protocol was approved by the ethics committee of Sophia University.

Experimental Design

The present study consists of 3 experiments under both low-pressure (LP) and

high-pressure (HP) conditions. To circumvent confounding effects of familiarization

with the experimental situation and stress induction, participants were different across

the experiments.

Experiment 1 was designed to test whether psychological stress influences the

perception of the duration of time interval based on auditory stimuli, the agility and

accuracy of movements, and sensory feedback control of movements in piano

performance. In total, 3 tests were carried out, which assessed the perception of the

auditory time interval ("*perception test*"), finger motor agility and accuracy ("*motor*

test"), and auditory feedback control of movements ("*sensorimotor test*") in the LP and

HP conditions. The perception test provided 2 sets of auditory stimuli, each of which

consisted of 2 successive tones. The first "reference" stimulus was 2 successive tones

with a tempo of 116 beats per minute (bpm) (i.e., the intertone interval was

approximately 129.3 ms), whereas the second stimulus was 2 tones with one of 7

different tempi ranging from 110 bpm to 122 bpm with a resolution of 2 bpm. At each

trial, each participant was asked to judge whether the second stimulus was the same or

different from the first stimulus in terms of the intertone interval. In total, 28 trials were

performed, the order of which was randomized across the participants. The motor test

asked the participants first to perform repetitive strikes of a piano key with each of five

digits of the right hand as fast as possible for 5 seconds ("the fastest single finger

tapping") [30] and second to play the first 8 tones of the piece of Op. 10 No. 8, composed

by Frédéric François Chopin, on the piano as fast as possible ("the fastest piano

performance"). Here, the second motor task consisted of tones of A5, G5, F5, C5, A4,

G4, F4, and C4, which were played with the fingering of the ring, middle, index, thumb,

ring, middle, index, and thumb, respectively. During the performance of these motor

tests, the piano tones were deprived (i.e., muted) so that auditory feedback could not

affect movement execution. These motor tests evaluated both the speed and the timing

precision of the sequential keystrokes with a single finger and with multiple fingers by

computing the mean and the standard deviation of the number of keystrokes per second

within a trial. The sensorimotor test asked the participants to play a short melody

extracted from Op. 10 No. 8, composed by Frédéric François Chopin, with only the

right hand (the first 14 bars) on an acoustic piano with MIDI sensors (YUS1SH,

Yamaha, Co.) at a tempo of 116 bpm (interkeystroke interval = 129.3 msec) and loud

(mezzo-forte). This piece was chosen first because it is technically challenging enough

to elicit anxiety for many pianists (note that this piece has been played at many piano

competitions and during the entrance exams for many musical conservatories) and

second because it requires high timing precision of the keystrokes during fast tempo

performance with little effects of emotional or aesthetic expression [9]. The target melody

was to be played with legato touch, meaning that a key was not released until the next

key was depressed. On the day of the experiment, participants were initially asked to

become familiar with the piano by practicing for 10 min, without using a metronome,

prior to the experiment. After the familiarization session, all participants were able to

play the target melody with the predetermined fingering at the target tempo without

pitch errors. During the sensorimotor test, prior to the task performance, 16 metronome

tones were provided to a cue of the tempo (i.e., 116 bpm), and then the pianists were

asked to play without the metronome sounds. While playing the target melody, the

timing of tone production was artificially delayed by 80 ms over 4 successive strikes

(i.e., delayed auditory feedback) as auditory perturbation [26,29,31], which occurred 3 times

within the melody. The notes with delayed tone production were not told to the pianists

in advance and were randomized across the pianists so that each pianist could not

predict which keystrokes elicited delayed tone production. For experiment 1, the order

of the perception, motor, and sensorimotor tests was randomized across the participants.

In the LP condition, the experimenter sat behind each pianist and out of sight. In the HP

session, another experimenter appeared and adopted a strict and unfriendly behavior;

this experimenter stood next to the participant and watched and evaluated the

performance. In addition, the performance was videotaped by a camera mounted in front

of the performer. Prior to initiating the HP session, the experimenter told the participants

that he/she would evaluate the performance. This experimental design was validated and

adopted in our previous studies [8,9].

Experiment 2 was designed to assess the effects of short-term piano practicing

with the provision of transiently delayed auditory feedback on the disruption of piano

performance following auditory perturbation. The experiment consisted of 4 successive

sessions in the following order: playing in the LP condition (LP_{pre}), intervention,

playing in the LP condition (LP_{post}), and playing in the HP condition (HP). Thirty

pianists were randomly assigned to 3 groups undergoing different interventions. At the

intervention session, they played the first 8 notes of the target melody over 200 trials.

The first control group was asked to play the 8 notes without any artificial delay in the

timing of the provision of auditory feedback ("normal" group). The second group was

instructed to play while ignoring the 80-ms delayed tone production, which occurred at

each of the last 4 notes ("ignore" group). The third group was instructed to play while

compensating for the delayed tone production at each of the last 4 notes so that the

tones could sound with consistent intertone intervals across the 8 notes ("compensate"

group). This means that the pianists in this group had to strike the keys 80 ms prior to

the timing at which the tone should be elicited. Prior to each trial, 4 metronome tones

were provided as a cue for the target tempo, and then the participants began playing

without the metronome cue. The intervention was designed based on previous studies in

which participants were instructed to either ignore or compensate for the altered

auditory feedback during movements [24,32,33].

Data Acquisition

We recorded MIDI data from the piano by using a custom-made script written

in JAVA [26]. This script allows for artificially delaying the timing of tone production

during piano playing by waiting for a predetermined duration until the MIDI signal is

elicited to the piano. Based on the MIDI data collected by this script during the

experiments, we derived the time at which each key was depressed and released. Based

on this information, the interkeystroke interval was computed. The heart rate of each

pianist was also recorded every second during the task performance in the experiments

(Mio Slice, Mio Global Inc.) and data were analyzed in an offline manner. In

experiment 1 only, the heart rate recordings from 3 pianists were contaminated by

unpredictable noises, resulting in missing data; therefore, the data from these pianists

were not used for the subsequent analyses.

Data Analysis

Using MIDI information during piano playing, we defined the timing error as

the temporal gap between two successive keypresses (i.e., between the measured and

ideal presses). Here, the interkeystroke interval of the ideal presses was defined based

on the tempo provided by the metronome. The timing error was computed for each of

the strikes and was then averaged across the strikes for each participant under each of

the HP and LP conditions. For the perception test of experiment 1, the perceptual

threshold was computed as 50% of the sigmoid function fitted to the time interval

detection probability curve [30].

Statistics

For datasets of the perception test and motor test (the fastest piano

performance) in experiment 1, a paired t-test was performed for a comparison between

the HP and LP conditions ($p = 0.05$). If the dataset did not follow the normal distribution

based on the Kolmogorov-Smirnov test, a nonparametric Wilcoxon test was performed.

For the heart rate in experiment 1, a two-way repeated-measures ANOVA was

performed with task (3 levels: perception, motor, sensorimotor tests) and condition (2

levels: LP and HP) as within-subject independent variables. For the fastest finger

tapping at the motor test in experiment 1, a two-way repeated-measures ANOVA was

performed with digit (5 levels: thumb, index, middle, ring, and little fingers) and

condition (2 levels: LP and HP) as within-subject independent variables. For the timing

error at the sensorimotor test in experiment 1, a two-way repeated-measures ANOVA

was performed with strike (4 levels: the first, second, third, and fourth strikes following

provision of the delayed tone production) and condition (2 levels: LP and HP) as

within-subject independent variables. With respect to experiment 2, a two-way

mixed-design ANOVA was performed with group (3 levels: normal, ignore, and

compensate) and strike (4 levels: the first, second, third, and fourth strikes following the

provision of delayed tone production) as independent variables. Mauchly's test was

used to test for sphericity prior to performing each ANOVA, and for nonspherical data,

the Greenhouse-Geisser correction was performed. Post hoc tests with correction for

multiple comparisons [34] were performed in the case of significance. These statistical

analyses were performed with R statistical software (ver. 3.2.3.).

RESULTS

Effects of psychological stress on perceptual, motor, and sensorimotor functions

Figure 1 illustrates the summarized results of experiment 1 that assessed the effects of psychological stress on heart rate and the perception of the time interval (i.e. duration perception), finger motor dexterity (i.e., motor function), and feedback control of movements based on sensory information (i.e., sensorimotor control) in the pianists.

Figure 1 inserted here

Figure 1A displays the group means of the average heart rate during each of the perception, motor, and sensorimotor control tasks in the pianists in the HP and LP conditions. During each of the three tasks, the heart rate was higher for the HP condition than for the LP condition. A two-way repeated-measures ANOVA demonstrated main effects of both task (3 levels: task 1-3) and condition (2 levels: HP and LP) on heart rate (task: $F(2,14) = 19.81$, $p = 8.27 \times 10^{-5}$, $\eta^2 = 0.103$; condition: $F(1, 7) = 11.03$, $p = 0.01$, $\eta^2 = 0.069$) but no interaction effect (task × condition: $F(2, 14) = 2.47$, $p = 0.12$, $\eta^2 = 0.008$).

Figure 1B illustrates the results of the psychophysics experiment that evaluated the threshold of the auditory perception of the duration of the time interval between two successive tones in pianists in the HP and LP conditions (i.e., *the perception test*). A paired t-test yielded no significant difference between the conditions ($t(10) = 0.024$, $p = 0.981$). This indicates no effects of psychological stress on the auditory perception of the time interval between the two tones.

Figure 1C displays the group means of the maximum tapping rate by each of the five digits in pianists during the HP and LP conditions (i.e. *the motor test*). A two-way repeated-measure ANOVA yielded neither an interaction effect between finger

and condition ($F(4, 40) = 0.154$, $p = 0.96$, $\eta^2 = 9.60 \times 10^{-5}$) nor a main effect of

condition ($F(1, 10) = 3.16 \times 10^{-4}$, $p = 0.99$, $\eta^2 = 5.57 \times 10^{-7}$) but a significant main effect

of finger ($F(4, 40) = 6.27$, $p = 5.14 \times 10^{-4}$, $\eta^2 = 0.10$). This indicates that finger tapping

ability did not differ irrespective of the pressure level. Similarly, with respect to the

variability of the number of keystrokes within a trial during the fastest finger tapping,

neither an interaction effect between finger and condition ($F(4, 40) = 0.74$, $p = 0.570$,

$\eta^2 = 0.015$) nor a condition effect ($F(1, 10) = 0.40$, $p = 0.539$, $\eta^2 = 0.002$) were observed.

This indicates no effect of psychological stress on the timing precision of the fastest

finger tapping movements.

Figure 1D displays the maximum number of keystrokes per second while

playing an excerpt of a melody that consists of 8 successive tones as fast as possible

without auditory feedback being provided (i.e., *the motor test*). A paired t-test yielded

no significant difference between the conditions ($t(10) = 0.05$, $p = 0.962$). Similarly,

with respect to the variability of the number of keystrokes within a trial while playing

this melody as fast as possible, a paired t-test identified no significant difference

between the conditions ($t(10) = -0.37$, $p = 0.721$). These results indicate no effects of

psychological stress on both the speed and the accuracy of playing a short melody.

Figure 1E illustrates the timing error of the piano keypresses following the

provision of the artificially delayed tone production in the pianists during the HP and LP

conditions (i.e., *the sensorimotor test*). A positive value indicates transient slow-down of

the local tempo following the perturbation. A two-way repeated-measure ANOVA

yielded significant main effects of both condition ($F(1, 9) = 15.49$, $p = 0.003$, $\eta^2 =$

0.059) and strike ($F(3, 27) = 5.96$, $p = 0.003$, $\eta^2 = 0.29$) but no interaction effect of

condition and strike ($F(3, 27) = 1.42$, $p = 0.260$, $\eta^2 = 0.04$). To see the main effect of the

condition, the average value across the 4 strikes was computed for each of the HP and

LP conditions (Fig. 1F). The effect of psychological stress on the timing error of the

keypresses was evident, as shown by the larger timing error in the HP condition

compared with the LP condition. A post hoc pairwise comparison identified a significant

difference in the timing error of the keypresses following delayed tone production

between the conditions ($p = 0.013$). These results indicate that pianists slowed down the

local tempo of piano playing following the transiently delayed tone production to a

larger extent in a higher pressure condition.

Effects of sensorimotor training

Experiment 2 tested whether practicing with the provision of delayed auditory

feedback would influence the rhythmic distortion of the keypresses following the

perturbation in piano performance under pressure. Figure 2 illustrates differences in the

effects of psychological stress on motor responses to transiently delayed tone

production among the 3 groups with different interventions (i.e. the normal, ignore, and

compensate groups). Here, the value at each strike indicates a difference in the timing

error of the keystrokes between the HP condition and the LP_{pre} condition, which was

computed to minimize the confounding effects of possible individual differences in the

interkeystroke interval across the groups. A two-way mixed-design ANOVA

demonstrated main effects of both group ($F_{(2, 27)} = 5.85$, $p = 0.008$, $\eta^2 = 0.07$) and

strike ($F_{(3, 81)} = 7.88$, $p = 1.12 \times 10^{-4}$, $\eta^2 = 0.20$) but no interaction effect between them

($F_{(6, 81)} = 1.17$, $p = 0.33$, $\eta^2 = 0.07$). A groupwise comparison with corrections for

multiple comparisons identified a significant group difference only between the ignore

and normal groups (inset of Fig. 2). Furthermore, there was no significant group effect

on the timing error of the keystrokes in either the LP_{pre} condition (F(2, 27) = 1.26, p =

0.30, η^2 = 0.02) or the LP_{post} condition (F(2, 27) = 2.90, p = 0.07, η^2 = 0.05). Together,

these findings indicate that the slow-down of the local tempo following delayed tone

production under pressure was reduced after short-term piano practice while ignoring

the delayed tone production.

315   *Figure 2 inserted here*

317   **DISCUSSION**

In the present study, we first tested whether psychological stress yields

perceptual, motor, or sensorimotor malfunctions in pianists. The results demonstrated

larger rhythmic disruption of the sequential finger movements following transient

perturbation of auditory feedback at the higher pressure condition, which provided

evidence supporting the abnormality in sensory feedback control of movements under

pressure. However, neither auditory perception of time interval nor finger motor

precision was affected by the induced psychological stress. These findings indicate that

an increase in the timing error of sequential movements under pressure is associated

with an abnormal elevation in the sensitivity of motor actions to perceived error (i.e., an

elevated feedback gain in sensorimotor control), but not with abnormalities of the

perception of the time interval or of fine motor control. In a nonstressful musical

performance, the feedback gain of sensorimotor control is usually low for pianists but

not for musically untrained individuals, as exemplified by a transient increase in

rhythmic error of movements following a transient delay in the timing of auditory

feedback during piano playing only in the latter individuals [26,29]. Therefore, these

findings suggest the de-expertise of sensorimotor control of skilled pianists under

pressure, which fits with the idea of choking, describing the failure of skill execution

under pressure [35]. In the subsequent experiment, we also tested whether sensorimotor

training normalizes the instability of sensorimotor control under pressure. The results of

the intervention experiment demonstrated that practicing the piano while ignoring

artificially delayed tone production resulted in the amelioration of the unstable

sensorimotor control under pressure. Intriguingly, this intervention effect was not

observed when playing without stress, which indicates specific training effects on

sensorimotor control during playing under stress. Furthermore, such an alleviation of

the stress effect was not observed after practicing with compensation for delayed tone

production and after practicing without the provision of delayed auditory feedback.

Taken together, our results indicated that state anxiety abnormally augmented reliance

on sensory feedback in the fine control of sequential finger movements during piano

playing, which can be restored specifically by ignoring auditory perturbation.

A question is why the pianists transiently slowed down the tempo following the

perturbed auditory feedback only in the anxiogenic situation. One may consider that the

feedback gain was elevated due to an increased uncertainty of the perception under

pressure. However, our results do not support this idea due to a lack of changes in

auditory perception of the time interval under pressure. Another possibility is that the

transient slow-down of the local tempo following the perturbation was to lower

accuracy demands on the performance by leveraging a tradeoff between the speed and

the accuracy of movements [36]. This idea is compatible with elevated coactivation

between the finger flexor and extensor muscles under pressure, which is considered a

coping strategy for accuracy demand [8,10]. A recent study also proposed a role of

muscular coactivation in feedback control of movements [37]. The shift from feedforward

to feedback control in the anxiogenic situation can affect reduced movement

individuation between the fingers under pressure [9] because the elevation of the muscular

coactivation increases the stiffness of the finger muscles connected with multiple

fingers and thereby increases biomechanical constraints on the individuated finger

movements [38]. Importantly, the reduced movement individuation between the fingers

was associated with an increase in the timing error of the piano performance under

pressure [9], which suggests that elevating feedback gain in the stressful condition played

no beneficial role in accurate musical performance. In addition, the transient slow-down

of the local tempo after delayed tone production under pressure was also unlikely to

contribute to accurate tempo control because the maintenance of the global tempo after

delayed tone production requires a transient speed-up of the local tempo to compensate

for the transient delay [29]. Taken together, the increased reliance on feedback control

under pressure is considered not as a strategy compensating for the external sensory

perturbation but as a mere response to an increase in anxiety.

Our observation indicating the increased gain of sensorimotor feedback control

of movements under pressure fits well with the explicit monitoring theory. The theory

proposes that after an appraisal of performance pressure, attention shifts exaggeratedly

toward the procedural process of motor performance, which eventually collapses skillful

performance. Previous studies reported that this attentional shift to the automatized skill

execution process is detrimental, particularly for skilled individuals who can perform

the well-learned procedural skill without explicitly attending to its underlying process

[5,11]. Although this theory explains well that skilled players are better at being aware of a

course of skill execution in the higher stressful situation [5], the present study provides

another but not exclusive viewpoint that a shift in sensorimotor control from

feedforward to feedback control mediates the deleterious effect of pressure on

performance. This idea is compatible with a recent finding of a pressure-induced

increase in the activation of the anterior cingulate cortex that is responsible for

monitoring motor performance [4].

Our intervention study clearly demonstrated that practicing the piano while

ignoring delayed tone production resulted in robust tempo control against artificially

delayed tone production in piano playing under pressure. Similarly, robust sensorimotor

control was observed when playing without explicit psychological stress. Therefore, the

abnormal elevation of the sensorimotor feedback gain under pressure can be normalized

through practicing in this way. The lack of a significant difference in the tempo error

between the group that compensated for the delayed tone production and the control

group further indicates a specific effect of practicing while ignoring the delayed tone

production on the performance fluctuation under pressure. This implicates the potential

of the practice of ignoring the auditory perturbation as a training method for stabilizing

tempo control in piano performance under pressure. Previous studies demonstrated that

skilled singers are better at ignoring the auditory perturbation in singing than

nonmusicians [24], similar to pianists [26]. The present sensorimotor training may enable

pianists to maintain low feedback gain in an uncertain and stressful situation.

There are several possible limitations of the present study. First, because a

repetition of playing under pressure potentially causes habituation to psychological

pressure, we did not additionally evaluate piano performance without the provision of

delayed tone production in the stressful condition. However, this limits the

understanding of a direct relationship of the loss of robustness in sensorimotor control

with temporal inaccuracy of piano performance in stressful situations, which needs to be

further explored. Second, although we evaluated the time interval perception at rest, we did not evaluate time perception while playing the piano (Pfordresher 2014), which might be affected by psychological stress, unlike the time perception at rest. Finally, we did not examine changes in finger muscular co-contraction under pressure through ignoring practice, which should be assessed in a future study.

REFERENCES

Beilock, S. L. & Carr, T. H. On the fragility of skilled performance: what governs choking under pressure? *J Exp Psychol Gen* **130**, 701-725 (2001).

Causer, J., Holmes, P. S., Smith, N. C. & Williams, A. M. Anxiety, movement kinematics, and visual attention in elite-level performers. *Emotion* **11**, 595-602, doi:10.1037/a0023225 (2011).

DeCaro, M. S., Thomas, R. D., Albert, N. B. & Beilock, S. L. Choking under pressure: multiple routes to skill failure. *J Exp Psychol Gen* **140**, 390-406, doi:10.1037/a0023466 (2011).

Ganesh, G., Minamoto, T. & Haruno, M. Activity in the dorsal ACC causes deterioration of sequential motor performance due to anxiety. *Nature communications* **10**, 4287, doi:10.1038/s41467-019-12205-6 (2019).

Gray, R. Attending to the execution of a complex sensorimotor skill: expertise differences, choking, and slumps. *Journal of experimental psychology. Applied* **10**, 42-54, doi:10.1037/1076-898X.10.1.42 (2004).

Higuchi, T., Imanaka, K. & Hatayama, T. Freezing degrees of freedom under stress: kinematic evidence of constrained movement strategies. *Hum Mov Sci* **21**, 831-846 (2002).

Bauerly, K. R., Jones, R. M. & Miller, C. Effects of Social Stress on Autonomic, Behavioral, and Acoustic Parameters in Adults Who Stutter. *J Speech Lang Hear Res* **62**, 2185-2202, doi:10.1044/2019_JSLHR-S-18-0241 (2019).

Ioannou, C. I., Furuya, S. & Altenmuller, E. The impact of stress on motor performance in skilled musicians suffering from focal dystonia: Physiological and psychological characteristics. *Neuropsychologia* **85**, 226-236, doi:10.1016/j.neuropsychologia.2016.03.029 (2016).

Kotani, S. & Furuya, S. State anxiety disorganizes finger movements during musical performance. *J Neurophysiol* **120**, 439-451, doi:10.1152/jn.00813.2017

(2018).

10   Yoshie, M., Kudo, K., Murakoshi, T. & Ohtsuki, T. Music performance anxiety
in skilled pianists: effects of social-evaluative performance situation on
subjective, autonomic, and electromyographic reactions. *Exp Brain Res* **199**,
117-126, doi:10.1007/s00221-009-1979-y (2009).

11   Beilock, S. L. & Gray, R. From attentional control to attentional spillover: a
skill-level investigation of attention, movement, and performance outcomes.
Hum Mov Sci **31**, 1473-1499, doi:10.1016/j.humov.2012.02.014 (2012).

12   Beilock, S. L., Wierenga, S. A. & Carr, T. H. Expertise, attention, and memory in
sensorimotor skill execution: impact of novel task constraints on dual-task
performance and episodic memory. *The Quarterly journal of experimental
psychology. A, Human experimental psychology* **55**, 1211-1240,
doi:10.1080/02724980244000170 (2002).

13   Lee, T. G., Acuna, D. E., Kording, K. P. & Grafton, S. T. Limiting motor skill
knowledge via incidental training protects against choking under pressure.
Psychonomic bulletin & review **26**, 279-290, doi:10.3758/s13423-018-1486-x
(2019).

14   Lee, T. G. & Grafton, S. T. Out of control: diminished prefrontal activity
coincides with impaired motor performance due to choking under pressure.
Neuroimage **105**, 145-155, doi:10.1016/j.neuroimage.2014.10.058 (2015).

15   Fayolle, S., Gil, S. & Droit-Volet, S. Fear and time: Fear speeds up the internal
clock. *Behav Processes* **120**, 135-140, doi:10.1016/j.beproc.2015.09.014 (2015).

16   Lake, J. I., LaBar, K. S. & Meck, W. H. Emotional modulation of interval timing
and time perception. *Neurosci Biobehav Rev* **64**, 403-420,
doi:10.1016/j.neubiorev.2016.03.003 (2016).

17   Sarigiannidis, I., Grillon, C., Ernst, M., Roiser, J. P. & Robinson, O. J. Anxiety
makes time pass quicker while fear has no effect. *Cognition* **197**, 104116,
doi:10.1016/j.cognition.2019.104116 (2020).

18   Christou, E. A., Jakobi, J. M., Critchlow, A., Fleshner, M. & Enoka, R. M. The
1- to 2-Hz oscillations in muscle force are exacerbated by stress, especially in
older adults. *Journal of applied physiology* **97**, 225-235,
doi:10.1152/japplphysiol.00066.2004 (2004).

19   Noteboom, J. T., Barnholt, K. R. & Enoka, R. M. Activation of the arousal
response and impairment of performance increase with anxiety and stressor
intensity. *Journal of applied physiology* **91**, 2093-2101,
doi:10.1152/jappl.2001.91.5.2093 (2001).

20 Kawato, M. Internal models for motor control and trajectory planning. *Current opinion in neurobiology* **9**, 718-727 (1999).

21 Maeda, R. S., Gribble, P. L. & Pruszynski, J. A. Learning New Feedforward Motor Commands Based on Feedback Responses. *Curr Biol*, doi:10.1016/j.cub.2020.03.005 (2020).

22 Osu, R. *et al.* Short- and long-term changes in joint co-contraction associated with motor learning as revealed from surface EMG. *J Neurophysiol* **88**, 991-1004, doi:10.1152/jn.2002.88.2.991 (2002).

23 Pfordresher, P. Q. Musical training and the role of auditory feedback during performance. *Ann N Y Acad Sci* **1252**, 171-178, doi:10.1111/j.1749-6632.2011.06408.x (2012).

24 Zarate, J. M. & Zatorre, R. J. Experience-dependent neural substrates involved in vocal pitch regulation during singing. *Neuroimage* **40**, 1871-1887, doi:10.1016/j.neuroimage.2008.01.026 (2008).

25 Kleber, B., Zeitouni, A. G., Friberg, A. & Zatorre, R. J. Experience-dependent modulation of feedback integration during singing: role of the right anterior insula. *J Neurosci* **33**, 6070-6080, doi:10.1523/JNEUROSCI.4418-12.2013 (2013).

26 van der Steen, M. C., Molendijk, E. B., Altenmüller, E. & Furuya, S. Expert pianists do not listen: the expertise-dependent influence of temporal perturbation on the production of sequential movements. *Neuroscience* **269**, 290-298, doi:10.1016/j.neuroscience.2014.03.058 (2014).

27 Fernholz, I. *et al.* Performance anxiety in professional musicians: a systematic review on prevalence, risk factors and clinical treatment effects. *Psychological medicine* **49**, 2287-2306, doi:10.1017/S0033291719001910 (2019).

28 Spahn, C. Treatment and prevention of music performance anxiety. *Prog Brain Res* **217**, 129-140, doi:10.1016/bs.pbr.2014.11.024 (2015).

29 Furuya, S. & Soechting, J. F. Role of auditory feedback in the control of successive keystrokes during piano playing. *Exp Brain Res* **204**, 223-237, doi:10.1007/s00221-010-2307-2 (2010).

30 Hosoda, M. & Furuya, S. Shared somatosensory and motor functions in musicians. *Scientific reports* **6**, 37632, doi:10.1038/srep37632 (2016).

31 Pfordresher, P. Q. Auditory feedback in music performance: Evidence for a dissociation of sequencing and timing. *Journal of experimental psychology. Human perception and performance* **29**, 949-964, doi:10.1037/0096-1523.29.5.949 (2003).

32   Behroozmand, R., Korzyukov, O., Sattler, L. & Larson, C. R. Opposing and
512   following vocal responses to pitch-shifted auditory feedback: evidence for
513   different mechanisms of voice pitch control. *J Acoust Soc Am* **132**, 2468-2477,
514   doi:10.1121/1.4746984 (2012).

515   33   MacDonald, E. N., Goldberg, R. & Munhall, K. G. Compensations in response
516   to real-time formant perturbations of different magnitudes. *J Acoust Soc Am* **127**,
517   1059-1068, doi:10.1121/1.3278606 (2010).

518   34   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical
519   and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol*
520   **57**, 289-300 (1995).

521   35   Baumeister, R. F. Choking under pressure: self-consciousness and paradoxical
522   effects of incentives on skillful performance. *Journal of personality and social*
523   *psychology* **46**, 610-620, doi:10.1037//0022-3514.46.3.610 (1984).

524   36   Fitts, P. M. The information capacity of the human motor system in controlling
525   the amplitude of movement. *Journal of experimental psychology* **47**, 381-391
526   (1954).

527   37   Saliba, C. M., Rainbow, M. J., Selbie, W. S., Deluzio, K. J. & Scott, S. H.
528   Co-contraction uses dual control of agonist-antagonist muscles to improve motor
529   performance. *bioRxiv* (2020).

530   38   Kimoto, Y., Oku, T. & Furuya, S. Neuromuscular and biomechanical functions
531   subserving finger dexterity in musicians. *Scientific reports* **9**, 12224,
532   doi:10.1038/s41598-019-48718-9 (2019).

Figure 1. Group means of the average heart rate during the perception, motor, and sensorimotor control tests (A); the perceptual threshold of time duration between two tones (B); the number of taps per second during the fastest tapping by each of five digits (C); the number of strikes per second by the fingers while playing the short melody excerpt as fast as possible without auditory feedback being provided (D); the timing error of the keypresses over 4 strikes following the provision of transiently delayed tone production (E); and their mean across four strikes (F) in the low-pressure (LP) and high-pressure (HP) conditions. *: $p < 0.05$. The blue and red boxplots indicate the LP and HP conditions, respectively.

Figure 2. Group means of differences in the timing error of the 4 successive keypresses following the provision of transiently delayed tone production between the high-pressure (HP) condition and low-pressure condition before running the intervention (LP_{pre}) for the normal (red), ignore (green), and compensate (blue) groups. x-axis: four successive strikes immediately after the transiently delayed tone production. The inset indicates the group means of the timing error across the 4 strikes for each of the normal, ignore, and compensate groups, which was depicted based on a significant group effect but not an interaction effect between group and strike (an error bar indicates one standard error). *: $p < 0.05$. The value 0 indicates no difference in the keypress timing error between the HP and LP_{pre} conditions.