Antimycobacterial potential of Mycobacteriophage under disease-mimicking conditions

Yeswanth Chakravarthy Kalapala[1], Pallavi Raj Sharma[1], Rachit Agarwal[1*]

[1]Centre for BioSystems Science and Engineering, Indian Institute of Science, Bangalore, India.

*Correspondence: rachit@iisc.ac.in

Running Title: Antimycobacterial potential of phage

Abstract

Antibiotic resistance continues to be a major global health risk with an increase in multi-drug resistant infections seen across nearly all bacterial diseases. Mycobacterial infections such as Tuberculosis (TB) and Non-Tuberculosis infections have seen a significant increase in the incidence of multi-drug resistant and extensively drug-resistant infections. With this increase in drug-resistant Mycobacteria, mycobacteriophage therapy offers a promising alternative. However, a comprehensive study on the infection dynamics of mycobacteriophage against their host bacteria and the evolution of bacteriophage (phage) resistance in the bacteria remains elusive. We aim to study the infection dynamics of a phage cocktail against Mycobacteria under various pathophysiological conditions such as low pH, low growth rate and hypoxia. We show that mycobacteriophages are effective against *M. smegmatis* under various conditions and the phage cocktail prevents emergence of resistance for long durations. Although the phages are able to amplify after infection, the initial multiplicity of infection plays an important role in reducing the bacterial growth and prolonging efficacy. Mycobacteriophages are effective against antibiotic-resistant strains of Mycobacterium and show synergy with antibiotics such as rifampicin and isoniazid. Finally, we also show that mycobacteriophages

are efficient against *M. tuberculosis* both under lag and log phase for several weeks. These findings have important implications for developing phage therapy for Mycobacterium.

1   Introduction

Prolonged and unsupervised use of antibiotics results in an increase of antibiotic resistance across all bacterial species which is considered a major threat to human health (Keshavjee and Farmer, 2012; Sprenger and Fukuda, 2016). The Mycobacterium genus with over 190 recognized species contains various pathogenic species including *M. tuberculosis*, *M. leprae*, *M. abscessus*, *M. avium* complex, etc. (Rastogi et al., 2001). These infections are often difficult to treat owing to their resilient cell walls which render them resistant to prolonged exposure to acidic and alkaline environments (Rastogi et al., 2001). In addition, several mycobacterial species such as *M. leprae, M. abscessus* and *M. tuberculosis* rapidly evolve resistance to antibiotics (Iseman, 1994; Cambau et al., 2018; Luthra et al., 2018; Geneva: World Health Organisation, 2019). Similarly, the *Mycobacterium avium* Complex (MAC) infections are noted to be on the rise, especially in AIDS and pulmonary disease patients (Horsburgh, 1997) and MAC infections are often associated with an increase in mortality (Horsburgh, 1997; Diel et al., 2018). The antibiotic treatment regimen is associated with the distinct and long-lasting changes to the gut flora, often resulting in the depletion of various species of commensal bacteria (Wipperman et al., 2017). Antibiotics are also associated with various adverse side-effects ranging from nausea, vomiting and fever to anaphylaxis, hypersensitivity and photodermatitis (Castells et al., 2019). In order to tackle this growing rate of antibiotic resistance and the side effects of antibiotic treatment regimens, various therapeutic strategies have been proposed, including phage therapy, that has shown tremendous potential against various bacterial pathogens (Kortright et al., 2019).

Phages are viruses that are host specific towards bacteria and are inherently non-pathogenic to humans. They are abundantly present in the natural microflora of the human body (De Paepe et al., 2014; Manrique et al., 2016). A recent non-comparative clinical trial showed that the intravenous delivery of anti-staphylococcal phages in patients with severe infection did not pose any challenges to tolerability and safety (Petrovic Fabijan et al., 2020). Moreover, phages remain infective and lyse antibiotic-resistant bacteria (Kortright et al., 2019). To circumvent the narrow-host range specificity and development of resistance (Labrie et al., 2010), multiple phages are used in phage therapy. Successful compassionate use of phages against antibiotic-resistant bacteria on patients in the USA and Europe has been on the rise and shows their safety and translation potential (Jennes et al., 2017; Schooley et al., 2017).

With over 11,000 mycobacteriophages isolated and more than 1800 sequenced (Russell and Hatfull, 2017), there is a large library of phages against Mycobacterium. Although the desired lytic phenotype of phages (Pimentel, 2014; Catalão and Pimentel, 2018) for phage therapy brings the number down, development of Bacteriophage Recombineering of Electroporated DNA (BRED) technique (Marinelli et al., 2008) allows rapid conversion of lysogenic phages to lytic phages for phage therapy. Recently, a three-phage cocktail has been successfully used in a clinic for the treatment of a cystic fibrosis patient suffering from *Mycobacterium abscessus* infection (Dedrick et al., 2019). Prophylactic phage (D29) therapy against TB also showed some success in a preclinical animal model in reducing the burden of infection (Carrigy et al., 2019). A study employing phage therapy in treating subcutaneously injected *M. tuberculosis* (H37Rv) infection in guinea pigs showed that mycobacteriophages have a therapeutic effect (Sula et al., 1981).

Phage therapy is a strong candidate for treating mycobacterial infections, however it presents

a few physiological hurdles that need to be taken into consideration. Various Mycobacterium

species (*M. tuberculosis*, *M. avium* complex, *M. bovis* and *M. leprae*) are found to reside

intracellularly at low pH conditions which poses a challenge in efficacy of anti-bacterials

(Sibling et al., 1987; Oh and Straubinger, 1996; Ehlers and Schaible, 2012; Benítez-Guzmán

et al., 2019). *M. tuberculosis* and *M. avium* complex also form granulomatous structures,

marked by a gradient of hypoxia, nutrient deprivation and acidity (Timm et al., 2003; Voskuil

et al., 2004; Lawn and Zumla, 2011; Ehlers and Schaible, 2012; Seto et al., 2020). For a phage

to successfully eliminate these bacilli, it needs to be active and efficient under such low pH

and hypoxic conditions and should be able to enter into the eukaryotic cells. Since most bacteria

enter a non-replicative stationary phase, the phage should also be able to infect and kill bacteria

in the stationary phase where most of the current antibiotics fail. Designing a phage therapy

that would be effective in such complex physiological environments would necessitate a better

understanding of the bacteria-phage dynamics under various conditions. However, to our

knowledge, no study has looked at the efficacy of phages over time in such different

environments. Hence, there is a need to study the efficacy of phage cocktails in the

physiological and pathophysiological conditions in which bacteria is known to survive.

In this study, we have performed phage-bacteria kinetics under various pathophysiological

conditions of mycobacterial diseases- low pH, hypoxia and stationary phase. We chose *M.*

smegmatis as our model organism in designing the experiments owing to its relatively fast

replication time and therefore, its ability to offer rapid screening and understanding of phage

kinetics in extreme host environments. We study the emergence of phage resistance in *M.*

smegmatis and the combinatory effect of antibiotics and phage therapy. We show that

mycobacteriophage cocktails are effective against Mycobacterium under the low pH, hypoxic

and stationary conditions and showed synergy with antibiotics. Finally, we also find that

mycobacteriophages are able to prevent growth of *Mycobacterium tuberculosis* which

encourages further exploration of phages for diseases caused by Mycobacterium.

104 2 Materials and Methods

105 2.1 Chemicals and Reagents

106 Middlebrook 7H9 Broth, Middlebrook 7H10 Agar and DMSO were purchased from Sigma (St

107 Louis, MO, USA). Middlebrook 7H11 agar was purchased from BD Difco (NJ, USA). ADC

108 growth supplement, Ziehl-Neelsen stain, rifampicin, and isoniazid were purchased from

109 HiMedia (India). Calcium Chloride, Magnesium Sulphate, Sodium Chloride, Tween 80 and

110 Glycerol were purchased from Fisher Scientific (MA, USA).

112 2.2 Bacterial cell culture and maintenance

Primary *Mycobacterium smegmatis* (mc^{2}155) or *Mycobacterium tuberculosis* (H37Ra)

cultures were grown in Middlebrook 7H9 broth supplemented with Glycerol, ADC and Tween-

80 (0.1% v/v). A log phase primary culture was inoculated into a secondary culture without

Tween-80 and supplemented with $CaCl_2$ (2 mM) to promote an efficient infection of phages.

Frozen stocks were prepared by adding 300 μL of a late log phase culture to 35% (v/v) final

concentration of glycerol and stored at -80°C until future use. All cultures were maintained at

37°C with rotary shaking at 180 rpm. For optical density (OD) measurements, 300 μL of

bacterial culture at various time intervals was diluted 10 times in media and pipetted several

times to obtain a uniform cell suspension. Readings were taken using a spectrophotometer

(Jenway 7205 UV/Visible Spectrophotometer) at 600 nm against a media blank.

124 2.3 Soft agar overlay and spot assays for phage enumeration

Soft agar overlay technique was used to detect and quantify phages. Briefly, soft agar was

prepared by adding agar (0.8%) to 7H9 media along with the required supplements. The

suspension was autoclaved and allowed to cool down to 42°C. 1 mL of late log phase bacterial

culture (OD>1) and/or 100 µL of phage dilutions was added to 5 mL of the soft agar and was

poured onto a 7H10 plate supplemented with $CaCl_2$. The soft agar was allowed to cool down

and solidify. The plate was incubated at 37°C overnight.

For verification of phage titers, spot assays were carried out in which late-log phase *M.*

smegmatis cultures (OD>1) were plated on a Middlebrook 7H10 media plate supplemented

with $CaCl_2$ (2 mM) and glycerol to form a bacterial lawn. Phage samples were serially diluted

and spotted on to the bacterial lawn. The plates were incubated at 37°C for 24 h and the number

of plaques formed at various dilutions was counted.

2.4   Phage amplification and maintenance

Five phages (D29, TM4, Che7, PDRPv (Bajpai et al., 2018) and PDRPxv (Sinha et al., 2020))

used in this study were amplified on *M. smegmatis*. Similarly, three phages (D29, TM4 and

DS6A) were amplified on *M. tuberculosis*. For liquid amplification, phage sample at 10 MOI

was added to a log phase culture (with OD between 1-2) and incubated for 12-24 h at 37°C

with rotary shaking at 180 rpm. The cultures were then centrifuged at 4000 g for 10 min and

the supernatant was collected and filtered through a 0.22 µm syringe filter (Biofil, FPV203030-

A). For plate amplification, a soft agar overlay containing log phase *M. smegmatis* (OD 1-2)

was prepared and the phage sample (10 MOI) was spotted on top of the soft agar. The plates

were incubated for 12-24 h at 37°C following which, the soft agar was collected into a 50 mL

falcon tube. An appropriate amount of Magnesium Sulphate- Tris Chloride- Sodium Chloride

Buffer (SM Buffer) was added to the falcon tube to completely immerse the soft agar and the

falcon tube was incubated for 2-4 h at 37°C or overnight at 4°C. The falcon tube was then

subjected to centrifuge at 4000 g for 10 min at 4°C and the supernatant was collected and

filtered using 0.22 μm syringe filters. The soft agar overlay was used to determine the phage

titers.

2.5　Bacterial growth kinetics

In order to study the bacterial growth kinetics, 7H9 broth supplemented with ADC and $CaCl_2$

was prepared and inoculated with 50 μL of log-phase secondary bacterial culture (OD 1-2) per

10 mL of the media. The cultures were then incubated at 37°C with rotary shaking at 180 rpm.

Periodic measurements of bacterial optical density (OD) were taken. For phage treated

samples, a 10 MOI (for *M. smegmatis*) or 1 for (*M. tuberculosis*) of each phage in the respective

phage solution was added at the appropriate time (either at the start of the experiment (lag

phase) or at a specified interval of mid-log phase).

2.6　Bacterial growth kinetics with phage – pH, stationary phase and hypoxia

To study the effects of acidic pH on phage infection dynamics, phage kinetics assay was

performed at varying the pH of the media. Briefly, the pH of the prepared media was reduced

from 6.8 prior to the experiments using HCl to either a pH of 6.0 or pH of 5.5. The media was

filtered and used for future growth kinetics experiments. Bacteria were inoculated into the

media at pH 6.8, 6.0 and 5.5. For phage treated samples, a 10 MOI for each phage in the 5-

phage cocktail was added at an appropriate time (either at the start of the experiment (lag phase)

or at a specified interval of mid-log phase. OD was measured at 600 nm.

For stationary phase experiments, *M. smegmatis* cultures were grown in 7H9 media

supplemented with Glycerol, ADC and $CaCl_2$ (2 mM) until an OD of over 10 was reached. The

stationary phase cultures were then either treated with the 5-phage cocktail at 10 MOI or left

untreated. OD values were obtained at regular intervals at 600 nm.

For the studies involving hypoxia, the cultures were grown in plastic K2-EDTA Vacutainers

(BD Cat No. 367525) to generate hypoxia. Briefly, K2-EDTA Vacutainers were obtained from

BD and washed with autoclaved sterile water over a period of 24 h to remove the EDTA

coating. The vacutainers were then washed with prepared media before experimentation. Upon

inoculation, the vacutainers were sealed with the rubber septa caps and parafilm to prevent

oxygen diffusion. Samples were collected through a syringe and 20G needle for

spectrophotometer readings. Oxygen levels were monitored by adding Methylene Blue

(0.01%) to the samples (Boshoff et al., 2004) and measuring the OD at 665 nm.

2.7   **Ziehl-Neelsen staining**

Ziehl-Neelsen staining was performed periodically to check for contamination and for the

confirmation of *Mycobacterium spp*. as described by the manufacturer. Briefly, the bacterial

suspension was spread evenly over the slide and air-dried for 30 min. The slide was then heat-

fixed on a hot plate for 15 min at 60°C and then flooded with Carbol Fuchsin stain at 85°C.

Care was taken not to overheat or boil the stain and the slide was frequently replenished with

additional Carbol Fuchsin stain. The slide was allowed to remain on the hot plate for 10 min

following which it was removed and allowed to incubate for an additional 5 min. The stain was

washed away under running tap water and flooded with decolorizer for 2 min or until the smear

was sufficiently decolorized. The slide was washed under running tap water and treated with

Methylene Blue for 30 s. The slide was subjected to one final washing under running tap water

to remove the Methylene Blue stain and was air-dried following which, it was observed under

a 100× oil immersion objective using an upright fluorescence microscope (Olympus BX53F)

for the presence of either pink rod-shaped acid-fast bacteria or blue stained contamination.

2.8   Antibiotic Minimum Inhibitory Concentration (MIC) determination

The MICs of both rifampicin and isoniazid were determined to establish the antimycobacterial

activities of the respective antibiotics against *M. smegmatis* strain mc^2 155. Antibiotic stock

solutions were prepared at concentration of 10 mg/mL in sterile deionized H_2O for isoniazid

and in DMSO for rifampicin. Stocks were stored at -20°C until further usage. Several dilutions

were prepared in a 96- well plate for each of the antibiotic stock solutions to obtain various

concentrations ranging from 1.5 μg/mL to 100 μg/mL. A total of 2×10^5 cells of *M. smegmatis*

were added to each of the wells. The 96-well plate was placed in a rotary shaker incubator at

37° C for 48 h and was observed for growth against control be measuring OD at 600 nm. The

lowest concentration of antibiotic without bacterial growth was termed as MIC for the

respective antibiotic.

2.9   Antibiotic growth kinetics

In order to study the effect of antibiotics- isoniazid and rifampicin on *M. smegmatis* growth,

7H9 broth supplemented with ADC and $CaCl_2$ was prepared and inoculated with 50 μL of log-

phase secondary bacterial culture per 10 mL of the media. The cultures were then incubated at

37°C with rotary shaking at 180 rpm. Periodic measurements of bacterial OD at 600 nm against

a media blank. For antibiotic-treated samples, a final concentration of rifampicin at 40 μg/mL

and 100 μg/mL and isoniazid at 40 μg/mL and 100 μg/mL were added at the appropriate time

(either at lag phase or at a specified interval of mid-log phase). In order to study the potential

synergistic relationship between antibiotics and phages, we co-treated bacterial cultures with

rifampicin at 20 μg/mL and phage cocktail (at MOIs 0.01, 0.05 and 10) at either the lag phase

or at a specified interval of mid-log phase. Periodic OD measurements were taken at 600 nm.

2.10   Phage kinetics against isoniazid resistant *M. smegmatis*

To study the effect of phage cocktail on antibiotic-resistant strain, 7H9 broth supplemented

with ADC, $CaCl_2$ and isoniazid (40 μg/mL) was prepared and inoculated with 50 μL of

isoniazid-resistant (mc^2 4XR1) (Mohan et al., 2015; Padiadpu et al., 2015, 2016) log phase

secondary bacterial culture per 10 mL of media. The cultures were then incubated at 37°C with

rotary shaking at 180 rpm. Periodic measurements of bacterial OD were taken using a

spectrophotometer at 600 nm against a media blank. For phage treated samples, the 5-phage

cocktail was added at 10 MOI at an appropriate time (either at lag phase or at a specified interval

of mid-log phase).

2.11   Checkerboard assay

To study the possible synergy between bacteriophages and rifampicin, checkerboard assays

were performed. Briefly, *M. smegmatis* bacterial cultures were seeded into 48-well plates at a

seeding density of 10^5 colony forming units/mL. Respective concentrations of antibiotic and

phages were added to the wells and the plates were incubated at 37°C with rotary shaking at

180 rpm for 48 h. OD readings were taken at 600 nm using a plate reader (Tecan Spark

Multimode Microplate Reader).

2.12   Phage stability assay

To study the stability of the 5-phage cocktail at various pH, we incubated a 10^8 solution at pH

6.8, 6.0 and 5.5 at 37°C. Samples were collected and titrated against *M. smegmatis* every 24

hours.

2.13   Statistical analysis

All experiments were performed on independent biological replicates. Statistical significance was determined for control and experimental groups using multiple t-test with Holm-Sidak method, with alpha = 0.05. For analysis of multiple groups, 2-way ANOVA was used with Sidak's multiple comparisons test, with alpha = 0.05. GraphPad (Prism) was used for all statistical analysis. Data points were excluded if contamination was identified using acid-fast staining.

3   Results

3.1   Mycobacteriophages reduce bacterial growth in *M. smegmatis*

Several lytic Mycobacteriophages were procured and amplified to titers greater than 10^9 pfu/mL. In order to test their efficacy, *M. smegmatis* bacterial cultures were infected with phages at the MOI of 10 either in the lag phase or in the mid-log phase. The optical density of the cultures at 600 nm was monitored using a spectrophotometer at various time intervals. Cultures treated with phages D29, TM4, Che7, PDRPv and PDRPxv individually showed a significant reduction in the optical density after 2-5 h when phages were added at log phase (time of addition shown by red arrows) (**Figure 1**). Similarly, when the phage infection was carried out at lag phase, no growth was observed for several hours. It is well known that bacteria evolve resistance to phages over time. *M. smegmatis* cultures infected with phages were also monitored for phage resistance. Phage resistance was observed in all phage treated *M. smegmatis* cultures. While D29 and Che7 treated cultures showed phage resistance at around 40 h, TM4 treated cultures showed phage resistance at 60 h; PDRPv and PDRPxv treated cultures showed phage resistance at 83 h and 101 h respectively (**Figure 1**). Bacterial cultures infected with phages at an MOI of 30 showed no significant increase in the onset of phage

resistance compared to MOI of 10 (**Supplementary Figure S1**). To test if the bacteria growing

after the phage challenge had any altered susceptibility to the phages, we titrated our individual

phage stocks on bacterial growths obtained after treatment with D29 and PDRPv. While D29

and PDRPv showed much reduced titers (2-3 log orders) against their respective regrowing

bacteria, no significant reduction in susceptibility to other phages was observed

(**Supplementary Figure S2**).

279
3.2   Phage cocktails show synergy in growth reduction

To test the effect of phage cocktails, *M. smegmatis* cultures infected with multiple phages (a

cocktail of 3 or a cocktail of 5) at 10 MOI of each phage were monitored for OD at 600 nm

(**Figure 2**). Phage cocktail treatment significantly reduced the optical density of the bacterial

cultures compared to controls 5 h post treatment. Phage resistance in cocktail treated cultures

prolonged to 105 h in case of 3 phage cocktail (D29, TM4 and Che7) (**Figure 2A**) and to 322

h in case of 5 phage cocktail (D29, TM4, Che7, PDRPv and PDRPxv) (**Figure 2B**). The

corresponding Colony Forming Units (CFU)/mL readings were obtained and the viable cell

count was observed to be correlating with OD readings (**Supplementary Figure S3A**). This

shows that the cocktail of phages can act synergistically to delay the evolution of resistance.

Periodic phage titer measurements showed increase in the phage titers as the bacterial OD

dropped and phages could be detected upto 400 h although a steady decline in the titer was

observed (**Supplementary Figure S7A**). Subsequent phage stock rechallenge on the growing

bacteria in the 5-phage cocktail kinetics showed that all phage stock titers had dropped by

>99% (2-3 log order) on the pre-exposed bacteria compared to untreated bacteria

(Supplementary Figure S3B) suggesting evolution of the bacteria to resist phage mediated

lysis. It is important to note that the phages used here belong to different genomic clusters:

cluster A (D29, subcluster: A2), cluster K (TM4, subcluster: K2) and cluster B (PDRPv and

PDRPxv, both in subcluster B1) (2–6). Cluster information for Che7 is not known. Hence, it is

expected that these phages will employ different mechanisms to infect the bacteria which is

consistent with delay in evolution of resistance (**Figure 2**) as more phages are added to the

cocktail. The 5-phage cocktail was selected and used for future experiments. To determine the

time it takes for OD of bacterial cultures to start reducing after phage addition, we took OD

readings every 15-20 mins. We found that OD of the bacterial cultures starts to drop only after

120-150 mins **(Supplementary Figure S4)** which is consistent with the previously reported

latency period of D29 (~60 min) (Samaddar et al., 2015; Bavda and Jain, 2020) and PDRPxv

(~135 min) in *M smegmatis* (Sinha et al., 2020).

To test for the efficacy of mycobacteriophages at various Multiplicity of Infection (MOI), *M.*

smegmatis cultures were infected with a MOI of 10, 1, 0.1 and 0.01 of each phage: bacteria in

either the lag phase or the mid-log phase. A 10 MOI **(Figure 2B)** significantly reduced the

growth of bacterial culture 3 h post treatment with phage resistance emerging at 322 h. A 1

MOI **(Figure 2C)** also resulted in a significant reduction of bacterial growth 3 h post treatment,

however, phage resistance emerged at around 222 h. While a 0.1 multiplicity of infection

(Figure 2D) resulted in a reduction in growth, resistance was observed at 102 h for mid-log

phase treated cultures. A 0.01 MOI did not result in any significant reduction in the bacterial

growth **(Figure 2E)**. For all future experiments, we have used the MOI of 10 to evaluate the

maximum lysing potential of phages against Mycobacterium.

3.3   Mycobacteriophages are effective in under low pH and hypoxia

Several species of Mycobacterium exist as intracellular pathogens in acidic and hypoxic

environments (Sibling et al., 1987; Oh and Straubinger, 1996; Rastogi et al., 2001; Houben et

al., 2006; Benítez-Guzmán et al., 2019; Seto et al., 2020). In order for phage therapy to be

efficient against such mycobacterial infections, phages need to retain their activity and

infectivity in these extreme environments. We, therefore, evaluated the efficiency of our 5-

phage cocktail under low pH conditions normally observed in infected phagosomes (ranges

from pH 4-6.5) of mammalian cells. *M. smegmatis* cultures did not grow at pH 5.0 and below.

Hence, we chose pH 5.5 and 6.0 to test the effectiveness of phages as these pH ranges are

known to exist in early and late endosomes (Rohde et al., 2007). We tested the phage stability

in 7H9 medium with pH 6.8, 6.0 and 5.5. Although there was a progressive loss of phage titers

over 300 h, no significant difference in phage activity was observed at lower pH compared to

pH of 6.8 (**Supplementary Figure S5**). Under low pH conditions of 6.0 and 5.5, the

mycobacteriophages remain efficient in infecting and killing *M. smegmatis* and showed

resistance around 300 h, similar to those observed at pH 6.8 (standard pH of 7H9 medium)

(**Figure 3A-B, Supplementary Figure S6A-B**).

Efficacy of the phages under hypoxic conditions was checked by culturing bacteria in sealed

vacutainers with septa. Since, there is no exchange of gases in this model, as the bacteria grows

over time, the oxygen present in the tube depletes creating a hypoxic environment. This was

confirmed by adding methylene blue to the media that turns colorless in the absence of oxygen

(**Supplementary Figure S8**) (Sumitani et al., 2004). We infected cultures with the 5-phage

cocktail, at either low hypoxia (intermediate methylene blue absorbance) or high hypoxia (at

lowest methylene blue absorbance). We observed efficient lysis of bacteria by the 5-phage

cocktail in both the hypoxic conditions 20-25 h post treatment. (**Figure 3D** and

Supplementary Figure S6D). Similar to pH experiments, the emergence of phage resistance

was observed in both low and high hypoxia phage treated samples in ~300 h.

3.4   Mycobacteriophages are effective in stationary phase of bacterial growth

Within the host environment, Mycobacterial species slow their growth rate and often enter a

non-replicative state (Voskuil et al., 2004; Houben et al., 2006; Ehlers and Schaible, 2012).

This reduced growth state poses a significant hurdle in the treatment of infections with

conventional antibiotics (Gutierrez et al., 2017). For phage therapy to be efficient against

mycobacterial infections, phages need to retain the infectivity against the slow growing

bacteria. To test the effect of phage cocktail against *M. smegmatis* in stationary phase, *M.*

smegmatis cultures were allowed to reach an OD>10. The stationary phase cultures were then

infected with the 5-phage cocktail without any supplementation with nutrients and periodic OD

measurements were taken at 600 nm. Significant reduction in OD and CFU were observed in

phage treated samples 24 h post treatment, demonstrating that the 5-phage cocktail is effective

in stationary phase conditions (**Figure 3C, Supplementary Figure S6C**). Interestingly, even

after 350 h of culture, no growth of bacteria was observed. In periodic phage titer

measurements, phages could be detected upto 400 h although a steady decline in the titer was

observed **(Supplementary Figure S7B)**.

361
3.5   Efficacy of isoniazid and rifampicin against *M. smegmatis*

Tuberculosis infections are conventionally treated by a combination of antibiotics, including

the first-line drugs, isoniazid and rifampicin (Geneva: World Health Organisation, 2019). To

compare our phage kinetics with these antibiotics, we first determined the minimal inhibitory

concentration of rifampicin (MIC: 25 µg/mL) and isoniazid (MIC: 15 µg/mL) on *M. smegmatis*

(mc^{2}155). We then treated the cultures with 40 µg/mL and 100 µg/mL of the respective

antibiotic at either lag phase or at mid-log phase. For 40 µg/mL antibiotic concentration, there

was no significant reduction of bacterial growth at log phase (**Supplementary Figure S9 A,**

B). For 100 µg/mL, we observed only a transient growth inhibition of *M. smegmatis* compared

to the phage treatment (**Supplementary Figure S9 C, D**). The cultures continue to grow even

in the presence of high concentration (several times the MIC values) of antibiotics. This growth

inhibition was much more pronounced in lag-phase cultures treated with antibiotics than log-

phase cultures treated with antibiotics. Antimicrobial susceptibility depends on multiple factors

such as inoculum size, metabolic state of the bacteria and cellular respiration rate (Lobritz et

al., 2015; Li et al., 2017; Lopatkin et al., 2019). All these factors change as bacterial cells

transition from lag phase to exponential growth and antibiotic efficacy may be variable even

at the same concentration for different growth states and bacterial numbers. Isoniazid interferes

with mycolic acid synthesis and hence is more effective on replicating Mycobacteria but due

to high numbers of bacteria in the exponential phase, its efficacy was lower than in the lag

phase. These results are also consistent with the previously observed results that isoniazid had

reduced efficacy on exponentially growing *M. smegmatis* cultures (Teng and Dick, 2003).

3.6   Testing synergy between Mycobacteriophages and antibiotics

To check whether treatment with the 5-phage cocktail enhances the efficacy of controlling

bacterial growth in antibiotic-treated cultures, we added 20 µg/mL of rifampicin (~MIC value)

along with a 5-phage cocktail solution at an MOI of 0.05, 0.01 or 10 during the lag phase or

the mid-log phase. For MOI of 0.01, we did not observe a significant reduction in growth in

only phage treated samples at both lag and log phases compared to untreated samples.

However, rifampicin and phage cotreated samples showed a significant growth reduction in

the log phase 6 h post treatment with resistance developing around 224 h **(Figure 4A)**. In the

lag-phase bacteria, phage addition at 0.01 MOI had no effect on growth and rifampicin treated

bacteria showed growth within 50 h while the lag-phase samples co-treated with both

rifampicin and phage cocktail showed no growth emergence upto 254 h. This prominent

reduction in bacterial growth in cultures co-treated with rifampicin and phage cocktail could

be attributed to antibiotic-phage synergy previously observed in various other bacterial

infections including *P. aeruginosa* infection treatment (Oechslin et al., 2017), *S. aureus*

infection treatment (Chhibber et al., 2013) and *A. baumannii* infection treatment (Jansen et al.,

2018). The effectiveness of a cotreatment at low MOI of phage cocktails and rifampicin

concentrations shows phage-antibiotic synergy. While rifampicin binds to RNA polymerase

that might interfere with phage amplification, a staggered treatment with phages followed by

rifampicin has been shown to be synergistic (Rahman et al., 2011; Tkhilaishvili et al., 2018).

Similar observations have been noted for antibiotics that target the DNA replication and

translational machinery (Torres-Barceló et al., 2014; Kim et al., 2018). It is possible that the

co-incubation of both anti-bacterial agents prevents the emergence of bacteria that is tolerant

to the individual anti-bacterials and hence show synergy. For cultures treated with a phage

cocktail MOI of 0.05, we observed a similar reduction in growth with phage resistance

emerging at 254 h in log-phase treated samples (**Supplementary Figure S10A**). Interestingly,

the phage-antibiotic synergy at log-phase became less pronounced with increase in phage MOI.

At MOI of 10, no synergy was observed in log phase cultures but a delay in emergence of

resistance in lag phase treated cultures was observed **(Supplementary Figure S10B).** To

further test the phage synergy with rifampicin, we performed experiments at different

concentration of phage-rifampicin through a checkerboard assay (**Figure 4B**) and found their

interaction to be synergistic in preventing bacterial growth (FIC index of 0.43 (<0.5) at phage

MOI 0.1 and rifampicin concentration of 10 μg/mL) (Hall et al., 1983).

3.7   Mycobacteriophages are able to lyse antibiotic-resistant *M. smegmatis*

Isoniazid-resistant infections are observed to be on the rise across the globe (Geneva: World

Health Organisation, 2019). We, therefore, looked at the efficacy of phage infection and

mycobacteriophages against isoniazid-resistant strain of *M. smegmatis* (Mohan et al., 2015;

Padiadpu et al., 2015, 2016). First, we tested whether individual phages have altered efficiency

in infecting and lysing the isoniazid-resistant strain. The phages retained their initial high titers

against isoniazid-resistant *M. smegmatis* and no significant drop in the efficiency of plating

was observed **(Supplementary Figure S11)**. Next, we observed that the 5-phage cocktail

remains effective in infecting and lysing antibiotic-resistant *M. smegmatis* (**Figure 4C**). We observed a sharp decrease in the OD of phage-treated log-phase cultures 5 h post treatment and an extended delay in the growth of lag-phase cultures. We also observed growth of phage treated bacterial cultures at 294 h, owing to the emergence of phage resistance similar to the previous experiments.

3.8   Mycobacteriophages are effective against *M. tuberculosis*

Next, we checked whether other slow-growing species of Mycobacteria also exhibit similar dynamics with Mycobacteriophages. Although several reports have shown that phages infect and kill *M. tuberculosis,* however, whether cocktail of phages can prevent growth of *M. tuberculosis* strains over long durations and how long it takes for evolution of resistance is not known. In our 5-phage cocktail, two phages (D29 and TM4) are known to infect *M. tuberculosis* strains. These two phages and another phage DS6A were amplified using *M. tuberculosis* (H37Ra) as a host because the phages expanded on *M. smegmatis* had low titer on H37Ra. For infection experiments, a phage cocktail was prepared using these three phages. Since our phage titers against *M. tuberculosis* strains were low (~10^7 PFU/mL) even after several rounds of amplification, we used a low inoculation density (~10^4 CFU/mL) of bacteria to perform infection experiments at 10 MOI for each phage. We proceeded to treat the cultures of *M. tuberculosis* (H37Ra strain) with this three-phage cocktail in both the lag phase (OD ~0.06) and in the log phase. Phage treatment was successful and efficient in preventing the growth of *M. tuberculosis* for several weeks (**Figure 5A).** Emergence of phage resistance was observed after 1368 h (57 days). We also observed reduction in bacterial CFU after phage treatment, consistent with our OD measurements **(Figure 5B**). These experiments show that phages can be also effective against slow growing Mycobacterium for long durations.

4   Discussion

Phage therapy has been successfully used in the treatment of various bacterial infections including *P. aeruginosa* urinary tract infection (Khawaldeh et al., 2011), multidrug-resistant *A. baumannii* infection (Schooley et al., 2017), *P. aeruginosa* aortic graft infection (Chan et al., 2018), and more recently, *Mycobacterium abscessus* infection of a 15- year old cystic fibrosis patient (Dedrick et al., 2019). Although phage therapy holds strong potential, there is little understanding of phage-host dynamics in disease environments. In this study, we studied the host-phage infection dynamics under various physiological and disease-mimicking conditions.

We chose mycobacterial species as our phage host, as persistent and multidrug-resistant mycobacterial infections have been increasing in incidence over the past decades and there is a need for new therapy against such infections (Iseman, 1994; Gygli et al., 2017; Aslam et al., 2018; Cambau et al., 2018; Luthra et al., 2018). For phage therapy to be successful and efficient against mycobacterial infections, the phages need to retain their infectious activity in these extreme environments and infect non-replicative stationary phase bacteria. D29 is a well-known phage against *M. tuberculosis* known to produce superoxide radicals in oxygen-rich environments, which synergizes with phage mediated lysis to accelerate the killing of Mycobacterium (Samaddar et al., 2016). However, it is observed to be ineffective in hypoxic conditions (Swift et al., 2014). TM4, on the other hand, is shown to be efficient in infecting *M. tuberculosis* during stationary phases and hypoxic conditions (Swift et al., 2014). Biochemical analysis revealed that a peptidoglycan motif in the tape measure protein of TM4 phage confers an advantage in infecting bacteria in stationary phase (Piuri and Hatfull, 2006). This complementary nature of phages suggests that a cocktail of phages would be effective in the elimination of mycobacterial infections across a broad range of environments and growth states. Although research has been carried out on individual mycobacteriophages under specific

environments, limited information exists which solidifies our understanding of phage cocktail

infection kinetics and evolution of phage resistance under different disease-mimicking

conditions.

In this study, we show that a 5-phage cocktail is efficient against *M. smegmatis* at various pH,

low oxygen availability and different growth rate of bacteria. Furthermore, for the first time,

our study shows the *M. tuberculosis* growth kinetics in presence of phage cocktail over long

durations. We found that a 3-phage cocktail of D29, TM4 and DS6A was successful in

preventing as well as reducing the growth of *M. tuberculosis* (H37Ra) for several weeks

(**Figure 5**).

We also found that two frontline TB antibiotics- isoniazid and rifampicin were limited in their

ability to control the growth of *M. smegmatis* in the exponential phase of bacterial growth

(**Supplementary Figure S9**). The more striking result from our data was the synergy of phages

with rifampicin and effectiveness of phages on isoniazid-resistant *M. smegmatis* (**Figure 4A-**

C). This finding stands in line with observations of synergy during phage-antibiotic

combination treatments for other bacterial systems (Comeau et al., 2007; Chhibber et al., 2013;

Oechslin et al., 2017; Jansen et al., 2018). Although phage therapy holds promise in treating

bacterial infections and antibiotic resistance, bacteria are known to develop resistance against

phages through various mechanisms (Labrie et al., 2010). However, unlike antibiotics which

have limited classes and require scrutinized drug design, phages are abundant in nature

(Edwards and Rohwer, 2005) and the mycobacteriophage repository alone is over 10,000 in

number (Russell and Hatfull, 2017), providing a much larger repertoire of arsenal against

bacterial infections. However, when designing a phage cocktail, the phage resistance needs to

be considered. The phage-bacterial dynamics and evolution of resistance described here are

some the first studies performed on mycobacterial species and provide useful information for

future exploration of phages for therapeutic use. We found that the bacteria eventually evolve

resistance to the phage or phage cocktail as is seen by regrowth of the bacteria after phage

challenge even though the phages can be detected in the medium at all times. Subsequent

challenge of our phage stocks on this regrowing population showed a drop of 2-3 log orders

(>99%) in titers suggesting evolution of the bacteria to resist phage mediated lysis. The

mechanisms of this resistance are not clear and need further investigation. Bacterial hosts are

known to develop resistance against phages through various mechanisms including genetic

mutations, hypermutable loci, the blockage of the injection of phage DNA, cleavage of injected

phage DNA, interference with phage assembly, abortive infection and other phage defense

systems such as the R-M system, DISARM system or the CRISPR-Cas system (Azam and

Tanji, 2019; Turkington et al., 2019). Whether *Mycobacterium spp.* utilize these mechanisms

to overcome the phage challenge is not clear and requires further research. In our studies, we

observed a strong correlation between the delay in the emergence of phage resistance and the

number of phages used in our phage cocktail.

Our phage infection at different MOI suggests that it is important to deliver a high number of

phages at the site of infection for therapy to be successful (**Figure 2C-D**). Several approaches

have been developed to deliver active phages to the deep lung such as vibrating mesh nebulizer,

soft mist inhalers and dry powder inhalers (Puapermpoonsiri et al., 2009; Vandenheuvel et al.,

2013; Liu et al., 2016; Carrigy et al., 2017; Agarwal et al., 2018). Similarly, delivery to

macrophages by direct phagocytosis of phages (Jończyk-Matysiak et al., 2015) or through

nanoparticles/microparticles can result in targeting of the intracellular niche of the bacteria

(Fenaroli et al., 2014; Nieth et al., 2015; Agarwal et al., 2018). However, the current

understanding of the diffusion of phages across the granuloma remains limited and merits

further studies in animal models.

A limitation to this study is that most experiments were performed on non-pathogenic, fast

growing *M. smegmatis* and may not translate to slow growing pathogenic species such as *M.*

tuberculosis. Although we show that phages can infect and lyse *M. tuberculosis* for long

durations, further studies are warranted in testing the efficiency of these phages on such slow

growing pathogenic species for longer durations and in acidic and hypoxic conditions.

However, efficacy of mycobacteriophages in lysing intracellular slow growing mycobacteria

in vitro as well as *in vivo* has been demonstrated by other groups. Preliminary reports on D29

incubation with H37Rv infected mouse macrophages show reduction in intracellular bacterial

burden, however penetration of D29 inside the macrophage was reported to be low (Lapenkova

et al., 2018). It is critical for the phages to directly encounter bacteria to be effective. Phage

delivery inside infected mammalian cells can be facilitated by using carriers, for instance by

adsorbing on non-pathogenic bacteria. This led to enhanced intracellular localization of phages

in vitro in *M. avium* and *M. tuberculosis* infected macrophages and reduction in bacterial load

in a mouse model of disseminated *M. avium* infection (Broxmeyer et al., 2002; Danelishvili et

al., 2006). Bactericidal efficiency of phages has also been shown *in vivo* in a *M. ulcerans*

infected mouse footpad model (Trigo et al., 2013). Another challenge for phage therapy is the

human immunological response. Although phages are known to elicit immune response (Ooi

et al., 2019), previous studies have shown that phages and host immune response can

synergistically result in the elimination of acute and chronic bacterial infections (Leung and

Weitz, 2017; Roach et al., 2017; Waters et al., 2017). This is further supported by the successful

treatment of several patients after phage administration in clinics.

Conclusion

In summary, we demonstrate the efficacy of phages under various pH ranges (6.8, 6.0 and

5.5), stationary phase as well as under hypoxic conditions. We show that phages are effective

in infecting and lysing antibiotic resistant bacteria and show synergy with antibiotics. We

further show that a three-phage cocktail is effective in preventing the growth of slow growing

M. tuberculosis for several weeks.

Author Contribution

RA conceived the idea. YCK and RA designed the research. YCK and PRS performed the

research. YCK and RA analysed the data and wrote the manuscript.

Funding

This study was supported by Ramanujan fellowship (SB/S2/RJN-036/2017, Department of

Science and Technology, India), Bill & Melinda Gates Foundation (OPP1210498), Indian

Institute of Science, Mr. Lakshmi Narayanan financial support and Department of

Biotechnology (Pallavi) for funding the project.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

Acknowledgements

We acknowledge Dr Sujoy K. Das (Bose Institute, Kolkata), Dr Urmi Bajpai (Acharya

Narendra Dev College, Delhi), Dr V.N. Azger Dusthackeer (National Institute for Research in

Tuberculosis, Chennai) and Dr Graham Hatfull (University of Pittsburgh) for providing us with

mycobacteriophages. We are grateful towards Dr Amit Singh (Indian Institute of Science,

Bengaluru) for providing bacterial strains. We thank Dr Nagasuma Chandra for kindly

providing us with the isoniazid-resistant *M. smegmatis*. We thank the Molecular Reproduction,

Development and Genetics, and Materials Engineering common facility for instrument access.

References

Agarwal, R., Johnson, C. T., Imhoff, B. R., Donlan, R. M., McCarty, N. A., and García, A. J.

(2018). Inhaled bacteriophage-loaded polymeric microparticles ameliorate acute lung

infections. *Nat. Biomed. Eng.* 2, 841–849. doi:10.1038/s41551-018-0263-5.

Aslam, B., Wang, W., Arshad, M. I., Khurshid, M., Muzammil, S., Rasool, M. H., et al.

(2018). Antibiotic resistance: a rundown of a global crisis. *Infect. Drug Resist.* 11,

1645–1658. doi:10.2147/IDR.S173867.

Azam, A. H., and Tanji, Y. (2019). Bacteriophage-host arm race: an update on the

mechanism of phage resistance in bacteria and revenge of the phage with the perspective

for phage therapy. *Appl. Microbiol. Biotechnol.* 103, 2121–2131. doi:10.1007/s00253-

019-09629-x.

Bajpai, U., Mehta, A. K., Eniyan, K., Sinha, A., Ray, A., Virdi, S., et al. (2018). Isolation and

characterization of bacteriophages from India, with lytic activity against Mycobacterium

tuberculosis. *Can. J. Microbiol.* 64, 483–491. doi:10.1139/cjm-2017-0387.

Bavda, V. R., and Jain, V. (2020). Deciphering the Role of Holin in Mycobacteriophage D29

Physiology. *Front. Microbiol.* 11, 883. doi:10.3389/fmicb.2020.00883.

Benítez-Guzmán, A., Esquivel-Solís, H., Romero-Torres, C., Arriaga-Díaz, C., and

Gutiérrez-Pabello, J. A. (2019). Genetically related Mycobacterium bovis strains

displayed differential intracellular growth in bovine macrophages. *Vet. Sci.* 6, 81.

doi:10.3390/vetsci6040081.

Boshoff, H. I. M., Myers, T. G., Copp, B. R., McNeil, M. R., Wilson, M. A., and Barry, C. E.

(2004). The transcriptional responses of Mycobacterium tuberculosis to inhibitors of

metabolism. Novel insights into drug mechanisms of action. *J. Biol. Chem.* 279, 40174–

40184. doi:10.1074/jbc.M406796200.

Broxmeyer, L., Sosnowska, D., Miltner, E., Chacón, O., Wagner, D., McGarvey, J., et al.

(2002). Killing of Mycobacterium avium and Mycobacterium tuberculosis by a

Mycobacteriophage Delivered by a Nonvirulent Mycobacterium: A Model for Phage

Therapy of Intracellular Bacterial Pathogens . *J. Infect. Dis.* 186, 1155–1160.

doi:10.1086/343812.

Cambau, E., Saunderson, P., Matsuoka, M., Cole, S. T., Kai, M., Suffys, P., et al. (2018).

Antimicrobial resistance in leprosy: results of the first prospective open survey

conducted by a WHO surveillance network for the period 2009–15. *Clin. Microbiol.*

Infect. 24, 1305–1310. doi:10.1016/j.cmi.2018.02.022.

Carrigy, N. B., Chang, R. Y., Leung, S. S. Y., Harrison, M., Petrova, Z., Pope, W. H., et al.

(2017). Anti-Tuberculosis Bacteriophage D29 Delivery with a Vibrating Mesh

Nebulizer, Jet Nebulizer, and Soft Mist Inhaler. *Pharm. Res.* 34, 2084–2096.

doi:10.1007/s11095-017-2213-4.

Carrigy, N. B., Larsen, S. E., Reese, V., Pecor, T., Harrison, M., Kuehl, P. J., et al. (2019).

Prophylaxis of Mycobacterium tuberculosis H37Rv Infection in a Preclinical Mouse

Model via Inhalation of Nebulized Bacteriophage D29. *Antimicrob. Agents Chemother.*

63, AAC.00871-19. doi:10.1128/AAC.00871-19.

Castells, M., Khan, D. A., and Phillips, E. J. (2019). Penicillin allergy. *N. Engl. J. Med.* 381,

2338–2351. doi:10.1056/NEJMra1807761.

Catalão, M. J., and Pimentel, M. (2018). Mycobacteriophage Lysis Enzymes: Targeting the

Mycobacterial Cell Envelope. *Viruses* 10, 428. doi:10.3390/v10080428.

Chan, B. K., Turner, P. E., Kim, S., Mojibian, H. R., Elefteriades, J. A., and Narayan, D.

(2018). Phage treatment of an aortic graft infected with Pseudomonas aeruginosa. *Evol.*

Med. Public Heal. 2018, 60–66. doi:10.1093/emph/eoy005.

Chhibber, S., Kaur, T., and Sandeep Kaur (2013). Co-Therapy Using Lytic Bacteriophage

and Linezolid: Effective Treatment in Eliminating Methicillin Resistant Staphylococcus

aureus (MRSA) from Diabetic Foot Infections. *PLoS One* 8, e56022.

doi:10.1371/journal.pone.0056022.

Comeau, A. M., Tétart, F., Trojet, S. N., Prère, M.-F., and Krisch, H. M. (2007). Phage-

Antibiotic Synergy (PAS): beta-lactam and quinolone antibiotics stimulate virulent

phage growth. *PLoS One* 2, e799–e799. doi:10.1371/journal.pone.0000799.

Danelishvili, L., Young, L. S., and Bermudez, L. E. (2006). Infection As Delivered by a

Nonvirulent Mycobacterium. *Microb. Drug Resist.* 12.

De Paepe, M., Leclerc, M., Tinsley, C. R., and Petit, M. A. (2014). Bacteriophages: An

underestimated role in human and animal health? *Front. Cell. Infect. Microbiol.* 5, 1–11.

doi:10.3389/fcimb.2014.00039.

Dedrick, R. M., Guerrero-Bustamante, C. A., Garlena, R. A., Russell, D. A., Ford, K., Harris,

K., et al. (2019). Engineered bacteriophages for treatment of a patient with a

disseminated drug-resistant Mycobacterium abscessus. *Nat. Med.* 25, 730–733.

doi:10.1038/s41591-019-0437-z.

Diel, R., Lipman, M., and Hoefsloot, W. (2018). High mortality in patients with

Mycobacterium avium complex lung disease: A systematic review. *BMC Infect. Dis.* 18,

206. doi:10.1186/s12879-018-3113-x.

Edwards, R. A., and Rohwer, F. (2005). Viral metagenomics. *Nat. Rev. Microbiol.* 3, 504–

510. doi:10.1038/nrmicro1163.

Ehlers, S., and Schaible, U. E. (2012). The granuloma in tuberculosis: Dynamics of a host-

pathogen collusion. *Front. Immunol.* 3, 1–9. doi:10.3389/fimmu.2012.00411.

Fenaroli, F., Westmoreland, D., Benjaminsen, J., Kolstad, T., Skjeldal, F. M., Meijer, A. H.,

et al. (2014). Nanoparticles as Drug Delivery System against Tuberculosis in Zebrafish

649 Embryos: Direct Visualization and Treatment. *ACS Nano* 8, 7014–7026.

650 doi:10.1021/nn5019126.

651 Geneva: World Health Organisation (2019). Global Tuberculosis Report. *Ann. ICRP* 39.

652 doi:10.1016/j.icrp.2009.12.007.

653 Gutierrez, A., Jain, S., Bhargava, P., Hamblin, M., Lobritz, M. A., and Collins, J. J. (2017).

654 Understanding and Sensitizing Density-Dependent Persistence to Quinolone Antibiotics.

655 *Mol. Cell* 68, 1147-1154.e3. doi:10.1016/j.molcel.2017.11.012.

656 Gygli, S. M., Borrell, S., Trauner, A., and Gagneux, S. (2017). Antimicrobial resistance in

657 Mycobacterium tuberculosis: Mechanistic and evolutionary perspectives. *FEMS*

658 *Microbiol. Rev.* 41, 354–373. doi:10.1093/femsre/fux011.

659 Hall, M. J., Middleton, R. F., and Westmacott, D. (1983). The fractional inhibitory

660 concentration (FIC) index as a measure of synergy. *J. Antimicrob. Chemother.* 11, 427–

661 433. doi:10.1093/jac/11.5.427.

662 Horsburgh, C. R. (1997). Epidemiology of Mycobacterium avium complex disease. *Am. J.*

663 *Med.* 102, 11–15. doi:10.1016/S0002-9343(97)00038-7.

664 Houben, E. N. G., Nguyen, L., and Pieters, J. (2006). Interaction of pathogenic mycobacteria

665 with the host immune system. *Curr. Opin. Microbiol.* 9, 76–85.

666 doi:10.1016/j.mib.2005.12.014.

667 Iseman, M. D. (1994). Evolution of drug-resistant tuberculosis: a tale of two species. *Proc.*

668 *Natl. Acad. Sci.* 91, 2428–2429. doi:10.1073/pnas.91.7.2428.

669 Jansen, M., Wahida, A., Latz, S., Krüttgen, A., Häfner, H., Buhl, E. M., et al. (2018).

670 Enhanced antibacterial effect of the novel T4-like bacteriophage KARL-1 in

671 combination with antibiotics against multi-drug resistant Acinetobacter baumannii. *Sci.*

672 *Rep.* 8, 14140. doi:10.1038/s41598-018-32344-y.

673 Jennes, S., Merabishvili, M., Soentjens, P., Pang, K. W., Rose, T., Keersebilck, E., et al.

(2017). Use of bacteriophages in the treatment of colistin-only-sensitive Pseudomonas

aeruginosa septicaemia in a patient with acute kidney injury-a case report. *Crit. Care* 21,

129. doi:10.1186/s13054-017-1709-y.

Jończyk-Matysiak, E., Łusiak-Szelachowska, M., Kłak, M., Bubak, B., Międzybrodzki, R.,

Weber-Dąbrowska, B., et al. (2015). The Effect of Bacteriophage Preparations on

Intracellular Killing of Bacteria by Phagocytes. *J. Immunol. Res.* 2015, 482863.

doi:10.1155/2015/482863.

Keshavjee, S., and Farmer, P. E. (2012). Tuberculosis, drug resistance, and the history of

modern medicine. *N. Engl. J. Med.* 367, 931–936. doi:10.1056/NEJMra1205429.

Khawaldeh, A., Morales, S., Dillon, B., Alavidze, Z., Ginn, A. N., Thomas, L., et al. (2011).

Bacteriophage therapy for refractory Pseudomonas aeruginosa urinary tract infection. *J.*

Med. Microbiol. 60, 1697–1700. doi:10.1099/jmm.0.029744-0.

Kim, M., Jo, Y., Hwang, Y. J., Hong, H. W., Hong, S. S., Park, K., et al. (2018). Phage-

Antibiotic Synergy via Delayed Lysis. *Appl. Environ. Microbiol.* 84, e02085-18.

doi:10.1128/AEM.02085-18.

Kortright, K. E., Chan, B. K., Koff, J. L., and Turner, P. E. (2019). Phage Therapy: A

Renewed Approach to Combat Antibiotic-Resistant Bacteria. *Cell Host Microbe* 25,

219–232. doi:10.1016/j.chom.2019.01.014.

Labrie, S. J., Samson, J. E., and Moineau, S. (2010). Bacteriophage resistance mechanisms.

Nat. Rev. Microbiol. 8, 317–327. doi:10.1038/nrmicro2315.

Lapenkova, M. B., Smirnova, N. S., Rutkevich, P. N., and Vladimirsky, M. A. (2018).

Evaluation of the Efficiency of Lytic Mycobacteriophage D29 on the Model of M.

tuberculosis-Infected Macrophage RAW 264 Cell Line. *Bull. Exp. Biol. Med.* 164, 344–

346. doi:10.1007/s10517-018-3986-0.

Lawn, S. D., and Zumla, A. I. (2011). Tuberculosis. *Lancet* 378, 57–72. doi:10.1016/S0140-

6736(10)62173-3.

Leung, C. Y. (Joey), and Weitz, J. S. (2017). Modeling the synergistic elimination of bacteria

by phage and the innate immune system. *J. Theor. Biol.* 429, 241–252.

doi:https://doi.org/10.1016/j.jtbi.2017.06.037.

Li, J., Xie, S., Ahmed, S., Wang, F., Gu, Y., Zhang, C., et al. (2017). Antimicrobial activity

and resistance: Influencing factors. *Front. Pharmacol.* 8, 1–11.

doi:10.3389/fphar.2017.00364.

Liu, K., Yang, W., Dong, X., Cong, L., Li, N., Li, Y., et al. (2016). Inhalation Study of

Mycobacteriophage D29 Aerosol for Mice by Endotracheal Route and Nose-Only

Exposure. *J. Aerosol Med. Pulm. Drug Deliv.* 29, 393–405.

doi:10.1089/jamp.2015.1233.

Lobritz, M. A., Belenky, P., Porter, C. B. M., Gutierrez, A., Yang, J. H., Schwarz, E. G., et

al. (2015). Antibiotic efficacy is linked to bacterial cellular respiration. *Proc. Natl. Acad.*

Sci. U. S. A. 112, 8173–8180. doi:10.1073/pnas.1509743112.

Lopatkin, A. J., Stokes, J. M., Zheng, E. J., Yang, J. H., Takahashi, M. K., You, L., et al.

(2019). Bacterial metabolic state more accurately predicts antibiotic lethality than

growth rate. *Nat. Microbiol.* 4, 2109–2117. doi:10.1038/s41564-019-0536-0.

Luthra, S., Rominski, A., and Sander, P. (2018). The role of antibiotic-target-modifying and

antibiotic-modifying enzymes in mycobacterium abscessusdrug resistance. *Front.*

Microbiol. 9, 1–13. doi:10.3389/fmicb.2018.02179.

Manrique, P., Bolduc, B., Walk, S. T., Van Oost, J. Der, De Vos, W. M., and Young, M. J.

(2016). Healthy human gut phageome. *Proc. Natl. Acad. Sci. U. S. A.* 113, 10400–

10405. doi:10.1073/pnas.1601060113.

Marinelli, L. J., Piuri, M., Swigoňová, Z., Balachandran, A., Oldfield, L. M., van Kessel, J.

C., et al. (2008). BRED: A Simple and Powerful Tool for Constructing Mutant and

Recombinant Bacteriophage Genomes. *PLoS One* 3, e3957.

doi:10.1371/journal.pone.0003957.

Mohan, A., Padiadpu, J., Baloni, P., and Chandra, N. (2015). Complete Genome Sequences

of a Mycobacterium smegmatis Laboratory Strain (MC2 155) and Isoniazid-Resistant

(4XR1/R2) Mutant Strains. *Genome Announc.* 3, 4–5. doi:10.1128/genomeA.01520-

14.Copyright.

Nieth, A., Verseux, C., Barnert, S., Süss, R., and Römer, W. (2015). A first step toward

liposome-mediated intracellular bacteriophage therapy. *Expert Opin. Drug Deliv.* 12,

1411–1424. doi:10.1517/17425247.2015.1043125.

Oechslin, F., Piccardi, P., Mancini, S., Gabard, J., Moreillon, P., Entenza, J. M., et al. (2017).

Synergistic Interaction Between Phage Therapy and Antibiotics Clears Pseudomonas

Aeruginosa Infection in Endocarditis and Reduces Virulence. *J. Infect. Dis.* 215, 703–

712. doi:10.1093/infdis/jiw632.

Oh, Y. K., and Straubinger, R. M. (1996). Intracellular fate of Mycobacterium avium: Use of

dual-label spectrofluorometry to investigate the influence of bacterial viability

opsonization on phagosomal pH phagosome-lysosome interaction. *Infect. Immun.* 64,

319–325. doi:10.1128/iai.64.1.319-325.1996.

Ooi, M. L., Drilling, A. J., Morales, S., Fong, S., Moraitis, S., MacIas-Valle, L., et al. (2019).

Safety and tolerability of bacteriophage therapy for chronic rhinosinusitis due to

staphylococcus aureus. *JAMA Otolaryngol. - Head Neck Surg.* 145, 723–729.

doi:10.1001/jamaoto.2019.1191.

Padiadpu, J., Baloni, P., Anand, K., Munshi, M. H., Thakur, C., Mohan, A., et al. (2016).

Identifying and tackling emergent vulnerability in drug-resistant mycobacteria. *ACS*

Infect. Dis. 2, 592–607. doi:10.1021/acsinfecdis.6b00004.

Padiadpu, J., Baloni, P., and Chandra, N. (2015). Gene expression profiles of wild-type and

isoniazid-resistant strains of Mycobacterium smegmatis. *Data Br.* 4, 186–189.

doi:10.1016/j.dib.2015.05.006.

Petrovic Fabijan, A., Lin, R. C. Y., Ho, J., Maddocks, S., Ben Zakour, N. L., Iredell, J. R., et

al. (2020). Safety of bacteriophage therapy in severe Staphylococcus aureus infection.

Nat. Microbiol. 5, 465–472. doi:10.1038/s41564-019-0634-z.

Pimentel, M. (2014). Genetics of Phage Lysis. *Microbiol. Spectr.* 2, MGM2—0017—2013.

doi:10.1128/microbiolspec.MGM2-0017-2013.

Piuri, M., and Hatfull, G. F. (2006). A peptidoglycan hydrolase motif within the

mycobacteriophage TM4 tape measure protein promotes efficient infection of stationary

phase cells. *Mol. Microbiol.* 62, 1569–1585. doi:10.1111/j.1365-2958.2006.05473.x.

Puapermpoonsiri, U., Spencer, J., and van der Walle, C. F. (2009). A freeze-dried

formulation of bacteriophage encapsulated in biodegradable microspheres. *Eur. J.*

Pharm. Biopharm. 72, 26–33. doi:https://doi.org/10.1016/j.ejpb.2008.12.001.

Rahman, M., Kim, S., Kim, S. M., Seol, S. Y., and Kim, J. (2011). Characterization of

induced Staphylococcus aureus bacteriophage SAP-26 and its anti-biofilm activity with

rifampicin. *Biofouling* 27, 1087–1093. doi:10.1080/08927014.2011.631169.

Rastogi, N., Legrand, E., and Sola, C. (2001). The Mycobacteria: An introduction to

nomenclature and pathogenesis. *OIE Rev. Sci. Tech.* 20, 21–54.

doi:10.20506/rst.20.1.1265.

Roach, D. R., Leung, C. Y., Henry, M., Morello, E., Singh, D., Di Santo, J. P., et al. (2017).

Synergy between the Host Immune System and Bacteriophage Is Essential for

Successful Phage Therapy against an Acute Respiratory Pathogen. *Cell Host Microbe*

22, 38-47.e4. doi:10.1016/j.chom.2017.06.018.

Rohde, K., Yates, R. M., Purdy, G. E., and Russell, D. G. (2007). Mycobacterium

tuberculosis and the environment within the phagosome. *Immunol. Rev.* 219, 37–54.

doi:10.1111/j.1600-065X.2007.00547.x.

Russell, D. A., and Hatfull, G. F. (2017). PhagesDB: the actinobacteriophage database.

Bioinformatics 33, 784–786. doi:10.1093/bioinformatics/btw711.

Samaddar, S., Grewal, R. K., Sinha, S., Ghosh, S., Roy, S., and Das Gupta, S. K. (2015).

Dynamics of Mycobacteriophage-Mycobacterial Host Interaction: Evidence for

Secondary Mechanisms for Host Lethality. *Appl. Environ. Microbiol.* 82, 124–133.

doi:10.1128/AEM.02700-15.

Samaddar, S., Grewal, R. K., Sinha, S., Ghosh, S., Roy, S., and Das Gupta, S. K. (2016).

Dynamics of mycobacteriophage-mycobacterial host interaction: Evidence for

secondary mechanisms for host lethality. *Appl. Environ. Microbiol.* 82, 124–133.

doi:10.1128/AEM.02700-15.

Schooley, R. T., Biswas, B., Gill, J. J., Hernandez-Morales, A., Lancaster, J., Lessor, L., et al.

(2017). Development and use of personalized bacteriophage-based therapeutic cocktails

to treat a patient with a disseminated resistant Acinetobacter baumannii infection.

Antimicrob. Agents Chemother. 61, e00954-17. doi:10.1128/AAC.00954-17.

Seto, S., Morimoto, K., Yoshida, T., Hiramatsu, M., Hijikata, M., Nagata, T., et al. (2020).

Proteomic Profiling Reveals the Architecture of Granulomatous Lesions Caused by

Tuberculosis and Mycobacterium avium Complex Lung Disease. *Front. Microbiol.* 10,

3081. doi:10.3389/fmicb.2019.03081.

Sibling, L. D., Franzblau, S. G., and Krahenbuhl, J. L. (1987). Intracellular fate of

Mycobacterium leprae in normal and activated mouse macrophages. *Infect. Immun.* 55,

680–685. Available at: https://www.ncbi.nlm.nih.gov/pubmed/3546136.

Sinha, A., Eniyan, K., Manohar, P., Ramesh, N., and Bajpai, U. (2020). Characterization and

genome analysis of B1 sub-cluster mycobacteriophage PDRPxv. *Virus Res.* 279,

197884. doi:https://doi.org/10.1016/j.virusres.2020.197884.

Sprenger, M., and Fukuda, K. (2016). New mechanisms, new worries. *Science (80-.).* 351,

1263–1264. doi:10.1126/science.aad9450.

Sula, L., Sulová, J., and Stolcpartová, M. (1981). Therapy of experimental tuberculosis in

guinea pigs with mycobacterial phages DS-6A, GR-21 T, My-327. *Czech. Med.* 4, 209–

14. Available at: http://www.ncbi.nlm.nih.gov/pubmed/7327068.

Sumitani, M., Takagi, S., Tanamura, Y., and Inoue, H. (2004). Oxygen Indicator Composed

of an Organic/Inorganic Hybrid Compound of Methylene Blue, Reductant, Surfactant

and Saponite. *Anal. Sci.* 20, 1153–1157. doi:10.2116/analsci.20.1153.

Swift, B. M. C., Gerrard, Z. E., Huxley, J. N., and Rees, C. E. D. (2014). Factors affecting

phage D29 infection: A tool to investigate different growth states of mycobacteria. *PLoS*

One 9, e106690. doi:10.1371/journal.pone.0106690.

Teng, R., and Dick, T. (2003). Isoniazid resistance of exponentially growing Mycobacterium

smegmatis biofilm culture. *FEMS Microbiol. Lett.* 227, 171–174. doi:10.1016/S0378-

1097(03)00584-6.

Timm, J., Post, F. A., Bekker, L. G., Walther, G. B., Wainwright, H. C., Manganelli, R., et al.

(2003). Differential expression of iron-, carbon-, and oxygen-responsive mycobacterial

genes in the lungs of chronically infected mice and tuberculosis patients. *Proc. Natl.*

Acad. Sci. U. S. A. 100, 14321–14326. doi:10.1073/pnas.2436197100.

Tkhilaishvili, T., Lombardi, L., Klatt, A. B., Trampuz, A., and Di Luca, M. (2018).

Bacteriophage Sb-1 enhances antibiotic activity against biofilm, degrades

exopolysaccharide matrix and targets persisters of Staphylococcus aureus. *Int. J.*

Antimicrob. Agents 52, 842–853. doi:10.1016/j.ijantimicag.2018.09.006.

Torres-Barceló, C., Arias-Sánchez, F. I., Vasse, M., Ramsayer, J., Kaltz, O., and Hochberg,

M. E. (2014). A window of opportunity to control the bacterial pathogen Pseudomonas

aeruginosa combining antibiotics and phages. *PLoS One* 9, 1–7.

doi:10.1371/journal.pone.0106628.

Trigo, G., Martins, T. G., Fraga, A. G., Longatto-Filho, A., Castro, A. G., Azeredo, J., et al.

(2013). Phage Therapy Is Effective against Infection by Mycobacterium ulcerans in a

Murine Footpad Model. *PLoS Negl. Trop. Dis.* 7. doi:10.1371/journal.pntd.0002183.

Turkington, C. J. R., Morozov, A., Clokie, M. R. J., and Bayliss, C. D. (2019). Phage-

Resistant Phase-Variant Sub-populations Mediate Herd Immunity Against

Bacteriophage Invasion of Bacterial Meta-Populations   . *Front. Microbiol.*   10, 1473.

Available at: https://www.frontiersin.org/article/10.3389/fmicb.2019.01473.

Vandenheuvel, D., Singh, A., Vandersteegen, K., Klumpp, J., Lavigne, R., and Van den

Mooter, G. (2013). Feasibility of spray drying bacteriophages into respirable powders to

combat pulmonary bacterial infections. *Eur. J. Pharm. Biopharm.* 84, 578–582.

doi:https://doi.org/10.1016/j.ejpb.2012.12.022.

Voskuil, M. I., Visconti, K. C., and Schoolnik, G. K. (2004). Mycobacterium tuberculosis

gene expression during adaptation to stationary phase and low-oxygen dormancy.

Tuberculosis 84, 218–227. doi:10.1016/j.tube.2004.02.003.

Waters, E. M., Neill, D. R., Kaman, B., Sahota, J. S., Clokie, M. R. J., Winstanley, C., et al.

(2017). Phage therapy is highly effective against chronic lung infections with

Pseudomonas aeruginosa. *Thorax* 72, 666–667. doi:10.1136/thoraxjnl-2016-209265.

Wipperman, M. F., Fitzgerald, D. W., Juste, M. A. J., Taur, Y., Namasivayam, S., Sher, A., et

al. (2017). Antibiotic treatment for Tuberculosis induces a profound dysbiosis of the

microbiome that persists long after therapy is completed. *Sci. Rep.* 7, 10767.

doi:10.1038/s41598-017-10346-6.

Figure 1. Efficacy of infection and evolution of phage resistance is dependent on individual Mycobacteriophages. A) Photograph showing spot clearance of a lawn of *M. smegmatis* after treatment with D29. Growth curves of *M. smegmatis* in liquid culture in the presence of **B)** D29, **C)** Che7, **D)** TM4, **E)** PDRPv and **F)** PDRPxv phage. Red arrow represents the time point at which phage solution was added to the cultures. These experiments were repeated several times (≥2) and the representative graphs were plotted. Multiple t-test with Holm-Sidak method was used to estimate the statistical significance. α represents a significant difference ($p<0.05$) between control and D29 treated (log phase) for all time points between 21 h and 73 h. β represents a significant difference ($p<0.05$) between control and Che7 treated (log phase) for all time points between 28 h and 44 h. χ represents a significant difference ($p<0.05$) between control and TM4 treated (log phase) for all time points between 21 h and 105 h. δ represents a significant difference ($p<0.05$) between control and PDRPv treated (log phase) for all time points between 28 h and 106 h. ε represents a significant difference ($p<0.05$) between control and PDRPxv treated (log phase) for all time points between 24 h and 106 h.

Figure 2. Phage cocktails show synergy compared to individual phage and their efficacy is MOI dependent. Growth kinetics of *M. smegmatis* cultures treated with **A)** 3-phage cocktail (D29, Che7 and TM4) and **B)** 5-phage cocktail (D29, TM4, Che7, PDRPv and PDRPxv) at MOI of 10 for each phage against a non-treated control. Growth kinetics of *M. smegmatis* cultures treated with a 5-phage cocktail at the MOI of **C)** 1, **D)** 0.1 and **E)** 0.01 for each phage in the phage cocktail. The experiments with MOI of 10 with 3- and 5-phage cocktails were repeated twice and the representative graphs are shown here. Red arrow represents the time point phage solution was added to the cultures. Multiple t-test with Holm-Sidak method was used to estimate the statistical significance (p<0.05). α represents a significant difference (p<0.05) between control and three-phage cocktail (D29, TM4 and Che7) treated (log phase) for all time points between 21 h and 105 h. β represents a significant difference (p<0.05) between control and five-phage cocktail (D29, TM4, Che7, PDRPv and PDRPxv) treated (log phase) for all time

points between 18 h and 346 h. χ represents a significant difference (p<0.05) between control and 5

phage cocktail treated (MOI 1, log phase) for all time points between 18 h and 270 h. δ represents a

significant difference (p<0.05) between control and 5 phage cocktail treated (MOI 0.1, log phase) for

all time points between 18 h and 174 h.

Figure 3. Mycobacteriophages are effective under various pathophysiological conditions. Growth

kinetics of *M. smegmatis* cultures treated with a 5-phage cocktail at **A)** pH condition of 6.0, **B)** pH

condition of 5.5, **C)** stationary phase and **D)** hypoxic conditions. For hypoxia experiment, solid lines

represent bacterial OD at 600 nm and dashed lines are the respective Methylene Blue readings at 665

nm. Red arrow represents the time point phage solution was added to the cultures. These experiments

were repeated twice and the representative graphs are shown here. For **A, B** and **C**, multiple t-test with

Holm-Sidak method was used to estimate the statistical significance (p<0.05). α represents a significant

difference (p<0.05) between control and phage cocktail treated (pH 6.0, log phase) for all time points

between 36 h and 338 h. β represents a significant difference (p<0.05) between control and phage

cocktail treated (pH 5.5, log phase) for all time points between 73 h and 338 h. χ represents a significant

difference (p<0.05) between control and phage cocktail treated (stationary phase) for all time points

between 24 h and 360 h. For **D,** 2-way ANOVA with Sidak's multiple comparisons test was used to

determine statistical significance (p<0.05). δ represents a significant difference between control and

phage (at low hypoxia) for all time points between 45 h and 432 h, and ε represents a significant

difference between control and phage (at high hypoxia) for all time points between 94 h and 432 h.

895
Figure 4. Phage cocktails show synergy with antibiotics and are effective against antibiotic-

resistant Mycobacterium. A) 5-phage cocktail (MOI 0.01) and rifampicin (20 µg/mL)- χ and δ

represent a significant difference between control and rifampicin + phage (log phase), and rifampicin +

phage (lag phase) for all time points between 30 h and 326 h. **B)** Checkerboard assay of M. smegmatis

cultures treated with rifampicin and 5-phage cocktail (D29, TM4, Che7, PDRPv and PDRPxv). **C)** Growth

kinetics of isoniazid-resistant M. smegmatis (4XR1) cultures treated with a 5-phage cocktail. represents

a significant difference between control and phage treated (log phase) for all time points between 40 h

and 310 h. Red arrow represents the time point antibiotic/phage solution was added to the cultures. For

A, 2-way ANOVA with Sidak's multiple comparisons test was used to determine statistical significance

(p<0.05). For C, multiple t-test with Holm-Sidak method was used to estimate the statistical significance

(p<0.05).

Figure 5. Phage cocktails are effective against M. tuberculosis. A) Growth kinetics of *M.*
tuberculosis (H37Ra) after treatment with a 3-phage cocktail (D29, Che7 and DS6A). Multiple t-test
with Holm-Sidak method was used to estimate the statistical significance (p<0.05). α represents a
significant difference between control and phage treated group (log phase) for all time points between
672 h and 1536 h. β represents a significant difference (p<0.05) between control and phage treated
group (lag phase) for all time points between 168 h and 1536 h. **B)** Colony Forming Unit (CFU)
measurements over time of *M. tuberculosis* cultures treated with a *3*-phage cocktail. χ represents a
significant difference (p<0.05) between control and phage (log phase) for all time points between 672
h and 1536 h. δ represents a significant difference (p<0.05) between control and phage (lag phase) for
all time points between 168 h and 1536 h. Multiple t-test with Holm-Sidak method was used to determine
the statistical significance. The detection limit for the assay is represented in black dotted line (10 CFU).