# EFF-1 promotes muscle fusion, paralysis and retargets infection by AFF-1-coated viruses in *C. elegans*

Anna Meledin[1], Xiaohui Li[1], Elena Matveev[1], Boaz Gildor[1], Ofer Katzir[1] and Benjamin Podbilewicz[1]*

[1]Department of Biology, Technion- Israel Institute of Technology, Haifa, 32000, Israel

*Correspondence to: podbilew@technion.ac.il

Running title: Abnormally fused muscles infected by fusexin-coated virus

**Graphical abstract:**

**A hallmark of muscle development is that myoblasts fuse to form myofibers. However,**
**smooth muscles and cardiomyocytes do not generally fuse. In *C. elegans*, the body wall**
**muscles (BWMs), the physiological equivalents of skeletal muscles, are mononuclear.**
**Here, to determine what would be the consequences of fusing BWMs, we express the**
**cell-cell fusogen EFF-1 in these cells. We find that EFF-1 induces paralysis and dumpy**
**phenotypes. To determine whether EFF-1-induced muscle fusion results in these**
**pathologies we injected viruses pseudotyped with AFF-1, a paralog of EFF-1, into the**
**pseudocoelom of *C. elegans*. When these engineered viruses encounter cells**
**expressing EFF-1 or AFF-1 they are able to infect them as revealed by GFP expression**
**from the viral genome. We find that AFF-1 viruses can fuse to EFF-1-expressing**
**muscles revealing multinucleated fibers that cause paralysis and abnormal muscle**
**morphogenesis. Thus, aberrant fusion of otherwise non-syncytial muscle cells may**
**lead to pathological conditions.**

**Significance statement**

Most cells are individual units that do not mix their cytoplasms. However, some cells fuse to become multinucleated in placenta, bones and muscles. In most animals, muscles are formed by myofibers that originate by cell-cell fusion. In contrast, in *C. elegans* the body wall muscles are mononucleated cells that mediate worm-like movement. EFF-1 and AFF-1 fusogens mediate physiological cell fusion in *C. elegans*. By ectopically expressing EFF-1 in body wall muscles we induce their fusion resulting in behavioral and morphological deleterious effects, revealing possible causes of congenital myopathies in humans. Using AFF-1-coated pseudoviruses we infect EFF-1-expressing muscle cells retargeting viral infection into these cells. We suggest that virus retargeting can be utilized to study myogenesis, neuronal regeneration, gamete fusion and screens for new fusogens in different organisms. In addition, our virus retargeting system can be used in gene-therapy, viral-based oncolysis and to study viral-host interactions.

**Introduction**

Vesicular Stomatitis Virus (VSV) is an enveloped negative strand RNA rhabdovirus. VSV utilizes its surface glycoprotein G (VSV-G) to infect vertebrates and invertebrates and lyses many cell lines tested to date [1–3]. VSV is widely used for pseudotyping other viruses and has a high transduction efficiency [1–4]. These properties turned VSVΔG into a promising vector for gene therapy, tissue regeneration, viral-based oncolysis, and VSV-based vaccines, some of which successfully completed phase III clinical trials [1,5]. Moreover, VSV-based vectors are useful for studying mechanisms of transcription and replication of RNA viruses, cellular trafficking, antiviral responses and fusion proteins (fusogens) [6–9]. However, the major bottleneck for both applied and fundamental research purposes is achieving an efficient and specific targeting of VSV-based vectors into desired cells/tissues of a live, multicellular organism [3]. Like other viruses, VSV lacks specificity for desired target cells as it suggestively enters cells through highly ubiquitous receptors such as the LDL receptor [10]. Retargeting VSV into particular tissues of interest therefore requires blocking the virus' natural interactions providing new, cell-specific interactions. Indeed, mutation or substitution of VSV-G with glycoproteins from other viruses or chimeric glycoproteins coupled with antibodies can retarget VSV to specific cells [3,4,11–13].

VSV-G is a class III viral fusogen whereas the *Caenorhabditis elegans* EFF-1 and AFF-1, which fuse somatic cells during development, are structural homologs of class II viral

proteins and gamete HAP2(GCS1) fusogens from the fusexin family [14–22]. EFF-1 and AFF-

1 also participate in maintaining neuronal architecture and neuronal reconnection following

injury [23–27]. Studied viral glycoproteins, including VSV-G, use a unilateral fusion

mechanism that depends on the expression of receptors only on the target cells [28,29]. In

contrast, EFF-1 and AFF-1 are bilateral fusogens- their presence is required on the

membranes of both apposing cells to mediate fusion [6,16,18,30,31]. These two fusogens can

act in either a homotypic or a heterotypic manner [6,16,18,21] and mediate heterotypic cell

fusion of Sf9 insect and Baby Hamster Kidney (BHK) cells [6,16]. Finally, VSV viruses

containing a *GFP* substituting the *VSV-G* coding sequence (VSVΔG) [32,33] that are coated

with AFF-1 (VSVΔG-AFF-1) specifically infect AFF-1 or EFF-1 expressing BHK cells [6]. Thus,

in contrast to the pseudotyped virus coated with the native, unilateral G glycoprotein (VSVΔG-

G), infection by VSVΔG-AFF-1 requires fusogen expression on the host cell membrane. To

date, however, VSVΔG coated with AFF-1, EFF-1 or any other non-viral fusogen have not

been tested for infection in a living organism. Recently VSVΔG-G was demonstrated to infect

living *C. elegans* [7,34]. Given that: (i) VSVΔG-AFF-1 requires a fusogen on the target cell for

infection, (ii) a detailed cellular atlas of AFF-1 and EFF-1 expression and function in *C. elegans*

is known and (iii) *C. elegans eff-1* and *aff-1* mutants are available, we test whether VSVΔG-

AFF-1 can be retargeted to specific cells in living *C. elegans*. We found that AFF-1-coated

viruses infect *C. elegans* and specifically target cells that express functional EFF-1 or AFF-1.

Furthermore, AFF-1-coated pseudoviruses can be redirected to mononucleated body wall

muscles (BWMs) that ectopically express EFF-1. Thus, the new delivery system enabled us

to observe that EFF-1-induced BWMs merger and formation of non-functional syncytial

muscle fibers, demonstrating the consequences of aberrant muscle fusion in an animal that

normally uses mononucleated muscles for gait. Based on our results, we propose that in *C.*

*elegans* a layer of longitudinal mononucleated muscles electrically coupled by gap junctions

efficiently mediate wavelike movement and ectopic fusion disrupts the morphology and

physiology of the muscular system. We also suggest additional applications for VSVΔG-AFF-

1 including finding new fusogens and fusogen-expressing tissues, studying fusogen-fusogen

interactions and fusogen-interacting proteins *in vivo,* studying neuronal regeneration

processes and using specific cell-delivery approaches for other viruses such as coronaviruses

and retroviruses in *C. elegans* and in other organisms.

99
100
**Figure 1. VSVΔG-AFF-1 and VSVΔG-G have different tropisms**
**(A)** BHK cells expressing AFF-1 (red pins) were infected with VSVΔG-G. The viral genome
encodes GFP replacing VSV-G (white pins). GFP(+) (green), infected cells. VSVΔG-AFF-1
pseudoviruses were harvested from the supernatant, titrated and microinjected into *C.*
*elegans*. **(B-F)** Confocal Z-stack projections of *drh-1* worms injected with VSVΔG-G (2300-
16000 IU) or VSVΔG-AFF-1 (33-240 IU) and imaged 48-72 h later (VSVΔG-G, n=65; VSVΔG-
AFF-1, n=46 from 5 and 6 independent experiments respectively). Images and their insets are
dot boxed. Arrowheads, hypodermal (HYP) nuclei. Arrows, infected cell. BWM-Body Wall
Muscle, UM-Uterine Muscle, SM-Stomatointestinal Muscle, EC-Excretory Cell, P-Pharynx.
Scale bars, 50 µm. **(G)** Distribution of worms with specified infected, GFP(+) cell types. **(H)**
Fraction of GFP(+) worms injected with indicated IU of VSVΔG-G or VSVΔG-AFF-1. Each dot
represents an independent experiment (n=10-28 worms per experiment). $R^2$ and the IU doses
producing 50% GFP(+) worms (ID50) are indicated. See also Figure S1.

**Results**

**VSV△G-AFF-1 and VSV△G-G infect mostly hypodermal and muscle cells respectively**

Wildtype, recombinant or pseudotyped VSV has been shown to infect rodents, fish, farm animals, and primates [1,35–37]. Recently, VSV∆G-G was shown to infect *C. elegans* [7]. Previously, pseudotyped VSV expressing the *C. elegans* somatic fusogen AFF-1 (VSV∆G-AFF-1) was shown to infect AFF-1 or EFF-1 expressing BHK cells [6], but was not tested in a living organism. To test whether pseudotyped VSV∆G-AFF-1 [6,33,38] can infect *C. elegans*, we prepared a recombinant VSV strain encoding the fluorescent reporter GFP, coated by AFF-1 fusion protein (VSV∆G-AFF-1). Briefly, BHK cells expressing AFF-1 were infected with VSV∆G-G helper virus (Fig. 1A). Newly generated virions were coated by plasma membrane-bound AFF-1, thereby producing VSV∆G-AFF-1 pseudotyped viruses. Then, we titered the viruses determining the number of viral Infective Units (IU)/ml in BHK-AFF-1 cells. Finally, VSV∆G-AFF-1 viruses were injected into worm's pseudocoelom; a body cavity filled with fluid that surrounds the internal organs. We expected that virus injected into the pseudocoelom will encounter different cells inside the worm (Figure 1A). Worms were injected with VSV∆G-AFF-1 that was pre-incubated with αVSV-G antibody (see materials and methods and Figure S4), VSV∆G-G (as a positive control) or DMEM medium (as negative control). We used *drh-1(-)* (Dicer Related Helicase -1) mutant worms, as they are more susceptible to VSV infection and do not have any observable phenotypes [7,34]. Taken that VSV-G works unilaterally, while AFF-1 and EFF-1 fusogens have to be present in both fusing membranes [6,16,18,30], we hypothesized that VSV∆G-G and VSV∆G-AFF-1 will produce different infection patterns. Hence, we characterized and compared the GFP positive (GFP(+)) cells infected by VSV∆G-G or VSV∆G-AFF-1. VSV∆G-G preferentially infected muscle cells, including BWM [7], uterine muscles (UM) and stomatointestinal muscles (SM, Figures 1B and 1C). VSV∆G-G also infected the epidermal cells (hypodermis, HYP), excretory canal cell (EC, Figure 1B), glia and neurons in the head (Figures S1A-S1C). In contrast, VSV∆G-AFF-1 infected mostly HYP cells (Figure 1D), and also excretory cell (Figure 1E) pharyngeal muscles (P) (Figure 1D), BWM (Figure 1F) and head neurons and glia (Figures S1D-S1F). VSV∆G-G infects muscles that do not express any known fusogen (67% of infected, GFP(+) worms had only BWM infection), whereas VSV∆G-AFF-1 mainly infects the EFF-1-expressing hypodermis (61% of GFP(+) worms had only hyp infection) (Figure 1G). Moreover, the calculated ID50 (infection dose producing 50% GFP(+) worms) for VSV∆G-G and VSV∆G-AFF-1 were approximately 1000 and 20 IU respectively (Figure 1H). Therefore, VSV∆G-AFF-1, a virus pseudotyped with a bilateral fusogen, can efficiently infect a multicellular organism such as *C. elegans*. Moreover, VSV∆G-AFF-1 and VSV∆G-G are specialized for different cellular targets and have different infectivity levels.

**Figure 2. VSVΔG-AFF-1 requires EFF-1 on target cells for infection**
**(A-C)** Confocal Z-stack projections of *C. elegans* infected with VSVΔG-AFF-1 (33-240 IU) and imaged 48 h later. *drh-1* (n=46), *drh-1;eff-1(-)* (n=15) and *drh-1;aff-1(-)* (n=10).
**(D)** Fraction of GFP(+) worms from (A-C).
**(E-H)** Confocal Z-stack projections of *C. elegans* infected with VSVΔG-G (2300-4700 IU) 48 h post-injection. *drh-1* (n=56), *drh-1;eff-1(-)* (n=39) or *drh-1;aff-1(-)* (n=15).
**(I)** Fraction of GFP(+) worms from (E-H).
Arrowheads, HYP nuclei. Arrows, infected cell. See Figure 1 for cell types. Scale bars, 100 μm. Error bars represent mean ± SEM. *p<0.05, ***p<0.001 (Student's t- test).
See also Figure S2.

**VSV△G-AFF-1 requires EFF-1 in host cells for infection**

We showed that cells infected by VSV△G-AFF-1 included hypodermis, excretory cell, neurons in the head, glia and pharyngeal cells. It is known that in adults pharyngeal cells, head neurons, glia and hypodermal cells express EFF-1 [14], while pharyngeal cells, excretory duct cell and glia express AFF-1 [18,21] and lastly, excretory cell, head neurons, hypodermal cells and pharyngeal cells express EFF-2, a recent duplication of EFF-1 with unknown function (BG, Oren-Suissa and BP unpublished data). Importantly, BWM cells that do not express any known fusogen in *C. elegans* were infected by VSV△G-AFF-1 in only 9% of the worms compared to 67% of the worms infected with VSV△G-G (Figure 1G). These findings suggest that infection of living *C. elegans* by VSV△G-AFF-1 follows a bilateral action mechanism, and that VSV△G-AFF-1 is capable of interacting with several members of the fusexin family on the target cell. Therefore, we hypothesized that EFF-1-expressing hypodermal cells could not be infected by VSV△G-AFF-1 in the absence of EFF-1, while cells expressing AFF-1 or EFF-2 would still be targeted. To test this hypothesis, we injected either VSV△G-G or VSV△G-AFF-1 into the pseudocoelom of wt, null *eff-1(ok1021)* mutants *((eff-1(-);*[14]), null *aff-1(tm2214)* mutants ((*aff-1(-)*;[18]) or putative null *eff-2(hy51)* (*eff-2(-)*; see materials and methods) worms, and quantified the percentage of worms with GFP(+) cells. We detected VSV△G-AFF-1 infected hypodermal cells in *eff-1(+)* (Figure 2A) but not in in *eff-1(-)* worms (Figure 2B), while GFP(+) excretory and glia cell were detected even in the absence of EFF-1 (Figure 2B). *aff-1(-)* worms had GFP(+) hypodermal cells as in wildtype (Figure 2C). Overall, the fraction of worms showing VSV△G-AFF-1-infected hypodermal cells was similarly high in wildtype and in *aff-1(-)* (87% and 92% for wt and *aff-1(-),* respectively) but was absent in *eff-1(-)* mutants (Figure 2D). In addition, muscles were inefficiently infected in wildtype and *aff-1(-)* mutants, and not infected in *eff-1(-)* animals (Figure 2D). The fraction of worms with GFP(+) excretory cells was not significantly different in all backgrounds (Figure 2D). Finally, the fraction of worms with GFP(+) glia/neuron cells was significantly higher in *eff-1(-)*, compared to wt and *aff-1(-)* animals (Figures S1D-S1F). Moreover, *eff-2(-)* and wildtype animals were similarly infected in their hypodermal, muscle and excretory cells (Figure S2). These results show that VSV△G-AFF-1 requires EFF-1 expressed on hypodermal cells, for their infection. Moreover, we show evidence for AFF-1-EFF-1 bilateral heterotypic interactions *in vivo*.

EFF-1 mediates formation of hyp7 syncytium by fusing 139 hypodermal cells during embryonic and larval development and *eff-1(-)* worms have mononucleated hypodermal cells instead of this syncytium [14]. To exclude the possibility that hypodermal cells of *eff-1(-)* worms have a reduced susceptibility to viral infection, we injected wt, *aff-1(-)* and *eff-1(-)* worms with VSV△G-G. We found that wild-type worms injected with VSV△G-G, presented a full body-length covered by infected hypodermal cells along with muscles and excretory cells (Figures

2E-2F), while *eff-1(-)* worms had only few GFP(+) hyp nuclei localized closer to the injection region- the head, as well as muscles and excretory cells (Figure 2G). *aff-1(-)* worms had hypodermal infection similar to wt animals (Figure 2H). Yet, the fraction of worms injected with VSVΔG-G that showed GFP(+) muscle and hypodermal cells was not significantly different in wt, *eff-1(-)* and *aff-1(-)* backgrounds (Figure 2I). Finally, we observed a small fraction of worms with GFP(+) glia/neurons in the head with no significant difference between all tested backgrounds (Figures S1A-S1C). Thus, hypodermal cells are not compromised for VSVΔG-G infection even in the absence of EFF-1. In addition, VSVΔG-G with its unilateral viral fusogen infects different cells including muscles, hypodermis, excretory cell, and glia/neurons in an EFF-1- and AFF-1-independent manner. This is in contrast to VSVΔG-AFF-1 that uses a bilateral mechanism and infects cells that normally express EFF-1 or AFF-1 in adult animals.

**VSVΔG-AFF-1-mediated infection depends on *eff-1* activity in target cells**

We have shown VSVΔG-AFF-1 fails to infect hypodermal cells in *eff-1(-)* animals (Figure 2). To study whether conditional induction of *eff-1* can trigger infection by VSVΔG-AFF-1 we varied the dosage of *eff-1* using temperature shifts in a conditional temperature sensitive (ts) mutant [14,25]. In *eff-1(hy21ts)* animals grown at the permissive 15°C, most epidermal cells fuse to form multinucleated hypodermis and vulva [14]. However, when *eff-1(hy21)* animals are grown at the restrictive 25°C, epidermal cells fail to fuse, producing phenotypes similar to the null *eff-1(ok1021)* worms [16]. Therefore, *eff-1(ts)* worms can serve as a temperature inducible system for modification of EFF-1 dosage *in vivo* [14,19,25]. To obtain varying expression of EFF-1, we maintained *eff-1(ts)* animals at 15°C, 25°C or performed 25°C to 15°C downshifts at different developmental stages and then injected VSVΔG-AFF-1 into young adult worms (Figure 3A). Wildtype worms in all experimental conditions had GFP(+) hyp nuclei spanning the worm's body from head to tail (Figures 3B-3E), while *eff-1(ts)* animals maintained at the restrictive temperature and then shifted down from 25°C to 15°C, had a gradual increase in infected GFP(+) hyp cells (Figures 3F-3I). In wt worms maintained in all conditions, the number of GFP(+) hypodermal nuclei was about 69, consistent with the number of hyp7 nuclei found on one side of the animals body [39,40]. In contrast, for *eff-1(ts)*, the longer time the worms developed at the permissive temperature (15°C), the more GFP(+) hyp nuclei were scored (Figure 3J). Thus, hypodermal infection by VSVΔG-AFF-1 increases with conditional induction of *eff-1*.

**Figure 3. Infection by VSVΔG-AFF-1 increases with induction of EFF-1 function**

**(A)** Wt or temperature sensitive *eff-1(ts)* animals were maintained at permissive (15°C) or restrictive (25°C) temperatures for different times during their development. Each row represents a different experimental condition. Blue needles indicate virus injection. L2, Larval stage 2; L3, Larval stage 3.

**(B-I)** Confocal Z-stack projections of worms infected with VSVΔG-AFF-1. *drh-1* or *drh-1;eff-1(ts)* animals were injected with VSVΔG-AFF-1 (7-125 IU) as adults, and imaged 48 h later. Arrowheads, HYP nuclei. Asterisks, patches of GFP(-) surrounded by GFP(+) hypodermal cells. Scale bar, 100 μm.

**(J)** Quantitation of experiments. Error bars average ± SEM. Student t-test. *p<0.05, **p<0.001. n=3-10 worms per condition. Triangle, dosage of *eff-1*.

**Ectopic EFF-1 expression in BWMs produces dumpy and uncoordinated worms**

We demonstrated that VSVΔG-AFF-1 specifically infects EFF-1 expressing cells and rarely infects muscle cells. In addition, EFF-1-dependent infection could be induced by EFF-1(ts) conditional expression in target cells (Figure 3). Hence, we propose that ectopic EFF-1 expression in host cells, such as body wall muscles (BWMs), could retarget VSVΔG-AFF-1 into these cells. *C. elegans* has 95 mononucleated rhomboid BWMs, arranged as staggered pairs and bundled in four quadrants that run along the worm's body [41–43]. While most vertebrates and invertebrates have syncytial striated muscles composed of long multinucleated myofibers, in *C. elegans* and other nematodes the BWMs have not been described to fuse and do not form multinucleated myofibers (https://www.wormatlas.org/hermaphrodite/muscleintro/MusIntroframeset.html). Therefore, we hypothesized that EFF-1 expression in BWMs can induce their ectopic fusion, alter their structure, produce muscle-related phenotypes and increase VSVΔG-AFF-1 infection of BWMs. First, we monitored animals with an extra-chromosomal array containing genomic EFF-1 under the muscle-specific *myosin-3* promoter (*myo-3p*::EFF-1), together with *myo-3p::*mCherry which labels cytoplasm and nuclei in BWMs, enteric, gonadal and vulval muscles. Most adult animals, including *myo-3p*::mCherry (+) and (-), were wild-type-like (Table S1, Figures 4A-4B and Movie S1), while about 10% of animals were both *myo-3p*::mCherry(+) and had an uncoordinated and dumpy (Unc+Dpy) phenotype (Table S1, Figures 4C-4D and Movie S1). Most *myo-3p*::mCherry(+), Unc+Dpy animals arrested during larval development (Table S1) with additional phenotypes including bridged BWMs behind the terminal bulb (0/12 wt-like worms and 9/15 Unc+Dpy worms), mCherry-labeled aggregates that could be observed with DIC (0/12 wt-like worms and 9/15 Unc+Dpy worms) (Figures 4E-4H), *myo-3p*::mcCherry(+) BWMs with clustered nuclei (3/12 wt-like worms and 9/15 Unc+Dpy worms) and elevated number of nuclei/ BWM cell (1.4±0.2 and 2.2±0.3 nuclei/BWM cell in wt-like vs Unc+Dpy larvae respectively) (Figures 4E-4I). While ectopic *hsp*::EFF-1 expression causes embryonic lethality [14,19,25], we found no difference between the fractions of *myo-3p*::mCherry (+) and (-) unhatched eggs (Table S1). Thus, EFF-1 expression in BWMs accounts for the Unc+Dpy phenotypes, larval arrest and clustered BWM nuclei, which supports the hypothesis that ectopic EFF-1 induces BWMs multinucleation by cell-cell fusion causing behavioral and morphological phenotypes.

**Figure 4. EFF-1 expression in BWMs produces Unc+Dpy worms with multinucleated cells**

**(A-H)** Images of fluorescent Z-stack projections and respective DIC of animals with extrachromosomal *myo-3p*::EFF-1*, myo-3p*::mCherry. (G) White arrowhead, bridge formed between 2 BWMs from opposing quadrants. Yellow arrows, *myo-3p*::mCherry accumulations also in DIC (H). Asterisks, clustered nuclei within one BWM. Scale bars, 100 µm.

**(I)** Number of nuclei per *myo-3p*::mCherry (+) BWM cell in L2s. wt-like (n=12); Unc+Dpy (n=15). Each dot represents the average number of nuclei/BWM cell/worm. Total average ± SEM for each phenotype. Two tailed Student's t-test p<0.05*.

See also Table S1.

**EFF-1 expression in BWMs induces their fusion and retargets VSVΔG-AFF-1 to muscles**

To determine whether EFF-1 expression in BWMs can induce their fusion, we imaged BWMs

expressing a membrane-bound YFP (MB::YFP), *myo-3p*::EFF-1 and *myo-3p*::mCherry, (see

materials and methods). BWMs of wt-like worms have a normal spindle-shaped morphology

with cellular membranes between adjacent BWMs showing different levels of cytoplasmic *myo-3p*::mCherry (as *myo-3p*::mCherry is an extrachromosomal array) (Figures 5A-5C). In contrast, Unc+Dpy worms had disordered muscle fiber structure, the membranes surrounding the BWMs were indistinguishable and the BWMs showed an evenly distributed cytoplasmic *myo-3p*::mCherry indicating fusion and content mixing between these cells during development (Figures 5D-5F). Thus, EFF-1 expression in BWMs induces disappearance of cellular membranes and cytoplasmic merger, suggesting that EFF-1-mediated cell-cell fusion alters BWM structure, likely resulting in Unc+Dpy and larval arrest phenotypes.

We showed that VSV△G-AFF-1 infects EFF-1-expressing hypodermal cells in >80% of the infected worms but infects muscle cells that do not express EFF-1 in about 20% of the infected animals (Figure 2D). Moreover, EFF-1 expression in BWMs induced their fusion, and generated Unc+Dpy worms (Table S1, Movie S1 and Figures 4A-4D and 5A-5F). Taken that EFF-1 and AFF-1 are bilateral fusogens and VSVG is a unilateral fusogen, we hypothesized that ectopic expression of EFF-1 in BWMs would increase their infection by VSV△G-AFF-1 but not by VSV△G-G. Therefore, we injected either VSV△G-AFF-1 or VSV△G-G into the pseudocoelom of worms ectopically expressing EFF-1 and mCherry in BWMs and imaged them. For VSV△G-AFF-1 but not for VSV△G-G, the fraction of infected BWMs was significantly higher in Unc+Dpy worms (expressing the *myo-3p*::mCherry marker) compared to wt-like (expressing the *myo-3p*::mCherry marker) and wt animals (Figure 6). Thus, EFF-1 expression in BWMs retargets VSV△G-AFF-1 into muscles but does not affect the natural VSV△G-G tropism for BWMs.

We showed that VSV△G-AFF-1 infects neuron and glia cells in the head (Figures S1D-S1F). In order to assess the infection of other fusogen-expressing neurons, we chose the arborized neuron pair PVD, previously shown to express EFF-1, which spans the length of the worm [25], VSV△G-AFF-1-infected PVD cells were not observed. Nevertheless, we found 3 wt animals (out of >100 infected worms), where PVD cells were infected with VSV△G-G following injections with high dosage of pseudotyped viruses (Movie S2). Moreover, we tested whether ectopic AFF-1 expression in PVD or ectopic EFF-1 expression in 12 mechanosensory and chemosensory neurons in *eff-1(ts)* background at the restrictive 25ºC, could induce VSV△G-AFF-1 infection in these cells. None of these treatments produced VSV△G-AFF-1 infection in the neurons ectopically expressing EFF-1 or AFF-1 (Figure S3 and Table S2). There are several possibilities to explain why VSV△G-AFF-1 did not infect these neurons; (i) relatively low titer of VSV△G-AFF-1 preps, (ii) a physical barrier such as hypodermis, sheath cells or glia cells that surround neurons or (iii) diffusion of the GFP signal in long neuronal processes.

**Figure 5. EFF-1 expression in BWMs induces their fusion**

(**A-C**) Confocal images of wt-like adult worms with membrane bound (*MB*) *myo-3p::MB::YFP* (cyan) and extrachromosomal array containing *myo-3p::*EFF-1, *myo-3p::*mCherry (magenta). (**D-F**) Confocal images of Unc+Dpy [*myo-3p::MB::YFP* (cyan); *myo-3p::EFF-1*, *myo-3p::mCherry].* Arrows, unfused BWMs with MB (cyan). Note only two unfused BWMs, all the others appear fused with no MB separating them. Insets correspond to white-dotted area. Scale bars 100 μm.

See also Movie S1.

**Figure 6. Retargeting of VSVΔG-AFF-1 to body wall muscle cells**
Wild-type worms and animals with extrachromosomal array containing *myo-3p::*EFF-1 and *myo-3p::*mCherry were injected with VSVΔG-AFF-1 (35-63 IU, red pins; n=39 wt, n=50 wt-like and n=27 Unc+Dpy worms) or VSVΔG-G ($3*10^5$ IU, white pins; n=30 wt-like and 14 Unc+Dpy) respectively. Wt worms injected with VSVΔG-G (2300-4700 IU, n=56) were taken from figure 2I. Animals were analysed by SDC microscopy. Data represents average percentage of worms with GFP(+) BWMs ± SEM. Student's t-test: *$p < 0.05$.

**EFF-1 expression in fused BWMs enables VSVΔG-AFF-1 and VSVΔG-G spreading**

Based on the aberrant muscle fusion phenotype seen in Unc+Dpy worms following EFF-1 expression in BWMs, we hypothesized that viral spreading through such novel BWM syncytia would be enhanced. To test this hypothesis, we injected VSVΔG-AFF-1 or VSVΔG-G into wt-like or Unc+Dpy worms and quantified the number of BWM cells that were either (i) mCherry(+) only (expressing EFF-1 and *myo-3p*::mCherry(+)), (ii) GFP(+) only (infected but not expressing EFF-1) or (iii) mCherry(+) and GFP(+) expression in the same cells (infected and expressing EFF-1). We found that for both VSVΔG-AFF-1 and VSVΔG-G, wt-like worms had individual GFP(+) BWM cells, that were mostly mCherry (-) (Figures 7A-7F, arrows), while Unc+Dpy animals had continuous GFP(+) BWMs that overlapped with mCherry along their body length (Figures 7G-7L, dashed lines and arrows). Compared to wt-like animals, the Unc+Dpy worms had about five-times fewer non-infected mCherry(+) only BWMs, a similar number of infected-GFP(+) only cells and about fifteen-fold more infected BWM cells that also express mCherry (Figures 7M-7N). Lastly, Unc+Dpy worms had ~2.5 nuclei per VSVΔG-AFF-1- or VSVΔG-G- infected BWMs, compared to wt-like animals with 1 nucleus per BWM cell (Figures 7O-7P). Therefore, EFF-1 expression in BWMs transforms mononucleated BWMs into syncytial muscle fibres enabling viral spreading within the fused cells and increasing the number of both VSVΔG-AFF-1- and VSVΔG-G-infected BWM cells.

**Figure 7. EFF-1 expression in BWM enables VSV△G-AFF-1 and VSV△G-G spreading along fused muscles**

**(A-L)** Z-stack projections of wt-like (A-F) and Unc+Dpy animals (G-L) expressing *myo-3p::*EFF-1 and *myo-3p::*mCherry infected with VSV△G-AFF-1 (35-63 IU red pins) or VSV△G-G (3*10⁵ IU; white pins). Insets and their corresponding images (yellow frames). Arrows, individual infected (cyan) BWMs. Dashed lines outline grouped BWMs that express *myo-3p*::mCherry and *myo-3p::*EFF-1 (magenta) and infected with virus (cyan) showing spreading of GFP. Scale bars, 100 μm. **(M-N)** Number of BWM cells/worm expressing EFF-1 (magenta cell), infected (cyan cell) or expressing EFF-1 and infected (magenta and cyan). wt-like (circles) and Unc+Dpy (triangles). Each point represents a single worm. **(O-P)** Quantitation of multinucleation of infected BWMs. Each dot represents an average number of nuclei/ GFP(+) BWM, calculated from 1-6 multinucleated BWMs of a single worm. (M and O) wt-like n=6 and Unc+Dpy n=10 animals. (N and P) wt-like n=9 and Unc+Dpy n=7 animals. Black horizontal lines, average ± SEM. Student's t-test, *** p<0.0001; n.s., not-significant.

**Discussion**

**Utilizing VSV△G-AFF-1 to study muscle fusion and muscle fusogens**

Myogenesis of striated muscle involves the formation of multinucleated myofibers by myoblast-myoblast fusion. Myoblast fusion is also essential for muscle tissue patterning, maintenance and damage repair [62,63]. In contrast, *C. elegans* BWMs do not fuse and exist as mononucleated cells, contracting in synchrony that mediate gait [64]. We used VSV△G-AFF-1 to label EFF-1-expressing BWMs and found that EFF-1-mediated BWMs fusion results in loss of coordination, although we did not determine whether the observed locomotory defects result from aberrant joining of separately-innervated muscles or from a loss of contractility. These results demonstrate that a locomotory circuit may be optimally adjusted for both separate (as in *C. elegans*) and syncytial (as in vertebrate) striated muscles. Indeed, failure of myoblast fusion in vertebrates may have equally severe phenotypes [47,48]. Despite the importance of muscle fusion, the studies of muscle fusion proteins are just emerging [55–58]. We suggest that VSV△G-AFF-1 can be used in screens to identify additional muscle fusogens in invertebrates (e.g. Drosophila). Alternatively, viruses coated with the bilateral Myomaker, can be a tool to screen for muscle-expressed Myomaker-interacting proteins in vertebrates.

**VSV△G-AFF-1 infects, and can be retargeted to muscles expressing EFF-1 or AFF-1**

Previously described viral-vector-based systems used viral unilateral fusogens to target desired cells. For instance, fusogens that are either native, modified or from different viral origin, change the specificity of the viruses targets and facilitate the delivery process [1–3]. In addition, a potent targeting of cancer cell lines was achieved with adenovirus (non-enveloped virus) armed with an adaptor composed from 2 designed ankyrin repeat proteins (DARPins), one binding to the viral fiber knob and the second binding to cancer-specific markers [44]. Our work presents a new approach to retarget viral vectors into muscle cells, based on viruses coated with a nematode bilateral fusogen, namely AFF-1 and expression of bilateral fusogens such as EFF-1/AFF-1 in desired host cells. VSV△G-AFF-1 has a high degree of specificity towards cells that express EFF-1/AFF-1 and can be redirected by expressing them on a certain cell (e.g. BWMs). Moreover, VSV△G-AFF-1-mediated infection can be manipulated quantitatively by changing the amount of a partner fusogen expressed on the target cells. VSV and VSV-pseudotypes successfully infect invertebrate and vertebrate cell-lines and model organisms [1,2,35–37]. Although EFF-1 and AFF-1 are not present, and have no known homologs in vertebrates, VSV△G-AFF-1 infects mammalian BHK cells expressing AFF-1/EFF-1 [6]. It will be interesting to test if VSV△G-AFF-1 can specifically target cells ectopically expressing AFF-1/EFF-1 in a vertebrate model. Similarly, pseudotyped VSV△G-SARS-CoV-

2-S-glycoprotein [4] could be targeted to humanized worms or mice expressing hACE2 to
study entry mechanisms, pathogenesis and potential treatments for viral infections, including
COVID19 in model organisms.

**Additional applications for VSVΔG-AFF-1**

In the recent decade, extracellular vesicles (EVs) emerged as intercellular communication

carriers, containing variable cargoes (e.g lipids, DNA, RNA, toxins and proteins) and involved

in different biological processes including cell-cell interactions, cancer, tissue and neuronal

regeneration. EVs can serve as biomarkers and are potent vehicles for gene and drug delivery

[26,45,46]. However, work with EVs-based vectors has to tackle the issues of specific

targeting, loading of desired cargo, presence of undesired/non-specific content and

quantification of produced EVs. Our system consisting of enveloped pseudotyped virus coated

with bilateral fusogens and host muscle cells expressing bilateral fusogens, demonstrates cell-

specific targeting. Moreover, VSVΔG-AFF-1 has the benefits of viral-based vectors, namely

loading a specific and tailored cargo including fluorescent markers such as GFP-coding RNA

sequence, simple vector quantification and cargo amplification within the host cell. Therefore,

VSVΔG-AFF-1 may serve as a paradigm for EVs-mediated transport. It can be utilized to

explore how EVs deliver their cargos and to study whether and how EVs are involved in

different biological processes.

Exoplasmic membrane fusion is a fundamental biological process involved in

myogenesis, fertilization, vulva development, bone formation and resorption, hypodermis

morphogenesis, placentation, tubulogenesis, neuronal regeneration and viral infection

[28,31,47,48]. Despite the importance of these processes, very few cell-cell fusogens have

been identified and characterized so far. These include: (i) Syncytins involved in placenta

formation [49–51]; (ii) Fusexins including EFF-1 and AFF-1 from *C. elegans* [14,17–22,31] and

HAP2/GCS1 which mediates gamete fusion in protists and flowering plants [38,52–54] and

(iii) Myomaker (TMEM8c) and myomerger (myomixer/minion) proteins that fuse muscle cells

in vertebrates [55–58]. We demonstrate that VSVΔG-AFF-1 injected into *C. elegans*

encounters and infects different EFF-1/AFF-1-expressing cells. Ectopic expression of EFF-1

in BWMs induces these mononucleated muscles to form syncytial myofibers, however, these

multinucleated muscles are pathological and the worms become deformed and paralyzed.

Moreover, bilateral heterotypic fusion has been described between EFF-1 and AFF-1 and also

between EFF-1 and sperm HAP2/GCS1 from *Arabidopsis* expressed on two populations of

BHKs [38]. Therefore, VSVΔG-AFF-1/EFF-1 may be exploited in screens to identify new

bilateral fusogens or fusogen-interacting proteins in different organisms and to understand

how and why some myoblasts fuse (e.g. skeletal muscles) while others function optimally as

individual mononucleated cells (e.g. smooth muscles and BWMs).

**Materials and methods**

**Nematode strains**

Unless otherwise stated, all nematodes were maintained at 20°C, according to standard

protocols [65,66]. *drh-1(tm1329)* served as the wild type background. Mutations and strains

that were used in this study are listed in Table S3.

**DNA constructs**

The *myo-3p*::EFF-1 plasmid was constructed by cloning the *myo-3* promoter region from *myo-*

*3p*::mCherry plasmid with Sal I (New England BioLabs Cat#R3138) and Nhe I (ThermoFisher

Cat# FD0974) and inserting it into the *hsp16-2*:: EFF-1 plasmid cut with the same enzymes to

replace the original heat shock promoter. To produce *eff-2 (hy51)* mutant worms with CRISPR,

the following DNA constructs were generated:

1. pBG115 plasmid encoding single guide targeting *eff-2* to insert *mNeonGreen* was

generated by cloning *eff-2* targeting sequence into the CAS9 plasmid pDD162 with BG84

and BG85 primers.

2. BG123 conversion oligonucleotide encoding wt *pha-1* fragment was PAGE purified.

3. PCR amplicon encoding for the mNeonGreen and unc-54 3'UTR flanked by homology

arms to the *eff-2* gene, generated by amplification of mNeonGreen from plasmid X, with

BG129 and BG130 primers containing a 50 bp homology arms to *eff-2*.

Sequences of primers and oligonucleotides mentioned above are found in Table S3.

**Transgenic animals**

For standard extrachromosomal transgenes, germline transformation was performed using

standard protocols [67]. Transgenic lines were kept as extrachromosomal arrays and

maintained by following the expression from either *myo-3p*::mCherry, *myo-2p*::GFP, or *mec-*

*4p*::dsRed, *odr-1p*::dsRed plasmids that were co-injected as transformation markers. *myo-*

*3p::*mCherry encodes mCherry expression in body wall muscle and vulva muscle cells. *myo-*

*2p*::GFP encodes GFP expression specifically expressed in pharyngeal muscles. *pmec-*

*4*::dsRed encodes dsRed expression in six touch receptor neurons. *odr-1p*::dsRed encodes

dsRed expression in two odor sensory neurons. Plasmids mentioned above and strains

containing these arrays are listed in Table S3. Transgenic lines with extrachromosomal arrays

were generated as follows:

• BP2126: *drh-1(tm1329);eff-1(hy21)* injected with 20ng/µl *pmyo-2::GFP* as
transformation marker and 0.1ng/µl *pdes-2::AFF-1*.

• BP2131-3: *drh-1(tm1329);eff-1(hy21)* injected 10ng/µl *pmec-4::dsRed* and 10ng/µl
*podr-1::dsRed* as transformation markers and to label sensory neurons, 1ng/µl *pmec-
4::EFF-1* and 1ng/µl *podr-1::EFF-1*.

• BP2137: *drh-1(tm1329)* injected 10ng/µl *myo-3p::mCherry* as transformation marker
and to label Body Wall Muscle cells, and 1ng/µl *myo-3p::EFF-1*.

*eff-2(hy51)* allele was generated by CRISPR/Cas9 [65,66] insertion of mNeonGreen into the
first exon of *eff-2* gene. *pha-1* was used as a conversion marker. *pha-1(e2123)* temperature
sensitive worms were injected with 50 ng/µl pBG115 targeting *eff-2* and containing CAS9, 50
ng/µl pJW1285 sgRNA plasmid against *pha-1* that contains the CAS9, 20 ng/µl BG123
conversion oligonucleotide encoding wt *pha-1* fragment and 20 ng/µl amplicon encoding for
the mNeonGreen and *unc-54* 3'UTR flanked by homology arms to the *eff-2* gene. Worms that
survived development at the restrictive temperature (25˚C) were screened and sequenced for
mutations in the *eff-2* gene. *hy51* is the result of an imprecise partial inverted insertion of the
PCR fragment into the *eff-2* locus. One bp was deleted at position +31 and 488bp of the unc-
54 3'UTR and mNeonGreen gene were inserted into *eff-2* coding region. The product of the
inverted insertion is a non-functional fluorescent protein that causes a frame shift in *eff-2* and
terminates its translation at Amino Acid #18. The *eff-2* targeting sequence was added to
primers (see Table S3) and inserted to the linearized vector using restriction-free cloning
technique.

**Live imaging of worms**
For imaging of viral infection and BWM fusion in *C. elegans*, worms were analyzed by
Nomarski optics and fluorescence microscopy using Nikon eclipse Ti inverted microscope with
Yokogawa CSU-X1 spinning disk confocal (SDC) as described previously [26,27]. Briefly,
animals were anesthetized in 0.01-0.05% tetramisole in M9 solution for 20-30 min and then
picked with an eyelash attached to a toothpick and transferred to a 5 µl droplet of M9 solution
placed on 3% agar slide. Images were acquired with Metamorph software, when using the
spinning disk confocal. Z-stacks were taken with Plan Fluor 40x NA=1.3 or Apochromat 60x
NA=1.4 objectives. Excitation of GFP was achieved with 488 nm wavelength laser (2-8%
intensity, 100 ms exposure time). mCherry was excited with 561 nm wavelength (15-20%
intensity, 100 ms exposure time). ~0.5 µm z-steps were recorded with iXon3 EMCCD camera
(Andor). Multidimensional data were reconstructed as maximum intensity projections using Fiji
software (NIH ImageJ). For live imaging of worms with fused BWM phenotypes, plates with

worms were placed on the stage of Zeiss stereo Discovery V8 stereo microscope. Images and

movies were captured at x8 magnification with additional magnification from PlanApo S 2.3X

objective and a Hamamatsu ORCA-ER camera controlled by micromanager software

(https://micro-manager.org). Figures were prepared using Fiji, Adobe Photoshop CS5 and

GraphPad Prism 8.

**DNA transformation and viral infection by microinjection**

Microinjections were performed as described [7,34] with some modifications. Shortly, late L4

or young adult worms were placed into droplet of halocarbon oil on 3% agarose pads. Pulled

capillary needles were secured onto a Nikon DIAPHOT 300 microscope equipped with a

micromanipulator and regulated pressure source (Narishige). For DNA transformation,

needles were loaded with 0.8 µl with DNA solution containing TE buffer, the target construct

DNA, a co-injection marker DNA and the required amount of with pKSI-1 (empty vector) DNA

to reach a total concentration of 100 ng/µl DNA. For experiments with viral infection, needles

were loaded 0.8 µl DMEM (mock-infections) or pseudo viruses in DMEM+5%FBS. Infection

doses of VSV are based on VSV titration on BHK cells and used 10nL volume as a single

microinjection dose [7]. Agar pads with worms were placed on stage of Nikon DIAPHOT 300

microscope. Worms were observed under x40 objective. For DNA transformation, animals

were injected into the gonad and immediately placed into droplet of M9 buffer placed on NGM

plates seeded with OP50-1 *E. coli*. For experiments with viral infection worms were injected

into pseudocoelom -behind the terminal bulb of the pharynx and were immediately placed into

droplet of M9 buffer placed on NGM plates with 50 µg/mL FUdR (Sigma), seeded with OP50-

1 *E. coli*. Unless otherwise stated, animals were maintained at 25°C until scoring of infection.

**Scoring viral infection**

Scoring viral infection using fluorescence assay was performed as described [7,34] with some

modifications. Briefly, 48-72 hours post injection worms were processed for live imaging as

described above. Animals that were unresponsive to prodding by a platinum wire worm pick

were considered dead and were removed from the experiment. Animals that crawled off the

plate or were lost during the experiment were censored. Worms with ≥1 GFP (+) cells were

considered as infected worms, while not injected or DMEM injected worms served as a

negative controls. For different experiments, either the number of GFP(+) cells/animals or the

type of infected tissues were observed and quantified.

**Temperature shift experiments**

Temperature sensitive *eff-1(hy21ts)* mutant worms were synchronized by hypochlorite
treatment of adult worms. The obtained eggs were left for overnight L1 hatching on NGM
plates without food at 20° C. L1 animals were then transferred to NGM plates with OP50
bacteria at 15° C or 25° C incubators until reaching the desired developmental stage. The
plates were either downshifted from 25° C to 15° C, or left at 25° C or at 15° C throughout the
experiment. The developmental stages were determined by analyzing gonadal size and
structure using Nomarski optics. Finally, worms were injected with VSVΔG-AFF-1, maintained
and imaged as described above.
565

**Counting overlapping cells**

To test if EFF-1 expressing BWMs were infected by VSVΔG-AFF-1, we utilized worms with an
extrachromosomal array expressing EFF-1 under the *myo-3* promoter and a plasmid
containing *myo-3p*::mCherry. We found that most mCherry (+) worms had a wt-like phenotype,
but a small fraction of these worms were Unc+Dpy. Importantly, all sibling mCherry(-) worms
were wt-like (Table S1). Hence, for the experiment we utilized a fluorescence
stereomicroscope and chose late L4/ young adult, wt-like and Unc+Dpy animals that had
continuous *myo-3p::mCherry*(+) BWMs (which could express EFF-1 and fuse with each other).
Next, these worms were injected with VSVΔG-AFF-1 or VSVΔG-G. 48-72h later, Z-stack
images of the worms were obtained with SDC microscope (as described in live imaging
section). Worms with ≥1 infected BWM were selected. For each worm, we counted the number
of BWMs that were mCherry(+) only ( express EFF-1), GFP(+) only (infected by virus) and
overlapping- with both mCherry(+) and GFP(+) (express EFF-1 and infected).
579

**Counting number of nuclei per BWM**

To test if ectopic EFF-1 expression in BWMs produces multinucleated BWMs, wt-like and
Unc+Dpy animals that had continuous *myo-3p::mCherry*(+) BWMs (which could express EFF-
1 and fuse with each other) were imaged by SDC microscope as described above. In one set
of experiments we imaged L2 worms and counted number of nuclei per *myo-3p::mCherry* cell.
In a second set of experiments, we imaged adult worms with BWMs infected by either VSVΔG-
AFF-1 or VSVΔG-G and counted number of nuclei per infected (GFP(+)) BWM cell. For both
sets of experiments, in worms with at least one multinucleated BWM we monitored all
multinucleated BWMs and calculated the average number of nuclei/BWM for each worm.
Worms in which multinucleated cells were not observed, were considered as having 1
nucleus/BWM.
591

**Counting phenotypes in worms expressing EFF-1 in BWMs**

To find whether ectopic expression of EFF-1 in BWM cells produces any special phenotypes,

we used BP2137 *drh-1(tm1329)IV*; hyEx375[*myo-3p*::EFF-1, *myo-3p*::mCherry, KS

bluescript]. We first isolated 4 wt-like *myo-3p::mCherry* L4 hermaphrodite worms, one worm

per NGM plate+Op50.  These worms were led to lay eggs and were transferred to a fresh

plate 1-2 times per day during 5 days. Progeny including eggs, larvae and adults were divided

into two groups, namely *myo-3p::mCherry*(+) and *myo-3p::mCherry*(-). We counted the total

number of progeny with a certain phenotype (e.g. wt-like worms, adult Unc+Dpy worms, larval

arrested Unc+Dpy worms and unhatched eggs) for each of the 4 mothers. Finally, we

calculated the average number of worms with certain phenotype ± SEM and the fraction ± SE

of worms with a certain phenotype in each group.

**Cell culture and preparation of pseudoviruses**

Baby Hamster Kidney cells (BHK), BHK-21(ATCC) were cultured in Dulbecco's Modified

Eagles Medium (DMEM) and recombinant viruses were prepared as described [6] with some

modifications. Briefly, BHK cells were grown to 70% confluence on 10 cm plates and then

transfected using Fugene HD + OptiMEM at ratio 1:4 (Fugene:DNA), with plasmids encoding

pOA20 (pCAGGS::*aff-1*::FLAG, 2µg/ml final concentration) [6] or pOA28 (pCAGGS::VSV-G

Indiana, 1µg/ml final concentration)[33]. Following 24 h incubation at 37°C in 5% $CO_2$, cells

were infected with VSVG-complemented VSV$\Delta$G recombinant virus (VSV$\Delta$G-G) at a

multiplicity of infection (MOI) of 5, for 1 hour at 37°C in a 5% $CO_2$ incubator in serum free

DMEM. Virus infected cells were washed 3-6 times with PBS (+ $Ca^{++}$ & $Mg^{++}$) to remove

unabsorbed VSV$\Delta$G-G virus. Following a 24 h incubation period at 37°C, the supernatant

containing the VSV$\Delta$G-G, or VSV$\Delta$G-AFF-1 pseudoviruses were harvested without scraping

the cells and centrifuged at 600 g for 10 min at 4°C to clear cell debris. VSV$\Delta$G-AFF-1 Virions

were filtered through 0.22 µm filter unit and then double concentrated. First, by pelleting at

100,000 g through a 20% sucrose cushion, and resuspension in 10%FBS DMEM, second, by

pelleting at 100,000 g through a 10% sucrose cushion, and final resuspension in 45µl DMEM.

Finally, VSV$\Delta$G-AFF-1 was incubated with anti-VSV-G antibody mAb diluted 1:1000 to inhibit

infection due to residual presence of VSV-G. The effective blocking of VSV-G was confirmed

by tittering pseudoviruses in BHK cells and by injection of VSV$\Delta$G-G incubated with anti-VSV-

G into worms (Figure S4).

**Titering VSV pseudotyped viruses**

Titering VSV pseudotyped viruses was performed as described [6] with some modifications.

Briefly, $5x10^3$ BHK cells were plated into each well of a 96 well tissue culture plate (NUNC,

cat# 167008). To determine the titer of VSV$\Delta$G-AFF-1, BHK cells were initially transfected with

2 µg/ml pOA20 (pCAGGS::AFF-1::FLAG). Cells transfected with empty vector served as

control. Eight serial x2 dilutions of the virus were performed and added to cells. After 18-24

hours of incubation, GFP expressing cells were counted in at least three dilutions using x20

objective of Zeiss Axiovert 200M fluorescence microscope. Inoculation was performed in the

presence of anti-VSV-G antibody mAb diluted 1:1000 to inhibit infection due to residual

presence of VSV-G.

**Statistical tests**

The specific tests used are described in the figure captions and the results section. The graphs

show mean ± SEM unless noted otherwise. For each experiment at least two biological

replicates were performed and the number of animals per experiment is stated in the figure

legends.

**References**

1.  Lichty, B.D., Power, A.T., Stojdl, D.F., and Bell, J.C. (2004). Vesicular stomatitis virus: Re-inventing the bullet. Trends Mol. Med. *10*, 210–216.

2.  van den Pol, A.N., Ozduman, K., Wollmann, G., Ho, W.S.C., Simon, I., Yao, Y., Rose, J.K., and Ghosh, P. (2009). Viral strategies for studying the brain, including a replication-restricted self-amplifying delta-G vesicular stomatis virus that rapidly expresses transgenes in brain and can generate a multicolor Golgi-like expression. J. Comp. Neurol. *516*, 456–481.

3.  Hastie, E., Cataldi, M., Marriott, I., and Grdzelishvili, V.Z. (2013). Understanding and altering cell tropism of vesicular stomatitis virus. Virus Res. *176*, 16–32.

4.  Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T., Erichsen, S., Schiergens, T.S., Herrler, G., Wu, N.-H., Nitsche, A., *et al.* (2020). SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell *181*, 271–280.

5.  Halperin, S.A., Das, R., Onorato, M.T., Liu, K., Martin, J., Grant-Klein, R.J., Nichols, R., Coller, B.-A., Helmond, F.A., and Simon, J.K. (2019). Immunogenicity, Lot Consistency, and Extended Safety of rVSVΔG-ZEBOV-GP Vaccine: A Phase 3 Randomized, Double-Blind, Placebo-Controlled Study in Healthy Adults. J. Infect. Dis. *220*, 1127–1135.

6.  Avinoam, O., Fridman, K., Valansi, C., Abutbul, I., Zeev-Ben-Mordehai, T., Maurer, U.E., Sapir, A., Danino, D., Grünewald, K., White, J.M., *et al.* (2011). Conserved eukaryotic fusogens can fuse viral envelopes to cells. Science. *332*, 589–592.

7.  Gammon, D.B., Ishidate, T., Li, L., Gu, W., Silverman, N., and Mello, C.C. (2017). The Antiviral RNA Interference Response Provides Resistance to Lethal Arbovirus Infection and Vertical Transmission in Caenorhabditis elegans. Curr. Biol. *27*, 795–806.

8.  Presley, J.F., Smith, C., Hirschberg, K., Miller, C., Cole, N.B., Zaal, K.J.M., and Lippincott-Schwartz, J. (1998). Golgi membrane dynamics. Mol. Biol. Cell *9*, 1617–1626.

9.  Ogino, T., and Green, T.J. (2019). RNA Synthesis and Capping by Non-segmented Negative Strand RNA Viral Polymerases: Lessons From a Prototypic Virus. Front. Microbiol. *10*, 1490.

10. Finkelshtein, D., Werman, A., Novick, D., Barak, S., and Rubinstein, M. (2013). LDL receptor and its family members serve as the cellular receptors for vesicular stomatitis virus. Proc. Natl. Acad. Sci. U. S. A. *110*, 7306–7311.

11. Bergman, I., Whitaker-Dowling, P., Gao, Y., Griffin, J.A., and Watkins, S.C. (2003). Vesicular stomatitis virus expressing a chimeric Sindbis glycoprotein containing an Fc antibody binding domain targets to Her2/neu overexpressing breast cancer cells. Virology *316*, 337–347.

12. Schnell, M.J., Johnson, J.E., Buonocore, L., and Rose, J.K. (1997). Construction of a novel virus that targets HIV-1-infected cells and controls HIV-1 infection. Cell *90*, 849–857.

13. Gao, Y., Whitaker-Dowling, P., Griffin, J.A., Barmada, M.A., and Bergman, I. (2009). Recombinant vesicular stomatitis virus targeted to Her2/neu combined with anti-CTLA4 antibody eliminates implanted mammary tumors. Cancer Gene Ther. *16*, 44–52.

14. Mohler, W.A., Shemer, G., Del Campo, J.J., Valansi, C., Opoku-Serebuoh, E., Scranton, V., Assaf, N., White, J.G., and Podbilewicz, B. (2002). The type 1 membrane protein EFF-1 is essential for development cell fusion. Dev. Cell *2*, 355–362.

15. Shemer, G., and Podbilewicz, B. (2002). LIN-39/Hox triggers cell division and represses EFF-1/fusogen-dependent vulval cell fusion. Genes Dev.

16. Podbilewicz, B. (2006). Cell fusion. WormBook *93*, 71–79.

17. Rasmussen, J.P., English, K., Tenlen, J.R., and Priess, J.R. (2008). Notch Signaling and Morphogenesis of Single-Cell Tubes in the C. elegans Digestive Tract. Dev. Cell *14*, 559–569.

695    18.    Sapir, A., Choi, J., Leikina, E., Avinoam, O., Valansi, C., Chernomordik, L. V., Newman,
696           A.P., and Podbilewicz, B. (2007). AFF-1, a FOS-1-Regulated Fusogen, Mediates
697           Fusion of the Anchor Cell in C. elegans. Dev. Cell *12*, 683–698.

698    19.    Shemer, G., Suissa, M., Kolotuev, I., Nguyen, K.C.Q., Hall, D.H., and Podbilewicz, B.
699           (2004). EFF-1 is sufficient to initiate and execute tissue-specific cell fusion in C.
700           elegans. Curr. Biol. *14*, 1587–1591.

701    20.    Abdus-Saboor, I., Mancuso, V.P., Murray, J.I., Palozola, K., Norris, C., Hall, D.H.,
702           Howell, K., Huang, K., and Sundaram, M. V. (2011). Notch and Ras promote sequential
703           steps of excretory tube development in C. elegans. Development *138*, 3545–3555.

704    21.    Soulavie, F., Hall, D.H., and Sundaram, M. V. (2018). The AFF-1 exoplasmic fusogen
705           is required for endocytic scission and seamless tube elongation. Nat. Commun. *9*,
706           1741.

707    22.    Stone, C.E., Hall, D.H., and Sundaram, M. V. (2009). Lipocalin signaling controls
708           unicellular tube development in the Caenorhabditis elegans excretory system. Dev.
709           Biol. *329*, 201–211.

710    23.    Ghosh-Roy, A., Wu, Z., Goncharov, A., Jin, Y., and Chisholm, A.D. (2010). Calcium
711           and cyclic AMP promote axonal regeneration in Caenorhabditis elegans and require
712           DLK-1 kinase. J. Neurosci. *30*, 3175–3183.

713    24.    Neumann, B., Coakley, S., Giordano-Santini, R., Linton, C., Lee, E.S. eun., Nakagawa,
714           A., Xue, D., and Hilliard, M.A. (2015). EFF-1-mediated regenerative axonal fusion
715           requires components of the apoptotic pathway. Nature *517*, 219–222.

716    25.    Oren-Suissa, M., Hall, D.H., Treinin, M., Shemer, G., and Podbilewicz, B. (2010). The
717           fusogen EFF-I controls sculpting of mechanosensory dendrites. Science. *328*, 1285–
718           1288.

719    26.    Oren-Suissa, M., Gattegno, T., Kravtsov, V., and Podbilewicz, B. (2017). Extrinsic
720           repair of injured dendrites as a paradigm for regeneration by fusion in Caenorhabditis
721           elegans. Genetics *206*, 215–230.

722    27.    Kravtsov, V., Oren-Suissa, M., and Podbilewicz, B. (2017). The fusogen AFF-1 can
723           rejuvenate the regenerative potential of adult dendritic trees by self-fusion. Dev. *144*,
724           2364–2374.

725    28.    Harrison, S.C. (2015). Viral membrane fusion. Virology *479–480*, 498–507.

726    29.    Podbilewicz, B. (2014). Virus and Cell Fusion Mechanisms. Annu. Rev. Cell Dev. Biol.
727           *30*, 111–139.

728    30.    Pérez-Vargas, J., Krey, T., Valansi, C., Avinoam, O., Haouz, A., Jamin, M., Raveh-
729           Barak, H., Podbilewicz, B., and Rey, F.A. (2014). Structural basis of eukaryotic cell-cell
730           fusion. Cell *157*, 407–419.

731    31.    White, J.M. (2007). The First Family of Cell-Cell Fusion. Dev. Cell *12*, 683–698.

732    32.    Robison, C.S., and Whitt, M.A. (2000). The Membrane-Proximal Stem Region of
733           Vesicular Stomatitis Virus G Protein Confers Efficient Virus Assembly. J. Virol. *74*,
734           2239–2246.

735    33.    Whitt, M.A. (2010). Generation of VSV pseudotypes using recombinant ΔG-VSV for
736           studies on virus entry, identification of entry inhibitors, and immune responses to
737           vaccines. J. Virol. Methods *169*, 365–374.

738    34.    Martin, A., Rex, E., Ishidate, T., Lin, R., and Gammon, D. (2017). Infection of
739           Caenorhabditis elegans with Vesicular Stomatitis Virus via Microinjection. BIO-
740           PROTOCOL *7*, e2617.

741    35.    Quiroz, E., Moreno, N., Peralta, P.H., and Tesh, R.B. (1988). A human case of
742           encephalitis associated with vesicular stomatitis virus (Indiana serotype) infection. Am.
743           J. Trop. Med. Hyg. *39*, 312–314.

744    36.    Guerra-Varela, J., Baz-Martínez, M., Da Silva-Álvarez, S., Losada, A.P., Quiroga, M.I.,
745           Collado, M., Rivas, C., and Sánchez, L. (2018). Susceptibility of Zebrafish to Vesicular
746           Stomatitis Virus Infection. Zebrafish *15*, 124–132.

37.    Barrera, J.D.C., and Letchworth, G.J. (1996). Persistence of Vesicular Stomatitis Virus

New Jersey RNA in Convalescent Hamsters. Virology *219*, 453–464.

38.  Valansi, C., Moi, D., Leikina, E., Matveev, E., Graña, M., Chernomordik, L. V., Romero,
H., Aguilar, P.S., and Podbilewicz, B. (2017). Arabidopsis HAP2/GCS1 is a gamete
fusion protein homologous to somatic and viral fusogens. J. Cell Biol. *216*, 571–581.

39.  Yochem, J., Gu, T., and Han, M. (1998). A new marker for mosaic analysis in
Caenorhabditis elegans indicates a fusion between hyp6 and hyp7, two major
components of the hypodermis. Genetics *149*, 1323–1334.

40.  Podbilewicz, B., and White, J.G. (1994). Cell fusions in the developing epithelia of C.
elegans. Dev. Biol. *161*, 408–424.

41.  Fukushige, T., Brodigan, T.M., Schriefer, L.A., Waterston, R.H., and Krause, M. (2006).
Defining the transcriptional redundancy of early bodywall muscle development in C.
elegans: Evidence for a unified theory of animal muscle development. Genes Dev. *20*,
3395–3406.

42.  Kim, J., Löwe, T., and Hoppe, T. (2008). Protein quality control gets muscle into shape.
Trends Cell Biol. *18*, 264–272.

43.  Frumkin, A., Dror, S., Pokrzywa, W., Bar-Lavan, Y., Karady, I., Hoppe, T., and Ben-Zvi,
A. (2014). Challenging muscle homeostasis uncovers novel chaperone interactions in
Caenorhabditis elegans. Front. Mol. Biosci. *6*, 21.

44.  Dreier, B., Honegger, A., Hess, C., Nagy-Davidescu, G., Mittl, P.R.E., Grütter, M.G.,
Belousova, N., Mikheeva, G., Krasnykh, V., and Plückthun, A. (2013). Development of
a generic adenovirus delivery system based on structure-guided design of bispecific
trimeric DARPin adapters. Proc. Natl. Acad. Sci. U. S. A. *5*, E869–E877.

45.  Raposo, G., and Stoorvogel, W. (2013). Extracellular vesicles: Exosomes,
microvesicles, and friends. J. Cell Biol. *200*, 373–383.

46.  Sdrimas, K., and Kourembanas, S. (2014). MSC microvesicles for the treatment of lung
disease: A new paradigm for cell-free therapy. Antioxidants Redox Signal. *21*, 1905–
1915.

47.  Segev, N., Avinoam, O., and Podbilewicz, B. (2018). Fusogens. Curr. Biol. *28*, R378-
R380.

48.  Brukman, N.G., Uygur, B., Podbilewicz, B., and Chernomordik, L. V. (2019). How cells
fuse. J. Cell Biol. *218*, 1436–1451.

49.  Blond, J.-L., Lavillette, D., Cheynet, V., Bouton, O., Oriol, G., Chapel-Fernandes, S.,
Mandrand, B., Mallet, F., and Cosset, F.-L. (2000). An Envelope Glycoprotein of the
Human Endogenous Retrovirus HERV-W Is Expressed in the Human Placenta and
Fuses Cells Expressing the Type D Mammalian Retrovirus Receptor. J. Virol. *74*,
3321–3329.

50.  Sha, M., Lee, X., Li, X. ping, Veldman, G.M., Finnerty, H., Racie, L., LaVallie, E., Tang,
X.Y., Edouard, P., Howes, S., *et al.* (2000). Syncytin is a captive retroviral envelope
protein involved in human placental morphogenesis. Nature *403*, 785–789.

51.  Kokkinos, M.I., Murthi, P., Wafai, R., Thompson, E.W., and Newgreen, D.F. (2010).
Cadherins in the human placenta - Epithelial-mesenchymal transition (EMT) and
placental development. Placenta *31*, 747–755.

52.  von Besser, K., Frank, A.C., Johnson, M.A., and Preuss, D. (2006). Arabidopsis HAP2
(GCS1) is a sperm-specific gene required for pollen tube guidance and fertilization.
Development *133*, 4761–4769.

53.  Liu, C., Zhou, Y., Zhang, X., Zhang, J., Zhou, Z., Weng, J., Li, X., and Wang, Z. (2019).
Natural variation in the THICK TASSEL DWARF1 (TD1) gene in the regulation of maize
(Zea mays L.) ear-related traits. Breed. Sci. *69*, 323–331.

54.  Cole, E.S., Cassidy-Hanley, D., Fricke Pinello, J., Zeng, H., Hsueh, M., Kolbin, D.,
Ozzello, C., Giddings, T., Winey, M., and Clark, T.G. (2014). Function of the male-
gamete-specific fusion protein HAP2 in a seven-sexed ciliate. Curr. Biol. *24*, 2168–
2173.

55.  Millay, D.P., O'Rourke, J.R., Sutherland, L.B., Bezprozvannaya, S., Shelton, J.M.,

Bassel-Duby, R., and Olson, E.N. (2013). Myomaker is a membrane activator of
myoblast fusion and muscle formation. Nature *499*, 301–305.

56.    Gamage, D.G., Leikina, E., Quinn, M.E., Ratinov, A., Chernomordik, L. V., and Millay,
D.P. (2017). Insights into the localization and function of myomaker during myoblast
fusion. J. Biol. Chem. *292*, 17272–17289.

57.    Mitani, Y., Vagnozzi, R.J., and Millay, D.P. (2017). In vivo myomaker-mediated
heterologous fusion and nuclear reprogramming. FASEB J. *31*, 400–411.

58.    Bi, P., Ramirez-Martinez, A., Li, H., Cannavino, J., McAnally, J.R., Shelton, J.M.,
Sánchez-Ortiz, E., Bassel-Duby, R., and Olson, E.N. (2017). Control of muscle
formation by the fusogenic micropeptide myomixer. Science (80-. ). *356*, 323–327.

59.    Zhu, T., Liang, X., Wang, X.M., and Shen, K. (2017). Dynein and EFF-1 control dendrite
morphology by regulating the localization pattern of SAX-7 in epidermal cells. J. Cell
Sci. *130*, 4063–4071.

60.    Buechner, M., Hall, D.H., Bhatt, H., and Hedgecock, E.M. (1999). Cystic canal mutants
in Caenorhabditis elegans are defective in the apical membrane domain of the renal
(excretory) cell. Dev. Biol. *214*, 227–241.

61.    Nelson, F.K., and Riddle, D.L. (1984). Functional study of the Caenorhabditis elegans
secretory-excretory system using laser microsurgery. J. Exp. Zool. *231*, 45–56.

62.    Gunage, R.D., Dhanyasi, N., Reichert, H., and VijayRaghavan, K. (2017). Drosophila
adult muscle development and regeneration. Semin. Cell Dev. Biol. *72*, 56–66.

63.    Sampath, S.C., Sampath, S.C., and Millay, D.P. (2018). Myoblast fusion confusion: The
resolution begins. Skelet. Muscle *8*, 3.

64.    Gjorgjieva, J., Biron, D., and Haspel, G. (2014). Neurobiology of caenorhabditis
elegans locomotion: Where do we stand? Bioscience, 476–486.

65.    Brenner, S. (1974). The genetics of Caenorabditis elegans. Genetics *77*, 71–94.

66.    Sulston, J., and Hodgkin, J. (1988). Methods in the nematode C. elegans. In Cold
Spring Harbor Laboratory Press, pp. 587–606.

67.    Mello, C., and Fire, A. (1995). DNA Transformation. Methods Cell Biol. *8*, 2159–2165.

68.    Takada, A., Robison, C., Goto, H., Sanchez, A., Murti, K.G., Whitt, M.A., and Kawaoka,
Y. (1997). A system for functional analysis of Ebola virus glycoprotein. Proc Natl Acad
Sci U S A *94*, 14764-14769.

69.    Stoker, M., and Macpherson, I. (1964). Syrian Hamster Fibroblast Cell Line Bhk21 and
Its Derivatives. Nature *203*, 1355-1357.

70.    Diaz-Balzac, C.A., Rahman, M., Lazaro-Pena, M.I., Martin Hernandez, L.A., Salzberg,
Y., Aguirre-Chen, C., Kaprielian, Z., and Bulow, H.E. (2016). Muscle- and Skin-Derived
Cues Jointly Orchestrate Patterning of Somatosensory Dendrites. Curr Biol *26*, 2397.

71.    Yang, J., Sun, B., Huang, H., Jiang, Y., Diao, L., Chen, B., Xu, C., Wang, X., Liu, J.,
Jiang, W., et al. (2014). High-efficiency scarless genetic modification in Escherichia coli
by using lambda red recombination and I-SceI cleavage. Appl Environ Microbiol *80*,
3826-3834.

72.    Frokjaer-Jensen, C., Davis, M.W., Hopkins, C.E., Newman, B.J., Thummel, J.M.,
Olesen, S.P., Grunnet, M., and Jorgensen, E.M. (2008). Single-copy insertion of
transgenes in Caenorhabditis elegans. Nat Genet *40*, 1375-1383.

73.    Dickinson, D.J., Ward, J.D., Reiner, D.J., and Goldstein, B. (2013). Engineering the
Caenorhabditis elegans genome using Cas9-triggered homologous recombination. Nat
Methods *10*, 1028-1034.

74.    Ward, J.D. (2015). Rapid and precise engineering of the Caenorhabditis elegans
genome with lethal mutation co-conversion and inactivation of NHEJ repair. Genetics
*199*, 363-377.

75.    Edelstein, A., Amodaj, N., Hoover, K., Vale, R., and Stuurman, N. (2010). Computer
control of microscopes using microManager. Curr Protoc Mol Biol *Chapter 14*, Unit14
20.

**Abbreviations**

| | |
|---|---|
| AFF-1 | Anchor cell Fusion Failure |
| BHK | Baby Hamster Kidney cells |
| BWM | Body Wall Muscle |
| Dpy | Dumpy phenotype |
| EC | Excretory Canal cell |
| EFF-1 | Epithelial Fusion Failure 1 |
| EFF-2 | Epithelial Fusion Failure 2 |
| FUSEXINS | FUSion proteins essential for sexual reproduction and Exoplasmic merger of plasma membranes |
| HYP | Hypodermis (epidermis) |
| SM | Somatointestinal Muscle |
| ts | temperature sensitive |
| UM | Uterine Muscle |
| Unc | Uncoordinated phenotype |
| VSV | Vesicular Stomatitis Virus |
| VSV-G | VSV-glycoprotein G |
| VSVΔG | pseudoviruses in which the glycoprotein G gene was deleted |
| VSVΔG-AFF-1 | pseudovirus coated with AFF-1 |
| VSVΔG-G | pseudovirus coated with G glycoprotein |

**Acknowledgements**

We thank Don Gammon for providing the *drh-1(-)* worms, Massimo Hilliard for providing *mec-4p*::mCherry, *mec-4p*::EFF-1, *odr-1p*::dsRed and *odr-1p*::EFF-1 plasmids and the Caenorhabditis Genetics Center for nematode strains. We thank Andy Fire for suggesting studying syncytial BWMs and their potential phenotypes. We also thank Sharon Inberg, Yael Iosilevskii, Rosina Giordano-Santini, Dan Cassel and Sivan Korenblit for helpful discussions and for critically reading the manuscript.

**Author contributions**

A.M. and B.P. conceived the project. A.M. performed all experiments unless otherwise specified. X.L. generated *myo-3p*::EFF-1 Plasmid and BP2171-3 nematode strains and performed imaging of worms in Figure 5. E.M. improved viral preparation protocol. B.G. generated EFF-2 CRISPR plasmid and O.K. produced *eff-2(hy51)* worms. A.M. and B.P analyzed data and wrote the manuscript, with input from all authors.

 **Online supplemental material**

**893  Figure S1: VSVΔG-AFF-1 and VSVΔG-G infect glia and neuron cells**
**(A-B)** SDC microscope Z-stack projections of *drh-1* and *drh-1; eff-1(-)* worms injected in the
pseudocoelom with 2300-4700 IU VSVΔG-G (white pins) and their heads imaged 48 h later.
**(C)** Fraction of GFP(+) worms infected with VSVΔG-G in neuron/glia cells in the specified
background. Animals were injected with VSVΔG-G and analyzed as in (A-B).
**(D-E)** SDC microscope Z-stack projections of *C. elegans* cells infected with 33-240 IU VSVΔG-
AFF-1 (red pins). *drh-1* or *drh-1; eff-1(-)* worms were injected  in the pseudocoelom and
imaged 48 h later.
**(F)** Fraction of GFP(+) worms infected with VSVΔG-AFF-1 in neuron/glia cells. Animals were
analyzed as in (D-E). For (A-B) and (D-E) arrowheads point to hypodermal nuclei (HYP) and
arrows point to indicated infected cell. CEPsh- Cephalic Sheath (glia), G- glia, N- neuron.
Scale bar, 50 µm. In (A, B and E) yellow dashed lines outline worm's anterior body (head). In
(C) and (F), bars represent average ± SEM. ***P<0.001 (Student's T- test). For VSVΔG-AFF-
1: n=46,15 and 10 for *drh-1,drh-1;eff-1(-)* and *drh-1;aff-1(-)* respectively, and for VSVΔG-G:
n= 56, 39 and 15 for *drh-1*, *drh-1;eff-1(-)* and *drh-1;aff-1(-)* respectively).

**Figure S2: VSVΔG-AFF-1 infects *eff-2(-)* worms**

**(A-B)** SDC microscope Z-stack projections of *drh-1* or *drh-1;eff-2(-)* worms injected with
VSVΔG-AFF-1 (35-67 IU) and imaged 48 h later.
**(C)** Fraction of GFP(+) worms infected with VSVΔG-AFF-1. Animals were injected with
VSVΔG-AFF-1 and analyzed as in (A-B). Arrowheads point to hypodermal (HYP) nuclei. Scale
bar, 100 μm. In (C) bars represent average ± SEM. n=23 and 20 for *drh-1* and *drh-1;eff-2(-)*
respectively. For all types of infected cells, there is no significant difference between % GFP(+)
worms from *drh-1* and *drh-1;eff-2 (-)* backgrounds with two-tailed Student's t-test (p<0.05).

**Figure S3. VSVΔG-AFF-1 does not infect PVD and other sensory neurons ectopically expressing AFF-1/EFF-1**

**(A-C)** SDC microscope Z-stack projections of animals infected with 82-103 IU VSVΔG-AFF-1 (red pins). (For genotypes and quantitation see Table S2). Scale bar, 100 μm.
(**A**) Young adult expressing mCherry in PVD.
(**B**) *eff-1(ts)* adult expressing AFF-1 in PVD.
(**C**) *eff-1(ts)* adult expressing EFF-1 and dsRed in 12 sensory neurons.
See also Table S2 and Movie S2.

**Figure S4. Anti-VSV-G antibody blocks VSVΔG-G infection in living worms**
**(A-B)** SDC microscope Z-stack projections of *drh-1* worms injected with 4700 IU VSVΔG-G that was either not preincubated (-αG), or preincubated with αVSV-G antibody (+αG) and imaged 48 h later. Scale bar, 50 μm.
**(C)** Fraction of worms that are infected or not infected with VSVΔG-G or VSVΔG-G +αG. Animals were treated as in (A-B). n=18 and 17 animals for VSVΔG-G and VSVΔG-G+αG respectively. Fisher exact test. p***<0.001.

 **Supplementary tables**

941 **Table S1. Phenotypes of worms with EFF-1 expressed in BWMs**

| Genotype | *drh-1(-); hyEx375[myo-3p*::mCherry*, myo-3p*::EFF-1*]* | | |
|---|---|---|---|
| Phenotype/ group | *myo-3p*::mCherry(+) | *myo-3p*::mCherry(-) | P-value |
| wt-like animals | 110 ± 12 (81±3%) | 60 ± 10 (87±6%) | 0.39258 |
| Adult Unc+Dpy | 3 ± 1 (2±1%) | 0 (0%) | 0.05316 |
| Larval-arrested Unc+Dpy | 15 ± 2 (11±1%) | 0 (0%) | 0.00029*** |
| Unhatched eggs | 7 ± 2 (5±1%) | 8 ± 3 (13±6%) | 0.23501 |

944 Phenotype count of total progeny from 4 hermaphrodites with extrachromosomal array
945 containing *myo-3p*::EFF-1 and *myo-3p*::mCherry. Data presented as average of the progenies
946 from 4 mothers ±SEM and the percentage out of total relevant group (e.g. *myo-3p*::mCherry(+)
947 or *myo-3p*::mCherry(-)). P-value calculated for certain phenotype fraction (%) in *myo-*
948 *3p::*mCherry(+) vs *myo-3p*::mCherry(-) groups. Two tailed Student's t-test. $p < 0.01$***.
949

950 **Table S2. AFF-1/EFF-1 expression in sensory neurons does not induce their infection**

951 **with VSV∆G-AFF-1**

| Strain | Genotype | Description | total # of worms | # VSV∆G-AFF-1-infected worms | # worms with GFP(+) cells overlapping with EFF-1/AFF-1 expressing cells |
|---|---|---|---|---|---|
| BP2126 | *drh-1(tm1329); eff-1(hy21); hyEx373 [pmyo-2::GFP, pdes-2::AFF-1(pME4), KS]* | AFF-1 expressed in PVD (Oren-Suissa et al 2017) , in *eff-1(ts)* background | 18 | 6 | 0 |
| BP2131 | *drh-1(tm1329); eff-1(hy21); hyEx374 [pmec-4::EFF-1, pmec-4::dsRed, podr-1::EFF-1,podr-1::dsRed, KS] line1* | EFF-1 expressed in 12 sensory neurons (kind gift from M. Hilliard lab), in *eff-1(ts)* background | 28 | 15 | 0 |
| BP2132 | *drh-1(tm1329); eff-1(hy21); hyEx374 [pmec-4::EFF-1, pmec-4::dsRed, podr-1::EFF-1,podr-1::dsRed, KS] line2* | EFF-1 expressed in 12 sensory neurons, in *eff-1(ts)* background | 22 | 7 | 0 |
| BP2133 | *drh-1(tm1329); eff-1(hy21); hyEx374 [pmec-4::EFF-1, pmec-4::dsRed, podr-1::EFF-1,podr-1::dsRed, KS] line3* | EFF-1 expressed in 12 sensory neurons, in *eff-1(ts)* background | 35 | 13 | 0 |

953
954

## Table S3. Reagents, strains, oligos, plasmids and software

| REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|---|---|---|
| **Antibodies** | | |
| Anti-VSV-G [8G5F11] Antibody | Kerafast | Cat#8G5F-11 |
| **Bacterial and Virus Strains and tissue culture cells** | | |
| Recombinant Vesicular Stomatitis Virus VSVΔG | [68] | VSVΔG |
| *E. coli*: OP50-1 | Caenorhabditis Genetics Center | OP50-1 |
| BHK-21(ATCC) | [69] | BHK-21(ATCC) |
| **Chemicals, Peptides, and Recombinant Proteins** | | |
| FUdR | Sigma | Cat#F0503-1G |
| Tetramisole | Sigma | Cat#L9756-5G |
| Dulbecco's Modified Eagles Medium (DMEM) High glucose, Gibco™ | Thermo Scientific | Cat#41965039 |
| Fugene HD | Promega | Cat#E2311 |
| OptiMEM | Gibco | Cat#31985070 |
| Dulbecco's Phosphate Buffered Saline (DPBS) With Calcium and Magnesium | Biological Industries | Cat#02-020-1A |
| Certified Fetal Bovine Serum (FBS), Heat Inactivated | Biological Industries | Cat#04-121-1A |
| ***C. elegans* strains** | | |
| *drh-1(tm1329) IV* | [7] | *drh-1 (tm1329)* |
| *eff-1(ok1021)II* | [14] | BP347 |
| *drh-1(tm1329) IV; eff-1(ok1021)II* | This paper | BP2122 |
| Strain *eff-1(hy21)II* | [14, 19] | BP75 |
| *drh-1(tm1329) IV; eff-1(hy21)II* | This paper | BP2123 |
| *drh-1(tm1329) IV; aff-1(tm2214)II* | This paper | BP2124 |
| *dzIS53[F49H124p::mCherry] Unknown chromosome* | [70] | EB2110 |
| *drh-1(tm1329) IV; dzIS53[F49H124p::mCherry]* | This paper | BP2121 |
| *trIs10 [myo-3p::MB::YFP + myo-2p::YFP + ceh-23::HcRed + unc-25::DsRed + unc-129nsp::CFP]* | Caenorhabditis Genetics Center | RP1 |
| *eff-2 (hy51) II; mcIs46[dlg-1::RFP;unc-119(+)]III* | This paper | BP2104 |
| *drh-1(tm1329) IV ; eff-2(hy51) II* | This paper | BP2125 |
| *drh-1(tm1329)IV; hyEx375 [myo-3p::EFF-1, myo-3p::mCherry, KS bluescript]* | This paper | BP2137 |
| *trIs10 [myo-3p::MB::YFP + myo-2p::YFP + ceh-23::HcRed + unc-25::DsRed + unc-129nsp::CFP];hyEx375 [myo-3p::EFF-1, myo-3p::mCherry, KS bluescript]* | This paper | BP2171-3 |
| *drh-1(tm1329) IV; eff-1(hy21) II; hyEx373 [myo-2p::GFP, des-2p::AFF-1, KS bluescript]* | This paper | BP2126 |
| *drh-1(tm1329) IV; eff-1(hy21) II; hyEx374 [mec-4p::EFF-1, pmec-4::dsRed, podr-1::EFF-1, odr-1p::dsRed, KS bluescript]* | This paper | BP2131-3 |
| **Oligonucleotides** | | |
| Primer: *Cloning myo-3p from myo3p::mCherry Plasmid with SalI restriction enzyme: ACGCGTCGAC AGTGATTATAGTCTCTGTTTTC* | This paper | LXH60 |
| Primer: *Cloning myo-3p from myo3p::mCherry Plasmid with NheI restriction enzyme: CTAGCTAGCCATTTCTAGATGGATCTAGTG* | This paper | LXH61 |
| Oligonucleotide: *Conversion oligonucleotide encoding wt pha-1 repair fragment for CRISPR: caaaatacgaatcgaagactcaaaaagagtatgctgtatgattacagatgttcatcaagttattcataaatcattgatag* | This paper | BG123 |
| Primer: *Addition of homology arms to eff-2 around mNeonGreen for CRISPR: ttcctgaagtttcatcaaattcgaagctggcGcgtattttcagaagcATGGTTTCCAAGGGAGAAGAGG* | This paper | BG129 |
| Primer: *Addition of homology arms to eff-2 around mNeonGreen for CRISPR: agattctaatgtgtcagatagtcttgagatcgCtgCgagagaatatactaGACTAGTAGGAAACAGTTATGTTTGG* | This paper | BG130 |
| Primer: *Production of sgRNA plasmid targeting eff-2 at locus AGAAGCAGAATAAACTGGGGAGG for CRISPR: AGAAGCAGAATAAACTGGGGGTTTTAGAGCTAGAAATAGC* | This paper | BG84 |
| Primer: *Production of sgRNA plasmid targeting eff-2 at locus AGAAGCAGAATAAACTGGGGAGG for CRISPR: CCCCAGTTTATTCTGCTTCTAAACATTTAGATTTGCAATTC* | This paper | BG85 |

| Recombinant DNA | | |
|---|---|---|
| Plasmid pCAGGS-Gind: *pCAGGS::VSV-G Indiana* | [68] | pCAGGS-Gind |
| Plasmid pOA20: *pCAGGS::aff-1::FLAG* | [6] | pOA20 |
| Plasmid pKSI-1: pBluescript empty vector | [71] | Addgene plasmid #51725 |
| Plasmid pXH20: *myo-3p::EFF-1* | This paper | pXH20 |
| Plasmid pME1: *hsp16.2::EFF-1* | [19] | pME1 |
| Plasmid pCFJ104: *myo-3p::mCherry* | [72] | Addgene plasmid #19328 |
| Plasmid pBG115: *CeU6p::eff-2_sgRNA for CRISPR* | This paper | pBG115 |
| Plasmid pDD162: *eft-3p::Cas9 + Empty sgRNA for CRISPR* | [73] | Addgene plasmid #47549 |
| Plasmid pJW1285: *eft-3p::Cas-9-SV40 NLSNLS::tbb-2 3'UTR + CeU6p::pha-1_sgRNA for CRISPR* | [74] | Addgene plasmid #61252 |
| Plasmid pNG1: *dlg-1p::Lifeact::mNeonGreen* | Kind gift of A. Hajnal | pNG1 |
| PCR amplicon: *mNeonGreen flanked by homology arms to the eff-2 gene for CRISPR* | This paper | N/A |
| Plasmid : *odr-1p::dsRed* | [24] | odr-1p::dsRed |
| Plasmid: *odr-1p::EFF-1* | [24] | odr-1p::EFF-1 |
| Plasmid: *mec-4p::EFF-1* | [24] | mec-4p::EFF-1 |
| Plasmid: *mec-4p::mCherry* | [24] | pmec-4::mCherry |
| Software | | |
| Fiji | NIH image | |
| GraphPad Prism 8 | GraphPad Software, Inc. | |
| Adobe Photoshop CS5 and CS6 | Adobe | |
| MetaMorph 7.8.1.0 | Molecular Devices | https://www.moleculardevices |
| Micro-Manager | [75] | https://micro-manager.org |

 **Data for movies**

**Movie S1. EFF-1 ectopic expression in BWMs results in Uncoordinated and Dumpy (Unc+Dpy) phenotypes**

Mixed population of worms with extrachromosomal *pmyo-3::mCherry* and *pmyo-3::EFF-1.* White arrowhead point to mCherry(+) worm that is Unc+Dpy. White arrow point to mCherry(+) worm that is wt-like. Yellow arrowhead points to a wt-like mCherry(-) worm, which left the frame within seconds. Elapsed time (seconds) indicated in top left corner.

**Movie S2. VSV∆G-G infects PVD neuron**

SDC microscope Z-stack of young adult worm injected with $10^6$ IU VSV∆G-G and imaged 48h

later. Arrowheads, infected PVD's candelabra/menorahs arborized branches. Arrows, infected

cell. BWM-Body Wall Muscle, EC-Excretory Cell, SE-Seam cell syncytium. Scale bars, 25 µm.

Note that there is a second worm (bottom, not indicated in images), also showing infected

PVD branches, EC, SE and a muscle cell.