Authors:

Masaru Sakai[1] ・ Katsuya Iwabuchi[2] ・ David Bauman[3,4]

Title:

Unique habitat and macroinvertebrate assemblage structures in spring-fed stream: a

comparison among clastic lowland tributaries and mainstreams in northern Japan

Affiliations:

[1] National Institute for Environmental Studies, Japan (Fukushima Branch), 10-2

Fukasaku, Miharu Town, Tamura District, Fukushima 963-7700, Japan

[2] Graduate School of Science and Engineering, Chuo University, 1-13-27 Kasuga,

Bunkyo-ku, Tokyo 112-8551, Japan

[3] Smithsonian Environmental Research Center, 647 Contees Wharf Road, Edgewater,

Maryland, 21037, USA

[4] Environmental Change Institute, School of Geography and the Environment,

University of Oxford, South Parks Road, Oxford, OX1 3QY, UK

✉ Masaru Sakai

sakai.masaru@nies.go.jp

ORCID

Masaru Sakai (0000-0001-5361-0978)

David Bauman (0000-0001-9115-6518)

Abstract

The stable flow and temperature regimes of spring-fed streams are distinct from the

dynamic regimes of other streams. We investigated differences in habitat and

macroinvertebrate assemblages among three stream types (spring-fed tributary,

non-spring-fed tributary and mainstream) in a clastic lowland of northern Japan.

Current velocity was the slowest in the spring-fed reach, where the percent of fine

sediment deposition was also 3.8–11.4 times higher than in the other stream types. The

standing stock of detritus was also greater in the spring-fed reach. These results

suggest that the stable flow regime in the spring-fed stream leads to the accumulation

of fine sediment and detritus on the streambed. Oligochaeta and chironomids, which

are burrower-gatherers, were remarkably abundant in the spring-fed reach. Total

macroinvertebrate abundance was 3.8–12.2 times greater in the spring-fed reach than

in the other stream types. Sprawler-grazer ephemeropterans were the most abundant in

the mainstream reaches, likely due to higher primary productivity. *Allomyia* sp, which

depend on cool spring-fed habitats, were found only in the spring-fed reach. The

indicator species analysis also indicated multiple taxa of detritivores and *Allomyia* sp.

for the spring-fed tributary. The macroinvertebrate assemblage in the spring-fed reach

was characterized by numerous burrowers, collector-gatherers, and crenobiont taxa,

highlighting the uniqueness and its contribution to enhance beta diversity in river

networks.

Keywords Community ecology · Functional feeding groups · Groundwater · Indicator

species · Life-form types · River ecosystem

Introduction

Groundwater flow into river ecosystems depends on underground flow pathways and aboveground topographic features (Allan and Castillo 2007). Although apportioning stream water to runoff and groundwater sources is generally difficult, streams that are sustained by a significant amount of permanent groundwater discharge from a spring mouth are typically referred to as spring-fed streams (Cantonati et al. 2010). Because groundwater flow is rarely affected by rapid fluctuations resulting from rainfall, snowmelt, or seasonal and diurnal changes in air temperature, spring-fed streams are characterized by stable water flow and temperature regimes (Mattson et al. 1995; Sear et al. 1999; Lusardi et al. 2016). Thus, spring-fed streams are distinct from the dynamic flow and temperature regimes that characterize non-spring-fed streams.

Flow regime is one of the most influential factors determining the structure of streambed substrates (Davis and Barmuta 1989). The frequency and magnitude of high flows are directly related to sediment runoff; stream substrate components are sequentially flushed from smaller to larger particles under high flows (Richards 2004). Streams with frequent high flows typically have a coarse streambed structure relative to those with less variable flow regimes. The substrates of spring-fed streams are dominated by fine sediments, owing to their stable flow regimes (Sear et al. 1999). Spring-fed streams in lowland regions may have particularly fine streambeds due to low channel gradients. Streambed particle size is a major determinant of the benthic macroinvertebrate community and its structure (Richards and Bacon 1994; Wood and Armitage 1997). For example, the relative density of burrowers and gatherers may increase with fine sediment deposition (Rabení et al. 2005). These groups are well

suited to depositional habitats because these habitats provide organic matter as well as

burrowing spaces. Lowland spring-fed streams may therefore house unique community

assemblages, dominated by burrowers and gatherers, relative to adjacent

non-spring-fed streams that have less fine sediment content.

Water temperature in spring-fed streams is highly stable; it is typically cold in

summer and warm in winter (Mattson et al. 1995). Previous studies have reported that

these stable water temperature regimes provide refugia for aquatic fauna to avoid

severe hot (Nakagawa et al. 2015) and cold seasonal conditions (Inoue and Ishigaki

1968). Moreover, Sun et al. (2020) suggested that most indicator species of spring-fed

habitats may be stenothermal. This further suggests that the narrow range of water

temperatures in spring-fed streams may lead to the formation of unique assemblages of

crenobiont and/or crenophilous species (Lusardi et al. 2016).

The stable flow and temperature regimes of spring-fed streams provide unique

habitat and macroinvertebrate assemblage structures and subsequently contribute to

beta diversity within river networks. However, studies on spring ecosystems have

advanced mainly in carbonate geologies (e.g., karst) with dense spring habitats

(Mattson et al. 1995; Wood et al. 2005; von Fumetti et al. 2017; von Fumetti and

Blattner 2017), and the ecosystem functions are poorly understood in other geologies.

This is perhaps related to sparse distributions of springs and difficulties to prepare

replicable sampling design in such geologies. Although more than 16,000 spring-fed

sites have been recognized in Japan (Ministry of Environment 2020), functional

aspects of spring-fed streams are still elusive in any geological conditions. We assessed

differences in habitat characteristics and macroinvertebrate assemblages between

spring-fed and adjacent non-spring-fed streams in a clastic lowland, one of the

dominant geological conditions in Japan. Our aim was to investigate potential

processes explaining these differences, detect potential indicator species of different

stream types, and to contribute to the understandings of faunal and functional

responses of macroinvertebrates to the unique habitats of spring-fed tributaries.

Materials and methods

Study area

We assessed streams in the Shubuto River system, located in Kuromatsunai, Hokkaido,

Japan (42.64°N, 140.34°E). The Shubuto River basin comprises 367 km^2 of montane

and lowland areas; our study focused on the lowland region (Fig. 1). Water quality is

suitable for most freshwater organisms: dissolved oxygen > 95% in degrees of

saturation, biochemical oxygen demand is 0.2-1.7 mg/L and ammonia concentration <

0.05mg/L (Miyazaki and Terui 2016). The geology is dominated by sandstone and

mudstone with Cainozoic fossil seashells and tuff. Mean annual precipitation and air

temperature, obtained from the Kuromatsunai automated meteorological data

acquisition system station (located 4 km northwest of the site) were 1615.8 mm and

7.5°C, respectively, between 2009 and 2018. Dominant tree species in the riparian

zones of the study area were *Salix* spp. and *Quercus crispula*, and dominant understory

plants were *Sasa kurilensis* and *Reynoutria sachalinensis*. High flows generally occur

in non-spring-fed streams during summer to autumn rainfalls and snowmelt season in

early spring (Fig. 2).

119   The lowland region of the Shubuto River basin includes areas of groundwater

discharge such as flowing wells, seepages, and springs. The spring-fed tributary

assessed in this study is fed by a significant amount of groundwater discharge

(approximately 0.1 m^3/s) that is directly connected to the Shubuto River (Sakai et al.

2020). This tributary has stable flow regime and water temperature compared to

adjacent mainstream (Fig. 2). We further selected a non-spring-fed tributary and three

mainstreams that also drained into the lowland area (Fig. 1). The spring-fed and

non-spring-fed tributaries were near to each other and had similar flow directions.

Three mainstream areas were identified below the confluence with the spring-fed

tributary (mainstream 1), between the confluences with the spring-fed and

non-spring-fed tributaries (mainstream 2), and below the confluence with the

non-spring-fed tributary (mainstream3) (Fig. 1). We used 40-m sections (reaches) in

each study site to assess macroinvertebrates and habitat characteristics. The

mainstream reaches had a typical fluvial geomorphology of Japanese midstream with

riffle-pool structures whereas the tributary reaches rarely had heterogeneous structures

like riffle-pools, step-pools, or cascades.

Although the river system has the only one typical spring-fed stream in the

lowland area, similarly to other river systems in clastic geology, assumable unique

habitat and community structures should be important for assessing local biodiversity

even if focal river systems have limited numbers of spring-fed streams. This limitation

may have hindered conceiving robust study design and advances in studies on

ecosystem functions of spring-fed streams draining outside carbonate geologies. This

study can therefore fill an important knowledge gap to understand the function of

sparsely distributed spring-fed streams in structuring local biodiversity.

Macroinvertebrate assemblage

We used Surber nets (25 × 25 cm, 0.5-mm mesh) to sample macroinvertebrates from

five randomly placed quadrats in each reach in June 2016. This period was chosen to

avoid seasons of snow accumulation, snowmelt, and primary insect emergences. Each

sample was collected along center lines of the streams to obtain representative

assemblages. The sampling points for the mainstream reaches categorized as riffles,

but those for the tributary reaches did not have any clear riffle nor pool structures.

Samples containing macroinvertebrates, detritus, and inorganic sediments were placed

in a tray and the macroinvertebrate and detritus portions were rinsed into unwoven

bags, and immediately preserved in 70% ethanol. Macroinvertebrates were sorted and

identified to the lowest possible taxonomic level using a stereomicroscope (SZ61,

Olympus, Tokyo, Japan) and keys published by Kawai and Tanida (2018) and Merritt

et al. (2008). Following identification, we classified macroinvertebrates by life-form

type (burrowers, sprawlers, clingers, and swimmers) and functional feeding group

(shredders, collector-gatherers, collector-filterers, predators, and grazers) based on

information from Takemon (2005) and Merritt et al. (2008) as well as our own

unpublished data.

Habitat characteristics

Stream width was measured with a measuring tape under base flow conditions at five

points in each study reach. We measured water depth and current velocity (repeated 5-s

measurements) using a folding scale and an electromagnetic current meter (VE10,

Kenek, Tokyo, Japan), respectively, at 15 points in each study reach during

macroinvertebrate sampling. Hemispherical photographs were taken in June 2016 at 10

m intervals along the 40 m reaches (five points) to estimate canopy openness.

Photographs were taken with a digital camera (D40, Nikon, Tokyo, Japan) equipped

with a fisheye lens (4.5 mm F2.8 EX DC Circular Fisheye HSM, Sigma, Kanagawa,

Japan) that was fixed horizontally at 1 m above the streambed. Canopy openness (%)

was estimated using CanopOn2 software (Takenaka 2009) from these photographs.

We applied the methods of Sakai et al. (2013) to estimate fine sediment

deposition in each study reach by visually estimating the percent cover of fine

sediment (< 2 mm) on the streambed in 40×40 cm quadrats with 20 rectangles

separated by wires (5% increments, ten quadrats per reach). We also estimated the

periphyton biomass on upper exposed surfaces of five submerged large cobbles in each

study reach. Periphyton was collected by brushing material from a 4 cm^2 patch on the

upper cobble surface and filtering the resulting slurry through pre-combusted

glass-microfiber filters (Whatman GF/F, Maidstone, United Kingdom). Prior to

periphyton sampling, each cobble was thoroughly rinsed in stream water to remove

attached fractions of nonalgal materials as much as possible. Periphyton dry mass and

ash-free dry mass (AFDM) were determined by drying the periphyton samples in an

oven (FC-610, Advantec, Tokyo, Japan) at 70°C for 24 h and then weighing them,

followed by combustion in a muffle furnace (FO300, Yamato Scientific Co., Ltd.,

Tokyo, Japan) at 500°C for 2 h and re-weighing. Detritus samples remained after

macroinvertebrate collections and identifications (five quadrat samples per reach) were

dried at ambient temperature for three days, followed by oven drying at 70°C for three

days and combustion in a muffle furnace at 500°C for 4 h. The AFDM of periphyton

and detritus samples were used as measures of periphyton biomass and standing stock

of detritus (mg/cm^2), respectively.

Statistical analyses

To test for differences in environmental variables, total abundance and taxa richness of

macroinvertebrates, and abundance of the most dominant taxa (> 1% of the individuals

in all samples) among the three stream types (spring-fed and non-spring-fed tributaries

and mainstream), we first used one-way analysis of variance (ANOVA) with

permutation test (Legendre 2007) to test for a significant difference of the response in

at least one stream type. When significant, we used the same permutation ANOVA test

for *post hoc* comparisons among the pairs of stream types. The permutation test used

allowed us to correctly handle the unbalanced design of the study (mainstream has

three times more observations than the tributaries). We adjusted *p*-values for multiples

tests using the false discovery rate correction (Benjamini and Hochberg 1995), to avoid

inflated type I error rates.

To visualize differences in macroinvertebrate assemblage structures among

stream types and associated environmental variables, we performed a principal

coordinate analysis (PCoA) based on the Chao index (Chao et al. 2005), followed by a

permutational multivariate analysis of variance (PERMANOVA), testing for significant

differences in macroinvertebrate assemblage structure among the stream types. Chao

index was calculated from the total numbers of individuals of each taxon in each

quadrat. We then superimposed *a posteriori* the environmental vectors (fine sediment

deposition, canopy openness, current velocity, stream width, water depth, standing

stock of detritus and periphyton biomass) using the weighted averages of the species,

and environmental variables (Borcard et al. 2018). The PERMANOVA was tested

using 999 permutations. In order to test for the presence of significant patterns of

spatial autocorrelation in the macroinvertebrate communities across our study reaches,

we generated spatial eigenvectors using Moran's eigenvector maps (MEM; Dray et al.

2006), a powerful multivariate approach allowing the detection of multiscale spatial

structures in ecological data. To obtain the spatial eigenvectors, we used a spatial

weighting matrix based on a Gabriel graph and weighted by a decreasing linear

function of the distance among streams (details in Bauman et al. 2018a), following

Bauman et al. (2018a, b). A redundancy analysis of the reach-scale community matrix

was performed against the whole set of spatial eigenvectors associated to positive

spatial autocorrelation patterns, for each spatial weighting matrix, and was tested by

permutations (9,999).

We additionally tested whether some of the macroinvertebrate taxa were

significant indicators of some of the stream types, that is, displaying both high fidelity

and specificity to one of the habitats. To do so, the *IndVal* index was calculated for all

species in the three stream types (Dufrêne and Legendre 1997) and was tested using

10,000 Monte Carlo permutations. All statistical analyses were performed in R version

3.6.3 (R Core Team 2020) using the packages *rcompanion* (Mangiafico 2020),

adespatial (Dray et al. 2020), *vegan* (Oksanen et al. 2019), *indicspecies* (Cáceres et al.

2020) and the function *anova.1way* (Legendre 2007).

Results

Stream width and water depth were significantly greater in mainstream reaches and

smallest in the non-spring-fed tributary (Table 1). Correspondingly, canopy openness

was relatively greater in the mainstreams compared to the tributaries. Current velocity

was significantly slower in the spring-fed tributary and faster in the mainstreams

(Table 1). The percent cover of fine sediment was significantly higher in the spring-fed

tributary; mean cover was approximately 11.4 times greater than those of the

mainstreams and 3.8 times greater than that of the non-spring-fed tributary (Table 1).

The periphyton biomass was not significantly different among the reaches, whereas the

standing stock of detritus was greatest in the spring-fed tributary (Table 1).

247   The total abundance and taxa richness of macroinvertebrates were both

significantly greater in the spring-fed tributary and smaller in the non-spring tributary

(Table 1; Supplementary Table S1). For life-form types, the four most abundant

burrower taxa (Oligochaeta, *Pagastia* spp., Chironomidae gen. spp. and Chironominae

spp.) and Tanypodinae sp. were more abundant in the spring-fed tributary. Sprawlers

were typically the most abundant group in the mainstreams, except for *Ordobrevia* sp.

(Table 1). For functional feeding groups, the four most dominant collector-gatherer

taxa (Oligochaeta, *Pagastia* spp., Chironomidae gen. spp. and Chironominae spp.)

were most abundant in the spring-fed tributary (Table 1). Excluding two trichopteran

grazers that were dominant in the spring-fed tributary (*Apatania* sp. and *Allomyia* sp.),

all grazer taxa reached their maximum abundance in the mainstreams. We noted that

Allomyia sp. was only found in the spring-fed tributary (Table 1). In this study, some

individuals of Chironomids could not be identified into subfamily or lower taxonomic

levels, so that Chironomidae gen. spp. may potentially influence the total number of

taxa, particularly in the spring-fed tributary that housed numerous Chironomid midges

(Table 1).

263   The MEM analysis did not detect any significant patterns of spatial

autocorrelation in the macroinvertebrate assemblage composition ($P = 0.68$), indicating

spatial independence among the communities. The first two axes of the PCoA captured

97.6% of the total variation in the community Chao index matrix (Fig. 3), and the

PERMANOVA indicated that macroinvertebrate assemblage structure significantly

differed among the stream types ($P = 0.001$). After overlapping the environmental

variables to the PCoA, the first PCoA axis indicated that the mainstreams presented

greater water depth, stream width, periphyton biomass, current velocity and canopy

openness than the spring-fed and non-spring-fed tributaries. The second PCoA axis

mostly separated the non-spring-fed tributary from the spring-fed tributary, the two

presenting very contrasted community composition as well as fine sediment deposition

and standing stock of detritus (higher for the spring-fed tributary). The indicator taxa

for the spring-fed tributary were mostly detritivores (shredders, collector-gatherers and

collector-filterers) of Chironomidae, Nemouridae, Ceratopogonidae, Lepidostomatidae,

Limnephilidae, Pisidiidae and Eusiridae, but also included one herbivorous

trichopteran (*Allomyia* sp.) (Table 2). The indicator taxa for the mainstreams were

mainly grazers (*Caenis* sp., *Epeorus* sp., *Drunella* sp., *Setodes* sp. and *Ecdyonurus*

viridis) and partly contained predators (*Drunella* sp. and *Rhyacophila nipponica*),

collector-gatherer (*Ephemerella setigera*) and collector-filterer (*Cheumatopsyche*

infascia) whereas those for the non-spring-fed tributary were collector-gatherer and/or

grazer (*Agapetus budoensis*) and collector-filterer (Simuliidae gen. spp.) (Table 2).

Discussion

Spring-fed streams in lowlands rarely have steep, deep channels (Sear et al. 1999) and

their flow regimes are stable (Mattson et al. 1995; Lusardi et al. 2016). Sakai et al.

(2020) showed that the spring-fed tributary assessed here maintained a stable flow

regime even during rainfall events, implying that sediment runoff during floods likely

occurs less in the tributary than in other non-spring-fed reaches. Because stable flow

regime can settle fine particles in streambed (e.g., Kiffney et al. 2000), the significantly

greater cover of fine sediment and higher standing stock of detritus observed in the

spring-fed tributary are attributable to its stable flow regime. The observed slower

current velocity in the spring-fed tributary could further contribute to the settlement of

fine particles. The relatively closed canopy openness in the spring-fed tributary might

additionally enhance accumulations of submerged detritus deriving from litter inputs.

Oligochaeta and most of the chironomid taxa were remarkably abundant in the

spring-fed tributary. As a result, the total macroinvertebrate abundance was 3.8–12.2

times greater in this tributary than in the other reaches. The relative abundance of

burrowers increases with fine sediment deposition (e.g. Zweig and Rabení 2001;

Rabení et al. 2005), and our findings suggest that the fine substrate found in spring-fed

streams provides attractive habitat for burrower species. Moreover, the stable flow

regimes of spring-fed streams may relate to less disturbance to macroinvertebrates

during flood events (e.g., Fritz and Dodds 2004; Robinson et al. 2004). Because the

burrower species that dominated in the spring-fed tributary typically gather and feed on

fine detritus (i.e., collector-gatherers), spring-fed environments clearly provide

adequate food resources for this group. The assumable long retention of submerged

detritus and associated consumptions and dominance of detritivores may thereby form

detritus-based food webs in spring-fed streams. Meanwhile, the strong variation of

macroinvertebrate abundances among the quadrats of a given stream type (e.g.,

Oligochaeta) suggest potential undetected microhabitat heterogeneity at finer spatial

scale, within the relatively homogeneous habitats found in the spring-fed tributary.

In contrast to our results, previous studies have reported that spring-fed

streams often have lush macrophytes (e.g., Sherwood et al. 2000; Katagiri et al. 2011)

and are accordingly dominated by herbivorous invertebrates (Takemon 2010).

Although obligate herbivores feeding on macrophytes are generally rare, macrophytes

may supplement food resources for grazers that usually consume algal masses and

provide spatially complex habitats. The streambed of the spring-fed stream we studied

lacked macrophytes perhaps because of limited light conditions. Food web and habitat

structures in spring-fed streams are probably site-specific and tightly associated with

environmental factors that affect the growth of macrophytes.

Wider streams with more open canopies generally induce higher primary

productivity (e.g., Vannote et al. 1980), but periphyton biomass was not significantly

different between the mainstreams and tributaries in this study. However,

ephemeropteran grazers significantly predominated in the mainstream reaches. This

contradiction is attributable to greater occupancy by grazers in reaches with higher

periphyton productivity. Feeding pressure of grazers is one of primary factors reducing

periphyton biomass (Feminella et al. 1989; Hill et al. 1995), and reaches with higher

productivity often have lower or similar periphyton biomass compared to those with

lower productivity (Feminella and Hawkins 1995). Therefore, grazing-based food webs

may be more pronounced in mainstreams and the dominance of grazer taxa and

associated consumption of periphyton may be responsible for the lack of differences in

periphyton biomass among reaches. The dominance of sprawlers in mainstreams is

attributable to their adaptability to fast-current habitats.

In the PCoA, the mainstream 2 had somewhat similar macroinvertebrate

assemblages to those in the spring-fed tributary. This result suggests that mainstream

reaches may have more heterogeneous habitats compared to tributaries, and partially

possess spring-fed-like macroinvertebrate communities even though the samples were

collected from riffles. For example, patches with upwelling water within reaches can

have such communities (Milner et al. 2001). Meanwhile, the macroinvertebrate

assemblages in the non-spring-fed tributary were clearly different from those in the

spring-fed tributary. Also, *Agapetus budoensis*, which was an indicator taxon of the

non-spring-fed tributary, was only found there. This indicates that the non-spring-fed

tributary also has unique habitats and macroinvertebrates assemblages compared to the

spring-fed tributary and mainstreams. Although taxa richness was lowest in the

non-spring-fed tributary, combinations of mainstream, spring-fed and non-spring-fed

tributaries can be important for maintaining the beta and gamma diversities of the river

network.

Given that the basal food resources and macroinvertebrate communities

differed between the spring-fed tributary and the adjacent mainstream reaches, we

suggest that the confluence of these streams may tangle the structure of the food web

such that the abundant macroinvertebrates in the spring-fed tributary may subsidize

predatory animals, such as fishes, inhabiting surrounding streams (cf. Uno and Power

2015). Moreover, the stable flow regime and water temperature in spring-fed streams

provide refugia during flood events (Sakai et al. 2020) and thermal stresses (Inoue and

Ishigaki 1968). Terui et al. (2018) reported that complex branches in river networks

can increase metapopulation stability for fish. Spring-fed streams may also contribute

to such stabilities in stream animal communities. Given that spring-fed habitats are

ubiquitous and distributed across Japan, examining the ecosystem functions of these

streams in nutrient cycling and as refugia is an important topic in river ecosystem

conservation. Given that the climate crisis relates to extreme runoff events (Yin et al.

2018) and severe thermal stress (van Vliet et al. 2013), these streams may be vital to

conservation efforts.

Two sprawler-grazer trichopteran species (*Allomyia* sp. and *Apatania* sp.) that

can be considered crenobiont and/or crenophilous were dominant in the spring-fed

tributary. In Japan, *Allomyia* species occur only in Hokkaido, and are preferentially

found in cool spring-fed habitats (Nishimoto and Kuhara 2001; Kawai and Tanida

2018). *Allomyia* sp. was found only in the spring-fed tributary in this study, similarly to

another study conducted in a montane spring-fed stream (Sueyoshi et al. 2014). At the

global scale, many of the crenophilous invertebrates are stenothermal, which is related

to their evolutionary origins (Sun et al. 2020), and the stable thermal environments of

spring-fed streams may provide refugia for these coldwater taxa (Lusardi et al. 2016).

The taxa that were dominant in the spring-fed tributary included stenothermal species,

implying that unique assemblages may be found in such streams. This may be an

important function of spring-fed streams; their unique habitats may contribute to the

beta diversity of focal river networks (Reiss et al. 2016).

Intensive researches conducted in carbonate geologies revealed that locations

of spring habitats are largely influential on local faunas. For example, faunal

characteristics tend to be similar between alpine springs and headwaters (Wigger et al.

2015), and substrate and macroinvertebrate assemblage structures vary with altitudinal

gradients (von Fumetti and Blatter 2017). Additionally, regions that receive small

amount of precipitations need to consider flow permanence of lotic systems. Some

studies evaluated faunal characteristics in between intermittent and perennial spring

habitats indicated similarity between them in high flow seasons (Wood et al. 2005;

White et al. 2018) but life cycle adaptations of some taxa in the intermittent habitats

(Wood et al. 2005). In lowland regions, Reiss et al. (2016) reported that crenobiont

taxa which can be considered as glacial relicts inhabit in spring habitats, corresponding

to this study. Although intermittent spring habitats may be relatively less under the

moist climate of Japan, a concern regarding impact of anthropogenic disturbance such

as land modifications and climate change on spring habitats (Cantonati et al. 2012) is

common across locations and geologies.

Our comparisons of spring-fed and non-spring-fed streams revealed unique

habitat characteristics and macroinvertebrate assemblage structure in the spring-fed

tributary. The macroinvertebrate assemblage was characterized by numerous burrowers

and collector-gatherers, and possibly crenobiont and/or crenophilous taxa. This unique

assemblage may therefore enhance the beta-diversity of this river network. Owing to

the high cover of fine sediments, the abundance of burrower taxa was remarkably high

in the spring-fed tributary. These taxa may further play significant roles as prey for

carnivores in the surrounding area. Although spring-fed streams are sparsely

distributed under clastic geology, ideal site selections that include multiple spring-fed

and non-spring-fed streams would further improve our understandings how spring-fed

habitats structure animal communities. Given that spring-fed streams may act as flow

and temperature refugia (Sakai et al. 2020, Lusardi et al. 2016), further studies that

examine availabilities of the refugia in spring-fed streams can inform biodiversity

conservation in river ecosystems. Here, we highlight the ecosystem function of

providing unique habitats and housing unique macroinvertebrate assemblages.

Acknowledgments Dr. Izumi Washitani provided invaluable comments on the study

plan. We thank the field assistance by Dr. Kosei Takahashi, Mr. Hitoshi Saito, Nobuo

Hatai and Kengo Ebihara.

Funding A portion of this study was supported by JSPS KAKENHI Grant numbers

2629181 and 19K20491, and Kuromatsunai Biodiversity Conservation Research Grant

(2017) to Masaru Sakai. David Bauman is supported by the Belgian American

Educational Foundation.

Competing interests The authors declare no conflict of interest.

Availability of data and materials The data that support the findings of this study are

available from the corresponding author upon reasonable request.

Authors' contributions MS, KI and DB conceived and designed the study. MS and KI

conducted the field investigations, and MS and DB performed the statistical analyses.

MS and KI wrote the first draft, and all authors contributed to the writing of the final

manuscript.

Ethical approval Not applicable.

Consent to participate Not applicable.

Consent for publication Not applicable

Code availability Not applicable.

References

Allan JD, Castillo MM (2007) Stream ecology: structure and function of running

waters. 2nd edition. Springer, Dordrecht, The Netherlands

Bauman D, Drouet T, Fortin MJ, Dray S (2018a) Optimizing the choice of a spatial

weighting matrix in eigenvector-based methods. Ecology 99:2159-2166

Bauman D, Drouet T, Dray S, Vleminckx J (2018b) Disentangling good from bad

practices in the selection of spatial or phylogenetic eigenvectors. Ecography

41:1638-1649

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and

powerful approach to multiple testing. J R Statist Soc B 57:289-300

Borcard D, Gillet F, Legendre P (2018) Numerical ecology with R. 2nd edition,

Springer, Dordrecht, The Netherlands

Cáceres MD, Jansen F, Dell N (2020) Package 'indicspecies'.

https://cran.r-project.org/web/packages/indicspecies/. Accessed 7 January 2021

Cantonati M, Lang-Bertalot H, Scalfi A, Angeli N (2010) *Cymbella tridentina* sp. nov.

(Bacillariophyta), a crenophilous diatom from carbonate springs of the Alps. J N

Am Benthol Soc 29:775-788

Cantonati M, Füreder L, Gerecke R, Jüttner I, Cox EJ (2012) Crenic habitats, hotspots

for freshwater biodiversity conservation: toward an understanding of their ecology.

Freshw Sci 31:463-480

Chao A, Chazdon RL, Colwell RK, Shen TJ (2005) A new statistical approach for

assessing similarity of species composition with incidence and abundance data.

Ecol Lett 8:148-159

Davis JA, Barmuta LA (1989) An ecologically useful classification of mean and

near-bed flows in streams and rivers. Freshw Biol 21:271-282

Dray S, Bauman D, Blanchet G, Borcard D, Clappe S, Guenard G, Jombart T,

Larocque G, Legendre P, Madi N, Wagner HH (2020) Package 'adespatial'.

https://cran.r-project.org/web/packages/adespatial/. Accessed 7 January 2021

Feminella JW, Power ME, Resh VH (1989) Periphyton responses to invertebrate

grazing and riparian canopy in three northern California coastal streams. Freshw

Biol 22:445-457

Feminella JW, Hawkins CP (1995) Interactions between stream herbivores and

periphyton: a quantitative analysis of past experiments. J N Am Benthol Soc

14:465-509

Fritz KM, Dodds WK (2004) Resistance and resilience of macroinvertebrate

assemblages to drying and flood in a tallgrass prairie stream system.

Hydrobiologia 527:99-112

Hill WR, Ryon MG, Schilling EM (1995) Light limitation in a stream ecosystem:

responses by primary producers and consumers. Ecology 76:1297-1309

Inoue S, Ishigaki K (1968) Notes on the biology of juvenile masu salmon

(*Oncorhynchus masou*) during winter in the Chihase River, Hokkaido. Jpn J

Limnol 29:27-36 (in Japanese with English abstract)

Katagiri K, Yabe K, Nakamura F, Sakurai Y (2011) Factors controlling the distribution

of aquatic macrophyte communities with special reference to the rapid expansion

of a semi-emergent *Phalaris arundinacea* L. in Bibi River, Hokkaido, northern

Japan. Limnology 12:175-185

Kawai T, Tanida K (2018) Aquatic insects of Japan: manual with keys and illustrations.

2nd edition. Tokai University Press, Kanagawa, Japan (in Japanese)

Kiffney PM, Richardson JS, Feller MC (2000) Fluvial and epilithic organic matter

dynamics in headwater streams of southwestern British Columbia, Canada. Arch

Hydrobiol 149:109-129

Legendre P (2007) One-way anova with permutation test.

http://adn.biol.umontreal.ca/~numericalecology/Rcode/. Accessed 7 January 2021

Lusardi RA, Bogan MT, Moyle PB, Dahlgren RA (2016) Environment shapes

invertebrate assemblage structure differences between volcanic spring-fed and

runoff rivers in northern California. Freshw Sci 35:1010-1022

Mangiafico S (2020) Package 'rcompanion'.

https://cran.r-project.org/web/packages/rcompanion/. Accessed 7 January 2021

Mattson RA, Epler JH, Hein MK (1995) Description of benthic communities in Karst,

spring-fed streams of northern central Florida. J Kansas Entomol Soc 68:18-41

Merritt RW, Cummins KW, Berg B (2008) An introduction to the aquatic insects of

North America. 4th edition. Kendall/Hunt Publishing Company, Iowa, USA

Milner AM, Brittain JE, Castella E, Petts GE (2001) Trends of macroinvertebrate

community structure in glacier-fed rivers in relation to environmental conditions:

a synthesis. Freshw Biol 46:1833-1847

Ministry of the Environment, Japan (MOE) (2020) Portal site for conservation of

springs. https://www.env.go.jp/water/yusui/. Accessed 7 January 2021 (in

Japanese)

Miyazaki Y, Terui Y (2016) Temporal dynamics of fluvial fish community caused by

marine amphidromous species in the Shubuto River, southwestern Hokkaido,

Japan. Ichthyol Res 63:173-179

Nakagawa H, Mishina T, Takemon Y (2015) Habitat use and nutritional conditions of

ayu (*Plecoglossus altivelis altivelis*) in the lower reaches of the Kamo River,

Kyoto, Japan. Ecol CivilEng 18:53-63 (in Japanese with English abstract)

Nishimoto H, Kuhara N (2001) Revision of the caddisfly *Allomyia* Banks of Japan

(Trichoptera: Apataniidae), with descriptions of seven new species. Entmol Sci

4:157-174

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin P R,

O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H (2019)

Package 'vegan'. https://cran.r-project.org/web/packages/vegan/. Accessed 7

January 2021

Rabení CF, Doisy KE, Zweig LD (2005) Stream invertebrate community functional

responses to deposited sediment. Aquat Sci 67:395-402

R Core Team (2020) R: a language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Accessed 7 January 2021

Reiss M, Martin P, Gerecke R, von Fumetti S (2016) Limno-ecological characteristics

and distribution patterns of spring habitats and invertebrates from the Lowlands to

the Alps. Environ Earth Sci 75:1033

Richards C, Bacon KL (1994) Influence of fine sediment on macroinvertebrate

colonization of surface and hyporheic stream substrates. Great Basin Nat

54:106-113

Richards KS (2004) Rivers: form and process in alluvial channels. Blackburn Press,

New Jersey, USA

Robinson CT, Aebischer S, Uehlinger U (2004) Immediate and habitat-specific

responses of macronvertebrates to sequential, experimental floods. J N Am

Benthol Soc 23:853-867

Sakai M, Natuhara Y, Fukushima K, Naito R, Miyashita H, Kato M, Gomi T (2013)

Responses of macroinvertebrate communities to 4 years of deer exclusion in first-

and second-order streams. Freshw Sci 32:563-575

Sakai M, Wakiya R, Hoshi G (2020) Spring-fed stream as a refugium: temporal

changes in fish assemblage before and after a rainfall event. *bioRxiv*

https://doi.org/10.1101/2020.04.26.062968.

Sear DA, Armitage PD, Dawson FH (1999) Groundwater dominated rivers. Hydrol

Process 13:255-276

Sherwood AR, Rintoul TL, Müller KM, Sheath RG (2000) Seasonality and distribution

of epilithic diatoms, macroalgae and macrophytes in a spring-fed stream system in

Ontario, Canada. Hydrobiologia 435:143-152

Sueyoshi M, Nakano D, Nakamura F (2014) The relative contributions of refugium

types to the persistence of benthic invertebrates in a seasonal snowmelt flood.

Freshw Biol 59:257-271

Sun Y, Takemon Y, Yamashiki Y (2020) Freshwater spring indicator taxa of benthic

invertebrates. Ecohydrol Hydrobiol 20:622-631

Takemon Y (2005) Life-type concept and functional feeding groups of benthos

communities as indicators of lotic ecosystem conditions. Jpn J Ecol 55:189-197

(in Japanese with English abstract)

Takemon Y (2010) Benthos community structure and characteristics in Kakita River.

In: Kakitagawa Ecosystem Workshop (Ed.), *The Nature of Kakita River* (in

Japanese)

Takenaka A (2009) CanopOn 2 program for analyzing hemispherical photographs.

http://takenaka-akio.org/etc/canopon2/. Accessed 7 January 2021

Terui A, Ishiyama N, Urabe H, Ono S, Finlay JC, Nakamura F (2018) Metapopulation

stability in branching networks. Proc. Natl. Acad. Sci. U. S. A. 115:E5963-E5969

Uno H, Power ME (2015) Mainstem-tributary linkages by mayfly migration help

sustain salmonids in warming river networks. Ecol Lett 18:1012-1020

Vannote RL, Minshall GW, Cummins KW, Sedell JR, Cushing CE (1980) The river

continuum concept. Can J Fish Aquat Sci 37:130-137

van Vliet MTH, Franssen WHP, Yearsley JR, Ludwig F, Haddeland I, Lettenmaier DP,

Kabat P (2013) Global river discharge and water temperature under climate

change. Glob Environ Change 23:450-464

von Fumetti S, Bieri-Wigger F, Nagel P (2017) Temperature variability and its

influence on macroinvertebrate assemblages of alpine springs. Ecohydrology

10:e1878

von Fumetti S, Blattner L (2017) Faunistic assemblages of natural springs in different

areas in the Swiss National Park: a small-scale comparison. Hydrobiologia

793:175-184

White JC, House A, Punchard N, Hannah DM, Wilding NA, Wood PJ (2018)

Macroinvertebrate community responses to hydrological controls and

groundwater abstraction effects across intermittent and perennial headwater

streams. Sci Total Environ 610-611:1514-1526

Wigger FW, Schmidlin L, Nagel P, von Fumetti S (2015) Macroinvertebrate

assemblages of natural springs along an altitudinal gradient in the Bernese Alps,

Switzerland. Ann Limnol – Int J Lim 51:237-247

Wood PJ, Armitage PD (1997) Biological effects of fine sediment in the lotic

environment. Environ Manage 21:203-217

Wood PJ, Gunn J, Smith H, Abas-Kutty A (2005) Flow permanence and

macroinvertebrate community diversity within groundwater dominated headwater

streams and springs. Hydrobiologia 545:55-64

Yin J, Gentine P, Zhou S, Sullivan SC, Wang R, Zhang Y, Guo S (2018) Large increase

in global storm runoff extremes driven by climate and anthropogenic changes. Nat

Commun 9:4389

Zweig LD, Rabení CF (2001) Biomonitoring for deposited sediment using benthic

invertebrates: a test on 4 Missouri streams. J N Am Benthol Soc 20:643-657

Table 1 Environmental characteristics, total abundance, taxa richness, and the

abundance of each dominant taxa in study reaches. LFT: life form type, B: burrower,

Sp: sprawler, C: clinger, Sw: swimmer. FFG: functional feeding group, C/G:

collector-gatherer, P: predator, G: grazer. Bold characters indicate statistically

significant difference among stream types by one-way analyses of variances with

permutation tests ($P < 0.05$), and different alphabetical letters indicate statistically

significant differences by *post hoc* pairwise tests ($P < 0.05$). *P*-values are corrected for

multiple tests (see Methods).

Variable	LFT	FFG	Spring-fed tributary			Non-spring-fed tributary			Mainstreams			F^*	P
			Mean	±	SD	Mean	±	SD	Mean	±	SD		
Environmental													
Stream width (m)	-	-	3.8	±	0.3^b	1.0	±	0.2^c	12.7	±	2.4^a	87.40	**<0.01**
Water depth (cm)	-	-	16.8	±	3.8^b	9.4	±	1.9^c	32.0	±	2.7^a	421.21	**<0.01**
Current velocity (cm/s)	-	-	12.7	±	3.0^c	24.5	±	5.5^b	41.8	±	16.0^a	32.90	**<0.01**
Canopy openness (%)	-	-	34.3	±	12.1^b	45.5	±	17.4^{ab}	55.1	±	15.0^a	3.77	**0.043**
Fine sediment deposition (%)	-	-	83.5	±	10.6^a	22.0	±	7.2^b	7.3	±	3.7^c	554.89	**<0.01**
Periphyton biomass (mg/cm^2)	-	-	1.4	±	0.4	1.5	±	0.3	1.6	±	0.4	0.53	0.589
Standing stock of detritus (mg/cm^2)	-	-	1.7	±	0.9^a	0.7	±	1.0^{ab}	0.3	±	0.2^b	9.14	**<0.01**
Macroinvertebrates													
Total abundance	-	-	416.4	±	210.9^a	34.0	±	18.2^c	108.5	±	53.4^b	22.46	**<0.01**
Taxa richness	-	-	23.8	±	3.8^a	10.0	±	3.9^c	17.9	±	3.2^b	20.03	**<0.01**
Oligochaeta	B	C/G	116.6	±	141.5	0.6	±	1.3	20.3	±	34.2	4.84	**0.018**
Pagastia spp.	B	C/G	116.2	±	52.9^a	0.0	±	0.0^b	0.4	±	0.7^b	52.81	**<0.01**
Drunella ishiyamana	Sp	G, P	2.6	±	2.1^b	0.2	±	0.4^c	31.8	±	20.3^a	10.53	**<0.01**
Apatania sp.	C	G	34.8	±	21.7^a	7.0	±	4.6^b	0.1	±	0.5^c	25.18	**<0.01**
Chironomidae gen. spp.	B	C/G	40.0	±	41.5^a	0.0	±	0.0	0.0	±	0.0^b	10.20	**<0.01**
Chironominae spp.	B	C/G	30.2	±	31.9^a	0.0	±	0.0	0.0	±	0.0^b	9.86	**<0.01**
Caenis sp.	Sp	C/G, G	0.0	±	0.0^b	0.0	±	0.0^b	8.7	±	8.3^a	5.23	**0.023**
Baetis sp.	Sw	G	3.0	±	3.2	2.6	±	3.4	4.9	±	4.2	0.86	0.440
Dicranota spp.	B	P	5.4	±	2.6	4.4	±	3.4	2.9	±	3.3	1.24	0.296

Allomyia sp.	C	G	18.4	±	11.7[a]	0.0	±	0.0[b]	0.0	±	0.0[b]	27.22	**<0.01**
Epeorus sp.	Sp	G	0.2	±	0.4	0.0	±	0.0	6.0	±	5.6	5.24	**0.013**
Ordobrevia sp.	Sp	C/G	1.8	±	1.3	4.2	±	1.3	4.1	±	5.8	0.49	0.625
Ephemerella setigera	Sp	C/G	0.4	±	0.5	0.0	±	0.0	5.6	±	6.4	3.33	**0.049**
Ecdyonurus viridis	Sp	G	0.4	±	0.5	0.0	±	0.0	3.9	±	4.4	3.45	**0.044**
Tanypodinae sp. 1	B	P	12.2	±	9.9[a]	0.0	±	0.0[b]	0.0	±	0.0[b]	16.59	**<0.01**
Drunella sp.	Sp	P, G	0.2	±	0.4	0.0	±	0.0	3.5	±	3.3	5.18	**0.018**
Orthocladiinae sp.	B	C/G	2.0	±	1.9	1.2	±	0.8	2.1	±	1.4	0.70	0.526
Paraleptophlebia japonica	B	C/G	2.4	±	3.7	0.6	±	1.3	1.7	±	1.7	0.88	0.422

*$F_{2,72}$ for water depth and current velocity, $F_{2,47}$ for fine sediment deposition, and $F_{2,22}$

for the other variables.

Table 2 Statistically significant indicator taxa for each stream type ($P < 0.05$; adjusted P

for multiple tests). LFT: life form type, B: burrower, C: clinger, Sp: sprawler, Sw:

swimmer. FFG: functional feeding group, C/G: collector-gatherer, G: grazer, P: predator,

S: shredder, C/F: collector-filterer.

Taxon	LFT	FFG	Stream type	*IndVal*	*P*
Chironomidae gen. spp.	B	C/G	Spring-fed tributary	1.000	0.0001
Allomyia sp.	C	G	Spring-fed tributary	1.000	0.0001
Tanypodinae sp. 1	B	P	Spring-fed tributary	1.000	0.0001
Pagastia spp.	B	C/G	Spring-fed tributary	0.998	0.0001
Nemoura sp.	Sp	S	Spring-fed tributary	0.995	0.0001
Chironominae spp.	B	C/G	Spring-fed tributary	0.894	0.0009
Tanypodinae sp. 2	B	P	Spring-fed tributary	0.894	0.0009
Ceratopogonidae gen. spp.	B	C/G	Spring-fed tributary	0.894	0.0009
Lepidostoma japonicum	Sp	S	Spring-fed tributary	0.824	0.0052
Nothopsyche sp.	Sp	C/G, S	Spring-fed tributary	0.775	0.0101
Pisidium japonicum	B	C/F	Spring-fed tributary	0.775	0.0080
Stenomoera yezoensis	Sw	C/G, S	Spring-fed tributary	0.764	0.0068
Agapetus budoensis	C	C/G, G	Non-spring-fed tributary	1.000	0.0002
Simuliidae gen. spp.	C	C/F	Non-spring-fed tributary	0.764	0.0124
Caenis sp.	Sp	C/G, G	Mainstream	0.966	0.0001
Epeorus sp.	Sp	G	Mainstream	0.950	0.0004
Drunella sp.	Sp	G, P	Mainstream	0.940	0.0006
Ephemerella setigera	Sp	C/G	Mainstream	0.899	0.0028
Setodes sp.	Sp	C/G, G	Mainstream	0.856	0.0044
Ecdyonurus viridis	Sp	G	Mainstream	0.852	0.0154
Cheumatopsyche infascia	C	C/F	Mainstream	0.775	0.0193
Rhyacophila nipponica	Sp	P	Mainstream	0.730	0.0464

Figure captions

Fig. 1 Study site location within the Shubuto River system in Kuromatsunai, Hokaido,

Japan. Open circles indicate the study reaches.

Fig. 2 Flow and temperature regimes of the spring-fed tributary and adjacent

mainstream. Water levels were initially set at 0 cm at the onset of the monitoring.

Fig. 3 Results of the principal coordinate analysis based on the abundance of all

macroinvertebrate taxa and the environmental variables. Significant indicator taxa for

each stream type are shown nearby the plots.

Kuromatsunai lowland

Caenis sp.
Epeorus sp.
Drunella sp.
Ephemerella setigera
Setodes sp.
Ecdyonurus viridis
Cheumatopsyche infascia
Rhyacophila nipponica

Agapetus budoensis
Simuliidae gen. spp.

Chironomidae gen. spp.
Allomyia sp.
Tanypodinae sp. 1
Pagastia spp.
Nemoura sp.
Chironominae spp.
Tanypodinae sp. 2
Ceratopogonidae gen. spp.
Lepidostoma japonicum
Nothopsyche sp.
Pisidium japonicum
Sternomoera yezoensis

PCoA 1 (54.0%)
PCoA 2 (43.6%)

●: spring-fed tributary
▲: non-spring-fed tributary
■: mainstream 1
◆: mainstream 2
⊠: mainstream 3

FSD: fine sediment deposition
CO: canopy openness
V: current velocity
WID: stream width
DEP: water depth
detr: standing stock of detritus
peri: periphyton biomass