1

Montane species and communities track recent warming more closely in the tropics

Running head: temperature tracking across latitudes

Benjamin G. Freeman[1,2*] , Yiluan Song[3], Kenneth J Feeley[4] , Kai Zhu[3]

[1] Biodiversity Research Centre, University of British Columbia, Vancouver, BC, Canada

V6T1Z4

[2] Department of Zoology, University of British Columbia, Vancouver, BC, Canada

V6T1Z4

[3] Department of Environmental Studies, University of California, Santa Cruz, CA 95064

USA

[4] Department of Biology, University of Miami, Coral Gables, FL 33156 USA

*Corresponding author: freeman@zoology.ubc.ca

Abstract

Tropical species have been hypothesized to be particularly sensitive to temperature

increases because of their physiological specialization to stable climates. Here we use

two independent datasets to test whether tropical montane species are tracking recent

warming via upslope range shifts better than temperate montane species. First, we

compiled published data documenting recent elevational shifts of plants and animals. For

this resurvey data, we analyzed temperature tracking for both entire communities (N =

83) and for individual species (N = 2871). Second, we calculated temperature tracking of

tree community structure using data from repeatedly surveyed forest inventory plots from

mountainous regions of the Americas (N = 300). We defined temperature tracking scores

as the ratio of observed changes vs. expected changes based on estimates of local

warming. Both datasets show that tropical montane taxa are tracking changing

temperatures significantly better than their temperate counterparts. On average, tropical

communities are tracking temperature 2.6 times more than temperate communities (0.84

± 0.15 vs. 0.32 ± 0.074), tropical species are tracking temperature 2.3 times more than

temperate species (0.91 ± 0.18 vs. 0.40 ± 0.083), and tropical tree plots are tracking

temperature 4.4 times more than temperate tree plots (0.44 ± 0.11 vs. 0.099 ± 0.042).

This latitudinal gradient in temperature tracking is consistent with accumulating evidence

that tropical montane biotas are disproportionately sensitive to ongoing global warming.

Keywords: global warming, latitudinal gradient, mountaintop extinction, range shift,

upslope shift

Introduction

One consequence of global warming is the upslope shift, or "migration", of many montane species' ranges to higher, cooler elevations (I.-C. Chen, Hill, Ohlemüller, Roy, & Thomas, 2011; Parmesan & Yohe, 2003). However, there is significant variation in the degree to which montane species are effectively tracking temperature increases via distributional shifts to higher elevations. For example, while moths in Borneo are rapidly shifting their ranges upslope at rates that approximate that of concurrent warming (I. C. Chen et al., 2009; Wu et al., 2019), the upslope shifts in the ranges of plants in the European Alps lag far behind the pace of concurrent warming (Rumpf et al., 2018), and some birds in northeastern North America are shifting their ranges *downslope* despite recent warming (DeLuca & King, 2017; Zuckerberg, Woods, & Porter, 2009).

Here we address the question "why are some species and communities effectively tracking changes in local temperatures along mountain slopes while others are not?" Specifically, we test the hypothesis that variation in temperature tracking is explained, at least in part, by the species' latitude. Tropical species are hypothesized to be disproportionately sensitive to rising temperatures, even though the absolute magnitude of recent warming is low in the tropics compared to higher latitudes. The explanation for greater sensitivity of tropical species is that they tend to live closer to their optimal temperatures and exhibit narrower thermal tolerances than their temperate counterparts (Deutsch et al., 2008; Laurance et al., 2011; Perez, Stroud, & Feeley, 2016; Sunday, Bates, & Dulvy, 2012). Several studies have reported that tropical montane species are moving rapidly upslope associated with recent warming (Sheldon, 2019). It is unknown, however, whether these scattered reports are general across taxa and continents.

We used two independent datasets to test the hypothesis that temperature tracking

in montane taxa is related to latitudinal position. First, we compiled a dataset of resurveys

that have reported elevational shifts for communities and species associated with recent

warming (Fig. 1a, Table S1, Datasets S1 and S2). By "communities", we mean a set of

species within a particular taxonomic group that lives within a particular montane region

(e.g., a bird community or a bee community). Second, we analyzed inventory data from

repeatedly surveyed forest plots located in montane areas across the Americas and that

have experienced significant warming over the past several decades (Fig. 1b). These two

datasets have complementary strengths for inferring latitudinal patterns in recent

temperature tracking. Resurveys have the advantage of explicitly measuring changes in

elevational distributions for individual species, with data available for many taxonomic

groups, but have the disadvantage that studies may use different methods to study

different taxa. In contrast, forest inventories use standardized methods and provide

comprehensive information about the composition of tree communities at given locations.

We calculated "temperature tracking" scores for resurvey and forest inventory

datasets. For resurveys, we analyzed data for both communities and individual species.

Note that not all studies that reported range shifts for communities also reported range

shifts for the individual species that comprised the community. We calculated

temperature tracking scores as the ratio of the observed elevational shift vs. the expected

elevational shift, where the expected elevational shift is based on estimates of local

warming between sampling events and adiabatic lapse rates. For forest inventory data, we

calculated community temperature indices (CTI) as the weighted mean of species'

temperature optima based on the species composition of plots at each sampling event. We

then calculated temperature tracking as the annualized change in CTI (the

thermophilization rate) divided by the observed rate of change in mean annual

temperature from 1979 to 2013. In both cases, temperature tracking scores of ~1 indicate

that observed changes closely match those expected based on concurrent warming (i.e.,

strong temperature tracking), and scores >1 indicate that species are shifting their ranges

faster than expected. Conversely, lower temperature tracking scores indicate little or no

change in the ranges of species or the composition of tree plots despite rising

temperatures. Negative scores indicate changes in species ranges or composition that are

in the "wrong" direction (e.g., downslope shifts despite warming temperatures).

We then tested the hypothesis that tropical montane species are tracking

temperature increases better than temperate montane species. We analyzed the resurvey

data at both the community-level (data = 146 estimates of temperature tracking from 83

communities, data provided in Dataset S1) and species-level (data = 5860 estimates of

temperature tracking from 2871 species from 65 communities; species-level data was

unavailable for 18 communities; see Fig. S1, data provided in Dataset S2). Our dataset of

forest inventory plots contained 256 plots from temperate montane forests from the

United States and 44 plots of tropical montane forests from Central and South America;

(see Fig. S2, data provided in Dataset S3). The combination of two independent datasets,

with complementary strengths, and large sample sizes within each dataset, allow us to

rigorously test the hypothesis that there is a latitudinal gradient in temperature tracking.

Methods

Resurvey data

Our aim was to compile a comprehensive list of studies that have measured recent range shifts for species in mountainous regions in response to recent warming. First, we conducted a Web of Science search on 11 July 2019 with the keywords "climate change" OR "global warming" AND "range shift" AND "mountain" OR "elevation" OR "altitude*". This search returned 1827 hits. We retained studies that met the following three criteria: (1) they measured recent range shifts at species' lower elevation limits, mean (or optimum) elevations, or upper elevational limits; (2) range shifts were reported for all species or the entire community, not just species with significant range shifts; and (3) range shifts were measured over a time period of >=10 years. Second, we located additional studies that met these three criteria by examining recent papers synthesizing the range shift literature (I.-C. Chen et al., 2011; Freeman, Lee-Yaw, Sunday, & Hargreaves, 2018; Lenoir & Svenning, 2015; Rumpf, Hülber, Zimmermann, & Dullinger, 2019; Wiens, 2016). There were three cases for which multiple publications reported elevational range shifts for the same community using the same underlying data (plants in France, and butterflies and birds in Great Britain). For these cases, we included only the study with a larger sample size of species.

For each study that met our criteria (see Table S1, full data provided in Dataset S1), we extracted the following key information: (1) taxon; (2) latitude of the study site; (3) duration over which range shifts were calculated (e.g., number of years elapsed between surveys); (4) spatial scale of study ("local" when studies were conducted along single elevational gradients or entirely within small montane regions; "regional" when the study was conducted within large political units such as the state of California or the

country of Spain); (5) number of species studied; (6) range shifts at lower elevational

limits, mean/optimum elevations, and upper elevational limits for the entire community

(i.e., for a community of 30 species, the mean range shift of these 30 species); (7) range

shifts at lower elevational limits, mean/optimum elevations, and upper elevational limits

for individual species (note that not all studies reported species-specific range shifts); (8)

temperature changes at the study site between surveys (because we were interested in

responses to warming, we did not include two studies that reported cooling temperatures,

both from the Appalachians in North America (Moskwik, 2014)); and, if reported, (9)

expected elevational range shifts based on local temperature changes and lapse rate.

There were two studies that did not report local temperature changes (Kusrini et al., 2017;

Moret, Aráuz, Gobbi, & Barragán, 2016). For these studies, we estimated local

temperature changes using data from high-resolution gridded data provided by the

Climatic Research Unit (Harris, Jones, Osborn, & Lister, 2013). When studies did not

report local adiabatic lapse rates, we assumed a lapse rate of 0.55°C / 100 m elevation,

which was the mean lapse rate from the 14 studies that reported locally measured lapse

rates. When data were presented only in figures, we used WebPlotDigitizer to extract

data from the published graphics (Rohatgi, 2017).

All statistics were done in R version 3.6.2 (R Development Core Team, 2020).

We first analyzed categorical latitudinal differences in temperature tracking between the

tropics and the temperate zone. To do so, we fit linear mixed-effects models using the

"lme4" package (Bates, Maechler, Bolker, & Walker, 2014). The response variable in

both community-level and species-level models was the temperature tracking score,

calculated as the ratio of the observed elevational shift to the expected elevational shift

given estimates of local warming and adiabatic lapse rate. We included latitudinal zone

(tropical vs. temperate) and four methodological covariates as fixed effects: (1)

distributional variable measured (lower limit vs. mean elevation vs. upper limit); (2)

spatial scale of study (local vs. regional); (3) number of species in the study (for the

community-level model only); and (4) duration of the study. We included the community

ID as a random effect, as multiple distributional variables were reported for many

communities.

We then analyzed latitudinal differences in temperature tracking as a continuous

function of position along the latitudinal gradient. These models were the same as those

described above, but with the absolute value of latitude as a fixed effect. To investigate

whether our results were driven by the inclusion of communities with few species, we fit

an additional model including data only from communities with >=10 species (125

estimates of temperature tracking from 70 communities). We did not include taxa as a

predictor variable in the community or species-level models because latitudinal sampling

was poor for most taxonomic groups.

Forest Inventory Data

We compiled our forest inventory plot dataset using Forest Inventory and Analysis (FIA)

plots from the United States and previously published inventory plot data from Central

and South America (Fadrique et al., 2018; Feeley, Hurtado, Saatchi, Silman, & Clark,

2013). Because we were interested in changes in mountains, we selected FIA plots

(Smith, 2002) that fell within mountainous areas using the global mountains raster map

derived from the 250 m global Hammond landforms product (Karagulle et al., 2017). We

included only FIA plots that had been surveyed at least twice. We collected the species

identity of all individual adult trees (diameter at breast height > 12.7 cm) for each survey

of each FIA plot. We then calculated the community temperature index (CTI) of the plot

as the weighted average of optimal temperature of each species, following the method of

Fadrique et al. (Fadrique et al., 2018). Specifically, we downloaded all georeferenced

plant location records available through the BIEN database (version 4.1.1 accessed in

November 2018 via BIEN package in R) for the New World (North America, Central

America, and South America, but excluding the Caribbean islands). The BIEN database

provides collated observation and collection data from multiple sources and provides a

base level of data filtering and standardization. We used BIEN's default download

preferences to exclude records of known introduced species and cultivated individuals.

We further filtered the records to include only those that were georeferenced and that list

the year of collection/observation as being between 1970 and 1980 (we restricted records

to just this 10-year window to minimize errors due to species migrating through time).

For each species, we then extracted the estimated mean annual temperature (BIOCLIM1)

at all retained collection coordinates from the CHELSA v1.2 raster of "current" (i.e.,

mean of 1979-2012) climate at 30 arc-second resolution and estimated the thermal optima

(MAT_{opt}) of each species as the mean MAT of all collections. If species had fewer than

10 records, we estimated MAT_{opt} at the genus-level using combined records of all

congeners. Genera with <10 records were excluded from subsequent analyses. We next

used the collection records to calculate the community temperature index (CTI) of each

plot in each census as the mean MAT_{opt} for species in the plot weighted by their relative

basal area.

We then calculated the thermophilization rate of each FIA plot as the linear trend

of CTI over time, and obtained previously published thermophilization rates for 10 sites

in Costa Rica and 63 sites in the tropical Andes (Fadrique et al., 2018; Feeley et al.,

2013). Using these data, we calculated the rate of temperature change for all sites using

monthly mean temperature estimates from 1980 to 2013 from the CHELSA time-series

dataset (Karger & Zimmermann, 2019). Specifically, we calculated the rate of

temperature change at each site as the linear trend of mean annual temperature over time.

As FIA plots are generally smaller in size compared to the plots in Central and South

America (168 m^2 for FIA plots compared to 1-hectare for most tropical plots), we

aggregated FIA plots into 1 degree diameter hexagons that contain approximately 30

plots each. We removed hexagons with <5 plots from subsequent analysis. As with the

resurvey data, we calculated the thermal tracking score for each plot as the ratio of

observed changes (rate of change in thermophilization rate) and expected changes (rate of

change of mean annual temperature).

In order to focus on the response of organisms to the warming climate, we

selected plots with a significant warming trend ($p < 0.05$) and examined the relationships

between absolute latitude and the temperature tracking score by fitting linear mixed-

effects models using the "spaMM" package in R (Rousset, 2017). We included both

absolute latitude and elevation as fixed effects. Due to the close proximity of plots and

spatial dependence in the residuals of non-spatial linear models, we modeled spatial

random effects that follow Gaussian covariance functions, and then evaluated the

significance of regression coefficients for fixed effects using the conditional standard

error of regression coefficients.

Results

Resurvey data show that tropical montane taxa are tracking temperature increases

"better" (i.e., have higher temperature tracking scores) than are temperate montane

species (Fig. 2a-2d, Fig. S3). We first categorized communities and species as being

"tropical" or "temperate" based on the location of the study, with tropical locations

defined as $< 23.4°$ absolute latitude. Tropical communities had temperature tracking

scores 2.6 times greater than temperate communities (Fig. 2a, Table S2; temperature

tracking scores for tropical and temperate communities $= 0.84 \pm 0.15$ vs. 0.32 ± 0.074, p

$= 0.0021$; estimates $\pm$ standard errors from mixed-effects models), and tropical species

had temperature tracking scores 2.3 times than temperate species (Fig, 2c, Table S3;

temperature tracking scores for tropical and temperate species $= 0.91 \pm 0.18$ vs. $0.40 \pm$

0.083, $p = 0.0082$).

Results were similar when we modeled temperature tracking from resurvey data

as a continuous function of position along the latitudinal gradient. Temperature tracking

scores decreased by an average of 0.16 ± 0.049 and 0.13 ± 0.053 per 10 degrees latitude

for communities and species, respectively (Figs. 2b and 2d, Tables S4 and S5). Hence,

tropical communities and species are closely tracking temperature changes while

temperate zone communities and species are not, though the explanatory power of

community models was much greater (marginal R^2 values from linear mixed-effects

models $= 0.12$ and 0.010 respectively). Methodological covariates included in models had

minimal explanatory power, with the exception that temperature tracking scores tended to

be larger at species' upper elevational limits compared to at their lower elevational limits

or mean elevations (Tables S2-S5). Results for communities all held when considering

only communities with 10 or more species (Fig. S4, Tables S6 and S7), indicating that the

results are not driven by the inclusion of depauperate communities.

Forest inventory data also show that tropical montane taxa are tracking

temperature increases better than temperate montane species (Fig. 2e and 2f, Fig S5,

Tables S8-S9). Tropical montane forest plots had higher temperature tracking scores

(0.44 ± 0.11) than temperate montane forest plots $(0.099 \pm 0.042; p = 0.0038;$ Table S8).

Latitude is a strong predictor in the linear mixed model: a 10-degree increase in latitude

corresponds to a 0.14 ± 0.039 decrease in the temperature tracking score (Table S9). The

greater temperature tracking in tropical inventory plots was due to faster changes in CTI

in the tropics, also known as the thermophilization rate. Tropical plots had

thermophilization rates of 0.095 ± 0.020 °C per decade vs. 0.033 ± 0.011 °C per decade

for temperate plots (Fig. S6). In contrast, the rate of warming was faster at high latitudes

in our dataset: 0.31 ± 0.0061 °C per decade for temperate plots vs. 0.22 ± 0.010 °C per

decade for tropical plots (Fig. S6).

Discussion

We found strong evidence that tropical montane species are tracking recent temperature

increases better than temperate montane species. Temperature tracking was greater in the

tropics in each of three datasets: (1) community-level data from resurvey studies, (2)

species-level data from resurvey studies, and (3) repeatedly-censused forest inventory

tree plots. Moreover, the effect of latitude is large, with tropical montane species tightly

tracking temperature increases (temperature tracking scores ~ 1, which indicates perfect

temperature tracking) and temperate montane species tracking temperature increases only

weakly (temperature tracking scores closer to 0; 0 would indicate a complete lack of

temperature tracking. Specifically, for resurvey data, estimates from linear mixed models

are that communities have a tracking score of 1.02 ± 0.20 at the equator, but 0.32 ± 0.074

at 45° latitude (averaged over levels of fixed effects; the equivalent values for the

species-level model are 0.99 ± 0.22 and 0.39 ± 0.86). For forest tree plot data, the model-

based estimate is that tree plots have a tracking score of 0.63 ± 0.0069 at the equator, but

-0.00080 ± 0.0068 at 45° latitude.

While both resurvey and forest inventory plot datasets show marked latitudinal

patterns in temperature tracking, differences in temperature tracking scores across

latitudes were higher for the resurvey dataset. This difference could reflect a biological

difference in generation time between species included in the datasets. Trees have long

generation times that lead to slow rates of community turnover and range shifts (Feeley,

Rehm, & Machovina, 2012; Lenoir, Gegout, Marquet, De Ruffray, & Brisse, 2008). In

comparison, the resurvey dataset consists primarily of taxa with much shorter generation

times such as birds, mammals, and herbaceous plants. An alternative explanation that

requires testing is that methodological differences between resurveys and forest tree

inventory plots, both in data collection and analysis, may explain why temperature

tracking scores differ between these datasets.

Our model explains the observed variation in temperature tracking at the level of

communities much better than for individual species. Individual species show a wide

variation in their temperature tracking scores (Freeman, Lee-Yaw, et al., 2018; Moritz et

al., 2008; Rumpf et al., 2019). Latitude is a significant predictor of temperature tracking

in the species-level model, but its explanatory power is low. Combined, these results

imply that despite a clear latitudinal pattern of temperature tracking for communities, we

still have only a limited ability to predict changes in individual species' elevational

distributions associated with warming temperatures (Angert et al., 2011). Range shifts at

the species-level may be more predictable in the marine realm, where most species

appear to closely track temperature (Lenoir et al., 2019; Pinsky, Eikeset, McCauley,

Payne, & Sunday, 2019).

Multiple mechanisms may explain why tropical species are tracking temperature

better than temperate species. The leading explanation is that tropical species are more

physiologically sensitive to climate change than are temperate species. This explanation

implies that the mean annual temperature is a more important driver of species'

elevational distributions in the tropics than at higher latitudes. An additional possibility is

that both tropical and temperate species are tracking recent warming, but that temperate

species are using phenological shifts to do so (Socolar, Epanchin, Beissinger, & Tingley,

2017). Seasonal temperature fluctuations in the tropics are minimal, meaning that tropical

species are unlikely to be able to track climate via phenological shifts. In other words,

temperate-zone species may track changing climate by shifting in time, while tropical

species track changing climate by shifting in space. If so, we may expect a decoupling of

biodiversity change and carbon dynamics in the temperate zone.

Several limitations of our study deserve explicit mention. First, we followed

previous analyses in calculating temperature tracking scores based on mean annual

temperature (I.-C. Chen et al., 2011). Analyses that incorporate temperature variability

(i.e., seasonality) and other climatic variables have also proven powerful (Crimmins,

Dobrowski, Greenberg, Abatzoglou, & Mynsberge, 2011; Tingley, Koo, Moritz, Rush, &

Beissinger, 2012), but are inherently more difficult to implement and interpret. Second,

future studies should address land use drivers of recent distribution changes in

conjunction with climate drivers (Guo, Lenoir, & Bonebrake, 2018; Larsen, 2012; Lenoir

& Svenning, 2015). Third, there remains a strong temperate zone bias in data on

elevational range shifts associated with recent climate change (Feeley, Stroud, & Perez,

2017). Tropical studies are indeed underrepresented in our datasets, but the general trends

we document are robust to the unbalanced samples. Fourth, the patterns we document are

strong but not without exceptions. For example, Puerto Rican frogs and birds are not

closely tracking temperature despite their tropical latitude (Campos-Cerqueira, Arendt,

Wunderle, & Aide, 2017; Campos-Cerqueira & Mitchell Aide, 2017), though this could

potentially reflect increasing forest cover on this island (Battey et al., 2019). Conversely,

some temperate zone communities are closely tracking recent warming (Kelly &

Goulden, 2008; Menéndez, González-Megías, Jay-Robert, & Marquéz-Ferrando, 2014).

The latitudinal gradient in temperature tracking we document has multiple

implications for the conservation of montane floras and faunas. The weak temperature

tracking of temperate montane species and communities indicates that acclimation and

adaptation, rather than elevational shifts, will likely be key processes helping temperate

montane species persist in the face of continued warming. It is an open question whether

adaptation and acclimation will be able to keep pace with rates of warming that are

unprecedented in recent evolutionary time. In contrast, the strong temperature tracking of

tropical montane species indicates that they may be able to persist despite warming

temperatures by shifting their distributions upslope, at least when protected elevational

corridors provide suitable habitats at higher elevations. On the other hand, this implies

that the "escalator to extinction"—wherein montane species respond to warming

temperatures by shifting upslope until they run out of available habitat on mountaintops

(Freeman, Scholer, Ruiz-Gutierrez, & Fitzpatrick, 2018)—runs faster in the tropics.

Notably, such mountaintop extinctions may occur well below the actual mountaintop, as

pervasive anthropogenic modifications of high-elevation tropical systems effectively

limit the ability of the tree line to shift upslope (Rehm & Feeley, 2015). Indeed, local

extinctions and range contractions associated with recent warming appear to be most

common in tropical montane species (Freeman, Lee-Yaw, et al., 2018; Wiens, 2016).

The sixth mass extinction in Earth's history is now underway(Ceballos, Ehrlich,

& Dirzo, 2017). The tropics have the highest species diversity of any biome, and tropical

mountains have the highest diversity of all (Rahbek et al., 2019). The relatively small

temperature changes in the tropics should minimize the impact of climate change, but the

disproportionately sensitive responses of tropical montane species have instead placed

this whole biota on an escalator to extinction. The degree to which predictions of

widespread species extinctions in tropical mountains (Sekercioglu, Schneider, Fay, &

Loarie, 2008) come true will depend on our ability to protect elevational corridors that

enable species to persist while shifting upslope (Feeley & Rehm, 2012), and, ultimately,

on whether humanity is able to slow global warming.

Acknowledgments

We thank the many scientists who have published resurvey data and measured forest

inventory plots. The Feeley lab and R. Yorque provided cogent comments that improved

this manuscript. BGF was supported from postdoctoral fellowships from Banting Canada

(379958), and the Biodiversity Research Centre, and YS was supported by a Regents'

fellowship from the University of California, Santa Cruz. KZ and KJF are supported by

the US National Science Foundation (NSF grants DEB 1926438 to KZ and DEB-

1350125 to KJF).

Author contributions

BGF compiled resurvey dataset; YS, KZ and KJF compiled forest inventory tree plot

dataset; BGF, YS and KZ conducted analyses; BGF led the writing process with

extensive input from KJF, YS and KZ.

Author contributions

The authors declare no competing interests.

References

Angert, A. L., Crozier, L. G., Rissler, L. J., Gilman, S. E., Tewksbury, J. J., & Chunco,

A. J. (2011). Do species' traits predict recent shifts at expanding range edges?

Ecology Letters, *14*, 677–689.

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2014). lme4: Linear mixed-effects

models using Eigen and S4.

Battey, C. J., Otero, L. M., Gorman, G. C., Hertz, P. E., Lister, B. C., García, A., …

Huey, R. B. (2019). Why Montane Anolis Lizards are Moving Downhill While

Puerto Rico Warms. *BioRxiv*, 751941. https://doi.org/10.1101/751941

Campos-Cerqueira, M., Arendt, W. J., Wunderle, J. M., & Aide, T. M. (2017). Have bird

distributions shifted along an elevational gradient on a tropical mountain? *Ecology*

and Evolution, *7*(23), 9914–9924. https://doi.org/10.1002/ece3.3520

Campos-Cerqueira, M., & Mitchell Aide, T. (2017). Lowland extirpation of anuran

populations on a tropical mountain. *PeerJ*, *2017*(11).

https://doi.org/10.7717/peerj.4059

Ceballos, G., Ehrlich, P. R., & Dirzo, R. (2017). Biological annihilation via the ongoing

sixth mass extinction signaled by vertebrate population losses and declines.

Proceedings of the National Academy of Sciences of the United States of America,

114(30), E6089–E6096. https://doi.org/10.1073/pnas.1704949114

Chen, I.-C., Hill, J. K., Ohlemüller, R., Roy, D. B., & Thomas, C. D. (2011). Rapid range

shifts of species associated with high levels of climate warming. *Science*, *333*(6045),

1024–1026. https://doi.org/10.1126/science.1206432

Chen, I. C., Shiu, H. J., Benedick, S., Holloway, J. D., Chey, V. K., Barlow, H. S., …

Thomas, C. D. (2009). Elevation increases in moth assemblages over 42 years on a

tropical mountain. *Proceedings of the National Academy of Sciences*, *106*(5), 1479–

1483. https://doi.org/10.1073/pnas.0809320106

Crimmins, S., Dobrowski, S., Greenberg, J., Abatzoglou, J., & Mynsberge, A. (2011).

Changes in climatic water balance drive downhill shifts in plant species' optimum

elevations. *Science*, *331*, 324–327.

DeLuca, W. V., & King, D. I. (2017). Montane birds shift downslope despite recent

warming in the northern Appalachian Mountains. *Journal of Ornithology*, *158*(2),

493–505. https://doi.org/10.1007/s10336-016-1414-7

Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak,

D. C., & Martin, P. R. (2008). Impacts of climate warming on terrestrial ectotherms

across latitude. *Proceedings of the National Academy of Sciences*, *105*(18), 6668–

6672. https://doi.org/DOI 10.1073/pnas.0709472105

Fadrique, B., Báez, S., Duque, Á., Malizia, A., Blundo, C., Carilla, J., … Feeley, K. J.

(2018). Widespread but heterogeneous responses of Andean forests to climate

change. *Nature*, *564*(7735), 207–212. https://doi.org/10.1038/s41586-018-0715-9

Feeley, K. J., Hurtado, J., Saatchi, S., Silman, M. R., & Clark, D. B. (2013).

Compositional shifts in Costa Rican forests due to climate-driven species

migrations. *Global Change Biology*, *19*(11), 3472–3480.

https://doi.org/10.1111/gcb.12300

Feeley, K. J., & Rehm, E. M. (2012). Amazon's vulnerability to climate change

heightened by deforestation and man-made dispersal barriers. *Global Change*

Biology, *18*(12), 3606–3614. https://doi.org/10.1111/gcb.12012

Feeley, K. J., Rehm, E. M., & Machovina, B. (2012). The responses of tropical forest

species to global climate change: acclimate, adapt, migrate, or go extinct? *Frontiers*

of Biogeography, *4*(2), 69–84.

Feeley, K. J., Stroud, J. T., & Perez, T. M. (2017). Most 'global' reviews of species'

responses to climate change are not truly global. *Diversity and Distributions*, *23*(3),

231–234. https://doi.org/10.1111/ddi.12517

Freeman, B. G., Lee-Yaw, J. A., Sunday, J. M., & Hargreaves, A. L. (2018). Expanding,

shifting and shrinking: The impact of global warming on species' elevational

distributions. *Global Ecology and Biogeography*, *27*, 1268–1276.

Freeman, B. G., Scholer, M. N., Ruiz-Gutierrez, V., & Fitzpatrick, J. W. (2018). Climate

change causes upslope shifts and mountaintop extirpations in a tropical bird

community. *Proceedings of the National Academy of Sciences of the United States*

of America, *115*(47), 11982–11987. https://doi.org/10.1073/pnas.1804224115

Guo, F., Lenoir, J., & Bonebrake, T. C. (2018). Land-use change interacts with climate to

determine elevational species redistribution. *Nature Communications*, *9*(1).

https://doi.org/10.1038/s41467-018-03786-9

Harris, I., Jones, P. D., Osborn, T. J., & Lister, D. H. (2013). Updated high-resolution

grids of monthly climatic observations–the CRU TS3. 10 Dataset. *International*

Journal of Climatology.

Karagulle, D., Frye, C., Sayre, R., Breyer, S., Aniello, P., Vaughan, R., & Wright, D.

(2017). Modeling global Hammond landform regions from 250-m elevation data.

Transactions in GIS, *21*(5), 1040–1060. https://doi.org/10.1111/tgis.12265

Karger, D. N., & Zimmermann, N. E. (2019). Climatologies at High resolution for the

Earth Land Surface Areas CHELSA V1 . 2: Technical specification. *Scientific Data*,

4(April), 170122. https://doi.org/10.1594/WDCC/CHELSA

Kelly, A. E., & Goulden, M. L. (2008). Rapid shifts in plant distribution with recent

climate change. *Proceedings of the National Academy of Sciences*, *105*(33), 11823–

11826.

Kusrini, M. D., Lubis, M. I., Endarwin, W., Yazid, M., Darmawan, B., Ul-Hasanah, A.

U., … Rachmadi, R. (2017). Elevation range shift after 40 years: The amphibians of

Mount Gede Pangrango National Park revisited. *Biological Conservation*, *206*, 75–

84. https://doi.org/10.1016/j.biocon.2016.12.018

Larsen, T. H. (2012). Upslope range shifts of Andean dung beetles in response to

deforestation: compounding and confounding effects of microclimatic change.

Biotropica, *44*(1), 82–89. https://doi.org/10.1111/j.1744-7429.2011.00768.x

Laurance, W. F., Carolina Useche, D., Shoo, L. P., Herzog, S. K., Kessler, M., Escobar,

F., … Thomas, C. D. (2011). Global warming, elevational ranges and the

vulnerability of tropical biota. *Biological Conservation*, *144*(1), 548–557.

https://doi.org/10.1016/j.biocon.2010.10.010

Lenoir, J., Bertrand, R., Comte, L., Bourgeaud, L., Hattab, T., Murienne, J., &

Grenouillet, G. (2019). Species better track the shifting isotherms in the oceans than

on land. *Https://Www.Biorxiv.Org/Content/10.1101/765776v1*.

Lenoir, J., Gegout, J. C., Marquet, P. A., De Ruffray, P., & Brisse, H. (2008). A

significant upward shift in plant species optimum elevation during the 20th century.

Science, *320*(5884), 1768–1771.

Lenoir, J., & Svenning, J. C. (2015). Climate-related range shifts - a global

multidimensional synthesis and new research directions. *Ecography*, *38*(1), 15–28.

https://doi.org/10.1111/ecog.00967

Menéndez, R., González-Megías, A., Jay-Robert, P., & Marquéz-Ferrando, R. (2014).

Climate change and elevational range shifts: evidence from dung beetles in two

European mountain ranges. *Global Ecology and Biogeography*, *23*(6), 646–657.

https://doi.org/10.1111/geb.12142

Moret, P., Aráuz, M. de los Á., Gobbi, M., & Barragán, Á. (2016). Climate warming

effects in the tropical Andes: first evidence for upslope shifts of Carabidae

(Coleoptera) in Ecuador. *Insect Conservation and Diversity*, *9*(4), 342–350.

484  https://doi.org/10.1111/icad.12173

485 Moritz, C., Patton, J. L., Conroy, C. J., Parra, J. L., White, G. C., & Beissinger, S. R.

486  (2008). Impact of a century of climate change on small-mammal communities in

487  Yosemite National Park, USA. *Science*, *322*(5899), 261–264.

488  https://doi.org/10.1126/science.1163428

489 Moskwik, M. (2014). Recent elevational range expansions in plethodontid salamanders

490  (Amphibia: Plethodontidae) in the southern appalachian mountains. *Journal of*

491  *Biogeography*, *41*(10), 1957–1966. https://doi.org/10.1111/jbi.12352

492 Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change

493  impacts across natural systems. *Nature*, *421*(6918), 37–42.

494  https://doi.org/10.1038/nature01286nature01286

495 Perez, T. M., Stroud, J. T., & Feeley, K. J. (2016). Thermal trouble in the tropics.

496  *Science*, *351*(6280), 1392–1393. https://doi.org/10.1126/science.aaf3343

497 Pinsky, M. L., Eikeset, A. M., McCauley, D. J., Payne, J. L., & Sunday, J. M. (2019).

498  Greater vulnerability to warming of marine versus terrestrial ectotherms. *Nature*,

499  *569*, 108–111. https://doi.org/10.1038/s41586-019-1132-4

500 R Development Core Team. (2020). R: A language and environment for statistical

501  computing. Vienna, Austria: R Foundation for Statistical Computing.

502 Rahbek, C., Borregaard, M. K., Antonelli, A., Colwell, R. K., Holt, B. G., Nogues-Bravo,

503  D., … Fjeldså, J. (2019). Building mountain biodiversity: Geological and

504  evolutionary processes. *Science*, *365*(6458), 1114–1119.

505  https://doi.org/10.1126/science.aax0151

506 Rehm, E. M., & Feeley, K. J. (2015). The inability of tropical cloud forest species to

invade grasslands above treeline during climate change: Potential explanations and

consequences. *Ecography*, *38*(12), 1167–1175. https://doi.org/10.1111/ecog.01050

Rohatgi, A. (2017). WebPlotDigitizer 3.11.

Rousset, F. (2017). An introduction to the spaMM package for mixed models, 1–35.

Rumpf, S. B., Hülber, K., Klonner, G., Moser, D., Schütz, M., Wessely, J., … Dullinger,

S. (2018). Range dynamics of mountain plants decrease with elevation. *Proceedings

of the National Academy of Sciences*, 201713936.

https://doi.org/10.1073/PNAS.1713936115

Rumpf, S. B., Hülber, K., Zimmermann, N. E., & Dullinger, S. (2019). Elevational rear

edges shifted at least as much as leading edges over the last century. *Global Ecology

and Biogeography*, *28*(4), 533–543. https://doi.org/10.1111/geb.12865

Sekercioglu, C. H., Schneider, S. H., Fay, J. P., & Loarie, S. R. (2008). Climate change,

elevational range shifts, and bird extinctions. *Conservation Biology*, *22*(1), 140–150.

https://doi.org/DOI 10.1111/j.1523-1739.2007.00852.x

Sheldon, K. S. (2019). Climate Change in the Tropics: Ecological and Evolutionary

Responses at Low Latitudes. *Annual Review of Ecology, Evolution, and Systematics*,

50(1), annurev-ecolsys-110218-025005. https://doi.org/10.1146/annurev-ecolsys-

110218-025005

Smith, W. B. (2002). Forest inventory and analysis: A national inventory and monitoring

program. *Environmental Pollution*, *116*(SUPPL. 1), 233–242.

https://doi.org/10.1016/S0269-7491(01)00255-X

Socolar, J. B., Epanchin, P. N., Beissinger, S. R., & Tingley, M. W. (2017). Phenological

shifts conserve thermal niches in North American birds and reshape expectations for

climate-driven range shifts. *Proceedings of the National Academy of Sciences*,

201705897. https://doi.org/10.1073/pnas.1705897114

Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2012). Thermal tolerance and the global

redistribution of animals. *Nature Climate Change*, *2*(9), 686–690.

https://doi.org/10.1038/nclimate1539

Tingley, M. W., Koo, M. S., Moritz, C., Rush, A. C., & Beissinger, S. R. (2012). The

push and pull of climate change causes heterogeneous shifts in avian elevational

ranges. *Global Change Biology*, *18*(11), 3279–3290. https://doi.org/10.1111/j.1365-

2486.2012.02784.x

Wiens, J. J. (2016). Climate-Related Local Extinctions Are Already Widespread among

Plant and Animal Species. *PLoS Biology*, *14*(12), 1–18.

https://doi.org/10.1371/journal.pbio.2001104

Wu, C., Holloway, J. D., Hill, J. K., Ho, C., Thomas, C. D., & Chen, I. (2019). Reduced

body sizes in climate-impacted Borneo moth assemblages are primarily explained by

range shifts. *Nature Communications*, 1–7. https://doi.org/10.1038/s41467-019-

12655-y

Zuckerberg, B., Woods, A. M., & Porter, W. F. (2009). Poleward shifts in breeding bird

distributions in New York State. *Global Change Biology*, *15*(8), 1866–1883.

Fig 1. Maps of resurvey studies that measured elevational shifts associated with recent

warming (a) and of forest inventory tree plots that have been repeatedly censused (b).

Locations of resurvey studies are jittered slightly to improve clarity. The Tropics of

Cancer and Capricorn (at 23.4° N and S, respectively) delimit the tropics, and are

illustrated with dashed lines.

Fig. 2. Tropical communities, species, and forest inventory tree plots have higher

temperature tracking scores than their temperate zone counterparts. (a, c, e) Temperature

tracking scores in tropical and temperate zones; (b, d, f) relationship between temperature

tracking and absolute latitude. Raw data are shown as points. Dashed lines illustrate both

perfect temperature tracking (temperature tracking = 1) and zero temperature tracking

despite warming temperatures (temperature tracking = 0). Trendlines illustrate

predictions from mixed models. (a) temperature tracking for communities in tropical and

temperate zones; (b) relationship between temperature tracking and absolute latitude (°)

for communities; (c) temperature tracking for species in tropical and temperate zones (~

0.5% of data has been removed for visualization); (d) relationship between temperature

tracking and absolute latitude (°) for species; (e) temperature tracking for forest inventory

tree plots in tropical and temperate zones; (f) relationship between temperature tracking

and absolute latitude (°) for forest inventory tree plots.

a resurvey data

b forest tree plot data

amphibian
arthropod
bird
fish
mammal
plant
reptile