Montane species and communities track rising temperatures better in the tropics

Benjamin G. Freeman[1,2*] , Yiluan Song[3], Kenneth J Feeley[4] , Kai Zhu[3]

[1] Biodiversity Research Centre, University of British Columbia, Vancouver, BC, Canada

[2] Department of Zoology, University of British Columbia, Vancouver, BC, Canada

[3] Department of Environmental Studies, University of California, Santa Cruz, CA USA

[4] Department of Biology, University of Miami, Coral Gables, FL 33156 USA

Abstract

Many species are responding to recent climate change by shifting their distributions

upslope to higher elevations, but the observed rates of shifts vary considerably among

studies. Here we test the hypothesis that latitudinal position in part explains this variation,

using two independent datasets—resurveys of species' elevational ranges, and repeatedly

surveyed tree plots. We find that communities are shifting 18.6 m per decade upslope, but

these upslope shifts are slightly faster in the tropics (25.1 m per decade on average) than

the temperate zone (13.1 m per decade on average). Faster upslope shifts in the tropics

are surprising because rates of warming are faster in the temperate zone. Consequently,

tropical communities are tracking changing temperatures 2.5 times (resurvey dataset) and

10.0 times (tree plot dataset) more than their temperate counterparts. For the amount of

warming expected to lead to a 100 m upslope shift, tropical communities have shifted

upslope by an average of 81 m while temperate communities have shifted upslope by an

average of only 32 m. Similarly, for the amount of warming expected to lead to a 1 °C

increase in CTI, tropical tree plots have increased in CTI by an average of 0.44 °C while

temperate tree plots have increased by an average of only 0.044 °C. This latitudinal

gradient in responsiveness to recent warming suggests tropical montane biotas are

disproportionately sensitive to ongoing global warming.

Introduction

One consequence of global warming is the upslope shift, or "migration", of many

montane species' ranges to higher, cooler elevations (Chen, Hill, Ohlemüller, Roy, &

Thomas, 2011; Parmesan & Yohe, 2003). However, there is significant variation in the

degree to which montane species are effectively tracking temperature increases via

distributional shifts to higher elevations. For example, while moths in Borneo are rapidly

shifting their ranges upslope at rates that approximately match local warming rates (Chen

et al., 2009; Wu et al., 2019), the upslope shifts in the ranges of plants in the European

Alps lag far behind the pace of warming (Rumpf *et al.* 2018), and some birds in

northeastern North America are shifting their ranges *downslope* despite recent warming

(Zuckerberg *et al.* 2009; DeLuca & King 2017).

Here we address the question "why are some species and communities effectively

tracking changes in local temperatures along mountain slopes while others are not?"

Specifically, we test the hypothesis that variation in temperature tracking is explained, at

least in part, by the species' latitude. Tropical species are hypothesized to be

disproportionately sensitive to rising temperatures, even though the absolute magnitude

of recent warming is lower in the tropics than in higher latitudes. The explanation for

greater sensitivity of tropical species is that they tend to live closer to their optimal

temperatures and exhibit narrower thermal tolerances than their temperate counterparts

(Deutsch *et al.* 2008; Laurance *et al.* 2011; Sunday *et al.* 2012; Perez *et al.* 2016). Several

studies have reported cases where tropical montane species are moving rapidly upslope

associated with recent warming (reviewed by Freeman & Class Freeman, 2014; Sheldon,

2019). It is unknown, however, whether these scattered reports are general across taxa

and continents.

We used two independent datasets to test the hypothesis that temperature tracking

in montane taxa is related to latitudinal position. First, we compiled a dataset of resurveys

that have reported elevational shifts for communities and species associated with recent

warming (Fig. 1a, Table S1, Datasets S1 and S2). By "communities" we mean a set of

species within a particular taxonomic group that lives within a particular montane region

(e.g., a bird community or a bee community). Second, we analyzed inventory data from

repeatedly surveyed forest plots located in montane areas across the Americas and that

have experienced significant warming over the past several decades (Fig. 1b). These two

datasets have complementary strengths for inferring latitudinal patterns in recent

temperature tracking. Resurveys have the advantage of explicitly measuring changes in

elevational distributions for individual species, with data available for many taxonomic

groups, but have the disadvantage that studies use different methods to study different

taxa. In contrast, forest inventories use standardized methods and provide comprehensive

information about the composition of tree communities at given locations.

We calculated "temperature tracking" scores for resurvey and forest inventory

datasets. For resurveys, we analyzed data for both communities and individual species.

We calculated temperature tracking scores as the ratio of the observed elevational shift

vs. the expected elevational shift, where the expected elevational shift is based on

estimates of local warming between sampling events and measured or estimated adiabatic

lapse rates. For forest inventory data, we calculated community temperature indices

(CTI) as the weighted mean of species' temperature optima based on the species

composition of plots at each sampling event. We then calculated temperature tracking as

the annualized change in CTI (i.e., the "thermophilization rate") divided by the observed

rate of change in mean annual temperature from 1979 to 2013. In both cases, temperature

tracking scores of ~1 indicate that observed changes closely match those expected based

on concurrent warming (i.e., strong temperature tracking), and scores >1 indicate that

species are shifting their ranges faster than expected. Conversely, temperature tracking

scores that are positive but closer to 0 indicate little or no change in the ranges of species

or the composition of tree plots despite rising temperatures. Negative scores indicate

changes in species ranges or composition that are in the "wrong" direction (e.g.,

downslope shifts despite warming temperatures).

We then tested the hypothesis that tropical montane species are tracking

temperature increases better than temperate montane species. We analyzed the resurvey

data at both the community-level (data = 146 estimates of temperature tracking from 83

communities, data provided in Dataset S1) and species-level (data = 5860 estimates of

temperature tracking from 2871 species from 65 communities; species-level data was

unavailable for 18 communities; see Fig. S1, data provided in Dataset S2). Our dataset of

forest inventory plots contained 212 plots from temperate montane forests from the

United States and 44 plots of tropical montane forests from Central and South America;

(see Fig. S2, data provided in Dataset S3). Our primary analyses focus on temperature

tracking, an index of relative response to climate change given local temperature

increase. However, latitudinal variation in the absolute responses to climate change is

also of interest, as greater warming at high latitudes could counter tighter temperature

tracking in the tropics and lead to similar rates of upslope shift across latitude. We

therefore also assess the latitudinal patterns in upslope range shifts (resurvey dataset) and

thermophilization rates (tree plot dataset). The combination of two independent datasets,

with complementary strengths, and large sample sizes within each dataset, allow us to

rigorously test the hypothesis that there is a latitudinal gradient in temperature tracking.

Materials and methods

Resurvey data

Our aim was to compile a comprehensive list of studies that have measured recent range

shifts for species in mountainous regions in response to recent warming (within the past ~

100 years). First, we conducted a Web of Science search on 11 July 2019 with the

keywords "climate change" OR "global warming" AND "range shift" AND "mountain"

OR "elevation" OR "altitude*". This search returned 1827 hits. We retained studies that

met the following three criteria: (1) they measured recent range shifts at species' lower

elevation limits, mean (or optimum) elevations, or upper elevational limits; (2) range

shifts were reported for all species or the entire community, not just species with

significant range shifts; and (3) range shifts were measured over a time period of >=10

years. Second, we located additional studies that met these three criteria by examining

recent papers synthesizing the range shift literature (Chen et al., 2011; Freeman, Lee-

Yaw, Sunday, & Hargreaves, 2018; Lenoir & Svenning, 2015; Rumpf, Hülber,

Zimmermann, & Dullinger, 2019; Wiens, 2016). There were three cases for which

multiple publications reported elevational range shifts for the same community using the

same underlying data (plants in France, and butterflies and birds in Great Britain). For

these cases, we included only the study with a larger sample size of species.

For each study that met our criteria (see Table S1, full data provided in Dataset

S1), we extracted the following key information: (1) taxon; (2) latitude of the study site;

(3) duration over which range shifts were calculated (e.g., number of years elapsed

between surveys); (4) spatial scale of study ("local" when studies were conducted along

single elevational gradients or entirely within small montane regions; "regional" when

the study was conducted within large political units such as the state of California or the

country of Spain); (5) number of species studied; (6) range shifts at lower elevational

limits, mean/optimum elevations, and upper elevational limits for the entire community

(i.e., for a community of 30 species, the mean range shift of these 30 species); (7) range

shifts at lower elevational limits, mean/optimum elevations, and upper elevational limits

for individual species (note that not all studies reported species-specific range shifts); (8)

temperature changes at the study site between surveys (because we were interested in

responses to warming, we did not include two studies that reported cooling temperatures,

both from the Appalachians in North America (Moskwik 2014)); and, if reported, (9)

expected elevational range shifts based on local temperature changes and lapse rate.

There were two studies that did not report local temperature changes (Moret *et al.* 2016;

Kusrini *et al.* 2017). For these studies, we estimated local temperature changes using data

from gridded data provided by the Climatic Research Unit (Harris *et al.* 2013). For

studies that did not report local adiabatic lapse rates, we used lapse rates reported for the

geographically nearest study within our dataset, following Chen et al. (2011). When data

were presented only in figures, we used WebPlotDigitizer to extract data from the

published graphics (Rohatgi 2017).

All statistics were done in R version 3.6.2 (R Development Core Team 2020). We

first analyzed categorical latitudinal differences in temperature tracking between the

tropics and the temperate zone. We categorized communities and species as being

"tropical" or "temperate" based on the location of the study, with tropical locations

defined as $< 23.4°$ absolute latitude. We then fit linear mixed-effects models using the

"lme4" package (Bates *et al.* 2014). The response variable in both community-level and

species-level models was the temperature tracking score, calculated as the ratio of the

observed elevational shift to the expected elevational shift given estimates of local

warming and adiabatic lapse rate. We included latitudinal zone (tropical vs. temperate)

and four methodological covariates as fixed effects: (1) distributional variable measured

(lower limit vs. mean elevation vs. upper limit); (2) spatial scale of study (local vs.

regional); (3) number of species in the study (for the community-level model only); and

(4) duration of the study. We included the community ID as a random effect, as multiple

distributional variables were reported for many communities.

We then analyzed latitudinal differences in temperature tracking as a continuous

function of position along the latitudinal gradient by replacing the factor

"tropical/temperate" with absolute latitude in models. We did not include taxa as a

predictor variable in the community or species-level models because taxonomic

differences predict minimal observed variation in recent range shifts (Chen *et al.* 2011;

Lenoir *et al.* 2019). Latitudinal sampling was also poor for most taxonomic groups. For

the one exception—birds—we repeated analyses after restricting our dataset to only bird

studies (6 from tropics, 12 from temperate zone). To investigate whether our results were

driven by the inclusion of communities with few species, we fit an additional model

including data only from communities with ≥10 species (125 estimates of temperature

tracking from 70 communities).

Last, to assess latitudinal variation in absolute response to recent warming, we

repeated analyses with elevational shift (m/decade) as the response variable instead of

temperature tracking.

Forest Inventory Data

We compiled our forest inventory plot dataset using Forest Inventory and Analysis (FIA)

plots from the United States and previously published inventory plot data from Central

and South America (Feeley *et al.* 2013; Fadrique *et al.* 2018). We filtered for FIA plots

that were fully forested and have not received observable treatment, including cutting,

site preparation, artificial regeneration, natural regeneration, and other silvicultural

treatment (Smith 2002). We also filtered for FIA plots that have not experienced

disturbances in at least five years. Because we were interested in changes in just

mountain forests, we selected FIA plots that fell within mountainous areas using the

global mountains raster map derived from the 250 m global Hammond landforms product

(Karagulle *et al.* 2017). We included only FIA plots that had been surveyed at least twice.

We collected the species identity of all individual adult trees (diameter at breast height >

12.7 cm, or 5 inches) for each survey of each FIA plot. We then calculated the

community temperature index (CTI) of the plot as the average of optimal temperature of

each species weighted by basal area, following the method of Fadrique et al. (Fadrique *et*

al. 2018). Specifically, we downloaded all georeferenced plant location records available

through the BIEN database (version 4.1.1 accessed in November 2018 via BIEN package

in R) for the New World (North America, Central America, and South America, but

excluding the Caribbean islands). The BIEN database provides collated observation and

collection data from multiple sources and provides a base level of data filtering and

standardization. We used BIEN's default download preferences to exclude records of

known introduced species and cultivated individuals. We further filtered the records to

include only those that were georeferenced and that list the year of collection/observation

as being between 1970 and 1980. We restricted records to just this 10-year window to

minimize errors due to the possibility of species changing their ranges through time. For

each species, we then extracted the estimated mean annual temperature (BIOCLIM1) at

all retained collection coordinates from the CHELSA v1.2 raster of "current" (i.e., mean

of 1979-2012) climate at 30 arc-second resolution and estimated the thermal optima

(MAT_{opt}) of each species as the mean MAT of all collections. If species had fewer than

10 records, we estimated MAT_{opt} at the genus-level using combined records of all

congeners. Genera with <10 records were excluded from subsequent analyses. We

additionally ran analyses using only species-level climatic optima measurements. We

next used the collection records to calculate the community temperature index (CTI) of

each plot in each census as the mean MAT_{opt} for species in the plot weighted by their

relative basal area.

We then calculated the thermophilization rate of each FIA plot as the linear trend

of CTI over time, and combined this dataset with previously published thermophilization

rates for 10 sites in Costa Rica and 63 sites in the tropical Andes (Feeley *et al.* 2013;

Fadrique *et al.* 2018).

For each plot, we calculated the rate of temperature change as the linear trend of

mean annual temperature over time, using monthly mean temperature estimates from

1980 to 2013 from the CHELSA time-series dataset (Karger & Zimmermann 2019). As

with the resurvey data, we calculated the thermal tracking score for each plot as the ratio

of observed changes (rate of change in thermophilization rate) and expected changes (rate

of change of mean annual temperature).

In order to focus on the response of organisms to the warming climate, we

selected plots with significant warming trends (plots with $p < 0.05$ in a regression of

mean temperature versus year). As FIA plots are generally smaller in size compared to

the plots in Central and South America (168 m^2 for FIA plots compared to 1-hectare for

most tropical plots), we aggregated FIA plots into 1 degree diameter hexagons that

contain approximately 30 plots each, averaging the thermophilization rate and

temperature tracking score weighted by total basal area. We removed hexagons with <5

plots from subsequent analysis. We examined the relationships between latitude and the

temperature tracking score by fitting linear mixed-effects models using the "nlme"

package in R (Pinheiro *et al.* 2017). We included both latitude (either the factor

"tropical/temperate", or absolute latitude) and elevation as fixed effects. Due to the close

proximity of hexagons/plots and spatial dependence in the residuals of non-spatial linear

models, we modeled spatial random effects that follow Gaussian covariance functions,

and then evaluated the significance of regression coefficients for fixed effects using the

conditional standard error of regression coefficients.

Results

Resurvey dataset

Resurvey data show that tropical montane taxa are tracking temperature increases

"better" (i.e., have higher temperature tracking scores) than are temperate montane

species (Fig. 2a-2d, Fig. S3). Tropical communities had temperature tracking scores 2.5

times greater than temperate communities (Fig. 2a, Table S2; temperature tracking scores

for tropical and temperate communities = 0.81 ± 0.16 vs. 0.32 ± 0.077, $p = 0.0058$;

estimates $\pm$ standard errors from mixed-effects models), and tropical species had

temperature tracking scores 2.2 times greater than temperate species (Fig, 2c, Table S3;

temperature tracking scores for tropical and temperate species = 0.86 ± 0.18 vs. $0.40 \pm$

0.085, $p = 0.018$).

Results were similar when we modeled temperature tracking from resurvey data

as a continuous function of position along the latitudinal gradient. Temperature tracking

scores decreased by an average of 0.15 ± 0.050 and 0.12 ± 0.054 per 10^{o} increase in

latitude for communities and species, respectively (Figs. 2b and 2d, Tables S4 and S5).

For resurvey data, estimates from linear mixed models are that communities at the

equator have a tracking score of 0.97 ± 0.21 but 0.32 ± 0.077 at 45° latitude (averaged

over levels of fixed effects; the equivalent values for the species-level model are $0.94 \pm$

0.23 and 0.39 ± 0.88). Hence, tropical communities and species are closely tracking

temperature changes while temperate zone communities and species are not, though the

explanatory power of community models was much greater (marginal R^2 values from

linear mixed-effects models = 0.10 and 0.0091 respectively).

Methodological covariates included in models had minimal explanatory power,

with the exception that temperature tracking scores tended to be larger at species' upper

elevational limits compared to at their lower elevational limits or mean elevations (Tables

S2-S5). Results for communities all held when considering only communities with 10 or

more species (Fig. S4, Tables S6 and S7), indicating that the results are not driven by the

inclusion of depauperate communities. In addition, all results held when subsetting the

resurvey dataset to only studies of birds (Fig. S5, Tables S8-S11).

Last, absolute responses to recent climate change tended to be greater in the

tropics, though this difference was not statistically significant (Tables S12-S15). On

average tropical communities have shifted upslope 25.1 meters per decade while

temperate communities have shifted upslope 13.1 meters per decade ($p = 0.093$); the

equivalent estimates for the species-level analysis are 20.8 meters per decade (tropics)

and 13.7 meters per decade (temperate; $p = 0.30$). Estimated rates of upslope shifts from

models that did not include latitude were 18.6 meters per decade (communities) and 14.5

meters per decade (species-level).

Forest inventory dataset

Tropical montane trees are also tracking temperature increases better than temperate

montane trees (Fig. 2e and 2f, Fig S6, Tables S16-S17). Tropical montane forest plots

had higher temperature tracking scores (0.44 ± 0.11) than temperate montane forest plots

(0.044 ± 0.044; $p = 0.00099$; Table S16). Latitude is a strong predictor in the linear mixed

model: a 10^{o} increase in latitude corresponds to a 0.13 ± 0.039 decrease in the

temperature tracking score (Table S17). The model-based estimate is that tree plots have

a tracking score of 0.56 ± 0.0043 at the equator, but -0.038 ± 0.0043 at 45° latitude.

Results were unchanged when using only species-level climatic optima data (Tables S18,

S19).

The greater temperature tracking in tropical inventory plots was due to faster

changes in CTI in the tropics, also known as the thermophilization rate. Tropical plots

had thermophilization rates of 0.095 ± 0.020 °C per decade vs. 0.017 ± 0.012 °C per

decade for temperate plots (Fig. S6). Thus, while the rate of warming was faster at high

latitudes in our dataset-- 0.31 ± 0.0065 °C per decade for temperate plots vs. 0.22 ± 0.010

°C per decade for tropical plots (Fig. S6), there were still greater absolute responses to

recent warming in the tropics.

Discussion

We found evidence that tropical montane species and communities are tracking recent

temperature increases better than temperate montane species. Temperature tracking was

greater in the tropics in each of three datasets: (1) community-level data from resurvey

studies, (2) species-level data from resurvey studies, and (3) repeatedly-censused forest

inventory tree plots. On average, tropical montane species track temperature increases,

while temperate montane species do not. This tighter temperature tracking in the tropics

means that absolute responses to recent warming are also greater in the tropics, despite

faster rates of warming in the temperate zone.

The overall rate of upslope shift for communities that we document of 18.6 m per

decade is 67% higher than the average overall shift for communities of 11.1 m per decade

reported nearly a decade ago (Chen et al., 2011), which was itself double the estimate of

6.1 m per decade reported nearly two decades ago (Parmesan & Yohe 2003). Hence, as

temperatures continue to warm and more resurvey datasets are published (e.g. 83

communities in the present study versus 30 in the Chen et al. 2011 study), estimated rates

of upslope shift continue to increase, counter to the more common pattern wherein effect

sizes tend to decline over time (Camerer *et al.* 2018). Further, we find evidence that rates

of upslope shift depend in part on latitude, with average upslope shifts for communities

of 25.1 m per decade in the tropics but only 13.1 m per decade in the temperate zone,

though this difference is marginally significant.

Our resurvey results appear to be robust to the heterogeneity present within our

datasets. Despite nearly two decades of research documenting elevational range shifts

associated with recent warming, the number of studies for most taxonomic groups

remains low, particularly in the tropics (Feeley *et al.* 2017). Birds are the only taxonomic

group with reasonable sampling across temperate and tropical zones, and we find strong

temperature tracking in tropical—but not temperate—birds. This means that we have

evidence for a latitudinal gradient in temperature tracking in both birds and trees

(comparing tropical birds to temperate birds, and tropical trees to temperate trees), while

data is lacking for other individual groups.

Resurvey and forest inventory plot datasets show marked latitudinal patterns in

temperature tracking, but differences in temperature tracking scores across latitudes were

higher for the resurvey dataset. This difference could reflect a biological difference in

generation time between species included in the datasets. Trees have long generation

times that lead to slow rates of community turnover and range shifts (Lenoir *et al.* 2008;

Feeley *et al.* 2012). In comparison, the resurvey dataset consists primarily of taxa with

much shorter generation times such as birds, mammals, and herbaceous plants. An

alternative explanation is that methodological differences between resurveys and forest

tree inventory plots, both in data collection and analysis, explain why temperature

tracking scores are higher in the resurvey dataset.

Latitude is a much better predictor of temperature tracking when considering

communities (a set of species aggregated together) than for individual species. Individual

species show a wide variation in their temperature tracking scores (Freeman et al., 2018;

Moritz et al., 2008; Rumpf et al., 2019). While latitude is a statistically significant

predictor of temperature tracking in the species-level model, its explanatory power is low.

Combined, these results imply that despite a clear latitudinal pattern of temperature

tracking for communities, we still have only a limited ability to predict changes in

individual species' elevational distributions associated with warming temperatures

(Angert *et al.* 2011). Range shifts at the species-level may be more predictable in the

marine realm, where most species appear to closely track temperature (Lenoir et al.,

2020; Pinsky, Eikeset, McCauley, Payne, & Sunday, 2019).

Multiple mechanisms may explain why tropical species are tracking temperature

changes better than temperate species. The leading explanation is that tropical species are

more physiologically sensitive to climate change than are temperate species. This

explanation implies that the mean annual temperature is a more important driver of

species' elevational distributions in the tropics than at higher latitudes. An additional

possibility is that both tropical and temperate species are tracking recent warming, but

that temperate species are using phenological shifts to do so (Socolar *et al.* 2017).

Seasonal temperature fluctuations in the tropics are minimal, meaning that tropical

species are unlikely to be able to track climate via phenological shifts. In other words,

temperate-zone species may track changing climate by shifting in time, while tropical

species track changing climate by shifting in space. If so, we may expect a decoupling of

biodiversity change and carbon dynamics in the temperate zone.

Several limitations of our study deserve explicit mention. First, we followed

previous analyses in calculating temperature tracking scores based on mean annual

temperature (Chen et al., 2011). Analyses that incorporate temperature variability (i.e.,

seasonality) and other climatic variables have also proven powerful (Crimmins *et al.*

2011; Tingley *et al.* 2012), but are inherently more difficult to implement and interpret.

Second, our analyses do not take into account variation in microclimate, which may be a

strong driver of range shifts or the lack thereof (Lembrechts *et al.* 2019; Zellweger *et al.*

2020). It is not clear how microclimate availability varies along a latitudinal gradient, but

a greater availability of microclimates and climate refugia in the temperate zone than the

tropics is an alternative explanation for our results. We were similarly unable to analyze

climatic factors that occur at intermediate spatial scales along mountain slopes, such as

cold-air pooling (Curtis *et al.* 2014). Third, future studies should address landscape-level

changes due to habitat loss or other disturbances (Larsen 2012; Lenoir & Svenning 2015;

Campos-Cerqueira *et al.* 2017; Guo *et al.* 2018). The resurveys and forest plots in our

dataset took place in landscapes that have not undergone intensive deforestation or other

land-use change. Given that highly modified landscapes predominate across most of the

globe, further tests of the interactions between landscape change and climate change are

needed. Fourth, the patterns we document are not without exceptions. For example,

Puerto Rican frogs and birds are not closely tracking temperature despite their tropical

latitude (Campos-Cerqueira & Mitchell Aide 2017; Campos-Cerqueira *et al.* 2017),

though this could potentially reflect increasing forest cover on this island (Battey *et al.*

2019). Conversely, some temperate zone communities are closely tracking recent

warming (Kelly & Goulden 2008; Menéndez *et al.* 2014).

Conclusions

The latitudinal gradient in temperature tracking we document has multiple implications

for the conservation of montane floras and faunas. The weak temperature tracking of

temperate montane species and communities indicates that acclimation and adaptation,

rather than elevational shifts, will likely be key processes in determining if temperate

montane species can persist in the face of continued warming. It is an open question

whether adaptation and acclimation will be able to keep pace with rates of warming that

are unprecedented in recent evolutionary time (Visser 2008; Feeley *et al.* 2012). In

contrast, the strong temperature tracking of tropical montane species indicates that they

may be able to persist despite warming temperatures by shifting their distributions

upslope, at least when protected elevational corridors provide suitable habitats at higher

elevations. On the other hand, this implies that the "escalator to extinction"—wherein

montane species respond to warming temperatures by shifting upslope until they run out

of available habitat on mountaintops (Freeman, Scholer, Ruiz-Gutierrez, & Fitzpatrick,

2018)—runs faster in the tropics. Notably, such mountaintop extinctions may occur well

below the actual mountaintop, as pervasive anthropogenic modifications of high-

elevation tropical systems effectively limit the ability of the tree line to shift upslope

(Rehm & Feeley 2015). Indeed, local extinctions and range contractions associated with

recent warming appear to be most common in tropical montane species (Freeman et al.,

2018; Wiens, 2016).

The sixth mass extinction in Earth's history is now underway (Ceballos *et al.*

2017). The tropics have the highest species diversity of any biome, and tropical

mountains have the highest diversity of all (Rahbek *et al.* 2019). The relatively small

temperature changes in the tropics should minimize the impact of climate change, but the

disproportionately sensitive responses of tropical montane species have instead placed

this whole biota on an escalator to extinction. The degree to which predictions of

widespread species extinctions in tropical mountains (Sekercioglu *et al.* 2008) come true

will depend on our ability to protect elevational corridors that enable species to persist

while shifting upslope (Feeley & Rehm 2012), and, ultimately, on whether humanity is

able to slow global warming.

Acknowledgements

We thank the many scientists who have published resurvey data and measured forest

inventory plots. The Feeley lab, Schluter lab, R. Yorque, J. Lembrechts and two

anonymous reviewers provided useful comments that improved this manuscript. BGF

was supported from postdoctoral fellowships from Banting Canada (379958), and the

Biodiversity Research Centre, and YS was supported by a Regents' fellowship from the

University of California, Santa Cruz. KZ and KJF are supported by the US National

Science Foundation (NSF grants DEB 1926438 to KZ and DEB-1350125 to KJF).

References

Angert, A.L., Crozier, L.G., Rissler, L.J., Gilman, S.E., Tewksbury, J.J. & Chunco, A.J.

(2011). Do species' traits predict recent shifts at expanding range edges? *Ecol. Lett.*,

14, 677–689.

Bates, D., Maechler, M., Bolker, B. & Walker, S. (2014). lme4: Linear mixed-effects

models using Eigen and S4.

Battey, C.J., Otero, L.M., Gorman, G.C., Hertz, P.E., Lister, B.C., García, A., *et al.*

(2019). Why Montane Anolis Lizards are Moving Downhill While Puerto Rico

Warms. *bioRxiv*, 751941.

Camerer, C.F., Dreber, A., Holzmeister, F., Ho, T.H., Huber, J., Johannesson, M., *et al.*

(2018). Evaluating the replicability of social science experiments in Nature and

Science between 2010 and 2015. *Nat. Hum. Behav.*, 2, 637–644.

Campos-Cerqueira, M., Arendt, W.J., Wunderle, J.M. & Aide, T.M. (2017). Have bird

distributions shifted along an elevational gradient on a tropical mountain? *Ecol.*

Evol., 7, 9914–9924.

Campos-Cerqueira, M. & Mitchell Aide, T. (2017). Lowland extirpation of anuran

populations on a tropical mountain. *PeerJ*, 2017.

Ceballos, G., Ehrlich, P.R. & Dirzo, R. (2017). Biological annihilation via the ongoing

sixth mass extinction signaled by vertebrate population losses and declines. *Proc.*

Natl. Acad. Sci. U. S. A., 114, E6089–E6096.

Chen, I.-C., Hill, J.K., Ohlemüller, R., Roy, D.B. & Thomas, C.D. (2011). Rapid range

shifts of species associated with high levels of climate warming. *Science*, 333,

1024–1026.

Chen, I.C., Shiu, H.J., Benedick, S., Holloway, J.D., Chey, V.K., Barlow, H.S., *et al.*

(2009). Elevation increases in moth assemblages over 42 years on a tropical

mountain. *Proc. Natl. Acad. Sci.*, 106, 1479–1483.

Crimmins, S., Dobrowski, S., Greenberg, J., Abatzoglou, J. & Mynsberge, A. (2011).

Changes in climatic water balance drive downhill shifts in plant species' optimum

elevations. *Science*, 331, 324–327.

Curtis, J.A., Flint, L.E., Flint, A.L., Lundquist, J.D., Hudgens, B., Boydston, E.E., *et al.*

(2014). Incorporating cold-air pooling into downscaled climate models increases

potential refugia for snow-dependent species within the Sierra Nevada Ecoregion,

CA. *PLoS One*, 9.

DeLuca, W. V. & King, D.I. (2017). Montane birds shift downslope despite recent

warming in the northern Appalachian Mountains. *J. Ornithol.*, 158, 493–505.

Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak,

D.C., *et al.* (2008). Impacts of climate warming on terrestrial ectotherms across

latitude. *Proc. Natl. Acad. Sci.*, 105, 6668–6672.

Fadrique, B., Báez, S., Duque, Á., Malizia, A., Blundo, C., Carilla, J., *et al.* (2018).

Widespread but heterogeneous responses of Andean forests to climate change.

Nature, 564, 207–212.

Feeley, K.J., Hurtado, J., Saatchi, S., Silman, M.R. & Clark, D.B. (2013). Compositional

shifts in Costa Rican forests due to climate-driven species migrations. *Glob. Chang.*

Biol., 19, 3472–3480.

Feeley, K.J. & Rehm, E.M. (2012). Amazon's vulnerability to climate change heightened

by deforestation and man-made dispersal barriers. *Glob. Chang. Biol.*, 18, 3606–

3614.

Feeley, K.J., Rehm, E.M. & Machovina, B. (2012). The responses of tropical forest

species to global climate change: acclimate, adapt, migrate, or go extinct? *Front.*

Biogeogr., 4, 69–84.

Feeley, K.J., Stroud, J.T. & Perez, T.M. (2017). Most 'global' reviews of species'

responses to climate change are not truly global. *Divers. Distrib.*, 23, 231–234.

Freeman, B.G. & Class Freeman, A.M. (2014). Rapid upslope shifts in New Guinean

birds illustrate strong distributional responses of tropical montane species to global

warming. *Proc. Natl. Acad. Sci.*, 111, 4490–4494.

Freeman, B.G., Lee-Yaw, J.A., Sunday, J.M. & Hargreaves, A.L. (2018a). Expanding,

shifting and shrinking: The impact of global warming on species' elevational

distributions. *Glob. Ecol. Biogeogr.*, 27, 1268–1276.

Freeman, B.G., Scholer, M.N., Ruiz-Gutierrez, V. & Fitzpatrick, J.W. (2018b). Climate

change causes upslope shifts and mountaintop extirpations in a tropical bird

community. *Proc. Natl. Acad. Sci. U. S. A.*, 115, 11982–11987.

Guo, F., Lenoir, J. & Bonebrake, T.C. (2018). Land-use change interacts with climate to

determine elevational species redistribution. *Nat. Commun.*, 9, 1-7.

Harris, I., Jones, P.D., Osborn, T.J. & Lister, D.H. (2013). Updated high-resolution grids

of monthly climatic observations–the CRU TS3. 10 Dataset. *Int. J. Climatol.*, 34,

623-642.

Karagulle, D., Frye, C., Sayre, R., Breyer, S., Aniello, P., Vaughan, R., *et al.* (2017).

Modeling global Hammond landform regions from 250-m elevation data. *Trans.*

GIS, 21, 1040–1060.

Karger, D.N. & Zimmermann, N.E. (2019). Climatologies at High resolution for the

Earth Land Surface Areas CHELSA V1 . 2: Technical specification. *Sci. Data*, 4,

170122.

Kelly, A.E. & Goulden, M.L. (2008). Rapid shifts in plant distribution with recent

climate change. *Proc. Natl. Acad. Sci.*, 105, 11823–11826.

Kusrini, M.D., Lubis, M.I., Endarwin, W., Yazid, M., Darmawan, B., Ul-Hasanah, A.U.,

et al. (2017). Elevation range shift after 40 years: The amphibians of Mount Gede

Pangrango National Park revisited. *Biol. Conserv.*, 206, 75–84.

Larsen, T.H. (2012). Upslope range shifts of Andean dung beetles in response to

deforestation: compounding and confounding effects of microclimatic change.

Biotropica, 44, 82–89.

Laurance, W.F., Carolina Useche, D., Shoo, L.P., Herzog, S.K., Kessler, M., Escobar, F.,

et al. (2011). Global warming, elevational ranges and the vulnerability of tropical

biota. *Biol. Conserv.*, 144, 548–557.

Lembrechts, J.J., Nijs, I. & Lenoir, J. (2019). Incorporating microclimate into species

distribution models. *Ecography*, 42, 1267–1279.

Lenoir, J., Bertrand, R., Comte, L., Bourgeaud, L., Hattab, T., Murienne, J., *et al.* (2020).

Species better track climate warming in the oceans than on land. *Nature Ecol. &*

Evol., 1-16.

Lenoir, J., Gegout, J.C., Marquet, P.A., De Ruffray, P. & Brisse, H. (2008). A significant

upward shift in plant species optimum elevation during the 20th century. *Science*,

320, 1768–1771.

Lenoir, J. & Svenning, J.C. (2015). Climate-related range shifts - a global

multidimensional synthesis and new research directions. *Ecography*, 38, 15–28.

Menéndez, R., González-Megías, A., Jay-Robert, P. & Marquéz-Ferrando, R. (2014).
Climate change and elevational range shifts: evidence from dung beetles in two
European mountain ranges. *Glob. Ecol. Biogeogr.*, 23, 646–657.

Moret, P., Aráuz, M. de los Á., Gobbi, M. & Barragán, Á. (2016). Climate warming
effects in the tropical Andes: first evidence for upslope shifts of Carabidae
(Coleoptera) in Ecuador. *Insect Conserv. Divers.*, 9, 342–350.

Moritz, C., Patton, J.L., Conroy, C.J., Parra, J.L., White, G.C. & Beissinger, S.R. (2008).
Impact of a century of climate change on small-mammal communities in Yosemite
National Park, USA. *Science*, 322, 261–264.

Moskwik, M. (2014). Recent elevational range expansions in plethodontid salamanders
(Amphibia: Plethodontidae) in the southern appalachian mountains. *J. Biogeogr.*, 41,
1957–1966.

Parmesan, C. & Yohe, G. (2003). A globally coherent fingerprint of climate change
impacts across natural systems. *Nature*, 421, 37–42.

Perez, T.M., Stroud, J.T. & Feeley, K.J. (2016). Thermal trouble in the tropics. *Science*,
351, 1392–1393.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Heisterkamp, S., Van Willigen, B., *et al.*
(2017). Nlme: Linear and Nonlinear Mixed Effects Models, version 3(1).

Pinsky, M.L., Eikeset, A.M., McCauley, D.J., Payne, J.L. & Sunday, J.M. (2019). Greater
vulnerability to warming of marine versus terrestrial ectotherms. *Nature*, 569, 108–
111.

R Development Core Team. (2020). R: A language and environment for statistical

531  computing.

532 Rahbek, C., Borregaard, M.K., Antonelli, A., Colwell, R.K., Holt, B.G., Nogues-Bravo,

533  D., *et al.* (2019). Building mountain biodiversity: Geological and evolutionary

534  processes. *Science*, 365, 1114–1119.

535 Rehm, E.M. & Feeley, K.J. (2015). The inability of tropical cloud forest species to invade

536  grasslands above treeline during climate change: Potential explanations and

537  consequences. *Ecography*, 38, 1167–1175.

538 Rohatgi, A. (2017). WebPlotDigitizer 3.11.

539 Rumpf, S.B., Hülber, K., Klonner, G., Moser, D., Schütz, M., Wessely, J., *et al.* (2018).

540  Range dynamics of mountain plants decrease with elevation. *Proc. Natl. Acad. Sci.*,

541  201713936.

542 Rumpf, S.B., Hülber, K., Zimmermann, N.E. & Dullinger, S. (2019). Elevational rear

543  edges shifted at least as much as leading edges over the last century. *Glob. Ecol.*

544  *Biogeogr.*, 28, 533–543.

545 Sekercioglu, C.H., Schneider, S.H., Fay, J.P. & Loarie, S.R. (2008). Climate change,

546  elevational range shifts, and bird extinctions. *Conserv. Biol.*, 22, 140–150.

547 Sheldon, K.S. (2019). Climate Change in the Tropics: Ecological and Evolutionary

548  Responses at Low Latitudes. *Annu. Rev. Ecol. Evol. Syst.*, 50, 303-333.

549 Smith, W.B. (2002). Forest inventory and analysis: A national inventory and monitoring

550  program. *Environ. Pollut.*, 116, 233–242.

551 Socolar, J.B., Epanchin, P.N., Beissinger, S.R. & Tingley, M.W. (2017). Phenological

552  shifts conserve thermal niches in North American birds and reshape expectations for

553  climate-driven range shifts. *Proc. Natl. Acad. Sci.*, 201705897.

Sunday, J.M., Bates, A.E. & Dulvy, N.K. (2012). Thermal tolerance and the global

redistribution of animals. *Nat. Clim. Chang.*, 2, 686–690.

Tingley, M.W., Koo, M.S., Moritz, C., Rush, A.C. & Beissinger, S.R. (2012). The push

and pull of climate change causes heterogeneous shifts in avian elevational ranges.

Glob. Chang. Biol., 18, 3279–3290.

Visser, M.E. (2008). Keeping up with a warming world; assessing the rate of adaptation

to climate change. *Proc. R. Soc. B Biol. Sci.*, 275, 649.

Wiens, J.J. (2016). Climate-Related Local Extinctions Are Already Widespread among

Plant and Animal Species. *PLoS Biol.*, 14, 1–18.

Wu, C., Holloway, J.D., Hill, J.K., Ho, C., Thomas, C.D. & Chen, I. (2019). Reduced

body sizes in climate-impacted Borneo moth assemblages are primarily explained by

range shifts. *Nat. Commun.*, 1–7.

Zellweger, F., De Frenne, P., Lenoir, J., Vangansbeke, P., Verheyen, K., Bernhardt-

römermann, M., *et al.* (2020). Forest microclimate dynamics drive plant responses to

warming. *Science*, 368, 772–775.

Zuckerberg, B., Woods, A.M. & Porter, W.F. (2009). Poleward shifts in breeding bird

distributions in New York State. *Glob. Chang. Biol.*, 15, 1866–1883.

573
574 **Figure 1.** Maps of resurvey studies that measured elevational shifts associated with

575 recent warming (a) and of forest inventory tree plots that have been repeatedly censused

576 (b). Locations of resurvey studies are jittered slightly to improve clarity. The Tropics of

577 Cancer and Capricorn (at 23.4° N and S, respectively) delimit the tropics, and are

578 illustrated with dashed lines.

Figure 2. Tropical communities, species, and forest inventory tree plots have higher

temperature tracking scores than their temperate zone counterparts. Raw data are shown

as points. Dashed lines illustrate both perfect temperature tracking (temperature tracking

= 1) and zero temperature tracking despite warming temperatures (temperature tracking =

0). Trendlines illustrate predictions from mixed models. (a) temperature tracking for

communities in tropical and temperate zones; (b) relationship between temperature

tracking and absolute latitude (°) for communities; (c) temperature tracking for species in

tropical and temperate zones (~ 0.5% of data has been removed for visualization); (d)

relationship between temperature tracking and absolute latitude (°) for species; (e)

temperature tracking for forest inventory tree plots in tropical and temperate zones; (f)

relationship between temperature tracking and absolute latitude (°) for forest inventory

tree plots.

Figure 3. Tropical communities and species have undertaken larger upslope shifts, and

forest inventory tree plots greater thermophilization, than their temperate zone

counterparts. Raw data are shown as points. Trendlines illustrate predictions from mixed

models. (a) elevational shifts for communities in tropical and temperate zones; (b)

relationship between elevational shifts and absolute latitude (°) for communities; (c)

elevational shifts for species in tropical and temperate zones (~ 0.5% of data has been

removed for visualization); (d) relationship between elevational shifts and absolute

latitude (°) for species; (e) thermophilization rate for forest inventory tree plots in tropical

and temperate zones; (f) relationship between thermophilization rate and absolute latitude

(°) for forest inventory tree plots.