1 **TITLE**
2 Cell type-specific effects of isoflurane on two distinct afferent inputs to cortical layer 1.
3

4 **AUTHORS AND AFFILIATIONS**
5 Caitlin A. Murphy[1]
6 Matthew I. Banks[1]
7 [1]Department of Anesthesiology, University of Wisconsin, Madison, Wisconsin
8
Corresponding Author:
Caitlin A. Murphy
caitlin.murphy@wisc.edu
Tel: (608) 263-5552

KEYWORDS
anesthesia; isoflurane; electrophysiology; neocortex; thalamus; optogenetics; interneurons;
pyramidal cells; patch-clamp techniques

ABSTRACT
While their behavioral effects are well-characterized, the mechanisms by which anesthetics
induce loss of consciousness are largely unknown. Anesthetics may disrupt integration and
propagation of information in corticothalamic networks. Recent studies have shown that
isoflurane diminishes synaptic responses of thalamocortical (TC) and corticocortical (CC)
afferents in a pathway-specific manner. However, whether the synaptic effects of isoflurane
observed in extracellular recordings persist at the cellular level has yet to be explored. Here,
we activate TC and CC layer 1 inputs in non-primary mouse neocortex in ex vivo brain slices
and explore the degree to which isoflurane modulates synaptic responses in pyramidal cells
and in two inhibitory cell populations, somatostatin-positive (SOM+) and parvalbumin-positive
(PV+) interneurons. We show that the effects of isoflurane on synaptic responses and intrinsic
properties of these cells varies among cell type and by cortical layer. Layer 1 inputs to L4
pyramidal cells were suppressed by isoflurane at both TC and CC synapses, while those to
L2/3 pyramidal cells and PV+ interneurons were not. TC inputs to SOM+ cells were rarely
observed at all, while CC inputs to SOM+ interneurons were robustly suppressed by isoflurane.
These results suggest a mechanism by which isoflurane disrupts integration and propagation
of incoming cortical signals.

INTRODUCTION

Anesthetics may ultimately influence both the level and contents of consciousness via actions

on corticothalamic circuits, disrupting integration of information throughout the cortical

hierarchy (Crick and Koch, 2003; Dehaene and Changeux, 2011; Koch et al., 2016). Activity in

higher order cortical areas is particularly sensitive to anesthetics (Nourski et al., 2017; Nourski

et al., 2018), as is corticocortical (CC) feedback connectivity (Ferrarelli et al., 2010; Boly et al.,

2012; Liu et al., 2012; Raz et al., 2014; Mashour and Hudetz, 2017; Murphy et al., 2019). In

addition to cortical effects, higher order thalamocortical (TC) connectivity is suppressed during

both sleep (Picchioni et al., 2014) and anesthesia (Langsjo et al., 2012; Liu et al., 2013; Akeju

et al., 2014). Activation of non-specific thalamic nuclei during anesthetic-induced

unconsciousness (Redinbaugh et al., 2020) and non-REM sleep (Honjoh et al., 2018)

promotes behavioral arousal, implicating higher order thalamic centers as enablers of

consciousness.

Recent studies from our lab have shown that in *ex vivo* slices, isoflurane diminishes

synaptic responses in a pathway-specific manner (Raz et al., 2014; Murphy et al., 2019).

Despite overlapping terminal fields in layer 1, CC feedback afferents to a non-primary cortical

area were preferentially suppressed by isoflurane compared to non-primary TC inputs.

However, extracellular recordings used in these investigations capture summed effects across

all synapses, leaving open questions as to whether observed effects are consistent across

post-synaptic targets. For example, differential effects on inhibitory interneurons compared to

excitatory cells may shift the balance of excitation and inhibition (Haider et al., 2013; Xue et al.,

2014), disrupting integration of inputs in pyramidal cell dendrites or constraining the output of

the cortical column. Superficial and deep layers may operate within functionally separate

domains, where cells in superficial layers primarily carry out intracortical interactions, and

those in deep layers facilitate interactions with lower cortical and subcortical targets in the

feedback direction (Douglas et al., 1989; Douglas and Martin, 2004; Douglas and Martin, 2007; Binzegger et al., 2009). Differential effects of anesthetics on cells residing in different layers may provide insight into the functional roles of discrete cortical networks and the mechanisms that subserve the breakdown of connectivity observed during anesthesia.

Here, we investigate the effect of isoflurane on independently activated TC and CC inputs to layer 1 of non-primary neocortex in brain slices. We evaluate the pathway- and cell type-specific effects of isoflurane on synaptic responses recorded in pyramidal cells in layers 2-5 and in two subpopulations of inhibitory cells, SOM+ and PV+ interneurons.

METHODS

To express the fluorescent reporter tdTomato in non-overlapping populations of GABAergic interneurons, homozyogous, Cre-dependent tdTomato male mice were bred with either SOM-Cre female or PV-Cre female mice; heterozygous offspring (both sexes) were used for all experiments. Viral constructs containing channelrhodopsin-2 (ChR2) and fluorescent reporter YFP (AAV2-hSyn-hChR2(H134R)-eYFP; provided by K. Deisseroth, Stanford University via University of North Carolina Vector Core) were injected into either posterior medial thalamus (AP: -2.25 mm, ML: 1.25 mm, DV: -3.35 mm; volume: 1 μL) or cingulate cortex (AP: 0.2 mm, ML: 0.3 mm, DV: 0.9 mm; volume: 1 μL) to independently activate non-primary TC and CC feedback afferents, respectively.

After allowing at least 3 weeks for expression of the virus, acute coronal brain slices containing were collected and recordings were made from medial secondary visual cortex. Intracellular whole cell patch clamp recordings were conducted in current clamp mode from pyramidal cells in layers 2-5, SOM+ interneurons, or PV+ interneurons; extracellular multichannel recordings were collected concurrently. Cell types were identified either by Cre-

dependent expression of td-Tomato reporter (for SOM+ and PV+ interneurons) or by

morphology (pyramidal cells). To further guide identification of cell types, each cell's response

to 600 msec current steps was evaluated as previously described (Krause et al., 2017). All

signals were low-pass filtered at 10 kHz and digitized at 40 kHz.

To activate either TC or CC afferents, blue light was delivered exclusively to layer 1 (150

μm diameter, 470 nm; Polygon400, Mightex Systems, Toronto, Ontario). For each trial,

synaptic responses were evoked using four brief pulses of light (2 msec; 10 Hz) at one of at

least five light intensities randomly selected from a range of intensities (0.022-2.2 mW). Each

trial was separated by ten seconds, and at least eight trials were conducted for each light

intensity, randomly interleaved.

Signals from each of 16 extracellular channels were filtered (bandpass filter, 1-300Hz) to

yield local field potentials used to calculate the current source density (CSD; (Mitzdorf, 1985))

with the spline inverse method (Pettersen et al., 2006). The signal in layer 1 with the shortest

latency current sink (inward transmembrane current) was used for further analysis. Intracellular

signals were filtered (bandpass filter, 0.1-1000 Hz), and responses were averaged across trials

of the same stimulus intensity. Responses to a single stimulus intensity were selected for

detailed analysis; the intensity selected was the minimum light intensity for which at least 50%

of trials (8-10 trials per intensity) evoked a response with <1 msec latency jitter.

Response amplitude was defined as the peak of the signal within 50 msec of the stimulus

(first pulse only). Paired pulse ratio was computed for intracellular recordings, calculated by

dividing the post-synaptic potential (PSP) amplitude of the second pulse by the PSP amplitude

of first pulse. TC- and CC-evoked responses were compared across control, isoflurane (0.24-

0.29 mM), and recovery conditions. To exclude the possibility that effects of isoflurane were

due to response degradation over time, the average of control and recovery measures were

used as a baseline for comparison to isoflurane.

A linear mixed effects model was used to evaluate the pathway- and cell type-dependent

effects of isoflurane on intrinsic and evoked response properties of individual cells. Fixed

effects were cell type (L2/3 Pyr, L4 Pyr, L5 Pyr, SOM+, or PV+) and drug condition (baseline or

isoflurane), as well as afferent pathway (TC or CC) for evoked responses. Interaction terms

between isoflurane and each of the other fixed effects were also included, as well as a three-

way interaction term representing interactions between cell type and pathway of drug effects;

random effects were slice experiment and cell with random slopes for drug condition. For PSP

amplitude and input resistance measures, heteroscedasticity was corrected by log

transforming response variables, and effects are reported as percent changes. To allow for

direct comparison among cell types and pathways, all isoflurane effects are reported relative to

baseline.

RESULTS

Activation of TC and CC inputs to layer 1 elicit short-latency synaptic responses

Synaptic responses from TC and CC afferents were elicited by activating channelrhodopsin in

afferent axon terminals in layer 1. Brief pulses of light evoked short latency current sinks in

layer 1 (Table 1). No effect of afferent pathway was observed for baseline amplitude ($t(29)$ =

0.41, p=0.69) or latency ($t(29)$ = 0.26, P = 0.80) of layer 1 current sinks. Evoked currents in

layer 1 precipitated PSPs in pyramidal cells in layers 2-5 (Table 2; Figure 2B,D), and in SOM+

and PV+ cells (Table 2; Figure 4B,D,F).

Main effects of pathway and cell type on amplitudes of synaptic responses were tested.

A 2-way ANOVA was not significant for effects of pathway ($F(1,47)$ = 0.07, P=0.79) nor target

cell population ($F(4,47)$ = 1.68, P=0.17), suggesting no differences in baseline (i.e., in the

absence of isoflurane) amplitudes among recorded cells. Similarly, no significant effect on

latency was observed for main effects of pathway ($F_{(1,47)}$ = 1.00, P=0.32) nor cell type

($F_{(4,47)}$ =1.68, P=0.17).

Isoflurane suppresses layer 1 current sinks elicited by CC inputs, but not TC inputs

Despite their overlapping terminal fields and similar synaptic dynamics under control

conditions, TC and CC inputs to layer 1 were differentially affected by isoflurane (Figure 1;

Table 1), consistent with previous results from our lab (Murphy et al., 2019). Amplitudes of

layer 1 current sinks were suppressed by isoflurane by 21.8% for CC inputs, while current

sinks evoked by TC inputs were relatively resistant to isoflurane (Figure 1; Table 1).

Isoflurane suppresses synaptic responses in a cell type-dependent manner

Current sinks isolated from current source densities represent averages of inward-going

transmembrane currents across all synapses. However, evidence suggests that the synaptic

properties – and therefore potential functional roles – of TC versus CC inputs to may be cell

type- and layer-dependent (Cruikshank et al., 2012; Sermet et al., 2019). Therefore, we sought

to investigate whether the effects of isoflurane on layer 1 current sinks were uniform for all

post-synaptic targets of each afferent pathway, or whether certain cell populations were

differentially sensitive to isoflurane.

We evaluated the degree to which isoflurane modulated the amplitudes of post-synaptic

potentials elicited by layer 1 afferents in $L^2/_3$, L4, and L5 pyramidal cells (Figure 2) as well as

SOM+ and PV+ cells (Figure 4). Independent of any interactions of cell type or afferent

pathway, isoflurane significantly suppressed evoked post-synaptic potentials of target cells

($\chi_{(1)}$ = 16.6, P<0.0001). Contrary to our findings for extracellularly recorded layer 1 current

sinks, we found no significant interaction of pathway on the effect of isoflurane ($\chi_{(1)}$ = 2.00,

P=0.16). However, the effect of isoflurane was significantly dependent on cell type ($\chi_{(4)}$ =

158 17.862, P=0.0013). The interaction between cell type and pathway was not significant ($\chi(1)$ =

159 6.52, P=0.16). In other words, the effect of isoflurane is cell type-dependent, and within each

160 cell type, no significant differences were observed between afferent pathways.

161   Amplitudes of synaptic responses in L2/3 cells were not significantly affected by

162 isoflurane for either TC or CC inputs (Figure 2B,D; Figure 3; Table 3). Conversely, amplitudes

163 of synaptic responses in L4 cells were significantly suppressed by isoflurane for both TC and

164 CC inputs, to approximately the same degree (Figure 2B,D; Figure 3; Table 3). Isoflurane also

165 suppressed CC inputs, but not TC inputs, to L5 pyramidal cells (Figure 2D; Figure 3; Table 3),

166 though this effect was not significantly different between pathways (P=0.19).

167   In addition to providing input to pyramidal cells, long-range TC and CC projections also

target inhibitory interneurons. We were also interested in the effect of isoflurane on evoked

responses in two subpopulations of inhibitory cells, SOM+ and PV+ interneurons, and whether

these effects were different than those observed for pyramidal cell populations. Consistent with

previous studies (Beierlein et al., 2003; Cruikshank et al., 2010; Audette et al., 2018) and our

own observations, TC inputs rarely elicited synaptic responses in SOM+ cells (n=2 cells); as

such, isoflurane effects on TC inputs to SOM+ cells are not reported. Activation of CC

afferents, however, reliably elicited synaptic responses in SOM+ cells (Figure 4). These

synapses were particularly sensitive to isoflurane; amplitudes of synaptic responses for CC

inputs to SOM+ cells were suppressed 35.8% during isoflurane compared to baseline (Table

3). Conversely, no effect of isoflurane on amplitudes of responses in PV+ cells was detected

for either afferent pathway (Table 3). Thus, even among subpopulations of GABAergic

interneurons, the effect of isoflurane on CC responses was significantly different (-35.8% for

SOM+ cells versus -10.0% for PV+ cells, P=0.014; Figure 5), suggesting distinct roles of each

cell type during anesthesia.

Pre- and post-synaptic effects of isoflurane are cell type-dependent

A multitude of cellular and synaptic dynamics may underlie changes in neural activity observed

during anesthesia. For example, effects of isoflurane on synaptic responses as we describe

here may manifest as a result of pre-synaptic effects of isoflurane (e.g., effects on calcium

influx or vesicle release) (Baumgart et al., 2015), on discrete intrinsic properties of post-

synaptic cells that give rise to changes in their excitability, or a combination of pre- and post-

synaptic effects. Although we cannot ascribe the contributions of specific synaptic or intrinsic

effects of isoflurane to observed effects on evoked responses, characterizing isoflurane-

induced changes in synaptic plasticity or intrinsic properties of post-synaptic cells may provide

insight into the complicated interactions of network components during anesthesia. To this

end, we evaluated the extent to which isoflurane modulated paired pulse ratios and intrinsic

membrane properties (resting membrane potential, input resistance, membrane time constant,

and spike threshold) of each cell. Baseline values for paired pulse ratio and intrinsic membrane

properties are displayed in Table 2 and Table 4, respectively; effects of isoflurane are

described below and in figures and tables as changes relative to baseline.

Isoflurane significantly enhanced the paired pulse ratio of CC inputs to SOM+ cells but

had no effect on paired pulse ratios for other pathway-cell type combinations (Figure 6; Table

5). Intrinsic membrane properties of L4 pyramidal cells were especially sensitive to isoflurane.

In L4 pyramidal cells, input resistance was 13.5% lower and membrane time constants 27.8%

shorter during isoflurane relative to baseline; isoflurane did not affect input resistance nor time

constants of any other cell type (Figure 7). Isoflurane also reduced resting membrane potential

in L4 and L5 pyramidal cells (Figure 7). Effects of isoflurane observed in L4 pyramidal cells

were accompanied by a decrease in spike threshold (Figure 7).

DISCUSSION

In these experiments, we describe the effects of isoflurane on synaptic responses and intrinsic

membrane properties of pyramidal cells in layers 2-5, as well as in SOM+ and PV+

interneurons. Previous investigations of the effects of isoflurane on synaptic responses have

suggested that CC feedback inputs are suppressed to a greater extent than TC and CC

feedforward inputs (Raz et al., 2014; Murphy et al., 2019). Here, we show that suppression of

extracellular synaptic responses is pathway-dependent, recapitulating previous findings using

a similar experimental paradigm (Raz et al., 2014; Murphy et al., 2019). We also demonstrate

that the suppression of synaptic responses by isoflurane is cell type- and layer-dependent,

providing insight into the mesocircuit-level mechanisms underlying the mechanisms of

anesthesia.

Resistance of PV+ interneurons to isoflurane suggests shift in E/I balance

We show that synaptic responses of PV+ interneurons to both TC and CC inputs are resistant

to suppression during isoflurane. TC input to PV+ cells provides stimulus-locked fast

feedforward-inhibition to pyramidal cells, restricting the window of opportunity for generating

spikes and improving spike timing among populations of pyramidal cells (Pouille and

Scanziani, 2001; Krause et al., 2017). We propose that the relative insensitivity of fast-spiking

PV+ neurons to isoflurane we observe here may underlie previous observations from our lab

and others. That is, early responses of the cortical network to TC inputs are relatively

preserved during isoflurane, while recurrent, propagating network activity is highly sensitive

(Hentschke et al., 2017; Murphy et al., 2017). Although we report activity of single cells only in

our study, the relative preservation of PV+ cell responses during isoflurane is likely to shift the

E/I balance in the network toward inhibition. PV+ cells are known to be important for mediating

E/I balance within cortical networks (Dichter and Ayala, 1987; Ferguson and Gao, 2018).

Pyramidal cells in L2/3, for example, receive PV+ cell-mediated inhibitory input proportional to

their excitation under normal conditions, and changes in input to pyramidal cells – as we

observe in L4 cells in our experiments – without commensurate changes in PV+ cells disrupts

this E/I balance (Xue et al., 2014). If these cell type-specific effects of isoflurane occur in

multiple cortical areas, shifts in local E/I balance may compound, contributing to the

progressive failure of information to be passed from one cortical area to the next.

Suppression of L4 pyramidal cells during isoflurane may disrupt feedforward processing

We show that synaptic responses of L4 pyramidal cells are suppressed during isoflurane for

responses evoked by both TC and CC inputs. That TC inputs to any cell type would be

affected by isoflurane is somewhat surprising, and that this effect would be exclusively

observed in L4 pyramidal cells is even more unexpected. L4 pyramidal cells are typically

considered mediators for propagating feedforward information, receiving sensory-specific

information from thalamus and lower areas (Cruikshank et al., 2010; Covic and Sherman,

2011; De Pasquale and Sherman, 2011; Viaene et al., 2011a; Viaene et al., 2011b) and

targeting L2/3 for output to higher areas (Xu et al., 2016; Meyer et al., 2018). Feedforward

circuits are relatively resistant to anesthetics (Lee et al., 2013; Raz et al., 2014; Murphy et al.,

2019) (though, see (Maksimow et al., 2014; Sanders et al., 2018)), and we show here that

extracellular recordings of current sinks evoked by activation of TC afferents were not affected

by isoflurane (Figure 1). Yet, interestingly, our results show that inputs to L4 cells are

preferentially suppressed by isoflurane, suggesting that suppression of TC and CC inputs

together may disrupt feedforward signaling.

While pre-synaptic mechanisms of isoflurane have been well-described (Herring et al.,

2009; Xie et al., 2013; Baumgart et al., 2015), our observations suggest that the mechanisms

underlying isoflurane effects in L4 pyramidal cells were not limited to pre-synaptic effects. No

changes in short-term synaptic plasticity were observed at these synapses (Figure 6), and TC

and CC inputs to L4 pyramidal cells were suppressed to a similar extent (Figure 3). However,

L4 pyramidal cells were significantly leakier during isoflurane, as evidenced by lower input

resistances and shorter membrane time constants relative to baseline (Figure 7). Not only

might these effects dampen voltage changes evoked by synaptic currents, but they are likely to

also contribute to the overall excitability of L4 cells, thereby attenuating their contributions to

processing of cortical signals.

Evidence for disruption of predictive coding schemes during anesthesia

In the present study, we find that isoflurane suppresses CC feedback afferents to L4 and L5

pyramidal cells (Figure 3) as well as those to dendrite-targeting SOM+ interneurons (Figure 5),

consistent with previous studies showing sensitivity of feedback connections during anesthesia

(Boly et al., 2012; Liu et al., 2012; Raz et al., 2014; Mashour and Hudetz, 2017; Murphy et al.,

2019). Feedback activation of the cortical column is mediated largely by intracortical afferents

to layer 1 that target distal dendrites of pyramidal cells. Dendritic calcium currents activated by

these inputs increase the gain of the input/output relationship of cortical pyramidal cells and

are highly correlated with sensory perception (Manita et al., 2015; Takahashi et al., 2016). This

process is tightly regulated by powerful disynaptic inhibition from SOM+ interneurons

(Murayama et al., 2009). Our observation that inputs to both SOM+ cells and infragranular

pyramidal cells are preferentially suppressed notably intersects with these previous findings in

the context of sensory coding. Suppression of intracortical feedback inputs to two key players

in this integration process may therefore disrupt the typical input/output function of pyramidal

cells.

Disruption of integration of inputs to L5 pyramidal cells is also consistent with what might be expected under anesthesia in the context of current theoretical models. The predictive coding hypothesis, for example, has often been used as a framework for understanding consciousness and sensory processing (Friston, 2005; Bastos et al., 2012; Clark, 2013). Under the predictive coding hypothesis, feedback signals must activate inhibition to offset incoming excitatory signals and prevent propagation of the excitatory, ascending error signal. Discrepancies between internally generated predictions and sensory input from external sources are propagated as excitatory "error signals". Our data provide further evidence that multiple components of the predictive coding architecture may be disrupted by anesthesia. First, integration of predictions carried by feedback afferents may be perturbed, as feedback inputs to both SOM+ cells and L5 pyramidal cells – key mediators of feedback modulation of sensory responses – are altered by isoflurane. Second, propagation of error signals in the feedforward direction may be disrupted, as both TC and CC inputs to L4 cells – which mediate intracortical feedforward signals – are suppressed by isoflurane. Our data may demonstrate physiological correlates of components of the predictive coding model and provide insight into the disruption of sensory processing during anesthesia.

Future Directions

In these experiments, we investigate the effect of isoflurane on discrete components of a higher order cortical circuit. Investigating the pathway-, cell type-, and layer-specific effects of isoflurane is imperative for understanding the larger systems in which these components operate. Our findings will inform future investigations of sensory processing and changes in consciousness during anesthesia. Although here we activated inputs to layer 1 independently, future experiments *in vivo* will provide insight into how overlapping inputs to layer 1 affect arousal and sensory processing. It is reasonable to expect that the combined effects of

isoflurane on TC and CC inputs, for example, may engender supralinear effects on underlying

circuit components, synergistically contributing to disruption of sensory processing. Moreover,

in vivo investigations, such as those that involve artificially manipulating activity in different cell

types during anesthesia, may allow for causal, detailed descriptions of the pathway- and cell

type-specific components that are necessary and/or sufficient for consciousness. Lastly, a

recent study by Suzuki and Larkum (2020) suggests that anesthetics may prevent integration

of feedback signals by targeting signal amplification in apical dendrites of pyramidal cells, and

implies that this process may be mediated by metabotropic glutamate and cholinergic

receptors(Suzuki and Larkum, 2020). Future investigations should probe the extent to which

distinct inputs to layer 1 engage metabotropic glutamate or cholinergic systems, and how the

synaptic effects of anesthetics on layer 1 contributes to disruption of input-output computations

by anesthetics.

ACKNOWLEDGEMENTS

This work was supported by the National Institutes of Health (R01 GM109086 to MIB) and the

Department of Anesthesiology, School of Medicine and Public Health, University of Wisconsin,

Madison, WI, USA. The authors would also like to thank Bryan Krause for insight provided for

the statistical analyses conducted in this study, and Sean Grady for technical support.

DISCLOSURES

The authors have nothing to disclose.

REFERENCES

Akeju O, Loggia ML, Catana C, Pavone KJ, Vazquez R, Rhee J, Contreras Ramirez V,
        Chonde DB, Izquierdo-Garcia D, Arabasz G, Hsu S, Habeeb K, Hooker JM, Napadow

V, Brown EN, Purdon PL (2014) Disruption of thalamic functional connectivity is a neural correlate of dexmedetomidine-induced unconsciousness. Elife 3:e04499.

Audette NJ, Urban-Ciecko J, Matsushita M, Barth AL (2018) POm Thalamocortical Input Drives Layer-Specific Microcircuits in Somatosensory Cortex. Cereb Cortex 28:1312-1328.

Bastos AM, Usrey WM, Adams RA, Mangun GR, Fries P, Friston KJ (2012) Canonical microcircuits for predictive coding. Neuron 76:695-711.

Baumgart JP, Zhou ZY, Hara M, Cook DC, Hoppa MB, Ryan TA, Hemmings HC, Jr. (2015) Isoflurane inhibits synaptic vesicle exocytosis through reduced Ca2+ influx, not Ca2+-exocytosis coupling. Proc Natl Acad Sci U S A 112:11959-11964.

Beierlein M, Gibson JR, Connors BW (2003) Two dynamically distinct inhibitory networks in layer 4 of the neocortex. Journal of Neurophysiology 90:2987-3000.

Binzegger T, Douglas RJ, Martin KA (2009) Topology and dynamics of the canonical circuit of cat V1. Neural Netw 22:1071-1078.

Boly M, Moran R, Murphy M, Boveroux P, Bruno MA, Noirhomme Q, Ledoux D, Bonhomme V, Brichant JF, Tononi G, Laureys S, Friston K (2012) Connectivity changes underlying spectral EEG changes during propofol-induced loss of consciousness. J Neurosci 32:7082-7090.

Clark A (2013) Whatever next? Predictive brains, situated agents, and the future of cognitive science. Behav Brain Sci 36:181-204.

Covic EN, Sherman SM (2011) Synaptic properties of connections between the primary and secondary auditory cortices in mice. CerebCortex 21:2425-2441.

Crick F, Koch C (2003) A framework for consciousness. Nat Neurosci 6:119-126.

Cruikshank SJ, Urabe H, Nurmikko AV, Connors BW (2010) Pathway-Specific Feedforward Circuits between Thalamus and Neocortex Revealed by Selective Optical Stimulation of Axons. Neuron 65:230-245.

Cruikshank SJ, Ahmed OJ, Stevens TR, Patrick SL, Gonzalez AN, Elmaleh M, Connors BW (2012) Thalamic Control of Layer 1 Circuits in Prefrontal Cortex. The Journal of Neuroscience 32:17813-17823.

De Pasquale R, Sherman SM (2011) Synaptic properties of corticocortical connections between the primary and secondary visual cortical areas in the mouse. JNeurosci 31:16494-16506.

Dehaene S, Changeux JP (2011) Experimental and theoretical approaches to conscious processing. Neuron 70:200-227.

Dichter MA, Ayala GF (1987) Cellular mechanisms of epilepsy: a status report. Science 237:157-164.

Douglas RJ, Martin KA (2004) Neuronal circuits of the neocortex. AnnuRevNeurosci 27:419-451.

Douglas RJ, Martin KAC (2007) Recurrent neuronal circuits in the neocortex. Current Biology 17:R496-R500.

Douglas RJ, Martin KAC, Whitteridge D (1989) A canonical microcircuit for neocortex. Neural Computation 1:480-488.

Ferguson BR, Gao W-J (2018) PV Interneurons: Critical Regulators of E/I Balance for Prefrontal Cortex-Dependent Behavior and Psychiatric Disorders. Frontiers in Neural Circuits 12.

Ferrarelli F, Massimini M, Sarasso S, Casali A, Riedner BA, Angelini G, Tononi G, Pearce RA (2010) Breakdown in cortical effective connectivity during midazolam-induced loss of consciousness. Proc Natl Acad Sci USA 107:2681-2686.

Friston K (2005) A theory of cortical responses. Philos Trans R Soc Lond B Biol Sci 360:815-836.

Haider B, Häusser M, Carandini M (2013) Inhibition dominates sensory responses in the awake cortex. Nature 493:97-100.

Hentschke H, Raz A, Krause BM, Murphy CA, Banks MI (2017) Disruption of cortical network activity by the general anesthetic isoflurane. Br J Anaesth 119:685-696.

Herring BE, Xie Z, Marks J, Fox AP (2009) Isoflurane inhibits the neurotransmitter release machinery. J Neurophysiol 102:1265-1273.

Honjoh S, Sasai S, Schiereck SS, Nagai H, Tononi G, Cirelli C (2018) Regulation of cortical activity and arousal by the matrix cells of the ventromedial thalamic nucleus. Nature Communications 9:2100.

Koch C, Massimini M, Boly M, Tononi G (2016) Neural correlates of consciousness: progress
        and problems. Nat Rev Neurosci 17:307-321.

Krause BM, Murphy CA, Uhlrich DJ, Banks MI (2017) PV+ Cells Enhance Temporal Population
        Codes but not Stimulus-Related Timing in Auditory Cortex. Cereb Cortex.

Langsjo JW, Alkire MT, Kaskinoro K, Hayama H, Maksimow A, Kaisti KK, Aalto S, Aantaa R,
        Jaaskelainen SK, Revonsuo A, Scheinin H (2012) Returning from oblivion: imaging the
        neural core of consciousness. J Neurosci 32:4935-4943.

Lee U, Ku S, Noh G, Baek S, Choi B, Mashour GA (2013) Disruption of frontal-parietal
        communication by ketamine, propofol, and sevoflurane. Anesthesiology 118:1264-1275.

Liu X, Lauer KK, Ward BD, Li SJ, Hudetz AG (2013) Differential effects of deep sedation with
        propofol on the specific and nonspecific thalamocortical systems: a functional magnetic
        resonance imaging study. Anesthesiology 118:59-69.

Liu X, Lauer KK, Ward BD, Rao SM, Li SJ, Hudetz AG (2012) Propofol disrupts functional
        interactions between sensory and high-order processing of auditory verbal memory.
        Hum Brain Mapp 33:2487-2498.

Maksimow A, Silfverhuth M, Langsjo J, Kaskinoro K, Georgiadis S, Jaaskelainen S, Scheinin H
        (2014) Directional connectivity between frontal and posterior brain regions is altered
        with increasing concentrations of propofol. PLoS One 9:e113616.

Manita S, Suzuki T, Homma C, Matsumoto T, Odagawa M, Yamada K, Ota K, Matsubara C,
        Inutsuka A, Sato M, Ohkura M, Yamanaka A, Yanagawa Y, Nakai J, Hayashi Y, Larkum
        ME, Murayama M (2015) A Top-Down Cortical Circuit for Accurate Sensory Perception.
        Neuron 86:1304-1316.

Mashour GA, Hudetz AG (2017) Bottom-Up and Top-Down Mechanisms of General
        Anesthetics Modulate Different Dimensions of Consciousness. Front Neural Circuits
        11:44.

Meyer JF, Golshani P, Smirnakis SM (2018) The Effect of Single Pyramidal Neuron Firing
        Within Layer 2/3 and Layer 4 in Mouse V1. Front Neural Circuits 12:29.

Mitzdorf U (1985) Current source-density method and application in cat cerebral cortex:
        investigation of evoked potentials and EEG phenomena. Physiol Rev 65:37-100.

Murayama M, Pérez-Garci E, Nevian T, Bock T, Senn W, Larkum ME (2009) Dendritic encoding of sensory stimuli controlled by deep cortical interneurons. Nature 457:1137-1141.

Murphy C, Krause B, Banks M (2019) Selective effects of isoflurane on cortico-cortical feedback afferent responses in murine non-primary neocortex. Br J Anaesth 123:488-496.

Murphy CA, Krause BM, Grady SM, Banks MI (2017) Effect of isoflurane on selectively activated afferent pathways in neocortex. In: Society for Neuroscience Annual Meeting. Washington D.C., USA.

Nourski KV, Steinschneider M, Rhone AE, Kawasaki H, Howard MA, 3rd, Banks MI (2018) Auditory Predictive Coding across Awareness States under Anesthesia: An Intracranial Electrophysiology Study. J Neurosci 38:8441-8452.

Nourski KV, Banks MI, Steinschneider M, Rhone AE, Kawasaki H, Mueller RN, Todd MM, Howard MA, 3rd (2017) Electrocorticographic delineation of human auditory cortical fields based on effects of propofol anesthesia. Neuroimage 152:78-93.

Pettersen KH, Devor A, Ulbert I, Dale AM, Einevoll GT (2006) Current-source density estimation based on inversion of electrostatic forward solution: effects of finite extent of neuronal activity and conductivity discontinuities. JNeurosciMethods 154:116-133.

Picchioni D, Pixa ML, Fukunaga M, Carr WS, Horovitz SG, Braun AR, Duyn JH (2014) Decreased connectivity between the thalamus and the neocortex during human nonrapid eye movement sleep. Sleep 37:387-397.

Pouille F, Scanziani M (2001) Enforcement of temporal fidelity in pyramidal cells by somatic feed- forward inhibition. Science 293:1159-1163.

Raz A, Grady SM, Krause BM, Uhlrich DJ, Manning KA, Banks MI (2014) Preferential effect of isoflurane on top-down versus bottom-up pathways in sensory cortex. Front Syst Neurosci 8.

Redinbaugh MJ, Phillips JM, Kambi NA, Mohanta S, Andryk S, Dooley G, Afrasiabi M, Raz A, Saalmann Y (2020) Thalamus Modulates Consciousness Via Layer-Specific Control of Cortex. Neuron 105.

Sanders RD, Banks MI, Darracq M, Moran R, Sleigh J, Gosseries O, Bonhomme V, Brichant JF, Rosanova M, Raz A, Tononi G, Massimini M, Laureys S, Boly M (2018) Propofol-induced unresponsiveness is associated with impaired feedforward connectivity in cortical hierarchy. Br J Anaesth 121:1084-1096.

Sermet BS, Truschow P, Feyerabend M, Mayrhofer JM, Oram TB, Yizhar O, Staiger JF, Petersen CCH (2019) Pathway-, layer- and cell-type-specific thalamic input to mouse barrel cortex. eLife 8:e52665.

Suzuki M, Larkum ME (2020) General Anesthesia Decouples Cortical Pyramidal Neurons. Cell 180:666-676.e613.

Takahashi N, Oertner TG, Hegemann P, Larkum ME (2016) Active cortical dendrites modulate perception. Science 354:1587-1590.

Viaene AN, Petrof I, Sherman SM (2011a) Synaptic properties of thalamic input to layers 2/3 and 4 of primary somatosensory and auditory cortices. JNeurophysiol 105:279-292.

Viaene AN, Petrof I, Sherman SM (2011b) Properties of the thalamic projection from the posterior medial nucleus to primary and secondary somatosensory cortices in the mouse. Proc Natl Acad Sci USA 108:18156-18161.

Xie Z, McMillan K, Pike CM, Cahill AL, Herring BE, Wang Q, Fox AP (2013) Interaction of anesthetics with neurotransmitter release machinery proteins. Journal of neurophysiology 109:758-767.

Xu X, Olivas ND, Ikrar T, Peng T, Holmes TC, Nie Q, Shi Y (2016) Primary visual cortex shows laminar-specific and balanced circuit organization of excitatory and inhibitory synaptic connectivity. Journal of Physiology 594:1891-1910.

Xue M, Atallah BV, Scanziani M (2014) Equalizing excitation–inhibition ratios across visual cortical neurons. Nature 511:596.

Figure 1

Figure 1. Effects of isoflurane on optogenetically evoked current sinks in layer 1 are pathway-specific. *A.* CSD color plots comparing evoked synaptic responses to a 2 msec light pulse (blue arrows) across control (*A*, left column), 0.24 mM isoflurane (*A*, middle column), and recovery (*A*, right column) for representative examples for TC (top row) and CC (bottom row) afferents; color scale: -5.0 to 5.0 µA mm^{-3}. *B.* Signals from layer 1 were used to compare synaptic effects across control (solid), 0.24 mM isoflurane (dotted), and recovery (dashed). Shaded regions indicate ± 1 SD among trials. *C.* Model fits of percent block of layer 1 sink by isoflurane with 95% confidence interval (vertical line). Responses evoked by stimulation of CC afferents to cortical layer 1 (orange) are suppressed to a greater extent under isoflurane than TC inputs (red). Pale markers reflect estimates of block by isoflurane at 0.24 mM based on results from Murphy et al., 2019. Raw values of CSD sink amplitudes under control conditions are shown in **Table 1.**

Figure 2

Figure 2. Example recordings from cortical pyramidal cells in response to activation of two distinct layer 1 afferents. *A.* Schematic demonstrating recordings conducted for panel *B*. TC afferents originating in the posterior medial nucleus of the thalamus were optogenetically activated while whole cell patch clamp recordings were conducted in pyramidal cells in L2/3, L4, or L5. *B.* Example recordings of TC-evoked responses from L2/3 (top), L4 (middle), or and L5 (bottom) cells during control (solid), 0.24 mM isoflurane (dotted), and recovery (dashed) conditions. Shaded regions indicate ± 1 SD among trials. *C-D.* Same as *A-B* for responses to CC inputs originating in cingulate cortex. Population results of isoflurane effects are shown in **Figure 3.** Raw values for control conditions are shown in **Table 2.**

Figure 3

Figure 3. Suppression of post-synaptic responses by isoflurane is cell type-specific. Model estimates of the percent block by isoflurane of post-synaptic responses evoked by TC (red) or CC (orange) inputs to L2/3, L4, and L5 cell populations. Estimates are represented by triangles; vertical lines span 95% confidence intervals for isoflurane effects for each pathway-cell type combination, where confidence intervals overlapping 0% block are not significant (P>0.05). Both TC and CC inputs to L4 pyramidal cells were suppressed by isoflurane, as were CC inputs to L5 pyramidal cells. Effects are also quantified in **Table 3.**

Figure 4

Figure 4. Example recordings from SOM+ and PV+ interneurons in response to activation of distinct layer 1 afferents. *A*. Schematic demonstrating recordings conducted for panel *B*. CC afferents to layer 1 were optogenetically activated while whole cell patch clamp recordings were conducted in SOM+ interneurons. *B*. Example recordings from CC-evoked responses in a SOM+ cell during control (solid), 0.24 mM isoflurane (dotted), and recovery (dashed) conditions. Shaded regions indicate ± 1 SD among trials. *C-D*. Same as *A-B* for CC inputs to PV+ cells. *E-F*. Same as A-B for TC inputs to PV+ cells. Note that examples of TC inputs to SOM+ cells are not shown, as TC inputs failed to evoke synaptic responses in SOM+ cells. Population results of isoflurane effects are shown in **Figure 5.** Raw values of PSP amplitude during control conditions are shown in **Table 2.**

Figure 5

Figure 5. CC feedback inputs to SOM+ interneurons are suppressed by isoflurane, while TC and CC inputs to PV+ interneurons are not. Model estimates of the percent block by isoflurane of post-synaptic responses evoked by TC (red) or CC (orange) inputs to SOM+ and PV+ interneuron populations. Estimates are represented by circles (SOM+) or squares (PV+); vertical lines span 95% confidence intervals for isoflurane effects for each pathway-cell type combination, where confidence intervals overlapping 0% block are not significant (P>0.05). CC inputs to SOM+ cells were suppressed by isoflurane, but no effect of isoflurane was detected in TC nor CC inputs to PV+ cells. Effects are also quantified in **Table 3.**

Figure 6

Figure 6. Isoflurane increases paired pulse ratios for CC inputs to SOM+ cells, suggest pre-synaptic mechanisms. *A.* Examples recordings from the four different pathway-cell type pairs that demonstrated suppression of synaptic responses by isoflurane. Responses for control (black) and isoflurane (either orange (CC) or red (TC)) conditions are normalized to the amplitude of the first pulse under control conditions to demonstrate paired pulse effects. Arrows show comparison of paired pulse effects between control (black) and isoflurane (either orange or red) conditions. *B.* Model estimates of the changes to paired pulse ratio during isoflurane, where positive values represent higher paired pulse ratios during isoflurane. Estimates of isoflurane effects on paired pulse ratios for responses evoked by TC (red) or CC (orange) inputs to all recorded populations are shown. Estimates are represented by shapes; vertical lines span 95% confidence intervals for isoflurane effects for each pathway-cell type combination, where confidence intervals overlapping 0 are not significant (P>0.05). Paired pulse ratios for CC inputs to SOM+ cells were enhanced by isoflurane, suggesting isoflurane blocks synaptic responses to SOM+ cells through pre-synaptic mechanisms. Raw values of PPR during control conditions are shown in **Table 2**, and isoflurane effects are quantified in **Table 5**.

Figure 7. Isoflurane alters intrinsic membrane properties of L4 pyramidal cells, suggesting a shared mechanism by which TC and CC inputs are blocked. *A.* Model estimates of percent change of input resistance during isoflurane, where negative values represent decreases in input resistance (i.e., increased conductance). *B.* Model estimates of shifts of resting membrane potential during isoflurane, where negative values represent hyperpolarization of membrane potentials during isoflurane compared to baseline values. *C.* Same as in *A* for percent changes in membrane time constants. *D.* Same as in *B* for spike thresholds. L4 pyramidal cells became leakier (i.e., lower input resistances and shorter time constants) during isoflurane relative to baseline. Raw values of intrinsic properties during control conditions are shown in **Table 4.**

Table 1

	Latency (msec)	L1 sink amplitude ($\mu A \cdot mm^{-3}$)	% Block by isoflurane [95% CI]	P-value
TC	3.81±0.52	6.18±1.64	+2.9 [-10.6 18.3]	0.69
CC	3.61±0.56	5.33±1.35	-21.8 [-31.1 -11.3]	0.00026

Table 1. Activation of cortical afferents evokes short latency current sinks in layer 1 that are sensitive to isoflurane in a pathway-dependent manner. Layer 1 current sink amplitudes evoked by activation of CC inputs were suppressed by isoflurane, while those evoked by TC inputs were not. Estimates are based on predicted values from linear mixed effects model (see Methods).

Table 2

	PSP amplitude	PSP latency	PPR
TC→			
L^2/$_3$ Pyr (n=5)	2.4±1.8	5.9±2.3	0.86±0.33
L4 Pyr (n=6)	4.1±2.3	5.6±1.6	0.64±0.23
L5 Pyr (n=5)	8.3±5.7	5.4±1.8	0.77±0.29
SOM+ (n=2)	5.1±4.5	3.0±0.5	0.61±0.04
PV+ (n=8)	6.4±2.3	4.1±0.7	0.68±0.25
CC→			
L^2/$_3$ Pyr (n=6)	2.0±1.4	4.6±1.2	0.57±0.30
L4 Pyr (n=6)	4.6±4.6	4.8±1.2	0.79±0.27
L5 Pyr (n=5)	2.6±1.7	6.5±1.8	1.05±0.34
SOM+ (n=6)	6.5±5.0	4.6±0.7	0.74±0.45
PV+ (n=8)	4.3±4.8	4.9±1.2	0.78±0.32

Table 2. Properties of evoked responses in whole cell patch clamp recordings conducted during control conditions in pyramidal cells and interneurons.
Values shown are mean ± standard deviation. Latency was calculated relative to onset of 2 msec light pulse. Paired pulse ratios were calculated as the amplitude of the 2nd post-synaptic potential (relative to the pre-response baseline) divided by the baseline subtracted amplitude of the 1st post-synaptic potential.

Table 3

	% Block	95% CI	P-value
TC→			
L^2/$_3$ Pyr	7.0	[-14.2 33.5]	0.54
L4 Pyr	-20.1	[-34.7 -2.3]	0.029
L5 Pyr	-1.1	[-20.7 23.3]	0.92
PV+	-6.6	[-21.6 11.3]	0.44
CC→			
L^2/$_3$ Pyr	-13.6	[29.4 5.7]	0.15
L4 Pyr	-20.3	[-34.8 -2.4]	0.028
L5 Pyr	-21.5	[-37.0 -2.0]	0.033
SOM+	-35.8	[-47.5 -21.4]	<0.0001
PV+	-10.0	[-24.5 7.2]	0.23

Table 3. Isoflurane affects synaptic responses in a cell type-dependent manner. Estimates are based on predicted values from linear mixed effects model (see Methods). Values represent the percent change in the amplitude of post-synaptic responses during isoflurane relative to baseline conditions. Negative values represent suppression of synaptic responses by isoflurane.

Table 4

	RMP (mV)	R_{in} (MΩ)	T_m (msec)	Spike Threshold (mV)
L^2/$_3$ Pyr (n=11)	-71.5±3.6	114.2±45.9	14.1±10.7	-43.1±4.0
L4 Pyr (n=12)	-68.9±5.3	183.2±103.8	19.0±13.2	-41.9±6.1
L5 Pyr (n=10)	-65.1±3.7	135.77±85.9	18.0±9.8	-41.8±5.3
SOM+ (n=8)	-60.3±6.2	317.3±138.8	38.2±38.0	-42.2±3.9
PV+ (n=16)	-63.5±4.0	166.8±47.1	9.6±2.5	-46.3±7.9

Table 4. Intrinsic properties of recorded cell types during control conditions.
Values are mean ± standard deviation. Properties were measured ~2 minutes after achieving whole cell access. A series of 10-20 current steps (600 msec) was delivered at progressive intervals (10-40 pA), beginning with a hyperpolarizing current injection. Resting membrane potentials (RMP) were calculated by averaging membrane potentials in the 100 msec prior to onset of all current pulses. Input resistances (R_{in}) were extrapolated from relationship between current and steady state voltages. Time constants (T_m) were estimated using the first hyperpolarizing current pulse. Spike thresholds were evaluated using the first trial to evoke an action potential, where the spike threshold was the voltage at which the second derivative exceeded twice the standard deviation of the pre-spike baseline value.

Table 5

	ΔPPR	95% CI	P-value
TC→			
L^2/$_3$ Pyr	+0.02	[-0.32 0.26]	0.90
L4 Pyr	+0.13	[-0.10 0.36]	0.27
L5 Pyr	-0.18	[-0.43 0.07]	0.15
PV+	+0.02	[-0.18 0.22]	0.84
CC→			
L^2/$_3$ Pyr	-0.07	[-0.30 0.16]	0.56
L4 Pyr	+0.13	[-0.10 0.36]	0.27
L5 Pyr	-0.18	[-0.42 0.08]	0.17
SOM+	+0.45	[0.22 0.68]	0.0002
PV+	-0.03	[-0.23 0.17]	0.76

Table 5. Isoflurane alters short-term synaptic plasticity for CC inputs to SOM+ interneurons. Estimates are based on predicted values from linear mixed effects model (see Methods) and represent the change in PPR (in mV) relative to baseline PPR. Negative values represent decreases in PPR (e.g., stronger paired pulse depression) during isoflurane.