Title: Interactions with Commensal and Pathogenic Bacteria Induce HIV-1 Latency in

Macrophages through Altered Transcription Factor Recruitment to the LTR

Authors: Gregory A. Viglianti[1], Vicente Planelles[2], Timothy M. Hanley[1,2,#]

Affiliations: [1]Department of Microbiology, Boston University School of Medicine, Boston, MA

02118

[2]Department of Pathology, University of Utah Health, Salt Lake City, UT 84105

Running Title: Induction of HIV-1 Latency in Macrophages by Bacteria

[#]Address correspondence to Timothy M. Hanley, timothy.hanley@hsc.utah.edu

ABSTRACT

Macrophages are infected by HIV-1 *in vivo* and contribute to both viral spread and

pathogenesis. Recent human and animal studies suggest that HIV-1-infected macrophages

serve as a reservoir that contributes to HIV-1 persistence during anti-retroviral therapy. The

ability of macrophages to serve as persistent viral reservoirs is likely influenced by the local

tissue microenvironment, including interactions with pathogenic and commensal microbes. Here

we show that the sexually transmitted pathogen *Neisseria gonorrhoeae* (GC) and the gut-

associated microbe *Escherichia coli* (*E. coli*), which encode ligands for both Toll-like receptor 2

(TLR2) and TLR4, repressed HIV-1 replication in macrophages and thereby induced a state

reminiscent of viral latency. This repression was mediated by signaling through TLR4 and the

adaptor protein TRIF and was associated with increased production of type I interferons.

Inhibiting TLR4 signaling, blocking type 1 interferon, or knocking-down TRIF reversed LPS- and

GC-mediated repression of HIV-1. Finally, the repression of HIV-1 in macrophages was

associated with the recruitment of interferon regulatory factor 8 (IRF8) to the interferon

stimulated response element (ISRE) downstream of the 5' HIV-1 long terminal repeat (LTR).

Our data indicate that IRF8 is responsible for repression of HIV-1 replication in macrophages in

response to TRIF-dependent signaling during GC and *E. coli* co-infection. These findings

highlight the potential role of macrophages as HIV-1 reservoirs as well as the role of the tissue

microenvironment and co-infections as modulators of HIV-1 persistence.

IMPORTANCE The major barrier toward the eradication of HIV-1 infection is the presence of a

small reservoir of latently infected cells, which include CD4+ T cells and macrophages that

escape immune-mediated clearance and the effects of anti-retroviral therapy. There remain

crucial gaps in our understanding of the molecular mechanisms that lead to transcriptionally

silent or latent HIV-1 infection of macrophages. The significance of our research is in identifying

microenvironmental factors, such as commensal and pathogenic microbes, that can contribute

to the establishment and maintenance of latent HIV-1 infection in macrophages. It is hoped that

identifying key processes contributing to HIV-1 persistence in macrophages may ultimately lead

to novel therapeutics to eliminate latent HIV-1 reservoirs *in vivo*.

INTRODUCTION

Macrophages are among the immune cells located within the gastrointestinal and genitourinary

mucosae thought to play a role in HIV-1 sexual transmission and pathogenesis (1-3). A number

of studies examining either HIV-1 infection of human vaginocervical or gastrointestinal tissue

explants or SIV_{mac} infection in rhesus macaque animal models have shown that macrophages

are among the earliest cells infected during mucosal transmission (2, 4, 5). Macrophages can

be productively infected with HIV-1 and are thought to be a source of virus persistence *in vivo*

(6). Given their role in transmission, pathogenesis, and viral persistence, it is important to

understand how the local mucosal microenvironment and cellular signaling pathways modulate

interactions between macrophages and HIV-1.

Sexually-transmitted infections (STIs) have been shown to be co-factors that enhance HIV-1

transmission (7). *Neisseria gonorrhoeae* (gonococcus, GC) is a non-ulcerative STI that is

thought to augment mucosal transmission of HIV-1 both by inducing inflammation and by

directly activating virus infection and replication (8-13). The role of GC in HIV-1 persistence is

less well understood. Several studies have implicated GC-encoded pathogen-associated

molecular patterns (PAMPs) as mediators of both inflammation and HIV-1 activation in target

cells such as macrophages; however, the interactions between GC and macrophages are

complex. GC encodes PAMPs capable of engaging Toll-like receptors (TLRs), including TLR2,

TLR4, and TLR9 (14, 15). While the effects of co-infection with live GC on HIV-1 replication in

macrophages have not been reported, purified lipooligosaccharide (LOS) as well as *Escherichia*

coli lipopolysaccharide (LPS) have been shown to repress virus replication through the

production of type 1 interferons (IFNs) (16, 17). In the case of LPS, repression is due to

undefined effects at the level of gene expression. Although it is not entirely clear how TLR2

signaling affects HIV-1 expression in macrophages, studies have shown that purified TLR2

ligands activate virus replication in macrophages (18) and latently-infected T cells (19).

Here, we demonstrate that co-infection with GC and *E. coli* repress HIV-1 expression in

macrophages. To investigate the underlying mechanism(s) responsible for this repression, we

examined the individual effects of TLR2 and TLR4 signaling on HIV-1 expression in

macrophages. TLR2 signaling activated HIV-1 expression in macrophages, whereas TLR4

signaling repressed virus expression. Importantly, TLR4 signaling overcame the activation

effects of TLR2 signaling in macrophages. The TLR4-mediated repression of HIV-1 in

macrophages co-infected with GC or *E. coli* was dependent on signaling through Toll/IL-1

receptor domain-containing adapter inducing interferon-β (TRIF) and required type 1 IFN

production. Finally, we showed that TLR4 signaling leads to the late-phase recruitment of IRF8

to the interferon-stimulated response element (ISRE) downstream of the 5' HIV-1 LTR in

infected macrophages. Taken together, our data suggest TRIF-mediated signaling represses

HIV-1 replication in response to GC or *E. coli* co-infection in an IRF8-dependent manner and

shifts macrophages from a state of robust HIV-1 expression to a state of persistent low-

level/latent infection.

MATERIALS AND METHODS

Ethics Statement. This research has been determined to be exempt by the Institutional Review

Board of the Boston University Medical Center since it does not meet the definition of human

subjects research.

Cell isolation and culture. Primary human CD14+ monocytes were isolated from the

peripheral blood mononuclear cells of healthy donors using anti-CD14 magnetic beads (Miltenyi

Biotec) per the manufacturer's instructions. Primary monocyte-derived macrophages (MDMs)

were generated by culturing CD14+ monocytes in the presence of 10% human AB serum and

10% FBS for 6 days. Following differentiation, MDMs were cultured in RPMI-1640

supplemented with 10% FBS, 100 U/ml penicillin, 100 μg/ml streptomycin, and 0.29 mg/ml L-

glutamine. The genetic sex of a subset of the donors was determined by PCR amplification of

the SRY gene located on the Y chromosome. PM1 cells were cultured in RPMI-1640

supplemented with 10% FBS, 100 U/ml penicillin, 100 μg/ml streptomycin, and 0.29 mg/ml L-

glutamine. 293T cells were cultured in DMEM supplemented with 10% FBS, 100 U/ml penicillin,

100 μg/ml streptomycin, and 0.29 mg/ml L-glutamine. MAGI-CCR5 cells were cultured in DMEM

supplemented with 10% FBS, 100U/ml penicillin, 100 μg/ml streptomycin, 0.29 mg/ml L-

glutamine, 500 μg/ml G418, 1 μg/ml puromycin, and 0.1 μg/ml hygromycin B. HEK293-

TLR2^{CFP}TLR1^{YFP} cells and HEK293-TLR4^{CFP}/MD-2/CD14 cells were cultured in DMEM

supplemented with 10% FBS, 10 μg/ml ciprofloxacin, 0.29 mg/ml L-glutamine, and 500 μg/ml

G418.

Bacterial culture. *Neisseria gonorrhea* (GC) strain FA1090B was a generous gift from Dr.

Caroline Genco. GC was cultured from a glycerol stock on GC agar plates supplemented with

IsoVitalex Enrichment (Becton Dickinson) in a humidified 37°C incubator with 5% CO_2. *E. coli*

strain DH5α was purchased from New England Biolabs and was cultured from a glycerol stock

on LB agar plates at 37°C. Where indicated, bacteria were heat inactivated (heat killed) by

incubation at 56°C for 2 hours. Heat inactivation was monitored by culture on GC or LB agar

plates as described above.

Flow cytometry. TLR expression on viable MDMs was assessed eight days after isolation

using antibodies against TLR2 (clone TL2.1) and TLR4 (clone HTA125) (both from eBioscience)

and eFluor 450 fixable viability dye (eBioscience). MDMs were stained in plates, washed with

phosphate-buffered saline (PBS), fixed using BD Cytofix (BD Biosciences), and then detached

after incubation in PBS supplemented with 20 mM EDTA for 1 hour at 4°C. Flow cytometric data

was acquired using a Becton-Dickenson FACScan II or LSRFortessa and data was analyzed

using FlowJo software.

TLR ligands, interferons, and chemical inhibitors. PAM3CSK4, FSL-1, *Salmonella*

typhimurium flagellin (FLA-ST), poly I:C, and *E. coli* K12 LPS were obtained from Invivogen.

TLR ligands were reconstituted in endotoxin-free H_2O. IFN-α and IFN-β were purchased from

PBL Interferon Source. B18R was purchased from Abcam. BAY 11-7082, celastrol, U0126,

PD95809, and SB203580 were purchased from Sigma and reconstituted in DMSO. Dynasore

was purchased from Tocris Bioscience and was reconstituted in DMSO.

Virus Production. Single-round replication-defective HIV-1 reporter viruses were generated by

packaging a luciferase expressing reporter virus, BruΔEnvLuc2, or an enhanced green

fluorescent protein expressing reporter virus, BruΔEnvEGFP3, with the envelope glycoproteins

from VSV (VSV-G). In these constructs, reporter gene expression is under the control of the 5'

LTR. Reporter virus stocks were generated by transfecting HEK293T cells using the calcium

phosphate method as described previously (18). Replication competent HIV-1$_{Ba-L}$ was generated

by infection of PM1 cells as described previously (18). Virus titers were determined using MAGI-

CCR5 cells, and p24^{gag} content was determined by ELISA as described previously (18).

Virus infections. To assess viral replication, macrophages (2.5 x 10^5 cells/well in 24-well

plates) were incubated with VSV-G-pseudotyped HIV-luciferase reporter virus at a multiplicity of

infection (MOI) of 0.1 for 4 hours at 37°C. Cells were washed four to five times with PBS to

remove unbound virus, and cultured in growth medium. Following 48 hours of culture, cells were

treated with TLR ligands or vehicle, as indicated in the text and figure legends. After 18 hours,

the cells were washed twice with PBS and lysed in PBS/0.02% Triton X-100. Luciferase activity

was measured using BrightGlo luciferase reagent (Promega) and a MSII luminometer.

HIV-1 transcription. Total cytoplasmic RNA was isolated from MDMs using the RNeasy Mini kit

(Qiagen). RNA (100 ng) was analyzed by reverse transcription-PCR (RT-PCR) using the

OneStep RT-PCR kit (Qiagen). RNA was reverse transcribed and amplified in a total volume of

50 µl containing 2.5 mM MgCl2, 400 µM concentrations of each deoxynucleoside triphosphate,

10 U of RNasin RNase inhibitor (Promega), 5 µCi of α-^{32}P dATP, and 0.6 µM HIV-1 specific

primers. RNA samples were reverse transcribed for 30 minutes at 50°C. After an initial

denaturing step at 95°C for 15 minutes, cDNA products were amplified for 25 cycles each

consisting of a 30-second denaturing step at 94°C, a 45-second annealing step at 65°C, and a

one-minute extension step at 72°C. The amplification concluded with a 10-minute extension

step at 72°C. Samples were resolved on 5% nondenaturing polyacrylamide gels, visualized by

autoradiography, and quantified in a Molecular Dynamics PhosphorImager SI using

ImageQuant software. Alternatively, HIV-1 RNA was analyzed using the QuantiTect SYBR

Green RT-PCR kit (Qiagen) in a LightCycler 480 (Roche). The HIV-1 primers were specific for

the R and U5 regions of the LTR and amplify both spliced mRNAs and genomic RNA. The HIV-

1 primers were sense primer 5'-GGCTAACTAGGGAACCCACTGC-3' and antisense primer 5'-

CTGCTAGAGATTTTCCACACTGAC-3'). α-tubulin primers were: sense primer 5'

CACCCGTCTTCAGGGCTTCTTGGTTT-3' and antisense primer, 5'

CATTTCACCATCTGGTTGGCTGGCTC-3'. RNA standards corresponding to 500, 50, and 5 ng

of RNA from PAM3CSK4-activated MDMs were included in each experiment to ensure that all

amplifications were within the linear range of the assay.

HIV-1 RNA stability assays. MDMs (2×10^6 cells/well in 6-well plates) were incubated with

VSV-G-pseudotyped HIV-luciferase reporter virus at an MOI of 0.1 for 4 h at 37°C. Cells were

washed four to five times with PBS to remove unbound virus and cultured in growth medium.

Following 48 h of culture, cells were treated with TLR ligands (PAM3CSK4 or LPS at 100 ng/ml)

or vehicle for 4 hours. Actinomycin D (10 µg/ml) was then added to cells to block *de novo* RNA

synthesis, and total cytoplasmic RNA was isolated at given times as described in the figure

legends. Viral RNA was analyzed using the QuantiTect SYBR Green RT-PCR kit (Qiagen) in a

LightCycler 480 (Roche) with primers specific for the R and U5 regions of the LTR as described

above.

Cytokine release assays. MDMs (2.5×10^5 cells/well) were treated with PAM3CSK4 (100

ng/ml), LPS (100 ng/ml), or GC (MOI of 10) for 24 hours. Cell-free culture supernatants were

collected and analyzed for TNF-α (eBioscience) or IFN-β (PBL Interferon Source) release by

commercially-available ELISA following the manufacturer's instructions.

Chromatin immunoprecipitation assays. 1.2×10^7 MDMs were incubated with VSV-G-

pseudotyped HIV-EGFP reporter virus at an MOI of 2 for four hours at 37°C. Cells were washed

four to five times with PBS to remove unbound virus and cultured in growth medium. Following

48 hours of culture, MDMs were treated with TLR ligands for various times as described in the

text. Cells were then fixed in 1% formaldehyde for 10 minutes at room temperature, quenched

with 125 mM glycine, and lysed in SDS lysis buffer (1% SDS, 10 mM EDTA, 50 mM Tris pH 8.1,

1 mM PMSF, 1 μg/ml aprotinin, 1 μg/ml pepstatin A). Cellular lysates were sonicated using a

cup horn (550 Sonic Dismembrator, Fisher Scientific) at a power setting of 5 with twenty-five 20-

second pulses on ice, which fragmented the chromatin to an average length of approximately

1000 bp. Samples were diluted and immunoprecipitated with antibodies against NF-κB p65,

IRF1, IRF2, IRF4, IRF8, rabbit IgG, or goat IgG (all from Santa Cruz Biotechnology). Purified

DNA samples from both ChIPs and input controls were resuspended in distilled H_2O and

analyzed by semi-quantitative PCR. PCR reactions contained 10 mm Tris-HCl pH 8.3, 50 mm

KCl, 1.5 mm $MgCl_2$, 100 pmol of each primer, 200 μm each dATP, dGTP, dCTP, and dTTP,

5μCi α^{32}P-dATP, and 2.5 units of Amplitaq Gold (Applied Biosystems) in a 50-μl reaction

volume. Following an initial denaturation step at 95°C for 15 minutes, DNAs were amplified for

30 cycles, each consisting of a 30-second denaturing step at 94°C, a 45-second annealing step

at 65°C, and a one-minute extension step at 72°C. Samples were electrophoresed on 5% non-

denaturing polyacrylamide gels, visualized by autoradiography, and quantified using a Molecular

Dynamics PhosphorImager SI using ImageQuant software. Alternatively, purified DNA from

ChIPs and input controls were analyzed using the PowerUp SYBR Green Master Mix (Applied

Biosystems) in a LightCycler 480 (Roche). Primers used to amplify specifically the HIV-1 5' LTR

and GLS were 5'- TGGAAGGGCTAATTTACTCCC -3' (sense) and 5'-

CATCTCTCCTTCTAGCCTC -3' (antisense). Control amplifications of a serial dilution of

purified genomic DNA from latently infected U1 cells were performed with each primer set to

ensure that all amplifications were within the linear range of the reaction. To calculate the

relative levels of association with the LTR, PhosphorImager data of the PCR products obtained

for immunoprecipitated chromatin samples were normalized against the PCR products obtained

for input DNA (% Input). Values were normalized across donors and expressed as relative

binding.

LTR mutant construction. The reported plasmid pLTR(Sp1)-luciferase was generated by PCR

amplification of pNL4-3 using the sense primer 5'-

CGGGGTACCCCGTGGAAGGGCTAATTTGGTCCC- 3' and the antisense primers 5'-

CCGCTCGAGCGGCATCTCTCTCCTTCTAGCCTC-3', digestion with KpnI and XhoI, and

ligation into KpnI/XhoI-digested pGL3-Basic (Promega). Mutations to the NF-κB and IRF binding

sites in pLTR(Sp1)-luciferase were generated using the QuickChange IIXL site-directed

mutagenesis kit (Stratagene). Primers used for site-directed mutagenesis are listed in Table 1.

The -158 LTR-luciferase construct was generated by deleting the LTR sequence upstream of

position -158 (relative to the start site of transcription) of pNL4-3, which includes the AP-1

binding sites located in the U3 portion of the 5' LTR, digestion of the resulting fragment with

KpnI and XhoI, and ligation into KpnI/XhoI-digested pGL3-Basic (Promega).

shRNA knock-down of MyD88, TRIF, and IRF8. MDMs (1.2×10^7) were transfected with

plasmids that encoded either a mixture of three to five shRNAs directed against MyD88, a

mixture of three to five shRNAs directed against TRIF, or a mixture of three to five control

shRNAs (Invivogen) and a blasticidin-resistance gene using Oligofectamine (Invitrogen) per the

manufacturer's instructions. Transfected cells were selected by culture in the presence of

blasticidin for 48 hours, and either used in HIV-1 replication assays or lysed for immunoblot

analysis to measure MyD88 and TRIF expression using a rabbit monoclonal antibody to MyD88

(Cell Signaling Technology), a rabbit polyclonal antibody to TRIF (Cell Signaling Technology), or

a mouse monoclonal antibody to β-actin (Sigma). Similarly, MDMs were transfected with

plasmids that encoded either a mixture of three to five shRNAs directed against IRF8 (Sigma) or

a mixture of control shRNAs (Sigma) and a puromycin-resistance gene using Oligofectamine

(Invitrogen) per the manufacturer's instructions. Transfected cells were selected by culture in

the presence of puromycin for 48 hours and either used in HIV-1 replication assays or lysed for

immunoblot analysis to measure IRF8 expression using a rabbit monoclonal antibody (Cell

Signaling Technology).

Overexpression of IRF8. MDMs (1.2×10^7) were transfected with a plasmid that encoded IRF8

(Origene) and a neomycin-resistance gene using Oligofectamine (Invitrogen) per the

manufacturer's instructions. Transfected cells were selected by culture in the presence of

neomycin for 48 hours and then used for HIV-1 replication assays or lysed for immunoblot

analysis to measure IRF8 expression using a rabbit monoclonal antibody (Cell Signaling

Technology).

Endocytosis/phagocytosis assays. MDMs (5×10^5/well) were treated with Dynasore (80 μM)

or DMSO and then incubated with pHrodo Green *E. coli* particles (Thermo Fisher) at 1 mg/ml for

2 hours at 37°C. The MDMs were then washed three times with PBS, incubated with eFluor 450

fixable viability dye (eBioscience) for 15 minutes at 4°C, and analyzed by flow cytometry. Flow

cytometric data was acquired using an Becton-Dickenson LSRFortessa, and data was analyzed

using FlowJo software.

Statistical analysis. Comparison between experimental samples was performed with a paired

one-tailed t-test with $p < 0.5$ denoting significant differences. Experiments were performed in

triplicate using cells from a minimum of four independent donors (unless otherwise indicated) to

control for interdonor variability.

RESULTS

HIV-1 gene expression in MDMs is enhanced or repressed in a TLR-specific manner. To

determine how purified TLR ligands affected HIV-1 gene expression, MDMs were infected with

a single-round infectious HIV-1 reporter virus, and then treated with ligands for TLR2, TLR3,

TLR4, or TLR5. Ligands that activated TLR2 or TLR5 enhanced HIV-1 replication, whereas

ligands for TLR3 or TLR4 repressed HIV-1 expression (Figure 1A). The effects of TLR ligands

on HIV-1 replication occurred at the level of transcription, as treatment with the TLR2/1 ligand

PAM3CSK4 led to an increase in HIV-1 mRNA accumulation, whereas treatment with the TLR4

ligand LPS led to a decrease in HIV-1 transcript levels (Figure 1C-1D). TLR treatment had no

effect of viral RNA stability, as viral RNA from LPS-treated MDMs had a similar decay rate to

that from untreated MDMs (Figure 1E). Recent studies have demonstrated that myeloid cells

from males and females have different susceptibilities to HIV-1 infection, largely due to

differential levels of innate immune responses and steroid hormones (20-22). We therefore

sought to determine whether there was a sex-based difference in the response to TLR ligand

treatment in MDMs. We found that TLR stimulation had similar effects on HIV-1 expression in

MDMs from both male and female donors (Supplemental Figure 1). These results indicate that

MyD88-dependent signaling enhances HIV-1 transcription whereas TRIF-dependent signaling

inhibits HIV-1 transcription in MDMs.

MyD88-dependent TLR signaling leads to the activation of both NF-κB and AP-1 transcription

factors, among others (23). The HIV-1 LTR contains binding sites for both NF-κB and AP-1. The

two NF-κB sites are thought to be essential for HIV-1 transcription (24, 25), whereas the AP-1

sites, while not essential, are thought to enhance HIV-1 transcription (26, 27). Previous studies

demonstrated that treatment of HIV-infected MDMs with the TLR2/TLR1 ligand PAM3CSK4 led

to an increased association of the p65 subunit of NF-κB and the c-fos subunit of AP-1 with the

5' LTR which, in turn, correlated with increased virus replication (18); however, the contributions

made by each pathway to TLR2-mediated activation have not been previously characterized.

To determine the roles of NF-κB and AP-1 in TLR2-activated HIV replication in MDMs, HIV-1-

infected cells were treated with either BAY 11-7082, an inhibitor of IκB kinase (28), celastrol, a

small molecule inhibitor of the IκB kinase complex (29), or inhibitors that disrupt AP-1 signaling.

As shown in Supplemental figure 2, BAY 11-7082 and celastrol treatment completely ablated

TLR2/1-enhanced HIV-1 expression. Similarly, the use of an LTR-based reporter construct with

mutations in the NF-κB binding sites did not result in increased gene expression in response to

TLR2 signaling (Supplemental Figure 2C). Treatment of HIV-infected macrophages with

inhibitors of kinases upstream of AP-1 activation, such as MEK1/2 (U0126, PD98509), and p38

(SB203580), resulted in modest, but reproducible, decreases in TLR2-mediated activation of

HIV-1 (Supplemental Figure 2D). Similarly, LTR reporter constructs lacking AP-1 binding sites

were activated in response to TLR2 signaling at levels similar to that of the WT construct, further

demonstrating the non-essential role of AP-1 in TLR2-mediated HIV-1 activation (Supplemental

Figure 2E). Although the regulation of HIV-1 transcription through multiple transcription factor

binding sites in and adjacent to the 5' LTR is complex, these data suggest that, in MDMs, TLR2-

activated HIV-1 expression is mediated primarily through NF-κB with a minor contribution from

AP-1 signaling.

Co-infection with *Neisseria gonorrhoeae* or *Escherichia coli* represses HIV-1 replication

in MDMs. Our preliminary studies using purified TLR ligands in isolation suggested that different

TLR signaling cascades had diverse effects on HIV-1 replication. Since most pathogens encode

multiple TLR ligands, we sought to determine the effects of intact pathogens on HIV-1

replication. We incubated HIV-infected MDMs with *N. gonorrhoeae* (GC), which expresses

ligands for TLR2, TLR4, and TLR9. We found that increasing amounts of GC led to a dose-

dependent decrease in HIV-1 replication in MDMs (Figure 1F). Bacterial replication was not

required for these effects, as heat-killed GC led to repression of HIV-1 replication in MDMs

(Supplemental Figure 3A). GC-mediated repression occurred at the level of viral transcription

(Figure 1G) and did not decrease viral RNA stability (Supplemental Figure 3B). Similar to what

we observed with purified LPS, the biological sex of the donors had no effect on GC- or *E. coli*-

mediated HIV-1 repression in MDMs (Supplemental Figure 3C). In addition, repression of HIV-1

replication is not specific for GC, but may be a generalized response to Gram-negative bacteria,

as co-infection with *E. coli* also repressed HIV-1 replication in MDMs in a manner similar to GC

(Figure 1H). Despite the presence of both activating (TLR2) and repressing (TLR4) TLR ligands,

both GC and *E. coli* mediated repression of HIV-1 replication in macrophages. This finding

raised several possibilities: 1) the dominance of TLR4 signaling over TLR2 signaling in MDMs;

2) different expression levels of TLR2, TLR4, and TLR4-associated molecules such as CD14

and MD-2 on MDMs; 3) different cytokine profiles produced in response to GC or *E. coli*; and/or

4) variable expression of signaling molecules downstream of TLRs. These scenarios were

further explored.

TLR4 signaling is dominant in MDMs. To determine whether certain TLR pathways are

dominant in MDMs, we performed co-treatments of HIV-infected MDMs with the TLR2 ligand

PAM3CSK4 and the TLR4 ligand LPS. We found that increasing the concentration of LPS

against a fixed concentration of PAM3CSK4 led to a reversal of TLR2-mediated activation of

HIV-1 and, eventually, to repression of HIV-1 replication (Figure 2A). Conversely, increasing the

concentration of PAM3CSK4 against a fixed concentration of LPS did not reverse LPS-mediated

repression of HIV-1 (Figure 2A). Flow cytometry was used to determine that the different

responses of MDMs were likely not due to receptor expression, as MDMs express both TLR2

and TLR4 (Figure 2B-C). In addition, MDMs produced both TNF-α and IFN-β in response to

LPS treatment, GC co-infection, and *E. coli* co-infection. Whereas treatment of MDMs with LPS

resulted in a similar cytokine profile to that of co-infection, treatment of MDMs with the TLR-

ligand PAM3CSK4 resulted in the production of TNF-α, but not appreciable levels of IFN-β

(Figure 2D-E). Taken together, our data suggest that TLR4 signaling, which negatively regulates

LTR-driven gene expression, is dominant in MDMs.

LPS- and GC-mediated repression of HIV-1 in MDMs is dependent on TRIF-mediated type

I IFN production. Since LPS and GC both induce type I IFN production, whereas the TLR2

ligand PAM3CSK4 does not, we wished to determine whether GC-stimulated production of IFN-

α/β contributes to repression of HIV-1 in MDMs. We found that treatment of HIV-infected MDMs

with the vaccinia virus-encoded soluble type I IFN receptor B18R reversed GC-mediated

inhibition of HIV-1 replication, suggesting that TLR4-mediated IFN production is required for

HIV-1 repression by GC (Figure 3A). Since both purified TLR4 ligands and GC, which encodes

ligands for TLR2, TLR4, and TLR9, repress HIV-1 replication in MDMs, we predicted that

downstream effector molecules of TLR4 signaling would contribute to the repression of HIV-1

replication in MDMs. First, we confirmed that TLR4 signaling was responsible for GC-mediated

HIV-1 repression in MDMs. Treatment with the TLR4-specific inhibitor TAK242 reversed the

LPS- and GC-dependent repression of HIV-1 in MDMs (Figure 3B). Treatment with TAK242 had

no effect on TLR2-mediated activation of HIV-1 replication in MDMs, consistent with reports that

TAK242 is specific for TLR4 (30).

It has been shown that TLR4, which can utilize both MyD88 and TRIF adaptor proteins, initiates

different signaling pathways dependent upon its cellular location. Cell-surface TLR4

engagement leads to MyD88-dependent signaling, whereas endosomal TLR4 engagement

leads to TRIF-dependent signaling (31). To examine whether TRIF-dependent signaling is

responsible for HIV-1 repression, we blocked dynamin-dependent endocytosis of TLR4 with

Dynasore, which prevents TRIF-dependent signaling while leaving MyD88-dependent signaling

intact. As shown, blocking endocytosis-mediated TLR4 internalization (Supplemental figure 4)

reversed GC-mediated repression of HIV-1 in MDMs (Figure 3C). Given the ability of GC to

signal through both TLR2-MyD88 and TLR4-TRIF, one might expect the inhibition of type I IFN

signaling by B18R or the inhibition of endocytosis by Dynasore to lead to augmented viral gene

expression through intact MyD88 signaling. However, we did not observe this, likely due to

incomplete inhibition of either IFN signaling or endocytosis.

To confirm the role of MyD88 in TLR2-mediated HIV-1 activation and TRIF in TLR4-mediated

HIV-1 repression, we used shRNAs to knock down the two molecules in HIV-infected MDMs

(Figure 3D). Knock-down of MyD88 led to a loss of TLR2-mediated HIV-1 activation, but had no

effect on LPS or GC-mediated HIV-1 repression (Figure 3E). In contrast, knock-down of TRIF

had no effect on TLR2-mediated HIV-1 activation, but reversed LPS- and GC-mediated

repression of HIV-1 replication (Figure 3E). Knock-down of either MyD88 or TRIF had no effect

on the activation of HIV-1 by the phorbol ester PMA, which signals directly through protein

kinase C, independently of TLRs (Figure 3E). These data suggest that the TLR4-TRIF-type I

IFN axis in MDMs leads to GC- and *E. coli*-mediated repression of HIV-1 replication.

TLR4 signaling leads to differential IRF recruitment to the HIV-1 LTR. Since type I IFN

production is critical for GC- and *E. coli*-mediated HIV-1 repression in MDMs, we examined the

role of interferon-stimulated genes (ISGs) in HIV-1 regulation. Previous studies have shown that

ISGs are temporally regulated in macrophages in response to innate immune sensors and type I

IFN signaling (32, 33). To determine whether the repressive effects of LPS were due to early or

late phase ISGs, HIV-1-infected MDMs were treated with the TLR2 ligand PAM3CSK4 or the

TLR4 ligand LPS and total cytoplasmic RNA was extracted at various times post treatment.

Treatment of HIV-infected MDMs with the TLR2 ligand PAM3CSK led to a continuous increase

in HIV-1 RNA levels (Figure 4A). In contrast, treatment of HIV-infected MDMs with the TLR4

ligand LPS led to an initial short-lived increase in HIV-1 RNA levels; however, levels steadily

declined thereafter (Figure 4A), indicating that HIV-1 transcription displays a biphasic response

to TLR4 stimulation in MDMs. This suggests that late-phase proteins induced by type I IFNs are

responsible for TLR4-mediated decreases in HIV-1 transcription. It is known that HIV-1 contains

an interferon-stimulated response element (ISRE) in the Gag-leader sequence (GLS),

immediately downstream from the 5' LTR. Because type I IFN is required for LPS- and GC-

mediated repression of HIV-1 in MDMs, we assessed the role of the ISRE in this process using

transient transfection assays with mutated LTR-reporter constructs in HEK293 cells expressing

TLR4, MD-2, and CD14. We found that LPS treatment repressed LTR-driven reporter-gene

expression in cells expressing WT ISRE elements, but not in cells transfected with an LTR-

luciferase construct containing a mutated ISRE (Supplemental Figure 5). This suggests that

transcription factor engagement of the ISRE governs TLR4-mediated HIV-1 repression.

Previous studies have shown that IRF1 and IRF2 both bind to this ISRE *in vitro* and that IRF1

and IRF2 expression are associated with enhanced HIV-1 transcription (34). Two other IRFs,

IRF4 and IRF8, are also expressed in macrophages (35) and have been shown to increase in

levels in response to type I IFN signaling and other signals (36, 37). Interestingly, IRF8 has

been implicated in maintaining HIV-1 latency in infected monocytic cell lines (34, 38, 39), and

IRF4 has been implicated in negative regulation of TLR signaling (40). We therefore

investigated whether various IRFs are recruited to the HIV-1 ISRE in response to LPS and GC

treatment in MDMs. Using chromatin immunoprecipitation analysis, we found that IRF1, IRF2,

IRF4, and IRF8 all are able to associate with the 5' LTR and GLS containing the ISRE in HIV-

infected MDMs (Figure 4C-D). Early after treatment with LPS, the levels of IRF1 associated with

this region of the viral promoter increased, whereas the levels of IRF2 and IRF4 decreased. By

24h post-treatment with LPS, the levels of IRF4 and IRF8 associated with this region increased.

Of particular note, the levels of IRF8 recruitment increased well above those seen in

unstimulated MDMs (Figure 4C). A similar pattern of IRF recruitment to the 5' LTR and GLS

occurred in GC-treated MDMs (4D), suggesting that repression of HIV-1 transcription in

response to LPS and GC treatment is due to enhanced IRF8 recruitment to the 5' LTR and

GLS. To confirm the central role of IRF8 in TLR4-mediated repression of HIV-1 expression in

MDMs, we used shRNA to knockdown IRF8 expression in MDMs (Figure 4E). Reducing IRF8

expression reversed TLR4-mediated HIV-1 repression in response to treatment with LPS or GC.

Knockdown of IRF8 led to activation of HIV-1 expression in cells treated with a combination of

PAM3CSK4 and LPS or GC, similar to that seen with treatment with PAM3CSK4 alone (Figure

4F). In contrast, overexpression of IRF8 in MDMs led to decrerased HIV-1 expression in

untreated MDMs and reversed the activation of HIV-1 expression in PAM3CSK4-treated MDMs,

but had no effect on LPS-mediated repression in MDMs (Figure 4G-H). There was a small, but

significant, enhancement of HIV-1 repression in MDMs treated with a combination of

PAM3CSK4 and LPS or with GC.

Taken together, these data suggest that both LPS and GC activate TLR4-mediated TRIF

signaling in MDMs, resulting in the production of type I IFNs. In turn, IFNs work in an autocrine

or paracrine fashion to induce the expression of IRF8, which then binds to the ISRE present in

the GLS of HIV-1 to repress viral transcription (Figure 6).

LPS and GC-treatment induces persistent low-level/latent HIV-1 infection in MDMs.

Recent studies in animals and human tissues demonstrate that HIV-1 can form persistent low-

level or latent infections in macrophages (41-44). Our data suggest that engagement of the

TLR4-TRIF-type I IFN axis in macrophages can repress virus replication and we wished to

determine whether signaling through this axis could contribute to the establishment of persistent

low-level or latent HIV-1 infection in macrophages. To this end, HIV-1-infected MDMs were

treated a single time with the TLR2 ligand PAM3CSK4, the TLR4 ligand LPS, heat-killed GC,

IFN-α, or IFN-β at day 3 post-infection. As shown in Figure 5, while there was a range of virus

replication in the various donors, we found that treatment with a single dose of LPS, heat-killed

GC, IFN-α, or IFN-β consistently led to a prominent, sustained decrease in HIV-1 replication in

MDMs, whereas treatment with PAM3CSK4 led to a transient increase in HIV-1 replication

followed by a slight decrease in replication. These data suggest that engagement of the TLR4-

TRIF-type I IFN axis can promote low-level persistent/latent HIV-1 infection in MDMs.

DISCUSSION

In these studies, we provide evidence that the interaction between commensal and pathogenic

bacteria can repress HIV-1 replication in macrophages by altering the recruitment of

transcription factors to the HIV-1 GLS, thereby inducing a state of proviral latency. We further

demonstrate that TLR2 ligands trigger MyD88-mediated signaling that increases virus

expression via the activation of NF-kB, whereas TLR4 ligands trigger TRIF-dependent

production of type I IFNs. Type I IFN signaling, in turn, is associated with the recruitment of

IRF8 to the ISRE located in the GLS and a shift to low-level or latent HIV-1 infection.

A number of studies have shown that IRFs play an important role in the regulation of HIV-1

replication. There is an ISRE located downstream from the 5' LTR in the GLS that is essential

for efficient viral replication (26, 45). This ISRE is typically bound by IRF1 and/or IRF2, leading

to activation of virus transcription (34, 46) through the recruitment of transcriptional co-

activators, such as the histone acetyl transferase (HAT) p300/CBP (47). IRF1 and IRF2 are

ubiquitously expressed in cells, thought they can be upregulated by type I IFNs (36) and, in the

case of IRF1, by TLR signaling (48, 49) and HIV-1 infection (45, 50), illustrating how HIV-1 can

co-opt the antiviral IFN response to augment its own replication. Once associated with the

ISRE, IRF1 can cooperatively bind to both NF-κB at the HIV-1 LTR and the viral transactivator

Tat at the HIV-1 TAR loop to augment viral transcription/elongation (34, 51). Our studies

demonstrate that both IRF1 and IRF2 associate with the HIV-1 ISRE in unstimulated MDMs

(Fig. 4). Upon stimulation with TLR4 ligands, IRF1recruitment to the HIV-1 ISRE is enhanced

(Fig. 4), consistent with the prevailing theory that TLR-MyD88 signaling can activate IRF1 (52).

This is accompanied by a concomitant decrease of IRF2 binding. These data suggest that IRF1

binding to the ISRE as either monomers or homodimers activates HIV-1 expression, whereas

IRF2 binding to the ISRE as monomers, homodimers, or heterodimers with IRF1 represses HIV-

1 expression. Unfortunately, ChIP analysis of HIV-infected MDMs using current tools does not

permit differentiating between the association of various homodimers and heterodimers with the

ISRE at a population level.

We demonstrate that at late time points after TLR4 engagement, IRF8 is recruited the the GLS

downstream from the 5' LTR (Figure 4), and that this is associated with decreased HIV-1

transcription (Figure 1). Macrophages express high basal levels of IRF8, although its expression

can be further enhanced in response to type I IFNs (36, 37) or TLR signaling (53, 54). IRF8 has

been shown to bind to IRF1, in addition to other transcription factors, and to serve as either a

transcriptional activator or a transcriptional inhibitor of other genes in a context-dependent

manner (55-57). Previous studies have shown that IRF8 can repress HIV-1 expression (34, 38,

39). In fact, the interaction between IRF8 and IRF1 has been shown to repress HIV-1

transcription in Jurkat cells (34). This may be due to IRF8-mediated disruption of the IRF1-Tat

interaction and/or the IRF1-NF-κB interaction (51) that increase viral replication. Based on our

data, we propose that changes in the IRF binding pattern to the ISRE in response to TLR

signaling have profound effects on HIV-1 replication. In unstimulated HIV-infected

macrophages, the ISRE is most likely occupied by IRF1/IRF2 heterodimers that allow for a low-

level of virus replication (Figure 6A). Early after stimulation of TLR4 with LPS, there is a switch

to IRF1 homodimers present at the ISRE that allow for high levels of virus replication due to

cooperative binding between IRF1, NF-κB, and HIV-1 Tat. (Figure 6B). At late time points after

TLR4 stimulation with LPS or GC, during the IFN feedback phase of the response, the ISRE is

occupied by IRF1/IRF8 heterodimers (Figure 6C). These IRF1/IRF8 heterodimers likely block

the cooperative interaction(s) between IRF1, NF-κB, and Tat, thereby repressing HIV-1

replication. Although we also demonstrate that IRF4 is recruited transiently to the HIV-1 ISRE

following treatment with LPS, the biological significance of this finding is uncertain. Prior studies

have provided evidence for an LPS/TLR4-mediated repression of HIV-1 expression through the

induction of type I IFNs and other mechanisms (16, 17, 58-62). Our data extend these findings

and demonstrate that LPS treatment, as well as infection with the sexually transmitted pathogen

GC or the gut-associated microbe *E. coli*, represses HIV-1 expression in MDMs through the

TLR4-mediated, TRIF-dependent production of type I IFNs and the subsequent recruitment of

IRF8 to the HIV-1 ISRE.

We have demonstrated that MyD88-dependent signaling activates HIV-1 expression in MDMs

by activating NF-κB, whereas TRIF-dependent signaling represses HIV-1 expression in these

cell types through the type I IFN-mediated recruitment of IRF8 to the HIV-1 ISRE. We have also

shown that TLR4-mediated, TRIF-dependent signaling is dominant in MDMs, both in the context

of treatment with purified TLR ligands as well as infection with the sexually transmitted pathogen

GC or the gut-associated microbe *E. coli*. Co-infection of virus-infected MDMs by pathogens

that express TLR4 ligands decreases HIV-1 replication through the recruitment of IRF8 to the

HIV-1 ISRE. Prior studies have provided evidence for an LPS/TLR4-mediated repression of

HIV-1 expression through the induction of type I IFNs and other mechanisms. These studies

employed HIV-1 infection of myeloid cells (16, 17, 58-61) or transfection using full-length viral

constructs (62). Our data extends these findings and demonstrates that LPS treatment

represses HIV-1 expression in MDMs through its interaction with TLR4. Specifically, we show

that LPS induces TLR4-mediated, TRIF-dependent production of type I IFNs, which in turn lead

to the repression of HIV-1 replication in MDMs through the recruitment of IRF8 to the HIV-1

ISRE.

Importantly, our data suggest that the microbial environment can influence the state of HIV-1

replication and the establishment of latency in human macrophages as part of the viral reservoir

in infected individuals under antiretroviral therapy (ART) regimens. Macrophages can be

productively infected with HIV-1 *in vivo* and viral replication can be modulated by co-pathogens

through their interactions with innate immune receptors such as TLRs (18, 63). We demonstrate

that productive infection of macrophages can be altered by TLR signaling in response to purified

ligands and bacterial co-infection, with TLR2- and TLR5-mediated signaling activating HIV-1

and TLR3- and TLR4-mediated signaling repressing HIV-1 replication in MDMs (Figure 1).

In addition to their role in HIV-1 production, macrophages also contribute to HIV-1 persistence *in*

vivo. Although CD4+ memory T cells are thought to constitute the majority of the HIV-1

reservoir, several studies have demonstrated that tissue resident macrophages in the lymph

nodes (64-66), gastrointestinal tract (5, 67), genitourinary tract (2, 42, 68), liver (69-71), and

lung (72-74), as well as perivascular macrophages and microglial cells in the brain (41, 75-80),

can serve as tissue reservoirs for HIV-1. In SHIV-infected rhesus macaques, *in vivo* viral

replication was sustained by tissue macrophages after depletion of CD4+ T cells (81).

Moreover, HIV-1 persistence in macrophages was confirmed in HIV-1 infected humanized

myeloid-only mice in which viral rebound was observed in a subset of the animals following

treatment interruption (43). These studies demonstrate that macrophages have the capacity to

serve as *bona fide* HIV-1 reservoirs *in vivo*. Our findings that pathogenic and commensal

bacteria, through engagement of TLRs, can influence HIV-1 replication in macrophages have

potential clinical significance. For example, sexually transmitted infections (STIs) that induce

robust type interferon production, such as GC or HSV-2, may repress virus replication in

genitourinary tract macrophages that harbor HIV-1 provirus and contribute to viral escape from

the immune system and from ART.

The major obstacle to the eradication of HIV-1 is the presence of a persistent viral reservoir that

that can resurface upon discontinuation of ART. The potential contribution of HIV-1 in tissue

macrophages to virus rebound with the cessation of ART is not entirely understood, but recent

primate studies suggest that the functional macrophage reservoir can contribute to viral rebound

upon treatment cessation (82-84). Our data demonstrate that interactions between

macrophages and pathogenic or commensal microorganisms within the genitourinary and

gastrointestinal tracts, such as GC and *E. coli*, may alter the ability of macrophages to serve as

reservoirs for viral persistence in the host. Our findings are consistent with independent studies

that demonstrate that repeated stimulation of M1-polarized MDMs with proinflammatory

cytokines (TNF-α) and/or type II IFNs (interferon-γ) induce a state akin to HIV-1 latency (85). In

addition, the oral pathogen *Porphyromonas gingivalis* has been shown to influence the

establishment and maintenance of persistent HIV-1 infection in MDMs (86). Finally, studies

have demonstrated that a subset of HIV-1-infected macrophages enter a state of viral latency

characterized by altered metabolic signatures (87) and apoptotic mechanisms (87). Taken

together, these studies demonstrate that co-infection, inflammatory stimuli, and metabolic

alterations can influence the establishment and maintenance of the HIV-1 reservoir in

macrophages. As an example, gastrointestinal macrophages constitute a major cellular

reservoir for HIV-1 (5, 88-90) and are frequently exposed to microbes and microbial products

either through luminal sampling (91) or microbial translocation, the latter of which is increased in

HIV-positive individuals (92). Our data suggest that interactions such as those between

intestinal macrophages and gut-associated microbes may have clinical significance for the

establishment and maintenance of the latent HIV-1 reservoir.

Our results demonstrating that Neisseria gonorrhoeae and E. coli repress HIV-1 replication in

macrophages by altering transcription factor recruitment to the HIV-1 GLS and induce a state of

viral latency confirm the need for further in vitro, ex vivo, and in vivo studies regarding the

effects of sexually-transmitted pathogens and commensal microbes on HIV-1 persistence.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge Dr. Laura Dickey for her scientific and editorial suggestions.

The following reagents were obtained through the AIDS Research and Reference Reagent

Program, Division of AIDS, NIAID, NIH: HIV IG from NABI and National Heart Lung and Blood

Institute (Dr. Luiz Barbosa); anti-p24gag monoclonal antibody clone 183-H12-5C from Dr. Bruce

Chesebro and Kathy Wehrly; and MAGI-CCR5 cells from Dr. Julie Overbaugh. The model figure

was created with Biorender.com.

This work was supported by funds obtained from the National Institutes of Health (www.nih.gov)

grants AI073149, (G.V.), AI143567-02 (V.P.), T32-AI07309 (T.M.H.), and T32-AI0764206

(T.M.H.), a Boston University Division of Graduate Medical Sciences Graduate Student

Research Fellowship award (T.M.H.), and a Resident Research Award from the Department of

Pathology at the University of Utah (T.M.H.).

DISCLOSURES

The authors have no conflicts of interest.

AUTHOR CONTRIBUTIONS

T.M.H. and G.A.V. designed the study. T.M.H and G.A.V. developed the methodology. T.M.H.

conducted experiments. T.M.H., V.P., and G.A.V. wrote the manuscript. T.M.H., V.P., and

G.A.V. acquired funds. T.M.H. and G.A.V. supervised the study.

FIGURE LEGENDS

Figure 1. Treatment with purified TLR ligands and intact bacteria alter HIV-1 replication at the level of transcription. (A) MDMs (2.5×10^5 cells/well) were infected with a single-round, replication-defective HIV-luciferase reporter virus at an MOI = 0.1. After four hours, unbound virus was removed by washing with PBS and cells were cultured in complete medium. Forty-eight hours after infection, cells were treated with the TLR2 ligand PAM3CSK4 (100 ng/ml), the TLR3 ligand poly I:C (25 μg/ml), the TLR4 ligand LPS (100 ng/ml), or the TLR5 ligand FLA-ST (100 ng/ml) for 18 hours. The cells were then lysed and assayed for luciferase activity. Bars represent the mean (± SD) of 11 donors, each donor tested in triplicate. (B-D) MDMs were infected as described above. Forty-eight hours after infection, cells were treated with PAM3CSK4 (100 ng/ml) or LPS (100 ng/ml) for six hours (B-C) or 24 hours (D). Cells were then lysed and assayed for viral RNA accumulation by RT-PCR. Shown are data from one representative donor (B) and composite data from eight donors at 6 hours (C) and four donors at 24 hours (D). (E) MDMs (1×10^6 cells/well) were infected as in (A). Forty-eight hours after infection, cells were treated with the TLR4 ligand LPS (100 ng/ml) for 4 h. Cells were then treated with actinomycin D (10 μg/ml) to inhibit transcription. Total cytoplasmic RNA was prepared from the treated cultures at the indicated time points following actinomycin D treatment and analyzed by RT-qPCR for the expression of HIV-1 RNA. The data are the means (± SD) from four donors. (F) MDMs were infected as in (A). Forty-eight hours after infection, the cells

were cultured overnight with increasing amounts of GC. Cells were then lysed and assayed for

luciferase activity. Bars represent mean (± SD) of seven donors, each donor tested in triplicate.

(G) MDMs were infected as described above. Forty-eight hours after infection, cells were

treated with GC at an MOI of 10 for 24 hours. Cells were then lysed and assayed for viral RNA

accumulation by RT-qPCR. Shown are data from four donors. (H) MDMs (2.5 x 10^5 cells/well)

were infected as in (A). Forty-eight hours after infection, the cells were cultured overnight with

increasing amounts of *E. coli*. Cells were then lysed and assayed for luciferase activity. Bars

represent mean (± SD) of four donors, each donor tested in triplicate. *, $p < 0.05$; **, $p < 0.01$;

***, $p < 0.001$; $p < 0.0001$.

Figure 2. TLR4 signaling is dominant in MDMs. (A) MDMs (2.5 x 10^5 cells/well) were infected

with a single-round, replication-defective HIV-luciferase reporter virus (MOI = 0.1). Forty-eight

hours after infection, cells were treated with a fixed concentration of PAM3CSK4 (100 ng/ml)

and increasing concentrations of LPS (1-1000 ng/ml, as indicated) or a fixed concentration of

LPS (100 ng/ml) and increasing concentrations of PAM3CSK4 (1-1000 ng/ml, as indicated) for

18 hours. Cells were then lysed and assayed for luciferase activity. The data are the mean (±

SD) of six donors, each donor tested in triplicate. (B-C) At eight days post-isolation, MDMs were

stained with antibodies against TLR2 or TLR4 or relevant isotype controls. Receptor expression

was assessed by flow cytometry. Histograms from one representative donor are shown in (B).

Grey, unstained cells; black line, isotype control; red line, TLR4; green line, TLR2. Mean

fluorescent intensity (MFI) ± SD from eight donors is depicted in (C). (D-E) MDMs (2.5 x 10^5

cells/well) were treated with the TLR2 ligand PAM3CSK4 (100 ng/ml), the TLR4 ligand LPS

(100 ng/ml), heat-killed GC (MOI = 10), or heat-killed *E. coli* (MOI = 10) for 18 hours. Cell

supernatant was harvested, filtered through a 0.2 μm filter, and analyzed by ELISA for TNF-α

(D) and IFN-β (E) production. Data represent mean ($\pm$ SD) of seven donors (four donors for

heat-killed *E. coli*). *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; $p < 0.0001$.

Figure 3. LPS- and GC-mediated repression of HIV-1 replication in MDMs requires TLR4,

TRIF, and type I IFNs. (A) MDMs (2.5×10^5 cells/well) were infected with a single-round,

replication-defective HIV-luciferase reporter virus (MOI = 0.1). Forty-eight hours after infection,

cells were treated with GC (MOI = 10) in the absence (white bars) or presence (black bars) of

B18R (100 ng/ml) for 18 hours. The cells were then lysed and assayed for luciferase activity.

The data are the mean ($\pm$ SD) of seven donors, each donor tested in triplicate. (B) MDMs (2.5 x

10^5 cells/well) were infected as above. Forty-eight hours after infection, cells were treated with

PAM3CSK4 (100 ng/ml), LPS (100 ng/ml), or heat-killed GC (MOI = 10) in the absence (white

bars) or presence (black bars) of TAK242 (1 μg/ml) for 18 hours. The cells were then lysed and

assayed for luciferase activity. The data are the mean ($\pm$ SD) of six donors, each donor tested in

triplicate. (C) MDMs (2.5×10^5 cells/well) were infected as above. Forty-eight hours after

infection, cells were treated with vehicle control (white bars) or with (black bars) the dynamin

inhibitor Dynasore (80 μM) for 15 minutes prior to treatment with heat-killed GC (MOI = 10) for

18 hours. The cells were then lysed and assayed for luciferase activity. The data are the mean

($\pm$ SD) of six donors, each donor tested in triplicate. (D-E) MDMs (2×10^6 cells/well) were

transfected with a control scrambled shRNA, shRNA targeting MyD88, or shRNA targeting

TRIF. Knock-down of protein expression was detected by western blot (D). Transfected MDMs

were infected with a single-round, replication-defective HIV-luciferase reporter virus at an MOI =

0.1. Forty-eight hours after infection, cells were treated with PAM3CSK4 (100 ng/ml), LPS (100

ng/ml), heat-killed GC (MOI = 10), or PMA (10 nM) for 18 hours. The cells were then lysed and

assayed for luciferase activity (E). The data are the mean ($\pm$ SD) of six donors. *, $p < 0.05$; **, p

< 0.01; ***, $p < 0.001$; $p < 0.0001$.

Figure 4. LPS- and GC-mediated repression of HIV-1 in MDMs is associated with changes

in IRF recruitment to the ISRE. (A) MDMs ($2x10^6$ cells/well) were infected with a single-round,

replication-defective HIV-luciferase reporter virus (MOI = 0.1). Forty-eight hours after infection,

cells were treated with PAM3CSK4 (100 ng/ml) or LPS (100 ng/ml). At various time points after

TLR stimulation, cells were harvested, lysed, and total cytoplasmic RNA was extracted. Viral

RNA accumulation was assessed by RT-PCR. The data are the mean ($\pm$ SD) of four donors. (B-

C) MDMs (1.2×10^7 cells/plate) were infected with a single-round replication-defective HIV-GFP

reporter virus at an MOI = 2. Forty-eight hours after infection, cells were treated with LPS (100

ng/ml). At either 1 or 24 hours after LPS treatment, cells were fixed with formaldehyde, lysed,

sonicated, and subjected to immunoprecipitation with antibodies against IRF1, IRF2, IRF4,

IRF8, or rabbit IgG (isotype control). Association with the HIV-1 ISRE was assessed by PCR

using HIV-1 specific primers. Data from one representative donor is depicted in (B). Composite

data representing the mean ($\pm$ SD) from five donors are showin in (C). (D) MDMs (1.2×10^7

cells/plate) were infected with a single-round replication-defective HIV-GFP reporter virus at an

MOI = 2. Forty-eight hours after infection, cells were treated with heat-killed GC (MOI = 10).

Twenty-four hours after GC treatment, cells were fixed with formaldehyde, lysed, sonicated, and

subjected to immunoprecipitation with antibodies against IRF1, IRF2, IRF4, IRF8, or rabbit IgG

(isotype control). Association with the HIV-1 ISRE was assessed by PCR using HIV-1 specific

primers. Composite data from five donors are shown. (E-F) MDMs ($2x10^6$ cells/well) transfected

with a controlled scrambled shRNA (white bars) or with shRNA targeting IRF8 (black bars) were

infected with a single-round, replication-defective HIV-luciferase reporter virus at an MOI = 0.1.

Knock-down of protein expression was detected by western blot (F). Forty-eight hours after

infection, cells were treated with PAM3CSK4 (100 ng/ml), LPS (100 ng/ml, a combination of

PAM3CSK4 and LPS (each at 100 ng/ml), or GC (MOI = 10) for 18 hours. Cells were then lysed

and assayed for luciferase activity (F). The experiment was performed using cells from five

different donors. (G-H) MDMs transfected with an empty vector (white bars) or a vector

encoding IRF8 (black bars) were infected with a single-round, replication-defective HIV-

luciferase reporter virus at an MOI = 0.1. IRF8 protein expression was detected by western blot

(G). Forty-eight hours after infection, cells were treated with PAM3CSK4 (100 ng/ml), LPS (100

ng/ml, a combination of PAM3CSK4 and LPS (each at 100 ng/ml), or GC (MOI = 10) for 18

hours. Cells were then lysed and assayed for luciferase activity (H). The experiment was

performed using cells from four different donors. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; $p <$

0.0001.

Figure 5. Treatment with LPS, GC, or type I IFNs induces a low-level persistent/latent

infection in MDMs. MDMs (2.5×10^5 cells/well) were infected with replication-competent HIV-1$_{Ba-}$

$_L$ (MOI of 0.1). At day 3 post-infection, the cells were treated with a single dose of PAM3CSK4

(100 ng/ml), LPS (100 ng/ml), GC (MOI = 10), IFN-α (1000 U/ml), or IFN-β (1000 U/ml). Cell-

free supernatants were harvested every three days, and virus production was monitored by p24

ELISA. Data from five independent donors, each tested in triplicate, are shown. *, $p < 0.05$; **, p

< 0.01; ***, $p < 0.001$; $p < 0.0001$.

Figure 6. Co-infection with GC or *E. coli* represses HIV-1 replication by altering IRF

recruitment to the HIV-1 GLS. (A) Upon engagement of TLR2 and/or TLR4 by GC or *E. coli* at

the cell surface, transcription factors such as NF-κB are phosphorylated and are subsequently

recruited the HIV-1 5' LTR to drive viral transcription in MDMs. Upon engagement of TLR4 in

the endosome, IRF3 and IRF7 are phosphorylated and recruited to the IFN-α and/or IFN-β

promoters to drive type I IFN expression. (B) During the early phase of the response, type I

IFNs drive the expression of ISGs, including IRFs. Signaling through endosomal TLR4 leads to

the activation of IRF1 and its recruitment to the HIV-1 GLS, further enhancing HIV-1

transcription. (C) During the late phase of the response there is a type I IFN-dependent

activation of IRF8 and subsequent recruitment to the HIV-1 GLS, thereby repressing HIV-1

transcription.

Supplemental Figure 1. Altered HIV-1 expression in MDMs in response to TLR

engagement is not sex-dependent. MDMs (2.5×10^5 cells/well) were infected with a single-

round, replication-defective HIV-luciferase reporter virus (MOI = 0.1). After four hours, unbound

virus was removed by washing with PBS and cells were cultured in complete medium. Forty-

eight hours after infection, cells were treated with the TLR2 ligand PAM3CSK4 (100 ng/ml), the

TLR3 ligand poly I:C (25 μg/ml), the TLR4 ligand LPS (100 ng/ml), or the TLR5 ligand FLA-ST

(100 ng/ml) for 18 hours. The cells were then lysed and assayed for luciferase activity. Bars

represent the mean (± SD) of five male donors and five female donors; each donor tested in

triplicate. Although virus replication was decreased overall in MDMs from female donors

compared to MDMs from male donors, it was activated by treatment with PAM3CSK4 and FLA-

ST, and repressed by Poly I:C and LPS, in a manner similar to that seen in MDMs from male

donors. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; $p < 0.0001$.

Supplemental Figure 2. TLR2-activated HIV-1 expression is mediated primarily through

NF-κB. (A-B) MDMs (2.5×10^5 cells/well) were infected with a single-round, replication-defective

HIV-luciferase reporter virus (MOI = 0.1). Forty-eight hours after infection, cells were treated

with PAM3CSK4 (100 ng/ml) in the presence or absence of 10 μM celastrol (A) or 10 μM BAY

11-7082 (B) for 18 hours. The cells were then lysed and assayed for luciferase activity. The data

are the mean (± SD) of six donors (celastrol) or three donors (BAY 11-7082); each donor tested

in triplicate. (C) HEK293-TLR2^{CFP}TLR1^{YFP} cells were transfected with HIV-1 LTR-luciferase

reporter constructs with intact NF-κB, mutated NF-κB, or deleted NF-κB binding sites. Following

transfection, cells were treated with PAM3CSK4 (100 ng/ml) for 18 hours and then harvested

and assayed for luciferase activity. Data represent the mean (± SD) of three independent

experiments, each performed in triplicate. (D) MDMs (2.5×10^5 cells/well) were infected as in

(A). Forty-eight hours after infection, cells were treated with PAM3CSK4 (100 ng/ml) in the

presence or absence of U0126 (10 μM), PD98059 (50 μM), or SB203580 (10 μM) for 18 hours.

Cells were then lysed and assayed for luciferase activity. The data are the mean (± SD) of six

donors; each donor tested in triplicate. (E) HEK293-TLR2^{CFP}TLR1^{YFP} cells were transfected with

HIV-1 LTR-luciferase reporter constructs with intact AP-1 sites (WT LTR) or deleted AP-1

binding sites (-158 LTR). Following transfection, cells were treated with PAM3CSK4 (100 ng/ml)

for 18 hours and then harvested and assayed for luciferase activity. Data are the mean (± SD)

of three independent experiments, each performed in triplicate. *, $p < 0.05$; **, $p < 0.01$; ***, $p <$

0.001; $p < 0.0001$.

Supplemental figure 3. Heat-killed GC represses HIV-1 replication in MDMs, but activates

it in MDDCs. (A) MDMs were infected with a single-round, replication-defective HIV-1 luciferase

reporter virus (MOI = 0.1). Forty-eight hours after infection, the cells were cultured with

increasing amounts of live or heat-killed (56°C treatment) GC overnight. The cells were then

lysed and luciferase activity was measured. The data are the mean (± SD) of three donors, each

donor tested in triplicate. (B) MDMs (1×10^6 cells/well) were infected as in (A). Forty-eight hours

after infection, cells were treated with heat-killed GC (MOI = 10) for 4 h. Cells were then treated

with actinomycin D (10 μg/ml) to inhibit transcription. Total cytoplasmic RNA was prepared from

the treated cultures at the indicated time points following actinomycin D treatment and analyzed

by RT-qPCR for the expression of HIV-1 RNA. The data are the means (± SD) from four donors.

(C) MDMs were infected as in (A). Forty-eight hours after infection, cells were treated with heat-

killed GC (MOI = 10) or heat-killed *E. coli* (MOI = 10) for 18 hours. The cells were then lysed

and assayed for luciferase activity. Bars represent the mean (± SD) of five male donors and five

female donors; each donor tested in triplicate. Although virus replication was decreased overall

in MDMs from female donors compared to MDMs from male donors, it was repressed by HKGC

and HKEC in a manner similar to that seen in MDMs from male donors. *, $p < 0.05$; **, $p < 0.01$;

***, $p < 0.001$; $p < 0.0001$.

Supplemental figure 4. Dynasore inhibits endocytosis/phagocytosis of labeled *E. coli*

particles by MDMs. (A-B) MDMs ($5x10^5$/well) were incubated with DMSO or Dynasore (80 μM)

for 15 minutes at 37°C. The cells were washed with PBS and incubated with pHrodo Green *E.*

coli (1 mg/ml) for 2 hours at 37°C. Endocytosis/phagocytosis was measured by flow cytometry.

Shown are data from one representative donor (A) and composite data from four donors (B). *,

$p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; $p < 0.0001$.

Supplemental figure 5. TLR4-mediated repression of HIV-1 requires the ISRE binding site

located in the GLS downstream of the 5' LTR. (A) HEK293-TLR4^{CFP}/MD-2/CD14 cells were

transfected with HIV-1 LTR/GLS-luciferase reporter constructs with an intact ISRE or mutated

ISRE binding site. Following transfection, cells were treated with LPS (100 ng/ml) for 18 hours

and then harvested and assayed for luciferase activity. Data are the mean (± SD) of three

independent experiments, each performed in triplicate. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; p

< 0.0001.

Table 1. Primers used for PCR-based mutagenesis.

Primer Name	Sequence

Forward mutI NF-κB	GGACTTTCCGCTGTCTACTTTCCAGG
Reverse mutI NF-κB	CCTGGAAAGTAGACAGCGGAAAGTCC
Forward mutII NF-κB	GCTTTCTACAATCTACTTTCCGCTGG
Reverse mutII NF-κB	CCAGCGGAAAGTAGATTGTAGAAAGC
Forward mutI&II NF-κB	GCTTTCTACAATCTACTTTCCGCTGTCTACTTTCCAGG
Reverse mutI&II NF-κB	CCTGGAAAGTAGACAGCGGAAAGTAGATTGTAGAAAGC
Forward delNF-κB	GCTGACATCGAGCTTTCTACAAAGGGAGGTGTGGCCTGGGCGGG
Reverse delNF-κB	CCCGCCCAGGCCACACCTCCCTTTGTAGAAAGCTCGATGTCAGC
Forward mutISRE	GCCCGAACAGGGACTTGCCCGCGCCCGTAAAGCCAGAGGAGATC
Reverse mutISRE	GATCCTCTGGCTTTACGGGCGCGGGCAAGTCCCTGTTCGGGC

1. Miller CJ, Shattock RJ. 2003. Target cells in vaginal HIV transmission. Microbes Infect
5:59-67.
2. Shen R, Richter HE, Clements RH, Novak L, Huff K, Bimczok D, Sankaran-Walters S,
Dandekar S, Clapham PR, Smythies LE, Smith PD. 2009. Macrophages in vaginal but not

intestinal mucosa are monocyte-like and permissive to human immunodeficiency virus
type 1 infection. J Virol 83:3258-67.

3.   McElrath MJ, Smythe K, Randolph-Habecker J, Melton KR, Goodpaster TA, Hughes SM,
Mack M, Sato A, Diaz G, Steinbach G, Novak RM, Curlin ME, Lord JD, Maenza J, Duerr A,
Frahm N, Hladik F, Network NHVT. 2013. Comprehensive assessment of HIV target cells
in the distal human gut suggests increasing HIV susceptibility toward the anus. J Acquir
Immune Defic Syndr 63:263-71.

4.   Shen R, Richter HE, Smith PD. 2011. Early HIV-1 target cells in human vaginal and
ectocervical mucosa. Am J Reprod Immunol 65:261-7.

5.   Zalar A, Figueroa MI, Ruibal-Ares B, Bare P, Cahn P, de Bracco MM, Belmonte L. 2010.
Macrophage HIV-1 infection in duodenal tissue of patients on long term HAART.
Antiviral Res 87:269-71.

6.   Wong ME, Jaworowski A, Hearps AC. 2019. The HIV Reservoir in Monocytes and
Macrophages. Front Immunol 10:1435.

7.   Galvin SR, Cohen MS. 2004. The role of sexually transmitted diseases in HIV
transmission. Nat Rev Microbiol 2:33-42.

8.   Malott RJ, Keller BO, Gaudet RG, McCaw SE, Lai CC, Dobson-Belaire WN, Hobbs JL, St
Michael F, Cox AD, Moraes TF, Gray-Owen SD. 2013. Neisseria gonorrhoeae-derived
heptose elicits an innate immune response and drives HIV-1 expression. Proc Natl Acad
Sci U S A 110:10234-9.

9.   Jarvis GA, Chang TL. 2012. Modulation of HIV transmission by Neisseria gonorrhoeae:
molecular and immunological aspects. Curr HIV Res 10:211-7.

10. Ferreira VH, Nazli A, Khan G, Mian MF, Ashkar AA, Gray-Owen S, Kaul R, Kaushic C. 2011.
Endometrial epithelial cell responses to coinfecting viral and bacterial pathogens in the
genital tract can activate the HIV-1 LTR in an NF{kappa}B-and AP-1-dependent manner. J
Infect Dis 204:299-308.

11. Ding J, Rapista A, Teleshova N, Mosoyan G, Jarvis GA, Klotman ME, Chang TL. 2010.
Neisseria gonorrhoeae enhances HIV-1 infection of primary resting CD4+ T cells through
TLR2 activation. J Immunol 184:2814-24.

12. Zhang J, Li G, Bafica A, Pantelic M, Zhang P, Broxmeyer H, Liu Y, Wetzler L, He JJ, Chen T.
2005. Neisseria gonorrhoeae enhances infection of dendritic cells by HIV type 1. J
Immunol 174:7995-8002.

13. Sanyal A, Shen C, Ding M, Reinhart TA, Chen Y, Sankapal S, Gupta P. 2019. Neisseria
gonorrhoeae uses cellular proteins CXCL10 and IL8 to enhance HIV-1 transmission across
cervical mucosa. Am J Reprod Immunol 81:e13111.

14. Mogensen TH, Paludan SR, Kilian M, Ostergaard L. 2006. Live Streptococcus
pneumoniae, Haemophilus influenzae, and Neisseria meningitidis activate the
inflammatory response through Toll-like receptors 2, 4, and 9 in species-specific
patterns. J Leukoc Biol 80:267-77.

15. Dobson-Belaire WN, Rebbapragada A, Malott RJ, Yue FY, Kovacs C, Kaul R, Ostrowski
MA, Gray-Owen SD. 2010. Neisseria gonorrhoeae effectively blocks HIV-1 replication by
eliciting a potent TLR9-dependent interferon-alpha response from plasmacytoid
dendritic cells. Cell Microbiol 12:1703-17.

16. Liu X, Mosoian A, Li-Yun Chang T, Zerhouni-Layachi B, Snyder A, Jarvis GA, Klotman ME.
2006. Gonococcal lipooligosaccharide suppresses HIV infection in human primary
macrophages through induction of innate immunity. J Infect Dis 194:751-9.
17. Kornbluth RS, Oh PS, Munis JR, Cleveland PH, Richman DD. 1989. Interferons and
bacterial lipopolysaccharide protect macrophages from productive infection by human
immunodeficiency virus in vitro. J Exp Med 169:1137-51.
18. Hanley TM, Viglianti GA. 2011. Nuclear receptor signaling inhibits HIV-1 replication in
macrophages through multiple trans-repression mechanisms. J Virol 85:10834-50.
19. Novis CL, Archin NM, Buzon MJ, Verdin E, Round JL, Lichterfeld M, Margolis DM,
Planelles V, Bosque A. 2013. Reactivation of latent HIV-1 in central memory CD4(+) T
cells through TLR-1/2 stimulation. Retrovirology 10:119.
20. Szaniawski MA, Spivak AM, Bosque A, Planelles V. 2019. Sex Influences SAMHD1 Activity
and Susceptibility to Human Immunodeficiency Virus-1 in Primary Human Macrophages.
J Infect Dis 219:777-785.
21. Ziegler S, Altfeld M. 2016. Sex differences in HIV-1-mediated immunopathology. Curr
Opin HIV AIDS 11:209-15.
22. Rodriguez-Garcia M, Biswas N, Patel MV, Barr FD, Crist SG, Ochsenbauer C, Fahey JV,
Wira CR. 2013. Estradiol reduces susceptibility of CD4+ T cells and macrophages to HIV-
infection. PLoS One 8:e62069.
23. Kawai T, Akira S. 2006. TLR signaling. Cell Death Differ 13:816-25.
24. Nabel G, Baltimore D. 1987. An inducible transcription factor activates expression of
human immunodeficiency virus in T cells. Nature 326:711-3.
25. Perkins ND, Edwards NL, Duckett CS, Agranoff AB, Schmid RM, Nabel GJ. 1993. A
cooperative interaction between NF-kappa B and Sp1 is required for HIV-1 enhancer
activation. EMBO J 12:3551-8.
26. Van Lint C, Amella CA, Emiliani S, John M, Jie T, Verdin E. 1997. Transcription factor
binding sites downstream of the human immunodeficiency virus type 1 transcription
start site are important for virus infectivity. J Virol 71:6113-27.
27. Canonne-Hergaux F, Aunis D, Schaeffer E. 1995. Interactions of the transcription factor
AP-1 with the long terminal repeat of different human immunodeficiency virus type 1
strains in Jurkat, glial, and neuronal cells. J Virol 69:6634-42.
28. Pierce JW, Schoenleber R, Jesmok G, Best J, Moore SA, Collins T, Gerritsen ME. 1997.
Novel inhibitors of cytokine-induced IkappaBalpha phosphorylation and endothelial cell
adhesion molecule expression show anti-inflammatory effects in vivo. J Biol Chem
272:21096-103.
29. Lee JH, Koo TH, Yoon H, Jung HS, Jin HZ, Lee K, Hong YS, Lee JJ. 2006. Inhibition of NF-
kappa B activation through targeting I kappa B kinase by celastrol, a quinone methide
triterpenoid. Biochem Pharmacol 72:1311-21.
30. Ono Y, Maejima Y, Saito M, Sakamoto K, Horita S, Shimomura K, Inoue S, Kotani J. 2020.
TAK-242, a specific inhibitor of Toll-like receptor 4 signalling, prevents endotoxemia-
induced skeletal muscle wasting in mice. Sci Rep 10:694.
31. Kagan JC, Su T, Horng T, Chow A, Akira S, Medzhitov R. 2008. TRAM couples endocytosis
of Toll-like receptor 4 to the induction of interferon-beta. Nat Immunol 9:361-8.

32. Nasr N, Alshehri AA, Wright TK, Shahid M, Heiner BM, Harman AN, Botting RA, Helbig KJ,
Beard MR, Suzuki K, Kelleher AD, Hertzog P, Cunningham AL. 2017. Mechanism of
Interferon-Stimulated Gene Induction in HIV-1-Infected Macrophages. J Virol 91.
33. Sheikh F, Dickensheets H, Gamero AM, Vogel SN, Donnelly RP. 2014. An essential role
for IFN-beta in the induction of IFN-stimulated gene expression by LPS in macrophages. J
Leukoc Biol 96:591-600.
34. Sgarbanti M, Borsetti A, Moscufo N, Bellocchi MC, Ridolfi B, Nappi F, Marsili G, Marziali
G, Coccia EM, Ensoli B, Battistini A. 2002. Modulation of human immunodeficiency virus
1 replication by interferon regulatory factors. J Exp Med 195:1359-70.
35. Yamamoto M, Kato T, Hotta C, Nishiyama A, Kurotaki D, Yoshinari M, Takami M, Ichino
M, Nakazawa M, Matsuyama T, Kamijo R, Kitagawa S, Ozato K, Tamura T. 2011. Shared
and distinct functions of the transcription factors IRF4 and IRF8 in myeloid cell
development. PLoS One 6:e25812.
36. Barber SA, Fultz MJ, Salkowski CA, Vogel SN. 1995. Differential expression of interferon
regulatory factor 1 (IRF-1), IRF-2, and interferon consensus sequence binding protein
genes in lipopolysaccharide (LPS)-responsive and LPS-hyporesponsive macrophages.
Infect Immun 63:601-8.
37. Gupta M, Shin DM, Ramakrishna L, Goussetis DJ, Platanias LC, Xiong H, Morse HC, 3rd,
Ozato K. 2015. IRF8 directs stress-induced autophagy in macrophages and promotes
clearance of Listeria monocytogenes. Nat Commun 6:6379.
38. Thornton AM, Buller RM, DeVico AL, Wang IM, Ozato K. 1996. Inhibition of human
immunodeficiency virus type 1 and vaccinia virus infection by a dominant negative
factor of the interferon regulatory factor family expressed in monocytic cells. Proc Natl
Acad Sci U S A 93:383-7.
39. Munier S, Delcroix-Genete D, Carthagena L, Gumez A, Hazan U. 2005. Characterization
of two candidate genes, NCoA3 and IRF8, potentially involved in the control of HIV-1
latency. Retrovirology 2:73.
40. Negishi H, Ohba Y, Yanai H, Takaoka A, Honma K, Yui K, Matsuyama T, Taniguchi T,
Honda K. 2005. Negative regulation of Toll-like-receptor signaling by IRF-4. Proc Natl
Acad Sci U S A 102:15989-94.
41. Abreu C, Shirk EN, Queen SE, Beck SE, Mangus LM, Pate KAM, Mankowski JL, Gama L,
Clements JE. 2019. Brain macrophages harbor latent, infectious simian
immunodeficiency virus. Aids 33 Suppl 2:S181-s188.
42. Ganor Y, Real F, Sennepin A, Dutertre CA, Prevedel L, Xu L, Tudor D, Charmeteau B,
Couedel-Courteille A, Marion S, Zenak AR, Jourdain JP, Zhou Z, Schmitt A, Capron C,
Eugenin EA, Cheynier R, Revol M, Cristofari S, Hosmalin A, Bomsel M. 2019. HIV-1
reservoirs in urethral macrophages of patients under suppressive antiretroviral therapy.
Nat Microbiol 4:633-644.
43. Honeycutt JB, Thayer WO, Baker CE, Ribeiro RM, Lada SM, Cao Y, Cleary RA, Hudgens
MG, Richman DD, Garcia JV. 2017. HIV persistence in tissue macrophages of humanized
myeloid-only mice during antiretroviral therapy. Nat Med 23:638-643.
44. Smith PD, Meng G, Salazar-Gonzalez JF, Shaw GM. 2003. Macrophage HIV-1 infection
and the gastrointestinal tract reservoir. J Leukoc Biol 74:642-9.

45.   Harman AN, Lai J, Turville S, Samarajiwa S, Gray L, Marsden V, Mercier SK, Jones K, Nasr
931   N, Rustagi A, Cumming H, Donaghy H, Mak J, Gale M, Jr., Churchill M, Hertzog P,
932   Cunningham AL. 2011. HIV infection of dendritic cells subverts the IFN induction
933   pathway via IRF-1 and inhibits type 1 IFN production. Blood 118:298-308.

934   46.   Marsili G, Borsetti A, Sgarbanti M, Remoli AL, Ridolfi B, Stellacci E, Ensoli B, Battistini A.
935   2003. On the role of interferon regulatory factors in HIV-1 replication. Ann N Y Acad Sci
936   1010:29-42.

937   47.   Marsili G, Remoli AL, Sgarbanti M, Battistini A. 2004. Role of acetylases and deacetylase
938   inhibitors in IRF-1-mediated HIV-1 long terminal repeat transcription. Ann N Y Acad Sci
939   1030:636-43.

940   48.   Hu Y, Park-Min KH, Yarilina A, Ivashkiv LB. 2008. Regulation of STAT pathways and IRF1
941   during human dendritic cell maturation by TNF-alpha and PGE2. J Leukoc Biol 84:1353-
942   60.

943   49.   Liljeroos M, Vuolteenaho R, Rounioja S, Henriques-Normark B, Hallman M, Ojaniemi M.
944   2008. Bacterial ligand of TLR2 signals Stat activation via induction of IRF1/2 and
945   interferon-alpha production. Cell Signal 20:1873-81.

946   50.   Sgarbanti M, Marsili G, Remoli AL, Ridolfi B, Stellacci E, Borsetti A, Ensoli B, Battistini A.
947   2004. Analysis of the signal transduction pathway leading to human immunodeficiency
948   virus-1-induced interferon regulatory factor-1 upregulation. Ann N Y Acad Sci 1030:187-
949   95.

950   51.   Sgarbanti M, Remoli AL, Marsili G, Ridolfi B, Borsetti A, Perrotti E, Orsatti R, Ilari R,
951   Sernicola L, Stellacci E, Ensoli B, Battistini A. 2008. IRF-1 is required for full NF-kappaB
952   transcriptional activity at the human immunodeficiency virus type 1 long terminal
953   repeat enhancer. J Virol 82:3632-41.

954   52.   Negishi H, Fujita Y, Yanai H, Sakaguchi S, Ouyang X, Shinohara M, Takayanagi H, Ohba Y,
955   Taniguchi T, Honda K. 2006. Evidence for licensing of IFN-gamma-induced IFN regulatory
956   factor 1 transcription factor by MyD88 in Toll-like receptor-dependent gene induction
957   program. Proc Natl Acad Sci U S A 103:15136-41.

958   53.   Tamura T, Ozato K. 2002. ICSBP/IRF-8: its regulatory roles in the development of
959   myeloid cells. J Interferon Cytokine Res 22:145-52.

960   54.   Zhao J, Kong HJ, Li H, Huang B, Yang M, Zhu C, Bogunovic M, Zheng F, Mayer L, Ozato K,
961   Unkeless J, Xiong H. 2006. IRF-8/interferon (IFN) consensus sequence-binding protein is
962   involved in Toll-like receptor (TLR) signaling and contributes to the cross-talk between
963   TLR and IFN-gamma signaling pathways. J Biol Chem 281:10073-80.

964   55.   Bovolenta C, Driggers PH, Marks MS, Medin JA, Politis AD, Vogel SN, Levy DE, Sakaguchi
965   K, Appella E, Coligan JE, et al. 1994. Molecular interactions between interferon
966   consensus sequence binding protein and members of the interferon regulatory factor
967   family. Proc Natl Acad Sci U S A 91:5046-50.

968   56.   Meraro D, Hashmueli S, Koren B, Azriel A, Oumard A, Kirchhoff S, Hauser H, Nagulapalli
969   S, Atchison ML, Levi BZ. 1999. Protein-protein and DNA-protein interactions affect the
970   activity of lymphoid-specific IFN regulatory factors. J Immunol 163:6468-78.

971   57.   Langlais D, Barreiro LB, Gros P. 2016. The macrophage IRF8/IRF1 regulome is required
972   for protection against infections and is associated with chronic inflammation. J Exp Med
973   213:585-603.

58.   Ahmed N, Hayashi T, Hasegawa A, Furukawa H, Okamura N, Chida T, Masuda T, Kannagi
M. 2010. Suppression of human immunodeficiency virus type 1 replication in
macrophages by commensal bacteria preferentially stimulating Toll-like receptor 4. J
Gen Virol 91:2804-13.
59.   Bernstein MS, Tong-Starksen SE, Locksley RM. 1991. Activation of human monocyte-
derived macrophages with lipopolysaccharide decreases human immunodeficiency virus
replication in vitro at the level of gene expression. J Clin Invest 88:540-5.
60.   Equils O, Salehi KK, Cornateanu R, Lu D, Singh S, Whittaker K, Baldwin GC. 2006.
Repeated lipopolysaccharide (LPS) exposure inhibits HIV replication in primary human
macrophages. Microbes Infect 8:2469-76.
61.   Simard S, Maurais E, Gilbert C, Tremblay MJ. 2008. LPS reduces HIV-1 replication in
primary human macrophages partly through an endogenous production of type I
interferons. Clin Immunol 127:198-205.
62.   Nordone SK, Ignacio GA, Su L, Sempowski GD, Golenbock DT, Li L, Dean GA. 2007. Failure
of TLR4-driven NF-kappa B activation to stimulate virus replication in models of HIV type
1 activation. AIDS Res Hum Retroviruses 23:1387-95.
63.   Appelberg KS, Wallet MA, Taylor JP, Cash MN, Sleasman JW, Goodenow MM. 2017. HIV-
1 Infection Primes Macrophages Through STAT Signaling to Promote Enhanced
Inflammation and Viral Replication. AIDS Res Hum Retroviruses 33:690-702.
64.   Embretson J, Zupancic M, Ribas JL, Burke A, Racz P, Tenner-Racz K, Haase AT. 1993.
Massive covert infection of helper T lymphocytes and macrophages by HIV during the
incubation period of AIDS. Nature 362:359-62.
65.   Orenstein JM, Fox C, Wahl SM. 1997. Macrophages as a source of HIV during
opportunistic infections. Science 276:1857-61.
66.   DiNapoli SR, Ortiz AM, Wu F, Matsuda K, Twigg HL, 3rd, Hirsch VM, Knox K, Brenchley
JM. 2017. Tissue-resident macrophages can contain replication-competent virus in
antiretroviral-naive, SIV-infected Asian macaques. JCI Insight 2:e91214.
67.   Yukl SA, Sinclair E, Somsouk M, Hunt PW, Epling L, Killian M, Girling V, Li P, Havlir DV,
Deeks SG, Wong JK, Hatano H. 2014. A comparison of methods for measuring rectal HIV
levels suggests that HIV DNA resides in cells other than CD4+ T cells, including myeloid
cells. AIDS 28:439-42.
68.   Ganor Y, Zhou Z, Bodo J, Tudor D, Leibowitch J, Mathez D, Schmitt A, Vacher-Lavenu MC,
Revol M, Bomsel M. 2013. The adult penile urethra is a novel entry site for HIV-1 that
preferentially targets resident urethral macrophages. Mucosal Immunol 6:776-86.
69.   Schmitt MP, Gendrault JL, Schweitzer C, Steffan AM, Beyer C, Royer C, Jaeck D, Pasquali
JL, Kirn A, Aubertin AM. 1990. Permissivity of primary cultures of human Kupffer cells for
HIV-1. AIDS Res Hum Retroviruses 6:987-91.
70.   Hufert FT, Schmitz J, Schreiber M, Schmitz H, Racz P, von Laer DD. 1993. Human Kupffer
cells infected with HIV-1 in vivo. J Acquir Immune Defic Syndr (1988) 6:772-7.
71.   Mosoian A, Zhang L, Hong F, Cunyat F, Rahman A, Bhalla R, Panchal A, Saiman Y, Fiel MI,
Florman S, Roayaie S, Schwartz M, Branch A, Stevenson M, Bansal MB. 2017. Frontline
Science: HIV infection of Kupffer cells results in an amplified proinflammatory response
to LPS. J Leukoc Biol 101:1083-1090.

72. Sierra-Madero JG, Toossi Z, Hom DL, Finegan CK, Hoenig E, Rich EA. 1994. Relationship
between load of virus in alveolar macrophages from human immunodeficiency virus
type 1-infected persons, production of cytokines, and clinical status. J Infect Dis 169:18-
27.
73. Jambo KC, Banda DH, Kankwatira AM, Sukumar N, Allain TJ, Heyderman RS, Russell DG,
Mwandumba HC. 2014. Small alveolar macrophages are infected preferentially by HIV
and exhibit impaired phagocytic function. Mucosal Immunol 7:1116-26.
74. Cribbs SK, Lennox J, Caliendo AM, Brown LA, Guidot DM. 2015. Healthy HIV-1-infected
individuals on highly active antiretroviral therapy harbor HIV-1 in their alveolar
macrophages. AIDS Res Hum Retroviruses 31:64-70.
75. Stoler MH, Eskin TA, Benn S, Angerer RC, Angerer LM. 1986. Human T-cell lymphotropic
virus type III infection of the central nervous system. A preliminary in situ analysis. JAMA
256:2360-4.
76. Wiley CA, Schrier RD, Nelson JA, Lampert PW, Oldstone MB. 1986. Cellular localization
of human immunodeficiency virus infection within the brains of acquired immune
deficiency syndrome patients. Proc Natl Acad Sci U S A 83:7089-93.
77. Fischer-Smith T, Croul S, Sverstiuk AE, Capini C, L'Heureux D, Regulier EG, Richardson
MW, Amini S, Morgello S, Khalili K, Rappaport J. 2001. CNS invasion by CD14+/CD16+
peripheral blood-derived monocytes in HIV dementia: perivascular accumulation and
reservoir of HIV infection. J Neurovirol 7:528-41.
78. Cosenza MA, Zhao ML, Si Q, Lee SC. 2002. Human brain parenchymal microglia express
CD14 and CD45 and are productively infected by HIV-1 in HIV-1 encephalitis. Brain
Pathol 12:442-55.
79. Ko A, Kang G, Hattler JB, Galadima HI, Zhang J, Li Q, Kim WK. 2019. Macrophages but not
Astrocytes Harbor HIV DNA in the Brains of HIV-1-Infected Aviremic Individuals on
Suppressive Antiretroviral Therapy. J Neuroimmune Pharmacol 14:110-119.
80. Micci L, Alvarez X, Iriele RI, Ortiz AM, Ryan ES, McGary CS, Deleage C, McAtee BB, He T,
Apetrei C, Easley K, Pahwa S, Collman RG, Derdeyn CA, Davenport MP, Estes JD, Silvestri
G, Lackner AA, Paiardini M. 2014. CD4 depletion in SIV-infected macaques results in
macrophage and microglia infection with rapid turnover of infected cells. PLoS Pathog
10:e1004467.
81. Igarashi T, Brown CR, Endo Y, Buckler-White A, Plishka R, Bischofberger N, Hirsch V,
Martin MA. 2001. Macrophage are the principal reservoir and sustain high virus loads in
rhesus macaques after the depletion of CD4+ T cells by a highly pathogenic simian
immunodeficiency virus/HIV type 1 chimera (SHIV): Implications for HIV-1 infections of
humans. Proc Natl Acad Sci U S A 98:658-63.
82. Abreu CM, Veenhuis RT, Avalos CR, Graham S, Parrilla DR, Ferreira EA, Queen SE, Shirk
EN, Bullock BT, Li M, Metcalf Pate KA, Beck SE, Mangus LM, Mankowski JL, Mac Gabhann
F, O'Connor SL, Gama L, Clements JE. 2019. Myeloid and CD4 T Cells Comprise the Latent
Reservoir in Antiretroviral Therapy-Suppressed SIVmac251-Infected Macaques. MBio 10.
83. Abreu CM, Veenhuis RT, Avalos CR, Graham S, Queen SE, Shirk EN, Bullock BT, Li M,
Metcalf Pate KA, Beck SE, Mangus LM, Mankowski JL, Clements JE, Gama L. 2019.
Infectious Virus Persists in CD4(+) T Cells and Macrophages in Antiretroviral Therapy-
Suppressed Simian Immunodeficiency Virus-Infected Macaques. J Virol 93.

84.   Andrade VM, Mavian C, Babic D, Cordeiro T, Sharkey M, Barrios L, Brander C, Martinez-
Picado J, Dalmau J, Llano A, Li JZ, Jacobson J, Lavine CL, Seaman MS, Salemi M,
Stevenson M. 2020. A minor population of macrophage-tropic HIV-1 variants is
identified in recrudescing viremia following analytic treatment interruption. Proc Natl
Acad Sci U S A 117:9981-9990.
85.   Graziano F, Aimola G, Forlani G, Turrini F, Accolla RS, Vicenzi E, Poli G. 2018. Reversible
Human Immunodeficiency Virus Type-1 Latency in Primary Human Monocyte-Derived
Macrophages Induced by Sustained M1 Polarization. Sci Rep 8:14249.
86.   Agosto LM, Hirnet JB, Michaels DH, Shaik-Dasthagirisaheb YB, Gibson FC, Viglianti G,
Henderson AJ. 2016. Porphyromonas gingivalis-mediated signaling through TLR4
mediates persistent HIV infection of primary macrophages. Virology 499:72-81.
87.   Castellano P, Prevedel L, Valdebenito S, Eugenin EA. 2019. HIV infection and latency
induce a unique metabolic signature in human macrophages. Sci Rep 9:3941.
88.   Josefsson L, von Stockenstrom S, Faria NR, Sinclair E, Bacchetti P, Killian M, Epling L, Tan
A, Ho T, Lemey P, Shao W, Hunt PW, Somsouk M, Wylie W, Douek DC, Loeb L, Custer J,
Hoh R, Poole L, Deeks SG, Hecht F, Palmer S. 2013. The HIV-1 reservoir in eight patients
on long-term suppressive antiretroviral therapy is stable with few genetic changes over
time. Proc Natl Acad Sci U S A 110:E4987-96.
89.   Moore AC, Bixler SL, Lewis MG, Verthelyi D, Mattapallil JJ. 2012. Mucosal and peripheral
Lin- HLA-DR+ CD11c/123- CD13+ CD14- mononuclear cells are preferentially infected
during acute simian immunodeficiency virus infection. J Virol 86:1069-78.
90.   Yukl SA, Shergill AK, Ho T, Killian M, Girling V, Epling L, Li P, Wong LK, Crouch P, Deeks
SG, Havlir DV, McQuaid K, Sinclair E, Wong JK. 2013. The distribution of HIV DNA and
RNA in cell subsets differs in gut and blood of HIV-positive patients on ART: implications
for viral persistence. J Infect Dis 208:1212-20.
91.   Schulz O, Jaensson E, Persson EK, Liu X, Worbs T, Agace WW, Pabst O. 2009. Intestinal
CD103+, but not CX3CR1+, antigen sampling cells migrate in lymph and serve classical
dendritic cell functions. J Exp Med 206:3101-14.
92.   Brenchley JM, Price DA, Schacker TW, Asher TE, Silvestri G, Rao S, Kazzaz Z, Bornstein E,
Lambotte O, Altmann D, Blazar BR, Rodriguez B, Teixeira-Johnson L, Landay A, Martin JN,
Hecht FM, Picker LJ, Lederman MM, Deeks SG, Douek DC. 2006. Microbial translocation
is a cause of systemic immune activation in chronic HIV infection. Nat Med 12:1365-71.
1093

Viglianti Figure 1A-D

Viglianti Figure 2

Viglianti Figure 3

Viglianti Figure 4 (A-C)

Vigilanti Figure 4 (D-H)

Viglianti Figure 5

A

TLR Engagement

B

Early Phase

C

Late Phase

Viglianti Supplemental Figure 2

Viglianti Supplemental Figure 3