Circadian clock mechanism driving mammalian photoperiodism

Wood S.H.[1,2,8], Hindle, M.M.[3,8], Mizoro, Y.[1], Cheng, Y. [4,5], Saer B.R.C.[1],

Miedzinska K.[3], Christian H.C.[6], Begley, N. [1], McNeilly J.[7], McNeilly A.S.[7],

Meddle, S.L. [3], Burt, D.W.[3,4], Loudon A.S.I.[1*]

[1] Centre for Biological Timing, Faculty of Life Sciences, University of

Manchester, Manchester M13 9PT, UK

[2] Arctic Chronobiology and Physiology research group, Department of Arctic

and Marine Biology, UiT – The Arctic University of Norway, Tromsø NO-9037,

Norway

[3] The Roslin Institute, and Royal (Dick) School of Veterinary Studies

University of Edinburgh, Roslin, Midlothian EH25 9PRG, UK

[4] UQ Genomics Initiative, The University of Queensland, Brisbane, QLD 4072,

Australia

[5] School of Life and Environmental Sciences, Faculty of Science, The

University of Sydney, New South Wales, Australia

[6] University of Oxford, Department of Physiology, Anatomy and Genetics, Le

Gros Clark Building, South Parks Road, Oxford, OX1 3QX, UK

[7] MRC Centre for Reproductive Health, Queen's Medical Research Institute,

Edinburgh EH16 4TJ, UK

[8] These authors contributed equally

*Corresponding author: andrew.loudon@manchester.ac.uk

Abstract

The annual photoperiod cycle provides the critical environmental cue synchronizing rhythms of life in seasonal habitats. In 1936, Bünning proposed a circadian-based coincidence timer for photoperiodic synchronization in plants. Formal studies support the universality of this so-called coincidence timer, but we lack understanding of the mechanisms involved. Here we show in mammals that long photoperiods induce the circadian transcription factor *BMAL2*, in the *pars tuberalis* of the pituitary, and triggers summer biology through the eyes absent / thyrotrophin (EYA3 / TSH) pathway. Conversely, long-duration melatonin signals on short photoperiods induce circadian repressors including *DEC1*, suppressing BMAL2 and the EYA3/TSH pathway, triggering winter biology. These actions are associated with progressive genome-wide changes in chromatin state, elaborating the effect of the circadian coincidence timer. Hence, circadian clock-pituitary epigenetic pathway interactions form the basis of the mammalian coincidence timer mechanism. Our results constitute a blueprint for circadian-based seasonal timekeeping in vertebrates.

INTRODUCTION

The annual photoperiod cycle provides a critical predictive environmental cue driving annual cycles of fertility, physiology and behaviour in most animal species. In 1936, Erwin Bünning proposed the external coincidence hypothesis for a circadian-based timing mechanism driving photoperiodic responses in plants [1]. The proposition was that photoperiod entrains a circadian rhythm of photosensitivity, and the expression of summer or winter biology depends on whether or not light coincides with the phase of high photosensitivity. An internal coincidence model has also been proposed where the role of light is to entrain two circadian oscillators and the phase relationship between the two oscillators determines the response to photoperiod [2,3]. In either case the role of the circadian clock is central and formal studies support the universality of a coincidence timer in animals [2–7], but we lack understanding of the mechanisms involved.

In mammals, the duration of the night-time pineal melatonin signal is sculpted by the photoperiod and is a critical regulator of annual cycles of reproduction, growth and metabolism [8–14]. The melatonin signal is decoded within the *pars tuberalis* (PT) of the pituitary gland [9,13,15–18]. Day-length dependent changes in hypothalamic thyroid hormone (TH) metabolism control the seasonal changes in physiology and are regulated via the PT, through altered secretion of thryrotrophin (TSH) [19–21]. This TSH circuit is specific to the PT, and distinct from that in the anterior pituitary, which controls normal thyroid gland function. LP-activation of PT-TSH depends on regulation of TSHβ subunit expression by the transcriptional co-activator, EYA3, which operates as a TSH on-switch. EYA3 co-activates the PAR bZIP Transcription Factor TEF (thyrotroph embryonic factor) via a D-box element on the *TSHβ* promoter [22–24] (Fig. 1a). This prior work places EYA3 at the centre of photoperiodic time measurement within the melatonin target tissue, but does not explain how expression is regulated by the seasonal clockwork.

In addition to acute circadian-based mechanisms an epigenetic basis to longer-term seasonal control in flowering in plants is well established. Specifically, the duration of cold temperatures during the winter season alters

chromatin accessibility at key genes, and is a requirement for full photoperiod
induced flowering (vernalisation) [25–27]. In animals, no evidence for dynamic
genome-wide seasonal epigenetic regulation of transcription has been
described.

Here, we show that coincidence timing depends on a flip-flop switch between
the expression of the circadian genes *BMAL2*, (a paralogue of the circadian
regulator *BMAL1*) and *DEC1,* and their respective activating or suppressive
effects on *EYA3* in the PT*.* The duration of the melatonin signal is key in
sculpting these molecular components. Additionally, the effect of the
coincidence timer is elaborated by progressive genome-wide changes in
epigenetic status at key seasonal gene promoters. Therefore, circadian clock
interactions with pituitary epigenetic pathways form the basis of the
mammalian coincidence timer mechanism.

RESULTS

Epigenetic regulation of the seasonal transcriptome

Seasonally breeding sheep are a well-established photoperiodic model for the study of neuroendocrine mechanisms underpinning seasonal physiology [14,28,29]. Using a study design that compared the effects of transfer from long photoperiod (LP) to short photoperiod (SP) with transfer from SP to LP (Fig. 1b), we collected *pars tuberalis* (PT) tissue at 1,7 and 28 days after transfer, with collections timed for 4-h after lights on (ZT4), when *EYA3* expression peaks under LP [22,24,29]. Changes in prolactin concentrations [29–31] confirmed photoperiodic hormone responses (Fig. 1b, n=28), as well as LP activation of *EYA3,* and inverse expression patterns of *TSHβ* (LP marker) and *CHGA* (SP marker) (Supplementary Fig. 1a-d, n=4), validating this paradigm.

Comparing LP day 28 to SP day 28 by electron microscopy we found an increased nuclear diameter, equating to an approximate doubling in volume of PT thyrotrophs, and a marked reduction in chromatin density on LP (Fig. 1c, Supplementary Fig. 1e & f). These morphological changes were not seen in *Pars distalis* (PD) somatotrophs but were observed in the PT follicular stellate (FS) cells to a lesser degree (Fig. 1c, Supplementary Fig. 1e & f). We hypothesized that increased chromatin accessibility may account for the photoinductive effects of LP and therefore changes in the seasonal transcriptome.

Comparing all seasonal time-points (Fig. 1b, red arrows) we performed ChIP-seq (histone marker H3K4me3, Supplementary Data 1 & 2, n=2) and RNA-seq (Supplementary Data 3, n=3) to screen for seasonal transcriptional activation. To determine if epigenetic changes in H3K4me3 marks were associated with transcriptional activation we first improved the sheep genome annotation of transcripts and transcription start sites (TSS) with PT RNA and a combination of Cap Analysis of Gene Expression (CAGE-seq, Supplementary Data 4) [32] and ISOSEQ long-read RNA-seq (SRA: PRJNA391103, see methods for details). Using this improved annotation we identified H3K4me3 peaks and found the distribution on genomic features to be similar to previous studies [33–35], validating our approach (see methods section).

Next we identified seasonally expressed genes, as defined by RNA-seq
analysis of differentially regulated genes (DEGs) in the SP to LP and LP to SP
transfers (Supplementary Data 3), and observed a strong correlation between
seasonal gene expression and H3K4me3 peaks around the transcription start
sites (TSS's)(Fig. 1d). Importantly, this correlation was absent in non-
seasonally regulated genes and we found the distributions between seasonal
and non-seasonally regulated genes to be significantly different (Fig. 1d, P
value < 0.001 Mann-Whitney U test). Histone modifications are precisely
balanced by methyltransferases (writers), demethylases (erasers) and effector
proteins (readers), therefore we checked the RNA expression of H3K4me3
readers, writers and erasers but found no seasonal changes (Supplementary
Data 5). This suggests that changes in protein activity of H3K4me3
modulators may be key in the observed seasonal alterations of H3K4me3
marks.

We noted that approximately 70% of seasonally DEGs (Supplementary Data
3) had more than one TSS compared to only ~20% of the PT genomic
background (Supplementary Fig. 1g, Supplementary Data 4). Next we took
genes that were up-regulated in either SP or LP and plotted the proportion of
genes with multiple TSS's, and repeated this for H3K4me3 marked TSS's, this
revealed that H3K4me3 marks are more likely to occur on genes with multiple
TSS's and highly expressed seasonal DEGs have a greater prevalence of
multiple TSSs than non-seasonal genes expressed at the same level (Fig. 1e).
Furthermore, this phenomenon was more pronounced in LP than SP. This
indicates that LP up-regulated genes are more likely to have multiple TSS's
and be associated with the H3K4me3 mark.

To investigate which promoter motifs may be actively transcribed we searched
for enriched transcription factor binding site motifs in the TSS's marked with
H3K4me3 in the LP day 28 vs SP day 28 comparison (Fig. 1f, Supplementary
Fig. 2a & b). This showed enrichment of D-box binding motifs, Basic Leucine
Zipper ATF-Like transcription related factors and POU domains for in both the
up and down-regulated genes (Fig. 1f).

TSHβ, an exemplar seasonal D-box regulated gene [22,23], revealed progressive

activation by LP, correlated with expression, and with no detectable H3K4me3

mark at either SP day 28 or 84 (R=0.961, p-value=0.002, Fig. 2a, b). In

contrast, CHGA revealed an inverse seasonal pattern (R=0.843, p-

value=0.009, Supplementary Fig. 2c, d). Global enrichment of D-box sites on

LP-activated genes indicates a potentially extensive role for TEF, SIX1 and

EYA3 co-activation in seasonal regulation of physiology, and strongly focuses

attention on the regulation of EYA3.

In line with our global analysis showing the presence of multiple TSSs in

seasonally expressed genes we identified two transcription start sites in *EYA3*,

an up-stream TSS with non-canonical E-boxes and a downstream TSS

containing two paired canonical E-Boxes[22] (Fig. 2c). CAGE analysis revealed

that only the down-stream TSS was actively transcribed on LP (Fig. 2c,

Supplementary Data 4). We correlated RNA expression with H3K4me3 marks

for both TSS's, which confirmed seasonal regulation specific to the down-

stream TSS (EYA3 downstream TSS: R=0.778, P=0.02, Fig. 2d, EYA3

upstream TSS: R=0.308, P=0.46, Supplementary Fig. 2e & f). Next, we cloned

each EYA3 TSS into luciferase reporters, and using COS7 cells transfected

the reporters along with known E-box regulators [22,23] (see methods for details).

This revealed significant activation specific to the downstream (seasonal) TSS

(Supplementary Fig. 2g), likely due to the presence of multiple canonical e-

box pairs.

BMAL2 an activator in a circadian-based coincidence timer

The E-box regulators *CLOCK* and *BMAL1* do not show significant changes in

amplitude in the PT under different photoperiods [22,36], therefore we aimed to

identify a candidate circadian E-box regulator of the EYA3 downstream TSS.

The prediction is that a circadian E-box regulator, would peak in expression

only when light falls on the photosensitive phase, as in the early light phase of

LP (approx. ZT4), and would be absent on SP where the photosensitive

phase is masked by darkness. The RNA-seq dataset revealed progressive up-

regulation by LP of multiple transcripts, and a slower inverse pattern on SP
(Fig. 3a, Supplementary Data 3).

Importantly these data identified *BMAL2*, an E-box regulator and paralogue of
BMAL1, as progressively up-regulated on LP (Fig. 3a & b), closely matching
the expression profile for *EYA3* (Fig. 3b). We tested whether BMAL2 was
capable of activating the downstream *EYA3* TSS (Supplementary Fig. 2h). In
the presence of CLOCK, or alone, BMAL2 had a weak non-significant effect,
but in the presence of BMAL1 and CLOCK, we observed significant 4 to 5-fold
activation (Supplementary Fig. 2h, Fig. 3c). We then tested whether this
depended on direct E-box binding by BMAL2, by mutating the BMAL2 DNA-
binding domain (arginine of basic helix-loop-helix domain to alanine R88A [37]),
and observed near-identical augmentation of *EYA3* expression (Fig. 3d). The
observation that transcription factors can act as co-activators [38] and that
bHLH circadian transcription factors require internal PAS domains for
functional protein-protein interactions [39], with PAS-B an essential domain for
BMAL1 protein-protein interactions [40] led us to mutate the PAS-B domain on
ovine BMAL2 (F427R_V439R) [40]. This significantly impaired *EYA3* activation,
blocking the co-activation effect with BMAL1 and CLOCK (Fig. 3e). This
suggests that BMAL2 operates as a co-activator of *EYA3*, in a
CLOCK/BMAL1-dependent manner, requiring PAS-B dependent protein-
protein interactions for the mechanism of action, confirmation of this will
require suitable antibodies to be raised for future studies.
We assessed the daily profile of *BMAL2* by in-situ hybridization, using
archived material [24]. This revealed strong LP-dependent induction of *BMAL2*
in the early light phase (ZT3, Fig. 3f), and flat low 24h expression on SP.
Collectively this reveals that BMAL2 operates as a co-activator driving
photoperiodic responses, and fulfils the criteria as a candidate activating arm
of the circadian-based coincidence timer.

Circadian clock repressors shut-down the LP transcriptome

A prediction from a coincidence timer model is that repressive mechanisms
would emerge on short photoperiods in the late night (i.e. photo-inducible

phase masked by darkness). To assess this we conducted a serial RNA-seq
experiment on PT tissue collected over at 4-hour intervals from both SP and
LP-housed animals (Day 28 Fig. 1b & Fig. 4a). Transcripts at the two ZT0
collections 24h apart (ZT0 and ZT24) in each photoperiod had virtually
identical RNA-seq profiles, validating dissection technique and down-stream
analyses (Supplementary Fig. 3a & Supplementary Data 6). We identified
transcripts that significantly changed across the day (diurnal genes) within
each photoperiod, and defined the phase, amplitude and period [41]
(Supplementary Data 6). More transcripts were rhythmic on SP compared to
LP (SP = 880, LP = 643 (Fig. 4b)), with striking asymmetry in the peak phase
of expression on both SP and LP (Fig. 4c). Large numbers of genes were
expressed on SP in late night (ZT20, 24) and on LP in the early light phase
(ZT4; Fig. 4c), and surprisingly relatively few genes common to both
photoperiods (250 transcripts, Fig. 4b & d), but shared genes included the
canonical clock genes (Supplementary Fig. 3b).

Amongst diurnal genes shared across photoperiods, dawn-peaking clusters
were enriched for the GO term cAMP binding activity, while the dark-onset
cluster was marked by the circadian suppressor *CRY1* [42–44] (Fig. 4d & e,
Supplementary Fig. 3c). Previous studies have shown *CRY1* is directly
induced by onset of melatonin secretion [17,45–47] and that the pattern of
induction does not alter between photoperiods. We also observed that
regulators of CRY1 protein stability, *FBXL3* and *FBXL21* [22,48,49], are
expressed in the PT, but here there were marked photoperiod-dependent
changes in phasing for *FBXL21* (Fig. 4e) .

BMAL2 emerged as the only transcription factor, which showed both phase
and amplitude changes in expression on LP (Fig. 4e, Supplementary Data 6).
On LP there was a 4 log-fold increase in expression of *BMAL2*, with bi-phasic
peaks at ZT4 and ZT20. In contrast, *BMAL1* showed no alteration in amplitude
of expression, on either SP or LP (Fig. 4e).

At SP ZT20 we noted a strong functional enrichment for the terms negative
regulation of transcription, E-box binding and co-repressor activity
(Supplementary Fig. 3c & d). Within this group, we identified 8 genes with

known circadian repressor function (*REVERB-α, DEC1, DEC2, CHRONO,*
FBXL21, KLF10, JUNB, GATA2 and *ERF,* Fig. 4e & Supplementary Fig. 3e).
Notably 6 are circadian clock components acting through E-boxes. Cistromic
analysis identified over-representation of circadian RORE, D-Box and E-Box
sites on the proximal promoter regions in late-night genes on SP (ZT20) and
on LP at the dark-light transition (Fig. 4f). Cluster analysis revealed 4 major
clusters of E-box motif-enriched genes associated with late night and dawn
transition on SP (ZT20) and LP (ZT4) respectively, with *BMAL2* contained
within the LP ZT4 cluster (Fig. 4g), and circadian repressor elements within
the SP ZT20 peaking cluster (Fig. 4g). This led us to ask; are SP ZT20
repressors more likely to interact, and therefore potentially repress, LP ZT4
up-regulated genes. We used curated experimental protein-protein interaction
(PPI) observations from the STRING database, which contain known protein-
protein interactions and functional associations [50]. We found that LP ZT4 up-
regulated genes are more enriched within the SP ZT20 repressor network (P-
value = 0.001) than down-regulated genes (Supplementary Fig 4a, b). We did
not find significant enrichment when considering genes that are differentially
up or down regulated across the whole day (P-value = 0.25) (Supplementary
Fig. 4b). This suggests that SP is defined by an up-regulation of
transcriptional repressors responsible for the suppression of the LP activated
transcriptome.

Melatonin-regulated circadian repression

Earlier studies have defined the role for melatonin both as an acute inducer of
CRY1 [17,36,45,47], but also determined clear melatonin duration-dependent
effects for photoperiodic responses [8,51]. Our data suggest that prolonged SP-
like signals may trigger late-night induction of a separate cohort of circadian
repressors genes. To test this, we maintained a cohort of animals on LP and
then transferred animals to constant light (LL), a regimen known to suppress
the endogenous rise in pineal melatonin secretion [52]. Simultaneously, and at
the time of normal lights off (i.e. expected onset of the melatonin signal) we
treated animal with an intradermal Regulin melatonin implant [47], which mimics
the endogenous dark-onset rise of this hormone [53], validated by RIA for
melatonin (Supplementary Fig. 5a, n=6). PT tissue was collected for in-situ

hybridization analysis at +1.5 hours (ZT17.5), +3.5 hours (ZT19.5), +6.5 hours
(ZT22.5) and +9.5h after hormone treatment, the latter time point being
equivalent to 1.5 hours into the predicted light-onset phase on LP (i.e. ZT1.5;
Fig. 5a, n=4).

CRY1 expression in the PT was completely suppressed by constant light, but
rapidly induced by melatonin treatment, as previously observed [17,45–47] (Fig.
5b). By ZT1.5, expression levels had dropped to basal. In LL conditions *DEC1,*
REVERBα and *CHRONO* were rapidly elevated in the early subjective dark
phase, and in an abnormal pattern when compared to endogenous profiles in
LP conditions (Fig. 4e vs Fig 5c-e). In marked contrast, exposure to melatonin
initially repressed all 3 genes, which peaked at +9.5h, equivalent to ZT1.5 on
LP (Fig.s 5c - e). For *DEC1* and *CHRONO*, this pattern closely resembled the
phase of the endogenous rise of these genes in animals housed in SP
photoperiods (Fig. 4e & 5f). From this, we conclude that multiple repressor
arms of the circadian clock are direct melatonin targets and act as a read-out
of long duration melatonin signals. This defines a molecular basis for a
hormone-regulated circadian-based coincidence timer mechanism, capable of
discriminating LP and SP responses within the melatonin target tissue.

DEC1 suppression of BMAL2-mediated EYA3 activation

DEC proteins are known E-box suppressors of the circadian clock [54,55], so we
selected DEC1 for further analysis. Quantitative in-situ hybridization confirmed
the temporal pattern of *DEC1* expression (Fig. 5f) in the PT, and using
immunohistochemistry of short-photoperiod derived PT tissue, we showed co-
localisation of DEC1 protein to αGSU expressing thyrotroph cells in the PT
(Fig. 5g). DEC1 appears to be specific to thyrotrophs, and was not detected in
a common cell-type in the PT, the folliculo-stellate cells. We cloned ovine
DEC1, and then tested action on BMAL2-mediated induction of *EYA3.* This
showed that DEC1 significantly suppressed the action of both CLOCK
/BMAL1 (Supplementary Fig. 5b & c) and CLOCK/BMAL1/BMAL2-mediated
expression of *EYA3* (Fig. 5h, Supplementary Fig. 5c). We also tested DEC2,
which was without effect (Supplementary Fig. 5d). Hence, DEC1 and BMAL2
exert mutually antagonistic effects on EYA3 expression.

Discussion

Our study reveals a circadian coincidence timer within the PT, sculpted by the
nocturnal melatonin signal that encodes the mammalian photoperiodic
response (Fig. 6). Under LP, *BMAL2* exhibits a high-amplitude peak timed
approximately 12h after the onset of the preceding dark phase, coincident with
expression of *EYA3* in the early light phase [22]. This 12h interval from dark
onset to the dawn peak is also remarkably close to the critical photoperiod
required to activate a long-day response in sheep [28,56]. We note that both
BMAL2 and EYA3 show a second peak on day 28 of LP, however, these
peaks are "dysynchonised" (ZT20 BMAL2 and ZT15/16 EYA3 [24]).
Furthermore, the second peak in EYA3 expression is absent in early
responses to LP [22] indicating that it is potentially not mechanisiticaly
important in a coincidence model. We show that BMAL2 acts as a co-activator
of *EYA3* with CLOCK and BMAL1, likely in a complex through the PAS-B
domain interaction [57]. Co-activators are recognized as important rapid-
response functional integrators of multiple transcription factors, driving distinct
biological programmes and environmental responses, including adaption to
cold, rapid diet change and disease [38]. Within the PT, BMAL2 and EYA3 may
therefore operate as a photoperiodic co-activator cascade It remains unclear
how *BMAL2* is regulated, and current models for the regulation of BMAL2 by
BMAL1 may not be applicable [58]. Our promoter motif analysis does indicate
the presence of E-boxes in BMAL2 but this presents a circular argument and
further work on the regulation and evolution of BMAL2 function is required.

While we cannot differentiate between an internal or external coincidence
model of photoperiodic time measurement in this study, generally the concept
of a photoperiodic coincidence timer predicts that on short photoperiods,
repressor activity would dominate in the late night. In line with this we show
that long-duration melatonin signals elicit repressor gene transcription in the
late night, timed approximately 12h after dark onset/rise of melatonin (ZT20
SP). Impressively, these repressors are directly implicated in the negative
regulation of LP ZT4 induced genes (Supplementary Fig. 4). Amongst these,
we show the circadian repressor DEC1 [54] blocks induction, by BMAL2, of
EYA3. Therefore, DEC1 and BMAL2 act as a circadian flip-flop switch for

photoperiodic time measurement (Fig. 6). It remains unclear whether DEC-
mediated repression is via a direct action on E-box sites occupied by BMAL2
or indirect, leading to modification of a co-activator complex. The discovery of
transcriptional repressors in the latter half of the night on SP contrasts with
previously described acute and photoperiod-independent induction of *CRY1*
by melatonin onset [17,36,45,47].   However, constant light and therefore CRY1
repression seems to induce the late-night repressor genes, suggesting that
these are direct targets of CRY1-mediated repression in the early to mid
nocturnal phase. This observation along with altered phasing by photoperiod
of the CRY1 protein stability regulator FBXL21 [48,49] indicate that the dynamics
of protein degradation of CRY1 could play a role in discriminating melatonin
signal duration.

Our study reveals a progressive increase in H3K4me3 marks at transcription
start sites of seasonally expressed genes, especially in long photoperiods (Fig.
1d). Furthermore, these long photoperiod induced genes are more likely to
have multiple transcription start sites marked with H3K4me3 (Fig. 1e,
Supplementary Fig. 1g). *EYA3* is a particular example of this showing
seasonal regulation of only one out of two transcription start sites, presumably
seasonally modulating the transcription of EYA3 via BMAL2. On transfer to
long photoperiods the *EYA3* and *TSHβ* locus show a progressive increase
over a number of weeks in H3K4me3 marks matching the increase in
expression (Fig. 2c, d). Our earlier work shows that at the individual cell level
the transition between winter and summer physiology is a binary, all-or-
nothing phenomenon [29,59]. Integrating these two findings, we suggest that
individual thyrotroph cells of the PT exhibit a distribution of critical day length
requirements/sensitivity for circadian triggering of the summer physiology
leading to a binary switch in cell phenotype, which in a whole tissue assay
would appear as a progressive change in epigenetic status.

We therefore envisage a photoperiodic time measurement model in which a
melatonin-regulated circadian-based interval timer interacts with the
underlying cellular chromatin state, rhythmically recapitulating the final
developmental stages of a thyrotroph endocrine cell in response to long days,
leading to an adaptive summer-like physiological state. Our data do indicate

morphological changes in another PT cell type, the FS cell, however these
cells lack melatonin receptors[60,61] therefore a role in a coincidence timer is
unclear. Defining how a flip-flop circadian timer (Fig. 6) interacts with the
chromatin state at the level of the single PT cell remains a future challenge.

In summary, the use of photoperiod to synchronise life history cycles in a
variable environment is an ancestral feature observed in a large majority of
species. Even amongst long day and short day breeders the EYA3-TSH
circuitry behaves similarly, demonstrating that although the downstream
reproductive responses to photoperiod are altered the mechanism of
photoperiodic time measurement is shared [9,13]. The EYA3-TSH circuitry is
conserved amongst vertebrates [9,13,16,21], as is the use of a
circadian/coincidence based system for photoperiodic time measurement [2,4–7].
Here, we have defined how photoperiodic changes in EYA3/TSH expression
stem from a molecular circadian coincidence timer mechanism. We expect
this model to be widely applicable across the vertebrate linage.

 Methods

 Animals & experimental design

All animal experiments were undertaken in accordance with the Home Office
Animals (Scientific Procedures) Act (1986), UK, under a Project License held
by A.S.I.L. Scottish blackface castrate males were housed in artificial light
dark cycles, either 8:16 h light ∕ dark cycle for short photoperiod (SP) or 16:8 h
light ∕ dark cycle for long photoperiod (LP).

Two separate photoperiod controlled studies were undertaken; 1. The
experiment presented in Fig. 1b (seasonal comparison) & Fig. 4a (diurnal
comparison at day 28), and, 2. The experiment presented in Fig. 5a
(melatonin implant study). Animals were blood sampled throughout the study
and terminally sampled at the indicated time-points (Figure 1b, Figure 4a,
Figure 5a). The seasonal experiment was designed to take into account the
effects of a photoperiodic switch from SP (SP day 84) to LP and the
progressive seasonal changes (LP day 1, 7, 28 and 112), followed by return to
SP (SP day 1, 7, 28). Animals were terminally sampled at ZT4 at all time-
points. The diurnal comparison was conducted on day 28 animals from this
study, they were sampled across the day at 4 hourly intervals for 24 hours.
The melatonin implant study was a separate experiment on pre-conditioned
LP animals (8 weeks).

All animals were killed by an overdose of barbiturate (Euthatal; Rhone
Merieux, Essex, UK) administered intravenously. Hypothalamic blocks with
the *pars tuberalis* (PT) and pituitary attached were collected for
immunohistochemistry (n=3 per group), electron microscopy (n=3 per group),
transcriptomics (n=3 per group), insitu-hybridization (n=4 per group) and
epigenomics (n=2 per group). From the omic analyses the PT was dissected
to minimise inclusion of transition zone and median eminence. The PT
samples were snap frozen on dry ice and stored at -80C.

Hormone assays

Ovine prolactin (oPRL) was measured as in our previous study [29] for 28
animals during the seasonal experiment (Fig. 1b). In brief, a competitive

ELISA using purified oPRL (ovine prolactin NIDDK-oPRL-21; AFP10692C;
from Dr. A Parlow, NHPP, Harbor-UCLA Torrance CA, USA) and a highly
specific rabbit anti-ovine prolactin (ASM-R50, produced by ASM) were used.
Plates were read at 450nm. The Coefficient of Variation for the assay on
control plasma samples was <10%.

Ovine melatonin was measured by radioimmunoassay as previously
described [47] for animals in the melatonin implant study (Fig. 5a, n=6 per
timepoint). In brief, using a rabbit antimelatonin antiserum (PF1288; P.A.R.I.S.,
Paris, France) and 2-Iodomelatonin (NEX236050UC; PerkinElmer, Boston,
Massachusetts) as tracer RIA was performed. All samples were assayed in a
single assay with an intra-assay coefficient of variation of 5% and a sensitivity
of 5 pg/mL.

Immunohistochemistry

Tissues (n=3 per group) were immersed in Bouin's fixative for 8 hours,
transferred to 70% ethanol, then dehydrated and embedded in paraffin wax.
Sagittal sheep brain sections were cut from paraffin embedded tissue at 5 μM,
floated onto Superfrost Plus slides (J1800 AMNZ, Thermo scientific), dried at
50°C overnight, then dewaxed and rehydrated. Triple immunoflourscence for
TSHb, CHGA and EYA3, and the double immunoflourscence for DEC1 /αGSU
was performed as in our previous study [29].

DEC1 primary antibody (CW27, gift from Prof. Adrian L Harris, Weatherall
Institute of Molecular Medicine, John Radcliffe Hospital, Oxford) [62] was used
at 1:2000. 1:1000 diluted horse radish peroxidase conjugated chicken anti-
rabbit IgG antibody was used as a secondary antibody (PI-1000, Vector
Laboratories). TSA Plus Cyanine 5 (NEL763001KT, Perkin Elmer) was used
to visualize. aGSU immunofluorescence was used at 1:2000 diluted, ASM-
HRSU, R20) and treated with TSA Plus Fluorescein (NEL741001KT, Perkin
Elmer). Nuclei were stained by Hoechst 33258 (ab228550, abcam) and cover
glasses were mounted by VectaMount AQ (H-5501, Vector Laboratories).
Images were collected on a Zeiss Axioimager.D2 upright microscope using a
40x / 0.7 Plan neofluar objective and captured using a Coolsnap HQ2 camera

(Photometrics) through Micromanager Software v1.4.23. Images were then
processed using Fiji ImageJ (http://imagej.net/Fiji/Downloads).

In Situ Hybridization (ISH) and Quantification of Signal

The OaTSHβ plasmid (XM_004002368.2) was kindly provided by David
Hazlerigg. The OaEya3 plasmid (NM_001161733.1) was cloned as previously
described [22]. The OaCHGA, OaDEC1, OaREVERB-alpha, CHRONO and
OaBMAL2 were cloned as 1,948-1,970 of XM_004017959.4, 407-767 of
NM_001129741.1, 1,012-1,411 of NM_001131029.1, 83-532 of
XM_027974329.1, and 1,518-1,906 of XM_027965976.1, respectively.

Frozen coronal ovine hypothalamic blocks (n=4 per group) for in-situ
hybridization were cut into 16µm sections using a cryostat (CM3050s Leica
Microsystems, Ltd., Milton Keynes, UK), and thaw mounted onto poly-l-lysine
coated slides (VWR International, Lutterworth, UK). Radiolabelled cRNA
riboprobes were prepared by plasmid linearization and transcribed using P33
α-UTP (Perkin-Elmer). Fixed sections were hybridized overnight at 60°C with
5 x 105 cpm of probe per slide. Hybridization signals were visualised on
autoradiographic film (Kodak Biomax MR Films, Kodak, USA) after one week
exposure at -80°C. Signal intensity was quantified by densitometry analysis of
autoradiographs using the image-Pro Plus 6.0 software (Media Cybernetics,
Inc., Marlow, UK).

Tissue processing and electron microscopy (EM)

Hypothalamo-pituitary tissue blocks were fixed by immersion in 3%
paraformaldehyde/0.05% paraformaldehyde in 0.1M phosphate buffer (pH
7.2) for 24 hours at room temperature and transferred to a 1:10 dilution of the
fixative in 0.1M phosphate buffer for storage at 4□C before processing (n=3
per group). Using a scalpel blade, areas from the medial PT and median
eminence were cut into 0.5mm3 pieces which were then stained with osmium
(1% in 0.1M phosphate buffer), uranyl acetate (2% w/v in distilled water),
dehydrated through increasing concentration of ethanol (70 to 100%),
followed by 100% acetone and embedded in Spurr's resin (TAAB laboratory
equipment, Aldermarston, UK). Ultrathin sections (50-80 nm) were prepared

using a Reichart-Jung Ultracut ultramicrotome and mounted on nickel grids
(Agar Scientific Ltd., Stanstead, UK). Sections were then counterstained with
lead citrate and uranyl acetate and examined on a JOEL 1010 transmission
electron microscope (JOEL USA Inc., Peabody, MA, USA). Sections from 3
animals per group were examined.

For analysis of PT cell morphology, twenty micrographs per animal (n=3
sheep per group) of individual PT cells were taken at a magnification of x
5,000. Negatives were scanned into Adobe Photoshop CS2 (Adobe Corp.,
San Jose, CA, USA) and analysed using Axiovison version 4.5 (Zeiss,
Oberkochen, Germany) image analysis software. The analyst was blind to the
sample code. For measurement of the cell and nuclear areas, margins were
drawn around the cell or nucleus respectively and the area was calculated.
All morphometric values represent the mean ± SEM (n=3 sheep per group).
Means were compared by one way analysis of variance (ANOVA) with post
hoc analysis by the Bonferroni test. $P<0.05$ was considered statistically
different.

RNA-seq

RNA was extracted from the pars tuberalis from the seasonal experiment
(including the diurnal samples)(Fig1b & Fig. 4a) using Qiagen's TissueLyser II
and RNeasy tissue kit (n=3 per group). The quality of the extracted RNA was
assessed using the Agilent 2100 Bioanalyser; all RNA integrity numbers
(RINs) were above 8, indicating that good quality RNA had been extracted.
Poly-A selection was used.

RNA was prepared with TruSeq Stranded mRNA Sample Preparation Guide,
(15031047 Rev. E, Oct 2013) and sequenced on HiSeq 2500 with 125 base
pair paired-end reads by Edinburgh Genomics.

The FASTQ files were trimmed with TrimGalore v0.4.0 and mapped (TopHat[63]
v2.1.0 and Bowtie[64] v2.3.5) to the 5[th] release of the sheep genome
(Oar_rambouillet_v1.0; assembly GCA_002742125.1). StringTie[65] was used
to combine RNASeq and IsoSeq full-length transcripts to generate the
genome transcript annotation (accessible in GEO, GSE144677). On average

90% (sdev: 8.46) of paired reads generated were mapped to the genome and

73% (stdev: 0.03) of these were assignable to genes using featureCount [66]

(Subread v1.6.3).

All sequence data have been submitted to SRA under the BioProject

accession PRJNA391103 and processed data to GEO under GSE144677.

A limma-voom [67] analysis pipeline was used to determine the statistical

significance of differential expressed genes. Voom was used to generate

normalized precision weighted counts per million (CPM) values which were

used in the following regression analyses.

Seasonal comparison

The effect of switching from SP to LP was assessed by comparing SP day 84

to LP day 1, 7 and 28, and the effect of switching from LP to SP was

assessed by comparing LP day 112 to SP day 1, 7 and 28 (Fig. 1b,

Supplementary Data 3). For each gene, we fit a least squared regression

model with limma that calculates a single f-test for significance across all

model coefficients (mitigating type I errors). Time (days) was treated as a

categorical independent variable model for all ZT4 observations in LP and SP

(photoperiod x day) in limma and which allowed us to extract from the model

the fold change and significance for each pairwise contrasts of interest in

limma (Supplementary Data 3; Figure 3a,b). Significance was determined by

an FDR < 0.05, >0 log2CPM and a >1 log2 absolute fold change.

Diurnal comparison

To test for diurnal changes, samples collected at day 28 in LP and SP at a 4hr

time resolution were used (n=3 per group, 7 time-points, Fig. 4a). We used a

polynomial regression model approach similar to that of maSigPro[68].Least

squared regression models were then fitted with orthogonal polynomials up to

the 5^{th} order for time in each photoperiod to identify significantly changing

genes. To test for rapid single time-point changes in gene expression a

categorical regression model was also fitted to this dataset. Diurnal genes that

were significantly changing across time were identified as FDR significance

<0.05, log2CPM > 0 and absolute log2 fold change > 1 (Supplementary Data

6; Figure 4b,c,d). FDR was calculated throughout using the Benjamini &
Hochberg method. Gene expression changes between photoperiods were
evaluated by fitting a photoperiod x time (orthogonal polynomials up to 5th
order) model and extracting the significance and effect size from photoperiod
coefficient of the linear model (Supplementary Data 6). In selecting the
polynomial we used Akaike information criterion (AIC) to investigate the
optimal model selection for expressed genes, balancing model overfitting and
underfitting (using the Oshlack and Gordon selectModel implementation in
limma). It is not possible to select a single model that is optimal across all
genes, however for genes > 0 log2CPM and with an amplitude > 1.5 we found
that including orthogonal polynomials up to 5th order was optimal for the most
genes in both SP and LP time-series. Again we used the thresholds FDR
significance <0.05, log2CPM > 0 and absolute log2 fold change > 1
(Supplementary Data 6). MetaCycle[69] v1.1.0 was used to evaluate gene
expression in the 24hr time series for periodicity. JTKCyle [41] and Lomb-
Scargle statistics were calculated for an assumed period of 24 hours
(Supplementary Data 6). Rayleigh tests for uniformity were performed with the
CircStats 0.2-6 package in R. We tested the uniformity of distribution of peak
expression times for genes containing each of the core clock motifs (canonical
EBOXs (CACGTG), DBOXs (TTA[CT][GA]TAA)   and RORE sites
(AANTAGGTCA)) within the H3K4me3 marked region proximal (within 500bp)
of the TSS.

For all gene cohorts described enrichment analysis of GO terms and
pathways was conducted using both consensusPathDB [70] and TopGO [71] 3.1.0.
Signicant terms extracted by a FDR < 0.05 from a fishers exact test. The
weight01 algorithm was used for GO term weighting in TopGO. Where gene
annotation did not yet exist for novel and unannotated transcripts OrthoMCL [72]
v5 was used to predict orthology from protein sequences and annotation,
which were then used to transfer annotation from cow, human, mouse and rat
genes. Protein sequences were predicted from novel transcripts using
TransDecoder v5.3.0 guided by Uniprot [73] and Pfam [74] best hits to rank coding
frames.

We clustered SP and LP time-series profiles using Partitioning Around
Medoids (PAM) (Fig. 4g) with the cluster 2.1.0 package in R. Day 28, 24 hour,
time-series were mean normalized and scaled and PAM clustered with
Euclidian distance. The Davies Bouldin index was used to evaluate the
optimum number of clusters (k=15 SP and k=9 LP). Motif enrichment of genes
clusters was evaluated using fishers two-way exact test against all PT
expressing genes as the background. Motifs we identified within H3K4me3
marked regions within 500bp of a candidate TSS assigned to a gene.

ChIP-seq

The method used for ChIP-seq was adapted from an ultra low cell number
native ChIP method [75]. In brief, nuclei were isolated from whole PT tissue
(n=2 per group) with a dounce homogenizer and sigma nuclear isolation
buffer. Mnase digestion was optimized to to give the best discrimination
between mono, di and tri histones. Immunoprecipitation was performed with
protein A and G beads (Invitrogen). Importantly the beads were pre-incubated
with the H3K4me3 antibody (active motif) to reduce background noise.
Stringent washing with salt buffers was also used to reduce the background
noise. The bound chromatin was elute and extracted using a phenol
chloroform method. Ampure bead purification was used to clean up the
samples. The qubit and tapestation were used to quantify, a minimum amount
of 5ng was required for the library preparation. The Manchester genomic
facility prepared the ChIP-seq libraries and sequenced them according to the
standard illumina protocol.

Massive parallel sequencing were performed by Illumina HiSeq4000 with 75
bp paired-end and converted fastqs by bcl2fastq (ver 2.17.1.14). Fastqs were
trimmed by trimmomatic (ver 0.36) [76] with the following parameters,
ILLUMINACLIP: TruSeq3-PE-2.fa:2:30:10 SLIDINGWINDOW:4:20
MINLEN:35, aligned to the sheep genome (Oar_rambouillet_v1.0) by BWA
MEM (ver 0.7.17) [77] with the default parameter and then converted to BAM
and sorted by samtools (ver 0.1.19). Peak calling were carried out by MACS2
(ver 2.1.0) [78] with following parameters, --format BAMPE –gsize 2.81e9 --
keep-dup 1 --broad --broad-cutoff 0.1 --bdg --SPMR --qvalue 0.05.

Read coverages of peaks were calculated by SICER (ver 1.1) [79] with following

parameters; window size=200, gap length=200, fdr=0.01. BED files of

replicate samples were merged in order to perform SICER analysis which

does not allow replicates. Peaks called by SICER were annotated by HOMER

(ver 4.10.3) [80] with default parameters. H3K4me3 peaks identified by SICER

were validated by monitoring the distributions on the sheep genome. By

HOMER annotation, each peak was described as promoter-TSS (1000 bp

from TSSs), exon, intron, TTS, intergenic and the distributions of H3K4me3

peaks were closely resembling to the previous reports [33,34] (Supplementary

Fig. 6a). Furthermore, H3K4me3 peaks were well-associated with CpG

islands (CGIs) on the sheep genome as described in the previous study

(Supplementary Fig. 6b) [35]. We use a standard definition of CpG islands [81];

nucleotides regions with > 50% GC content, extending to > 200 bp and with

an observed vs expected CpG ratio >6.5, and detected them using

CgiHunterLight 1.0 on Oar_rambouillet_v1.0 (assembly GCA_002742125.1).

H3K4me3 peaks of each sampling day were shuffled by bedtools shuffle (ver

2.27.1) [82] with -noOverlapping as negative controls. For correlation analysis

with RNA expression, ChIP read counts of peaks overlapped in ± 200 bp from

TSSs were used. A workflow diagram can be found in Supplementary Fig. 7.

ISO-seq

For gene annotation, five tissue samples were sequenced over two

experimental runs using PacBio Iso-Seq. In the first run PT and PD samples

were sequenced from an RNA pool of SP and LP Scottish blackface sheep

(N=1) and a pineal from a commercial mule sheep from Manchester, UK. This

RNA was sent to GATC Biotech (Konstanz, Germany) for cDNA library

preparation using their in-house method with mRNA 5' cap and poly(A) tail

selections and sequencing on a PacBio RSII system. GATC made full length

normalized RNA libraries.size selected for <2kb, 2kb-4kb, >4kb. sequenced

across 75 PacBio RS II SMRT cells [SRX7688275]. In a second run, PT from

a pool of sheep in LP, and SP (n=3) were sequenced. RNA was extracted

using RNeasy Mini Kit (Qiagen) with on-column Dnase digestion. A full-length

cDNA library was constructed for each sample using the TeloPrime Full-

Length cDNA Amplification Kit V1 (Lexogen) and amplified using PrimeSTAR

GXL DNA Polymerase (Takara Bio) with 22 PCR cycles of 98 °C denaturation
for 10 seconds, 60 °C annealing for 15 seconds, and 68 °C extension for 10
minutes. PacBio SMRTbell libraries were prepared using SMRTbell Template
Prep Kit 1.0 and each library was sequenced on two SMRT Cells v2 LR using
20-hour movies on a Sequel platform at the IMB Sequencing Facility
(University of Queensland, [SRX7688271]). All Iso-Seq data was first
processed using software IsoSeq v3.1 to obtain full-length non-concatemer
reads with at least 3 full sequencing passes, which were then mapped to the
sheep reference genome GCA_002742125.1 using GMAP version 2018-05-
30. TAMA Collapse from the TAMA tool kit [83] was used to generate unique
gene and transcript models, which were further merged with RNAseq-based
annotation data using TAMA Merge to incorporate any transcript models that
were identified by RNAseq but not Iso-Seq. Functional annotation of
transcripts was carried out using Trinotate (v3.1.1).

Where multiple transcripts were present for an expressed gene, with more
than one transcription start sites (TSS) candidate, the active proximal
promoter regions were inferred by selecting contiguous H3K4me3 marked
regions within 100bp of a TSS.

CAGE-seq

We applied cap analysis gene expression (CAGE) to identify the location and
relative expression of TSS regions of the PT across both LP and SP. When
combined with IsoSeq and RNASeq derived transcript annotation this
provided a comprehensive identification of TSS in the genome which allowed
us to more accurately apply DNA binding motif analysis to promoter regions.
Libraries were prepared according to Hazuki *et. al* [32] and sequenced on an
Illumina HiSeq 2500 using V4 chemistry on a 50 cycle Single end sequencing
run. We sequenced archived RNA samples from the PT in both SP and LP
(ZT4, week 12) [29]. We also sequenced RNA from PD (both SP and LP), and
Pineal for comparison as outgroups. Reads were trimmed using fastx toolkit
0.0.14 and cutadapt 1.4. Reads were mapped using BWA 0.7.17 to the 5[th]
release of the sheep genome (Oar_rambouillet_v1.0; assembly
GCA_002742125.1). CAGEr 1.26.0 was used for processing and cluster

analysis of TSS (Supplementary Data 4). We filtered reads for a mapping
quality > 30 and sequencing quality > 20. Tag counts were normalised using
the power law method with an alpha of 1.12 and T of 10^6 (deterimined by
plotting the reverse cumulatives of PT samples). We clustering TSS with >1
TPM together using the distclu methods allowing a max distance between
TSS of 20 nucleotides.

Transcription factor binding site analysis

Transcription Factor binding motifs were identified using the FIMO tool from
MEME v4.11.4 with a p-value threshold of $<1\times10^{-7}$. for Jasper 2018 core
vertebrate database. Fishers one-way exact tests for enrichment were used to
identify the significance of motif enrichment within active promoter regions of a
gene cohort compared to a background of all PT expressed genes (> 0 CPM).
Fishers two-way exact tests were applied to evaluate enrichment and
depletion of motifs within active proximal promoter regions of genes with the
use of SP and LP H3K4me3 marked regions within 500bp of candidate TSSs.

Protein-protein interaction networks

Using experimentaly evidenced protein-protein interaction (PPI) annotation
from the STRING [50] v10 database for cow, sheep, rat, mouse and human we
integrated protein interactions in Cytoscape [84] with significantly changing
genes in the ZT 4 SP vs ZT20 LP from the 24 hr categorical contrast, which is
12 hrs from dark/melatonin onset (Supplementary Data 6 & Supplementary
Figure 4). A threshold of >0.4 for confidence score in experimental evidence
and a combined score of > 0.7 was applied, and orphaned proteins removed
from the network. The significance of enrichment of PPI repressor connected
genes within up-regulated vs down-regulated genes was evaluated using
fishers two-way exact tests.

Cloning and constructs

Expression plasmids: PCR fragments of the expected sizes were extracted
using a gel extraction kit (Qiagen) and cloned in pGEM-T easy vector

(Promega); Four to six positive clones were sequenced (MWG, United Kingdom). To generate expression constructs, a second round of PCR was performed using primers flanked by adequate restriction sites and the pGEM-T clone as template. PCR fragments were extracted as described above, digested by the adequate restriction enzymes, purified with a PCR purification kit (Qiagen) and cloned in the expression vector backbone (pCS2-HIS). In order to generate the mutant expression plasmids for BMAL2 we used the QuikChange Lightning Multi Site-Directed Mutagenesis Kit (210515, Agilent). The bHLH mutant was generated by converting an arginine to alanine (OaBMAL2_R88A) based on a mouse mutagenesis study on BMAL1[37]. The PAS-B mutant was made by converting a phenylalanine to arginine, and a valine to arginine (OaBMAL2_F427R_V439R). Based on a on a mouse mutagenesis study on BMAL1 [40].

Sanger sequencing of clones are available in Genbank for BMAL2 cds constructs (Genbank: [MT001920]), DEC1 cds constructs (Genbank: [MT019539]), DEC2 cds constructs (Genbank: [MT019540]), PAS-B-mutated BMAL2 cds constructs (Genbank: [MT019541]), bHLH-mutated BMAL2 cds constructs (Genbank: [MT019542]),

Promoter reporter constructs: a strategy identical to that described above was applied and fragments were cloned into the pGL4 basic backbone (Promega) digested with the appropriate restriction enzymes. Sequencing was performed to check accuracy of all re-amplified cloned fragments. EYA3 generic and seasonal promoter construct sequences are available on genebank ([MT001921] and [MT001924] respectively).

Cell culture, transfection and luciferase reporter assays

The procedure was as previously reported [22]. In brief, COS-7 cells were grown in Dulbecco's modified eagle's medium supplemented with 10% fetal bovin serum, 1% penicillin/streptomycin at 5% CO_2 and 37□. Cells were plated in 24-well plates at a density of 4x104 cells per ml and incubated for 24 hours prior to transfection. Transfection was performed using Genejuice (Novagen) and the concentration was optmised to transfect the greatest number of cells without compromising cell survival, this was assessed using a

luciferase positive control pGL3 containing SV40 (Promega) and trypan blue
staining. We recorded a 90% cell survival and a high transfection efficiency.
The EYA3 promoter constructs were used at 50ng per well, as in a previous
study [22]. The expression plasmids were used at different doses based on a
previous study and optimization of the assay: TEF = 12.5ng, DEC1 = 25ng,
CLOCK, BMAL1, BMAL2 and mutant BMAL2 were all used at 50ng, unless
otherwise stated. The total transfected DNA amount was set to an equal
amount between all conditions by addition of the corresponding empty vector.
The luciferase assays were performed 48 hours after transfection using the
luciferase assay kit (Promega) and the Glomax luminometer (Promega).
Thetotal protein per well, assessed by Bradford assay was used to normalize
the values to total protein content (a proxy for cell number). All data (in
Relative Luminescence Units, RLU) represent fold induction once normalized
to total protein content and relative to an inert control transfection. Each
experiment contained 4 replicate wells and was repeated 4 times giving
similar results. An one-way ANOVA using Tukey's multiple comparisons test
was performed for each separate experiment conducted in Graphpad prism
7.05. Representitive plots (n=4) are shown.

Data availability statement

All sequence data supporting the findings of this study have been deposited in
the Sequence Read Archive (SRA) under the BioProject accession
[PRJNA391103] and processed data to GEO under [GSE144677]. Chip-seq
raw fastq is available in SRA [SRP110487] and processed bigWig data for in
GEO [GSE144515]. Additionally, we have made the bigWig data for
rambouillet 1.0 genome [GCF_002742125.1] available as a track hub for use
with   NCBI   browser   at   https://data.cyverse.org/dav-
anon/iplant/home/mhindle/hub.txt that can be loaded in NCBI Genome Data
Viewer. CAGE raw data is available in SRA [SRP110487] and as BigWig
annotations in the public trackhub. Source data for Figure 1b, Figure 1c,
Figure 3c, Figure 3d, Figure 3e, Figure 3f, Figure 5c, Figure 5d, Figure 5e,
Figure 5f and Figure 5h are provided with the paper.

Code availability statement

Code used is available from the authors on request.

References

1. Bünning, E. Die endogene Tagesperiodik als Grundlage der
photoperiodischen Reaktion. *Ber Dtsch Bot Ges* **54**, 590–608 (1936).

2. Pittendrigh, C. S. & Minis, D. H. The entrainment of circadian
oscillations by light and their role as photoperiodic clocks. *Am. Nat.* **98**,
261–299 (1964).

3. Nanda, K. K. & Hamner, K. . Studies on the nature of the endogenous
rhythm affecting photoperiodic response of Biloxi soybean. *Bot. Gaz.*
120, 14–28 (1958).

4. Follett, B. K., Mattocks, P. W. & Farner, D. S. Circadian function in the
photoperiodic induction of gonadotropin secretion in the white-crowned
sparrow, Zonotrichia leucophrys gambelii. *Proc. Natl. Acad. Sci. U. S. A.*
71, 1666–9 (1974).

5. Goldman, B. D. Mammalian photoperiodic system: formal properties
and neuroendocrine mechanisms of photoperiodic time measurement. *J.*
Biol. Rhythms **16**, 283–301 (2001).

6. Pittendrigh, C. S., Elliott, J. & Takamura, T. The Circadian Component
in Photoperiodic Induction. in 26–47 (Wiley-Blackwell, 2008).
doi:10.1002/9780470720851.ch4

7. Ikegami, K. & Yoshimura, T. Circadian clocks and the measurement of
daylength in seasonal reproduction. *Mol. Cell. Endocrinol.* **349**, 76–81
(2012).

8. Bartness, T. J., Powers, J. B., Hastings, M. H., Bittman, E. L. &
Goldman, B. D. The timed infusion paradigm for melatonin delivery:
what has it taught us about the melatonin signal, its reception, and the
photoperiodic control of seasonal responses? *J. Pineal Res.* **15**, 161–90
(1993).

9. Wood, S. & Loudon, A. Clocks for all seasons: unwinding the roles and
mechanisms of circadian and interval timers in the hypothalamus and

799   pituitary. *J. Endocrinol.* **222**, R39-59 (2014).

800 10. Woodfill, C. J. I., Wayne, N. L., Moenter, S. M. & Karsch, F. J.
801   Photoperiodic Synchronization of a Circannual Reproductive Rhythm in
802   Sheep□: Identification of Season-Specific Time Cues ' Reproductive
803   Sciences Program and Department of Physiology , University of
804   Michigan. *Biol. Reprod.* **50**, 965–976 (1994).

805 11. Lincoln, G. A. & Ebling, F. J. P. Effect of constant-release implants of
806   melatonin on seasonal cycles in reproduction, prolactin secretion and
807   moulting in rams. *Reproduction* **73**, 241–253 (1985).

808 12. Bittman, E. L., Dempsey, R. J. & Karsch, F. J. Pineal melatonin
809   secretion drives the reproductive response to daylength in the ewe.
810   *Endocrinology* **113**, 2276–83 (1983).

811 13. Dardente, H., Wood, S., Ebling, F. & Sáenz de Miera, C. An integrative
812   view of mammalian seasonal neuroendocrinology. *J. Neuroendocrinol.*
813   **31**, e12729 (2019).

814 14. Lincoln, G. A. Melatonin Entrainment of Circannual Rhythms.
815   *Chronobiol. Int.* **23**, 301–306 (2006).

816 15. Wood, S. & Loudon, A. The pars tuberalis: The site of the circannual
817   clock in mammals? *Gen. Comp. Endocrinol.* **258**, 222–235 (2017).

818 16. West, A. C. & Wood, S. H. Seasonal physiology: making the future a
819   thing of the past. *Curr. Opin. Physiol.* **5**, 1–8 (2018).

820 17. Johnston, J. D. *et al.* Multiple Effects of Melatonin on Rhythmic Clock
821   Gene Expression in the Mammalian Pars Tuberalis. *Endocrinology* **147**,
822   959–965 (2006).

823 18. West, A. *et al.* Npas4 is activated by melatonin, and drives the clock
824   gene Cry1 in the ovine pars tuberalis. *Mol. Endocrinol.* **27**, 979–989
825   (2013).

826 19. Hanon, E. A. *et al.* Ancestral TSH mechanism signals summer in a
827   photoperiodic mammal. *Curr. Biol.* **18**, 1147–52 (2008).

   29

20. Nakao, N. *et al.* Thyrotrophin in the pars tuberalis triggers photoperiodic
response. *Nature* **452**, 317–22 (2008).

21. Nakane, Y. & Yoshimura, T. Universality and diversity in the signal
transduction pathway that regulates seasonal reproduction in
vertebrates. *Front. Neurosci.* **8**, 115 (2014).

22. Dardente, H. *et al.* A molecular switch for photoperiod responsiveness
in mammals. *Curr. Biol.* **20**, 2193–8 (2010).

23. Masumoto, K. *et al.* Acute Induction of Eya3 by Late-Night Light
Stimulation Triggers TSHβ Expression in Photoperiodism. *Curr. Biol.* **20**,
2199–2206 (2010).

24. Dupré, S. M. *et al.* Identification of Eya3 and TAC1 as long-day signals
in the sheep pituitary. *Curr. Biol.* **20**, 829–35 (2010).

25. Angel, A., Song, J., Dean, C. & Howard, M. A Polycomb-based switch
underlying quantitative epigenetic memory. *Nature* **476**, 105–8 (2011).

26. Song, J., Irwin, J. & Dean, C. Remembering the prolonged cold of winter.
Curr. Biol. **23**, R807-11 (2013).

27. Satake, A. & Iwasa, Y. A stochastic model of chromatin modification:
Cell population coding of winter memory in plants. *J. Theor. Biol.* **302**,
6–17 (2012).

28. Dardente, H., Hazlerigg, D. G. & Ebling, F. J. P. Thyroid hormone and
seasonal rhythmicity. *Front. Endocrinol. (Lausanne).* **5**, 19 (2014).

29. Wood, S. H. *et al.* Binary switching of calender cells in the pituitary
defines the phase of the circannual cycle in mammals. *Curr. Biol.* **25**,
(2015).

30. Karsch, F. J., Robinson, J. E., Woodfill, C. J. & Brown, M. B. Circannual
cycles of luteinizing hormone and prolactin secretion in ewes during
prolonged exposure to a fixed photoperiod: evidence for an endogenous
reproductive rhythm. *Biol. Reprod.* **41**, 1034–46 (1989).

31.   Lincoln, G. A., Clarke, I. J., Hut, R. A. & Hazlerigg, D. G. Characterizing
a mammalian circannual pacemaker. *Science* **314**, 1941–4 (2006).

32.   Takahashi, H., Kato, S., Murata, M. & Carninci, P. CAGE (Cap analysis
of gene expression): A protocol for the detection of promoter and
transcriptional networks. *Methods Mol. Biol.* **786**, 181–200 (2012).

33.   Hussey, S. G., Loots, M. T., van der Merwe, K., Mizrachi, E. & Myburg,
A. A. Integrated analysis and transcript abundance modelling of
H3K4me3 and H3K27me3 in developing secondary xylem. *Sci. Rep.* **7**,
3370 (2017).

34.   Ershov, N. I. *et al.* Consequences of early life stress on genomic
landscape of H3K4me3 in prefrontal cortex of adult mice. *BMC*
Genomics **19**, 93 (2018).

35.   Rose, N. R. & Klose, R. J. Understanding the relationship between DNA
methylation and histone lysine methylation. *Biochim. Biophys. Acta -*
Gene Regul. Mech. **1839**, 1362–1372 (2014).

36.   Lincoln, G., Messager, S., Andersson, H. & Hazlerigg, D. Temporal
expression of seven clock genes in the suprachiasmatic nucleus and
the pars tuberalis of the sheep: evidence for an internal coincidence
timer. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 13890–5 (2002).

37.   Hosoda, H., Motohashi, J., Kato, H., Masushige, S. & Kida, S. A BMAL1
mutant with arginine 91 substituted with alanine acts as a dominant
negative inhibitor. *Gene* **338**, 235–41 (2004).

38.   Spiegelman, B. M. & Heinrich, R. Biological Control through Regulated
Transcriptional Coactivators. *Cell* **119**, 157–167 (2004).

39.   Guo, Y., Scheuermann, T. H., Partch, C. L., Tomchick, D. R. & Gardner,
K. H. Coiled-coil Coactivators Play a Structural Role Mediating
Interactions in Hypoxia-inducible Factor Heterodimerization. *J. Biol.*
Chem. **290**, 7707–7721 (2015).

40.   Huang, N. *et al.* Crystal Structure of the Heterodimeric CLOCK:BMAL1

Transcriptional Activator Complex. *Science (80-.).* **337**, 189–194 (2012).

41.   Hughes, M. E., Hogenesch, J. B. & Kornacker, K. JTK-CYCLE: An
efficient nonparametric algorithm for detecting rhythmic components in
genome-scale data sets. *J. Biol. Rhythms* **25**, 372–380 (2010).

42.   Xu, H. *et al.* Cryptochrome 1 regulates the circadian clock through
dynamic interactions with the BMAL1 C terminus. *Nat. Struct. Mol. Biol.*
22, 476–484 (2015).

43.   Maywood, E. S. *et al.* Translational switching of Cry1 protein expression
confers reversible control of circadian behavior in arrhythmic Cry-
deficient mice. *Proc. Natl. Acad. Sci.* **115**, E12388–E12397 (2018).

44.   Partch, C. L., Green, C. B. & Takahashi, J. S. Molecular architecture of
the mammalian circadian clock. *Trends Cell Biol.* **24**, 90–99 (2014).

45.   Dardente, H. *et al.* Melatonin induces Cry1 expression in the pars
tuberalis of the rat. *Mol. Brain Res.* **114**, 101–106 (2003).

46.   Dupré, S. M. *et al.* Identification of melatonin-regulated genes in the
ovine pituitary pars tuberalis, a target site for seasonal hormone control.
Endocrinology **149**, 5527–39 (2008).

47.   West, A. *et al.* Npas4 is activated by melatonin, and drives the clock
gene Cry1 in the ovine pars tuberalis. *Mol. Endocrinol.* (2013).
doi:10.1210/me.2012-1366

48.   Hirano, A. *et al.* FBXL21 Regulates Oscillation of the Circadian Clock
through Ubiquitination and Stabilization of Cryptochromes. *Cell* **152**,
1106–1118 (2013).

49.   Dardente, H., Mendoza, J., Fustin, J.-M., Challet, E. & Hazlerigg, D. G.
Implication of the F-Box Protein FBXL21 in Circadian Pacemaker
Function in Mammals. *PLoS One* **3**, e3530 (2008).

50.   Szklarczyk, D. *et al.* The STRING database in 2017: quality-controlled
protein–protein association networks, made broadly accessible. *Nucleic*
Acids Res. **45**, D362–D368 (2017).

51.   Maywood, E. S. *et al.* The effect of signal frequency on the gonadal
915   response of male Syrian hamsters to programmed melatonin infusions.
916   *J. Neuroendocrinol.* **4**, 37–44 (1992).

917   52.   Ebling, F. J., Lincoln, G. A., Wollnik, F. & Anderson, N. Effects of
918   constant darkness and constant light on circadian organization and
919   reproductive responses in the ram. *J. Biol. Rhythms* **3**, 365–84 (1988).

920   53.   Staples, L. D., McPhee, S., Kennaway, D. J. & Williams, A. H. The
921   influence of exogenous melatonin on the seasonal patterns of ovulation
922   and oestrus in sheep. *Anim. Reprod. Sci.* **30**, 185–223 (1992).

923   54.   Honma, S. *et al.* Dec1 and Dec2 are regulators of the mammalian
924   molecular clock. *Nature* **419**, 841–844 (2002).

925   55.   Dardente, H., Fustin, J.-M. & Hazlerigg, D. G. Transcriptional feedback
926   loops in the ovine circadian clock. *Comp. Biochem. Physiol. Part A Mol.*
927   *Integr. Physiol.* **153**, 391–398 (2009).

928   56.   Hazlerigg, D., Lomet, D., Lincoln, G. & Dardente, H. Neuroendocrine
929   correlates of the critical day length response in the Soay sheep. *J.*
930   *Neuroendocrinol.* e12631 (2018). doi:10.1111/jne.12631

931   57.   Gustafson, C. L. & Partch, C. L. Emerging Models for the Molecular
932   Basis of Mammalian Circadian Timing. *Biochemistry* **54**, 134–149
933   (2015).

934   58.   Bunger, M. K. *et al.* Mop3 Is an Essential Component of the Master
935   Circadian Pacemaker in Mammals. *Cell* **103**, 1009–1017 (2000).

936   59.   Wood, S. H. How can a binary switch within the pars tuberalis control
937   seasonal timing of reproduction? *J. Endocrinol.* **239**, R13–R25 (2018).

938   60.   Klosen, P. *et al.* The mt1 melatonin receptor and RORbeta receptor are
939   co-localized in specific TSH-immunoreactive cells in the pars tuberalis
940   of the rat pituitary. *J. Histochem. Cytochem.* **50**, 1647–57 (2002).

941   61.   Dardente, H., Klosen, P., Pévet, P. & Masson-Pévet, M. MT1 melatonin
942   receptor mRNA expressing cells in the pars tuberalis of the European

hamster: effect of photoperiod. *J. Neuroendocrinol.* **15**, 778–86 (2003).

62. Turley, H. *et al.* The hypoxia-regulated transcription factor DEC1(Stra13,
SHARP-2) and its expression in human tissues and tumours. *J. Pathol.*
203, 808–813 (2004).

63. Kim, D. *et al.* TopHat2: accurate alignment of transcriptomes in the
presence of insertions, deletions and gene fusions. *Genome Biol.* **14**,
R36 (2013).

64. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with
Bowtie 2. *Nat. Methods* **9**, 357–359 (2012).

65. Pertea, M., Kim, D., Pertea, G. M., Leek, J. T. & Salzberg, S. L.
Transcript-level expression analysis of RNA-seq experiments with
HISAT, StringTie and Ballgown. *Nat. Protoc.* **11**, 1650–1667 (2016).

66. Liao, Y., Smyth, G. K. & Shi, W. FeatureCounts: An efficient general
purpose program for assigning sequence reads to genomic features.
Bioinformatics **30**, 923–930 (2014).

67. Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: precision weights
unlock linear model analysis tools for RNA-seq read counts. *Genome*
Biol. **15**, R29 (2014).

68. Nueda, M. J., Tarazona, S. & Conesa, A. Next maSigPro: updating
maSigPro bioconductor package for RNA-seq time series.
Bioinformatics **30**, 2598 (2014).

69. Wu, G., Anafi, R. C., Hughes, M. E., Kornacker, K. & Hogenesch, J. B.
MetaCycle: An integrated R package to evaluate periodicity in large
scale data. *Bioinformatics* **32**, 3351–3353 (2016).

70. Kamburov, A., Stelzl, U., Lehrach, H. & Herwig, R. The
ConsensusPathDB interaction database: 2013 update. *Nucleic Acids*
Res. **41**, D793-800 (2013).

71. Alexa, A., Rahnenführer, J. & Lengauer, T. Improved scoring of
functional groups from gene expression data by decorrelating GO graph

structure. *Bioinformatics* **22**, 1600–1607 (2006).

72.   Li, L., Stoeckert, C. J. & Roos, D. S. OrthoMCL: Identification of ortholog
groups for eukaryotic genomes. *Genome Res.* **13**, 2178–2189 (2003).

73.   UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res.* **47**,
D506–D515 (2019).

74.   El-Gebali, S. *et al.* The Pfam protein families database in 2019. *Nucleic*
Acids Res. **47**, D427–D432 (2019).

75.   Brind'Amour, J. *et al.* An ultra-low-input native ChIP-seq protocol for
genome-wide profiling of rare cell populations. *Nat. Commun.* **6**, 6033
(2015).

76.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer
for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

77.   Li, H. & Durbin, R. Fast and accurate short read alignment with
Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

78.   Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome*
Biol. **9**, R137 (2008).

79.   Zang, C. *et al.* A clustering approach for identification of enriched
domains from histone modification ChIP-Seq data. *Bioinformatics* **25**,
1952–1958 (2009).

80.   Heinz, S. *et al.* Simple Combinations of Lineage-Determining
Transcription Factors Prime cis-Regulatory Elements Required for
Macrophage and B Cell Identities. *Mol. Cell* **38**, 576–589 (2010).

81.   Ioshikhes, I. P. & Zhang, M. Q. Large-scale human promoter mapping
using CpG islands. *Nat. Genet.* **26**, 61–63 (2000).

82.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for
comparing genomic features. *Bioinformatics* **26**, 841–842 (2010).

83.   Kuo, R. I., Cheng, Y., Smith, J., Archibald, A. L. & Burt, D. W.
Illuminating the dark side of the human transcriptome with TAMA Iso-

Seq analysis. *bioRxiv* 780015 (2019). doi:10.1101/780015

84.   Shannon, P. *et al.* Cytoscape: A software Environment for integrated

models of biomolecular interaction networks. *Genome Res.* **13**, 2498–

2504 (2003).

1004

Acknowledgements

The authors thank Joan Docherty and the staff at the Marshall Building, Roslin, Edinburgh for expert care of their research animals. Past members of the Manchester/Edinburgh team have made contributions to discussions and offered advice, including Sandrine Dupre, and Alex West. We thank David Hazlerigg, Carrie Partch and Jean Michel Fustin for their critical review of earlier drafts of this work. The work was supported by grants from the Biotechnology and Biological Sciences Research Council UK (BB/N015584/1, BB/N015347/1, BBP013759/1) and a Human Sciences Frontier Programme Grant RGP0030/2015 Evolution of Seasonal Timers awarded to ASIL and DWB. ASIL acknowledges the support of the Wellcome Trust, Grant 107851/Z/15/Z

Author contributions

SHW - designed the experiments, collected samples, performed RNA preparation, chromatin immunoprecipitation preparation, luciferase reporter assays, cloning and mutagenesis, analysed/interpreted data, prepared the manuscript and figures. **MH** – Bioinformatic analysis, data analysis and submission, sequencing, prolactin assay, figure preparation and revised the manuscript. **YM** - Bioinformatic analysis, immunohistochemistry, luciferase reporter assays, cloning and mutagenesis, figure preparation and revised manuscript. **YC** – ISO-seq and bioinformatic analysis **KM** – immunohistochemistry and analysis. **HC** – performed EM and analysis, revised manuscript. **BRCS** - collected samples and performed the in situ hybridization, luciferase reporter assays, cloning and mutagenesis. **NB** – collected samples and provided lab support. **JM** - developed the novel prolactin assay and prepared antibodies. **ASM** - collected samples, designed experiments, developed the prolactin assay, and revised the manuscript. **SM** - collected samples, designed experiments, prolactin assay, and revised manuscript. **DWB** - designed experiments, bioinformatics analysis, and revised the manuscript. **ASIL** – conceived the study, designed experiments, collected samples, analysed/interpreted data, and prepared the manuscript.

1037

Competing interests

1039 The authors have no competing interests to declare.

1040

Figure legends

Fig. 1: Photoperiod dependent epigenetic regulation of transcription in the
Pars tuberalis

a. Current model for the mammalian photoperiodic circuitry.

b. Prolactin concentrations during the experiment (black), n = 28
biologically independent animals per time-point. Error bars represent the
SEM. Grey line: number of hours of light the animals received in 24 hours.
Red arrows show terminal sampling points (ZT4), with the exception of day
28 where animals were collected across the day at 4 hour intervals.

c. EM images of PT thyrotrophs and PD somatotrophs at LP day 28 and
SP day 28. Dense chromatin in the nucleus is false coloured in purple.
Black scale bars = 2um.

d. Histogram revealing frequency distributions of Pearson correlation
coefficients between RNA expression ($\log_2$ CPM) and H3K4me3 peak read
counts ±200 bp from TSSs ($\log_2$ read counts). Red bars are seasonally
expressed genes ($\log_2$ fold change ≥ 1 or ≤ -1 and adjusted p value < 0.05
of short photoperiod (SP) day 84 vs long photoperiod (LP) day 1, 7, 28 and
LPday112 vs SPday1, 7, 28 differentially expressed genes (DEGs)=480)
and black bars are non-seasonally expressed genes ($\log_2$ fold change ≤
0.1 and ≥ -0.1 from the same comparisons, (genes=218)).

e. Prevalence of multiple (>1) TSS's in LP day28 up-regulated DEGs (solid
red), SP day 28 up-regulated DEGs (solid blue) and all PT expressed
genes as the background (solid black) as a percentage of the uppermost
expressed genes (i.e. increasing thresholds for the upper quantile of gene
expression). Log2CPM values for upper quantiles on upper x-axis. Dashed

lines indicate the proportion of gene in the cohorts with multiple H3K4me3
(>1) marked TSS.

f. Over-represented motifs LP day 28 up and down-regulated genes,
compared to SP day 28. The percentage of H3K3me3 marked TSS (+/-
100bp) containing a specific motif is given. The black triangles represents
the motif abundance in all the genes expressed (>0 CPM) in the PT and
the percentage of marked with H3k4me3 in LP (x-axis) or SP (y-axis). The
circles show the motif abundance and percentage H3K4me3 marked
TSS's in LP up-regulated (blue) and down-regulated (red) genes (FDR <
0.05; fishers two-way exact test).

Fig. 2: *TSH*β and EYA3 are epigenetically regulated over the season

a. ChIP-seq tracks for *TSH*β gene H3K4me3 peaks across all
experimental time-points. Chromsome 1 region is shown. CAGE-seq track
identifying active TSS's in LP day 28 vs SP day 28. Pink represents
samples in long photoperiod and marine green represents short
photoperiod. Solid green boxes are canonical E-box motifs. Blue boxes are
D-box motifs.

b. Correlation plot for *TSH*β log2 H3k4me3 peaks from ChIP-seq versus
TSHb log2 counts per million (CPM) from RNA-seq. Red symbols are LP
sampling points, green are SP sampling points. Correlation coefficient R is
shown. R=0.961, p-value=0.002. Note: SP day 28 and 84 are not included
because the H3k4me3 peaks are very low.

c. ChIP-seq tracks for *EYA3* gene H3K4me3 peaks. Chromsome 2 region
is shown. CAGE-seq track identifying active TSS's in LP day 28 vs SP day
28. Zoom in box provided to identify the downstream promoter.

d. Correlation plot for *EYA3* downstream TSS log2 H3K4me3 peaks from
ChIP-seq versus *EYA3* log2 counts per million (CPM) from RNA-seq. Red
symbols are LP sampling points, green are SP sampling points.
Correlation coefficient R is shown. R=0.778, p-value=0.02.

Fig. 3: Identification of BMAL2 as a seasonally regulated gene

a. Volcano plots showing the number of genes up (yellow) and
downregulated (blue) in pairwise comparisons. The design for the
comparisons is shown on the left. These plots demonstrate the
consistency of the BMAL2 seasonal signal. Numbers in grey boxes are the
differentially expressed genes in the pairwise comparisons
(Supplementary Data 3).

b. Z-score normalized log2CPM RNA-seq plots for EYA3 (blue) and
BMAL2 (orange) across the whole experiment. N= 3, individual animal
values are shown. Statistcal significances across the season are found in
Supplementary Data 3.

c. Transactivation of the EYA3-downstream-TSS-luc reporter by TEF,
CLOCK, BMAL1 and BMAL2. The experiment was repeated 4 times (n=4
per experiment), plot displayed is a representative result. A one-way
ANOVA was performed on each individual experiment using Tukey's
multiple comparisons test. Different letters indicate significant differences
between groups (P < 0.01). Error bars SD.

d. Transactivation of the EYA3-downstream-TSS-luc reporter by TEF,
CLOCK, BMAL1 and BMAL2. The lack of effect of mutating the DNA
binding domain of BMAL2 is shown. The experiment was repeated 4 times
(n=4 per experiment), plot displayed is a representative result. A one-way
ANOVA was performed on each individual experiment using Tukey's
multiple comparisons test. Different letters indicate significant differences
between groups (P < 0.01). Error bars SD.

e. Transactivation of the EYA3-downstream-TSS-luc reporter by TEF,
CLOCK, BMAL1 and BMAL2. The effect of mutating the PAS-B domain of
BMAL2 is shown. The experiment was repeated 4 times (n=4 per
experiment), plot displayed is a representative result. A one-way ANOVA
was performed on each individual experiment using Tukey's multiple
comparisons test. Different letters indicate significant differences between
groups (P < 0.01). Error bars SD.

f. In situ hybridization and quantification for BMAL2 mRNA from archived
material from collected every 4 hours from SP and LP. Representative
images are shown (n = 4, three independent observations per animal).
Error bars represent the SD. Statistical analysis by one-way ANOVA
performed, different letters indicate significant differences between groups
(P < 0.01).

Fig. 4: Short photoperiods are defined by the expression of transcriptional
repressors

a. Experimental design: SP day 28 and LP day 28 the pars tuberalis was
collected at 4 hour intervals for 24 hours. N= 3 per time-point.

b. Venn diagram showing the number of diurnal genes (significantly
changing throughout the day, FDR <0.05) in the PT in LP and SP
(Supplementary Data 4).

c. Rose plot showing the significant diurnal genes peaking at a particular
phase in LP and SP. Grey bars indicate genes shared between
photoperiods and yellow indicates they are unique to a photoperiod.
Around the outside of the plot the durations of light and dark the animals
received is plotted in grey and black respectively, the ZT times are given.

d. Heatmap showing the expression profile of the genes diurnal in both SP
and LP (250 genes) for each individual (n=3 per timepoint). Data are
ordered by SP peak phase. Yellow is up-regulated and blue down-
regulated, the data are scaled. Light dark bars are ZT times are given at
the bottom of the heatmap.

e. RNA-seq log2 CPM plots, n= 3, individual data shown. Grey shading
indicates lights off. Statistical significances are in Supplementary Data 6.

f. Transcription factor binding site analysis by peak phase in SP and LP.
Grey and white shading show transition from dark and light. The arrows
represent enrichment of either RORE (blue), E-boxes (pink), or D-boxes
(red). The direction of arrows indicate the mean expression peak of genes

containing the motif in their promoter and the length of the arrow is the -

log10 p-value from a Rayleigh tests of uniformity.

g. Cluster analysis of the RNA-seq data identified ZT20 SP peaking and

ZT4 LP peaking gene clusters that were enriched for E-boxes. Plots shown

are mean normalised log2CPM of expression profiles to visualise cluster

trends. The number of genes in each cluster is indicated in the bottom left.

The broken red line represents the most represenive medoid gene in the

cluster.

Fig. 5: Melatonin duration defines the expression of repressor genes

a. Study design. Animals were sampled at 1.5 hour intervals with either a

sham implant or melatonin implant. LL= constant light.

b. In situ hybridization and quantification for *CRY1* mRNA. Red bars are

the value in the presence of melatonin, black bars are in constant light with

a sham implant (n = 4). Representative images are shown. Statistical

significances from a one-way ANOVA and are between the melatonin and

sham implant at each time-point. Error bars represent the SD. P-value; * =

<0.01, **=<0.001, ***=<0.0001, ****=<0.00001

c. In situ hybridization and quantification for *REVERb-alpha* mRNA. Red

bars are the value in the presence of melatonin, black bars are in constant

light with a sham implant (n = 4). Representative images are shown.

Statistical analysis by one-way ANOVA performed, different letters indicate

significant differences between groups (P < 0.01). Error bars represent the

SD.

d. As in c for *CHRONO* mRNA.

e. As in c for *DEC1* mRNA.

f. In situ hybridization and quantification for *DEC1* mRNA from archived

material from collected every 4 hours from SP and LP (n = 4).

Representative images are shown. Error bars represent the SD. Statistical

analysis by one-way ANOVA performed, different letters indicate

significant differences between groups (P < 0.01).

g. Double immunohistochemistry for DEC1 (red), aGSU (green) and dapi
nuclear stain (blue). White arrow heads indicate FS cells. Scale bar 20um.

h. Transactivation of the EYA3-downstream-TSS-luc reporter by TEF,
CLOCK, BMAL1 and BMAL2 is repressed by DEC1. The experiment was
repeated 4 times (n=4 per experiment), plot displayed is a representative
result. A one-way ANOVA was performed on each individual experiment
using Tukey's multiple comparisons test. Different letters indicate
significant differences between groups ($P < 0.01$). Error bars SD.

Fig. 6: Mammalian photoperiodism model: coincidence timing and epigenetic
regulation

The photoperiod sculpts the duration of the melatonin signal, long duration on
short photoperiods (SP, winter) and short duration on long photoperiods (LP,
summer). In the pars tuberalis approximately 12 hours after the onset of
darkness there is a photosensitive phase. When this phase coincides with
light BMAL2 is expressed. When this phase is coincident with darkness DEC1
is expressed. This forms a photoperiodic flip-flop switch between two stable
states. The role of BMAL2, CLOCK and BMAL1 is to co-activate EYA3 and
subsequently seasonal physiological changes. DEC1 suppresses the E-box
based activation of EYA3. Also occurring in the dark phase of SP a number of
repressors are expressed which target appear to target LP induced genes
potentially inhibiting the expression of summer physiology. Aside from this
acute photoperiodic flip-flop switch there is a progressive photoperiod
dependent epigenetic regulation over a number of weeks augmenting the
coincidence based timer drive on summer physiology.