1   Individual differences in spatial learning are correlated across cognitive tasks but not associated with

2   stress response behaviour in the Trinidadian guppy

4   Pamela M. Prentice[1], Chloe Mnatzaganian[1], Thomas M. Houslay[1], Alex Thornton[1], Alastair J. Wilson[1]

[1]Centre for Ecology and Conservation, University of Exeter, Penryn Campus, Cornwall, TR10 9FE, UK

Acknowledgements

We would like to thank technical staff in the Penryn Aquatic facility for help with technical and

husbandry support. This work was supported by the Biotechnology and Biological Sciences Research

Council (grant BB/L022656/1) and by a BBSRC studentship to PMP. We are also grateful to Neeltje

Boogert and John Quinn for constructive comments on an earlier version of this MS.

Abstract

Cognition is vital for carrying out behaviours required for survival and reproduction. In animals, we

now know that cognitive performance varies not just among species, but also among individuals within

populations. While this variation is a prerequisite for contemporary natural selection, it is also true

that selection does not act on traits in isolation. The extent to which cognitive traits covary with other

aspects of phenotype (e.g. personality traits) is therefore expected to be an important factor in

shaping evolutionary dynamics. Here we adopt a multivariate approach to test for spatial learning

ability in a captive population of male Trinidadian guppies (*Poecilia reticulata*), and ask whether

differences in cognitive performance are associated with (repeatable) differences in stress response

behaviour. We focus on stress response for two reasons. First, functional links between cognitive traits

and 'stress coping style' have been hypothesised. Second, individual-level studies of cognitive

performance typically rely on multiple testing paradigms that may themselves be a stressor. Thus,

there is a risk that variation in stress responsiveness is itself a cause of apparent, but artefactual

variance in cognitive ability. Using a set of fish exposed repeatedly to two distinct spatial learning tasks

(maze layouts), and an acute stress response test (open field trial), we find differences among-

individuals in task performance that are repeatable within- and across maze layouts. On average

performance improves with experience in the first maze, consistent with spatial learning, but not in

the second. In both mazes, there is among-individual variation in the trajectory of mean performance

with trial number suggesting individuals differ in 'learning rate'. Acute stress response behaviour is

repeatable but predicts neither average time to solve the maze nor learning rate. We thus find no

support for among-individual correlation between acute stress response and cognitive performance.

However, we highlight the possibility that cumulative, chronic stress effects may nonetheless cause

observed declines in performance across repeats for some individuals (leading to lack of improvement

in mean time to solve the second maze). If so, this may represent a pervasive but difficult challenge

for our ability to robustly estimate learning rates in studies of animal cognition.

Introduction

Cognition is defined as the set of mechanisms by which animals acquire, process, store and use information from the environment (Healy & Rowe, 2010; Shettleworth, 2010), and is vital for carrying out day-to-day behaviours needed for survival and reproduction. While differences in cognitive performance among-species have long been studied in comparative psychology (for a review see Healy 2019), a more recent focus in behavioural ecology has been the characterisation of among-individual variation within populations of non-human animals (Lucon-Xiccato & Bisazza, 2017a; Ashton et al., 2018; Boogert et al., 2018). This among-individual variation is interesting from an evolutionary perspective, as it is a pre-requisite for natural selection and genetic variation – both of which are fundamental for adaptive evolution to occur (Wilson et al., 2010). Though selection does not act on traits in isolation, so quantifying relationships between cognitive variation and other aspects of phenotype is important. For example, functional links between cognitive performance and other aspects of behaviour (including, for example neophobia, boldness and stress responsiveness) have been widely hypothesised (Quinn et al., 2012; Sweis et al., 2013; Griffin et al., 2015; Medina-García et al., 2017). However, robustly testing these relationships is often challenging, requiring multivariate data collection and analyses to detect and describe patterns of variation between associated traits at the appropriate level (e.g., among-individual and/or among genotype; Dingemanse & Dochtermann, 2013). Nonetheless, such efforts are important if we hope to understand the adaptive evolution of cognition in the context of the wider phenotype (Thornton & Wilson, 2015). Here we address this broad goal in the more specific context of testing hypothesised links between cognitive performance and a behavioural stress-response (Øverli et al., 2007; Gibelli et al., 2019) in Trinidadian guppies (*Poecilia reticulata*).

Quantifying patterns of among-individual variation in cognitive traits is still in its infancy (Rowe & Healy, 2014; Thornton et al., 2014; Boogert et al., 2018), and empirical studies therefore remain somewhat limited (but see Ashton et al. 2018; Tyrone Lucon-Xiccato & Bisazza 2017; Niemelä et al.

2013 for examples). However it is now abundantly clear that populations typically harbour high levels

of among-individual variation in behavioural traits more generally (Dingemanse & Réale, 2005).

Individual differences in (mean) behaviours, commonly referred to as personality, can manifest as, for

instance, variation in aggressiveness or sociability towards conspecifics, or differences in response

when faced with predators or other sources of perceived risk (Bridger et al. 2015; Réale et al. 2007).

Since strong directional or stabilising selection is usually predicted to erode variation (Roff, 2002), it is

hypothesised that variation in personality traits is maintained by fitness trade-offs of some kind

(Dingemanse et al., 2004; Quinn et al., 2016). For example, bolder individuals may be better at

acquiring resources to invest in life history traits (e.g., growth, reproduction) but their behaviour may

also expose them to greater predation risk. In this way personalities can themselves be viewed as

components of life history strategies, leading to an expectation that they will be correlated with – and

trade-off against - other aspects of physiological, reproductive, and behavioural phenotype (Réale et

al. 2010; Sih, Bell, & Johnson 2004; Wolf et al. 2007). Certainly, arguments that trade-offs can maintain

variation in cognitive performance parallel explanations made for widespread presence of personality.

These could be trade-offs among cognitive domains, or alternatively between – for instance – an

overall cognitive performance trait ('general intelligence' (Plomin & Spinath, 2002; Galsworthy et al.,

2005; Burkart et al., 2017)) and other aspects of the phenotype.

Variation in an animal's stress physiology may provide one putative source of among-

individual differences in both personality traits and cognitive performance (Raoult et al., 2017; Gibelli

et al., 2019). The widely used concept of stress coping style model predicts that individuals will vary -

both behaviourally and physiologically- along a proactive/ reactive continuum (Coppens et al., 2010;

Koolhaas et al., 1999; Sih, Bell, & Johnson 2004). As originally posited, the model predicts proactive

coping styles will express more 'fight or flight' type behavioural responses induced by adrenaline-

response to stressors. At the other extreme, reactive coping styles will be more behaviourally 'passive'

(e.g., freezing or hiding) and show high HPA(I) activity leading to cortisol response (Øverli et al., 2007;

Carere et al., 2014). Various links to cognitive performance variation have been suggested. For

instance, proactive styles are associated with 'bold', exploratory, risk-taking personalities that may

present with more opportunities to learn initially. Conversely, greater behavioural flexibility

associated with reactive coping styles (Coppens et al., 2010) may be important for tasks such as

reversal learning, that require an ability to acquire (and use) new information under changing

environmental conditions (Koolhaas et al., 1999; Sih & Del Giudice, 2012; Griffin et al., 2015). More

generally, sensitivity to external stressors or challenges could impact performance in cognitive assays

if more stressed individuals are simply more or less motivated and/or are focused on sources of risk

rather than environmental cues of rewards.

Although hypothesised links between stress responsiveness (or coping style) and cognitive

performance seem intuitive, empirical evidence is still limited to a small number of studies (Lukowiak

et al., 2014; Mesquita et al., 2015; Bebus et al., 2016; Bensky et al., 2017; Brust & Guenther, 2017;

Mazza et al., 2018). There are also contrasting studies in which either a weak or no relationship was

detected (Cole et al., 2011; Carazo et al., 2014; Guillette et al., 2015). It is also possible that

relationships are variable across different aspects of cognition. For instance in sailfin mollies (*Poecilia*

latipinna), individual fish displaying less thigmotaxic behaviour (an anxiety related behaviour in fish)

performed better in a discrimination learning task than highly anxious individuals, whereas the

opposite was found in a reversal learning task (Gibelli et al., 2019). Clearly, there is need for more

empirical work before a clear picture of the complex relationship between variation in cognitive

performance and stress responsiveness/coping style is understood. Here we address this broad goal

by testing the hypothesis that individual differences in cognitive performance and stress

responsiveness are correlated in male Trinidadian guppies (*Poecilia reticulata*).

The guppy is a freshwater poeciliid fish that is widely used as a model in behavioural and

evolutionary ecology. Methods for assaying among-individual 'personality' variation are well

established in this species generally (Burns & Rodd, 2008; White et al., 2016), while guppies have been

used in cognitive studies that target learning colour discriminations (Trompf & Brown, 2014; Buechel

et al., 2018), numerical discriminations (Kotrschal et al., 2013; Lucon-Xiccato & Bisazza, 2017b),

reversal learning (Buechel et al., 2018), spatial learning (Lucon-Xiccato & Bisazza, 2017c) and

inhibitory control (e.g. Lucon-Xiccato & Bisazza, 2016). Here, we investigate the relationship between

behavioural stress response and performance in a spatial learning task in which male guppies

repeatedly navigated a maze to access females as a reward. The cognitive task was repeated using a

second, differently structured maze in order that we could assess not just variation in learning within

a single spatial context, but also ask whether – for instance – individuals displaying greater

performance in trials using the first maze subsequently also performed better in the second. In the

wild, male guppies usually utilize large home ranges during mate search and foraging (Croft et al.,

2003), and as such spatial learning is expected to be an ecologically relevant trait (Brown et al., 2005).

For our measure of stress responsiveness, we utilise 'Open Field Trials' (OFT). Widely used across

species as a paradigm for characterising behavioural differences related to exploration, activity, and

'shy-bold' type variation (Gosling, 2001; Bell et al., 2009), previous studies on this captive population

of guppies have highlighted the utility of OFTs for assaying behavioural stress response (see e.g.,

Prentice et al., 2020). Observed behaviours expressed in the OFT are both repeatable and plastic with

respect to experimentally-manipulated stressor severity (specifically perceived predation risk)

(Houslay et al., 2018). We also know from pedigree-based quantitative genetic studies that individual

(mean) behaviours and their predictability (defined as within-individual variance) are heritable (White,

2019, 2019; Prentice, 2020). Furthermore, there is evidence of genetic integration between OFT

behaviour and cortisol expression, strengthening the view that the OFT provides an appropriate assay

of behavioural stress response (Houslay et al., 2019).

In what follows we: i) test for evidence of learning in naïve guppies repeatedly exposed to a

spatial learning task (maze), ii) ask whether individuals differ in cognitive performance across

repeated trials and if so; iii) whether performance in the first maze predicts performance in a second

spatial context (i.e. reconfigured maze). We predict that time to complete the mazes (our proxy of

cognitive performance) will, on average, improve with experience consistent with spatial "learning",

but that individuals will differ consistently in cognitive performance within each maze. We also predict

that individual performance in the first maze will be positively correlated with performance in the

second, suggesting stable differences in cognitive ability, although we acknowledge proactive

interference (difficulty inhibiting memory or previously learnt associations; Shettleworth, 2009a) may

affect performance in the second maze. Finally, iv) we test the hypothesis that individual differences

in cognitive performance will be associated with differences in stress responsiveness. Although

empirical evidence suggests potential relationships in both directions between stress responsiveness

and cognitive performance, with the current absence of specific models, we make no *a priori*

predictions about the sign of the relationship here.

Methods

Study site and housing

All behavioural assays were carried out on guppies from a captive population (derived from wild fish

collected in the Aripo River, Trinidad in 2008) housed at the University of Exeter's Penryn campus.

Adult males (n = 64) were randomly sampled from the stock population, and housed in groups of 8 in

separate home tanks (15 l, 18.5 × 37 × 22 cm) maintained at 23–24°C on a 12:12 light/dark cycle. The

tanks shared a recirculating sump water supply which underwent a 25% water change once per week.

All fish were fed to satiation twice daily on commercial flake food and live brine shrimp (*Artemia*

salina) to control as much as possible for energetic and nutritional states prior to testing. We elected

to focus on males only for several reasons. First, pilot studies showed a high occurrence of 'freezing'

behaviour in females (relative to males) when introduced to the maze. While freezing can be a

component of the behavioural stress response (Houslay et al., 2018), we considered that frequent

occurrence during the cognitive assay would complicate data interpretation. Second, males show

consistent sexual reproductive motivation towards females (Burns & Rodd, 2008), enabling the use of

females as a 'reward' for males solving the maze (Kotrschal et al., 2015). Third, male guppies exhibit

distinctive markings and colouration on body and fins. By recording and sketching these for each fish

we were able identify individuals within groups without the need to subject individuals to invasive

tagging.

Ethics

This work was conducted under the auspices of the Animals (Scientific Procedures Act) out with

approval of the University of Exeter research ethics committee, under licence from the Home Office

(UK) (Licence Number PPL30/3256). Experimental procedures and behavioural assays were developed

in accordance with the principles of the three Rs and ASAB guidelines (Buchanan et al., 2020) for use

of animals. All periods of handling and emersion were kept to a minimum and only fish deemed

healthy and exhibiting normal behaviour were used in trials. At the end of the experiment, fish were

returned to a designated 'retirement' tank (containing females as well as males) and not used in any

further experiments.

<u>*Overview of behavioural testing scheme*</u>

We used a repeated measures approach to test for among-individual (co)variation in spatial learning

performance and stress responsiveness. Spatial learning was first assessed by repeatedly trialling

individuals in a maze apparatus (Figure 1). Each individual fish was tested once per day for 11

consecutive days with reduction in time to complete the maze interpreted as 'learning'. This is

consistent with previous studies using either time to complete an objective or to perform a particular

task to investigate variation in cognitive performance among-individuals (Guillette et al., 2015; Lucon-

Xiccato & Bisazza, 2016; Mazza et al., 2018; Zidar et al., 2018). We acknowledge that this

interpretation strictly requires the implicit assumption that the contribution of any other factors to

among-individual variation (e.g., motivation, energetic state, experience previous to the experiment;

Rowe & Healy 2014) is negligible relative to differential cognitive performance. We attempted to

mitigate against other sources of among-individual variation as far as possible using standardised

housing and husbandry conditions.

Following completion of 11 spatial learning trials using Maze A, individuals were tested for

stress responsiveness three times each over a three-week period using Open Field Trials (OFT) with a

mean (range) of 4 (1-5) days between successive trials. Fish were then retested in a second maze

(Maze B) with a different layout, with repeat trials conducted (as before) one per day for 11

consecutive days. Thus, in total, the design called for all individuals to complete 22 spatial learning

trials, 11 on each of two different maze layouts (distributed across two different mazes) and three OFT

over a total testing period of 43 days. Note that the sample size declined slightly across the

experiment as (i) a few mortalities occurred naturally within the testing period and, (ii) we proactively

'retired' any fish not deemed to be feeding well and behaving normally in their home tanks as a

precaution against cumulative adverse effects arising. Thus 63 fish experienced Maze A, which

declined to n=60 at trial 11 and OFT testing. Five fish were then removed prior to experiencing Maze

B (n= 55 at trial 1 and n=53 at trial 11).

<u>Spatial Learning Trials</u>

In order to facilitate more rapid data collection, a single aquarium (25 x 45 x 25cm) was divided into

two, with each half containing an identical version of maze A (A1, A2). Two replicates of maze B were

similarly constructed (Figure 1). This allowed two fish to be tested concurrently during trials. Each

maze consisted of 6 opaque Perspex panels (8 cm), spaced 5cm apart (Figure 1). A visually transparent

perforated panel at one end of each maze was used to separate a small holding area (12.5 x 10 x 25

cm) contain two adult females selected randomly from stock. During trials the experimental maze

tanks were lit from below by one fluorescent lamp and filled to a depth of 8 cm with room temperature

water (approx. 23-24 °C). The water was taken from the same recirculating system used to house the

male groups and was changed between each housing group (i.e. after every 4 runs with two fish

trialled per run). Stimulus females were also changed at the same time.

At each trial, two males were individually netted from their home tank and quickly identified

from natural markings. Each was randomly allocated to one of the two maze replicates and carefully

placed within a perforated plastic tube at the start of the maze (Figure 1). They were given 60 s to

acclimate before the plastic tubes were removed. A Sunkwang C160 video camera mounted above the

tank   allowed   the   fish   to   be   observed   without   disturbance.   Tracking   software

(http://www.biobserve.com) was used to determine *maze time*, measured as the total time taken to

complete the maze after fish were released from the perforated plastic tube (with completion defined

as reaching the 'end' zone; Figure 1). On reaching the 'end' zone, individuals were given 60 s

undisturbed visual access to the females before an opaque plastic sheet was inserted to obstruct

females from view. Following the 60 s reward period, fish were netted and returned to the home

tank. To ensure standardized exposure to the reward stimulus, individuals that did not complete the

maze within 480s of being released from the tube were gently guided through the maze to the end

zone using a net behind them and then experienced 60 s visual access to the females. Following the

60 s reward period, fish were netted and returned to the home tank. These fish were assigned a right

censored value of 480 second for *maze time*.

<u>Open Field Trial (OFT)</u>

OFTs to characterise stress responsiveness closely followed the protocol described in White et al.

(2016). For each trial, a single individual was netted from the home tank, quickly identified from

natural markings and introduced gently into the centre of an open arena (a 30 × 20 cm tank filled to 5

cm water placed on a light-box). A cardboard screen around the tank prevented visual disturbance

and a Sunkwang C160 video camera mounted above the arena again allowed movement to be tracked.

Following a 30 s acclimation period, individuals' movements were tracked for 4 minutes and 30 s to

determine *track length* (total distance swum (cm)) and *area covered* (percent of tank area covered).

These two observed behaviours which are known to be repeatable and heritable in this population

(Houslay, 2018; White, 2019, 2019), were used to calculate the derived trait of *relative area* following

Houslay et al. (2019). *Relative area* is the observed area covered in the trial minus the expected area

covered under a simulated 'random swim' of length equal to the observed track length (see Houslay

et al. (2019) for further detail on simulations). Low values of *relative area* result from a 'flight type'

behavioural stress response in which individuals swim rapidly (yielding a high track length) but largely

cover the same small area, thus covering relatively little area of tank arena. Low values of relative area

are strongly correlated with thigmotaxis in this species, with fish swimming repeatedly across sections

next to the tank walls and seeking escape from the tank arena. In contrast, high values of *relative area*

correspond to efficient exploration (i.e. a high proportion of the arena covered given distance swum),

by putatively less stressed fish.

Statistical Analysis

Data from both types of behavioural assay were analysed using univariate and multivariate linear

mixed effect models fitted by REML (restricted maximum likelihood) using ASReml within R

(http://www.vsni.com) (Gilmour et al., 2009). By including individual identity as a random effect in

these models we test for and characterise among-individual (co)variation. Traits were mean centred

and scaled to standard deviation units to ease interpretation of results and facilitate convergence of

multivariate models. For *maze time* we did this using the overall mean and standard deviation of

observations from both mazes in order to preserve any differences in the distributions of performance

times between A and B. With traits in standard deviation units (sdu), estimates of among-individual

variance (V_{ind}) can be interpreted as repeatabilities (i.e. proportion of the observed phenotypic

variance explained by among-individual differences). However, we also calculate estimates of

adjusted repeatability (R), the proportion of phenotypic variance explained by consistent among-

individual differences, after controlling for fixed effects on the mean (Nakagawa & Schielzeth, 2010).

Thus $R=V_{ind}/(V_{ind}+V_R)$ where V_R is the residual (within-individual) variance estimated from each model.

The significance of random effects was tested using likelihood ratio tests (LRT), while fixed effects

(included in the various models as described below) were tested using conditional F-statistics. All

models assumed Gaussian error structures, deemed acceptable based on visual inspection of the

model residuals.

<u>*Univariate analyses of maze performance and spatial learning*</u>

We use *maze time* as our observed measure of performance. Here we describe in full the

univariate analysis of data collected in maze A (subsequently *maze time$_A$*). Identical procedures were

then applied to data from maze B. First, we visualised the distribution of *maze time$_A$* across repeat

using box plots and also plotted the proportion of mazes completed as a function of repeat to see if a

pattern of increasing average performance (i.e. decreasing *maze time* and/or increasing proportion of

successful completion) was immediately apparent. Next a series of three nested models with identical

fixed effects but differing random effect structure were fitted to the centred and scaled *maze time$_A$*

data. All models included a fixed effect of *trial number* (the cumulative number of trials experienced

by an individual, treated as a continuous variable), allowing us to test for improvement in the mean

(indicative of learning). Additional fixed effects were included as statistical controls for potential

sources of variance not relevant to hypotheses being tested here. These included time of day (in

minutes after 9 am), maze replicate (as a factor denoting position 1 or 2 in maze tank), and order

caught from the home tank. The latter was to account for any cumulative disturbance effect of

removing fish sequentially from the home tank and/or build-up of chemical cues in the maze between

water changes.

The first univariate model of *maze time* contained no random effects, while the second

contained a random intercept of individual identity. Likelihood ratio test (LRT) comparison of these

models was conducted to test the hypothesis that individuals differ in their average performance

(*maze time$_A$*) across the 11 repeats, and we estimated the (adjusted) repeatability of performance

under the second model. For the LRT we assume twice the difference in model log-likelihoods is

distributed as a 50:50 mix of X^2_1 and X^2_0 following Stram & Lee (1994). The third model was a first

order random regression (i.e. a random slope and intercept model) in which each individual's

deviation from the fixed effect mean *maze time* can change as a linear function of *trial number* (1-11).

Variation in random slopes means that there is among-individual variation around the mean *maze*

time$_A$ - trial number relationship. Thus, LRT comparison of the second and third models thus provides

a test for among-individual variation in learning rate. This comparison is conducted assuming the test

statistic is distributed as X^2_2 since the third model has two extra parameters (a slope variance and a

slope-intercept covariance). Note that among-individual variance in slopes cannot be scaled to a

repeatability as within individual variance in slope is not estimable (using data from a single maze; see

below). Nor is its magnitude directly comparable to random intercept variance since slopes and

intercepts are in different units. However, under the third model, among-individual variance in

learning (slope) means that among-individual variance *maze time$_A$* changes with *trial number*

(Supplementary Information Figure S1). Thus, to understand the biological effect size of estimated

variance in slopes, we use the third model to predict among-individual variance (V_{ind}) and adjusted

repeatability (R) of *maze time$_A$* at both initial (trial 1) and final (trial 11) performance (following e.g.,

Nussey et al. (2007); see also Supplementary Information Table S3 for a didactic explanation of the

linear algebra behind this). We note that among-individual variation at final performance has been

used to infer differences in cognitive ability in studies adopting similar repeated measures designs

(e.g. Langley et al. 2020) and so also has a useful biological interpretation here.

Univariate analysis of relative area

To verify our expectation that individuals would show consistent differences in stress responsiveness,

we fit a simple random intercepts model to (scaled and centred) *relative area*. This model included

fixed effects of trial number (1-3), and time of day (in minutes after 9 am in which each trial took place)

as well as a random effect of individual identity. Adjusted repeatability (R) of *relative area* was

calculated and the significance of among individual variance tested by LRT comparison to a simplified

model with no random effect (assuming the test statistic was distributed as a 50:50 mix of X^2_1 and X^2_0

as above).

Multivariate modelling of Maze A, Maze B and OFT data combined

Finally, to test the predicted correlation structure between cognitive performance and stress

responsiveness, we formulated a trivariate mixed model in which the three response variables were

maze time$_A$, maze time$_B$ and *relative area*. Fixed effects were exactly as described above on all three

traits. Random effects were also as described above (i.e. individual level random intercepts and slopes

for *maze time$_A$* and *maze time$_B$* but a random intercept only for *relative area*) but the multivariate

formulation allowed us to estimate the full 5x5 among-individual covariance matrix (**ID**) among these

effects. Since each observation of a fish provided data on a single trait only, residual covariances

among traits were fixed to zero. After fitting the model, we compared it to a simplified fit in which all

among-trait covariance elements in **ID** were constrained to zero. This provides a global test of

individual covariance between traits. We then scaled estimated pairwise covariances in **ID** to their

corresponding correlations for easier interpretation (noting for a pair of effects x,y the correlation r_{xy}

$= COV_{xy}/(V_xV_y)^{0.5}$. This allowed us to scrutinise the correlation structure between stress responsiveness

and cognitive performance in both mazes A and B, using both final performance and learning rate (i.e.

random regression slope) as measures of cognition.

This model also yields an estimate the individual level correlation in cognitive performance

measures (final *maze time* performance, learning) across mazes. These are not strictly equivalent to

individual repeatabilities of cognitive performance measures across mazes (as opposed to individual

repeatability of *maze time* across trials within mazes) because estimates could be negative (Barbosa

& Morrissey, 2021). However, they can be readily interpreted in those terms; a strong positive

correlation between, for example, individual *learning* in maze A and maze B means this latent variable

is highly repeatable across mazes. Conversely, a negative correlation means that individuals learning

faster in maze A tend to learn more slowly in maze B (and *vice versa*).

Results

Performance in Maze A

Plots of the raw data suggest that average time to complete Maze A decreases across trials (Figure 2),

and this pattern is qualitatively consistent with expectations if (average) performance improves as a

consequence of learning. The mixed model analysis of *maze time$_A$* confirms statistical support for this

with a significant negative effect of trial repeat number (based on the full random slope and intercept

model; coefficient = -0.058 (0.014) sdu, $F_{1,59.8}$ = 17.890, P < 0.001. This effect size equates to an

estimated decrease of 102.39 seconds in average *maze time* over the 11 trials. Other fixed effects of

order caught and maze position were non-significant (see Supplementary Information Table S1).

Likelihood ratio tests (LRT) confirmed among-individual variation in *maze time$_A$* (comparison of null

and random intercept models; $\chi^2_{0,1}$ = 184.713 P < 0.001). Under the random intercept model,

repeatability of *maze time$_A$* conditional on fixed effects was estimated as R_A = 0.379 (0.052).

367   LRT comparison of the random intercept and first order random regression models showed

the latter to be a significantly better fit to the data (χ^2_2 = 26.990, P < 0.001). This comparison provides

evidence for among-individual variance in the rate of change of *maze time$_A$* across repeated trials. We

interpret this (with caveats discussed below) as variation in the rate of spatial learning. Among-

individual variance in intercepts (int) and slope (slp) were estimated as $V_{ind_{int}}$ = 0.402 (0.104) and

$V_{ind_{slp}}$ = 0.006 (0.032) respectively while the among-individual intercept –slope correlation was

estimated as ($r_{ind_{int},ind_{slp}}$= -0.375 (0.168)). Biological interpretation of these parameters is not

completely straightforward. Given the scaling of *trial number* in the random effect structure of the

model (see Supplementary Information Table S3) $V_{ind_{int}}$ is interpretable as among individual variance

in *maze time$_A$* at first trial. While slope variance is in different units and thus not of directly comparable

magnitude, variation in slopes actually means that among-individual variance in the observed trait

(V_{ind} for *maze time$_A$*) changes with trial repeat number. Here the random regression model predicts

values of $V_{ind_{A1}}$ = 0.402 (0.104), and $V_{ind_{A11}}$ = 0.677 (0.157) at first and last trial in maze A

respectively, suggesting more among individual variation in performance at the end of trials than at

the beginning. The negative intercept-slope correlation ($r_{ind_{A.int,A.slp}}$= -0.375 (0.168)) means that

individuals with higher intercepts (high *maze time*$_A$ at trial 1) tended to have lower (i.e., more negative)

slopes indicative of faster learning. The corresponding predictions of repeatability at first and last

observed trial are R_{A1} = 0.429 (0.070) and R_{A11} = 0.560 (0.063). These patterns are represented visually

in Figure 3a, which shows the individual reaction norms predicted from the best linear unbiased

predictions (BLUPs) of random intercept and slope for each fish (following e.g., Houslay & Wilson

(2017)).

Performance in Maze B

In contrast to Maze A, plotting *maze time*$_B$ data reveals no clear increase in performance (i.e. decrease

in time) across trials. The mixed model analysis confirms the lack of improvement in the mean *maze*

time$_B$, with a (non-significant) positive estimate of the trial repeat number effect (from random slope

and intercept model; coefficient =0.019 (0.014), $F_{1,53.8} = 2.054$, $P = 0.175$). Effects of order

caught and maze position were not significant (Supplementary Information Table S1). Likelihood ratio

tests (LRT) between the univariate random intercept model and the null model with no random effect,

shows the presence of significant among-individual variation for *maze time*$_B$ ($\chi^2_{0,1}$= 200.048 P < 0.001),

with a corresponding repeatability estimate of R_B = 0.423 (0.056). The random slope model was a

significantly better fit again (χ^2_2 = 15.926 P = 0.001) providing evidence of among-individual variation

in the performance-trial number relationship. Among-individual variance in intercepts (int) and slope

(slp) were estimated as 0.472 (0.125) and 0.005 (0.034) respectively. The intercept –slope correlation

was negative as in Maze A ($r_{ind_{B.int,B.slp}}$= -0.332 (0.190)). These estimates mean predicted values of

$V_{ind_{B_1}} = 0.472$ (0.125) and $V_{ind_{B_{11}}}$ = 0.659 (0.162)) which correspond to repeatabilities of R_{B1} =

0.471 (0.072) and R_{B11} = 0.554 (0.067). Although there is no (significant) effect of trial number on

mean *maze time*$_B$ the presence of among-individual variance in slope suggest that some individuals

are improving (consistent with learning) while for others performance is tending to get worse across

repeats in Maze B (Figure 3b).

Among-individual differences in OFT behaviour

We found evidence of significant among-individual variation in *relative area*,

$\left(repeatability\,(with\,SE), R = 0.465\,(0.089), \chi^2_{0,1} = 20.421, P < 0.001\right)$. This replicates previous

findings in the same population (Prentice et al., 2020) though the current estimate of repeatability is

somewhat higher, likely due to differences in study design (e.g. the current study used a shorter inter-

observation interval and was limited to males only). Fixed effects from the OFT behaviour models are

presented in the Supplementary Information Table S2 for completeness, although are not directly

relevant to our hypotheses in this study.

Multivariate model

The full multivariate model of *maze time_A, maze time_B* and *relative area* provides evidence of some

significant covariance structure among traits at the individual level (comparison of the full model to

one in which all among-individual between trait covariances are fixed to zero; $\chi^2_8 = 48.844$, P < 0.001).

Examination of the estimated covariances and correlations (Table 1) suggests this result is largely

driven by a strong positive correlation between the individual intercepts for *maze time_A* and *maze*

time_B $\left(r_{ind_{A.int,B.int}} = 0.704\,(0.127)\right)$. Biologically, this means that individual performance at first trial

is strongly positively correlated across mazes. At final trial (i.e. 11), the individual correlation of

performance across mazes is estimated at $r_{ind_{A11,B11}} = 0.629\,(0.119)$. Thus, our results show strong

positive correlations of individual performance as measured by maze time across trials and mazes.

This is not just true at first and last trial, but also for intermediate trial numbers within and across

mazes (see Supplementary Table S3 for estimates and an explanation of their derivation from the

random regression model output). However, taking a reaction norm interpretation of results, we do

not find strong support for repeatable variation in learning rate across mazes. The individual

correlation of reaction norm slopes is positive ($r_{ind_{A.slp,B.slp}}$ = 0.186 (0.246)), but relatively weak and

not statistically significant (assuming approximate 95% confidence intervals of r±1.96SE). Nor do we

find statistical support for among-individual correlation between maze performance intercepts or

slopes (for either maze) and relative area (Table 1).

Discussion

Here, we show evidence of among-individual differences in performance – measured as time to complete a maze – in guppies exposed to a spatial learning test paradigm. Performance of individuals is repeatable both within, and across, the two spatial learning tasks (i.e. mazes) presented. However, the question of whether there is robust evidence of learning, on average or by individual fish, is somewhat less clear cut. In particular, in the first maze used (A) we find evidence of improvement in mean performance consistent with learning (on average). We also find among-individual variation in this rate of improvement, and so – putatively – their rate of learning. However, the same fish exposed to maze B show (on average) no increase in performance across successive trials. We found among-individual correlation structure between performances (i.e. time in the maze) but not learning (i.e. rate of improvement) across the 2 spatial learning tasks. We did not find any significant association between individual differences in maze performance (or learning) and repeatable stress responsiveness as measured in the open field trials. In what follows we describe each of these findings in more detail and discuss them in the wider context of the cognitive literature.

The data from Maze A show that, on average, time to complete the maze improves across repeated trials. This improvement suggests that spatial learning is occurring in the guppies, a finding consistent with previous studies of this species (Kotrschal et al., 2015; Lucon-Xiccato & Bisazza, 2017c; Fong et al., 2019). We also see evidence of consistent, repeatable differences among-individuals in performance in Maze A. This is shown in our reaction norm models as significant among-individual variance in intercepts, which can be understood as performance at first trial. However, using among-individual variation in intercepts and slope to predict the corresponding variance at, and correlation among-, all trials (see Supplementary Information Table S3 for derivation and presentation of these estimates) reveals that in fact individual performance is positively correlated across all trials from 1 to 11. In simple terms, fish that are faster than average at completing Maze A in their first trial, tend to be faster than average across all subsequent trials too. Predicted repeatability of *maze time* is

moderately high relative to many behavioural studies (e.g., 43% at trial 1, 51% at trial 11) but broadly

comparable to estimates reported from similar assays designed to test cognitive variation (see

Cauchoix et al., 2017 for an overview). We note that a contributing factor is likely to be short inter-

observation period (here 24 hrs) typical of cognitive studies, since behavioural repeatabilities

generally decline as this increases (Boulton et al., 2014).

Accepting that improvement across repeated trials can be interpreted as learning (caveats to this

are discussed below), our random regression model also provides evidence for among-individual

variation in spatial learning in Maze A. Usefully, our modelling strategy allowed all observations to

contribute to estimating variance in the latent cognitive trait (learning) while avoiding statistically

problematic 'two-step' analysis (Houslay & Wilson, 2017). Although this strategy is now widely used

in studies of behavioural plasticity, it has not yet been widely adopted by researchers focussing

specifically on animal cognition (but see e.g., Langley et al., 2020). In addition to finding variance in

slopes (learning), we estimated a negative among-individual intercept-slope correlation using the

Maze A data; individuals with higher intercepts (i.e. *maze time* at first trial) tend to have lower (more

negative) slopes. While it is therefore the case that those fish performing poorly initially exhibit higher

rates of learning, it is also true - as noted above - that individual performance (*maze time*) is positively

correlated across trials 1-11. These two results are compatible because differences in learning (slope)

are not sufficiently pronounced that initially poor performing (but fast learning) fish 'overtake' initially

good performing (but slow learning) individuals by trial 11. We cannot comment on what fitness

consequences, if any, the variation detected here would have in wild fish. Nonetheless, this finding

does highlight a danger with any common presumptions that cognitive abilities may be under positive

selection. Here, if we assumed that fitness benefits were accrued by rapidly achieving a spatial task

(e.g. locating a resource) regardless of mechanism, it would be the slower learners that were

advantaged. Thus, while it is tempting to assume fast learners will achieve better outcomes, they may

sometimes simply be those with the 'most room for improvement'.

Thus, findings from Maze A are consistent with our initial predictions that time to complete the

maze would improve (on average) with experience due to spatial learning, but that individuals would

also vary in both performance (*maze time*) and learning (rate of change in performance with

experience). We also found that individuals that were quicker (over all trials) to complete Maze A,

tended to be quicker (over all trials) to complete Maze B. While this could be attributable to cognitive

differences, there are certainly other possibilities. For instance more explorative and/or less

neophobic individuals may be generally faster at solving tasks (Boogert et al., 2006; Bousquet et al.,

2015; Zidar et al., 2018). Similarly there could be among-individual variation in perceived cue salience

(Meyer et al., 2012), individual physiology (Bókony et al., 2014), or motivation (van Horik & Madden,

2016). Regardless of these unknowns, an important difference between Maze A and Maze B was that

we found no evidence of learning on average in the latter. In fact, for Maze B the mean *maze time*

actually increased slightly, though not significantly, across trials. Thus, there is among-individual

variation in intercept (*maze time* at trial 1) and also in slope. Given that there is no (significant) change

in mean performance, but there is significant variation in slopes, we conclude that some individuals

are improving (learning) in Maze B while others are getting worse with experience. We also note that,

as in Maze A, slope variance is present, but not sufficiently high to break down the positive correlation

structure of individual performance (*maze time*) across trials 1-11.

Although we did not formally test whether the average rate of learning across repeated trials

differs between maze A and B it is reasonable to conclude it does (given no overlap of approximate

95% confidence intervals calculated as the slope ±1.96SE). Several possibilities may explain the finding

of spatial learning on average in A but not B. First, the results from maze A may be a false positive

(Sterne & Smith, 2001; Fraser et al., 2018). However coinciding with previous studies which show this

species is capable of learning an initial spatial learning task (Kotrschal et al., 2015; Lucon-Xiccato &

Bisazza, 2017b; Fong et al., 2019), we consider this unlikely. Second, it may be that the layout of maze

B was more challenging to learn. This could certainly be true if, for instance learning to navigate a new

maze following the acquisition of a previously learnt layout poses a more challenging task, for example

due to proactive interference (difficulty inhibiting memory; Shettleworth, 2009a). In this case the

second maze may require more trials to detect improvement. There is some evidence for such effects

in guppies. For instance, Lucon-Xiccato & Bisazza (2014) found that on average guppies took 14.61

trials to learn a reversed colour cue association, while Fong et al., (2019) found that on average, 15.30

trials were required for guppies to learn a reversed maze layout.

A third possible explanation could be that learning is occurring in Maze B for some individuals, but

that others experience some form or 'trial fatigue' or change in 'state' that reduces cognitive

performance and/or motivation. For example, chronic stress effects may be arising from repeated

capture and handling experienced in the experimental design (Huntingford et al., 2006; Warren &

Callaghan, 1976; Wong et al., 2008). Such an effect could manifest in our Maze B data as an increase

in the proportion of maze trials not successfully completed in <480 seconds (assuming trial fatigue

increases the probability of not completing), coupled with a decline in average time across trial repeats

for those trials that were successfully finished (assuming that better performing fish are learning).

However, visual inspection of the data revealed no obvious trend in proportion of trials completed in

<480 seconds, and post hoc analysis of maze B trials with censored records excluded provided no

evidence of improvement either (results not shown). Empiricists obviously seek to minimise the

possibility of chronic stress confounding conclusions from cognitive studies but it can be difficult to

validate the implicit assumption that individuals remain (equally) 'unstressed' over experimental

periods requiring repeated observations. Here we suggest this is a plausible hypothesis but not one

we can currently test.

While the influence, or otherwise, of chronic stress on our results necessarily remains speculative,

our experiment does confirm among-individual variation in acute stress behaviour as measured by

relative area in the OFT. This replicates earlier results using independent data sets of fish from the

same captive population (White, 2016; Houslay, 2019; Prentice, 2020). Acute stressor exposure can

affect cognitive performance in spatial learning tasks in both mammals and fish (Gaikwad et al., 2011;

Wong et al., 2019). At the individual level, there is also evidence that acute stress responses can

predict outcomes under longer term chronic and/or repeated stressor exposure (Segerstrom & Miller,

2004; Salak-Johnson & McGlone, 2007; Øverli et al., 2007). However, we find no evidence of strong

relationships between acute stress behaviour and performance or learning in either maze. Thus, we

find no support for the prediction, made under the stress coping style model, that (acute) stress

responsiveness will covary with cognitive performance (Coppens et al., 2010; Sih & Del Giudice, 2012;

Griffin et al., 2015).

In summary, here we have evidence of consistent differences among-individuals in spatial task

performance in the guppy *P. reticulata*. Individual performance is repeatable across trials within- and

between two different spatial tasks (i.e. maze layouts). This among-individual variation in performance

is consistent with underlying differences at cognitive factors but differences in 'personality' (e.g.

neophobia, exploratory tendency) may also contribute. We also find evidence of improved

performance with experience, consistent with spatial learning. In both tasks, variation around the

trajectory of mean performance across trial number was present. While this means individuals can be

considered as differing in 'spatial learning rate' it is important to note that performance declines for

some individuals, especially in the second maze where there was no improvement in average time

across 11 trials. We show here that an individual's (repeatable) behavioural response to an acute

stress stimulus does not predict either average performance in the maze or learning rate. However,

we suggest the possibility that cumulative, chronic stress effects may contribute to declining

performance (or reduced improvement) in our study. If individuals generally differ in susceptibility to

chronic stress, this may represent a widespread but currently poorly acknowledged challenge for

characterisation of cognitive variation in animal studies.

Figure 1. Aerial view of the maze designs used in the experiments (A and B), each tank was split into two identical mazes (1 and 2). Blue circles indicate perforated plastic tubes where individual fish started each trial, and time taken to reach the 'end zone' was recorded each trial.

a)   b)

c)   d)

Figure 2. Plots of raw data of *maze time* across both maze designs. Boxplots (a) and (c) show the data

distributions for time to complete Maze A and Maze B respectively across the 11 trials. Grey boxes

display data of all individuals and red boxes represent only those individuals that successfully

completed the task within 480 s. Horizontal lines within box correspond to behavioural medians, box

boundaries correspond to first and third quartiles. When present, whiskers correspond to 10th and

90th percentiles, and points correspond to outliers. Plots (b) and (d) represent mean and standard

errors for time to complete Maze A and B respectively. Colours represent the same groups; black error

bars represent mean and standard errors of *maze time* for all individuals, and red represent only those

individuals that successfully completed the maze in the allocated time.

a)

b)

Figure 3. Spatial learning traits across Maze A and Maze B as a function of trial number, *maze time$_A$*

(a), and *maze time$_B$* (b). Grey lines represent individual predicted reaction norms (BLUPs) from

univariate random slope models for each trait. Coloured lines are used to illustrate reaction norms for

a small random set of arbitrarily chosen individuals tested in both mazes (colours represent the same

individuals across panels). The black dashed line represents the trend in fixed effect mean *maze time*

across repeat trials

588 **Table 1.** Among individual variance–covariance–correlation matrix from the final trivariate model of *maze time$_A$*, *relative area* and *maze time$_B$*. Variances are

589 shown on the diagonal (dark grey shading), with covariances above and correlations below. Light grey shading denotes within trait covariance/correlation

590 estimates (i.e. between reaction norm intercepts and slopes). Standard errors are shown in parentheses and bold font denotes nominally significant pairwise

591 estimates assuming approximate 95% CI of ± 1.96SE).

	Maze time$_A$		Relative area	Maze time$_B$	
	intercept$_A$	slope$_A$		intercept$_B$	slope$_B$
intercept$_A$	**0.402 (0.104)**	-0.019 (0.012)	0.096 (0.079)	**0.305 (0.089)**	-0.005 (0.010)
slope$_A$	**-0.374 (0.168)**	**0.007 (0.002)**	-0.002 (0.011)	0.005 (0.011)	0.001 (0.001)
Relative area	0.226 (0.178)	-0.034 (0.202)	**0.452 (0.118)**	0.035 (0.086)	0.012 (0.011)
intercept$_B$	**0.704 (0.127)**	0.094 (0.203)	0.075 (0.186)	**0.467 (0.122)**	-0.016 (0.012)
slope$_B$	-0.103 (0.226)	0.186 (0.246)	0.255 (0.221)	-0.335 (0.189)	**0.005 (0.002)**

References

Ashton, B. J., Ridley, A. R., Edwards, E. K., & Thornton, A. (2018). Cognitive performance is linked to group size and affects fitness in Australian magpies. *Nature*, *554*(7692), 364–367. https://doi.org/10.1038/nature25503

Barbosa, M., & Morrissey, M. B. (2021). The distinction between repeatability and correlation in studies of animal behaviour. *Animal Behaviour*, *175*, 201–217. https://doi.org/10.1016/J.ANBEHAV.2021.03.008

Bebus, S. E., Small, T. W., Jones, B. C., Elderbrock, E. K., & Schoech, S. J. (2016). Associative learning is inversely related to reversal learning and varies with nestling corticosterone exposure. *Animal Behaviour*, *111*, 251–260. https://doi.org/10.1016/j.anbehav.2015.10.027

Bell, A. M., Hankison, S. J., & Laskowski, K. L. (2009). The repeatability of behaviour: a meta-analysis. In *Animal Behaviour* (Vol. 77, Issue 4, pp. 771–783). https://doi.org/10.1016/j.anbehav.2008.12.022

Bensky, M. K., Paitz, R., Pereira, L., & Bell, A. M. (2017). Testing the predictions of coping styles theory in threespined sticklebacks. *Behavioural Processes*, *136*, 1–10. https://doi.org/10.1016/j.beproc.2016.12.011

Bókony, V., Lendvai, Á. Z., Vágási, C. I., Pătraş, L., Pap, P. L., Németh, J., Vincze, E., Papp, S., Preiszner, B., Seress, G., & Liker, A. (2014). Necessity or capacity? Physiological state predicts problem-solving performance in house sparrows. *Behavioral Ecology*, *25*(1), 124–135. https://doi.org/10.1093/beheco/art094

Boogert, N. J., Madden, J. R., Morand-Ferron, J., & Thornton, A. (2018). Measuring and understanding individual differences in cognition. In *Philosophical Transactions of the Royal Society B: Biological Sciences* (Vol. 373, Issue 1756, p. 20170280). https://doi.org/10.1098/rstb.2017.0280

Boogert, N. J., Reader, S. M., & Laland, K. N. (2006). The relation between social rank, neophobia and

individual learning in starlings. *Animal Behaviour*, *72*(6), 1229–1239.

https://doi.org/10.1016/j.anbehav.2006.02.021

Boulton, K., Grimmer, A. J., Rosenthal, G. G., Walling, C. A., & Wilson, A. J. (2014). How stable are

personalities? A multivariate view of behavioural variation over long and short timescales in

the sheepshead swordtail, Xiphophorus birchmanni. *Behavioral Ecology and Sociobiology*,

68(5), 791–803. https://doi.org/10.1007/s00265-014-1692-0

Bousquet, C. a. H., Petit, O., Arrivé, M., Robin, J. P., & Sueur, C. (2015). Personality tests predict

responses to a spatial-learning task in mallards, Anas platyrhynchos. *Animal Behaviour*, *110*,

145–154. https://doi.org/10.1016/j.anbehav.2015.09.024

Bridger, D., Bonner, S. J., & Briffa, M. (2015). Individual quality and personality: bolder males are less

fecund in the hermit crab Pagurus bernhardus. *Proc Biol Sci*, *282*(1803), 20142492.

https://doi.org/10.1098/rspb.2014.2492

Brown, C., & Braithwaite, V. A. (2005). Effects of predation pressure on the cognitive ability of the

poeciliid Brachyraphis episcopi. *Behavioral Ecology*, *16*(2), 482–487.

https://doi.org/10.1093/beheco/ari016

Brust, V., & Guenther, A. (2017). Stability of the guinea pigs personality – cognition – linkage over

time. In *Behavioural Processes* (Vol. 134, pp. 4–11).

https://doi.org/10.1016/j.beproc.2016.06.009

Buchanan, K., Perera, T. B. de, … C. C.-A., & 2012, U. (2020). Guidelines for the treatment of animals

in behavioural research and teaching. *Animal Behaviour*, *159*, I–XI.

https://doi.org/10.1016/j.anbehav.2019.11.002

Buechel, S. D., Boussard, A., Kotrschal, A., van Der Bijl, W., & Kolm, N. (2018). Brain size affects

performance in a reversal-learning test. *Proceedings of the Royal Society B: Biological Sciences*,

285(1871). https://doi.org/10.1098/rspb.2017.2031

Burkart, J. M., Schubiger, M. N., & Van Schaik, C. P. (2017). The evolution of general intelligence.

Behavioral and Brain Sciences, *40*, e195. https://doi.org/10.1017/S0140525X16000959

Burns, J. G., & Rodd, F. H. (2008). Hastiness, brain size and predation regime affect the performance

of wild guppies in a spatial memory task. *Animal Behaviour*, *76*(3), 911–922.

https://doi.org/10.1016/j.anbehav.2008.02.017

Carazo, P., Noble, D. W. A., Chandrasoma, D., & Whiting, M. J. (2014). Sex and boldness explain

individual differences in spatial learning in a lizard. *Proceedings of the Royal Society B:*

Biological Sciences, *281*(1782), 20133275. https://doi.org/10.1098/rspb.2013.3275

Carere, C., Maestripieri, D., Caramaschi, D., Carere, C., Sgoifo, A., & Koolhaas, J. M. (2014).

Neuroendocrine and Autonomic Correlates of Animal Personalities. In *Animal Personalities* (pp.

353–377). https://doi.org/10.7208/chicago/9780226922065.003.0013

Cauchoix, M., Hermer, E., Chaine, A. S., & Morand-Ferron, J. (2017). Cognition in the field:

Comparison of reversal learning performance in captive and wild passerines. *Scientific Reports*,

7(1), 12945. https://doi.org/10.1038/s41598-017-13179-5

Cole, E. F., Cram, D. L., & Quinn, J. L. (2011). Individual variation in spontaneous problem-solving

performance among wild great tits. *Animal Behaviour*, *81*(2), 491–498.

https://doi.org/10.1016/j.anbehav.2010.11.025

Coppens, C. M., de Boer, S. F., & Koolhaas, J. M. (2010). Coping styles and behavioural flexibility:

towards underlying mechanisms. *Philosophical Transactions of the Royal Society of London.*

Series B, Biological Sciences, *365*(1560), 4021–4028. https://doi.org/10.1098/rstb.2010.0217

Croft, D. P., Albanese, B., Arrowsmith, B. J., Botham, M., Webster, M., & Krause, J. (2003). Sex-biased

movement in the guppy (Poecilia reticulata). *Oecologia*, *137*(1), 62–68.

https://doi.org/10.1007/s00442-003-1268-6

Dingemanse, N. J., Both, C., Drent, P. J., & Tinbergen, J. M. (2004). Fitness consequences of avian

personalities in a fluctuating environment. *Proceedings of the Royal Society of London, Series B:*

Biological Sciences, *271*(1541), 847–852.

Dingemanse, N. J., & Dochtermann, N. A. (2013). Quantifying individual variation in behaviour:

Mixed-effect modelling approaches. *Journal of Animal Ecology*, *82*(1), 39–54.

https://doi.org/10.1111/1365-2656.12013

Dingemanse, N. J., & Réale, D. (2005). Natural Selection and Animal Personality. *Behaviour*,

142(9/10), 1159–1184. https://doi.org/10.1163/156853905774539445

Fong, S., Buechel, S. D., Boussard, A., Kotrschal, A., & Kolm, N. (2019). Plastic changes in brain

morphology in relation to learning and environmental enrichment in the guppy (Poecilia

reticulata). *Journal of Experimental Biology*, *222*(10). https://doi.org/10.1242/jeb.200402

Fraser, H., Parker, T., Nakagawa, S., Barnett, A., & Fidler, F. (2018). Questionable research practices

in ecology and evolution. *PLoS ONE*, *13*(7), e0200303.

https://doi.org/10.1371/journal.pone.0200303

Gaikwad, S., Stewart, A., Hart, P., Wong, K., Piet, V., Cachat, J., & Kalueff, A. V. (2011). Acute stress

disrupts performance of zebrafish in the cued and spatial memory tests: The utility of fish

models to study stress-memory interplay. *Behavioural Processes*, *87*(2), 224–230.

https://doi.org/10.1016/j.beproc.2011.04.004

Galsworthy, M. J., Paya-Cano, J. L., Liu, L., Monleón, S., Gregoryan, G., Fernandes, C., Schalkwyk, L.

C., & Plomin, R. (2005). Assessing reliability, heritability and general cognitive ability in a

battery of cognitive tasks for laboratory mice. *Behavior Genetics*, *35*(5), 675–692.

https://doi.org/10.1007/s10519-005-3423-9

Gibelli, J., Aubin-Horth, N., & Dubois, F. (2019). Individual differences in anxiety are related to

differences in learning performance and cognitive style. *Animal Behaviour*, *157*, 121–128.

https://doi.org/10.1016/j.anbehav.2019.09.008

Gilmour, A. R., Gogel, B. J., Cullis, B. R., Thompson, R., & Butler, D. (2009). ASReml user guide release

3.0. *VSN International Ltd, Hemel Hempstead, UK*.

Gosling, S. D. (2001). From mice to men: What can we learn about personality from animal research?

Psychological Bulletin, *127*(1), 45–86. https://doi.org/10.1037/0033-2909.127.1.45

Griffin, A. S., Guillette, L. M., & Healy, S. D. (2015). Cognition and personality: An analysis of an

emerging field. In *Trends in Ecology and Evolution* (Vol. 30, Issue 4, pp. 207–214).

https://doi.org/10.1016/j.tree.2015.01.012

Guillette, L. M., Hahn, A. H., Hoeschele, M., Przyslupski, A. M., & Sturdy, C. B. (2015). Individual

differences in learning speed, performance accuracy and exploratory behaviour in black-

capped chickadees. *Animal Cognition*, *18*(1), 165–178. https://doi.org/10.1007/s10071-014-

0787-3

Healy, S. D. (2019). The face of animal cognition. In *Integrative Zoology* (Vol. 14, Issue 2, pp. 132–

144). John Wiley & Sons, Ltd. https://doi.org/10.1111/1749-4877.12361

Healy, S. D., & Rowe, C. (2010). Information Processing: The Ecology and Evolution of Cognitive

Abilities. In *Evolutionary Behavioral Ecology* (pp. 162–174). https://eprints.ncl.ac.uk/139823

Houslay, T. M., Earley, R. L., White, S. J., Lammers, W., Grimmer, A. J., Travers, L. M., Johnson, E. L.,

Young, A. J., & Wilson, A. J. (2019). Genetic integration of the stress response. *BioRxiv*, 770586.

https://doi.org/10.1101/770586

Houslay, T. M., Vierbuchen, M., Grimmer, A. J., Young, A. J., & Wilson, A. J. (2018). Testing the

stability of behavioural coping style across stress contexts in the Trinidadian guppy. *Functional*

Ecology, *32*(2), 424–438. https://doi.org/10.1111/1365-2435.12981

Houslay, T. M., & Wilson, A. J. (2017). Avoiding the misuse of BLUP in behavioural ecology.

Behavioral Ecology, *28*(4), 948–952. https://doi.org/10.1093/beheco/arx023

Huntingford, F. A., Adams, C., Braithwaite, V. A., Kadri, S., Pottinger, T. G., Sandøe, P., & Turnbull, J.

F. (2006). Current issues in fish welfare. In *Journal of Fish Biology* (Vol. 68, Issue 2, pp. 332–

372). https://doi.org/10.1111/j.0022-1112.2006.001046.x

Koolhaas, J. M., Korte, S. M., De Boer, S. F., Van Der Vegt, B. J., Van Reenen, C. G., Hopster, H., De

Jong, I. C., Ruis, M. A. W., & Blokhuis, H. J. (1999). Coping styles in animals: Current status in

behavior and stress- physiology. *Neuroscience and Biobehavioral Reviews*, *23*(7), 925–935.

https://doi.org/10.1016/S0149-7634(99)00026-3

Kotrschal, A., Corral-Lopez, A., Amcoff, M., & Kolm, N. (2015). A larger brain confers a benefit in a

spatial mate search learning task in male guppies. *Behavioral Ecology*, *26*(2), 527–532.

https://doi.org/10.1093/beheco/aru227

Kotrschal, A., Rogell, B., Bundsen, A., Svensson, B., Zajitschek, S., Brännström, I., Immler, S.,

Maklakov, A. A., & Kolm, N. (2013). Artificial selection on relative brain size in the guppy reveals

costs and benefits of evolving a larger brain. *Current Biology*, *23*(2), 168–171.

Langley, E. J. G., Adams, G., Beardsworth, C. E., Dawson, D. A., Laker, P. R., van Horik, J. O.,

Whiteside, M. A., Wilson, A. J., & Madden, J. R. (2020). Heritability and correlations among

learning and inhibitory control traits. *Behavioral Ecology*.

https://doi.org/10.1093/beheco/araa029

Lucon-Xiccato, T., & Bisazza, A. (2014). Discrimination reversal learning reveals greater female

behavioural flexibility in guppies. *Biology Letters*, *10*(6), 20140206.

Lucon-Xiccato, T., & Bisazza, A. (2016). Male and female guppies differ in speed but not in accuracy

in visual discrimination learning. *Animal Cognition*, *19*(4), 733–744.

https://doi.org/10.1007/s10071-016-0969-2

Lucon-Xiccato, T., & Bisazza, A. (2017a). Individual differences in cognition among teleost fishes. In

Behavioural Processes (Vol. 141, pp. 184–195). https://doi.org/10.1016/j.beproc.2017.01.015

Lucon-Xiccato, T., & Bisazza, A. (2017b). Sex differences in spatial abilities and cognitive flexibility in

the guppy. *Animal Behaviour*, *123*, 53–60. https://doi.org/10.1016/j.anbehav.2016.10.026

Lucon-Xiccato, T., & Bisazza, A. (2017c). Complex maze learning by fish. *Animal Behaviour*, *125*, 69–

75. https://doi.org/10.1016/j.anbehav.2016.12.022

Lukowiak, K., Orr, M., de Caigny, P., Lukowiak, K. S., Rosenegger, D., Han, J. Il, & Dalesman, S. (2014).

Ecologically relevant stressors modify long-term memory formation in a model system.

Behavioural Brain Research, *214*(1), 18–24. https://doi.org/10.1016/j.bbr.2010.05.011

Mazza, V., Eccard, J. A., Zaccaroni, M., Jacob, J., & Dammhahn, M. (2018). The fast and the flexible:

cognitive style drives individual variation in cognition in a small mammal. *Animal Behaviour*,

137, 119–132. https://doi.org/10.1016/j.anbehav.2018.01.011

Medina-García, A., Jawor, J. M., & Wright, T. F. (2017). Cognition, personality, and stress in

budgerigars, Melopsittacus undulatus. *Behavioral Ecology*, *28*(6), 1504–1516.

https://doi.org/10.1093/beheco/arx116

Mesquita, F. O., Borcato, F. L., & Huntingford, F. A. (2015). Cue-based and algorithmic learning in

common carp: A possible link to stress coping style. *Behavioural Processes*, *115*, 25–29.

https://doi.org/10.1016/j.beproc.2015.02.017

Meyer, P. J., Lovic, V., Saunders, B. T., Yager, L. M., Flagel, S. B., Morrow, J. D., & Robinson, T. E.

(2012). Quantifying individual variation in the propensity to attribute incentive salience to

reward cues. *PLoS ONE*, *7*(6). https://doi.org/10.1371/journal.pone.0038987

Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: A practical

guide for biologists. In *Biological Reviews* (Vol. 85, Issue 4, pp. 935–956).

https://doi.org/10.1111/j.1469-185X.2010.00141.x

Niemelä, P. T., Vainikka, A., Forsman, J. T., Loukola, O. J., & Kortet, R. (2013). How does variation in

the environment and individual cognition explain the existence of consistent behavioral

differences? *Ecology and Evolution*, *3*(2), 457–464. https://doi.org/10.1002/ece3.451

Nussey, D. H., Wilson, A. J., & Brommer, J. E. (2007). The evolutionary ecology of individual

phenotypic plasticity in wild populations. *Journal of Evolutionary Biology*, *20*(3), 831–844.

https://doi.org/10.1111/j.1420-9101.2007.01300.x

Øverli, Ø., Sørensen, C., Pulman, K. G. T., Pottinger, T. G., Korzan, W., Summers, C. H., & Nilsson, G. E.

(2007). Evolutionary background for stress-coping styles: Relationships between physiological,

behavioral, and cognitive traits in non-mammalian vertebrates. In *Neuroscience and*

Biobehavioral Reviews (Vol. 31, Issue 3, pp. 396–412). Pergamon.

https://doi.org/10.1016/j.neubiorev.2006.10.006

Plomin, R., & Spinath, F. M. (2002). Genetics and general cognitive ability (g). In *Trends in Cognitive*

Sciences (Vol. 6, Issue 4, pp. 169–176). https://doi.org/10.1016/S1364-6613(00)01853-2

Prentice, P. M., Houslay, T. M., Martin, J. G. A., & Wilson, A. J. (2020). Genetic variance for

behavioural 'predictability' of stress response. *Journal of Evolutionary Biology*, *33*(5), 642–652.

https://doi.org/10.1111/jeb.13601

Quinn, J. L., Cole, E. F., Bates, J., Payne, R. W., & Cresswell, W. (2012). Personality predicts individual

responsiveness to the risks of starvation and predation. *Proceedings of the Royal Society B:*

Biological Sciences, *279*(1735), 1919–1926. https://doi.org/10.1098/rspb.2011.2227

Quinn, J. L., Cole, E. F., Reed, T. E., & Morand-Ferron, J. (2016). Environmental and genetic

determinants of innovativeness in a natural population of birds. *Philosophical Transactions of*

the Royal Society B: Biological Sciences, *371*(1690). https://doi.org/10.1098/rstb.2015.0184

Raoult, V., Trompf, L., Williamson, J. E., & Brown, C. (2017). Stress profile influences learning

approach in a marine fish. *PeerJ*, *2017*(6). https://doi.org/10.7717/peerj.3445

Réale, D., Garant, D., Humphries, M. M., Bergeron, P., Careau, V., & Montiglio, P. O. (2010).

Personality and the emergence of the pace-of-life syndrome concept at the population level.

Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences,

365(1560), 4051–4063. https://doi.org/10.1098/rstb.2010.0208

Réale, D., Reader, S. M., Sol, D., McDougall, P. T., & Dingemanse, N. J. (2007). Integrating animal

temperament within ecology and evolution. *Biological Reviews*, *82*(2), 291–318.

Roff, D. A. (2002). *Life history evolution* (p. 527). Sinauer Associates.

Rowe, C., & Healy, S. D. (2014). Measuring variation in cognition. In *Behavioral Ecology* (Vol. 25, Issue

6, pp. 1287–1292). https://doi.org/10.1093/beheco/aru090

Salak-Johnson, J. L., & McGlone, J. J. (2007). Making sense of apparently conflicting data: stress and

immunity in swine and cattle. In *Journal of animal science* (Vol. 85, Issue 13 Suppl, pp. 81–88).

https://doi.org/10.2527/jas.2006-538

Segerstrom, S. C., & Miller, G. E. (2004). Psychological stress and the human immune system: A

meta-analytic study of 30 years of inquiry. *Psychological Bulletin*, *130*(4), 601–630.

https://doi.org/10.1037/0033-2909.130.4.601

Shettleworth, S. J. (2009). Cognition, Evolution, and Behavior. *Oxford University Press*.

Shettleworth, S. J. (2010). Getting Around: Spatial Cognition. In *Cognition, Evolution, and Behavior*

(second, pp. 261–312). Oxford University Press.

Sih, A., Bell, A., & Johnson, J. C. (2004). Behavioral syndromes: an ecological and evolutionary

overview. *Trends in Ecology & Evolution*, *19*(7), 372–378.

Sih, A., Bell, A. M., Johnson, J. C., & Ziemba, R. E. (2004). Behavioral Syndromes: An Integrative

Overview. *The Quarterly Review of Biology*, *79*(3), 241–277. https://doi.org/10.1086/422893

Sih, A., & Del Giudice, M. (2012). Linking behavioural syndromes and cognition: a behavioural

ecology perspective. *Philosophical Transactions of the Royal Society B-Biological Sciences*,

367(1603), 2762–2772. https://doi.org/10.1098/rstb.2012.0216

Sterne, J. A. C., & Smith, G. D. (2001). Sifting the evidence—what's wrong with significance tests?

Physical Therapy, *81*(8), 1464–1469. https://doi.org/10.1093/ptj/81.8.1464

Stram, D. O., & Lee, J. W. (1994). Variance Components Testing in the Longitudinal Mixed Effects

Model. *Biometrics*, *50*(4), 1171. https://doi.org/10.2307/2533455

Sweis, B. M., Veverka, K. K., Dhillon, E. S., Urban, J. H., & Lucas, L. R. (2013). Individual differences in

the effects of chronic stress on memory: Behavioral and neurochemical correlates of resiliency.

Neuroscience, *246*, 142–159. https://doi.org/10.1016/j.neuroscience.2013.04.052

Thornton, A., Isden, J., & Madden, J. R. (2014). Toward wild psychometrics: Linking individual

cognitive differences to fitness. *Behavioral Ecology*, *25*(6), 1299–1301.

https://doi.org/10.1093/beheco/aru095

Thornton, A., & Wilson, A. J. (2015). In search of the Darwinian Holy Trinity in cognitive evolution: A

comment on Croston et al. In *Behavioral Ecology* (Vol. 26, Issue 6, pp. 1460–1461).

https://doi.org/10.1093/beheco/arv119

Trompf, L., & Brown, C. (2014). Personality affects learning and trade-offs between private and social

information in guppies, Poecilia reticulata. *Animal Behaviour*, *88*, 99–106.

van Horik, J. O., & Madden, J. R. (2016). A problem with problem solving: Motivational traits, but not

cognition, predict success on novel operant foraging tasks. *Animal Behaviour*, *114*, 189–198.

https://doi.org/10.1016/j.anbehav.2016.02.006

Warren, E. W., & Callaghan, S. (1976). The response of male guppies (Poecilia reticutata, Peters) to

repeated exposure to an open field. *Behavioral Biology*, *18*(4), 499–513.

https://doi.org/10.1016/S0091-6773(76)92524-4

White, S. J., Houslay, T. M., & Wilson, A. J. (2019). Evolutionary genetics of personality in the

Trinidadian guppy II: sexual dimorphism and genotype-by-sex interactions. *Heredity*, *122*(1),

15–28. https://doi.org/10.1038/s41437-018-0083-0

White, S. J., Kells, T., & Wilson, A. J. (2016). Metabolism, personality and pace of life in the

Trinidadian guppy, Poecilia reticulata. *Behaviour*, *153*(13–14), 1517–1543.

https://doi.org/10.1163/1568539X-00003375

White, S. J., & Wilson, A. J. (2019). Evolutionary genetics of personality in the Trinidadian guppy I:

maternal and additive genetic effects across ontogeny. *Heredity*, *122*(1), 1–14.

https://doi.org/10.1038/s41437-018-0082-1

Wilson, A. J., Réale, D., Clements, M. N., Morrissey, M. M., Postma, E., Walling, C. A., Kruuk, L. E. . B.,

& Nussey, D. H. (2010). An ecologists guide to the animal model. In *J Anim Ecol* (Vol. 79, Issue 1,

pp. 13–26). http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2656.2009.01639.x/full

Wolf, M., Van Doorn, G. S., Leimar, O., & Weissing, F. J. (2007). Life-history trade-offs favour the

evolution of animal personalities. *Nature*, *447*(7144), 581–584.

Wong, R. Y., French, J., & Russ, J. B. (2019). Differences in stress reactivity between zebrafish with

alternative stress coping styles. *Royal Society Open Science*, *6*(5), 181797.

https://doi.org/10.1098/rsos.181797

Wong, S. C., Dykstra, M., Campbell, J. M., & Earley, R. L. (2008). Measuring water-borne cortisol in

convict cichlids (Amatitlania nigrofasciata): Is the procedure a stressor? *Behaviour*, *145*(10),

1283–1305. https://doi.org/10.1163/156853908785765863

Zidar, J., Balogh, A., Favati, A., Jensen, P., Leimar, O., Sorato, E., & Løvlie, H. (2018). The relationship

between learning speed and personality is age- and task-dependent in red junglefowl.

Behavioral Ecology and Sociobiology, *72*(10), 168. https://doi.org/10.1007/s00265-018-2579-2

852