Examining Blunted Initial Response to Reward and Recent Suicidal Ideation in

Children and Adolescents Using Event-Related Potentials: Failure to Conceptually

Replicate Across Two Independent Samples

Austin J. Gallyer, M.S.[1], Kreshnik Burani, M.S.[1], Elizabeth M. Mulligan, M.S.[1], Nicholas

Santopetro, B.S.[1], Sean P. Dougherty, M.S.[1], Min Eun Jeon, M.A.[1], Brady D. Nelson, Ph.D.[2],

Thomas E. Joiner, Ph.D.[1], and Greg Hajcak, Ph.D.[1, 3]

[1]Department of Psychology, Florida State University

[2]Department of Psychology, Stony Brook University

[3]Department of Biomedical Sciences, Florida State University

Author Note

This work was supported by grants from the National Institute of Mental Health

(T32MH093311; R01MH106477; R01MH097767). This work was also supported in part by

the Military Suicide Research Consortium (MSRC), an effort supported by the Assistant

Secretary of Defense for Health Affairs under Award No. (W81XWH-16-2-0004). Opinions,

interpretations, conclusions, and recommendations are those of the author and are not

necessarily endorsed by the MSRC or the Department of Defense.

Correspondence concerning this article should be addressed to Austin J. Gallyer,

M.S., 1107 West Call St., Tallahassee, FL, 32306. Email: agallyer@neuro.fsu.edu.

Abstract

A recent study by Tsypes and colleagues (2019) found that children with recent suicidal ideation had blunted neural reward processing, as measured by the reward positivity (RewP), compared to matched controls, and that this difference was driven by reduced neural responses to monetary loss, rather than by blunted neural response to monetary reward. Here, we aimed to conceptually replicate and extend these findings in two samples ($n = 275$, 27 with suicidal ideation; and $n = 235$, 48 with suicidal ideation at T1) of children and adolescents (11 to 15 years and 8 to 15 years, respectively). Results from both samples showed no evidence that children and adolescents with suicidal ideation have abnormal reward or loss processing, nor that reward processing predicts suicidal ideation two years later. The results highlight the need for greater statistical power, as well as continued research examining the neural underpinnings of suicidal thoughts and behaviors.

Keywords: suicidal ideation, reward, ERP, children, reward positivity, RewP, feedback negativity, suicide, STBs

Examining Blunted Initial Response to Reward and Recent Suicidal Ideation in

Children and Adolescents Using Event-Related Potentials: Failure to Conceptually

Replicate Across Two Independent Samples

Suicide is a growing problem among children and adolescents. In the United States from 1999 to 2018, the suicide rate increased from 1.2 to 2.9, and from 8.0 to 11.4, per 100,000 for those 10–14 and 15–19 years old, respectively, primarily among females (Centers for Disease Control, 2018). Moreover, suicide is the second leading cause of death for children and adolescents between 10 and 19 years of age (Centers for Disease Control and Prevention, 2018). Though death by suicide is understandably the focus of suicide prevention efforts, suicidal ideation is itself a serious public health issue among children and adolescents and one deserving of inquiry (Jobes & Joiner, 2019). The necessity of addressing suicidal ideation itself is underscored by the prevalence of suicidal ideation in this age range, with a recent survey of high school students in the United States finding that 18.8% reported seriously considering suicide during the past year (Ivey-Stephenson et al., 2020). Unfortunately, recent meta-analytic work shows that, while treatments for suicidal ideation do have an effect, effect sizes are small and are least effective in younger patients, such as children and adolescents ($g = -.09$; Fox et al., 2020). It has been suggested that this relative lack of treatment effectiveness may be due to failures to determine the causes of suicidal thoughts and behaviors and an overemphasis on risk factors that have modest longitudinal associations with suicidal ideation and other suicide-related outcomes (Fox et al., 2020; Franklin et al., 2017, 2018).

One approach that has been offered as a solution to developing more effective treatments for suicidal ideation is to study its purported causes from a transdiagnostic perspective grounded in neuroscience (Glenn et al., 2017). In particular, the Research Domain Criteria (RDoC), a framework developed by the National Institute of Mental Health

that aims to integrate multiple levels of analysis to identify the underlying neural mechanisms of psychopathology, has been proposed as a way to improve the understanding of suicidal ideation (Glenn et al., 2017; Insel et al., 2010). One of the systems outlined by RDoC that has been suggested to be important for understanding suicidal ideation is the positive valence system. Specifically, differences in brain areas involved in reward processing have been proposed as important for the development of suicide-related outcomes (Auerbach et al., 2021; Schmaal et al., 2019), including suicidal ideation (Albanese & Hajcak, 2021). One strong, indirect source of support for this hypothesis comes from a meta-analysis that found that anhedonia is positively associated with suicidal ideation, independent of depression (Ducasse et al., 2017). However, the application of this finding in the treatment of suicidal ideation in children and adolescents requires evidence demonstrating differences in the reward system are associated with suicidal ideation in children and adolescents. Consequently, doing so requires an understanding of the brain structures involved in the reward system and the reward system's development in late childhood and early adolescence.

The reward system and its development in children and adolescents

The mesolimbic dopamine pathway plays a key role in the reward system (Berridge & Robinson, 1998) and consists of structures such as the ventral tegmental area, nucleus accumbens, and the amygdala. The reward system also includes cortical areas, such as the anterior cingulate cortex and the orbitofrontal cortex, among others (Haber & Knutson, 2010). Though what exactly this circuit is signaling about reward is debated (particularly the dopamine signal in the ventral tegmental area and ventral striatum; see Berridge, 2012; Schultz, 2016), it is scientific consensus that this circuit does not primarily mediate hedonic "liking" or pleasure (Berridge & Robinson, 1998, 2003). Research has converged on the theory that, beginning in late childhood and into early adolescence, the reward system undergoes significant changes that lead to increases in reward sensitivity (Galvan, 2010). For

example, in rats, dopamine levels in the ventral striatum increase into adolescence, with levels dropping off in adulthood (Teicher et al., 1993). Dopamine receptor binding has also been shown to peak in adolescent rodents (Tarazi et al., 1998, 1999). Functional magnetic resonance imaging (fMRI) studies have also found that, compared to children (aged 7–12) and adults (aged 18–29), adolescents show hyperresponsive ventral striatal activity to receiving rewards (Galvan et al., 2006; Van Leijenhorst et al., 2010). There is also evidence from work using event-related potentials (ERPs) that reward sensitivity increases from late childhood into early adolescence.

Briefly, ERPs are direct measures of the neural response to specific events (e.g., onset of feedback) recorded using the electroencephalogram. The reward positivity (RewP) is an ERP that indexes the neural response to reward receipt, typically measured as the difference between monetary rewards and losses (Proudfit, 2015). The RewP is related to self-report and behavioral measures of reward processing (Bress & Hajcak, 2013; Pizzagalli et al., 2005) and is correlated with activation in reward-related brain regions, including the ventral striatum (Carlson et al., 2011). Recent work found that the RewP increases across early childhood into adolescence, with the youngest children seeing the greatest increases in the RewP at a two-year follow-up (Burani et al., 2019). Together, these findings illustrate that late childhood and early adolescence are characterized by changes in the reward system leading to increased reward sensitivity, particularly in the ventral striatum. However, do differences in the reward system during this developmental period characterize children and adolescents who experience suicidal ideation?

Reward system and suicidal thoughts and behaviors

A recent review claims that "blunted striatal activation characterize[s]" suicidal thoughts and behaviors (Auerbach et al., 2021). However, the evidence for this claim is relatively weak, particularly pertaining to whether this work has any bearing on suicidal

ideation. For example, one study used a Cyberball peer-interaction task and found that

adolescents with "high suicidal ideation" had blunted activity in the putamen, rather than the

ventral striatum, compared to adolescents with "low suicidal ideation" to social rewards

(Harms et al., 2019). Of note, adolescents in the "high suicidal ideation" group included those

who had previously attempted suicide, limiting the applicability of the study's findings to

suicidal ideation research. Theoretically and empirically, risk factors for suicidal ideation and

those for transitioning to suicidal behavior do not necessarily overlap (Joiner, 2005; May &

Klonsky, 2016; O'Connor, 2011; Van Orden et al., 2010). Thus, it is possible that the blunting

of the putamen was a result of adolescents who had previously attempted suicide, rather than

a result of suicidal ideation per se. Similarly, another study found blunted striatal activity

during a monetary incentive delay task in a group that had many adolescents who reported

suicidal ideation, compared to a control group; however, the inclusion criteria for this group

included engagement in non-suicidal self-injury (Sauder et al., 2016). While non-suicidal

self-injury and suicidal thoughts and behaviors are related, factor analytic work has

established that they are clearly distinguishable constructs with unique correlates (Evans &

Simms, 2019), further complicating the distinction of the blunter striatal activity as a

correlate of suicidal ideation as opposed to non-suicidal self-injury. Thus, these findings

cannot address whether children/adolescents experiencing suicidal ideation have a blunted

reward response compared to controls.

  Work leveraging the RewP is also unclear. For example, one study found an *increased*

RewP in depressed adolescents experiencing suicidal thoughts and behaviors compared to

depressed controls (Pegg et al., 2020). However, in this study, suicidal thoughts and behaviors

were conflated, again making it difficult to determine whether this difference in the RewP

was driven by suicidal thoughts, suicidal behaviors, or both. Perhaps the best test of whether

the reward system is blunted in children experiencing suicidal ideation was conducted by

Tsypes and colleagues (2019). In this study, the authors found that children with recent

suicidal ideation had a reduced RewP, compared to demographically and clinically matched

controls (Tsypes et al., 2019). Moreover, the authors found that this difference was driven by

an enhanced (i.e., more positive) neural response to losses. These results suggest that children

with suicidal ideation have blunted reward processing, compared to children without suicidal

ideation, implying that RewP differences may help identify children with suicidal ideation.

These results also point to neural responses to losses as a potential marker of suicidal ideation

among children.

Given the significance and implication of Tsypes and colleagues' findings to the

field's understanding of suicidal ideation in late childhood, we sought to conceptually

replicate and extend their findings. Specifically, we examined whether the RewP was blunted

in children and adolescents with recent suicidal ideation, compared to controls, using nearly

identical methods, measures, and processing procedures as those used by Tsypes and

colleagues (see Methods) in two large samples. Moreover, we sought to expand on Tsypes

and colleagues' (2019) findings by examining whether the RewP is not only related to

suicidal ideation cross-sectionally but also whether the RewP could predict suicidal ideation

two years later. In our first sample, consistent with Tsypes et al. (2019), we hypothesized that

children and adolescents with recent suicidal ideation would have a significantly smaller

RewP than those without recent suicidal ideation and that this effect would be driven by a

blunted neural response to loss trials. For our second sample, we hypothesized the following:

(1) Children and adolescents with recent suicidal ideation would have a significantly smaller

RewP than those without recent suicidal ideation cross-sectionally at both Time 1 and at Time

2; and (2) RewP at Time 1 would be negatively associated with suicidal ideation at Time 2,

two years later

Study 1: Conceptual Replication

Methods

Participants

Our first sample consisted of 275 children and adolescents. Participants ranged from 11.01 to 14.98 years of age (M = 12.91, SD = 1.14). Our sample consisted of slightly more males than females (54.9% male). In this sample, 72.36% identified as white, 10.18% as African American, 6.18% as Hispanic, 4.73% as Asian, and 6.55% identified their race/ethnicity as "other." Notably, our sample's racial and gender demographics were similar to Tsypes and colleagues' (2019) sample, but our sample was slightly older. Based on the present study's sample size and an alpha of .05, we were powered at .84 to detect an effect size of Hedges' g = 0.60 (calculation based on information in Tsypes et al. [2019]) for our difference in the RewP between those with recent suicidal ideation and controls. Participants and their parents provided informed consent and assent. The study was approved by the Institutional Review Broad at Florida State University.

Measures

Suicidal Ideation Measure. To assess for the presence of recent suicidal ideation (i.e., within the past two weeks), we used item 9[1] of the original Children's Depression Inventory (CDI; Kovacs, 1981) and the Schedule for School-Age Children—Present and Lifetime Version (K-SADS-PL; Kaufman et al., 1997). Individuals who endorsed a "1" (*I think about killing myself but I would not do it*) or "2" (*I want to kill myself*) on the CDI were grouped as having recent suicidal ideation. If individuals answered "1" or "2" on the CDI, then they were assessed for suicidal thoughts within the last two weeks on the K-SADS-PL interview, during which the children and their parents were asked, "Sometimes children who get upset or feel bad think about dying or even killing themselves. Have you/your child ever

---

[1]   Note that we used the original CDI. In the revised CDI, the suicide item is item 8.

had these types of thoughts?" All children who endorsed suicidal ideation on the CDI also

endorsed suicidal ideation during the K-SADS-PL. This is a small but important difference

between this sample and the study by Tsypes and colleagues (2019), who grouped children as

having recent suicidal ideation if they endorsed it on the CDI *or* the K-SADS-PL. Using our

approach, 27 (9.81%) participants endorsed having recent suicidal ideation in this sample.

For a breakdown of demographics between those with and without recent suicidal ideation,

see Table 1.

Doors Task. To measure neural responses to reward, we used the door task, which is

the same monetary guessing task used by Tsypes and colleagues (2019). The doors task is

frequently used in ERP studies of reward processing (Bress et al., 2012, 2013; Burani et al.,

2019; Klawohn et al., 2020). During this task, participants are presented with two doors and

instructed to guess which door has the monetary prize. To make their selection, the

participants use left and right mouse buttons corresponding with the door on the left and

right, respectively. After selecting a door, visual feedback is given consisting of either a green

arrow pointing upward, indicating that the participant won $0.50, or a red arrow pointing

downward, indicating that the participant lost $0.25. Feedback was presented for 2,000 ms;

then, the message "Click for next round" was presented. Our task consisted of 60 trials

presented in two blocks of 30 trials. For this task, 30 win and 30 loss trials were presented in

random order.

ERP Data Collection and Processing. During the doors task, electroencephalogram

(EEG) was recorded with an active electrode EEG-system (ActiCHamp, Brain Products

GmbH) with 32 electrodes positioned in accordance with the 10/20-system (ActiCAP, Brain

Products GmbH). Electrode Cz served as the online reference, a ground electrode was placed

on the forehead, and two electrodes were placed on the mastoids. Electrooculogram was

recorded from four additional electrodes: two 1 cm above and below the left eye and two at

the outer canthi of both eye. Continuous EEG waves were recorded at a sampling rate of

1000 Hz using a bandpass recording filter of 0.01–100 Hz. EEG data were processed using

Brain Vision Analyzer, Version 2.1 (Brain Products, Gilching, Germany). For some

participants ($n = 14$), at least one channel had to be interpolated using surrounding channels.

Then, for all participants, data were re-referenced to the average of the mastoid electrodes

and filtered from 0.01 to 30 Hz (Butterworth, 4th order). We then extracted 1000-ms

feedback-locked epochs, starting 200 ms before feedback presentation. Next, we corrected for

eye movement artifacts using the algorithm developed by Gratton, Coles, and Donchin

(1983). After this correction, segments that contained voltage steps > 50 mV between sample

points, a voltage difference of 175 mV within a 400-ms interval, or a maximum voltage

difference of < 0.5 mV within 100-ms intervals were automatically rejected.

Win and loss trials were averaged separately, and a baseline correction was applied

using the 200-ms preceding feedback onset. Then, in line with Tsypes and colleagues' (2019)

procedures, we exported these averages for temporospatial principal component analysis

(PCA), a factor analytic approach used to parse the ERP waveform into underlying

constituent components. PCA was conducted using the ERP PCA Toolkit (Version 2.82; Dien,

2010a). In line with evidence-based recommendations (Dien, 2010b), we first conducted a

temporal PCA using promax rotation. Covariance matrix and Kaiser normalization were used

for this PCA (Dien et al., 2005). Based on the resulting scree plot, 15 factors were extracted

in the temporal domain. We then analyzed the spatial distributions of these temporal factors

by conducting a spatial PCA using infomax rotation (Dien, 2010b). Based on the averaged

scree plot for all 15 temporal factors, we extracted four spatial factors, yielding 60 factor

combinations. Twenty factors each accounted for more than 0.50% of the variance and were

retained for further inspection (Kaiser, 1960). After examining the factors, TF03SF1 (i.e.,

temporal factor 3, spatial factor 1), which accounted for 11.20% of the variance, most closely

resembled the RewP: this factor was maximal at the Cz electrode site at 235 ms following

feedback onset and was more positive following win than loss trials.

Data Analytic Plan

All statistical analyses were conducted in R (Version 3.5.2; R Core Team, 2018),

using the following packages: "tidyverse" (Version 1.3.0; Wickham et al., 2019), "haven"

(Version 2.2.0; Wickham & Miller, 2019), "afex" (Version 0.23-0; Singmann et al., 2019),

"here" (Version 0.1; Müller, 2017), "DataExplorer" (Version 0.7.0; Cui, 2018), "emmeans"

(Version 1.3.2; Lenth, 2019), and "psych" (Version 1.8.4; Revelle, 2018). To test our

hypotheses, we conducted a 2x2 (condition: win, loss; ideation: no SI, recent SI) mixed-

measures ANOVA[2]. To rule out alternative explanations for our results, we reran this ANOVA

adding age as a covariate. We also reran this ANOVA adding household income as a

covariate. Though we view our relatively larger sample size as a strength of our study, similar

to Tsypes and colleagues' (2019) study, we also reran our 2x2 mixed-measures ANOVA after

selecting controls using a matching procedure. Specifically, we used a genetic matching

algorithm to select two controls for every case with recent suicidal ideation using the

"Matching" package (Version 4.9-7; Diamond & Sekhon, 2013; Sekhon, 2011). Similar to

Tsypes and colleagues (2019), we also matched by age, sex, race, household income, lifetime

history of major depressive disorder, lifetime history of any anxiety disorder, and current

symptoms of depression, anxiety, and externalizing disorders. Last, given the results of our

primary analyses and that traditional null hypothesis testing cannot provide confirmatory

evidence for the null hypothesis, we decided to conduct two equivalence tests using the

"TOSTER" package (Version 0.3.4; Lakens, 2017). These analyses require specifying an

effect size range that we deem to be uninteresting or "effectively zero." A significant

equivalence test provides evidence that the effect is smaller than the specified effect size

---

[2]   We also conducted these analyses using an area measure (250–350 ms) of the RewP at FCz, without using
          temporospatial PCA. The pattern of results remained the same.

range. We chose two effect size ranges, resulting in two separate equivalence tests: (1) lower

bound $d = -0.24$ and upper bound $d = 0.24$ and (2) lower bound $d = -0.60$ and upper bound d

$= 0.60$. Our first effect size estimate was based on recent meta-analytic evidence that the

upper bound on the relationship between ERPs and suicidal ideation is Hedges' $g = 0.24$, and

the second effect size range was based on the effect reported in the study we sought to

conceptually replicate (Gallyer et al., 2021; Tsypes et al., 2019). For our equivalence tests, we

used the PCA ΔRewP as our dependent variable and ideation group (no SI, recent SI) as the

independent variable.

Results

The means, standard deviations, and sample sizes of study variables across Study 1

and Study 2 are presented in Table 2. The mixed-measures ANOVA revealed an independent

effect of condition ($F[1, 262] = 45.61$, $p < .001$, $\eta^2 = .024$), such that wins ($M = 17.9$, 95% CI

[16.80, 19.00]) were greater than losses ($M = 13.80$, 95% CI [12.80, 14.9]). There was no

independent effect of ideation ($F[1, 262] = 1.66$, $p = .200$, $\eta^2 = .005$) or interaction between

ideation and condition ($F[1, 262] = 0.08$, $p = .773$, $\eta^2 < .001$; see Figure 1). Our ANCOVA

with age as a covariate revealed no statistically significant effects, including the interaction

between ideation and condition ($F[1, 260] = 0.28$, $p = .601$, $\eta^2 < .001$). Our analysis adding

household income as a covariate revealed a significant difference between wins and losses

($F[1, 232] = 6.87$, $p = .009$, $\eta^2 = .004$) but no other interactions or main effects. Our ANOVA

using our matched controls based on the genetic matching algorithm also did not find a main

effect of ideation ($F[1, 64] = 2.49$, $p = .120$, $\eta^2 = 0.033$) or an interaction between ideation

and condition ($F[1, 64] = 0.25$, $p = .617$, $\eta^2 < 0.001$). The result of our first equivalence test,

using effect size bounds of $d = -0.24$ and $d = 0.24$, was not significant ($t[34.47] = -0.93$, p

$= .179$). In contrast, our second equivalence test, using effect size bounds of $d = -0.60$ and $d =$

0.60, was significant ($t[34.47] = -2.82$, $p = .004$), suggesting that, if there is an effect, it is

smaller than that previously reported by Tsypes and colleagues (2019).

Study 2: Extension

Method

Participants

      Our second sample consisted of 325 child and adolescent females who ranged from 8.01 to 15.04 years of age (M = 12.42, SD = 1.80) at Time 1. At the two-year follow-up, the sample consisted of 235 participants (M = 14.41, SD = 1.82). Notably, there were no differences in race (χ^2[12] = 4.85, p = .963) between those who dropped out of the study and those who returned at Time 1 for the follow-up, but those who dropped out of the study were on average 0.71 years older than those who remained in the study (t[129.77] = 3.09, p = .013). In this sample, 81.54% identified their race as White/Caucasian, 5.85% as Black/African American, 0.31% as Native Hawaiian/Pacific Islander, 0.31% as American Indian/Alaskan Native, and 6.15% identified their race as "Other." As in Study 1, this racial breakdown is similar to the sample in Tsypes and colleagues (2019), and our sample was slightly older than the participants recruited by Tsypes and colleagues. Participants were recruited from the community through a commercial mailing list, using flyers, and using word-of-mouth. Based on the present study's sample size and an alpha of .05, we were powered at .80 to detect an effect size of Hedges' g = .60 (calculation based on information in Tsypes et al. [2019]) for a difference in the RewP between those with recent suicidal ideation and controls at Wave 1, and .80 at Wave 2. All participants and their parents provided informed consent and assent, as approved by the Institutional Review Board at Stony Brook University.

Measures

      Suicidal Ideation Measure. We used item 9 of the CDI and the K-SADS-PL interview to assess recent suicidal ideation, and participants who endorsed above "0" and/or

endorsed suicidal ideation during the K-SADS-PL were categorized as having recent suicidal

ideation. In contrast to Study 1, this coding is in line with Tsypes and colleagues' (2019)

approach. Based on this approach, 48 participants (14.8%) endorsed recent suicidal ideation

at Wave 1, and 33 participants (14.0%) endorsed recent suicidal ideation at Wave 2.

Doors Task. We used the same doors task that was used in Study 1.

ERP Data Collection and Processing. During the doors task, continuous EEG was

recorded using the ActiveTwo BioSemi system (BioSemi, Amsterdam, Netherlands) with a

cap containing 34 electrodes placed according to the 10/20 system (i.e., 32 channels plus FCz

and Iz). We also placed electrodes above and below the left eye, as well as near the outer

canthi of both eyes, to measure vertical and horizontal electrooculographic activity,

respectively. Two electrodes were also placed on the left and right mastoids. The EEG signal

was preamplified at the electrode, and data were digitized at a 24-bit resolution with a

sampling rate of 1024 Hz using a low-pass fifth-order sinc filter with a half-power cutoff of

204 Hz. Active electrodes were measured online with reference to a Common Mode Sense

active electrode constructing a monopolar channel.

Data were processed using BrainVision Analyzer, Version 2.1 (Brain Products,

Gliching, Germany). The data processing procedures were largely the same as in Study 1,

with small differences in artifact detection: Segments that contained voltage steps > 50 mV

between sample points, a voltage difference of 300 mV within a segment, or a maximum

voltage difference of < 0.5 mV within 100-ms intervals were automatically rejected. As in

Study 1, we conducted temporospatial PCA using ERP PCA Toolkit (Dien, 2010a). To do so,

we first conducted a temporal PCA with promax rotation. A covariance matrix and Kaiser

normalization were used for this PCA (Dien et al., 2005). Based on the resulting scree plot,

22 temporal factors were extracted in the temporal domain. We then analyzed the spatial

distributions of these temporal factors by conducting a spatial PCA using infomax rotation

(Dien, 2010b). Based on the averaged scree plot for all 22 temporal factors, we extracted 4

spatial factors, yielding 88 factor combinations. Twenty-four factors each accounted for more

than 0.50% of the variance and were retained for further inspection (Kaiser, 1960).

TF07SF1[3], which accounted for 0.8% of the variance, most closely resembled the RewP, as it

was maximal at the FCz electrode site at 346 ms and more positive for win trials than for loss

trials. Thus, factor scores from this PCA factor were used as the RewP measure for Wave 1.

We repeated this same process for Wave 2. Our temporal PCA resulted in 19 factors,

and our spatial PCA resulted in three factors, yielding 57 factor combinations together.

Eighteen factors each accounted for more than 0.50% of the variance and were retained for

further inspection (Kaiser, 1960). The TF06SF1 factor, which accounted for 1.5% of the

variance, most closely resembled the RewP, as it was maximal at the FCz electrode site at 278

ms and more positive for win trials than for loss trials.

Data Analytic Plan

We used the same statistical software and packages that were used in Study 1. To test

our hypotheses, we conducted the same analyses we used in Study 1 for Wave 1 and Wave 2

separately. One notable difference is that, for our genetic matching procedure, we did not

match based on gender, given that the entire sample in this study was female. To extend on

Tsypes and colleagues (2019) and other work (Pegg et al., 2020) that found a cross-sectional

relationship between the RewP and suicidality in children and adolescents, we also conducted

a series of logistic regression analyses. Specifically, we examined whether ΔRewP at Time 1

predicted ideation status at Time 2. Given work showing that childhood and adolescence are

associated with significant development in reward processing, we also reran this analysis

including age as a covariate (Burani et al., 2019).

---

[3]   Notably, another factor was also considered that explained more of the variance. We opted for this factor
        because: (1) The time window for the alternative factor was much earlier/outside of the traditional range of
        the RewP, and (2) the alternative factor did not display sufficient difference between wins and losses.
        Importantly, we also ran all of our analyses using the alternative factor; the results were the same: We found
        no differences in the RewP between the groups.

Results

Wave 1 Analyses

As expected, our 2x2 mixed-measures ANOVA revealed a main effect of condition (F[1, 309] = 95.56, p < .001, η^2 = .067), such that wins (M = 3.91, 95% CI [3.16, 4.66]) were greater than losses (M = -0.59, 95% CI [-1.34, 0.16]). In line with our Study 1 results, we did not find evidence of a main effect of ideation (F[1, 309] = 1.11, p = .293, η^2 = .003) or an interaction between ideation and condition (F[1, 309] = 1.11, p = .294, η^2 = .001). Our ANCOVA with age as a covariate revealed a main effect of age (F[1, 307] = 7.36, p = .007, η^2 = .018) but no other statistically significant effects, including the interaction between ideation and condition (F[1, 307] = 1.15, p = .285, η^2 = .001). Our ANCOVA adding income as a covariate revealed the expected main effect of condition (F[1, 262] = 9.06, p = .003, η^2 = .008), but no other effects were significant. Our analysis using our matched controls revealed a main effect of condition (F[1, 136] = 46.66, p < .001, η^2 = .076), but our main effect of ideation (F[1, 136] = 0.78, p = .378, η^2 = .004) and the interaction between ideation and condition (F[1, 136] = 0.67, p = .416, η^2 = .001) were not significant. The results of our first equivalence test, using effect size bounds of d = -0.24 and d = 0.24, were not significant (t[72.37] = -0.93, p = 0.259). In contrast, our second equivalence test, using effect size bounds of d = -0.60 and d = 0.60, was significant (t[72.37 = 2.87, p = .003), again suggesting that, if there is an effect, it is smaller than previously reported by Tsypes and colleagues (2019).

Wave 2 Analyses

Our 2x2 mixed-measures ANOVA revealed a main effect of condition (F[1, 253] = 45.59, p < .001, η^2 = .056) such that wins (M = 6.70, 95% CI [5.87, 7.54]) were greater than losses (M = 2.56, 95% CI [1.72, 3.39]). In line with our Study 1 and Study 2 Wave 1 results, we did not find evidence of a main effect of ideation (F[1, 253] = 0.11, p = .739, η^2 < .001) or

an interaction between ideation and condition (F[1, 253] = 0.11, p = .736, η^2 = .001). Our

ANCOVA with age as a covariate revealed no statistically significant effects, including the

interaction between ideation and condition (F[1, 251] < 0.01, p = .980, η^2 < .001). Our

ANCOVA adding income as a covariate revealed the expected main effect of condition (F[1,

228] = 9.06, p = .003, η^2 = .013), but no other effects were significant. Our analysis using our

matched controls revealed a main effect of condition (F[1, 91] = 20.79, p < .001, η^2 = .056),

but our main effect of ideation (F[1, 91] = 0.01, p = .927, η^2 < .001) and the interaction

between ideation and condition (F[1, 91] = 0.45, p = .505, η^2 = .001) were not significant.

The results of our first equivalence test, using effect size bounds of d = -0.24 and d = 0.24,

were not significant (t[48.73] = 1.19, p = 0.120). In contrast, our second equivalence test,

using effect size bounds of d = -0.60 and d = 0.60, was significant (t[48.73] = 3.14, p = .002),

again suggesting that, if there is an effect, it is smaller than previously reported by Tsypes and

colleagues (2019).

Wave 1 ΔRewP Predicting Wave 2 Ideation

Our first logistic regression predicting Wave 2 ideation status using Wave 1 ΔRewP as

the predictor was not significant (*OR* = 0.98, 95% CI [0.92, 1.04], p = 0.479). Our second

logistic regression adding Wave 1 age as a covariate showed that Wave 1 age was a

significant predictor of Wave 2 suicidal ideation status (*OR* = 1.62, 95% CI [1.26, 2.17], *p*

< .001), but Wave 1 ΔRewP was still not significant (*OR* = 0.96, 95% CI [0.90, 1.02], *p* =

0.209).

Discussion

The purpose of this study was to test whether children and adolescents with recent

suicidal ideation have blunted neural response to gains and/or losses cross-sectionally in a

large sample of males and females, and longitudinally in a sample of females. A previous

cross-sectional study found that children with suicidal ideation exhibit aberrant reward

processing, compared to matched controls, and this difference appeared to be driven by neural responses to loss trials, rather than to gain trials (Tsypes et al., 2019). However, in our conceptual replication using nearly identical methods, we did not find any evidence that children and adolescents with recent suicidal ideation had different neural reward responses than those without recent suicidal ideation. We attempted to rule out other explanations by conducting additional analyses, including controlling for age, controlling for household income, and using matched samples. Nevertheless, in not a single analysis were we able to find a main effect of suicidal ideation or an interaction between suicidal ideation group and wins/losses. To further investigate these null findings, we also conducted equivalence tests using two different cutoffs (viz., $d = .60$ and $d = .24$) and only found statistical equivalence when we used the larger cutoff. These results suggest that, *if* there is a difference in reward processing in children and adolescents within this age range, it is likely smaller than the effect originally reported by Tsypes and colleagues (2019).

Our results are in contrast to previous work that has found differences in striatal activation in adolescents with high suicidal ideation (Auerbach et al., 2021). However, aside from the fact that our study leveraged the RewP, rather than fMRI, there are some key methodological differences between our study and the broader literature that uses fMRI to study reward processing in suicidal thoughts and behaviors. First, many fMRI studies did not distinguish among suicidal ideation, suicidal behavior, and nonsuicidal self-injury. Moreover, many fMRI studies examine other types of rewards, such as social reward (e.g., Harms et al., 2019). Therefore, it is possible that these methodological differences explain the discrepancy between our results and the general fMRI literature. However, a recent, well-powered fMRI study ($N = 4,074$) failed to find any differences in any phase of reward processing (e.g., reward anticipation, reward receipt) in any brain region in children (9–10 years old) experiencing suicidal ideation, compared to controls (Vidal-Ribas et al., 2021). In fact, Vidal-

Ribas and colleagues (2021) also used equivalence testing and found that these effects were statistically equivalent to their smallest effect size of interest (i.e., *d* = .15). Thus, our results are in line with one of the most well-powered fMRI studies to date on reward processing in children with suicidal ideation.

Regarding the study by Tsypes and colleagues (2019) and why we failed to find the same results that they did, we believe there are at least three explanations. First, there are some subtle but key demographic differences between our samples and the sample reported by Tsypes and colleagues. Namely, the average age of both of our samples was about two-to-three years older, and the median household income in our samples was much higher ($85,000–$125,000, depending on sample and wave of data collection), compared to the sample collected by Tsypes and colleagues (~$20,000). The differences in age are especially key, as the reward system, and the RewP in particular, undergo substantial changes in late childhood and early adolescence, with reward sensitivity increasing over time (Burani et al., 2019; Galvan, 2010; Teicher et al., 1993). Thus, it is possible that reward processing in children with suicidal ideation only differs from that of children without suicidal ideation in late childhood, and that this effect diminishes as children progress into adolescence. Moreover, given that the median incomes in our samples were much higher than that in Tsypes and colleagues' sample, it is possible that environmental stressors such as poverty moderate the difference in reward processing between children and adolescents experiencing suicidal ideation. The potential for this type of effect is bolstered by research showing that poverty can influence the relationship between neural processing and other processes, such as inflammation (Miller et al., 2021).

A second explanation for the contrast in our findings is that differences in methodology may have contributed to the results, even despite our efforts to replicate those used by Tsypes and colleagues as closely as possible. Most notably, because Tsypes and

colleagues' study used PCA, it is difficult, if not impossible, to exactly replicate the weightings of time and space in the quantification of the RewP. Though we chose the PCA factors that most closely resembled the RewP in each of our samples, our Study 1 PCA factor and Study 2 Wave 1 factor peaked at slightly different times than is traditionally attributed to the RewP (i.e., 235 ms and 346 ms vs. 250–350 ms). Therefore, it is possible that factor selection influenced our results. Nonetheless, we found null results across Study 1 and Study 2, even when we used area measures (250–350 ms) without conducting PCA. Therefore, our results appear to be robust to other methodological choices. We also determined the groups slightly differently in Study 1: whereas Tsypes and colleagues (2019) determined children to have recent suicidal ideation if they endorsed suicidal ideation on the CDI *or* the K-SADS-PL, in Study 1 the suicidal ideation portion of the K-SADS-PL was only administered if participants first endorsed suicidal ideation on the CDI. In practice, this methodological difference means that it is possible we missed participants who would have endorsed suicidal ideation on the K-SADS-PL but not on the CDI; however, recent work among adults shows that this group is expected to be extremely small, given that increased anonymity is associated with a greater likelihood to endorse suicidal thoughts and behaviors (Deming et al., 2021). Given this evidence and the fact that we used the same approach to determine our recent suicidal ideation group as Tsypes and colleagues (2019) in our second sample and still did not find differences in the RewP between groups, it seems unlikely that this methodological difference drove our null results in our first sample.

The third potential explanation for the discrepancy between our findings and Tsypes and colleagues' findings is that our study is a false negative (i.e., there is a true effect, but we failed to find it) or that the study by Tsypes and colleagues was a false positive. Only considering our study and the study by Tsypes and colleagues, it is difficult to make a determination between these two scenarios. We recently conducted a meta-analysis on the

relationship between ERPs and suicidal ideation and found a very small (viz., Hedges' *g* =

-.06) blunting of the RewP in individuals experiencing suicidal ideation compared to controls

(Gallyer et al., 2021). For reasons we discuss in that meta-analytic review, including the small

effect size and the high correlation between depression and suicidal ideation (Rogers et al.,

2016), it cannot be ruled out that this blunting of the RewP is a false positive. In isolation,

this possibility would appear to suggest that there are no differences in reward processing

between those with versus without suicidal ideation. However, it is still possible that our

results, and the results of our meta-analysis, are due to insufficient sample sizes. Recent work

has shown that, across both fMRI and ERP studies, sample sizes tend to be too small to detect

realistic effect sizes in individual differences (Button et al., 2013; Clayson et al., 2019; Elliott

et al., 2020; Nord et al., 2017)[4].

What do our results mean about future work on reward processing in children and

adolescents experiencing suicidal ideation? First, we recommend better measurement

practices in the study of the neural correlates of suicidal thoughts and behaviors. As was used

in our study, it is common practice to use single-item measures of suicidal thoughts and

behaviors and to dichotomize variables into a group experiencing any suicidal thoughts

and/or behaviors and another group experiencing no suicidal thoughts and/or behaviors. The

use of a single-item assessment is particularly problematic given evidence that a non-

negligible number of participants who endorse suicidal thoughts/behaviors on a single item

do not meet standardized criteria for suicidal thoughts/behaviors if asked follow-up interview

questions (Hom et al., 2016; Millner et al., 2015). Future work should ideally use validated

multi-item interviews that allow for clarifying questions (e.g., Chu et al., 2015; Gallyer et al.,

2020; Gratch et al., 2021; Joiner et al., 1999; Nock et al., 2007).

Second, we recommend that researchers carefully consider the ideation-to-action

---

[4]   It is important to remember that low statistical power increases the probability of false negatives *and* false
       positives (see Button et al., 2013).

framework when studying suicidal thoughts and behaviors (Klonsky et al., 2018). A large

portion of the fMRI and ERP literature was not directly comparable to our own study because

many studies conflate suicidal thoughts, suicidal behaviors, and nonsuicidal self-injury (e.g.,

Harms et al., 2019; Pegg et al., 2020; Santamarina-Perez et al., 2019; Sauder et al., 2016).

Distinguishing among suicidal ideation, suicidal behaviors, and nonsuicidal self-injury will

aid in interpreting which neural correlates are, or are not, associated with different phases of

the ideation-to-action framework.

Third, researchers may consider examining fMRI and ERPs in the same participants

looking at different phases of the reward response. Such a study, if it were sufficiently

powered, would provide an excellent test of whether reward processing plays a role in

suicidal ideation in children and adolescents.

The current study has notable strengths. First, our study largely used the same

methods as the study we attempted to conceptually replicate, including the same task, the

same measure of recent suicidal ideation, and similar data processing and data analytic

procedures. The present study also used relatively large samples that are not typically seen in

ERP research, making our study well-powered to detect medium effects. Our study also used

two independent samples and extended on previous work by using an additional longitudinal

sample, with EEG and suicidal ideation measurements at both timepoints. As such, the

current study is the first to our knowledge to attempt to prospectively predict suicidal ideation

using an ERP—here, the RewP.

Despite these strengths, the current study also has some limitations. First, though our

study was well-powered to detect the effect size previously found by Tsypes and colleagues,

much larger samples are needed to detect smaller effects. Due to measurement error, it is

expected that ERPs will have small-to-moderate effects with behavioral and self-report

outcomes. Given the low base rate of suicidal ideation and other suicidal outcomes, large

samples are crucial for elucidating the relationship between the RewP and suicidal thoughts and behaviors. Second, our second sample consisted only of female participants. Though this potential demographic limitation is mitigated by our first sample, which has a fairly even split of males and females, this fact underscores that our longitudinal analyses may not apply to samples that include males. Thus, examing whether our longitudinal results generalize to other genders, as well as studying any differences in response to reward and its relation to future suicidal ideation across genders, is an important area for future research.

Conclusions

In summary, the present study attempted to conceptually replicate and extend previous findings that children with recent suicidal ideation had blunted neural reward processing, compared to controls. In two samples, we did not find evidence of this effect. We also did not find evidence that blunted reward processing was associated with future suicidal ideation. Our results highlight the need for larger sample sizes and multiple studies when investigating the relationship between neural reward processing and suicidal ideation. Our hope is that the present results will spur further research into the neural underpinnings of suicidal thoughts and behaviors and that greater collaboration and larger sample sizes will begin to dominate this field. Such approaches will greatly enhance our ability to contribute to the knowledge of the neurophysiology of suicidal thoughts and behaviors and will allow us to explore the potential clinical applications of event-related potentials to suicidal thoughts and behaviors.

References

Albanese, B. J., & Hajcak, G. (2021). Reward Processing Abnormalities and Promising New

Directions for Understanding Suicide Vulnerability. *Biological Psychiatry: Cognitive*

Neuroscience and Neuroimaging, *6*(1), 8–9.

https://doi.org/10.1016/j.bpsc.2020.11.002

Auerbach, R. P., Pagliaccio, D., Allison, G. O., Alqueza, K. L., & Alonso, M. F. (2021).

Neural Correlates Associated With Suicide and Nonsuicidal Self-injury in Youth.

Biological Psychiatry, *89*(2), 119–133.

https://doi.org/10.1016/j.biopsych.2020.06.002

Berridge, K. C. (2012). From prediction error to incentive salience: Mesolimbic computation

of reward motivation: From prediction error to incentive salience. *European Journal*

of Neuroscience, *35*(7), 1124–1143. https://doi.org/10.1111/j.1460-9568.2012.07990.x

Berridge, K. C., & Robinson, T. E. (1998). What is the role of dopamine in reward: Hedonic

impact, reward learning, or incentive salience? *Brain Research Reviews*, *28*(3), 309–

369. https://doi.org/10.1016/S0165-0173(98)00019-8

Berridge, K. C., & Robinson, T. E. (2003). Parsing reward. *Trends in Neurosciences*, *26*(9),

507–513. https://doi.org/10.1016/S0166-2236(03)00233-9

Bress, J. N., Foti, D., Kotov, R., Klein, D. N., & Hajcak, G. (2013). Blunted neural response

to rewards prospectively predicts depression in adolescent girls: Feedback negativity

predicts depression. *Psychophysiology*, *50*(1), 74–81. https://doi.org/10.1111/j.1469-

8986.2012.01485.x

Bress, J. N., & Hajcak, G. (2013). Self-report and behavioral measures of reward sensitivity

predict the feedback negativity: Feedback negativity, self-report, and behavior.

Psychophysiology, *50*(7), 610–616. https://doi.org/10.1111/psyp.12053

Bress, J. N., Smith, E., Foti, D., Klein, D. N., & Hajcak, G. (2012). Neural response to reward

and depressive symptoms in late childhood to early adolescence. *Biological*

Psychology, *89*(1), 156–162. https://doi.org/10.1016/j.biopsycho.2011.10.004

Burani, K., Mulligan, E. M., Klawohn, J., Luking, K. R., Nelson, B. D., & Hajcak, G. (2019).

Longitudinal increases in reward-related neural activity in early adolescence:

Evidence from event-related potentials (ERPs). *Developmental Cognitive*

Neuroscience, *36*, 100620. https://doi.org/10.1016/j.dcn.2019.100620

Button, K. S., Ioannidis, J. P. A., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S. J., &

Munafò, M. R. (2013). Power failure: Why small sample size undermines the

reliability of neuroscience. *Nature Reviews Neuroscience*, *14*.

https://doi.org/10.1038/nrn3475

Carlson, J. M., Foti, D., Mujica-Parodi, L. R., Harmon-Jones, E., & Hajcak, G. (2011).

Ventral striatal and medial prefrontal BOLD activation is correlated with reward-

related electrocortical activity: A combined ERP and fMRI study. *NeuroImage*, *57*(4),

1608–1616. https://doi.org/10.1016/j.neuroimage.2011.05.037

Centers for Disease Control. (2018). *WISQARS: Web-Based Injury Statistics Query and*

Reporting System [Government Database]. www.cdc.gov/injury/wisqars

Chu, C., Klein, K. M., Buchman-Schmitt, J. M., Hom, M. A., Hagan, C. R., & Joiner, T. E.

(2015). Routinized assessment of suicide risk in clinical practice: An empirically

informed update. *Journal of Clinical Psychology*, *71*(12), 1186–1200.

https://doi.org/10.1002/jclp.22210

Clayson, P. E., Carbine, K. A., Baldwin, S. A., & Larson, M. J. (2019). Methodological

reporting behavior, sample sizes, and statistical power in studies of event-related

potentials: Barriers to reproducibility and replicability. *Psychophysiology*.

https://doi.org/10.1111/psyp.13437

Cui, B. (2018). *DataExplorer: Data explorer*.

https://CRAN.R-project.org/package=DataExplorer

Diamond, A., & Sekhon, J. S. (2013). Genetic Matching for Estimating Causal Effects: A

General Multivariate Matching Method for Achieving Balance in Observational

Studies. *Review of Economics and Statistics*, *95*(3), 932–945.

Dien, J. (2010a). Evaluating two-step PCA of ERP data with Geomin, Infomax, Oblimin,

Promax, and Varimax rotations. *Psychophysiology*, *47*(1), 170–183.

https://doi.org/10.1111/j.1469-8986.2009.00885.x

Dien, J. (2010b). The ERP PCA Toolkit: An open source program for advanced statistical

analysis of event-related potential data. *Journal of Neuroscience Methods*, *187*(1),

138–145. https://doi.org/10.1016/j.jneumeth.2009.12.009

Dien, J., Beal, D. J., & Berg, P. (2005). Optimizing principal components analysis of event-

related potentials: Matrix type, factor loading weighting, extraction, and rotations.

Clinical Neurophysiology, *116*(8), 1808–1825.

https://doi.org/10.1016/j.clinph.2004.11.025

Ducasse, D., Loas, G., Dassa, D., Gramaglia, C., Zeppegno, P., Guillaume, S., Olié, E., &

Courtet, P. (2017). Anhedonia is associated with suicidal ideation independently of

depression: A meta-analysis. *Depression and Anxiety*, 1–11.

https://doi.org/10.1002/da.22709

Elliott, M. L., Knodt, A. R., Ireland, D., Morris, M. L., Poulton, R., Ramrakha, S., Sison, M.

L., Moffitt, T. E., Caspi, A., & Hariri, A. R. (2020). What Is the Test-Retest Reliability

of Common Task-Functional MRI Measures? New Empirical Evidence and a Meta-

Analysis. *Psychological Science*, *31*(7), 792–806.

https://doi.org/10.1177/0956797620916786

Evans, C. M., & Simms, L. J. (2019). The latent structure of self-harm. *Journal of Abnormal

Psychology*, *128*(1), 12–24. https://doi.org/10.1037/abn0000398

Fox, K. R., Huang, X., Guzmán, E. M., Funsch, K. M., Cha, C. B., Ribeiro, J. D., & Franklin,

J. C. (2020). Interventions for suicide and self-injury: A meta-analysis of randomized

controlled trials across nearly 50 years of research. *Psychological Bulletin*, *146*(12),

1117–1145. https://doi.org/10.1037/bul0000305

Franklin, J. C., Huang, X., Fox, K. R., & Ribeiro, J. D. (2018). What suicide interventions

should target. *Current Opinion in Psychology*, *22*, 50–53.

https://doi.org/10.1016/j.copsyc.2017.08.002

Franklin, J. C., Ribeiro, J. D., Fox, K. R., Bentley, K. H., Kleiman, E. M., Huang, X.,

Musacchio, K. M., Jaroszewski, A. C., Chang, B. P., & Nock, M. K. (2017). Risk

factors for suicidal thoughts and behaviors: A meta-analysis of 50 years of research. *Psychological Bulletin, 143*(2), 187–232. https://doi.org/10.1037/bul0000084

Gallyer, A. J., Chu, C., Klein, K. M., Quintana, J., Carlton, C., Dougherty, S. P., & Joiner, T. E. (2020). Routinized categorization of suicide risk into actionable strata: Establishing the validity of an existing suicide risk assessment framework in an outpatient sample. *Journal of Clinical Psychology*, jclp.22994. https://doi.org/10.1002/jclp.22994

Gallyer, A. J., Dougherty, S. P., Burani, K., Albanese, B. J., Joiner, T. E., & Hajcak, G. (2021). Suicidal thoughts, behaviors, and event-related potentials: A systematic review and meta-analysis. *Psychophysiology*, 20.

Galvan. (2010). Adolescent development of the reward system. *Frontiers in Human Neuroscience*. https://doi.org/10.3389/neuro.09.006.2010

Galvan, A., Hare, T. A., Parra, C. E., Penn, J., Voss, H., Glover, G., & Casey, B. J. (2006). Earlier Development of the Accumbens Relative to Orbitofrontal Cortex Might Underlie Risk-Taking Behavior in Adolescents. *Journal of Neuroscience, 26*(25), 6885–6892. https://doi.org/10.1523/JNEUROSCI.1062-06.2006

Glenn, C. R., Cha, C. B., Kleiman, E. M., & Nock, M. K. (2017). Understanding suicide risk within the research domain criteria (RDoC) framework: Insights, challenges, and future research considerations. *Clinical Psychological Science, 5*(3), 568–592.

Gratch, I., Fernandes, S. N., Bell, K.-A., Pollak, O. H., Fox, K. R., Tezanos, K., Ebo, T., & Cha, C. B. (2021). Self-Injurious Thoughts and Behaviors Interview–Revised (SITBI-R): Reliability, Validity, and Inter-Informant Agreement in an Adolescent Sample. *Journal of Clinical Child & Adolescent Psychology*, 1–11. https://doi.org/10.1080/15374416.2021.1901229

Gratton, G., Coles, M. G. H., & Donchin, E. (1983). A new method for off-line removal of ocular artifact. *Electroencephalography and Clinical Neurophysiology, 55*, 468–484.

Haber, S. N., & Knutson, B. (2010). The Reward Circuit: Linking Primate Anatomy and Human Imaging. *Neuropsychopharmacology, 35*(1), 4–26. https://doi.org/10.1038/npp.2009.129

Harms, M. B., Casement, M. D., Teoh, J. Y., Ruiz, S., Scott, H., Wedan, R., & Quevedo, K. (2019). Adolescent suicide attempts and ideation are linked to brain function during peer interactions. *Psychiatry Research: Neuroimaging*, *289*, 1–9. https://doi.org/10.1016/j.pscychresns.2019.05.001

Hom, M. A., Joiner, T. E., & Bernert, R. A. (2016). Limitations of a single-item assessment of suicide attempt history: Implications for standardized suicide risk assessment. *Psychological Assessment, 28*(8), 1026–1030. https://doi.org/10.1037/pas0000241

Insel, T., Cuthbert, B., Garvey, M., Heinssen, R., Pine, D. S., Quinn, K., Sanislow, C., & Wang, P. (2010). Research domain criteria (RDoC): Toward a new classification framework for research on mental disorders. *American Journal of Psychiatry, 167*(7), 748–751.

Ivey-Stephenson, A. Z., Demissie, Z., Crosby, A. E., Stone, D. M., Gaylor, E., Wilkins, N., Lowry, R., & Brown, M. (2020). Suicidal Ideation and Behaviors Among High School Students — Youth Risk Behavior Survey, United States, 2019. *MMWR Supplements, 69*(1), 47–55. https://doi.org/10.15585/mmwr.su6901a6

Jobes, D. A., & Joiner, T. E. (2019). Reflections on Suicidal Ideation. *Crisis, 40*(4), 227–230. https://doi.org/10.1027/0227-5910/a000615

Joiner, T. E. (2005). *Why people die by suicide*. Harvard University Press.

Joiner, T. E., Walker, R. L., Rudd, M. D., & Jobes, D. A. (1999). Scientizing and routinizing the assessment of suicidality in outpatient practice. *Professional Psychology: Research and Practice, 30*(5), 447–453.

Kaiser, H. F. (1960). The Application of Electronic Computers to Factor Analysis. *Educational and Psychological Measurement, 20*(1), 141–151. https://doi.org/10.1177/001316446002000116

Kaufman, J., Birmaher, B., Brent, D., Rao, U., Flynn, C., Moreci, P., Williamson, D., & Ryan, N. (1997). Schedule for affective disorders and schizophrenia for school-age children-present and lifetime version (K-SADS-PL): Initial reliability and validity data. *Journal of the American Academy of Child & Adolescent Psychiatry, 36*(7), 980–988.

Klawohn, J., Burani, K., Bruchnak, A., Santopetro, N., & Hajcak, G. (2020). Reduced neural

    response to reward and pleasant pictures independently relate to depression.

    Psychological Medicine, 1–9. https://doi.org/10.1017/S0033291719003659

Klonsky, E. D., Saffer, B. Y., & Bryan, C. J. (2018). Ideation-to-action theories of suicide: A

    conceptual and empirical update. *Current Opinion in Psychology*, *22*, 38–43.

    https://doi.org/10.1016/j.copsyc.2017.07.020

Lakens, D. (2017). Equivalence Tests: A Practical Primer for *t* Tests, Correlations, and Meta-

    Analyses. *Social Psychological and Personality Science*, *8*(4), 355–362.

    https://doi.org/10.1177/1948550617697177

Lenth, R. (2019). *emmeans: Estimated marginal means, aka least-squares means*.

    https://CRAN.R-project.org/package=emmeans

May, A. M., & Klonsky, E. D. (2016). What distinguishes suicide attempters from suicide

    Ideators? A meta-analysis of potential factors. *Clinical Psychology: Science and

    Practice*, *23*(1), 5–20. https://doi.org/10.1111/cpsp.12136

Miller, G. E., White, S. F., Chen, E., & Nusslock, R. (2021). Association of Inflammatory

    Activity With Larger Neural Responses to Threat and Reward Among Children Living

    in Poverty. *American Journal of Psychiatry*, *178*(4), 313–320.

    https://doi.org/10.1176/appi.ajp.2020.20050635

Millner, A. J., Lee, M. D., & Nock, M. K. (2015). Single-item measurement of suicidal

    behaviors: Validity and consequences of misclassification. *PLoS ONE*, *10*(10).

    http://journals.plos.org/plosone/article/file?id=10.1371/journal.pone.0141606&type=p

    rintable

Müller, K. (2017). *here: A simpler way to find your files*.

    https://CRAN.R-project.org/package=here

Nock, M. K., Holmberg, E. B., Photos, V. I., & Michel, B. D. (2007). Self-injurious thoughts

    and behaviors interview: Development, reliability and validity in an adolescent

    sample. *Psychological Assessment*, *19*(3), 309–317. https://doi.org/10.1037/1040-

    3590.19.3.309

Nord, C. L., Valton, V., Wood, J., & Roiser, J. P. (2017). Power-up: A Reanalysis of "Power Failure" in Neuroscience Using Mixture Modeling. *The Journal of Neuroscience*, *37*(34), 8051–8061. https://doi.org/10.1523/JNEUROSCI.3592-16.2017

O'Connor, R. C. (2011). Towards an integrated motivational-volitional model of suicidal behavior. In R. C. O'Connor, S. Platt, & J. Gordon (Eds.), *International Handbook of Suicide Prevention: Research, Policy and Practice* (pp. 181–198). John Wiley & Sons, Ltd. www.wiley.com/wiley-blackwell.

Pegg, S., Dickey, L., Green, H., & Kujawa, A. (2020). Differentiating clinically depressed adolescents with and without active suicidality: An examination of neurophysiological and self-report measures of reward responsiveness. *Depression and Anxiety*. https://doi.org/10.1002/da.23012

Pizzagalli, D. A., Jahn, A. L., & O'Shea, J. P. (2005). Toward an objective characterization of an anhedonic phenotype: A signal-detection approach. *Biological Psychiatry*, *57*(4), 319–327. https://doi.org/10.1016/j.biopsych.2004.11.026

Proudfit, G. H. (2015). The reward positivity: From basic research on reward to a biomarker for depression: The reward positivity. *Psychophysiology*, *52*(4), 449–459. https://doi.org/10.1111/psyp.12370

R Core Team. (2018). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. https://www.r-project.org/

Revelle, W. (2018). *psych: Procedures for psychological, psychometric, and personality research*. https://CRAN.R-project.org/package=psych

Rogers, M. L., Stanley, I. H., Hom, M. A., Chiurliza, B., Podlogar, M. C., & Joiner, T. E. (2016). Conceptual and empirical scrutiny of covarying depression out of suicidal ideation. *Assessment*. https://doi.org/10.1177/1073191116645907

Santamarina-Perez, P., Romero, S., Mendez, I., Leslie, S. M., Packer, M. M., Sugranyes, G., Picado, M., Font, E., Moreno, E., Martinez, E., Morer, A., Romero, M., & Singh, M. K. (2019). Fronto-Limbic Connectivity as a Predictor of Improvement in Nonsuicidal Self-Injury in Adolescents Following Psychotherapy. *Journal of Child and Adolescent*

Psychopharmacology, *29*(6), 456–465. https://doi.org/10.1089/cap.2018.0152

Sauder, C. L., Derbidge, C. M., & Beauchaine, T. P. (2016). Neural responses to monetary incentives among self-injuring adolescent girls. *Development and Psychopathology*, *28*(1), 277–291. https://doi.org/10.1017/S0954579415000449

Schmaal, L., van Harmelen, A. L., Chatzi, V., Lippard, E. T. C., Toenders, Y. J., Averill, L. A., Mazure, C. M., & Blumberg, H. P. (2019). Imaging suicidal thoughts and behaviors: A comprehensive review of 2 decades of neuroimaging studies. *Molecular Psychiatry*. https://doi.org/10.1038/s41380-019-0587-x

Schultz, W. (2016). Dopamine reward prediction-error signalling: A two-component response. *Nature Reviews Neuroscience*, *17*, 183–195.

Sekhon, J. S. (2011). Multivariate and Propensity Score Matching Software with Automated Balance Optimization: The Matching Package for R. *Journal of Statistical Software*, *42*(7), 1–52.

Singmann, H., Bolker, B., Westfall, J., & Aust, F. (2019). *afex: Analysis of factorial experiments*. https://CRAN.R-project.org/package=afex

Tarazi, F. I., Tomasini, E. C., & Baldessarini, R. J. (1998). Postnatal development of dopamine D4-like receptors in rat forebrain regions: Comparison with D2-like receptors. *Developmental Brain Research*, *110*(2), 227–233. https://doi.org/10.1016/S0165-3806(98)00111-4

Tarazi, F. I., Tomasini, E. C., & Baldessarini, R. J. (1999). Postnatal Development of Dopamine D_1-Like Receptors in Rat Cortical and Striatolimbic Brain Regions: An Autoradiographic Study. *Developmental Neuroscience*, *21*(1), 43–49. https://doi.org/10.1159/000017365

Teicher, M. H., Barber, N. I., Gelbard, H. A., Gallitano, A. L., Campbell, A., Marsh, E., & Baldessarini, R. J. (1993). Developmental Differences in Acute Nigrostriatal and Mesocorticolimbic System Response to Haloperidol. *Neuropsychopharmacology*, *9*(2), 147–156. https://doi.org/10.1038/npp.1993.53

Tsypes, A., Owens, M., & Gibb, B. E. (2019). Blunted Neural Reward Responsiveness in

Children With Recent Suicidal Ideation. *Clinical Psychological Science*, 216770261985634. https://doi.org/10.1177/2167702619856341

Van Leijenhorst, L., Zanolie, K., Van Meel, C. S., Westenberg, P. M., Rombouts, S. A. R. B., & Crone, E. A. (2010). What Motivates the Adolescent? Brain Regions Mediating Reward Sensitivity across Adolescence. *Cerebral Cortex, 20*(1), 61–69. https://doi.org/10.1093/cercor/bhp078

Van Orden, K. A., Witte, T. K., Cukrowicz, K. C., Braithwaite, S. R., Selby, E. A., & Joiner, T. E. (2010). The interpersonal theory of suicide. *Psychological Review, 117*(2), 575–600. https://doi.org/10.1037/a0018697

Vidal-Ribas, P., Janiri, D., Doucet, G. E., Pornpattananangkul, N., Nielson, D. M., Frangou, S., & Stringaris, A. (2021). Multimodal Neuroimaging of Suicidal Thoughts and Behaviors in a U.S. Population-Based Sample of School-Age Children. *American Journal of Psychiatry*, appi.ajp.2020.2. https://doi.org/10.1176/appi.ajp.2020.20020120

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open Source Software, 4*(43), 1686. https://doi.org/10.21105/joss.01686

Wickham, H., & Miller, E. (2019). *haven: Import and export "SPSS", "stata" and "SAS" files*. https://CRAN.R-project.org/package=haven

Author Contributions

Testing and data collection were performed by NS and KB. AJG performed the data analysis and interpretation under the supervision of TEJ and GH. AJG drafted the paper, and all other authors provided critical revisions. All authors approved the final version of the paper for submission.

Table 1

Demographics of Groups

	Study 1	
Variable	No SI (n = 237)	Recent SI (n = 27)
Age	M = 12.86, SD = 1.14	M = 13.27, SD = 1.17
Sex (% female)	44.30%	51.85%
Race (% White)	74.26%	59.26%
Household Income (Median)	$85,250	$80,500
	Study 2 Wave 1	
Variable	No SI (n = 263)	Recent SI (n = 48)
Age	M = 12.29, SD = 1.80	M = 13.09, SD = 1.60
Sex (% female)	100%	100%
Race (% White)	85.17%	81.25%
Household Income (Median)	$105,000	$127,500
	Study 2 Wave 2	
Variable	No SI (n = 222)	Recent SI (n = 33)
Age	M = 14.21, SD = 1.85	M = 15.51, SD = 1.20
Sex (% female)	100%	100%
Race (% White)	83.78%	78.79%
Household Income (Median)	$125,000	$90,000

Note. SI = suicidal ideation.

Table 2

Means and Standard Deviations of Study Variables

	Study 1	
	M(SD)	*M(SD)*
Variable	No SI (*n* = 27)	Recent SI (*n* = 237)
Wins	18.20(8.18)	16.09(7.91)
Losses	13.95(7.91)	12.18(7.44)
ΔRewP	4.26(6.03)	3.91(5.20)
	Study 2 Wave 1	
Variable	No SI (*n* = 263)	Recent SI (*n* = 48)
Wins	3.81(6.45)	3.41(5.12)
Losses	-0.21(6.03)	-1.58(5.40)
ΔRewP	4.02(5.99)	4.99(5.14)
	Study 2 Wave 2	
Variable	No SI (*n* = 222)	Recent SI (*n* = 33)
Wins	6.64(6.14)	6.55(4.52)
Losses	2.70(5.70)	2.20(4.43)
ΔRewP	3.94(6.75)	4.36(5.32)

Note. SI = suicidal ideation; wins, losses, and ΔRewP all refer to the reward positivity (RewP), an event-related potential that indexes neural responses to reward receipt.

Figure 1

Study 1 Stimulus-Locked Event-Related Potentials
[insert figure 1]
Note. Waveforms represent event-related potentials to wins and losses at Cz without conducting temporosptatial PCA. SI = suicidal ideation.

Figure 2

Study 1 ΔRewP of Children and Adolescents With and Without Recent SI

[insert figure 2]
Note. Red point represents mean ΔRewP in each group. RewP = reward positivity, an event-related potential that indexes neural responses to reward receipt; SI = suicidal ideation.

Figure 3

Study 2 ΔRewP of Children and Adolescents Across Wave 1 and Wave 2

[insert figure 3]

Note. Red points represent mean ΔRewP across groups. A = Wave 1; B = Wave 2. RewP = reward positivity, an event-related potential that indexes neural responses to reward receipt; SI = suicidal ideation.