Article Type: Letter

High-altitude adaptation in rhesus macaques

Zachary A. Szpiech[†]
Department of Biological Sciences
Auburn University
Auburn, AL 36842, USA
zas0020@auburn.edu

*Taylor E. Novak**
Department of Biological Sciences
Auburn University
Auburn, AL 36842, USA
tep0007@auburn.edu

*Nicholas P. Bailey**
Department of Biological Sciences
Auburn University
Auburn, AL 36842, USA
npb0015@auburn.edu

Laurie S. Stevison[†]
Department of Biological Sciences
Auburn University
Auburn, AL 36842, USA
lss0021@auburn.edu

[†]*Correspondence: zas0020@auburn.edu, lss0021@auburn.edu*
**Equal Contribution*

Running Title: **High-altitude adaptation in rhesus macaques**

Keywords: Adaptation, high-altitude, EGLN1, selection, macaque, macaca mulatta

Abstract Word Count (300 max): 216
Total Word Count (5000 max): 3813

Abstract

When natural populations split and migrate to different environments, they may experience different selection pressures that can lead to local adaptation. For aerobic life, the low atmospheric oxygen content of high altitude living presents a special challenge and a strong selection pressure. Searching for evidence of adaptation to high altitude, we compare the whole genomes of 23 wild rhesus macaques captured at high altitude (mean altitude > 4000m above sea level) to 22 wild rhesus macaques captured at low altitude (mean altitude < 500m above sea level). To capture the genomic patterns of a positive selective sweep, we develop XP-nSL, a haplotype-based genomic scan for differential local adaptation with power to detect ongoing and recently completed hard and soft sweeps. We find evidence of local adaptation in the high-altitude population at or near 303 known genes and several unannotated regions. We find the strongest signal for adaptation at EGLN1, a classic target for convergent evolution in several species living in low oxygen environments. Furthermore, many of the 303 genes are involved in processes related to hypoxia, regulation of ROS, DNA damage repair, synaptic signaling, and metabolism. These results suggest that, beyond adapting via a beneficial mutation in one single gene, adaptation to high altitude in rhesus macaques is polygenic and spread across numerous important biological systems.

Impact Summary

Extreme environments pose a challenge to life on multiple fronts. Very high-altitude environments are one such example, with low atmospheric oxygen, increased ultraviolet light exposure, harsh temperatures, and reduced nutrition availability. In spite

of these challenges, many plants and animals, including humans, have genetically

adapted to cope with these hardships. Here we study a population of rhesus macaques

living at high altitude and compare their genomic patterns with those of a population

living much closer to sea level, searching for evidence of genetic changes that are

indicative of adaptation to their environment.

When positive selection is ongoing or a beneficial mutation has recently fixed in a

population, genetic diversity is reduced in the vicinity of the adaptive allele, and we

expect to observe long homozygous haplotypes at high frequency. Here we develop a

statistic that summarizes these expected patterns and compares between two

populations in order to search for evidence of adaptation that may have occurred in one

but not the other. We implement this statistic in a popular and easy-to-use software

package.

We find evidence for adaptation at a critical gene that helps control physiological

response to low-oxygen, one that has been the target of repeated convergent evolution

across many species. We also find evidence for positive selection across a range of

traits, including metabolic and neurological. This work helps to explain the evolutionary

history of the rhesus macaque and furthers our understanding about the ways

organisms genetically adapt to high altitude environments.

Introduction

Selective sweeps produce regions of reduced genetic diversity in the vicinity of an

adaptive mutation. These patterns manifest as long extended regions of homozygous

haplotypes segregating at high frequency (Przeworski 2002; Sabeti *et al.* 2002; Kim and

Nielsen 2004; Garud *et al.* 2015). In the event of a *de novo* mutation that is adaptive in a

population, we expect the haplotype it resides on to rapidly rise in frequency in the

population (called a 'hard' sweep). On the other hand, if an ancestrally segregating

neutral or mildly deleterious allele turned out to be adaptive in a new environment, it

would likely reside on two or more haplotypes, which would rapidly rise in frequency in

the population (called a 'soft' sweep) (Hermisson and Pennings 2005; Pennings and

Hermisson 2006). As both of these processes happen on a time scale faster than

mutation or recombination can act to break up the sweeping haplotypes, we expect to

observe long and low diversity haplotypes at high frequency in the vicinity of an adaptive

mutation. However, if this mutation either does not exist or is not adaptive in a sister

population, we would not expect a sweep to occur and thus we would not expect to

observe similar haplotype patterns.

To capture these haplotype patterns and contrast them between a pair of

populations, we develop XP-nSL, a haplotype-based statistic with good power to detect

partial, fixed, and recently completed hard and soft sweeps by comparing a pair of

populations. XP-nSL is an extension nSL (Ferrer-Admetlla *et al.* 2014) and does not

require a genetic recombination map for computation. The lack of dependence on a

recombination map is important, as other statistics for identifying positive selection are

biased towards low-recombination regions (O'reilly *et al.* 2008), but the approach taken

by nSL has been shown to be more robust (Ferrer-Admetlla *et al.* 2014). Both nSL and

XP-nSL summarize haplotype diversity by computing the mean number of variant sites

in a region that are identical-by-state across all pairs of haplotypes. Whereas nSL

contrasts between haplotype pools carrying an ancestral or a derived allele in a single

population, XP-nSL contrasts between haplotype pools in two different populations,

allowing it to detect differential local adaptation.

An extreme example of adaptation to a local environment is the transition to high

altitude living. Organisms living at high altitude are confronted with many challenges,

including a low-oxygen atmosphere and increased ultraviolet light exposure, and these

harsh environments inevitably exert strong selection pressure. Indeed, adaptation to

high altitude living has been studied extensively across many organisms from plants,

including monocots (Gonzalo-Turpin and Hazard 2009; Ahmad et al. 2016) and dicots

(Kim and Donohue 2013; Liu et al. 2014; Munne-Bosch et al. 2016; Guo et al. 2018), to

numerous animals including amphibians (Yang et al. 2016), canids (Li et al. 2014; Wang

et al. 2014; Wang et al. 2020), humans (Bigham et al. 2009; Bigham et al. 2010; Xu et

al. 2010; Yi et al. 2010; Peng et al. 2011; Huerta-Sanchez et al. 2013; Huerta-Sanchez

et al. 2014; Jeong et al. 2014), yaks (Qiu et al. 2012), avians (Cai et al. 2013; Qu et al.

2013; Wang et al. 2015; Graham and Mccracken 2019), boars (Li et al. 2013), mice

(Storz et al. 2007; Cheviron et al. 2012; Schweizer et al. 2019; Storz et al. 2019; Velotta

et al. 2020), moles (Campbell et al. 2010), antelope (Ge et al. 2013), and horses

(Hendrickson 2013). Liu et al. (2018) recently sequenced and published the whole

genomes of 79 wild-born Chinese rhesus macaques collected from multiple sites in

China. Among these animals, 23 were sampled from far western Sichuan province in a

region with mean altitude > 4000 m above sea level (Liu et al. 2018), providing an

opportunity to study the genetics of local adaption to high altitude in rhesus macaques.

Rhesus macaques are the second most widely distributed primate, with a range

extending from Afghanistan to Vietnam and from a latitude of 15 to 38 degrees north

(Fooden 2000). Early ancestors of the macaque migrated out of Africa to the Eurasian

continent approximately 7 mya—the earliest catarrhine fossils on the continent are

macaque-like (Stewart and Disotell 1998). Modern rhesus macaques trace a recent

origin to Southeast Asia, with a major migratory split occurring approximately 162 kya

separating the ancestors of modern Indian and Chinese rhesus macaques (Hernandez

et al. 2007). Macaques have proven to be quite evolutionarily successful, demonstrating

ecological flexibility and adaptability via developmental plasticity and behavioral changes

(Richard *et al.* 1989; Madrid *et al.* 2018). Other studies have looked at how the rhesus

macaque radiation has led to population-level adaptation to climate and food availability

(Liu *et al.* 2018).

Here we test and evaluate our XP-nSL statistic and apply it to study the genomic

consequences of high altitude living in the rhesus macaque. We use it to compare the

haplotype patterns of the 23 animals from the high-altitude population with another 22

that Liu *et al.* (2018) sampled in lower-lying regions in eastern China with a mean

altitude < 500 m above sea level.

Methods

Data Preparation

Liu *et al.* (2018) generated whole genome sequencing data for 79 Chinese

macaques and called all biallelic polymorphic sites according to GATK best practices

using rheMac8, identifying 52,534,348 passing polymorphic autosomal sites. We then

filter all loci with > 10% missing data leaving 35,639,395 biallelic sites. Next, the

program SHAPEIT v4.1.2 (Delaneau *et al.* 2019) was used to phase haplotypes in the

full data set with a genetic map that was available for rheMac8 (Bcm-Hgsc 2020).

SHAPEIT performs imputation during phasing for any missing genotypes. Our analyses

here focus on 45 of the 79 samples from Liu *et al.* (2018), representing 23 from high-

altitude regions of China (*M. m. lasiotis*) and 22 from low-altitude regions of China (*M. m.*

littoralis), based on capture location information. See Table S1 for individual IDs used.

Liu *et al.* (2018) also inferred joint demographic histories for their five populations,

and we extract the demographic parameters for our two of interest. This demographic

history is recapitulated in Fig. 1 with detailed parameters given in Table 1, which are

then used for simulations to test XP-nSL.

A Statistic for Detecting Local Adaptation

We develop a cross-population haplotype-based statistic, XP-nSL, to scan for

regions of the genome implicated in differential local adaptation between two

populations by extending nSL (Ferrer-Admetlla *et al.* 2014), each of which is defined

below.

Consider the sets $A(k)$ and $D(k)$, representing the set of haplotypes at site k

carrying the ancestral or derived allele, respectively, and let $n_A(k) = |A(k)|$ and

$n_D(k) = |D(k)|$. Next $L_{ij}(k)$ is defined as the number of consecutive sites at which

haplotype i and j are identical-by-state (IBS) in the interval containing site k. Then nSL

at site k is $nS_L(k) = \log \frac{SL_A(k)}{SL_D(k)}$, where $SL_A(k) = \binom{n_A(k)}{2} \sum_{i<j \in A(k)} L_{ij}(k)$ and $SL_D(k) =$

$\binom{n_D(k)}{2} \sum_{i<j \in D(k)} L_{ij}(k)$. $SL_A(k)$ and $SL_D(k)$ represent the mean $L_{ij}(k)$ over all pairs of

haplotypes carrying either the ancestral or derived allele at locus k, respectively. nSL

scores are then normalized genome-wide in frequency bins either with respect to the

empirical background or neutral simulations with a matching demographic history. nSL is

implemented in `nsl` (Ferrer-Admetlla *et al.* 2014) and `selscan` v1.1.0+ (Szpiech and

Hernandez 2014).

XP-nSL is defined similarly, except instead of comparing pools of haplotypes

containing ancestral or derived alleles, it compares pools of haplotypes between two

different populations. Let $P_1(k)$ and $P_2(k)$, represent the set of haplotypes at site k in

population 1 and population 2, respectively, and let $n_{P_1}(k) = |P_1(k)|$ and $n_{P_2}(k) =$

$|P_2(k)|$. Then XP-nSL at site k is $XPnS_L(k) = \log\frac{SL_{P_1}(k)}{SL_{P_2}(k)}$, where $SL_{P_1}(k) =$

$\binom{n_{P_1}(k)}{2}\sum_{i<j\in P_1(k)} L_{ij}(k)$ and $SL_{P_2}(k) = \binom{n_{P_2}(k)}{2}\sum_{i<j\in P_2(k)} L_{ij}(k)$. $SL_{P_1}(k)$ and $SL_{P_2}(k)$

represent the mean $L_{ij}(k)$ over all pairs of haplotypes in population 1 or population 2 at

locus k, respectively. XP-nSL scores are then normalized genome-wide either with

respect to the empirical background or neutral simulations with a matching demographic

history. XP-nSL is implemented in `selscan` v1.3.0+ (Szpiech and Hernandez 2014). It

is worth noting that XP-nSL is analogous to XP-EHH (Sabeti *et al.* 2007) as nSL is

analogous to iHS (Voight *et al.* 2006).

The goal of these statistics is to capture a signal of extended regions of low

diversity on sweeping haplotypes (emblematic of an ongoing or recently completed

selective sweep) within a population (nSL) or on sweeping haplotypes in one population

versus another (XP-nSL). When XP-nSL scores are positive this suggests evidence for

a hard or soft sweep in population 1, and when XP-nSL scores are negative this

suggests evidence for a hard or soft sweep in population 2.

Simulations

In order to test the ability of XP-nSL to detect ongoing and recently completed

hard and soft sweeps, coalescent simulations were performed conditional on an allele

frequency trajectory with the program `discoal` (Kern and Schrider 2016). `discoal`

simulates an allele frequency trajectory for a single non-neutral allele backwards in time

and then simulates a neutral coalescent process conditional on this trajectory. This

takes advantage of the speed and efficiency of the coalescent while still being able

simulate genetic diversity patterns in the vicinity of a non-neutral locus.

All simulations were run with a two-population divergence demographic history

(Fig. 1 and Table 1), as inferred by Liu *et al.* (2018), and given by the following discoal

command line arguments `-p 2 46 44 -en 0 1 0.230410476572876 -en`

`0.071964666275442 1 0.278345739259351 -en 0.086558359329153 1`

`1.282885999320505 -ed 0.146214905642895 1 0 -en 0.146214905642895`

`0 4.089940389782871`. Here 46 haplotypes were sampled from population 0, which

has the demographic history of the high-altitude population, and 44 haplotypes were

sampled from population 1, which has the demographic history of the low-altitude

population. A mutation rate of $\mu = 2.5 \times 10^{-8}$ (Fan *et al.* 2018) was used, along with a

recombination rate of $r = 5.126 \times 10^{-9}$, which was computed as the genome-wide mean

rate from the rheMac8 recombination map (Bcm-Hgsc 2020). A 500 kb region was

simulated, thus giving a scaled mutation rate and scaled recombination parameters for

discoal as `-t 809.425 -r 165.967`. 5,349 replicates of neutral sequence was

simulated under this model, representing approximately the entire macaque genome

minus 500 kb.

228   For non-neutral simulations, sweep scenarios were simulated with a positive

229 additive selection coefficient $s \in \{0.01, 0.02, 0.05\}$ (corresponding to `discoal` flag `-a`

230 selected from $\{323.77, 647.54, 1618.85\}$). Soft sweeps are simulated as a mutation that

231 arose neutral and turned beneficial at a particular establishment frequency $e \in$

232 $\{0.01, 0.02, 0.03, 0.04, 0.05, 0.10, 0.20\}$ (`discoal` flag `-f`). Hard sweeps are simulated from

233 a *de novo* mutation that was never neutral, i.e. $e = 0$. Finally, sweeps were conditioned

234 on having either reached a certain frequency in the population at the time of sampling,

235 $f \in \{0.7, 0.8, 0.9, 1.0\}$ (`discoal` flag `-c`), or that the adaptive mutation reached fixation

236 some number of generations prior to sampling, $g \in \{50, 100, 200\}$ (`discoal` flag `-ws`

237 selected from $\{7.7215 \times 10^{-4}, 1.5443 \times 10^{-3}, 3.0886 \times 10^{-3}\}$). For the sake of being

238 conservative in our estimation of power (Voight *et al.* 2006; Sabeti *et al.* 2007), it was

239 assumed the actual adaptive mutation remains unsampled (`discoal` flag `-h`) even

240 though whole genome sequencing data are being analyzed. For each combination of

241 parameter values, 500 replicates were simulated.

242 **Detecting Local Adaptation in Real Data**

243   From the phased data set, animals captured at high altitude (n = 23) and animals

244 captured at low altitude (n = 22) were subset (see Table S1). Using `selscan` v1.3.0

245 (Szpiech and Hernandez 2014) to compute raw XP-nSL scores across the genome,

246 scores were then normalized using the genome-wide empirical background. The low-

247 altitude population was used as the reference population, so positive XP-nSL scores

248 correspond to long homozygous haplotypes and a possible sweep in the high-altitude

population compared to the low-altitude population, and vice versa for negative XP-nSL

scores.

In order to identify regions implicated as potentially adaptive, we search for

clusters of extreme scores along a chromosome. Using `selscan`'s companion program

`norm`, the genome is divided into non-overlapping 100kb regions and both the maximum

XP-nSL score and the fraction of XP-nSL scores > 2 are computed. `norm` then creates

10 quantile bins for windows and identifies the top 1% of windows with the highest

fraction of extreme scores. Each window is then annotated with the ensembl rheMac8

gene list, and a maximum XP-nSL score is assigned to a given gene based on the max-

score in the 100 kb window with which it overlaps. If a gene overlaps more than one 100

kb window, it is assigned the top max-score from among the windows.

Results

Power Analysis of XP-nSL

First, we evaluate the performance of XP-nSL based on simulations. After

computing XP-nSL for all sites in all simulations, scores were normalized by subtracting

the mean and dividing by standard deviation of the neutral simulations, giving the neutral

scores an approximately N(0,1) distribution (Fig. S1).

We consider the maximum score in a 100 kb interval as way to identify regions

under positive selection similar to (Voight *et al.* 2006). To get the null distribution of max-

scores, the maximum score is computed in the central 100 kb of all neutral simulations.

The distribution of max-scores in neutral simulations had a median of 2.093 with 95% of

the mass between 0.804 and 3.492, which is represented in Fig. 2 as a solid horizontal

black line (median) and two dashed horizontal black lines (95% interval). Next, the

maximum score is computed in the central 100 kb of all non-neutral simulations, and the

median and 95% intervals are plotted for each parameter combination. Fig. 2 shows

good separation between the neutral distribution of max-scores and the distribution of

max-scores for a range of non-neutral parameters, suggesting that our statistic can

distinguish between neutral and non-neutral scenarios. Note that soft sweeps that start

at 0.1 or 0.2 frequency see the least separation from the neutral distribution. To evaluate

the power of the max-score statistic, the 99[th] percentile of the max-score distribution is

computed in the neutral distribution ($neutral_{99} = 3.792$) and power is calculated as the

mass of each non-neutral max score distribution $> neutral_{99}$. This fixes the false

positive rate at 1%. Results are plotted in Fig. 3A, which shows good power to detect

both incomplete and completed sweeps. For soft sweeps that start at a frequency $\leq$

0.03, power is $> 75\%$ when the sweep is near or just past fixation; the ability to detect

soft sweeps falls off for sweeps that start > 0.03 frequency.

Next, we consider that due to linkage disequilibrium consecutive scores will be

correlated, and we should therefore expect clusters of extreme scores in true non-

neutral regions. We thus consider a window-based approach to identify selected regions

similar to (Voight *et al.* 2006). Taking the central 100 kb region of our simulations, the

fraction of XP-nSL scores > 2 is computed, representing approximately the highest 2%

of all neutral scores. Since each 100 kb window has different numbers of scores within

it, 10 quantile bins are created for windows with similar numbers of scores. Then the top

1% of windows in each bin of windows with the highest fraction of extreme scores are

taken as putatively selected regions. This approach also has the effect of ensuring the

false positive rate is at most 1%. The proportion of times the central 100 kb of each non-

neutral simulation is computed in order to get power. The results are plotted in Fig. 3B,

which shows improved power over the max-score approach across a wider range of

parameters. Indeed, using the window-based method, power to detect soft sweeps

improves substantially across the parameter space, with $> 75\%$ power to detect soft

sweeps at or near fixation that started at frequency ≤ 0.05.

Identifying Evidence for Adaptation to High Altitude in Rhesus Macaques

Next we analyze the pair of rhesus macaque populations using XP-nSL,

searching for evidence of local adaptation in the high-altitude population. Using the low-

altitude population as the reference population, normalized $XPnS_L > 0$ corresponds to

evidence for adaptation in the high-altitude population. Dividing the genome into 100 kb

windows, the maximum XP-nSL score of that region is assigned to each gene in it (see

Methods).

Using the per-gene max-scores, PANTHER (Mi *et al.* 2019) gene ontology

categories were tested for enrichment of high scores, where significance suggests an

enrichment of signals of positive selection among genes involved. Significant terms

related to regulation of ion transport and synaptic signaling (Table 2), each of which are

affected by hypoxic conditions (Karle *et al.* 2004; Corcoran and O'connor 2013).

From the genomic regions that were identified to contain a high proportion of

extreme positive scores (see Methods), 303 annotated genes were found. These

regions, their characteristics, and the genes contained therein are given in Table S2. A

PANTHER (Mi *et al.* 2019) gene ontology overrepresentation test indicates a 9.04-fold

enrichment of genes associated with monooxygenase activity ($FDR = 4.47 \times 10^{-2}$).

The monooxygenases in the selected regions include FMO2, FMO5, CYP2C8,

CYP2C9, CYP2C93, and ENSMMUT00000011129. These genes are important for the

metabolism of oxygen and the generation of reactive oxygen species (ROS) (Krueger

and Williams 2005). Under the physiological stress of a low-oxygen environment, ROS

levels increase and cause oxidative damage, and, in humans, long-term adaptation to

high altitudes includes adaptation to oxidative damage (Janocha *et al.* 2017). Indeed,

AOX1 is also identified in our top regions, mutations in which have been shown to affect

ROS levels in humans (Foti *et al.* 2017).

The genome-wide top ten 100 kb windows based on the percentage of extreme

XP-nSL scores are summarized in Table 3, and these windows overlap several genes,

including EGLN1. The EGLN1 locus is single strongest selection signal identified in the

entire genome (Fig. 4). This locus has the third highest cluster of extreme scores (Table

3), contains the highest XP-nSL score in the entire genome (chr1:207,698,003, $XPnS_L =$

6.54809), and contains six of the top ten genome wide XP-nSL scores (colored dark red

in Fig. 4). EGLN1 encodes Hypoxia-Inducible Factor Prolyl Hydroxylase 2 and acts as

an oxygen sensor by catalyzing the formation of HIF1α. The EGLN1 locus is a classic

target for adaptation to low-oxygen levels, having repeatedly been the target of

adaptation in numerous organisms living at high altitude around the world (Bigham *et al.*

2009; Bigham *et al.* 2010; Jeong *et al.* 2014; Graham and Mccracken 2019). In addition

to EGLN1, other genes related to lung function, oxygen use, and angiogenesis had

evidence for differential local adaptation between low- and high-altitude populations:

TRPM7, PAXIP1, RBPJ, and ENSMMUT00000040566 (Table S2). TRPM7 has been

found to be downregulated in a hypoxia-induced rat model and associated with

pulmonary hypertension (PAH) (Xing *et al.* 2019). PAXPIP1 may be involved in vascular

development and has also been associated with PAH (Jandl *et al.* 2019).

ENSMMUT00000040566 is a MAPK6 ortholog, which interacts with EGLN3 (Rodriguez

et al. 2016), and both it and RBPJ involved in angiogenesis (Ramasamy *et al.* 2014).

Due to the reduced oxygen levels at high altitudes, we expect genes involved in

metabolism and respiration may be under positive selection. Indeed, MDH1 encodes a

critical enzyme in the citrate cycle (Tanaka *et al.* 1996) and is found in the top ten

genome-wide regions (Table 3). A paralog of MDH1, MDH1B, has been previously

identified as a target of selection in humans living at high altitude (Yi *et al.* 2010).

ACADM and COX15 are also found in putatively adaptive regions (Table S2). Mutations

in and differing expression levels of ACADM are related to oxidative stress and

mitochondrial dysfunction in human disease (Xu *et al.* 2018). COX15 is involved in

oxidative phosphorylation (Alston *et al.* 2017), and cytochrome c oxidase (COX) genes

have previously been identified as under selection primates relative to other mammals

(Osada and Akashi 2012).

STXBP5L also appears in the top ten regions (Table 3) and is involved in

vesicular trafficking and neurotransmitter release (Kumar *et al.* 2015). As primate brains

use large amounts of oxygen and energy to function (Osada and Akashi 2012)

signatures of selection on neurological genes may be expected across populations living

at altitudes with differing oxygen levels. In addition to STXBP5L, several genes related

to neural development and synaptic formation (JAG2, TRPM7, DOCK7, NSG2, AUTS2)

were identified (Table S2). JAG2 is involved in the Notch signaling pathway that is

crucial for neuronal differentiation in the mammalian brain (Cardenas *et al.* 2018).

TRPM7, in addition to its association with PAH, plays a role in hypoxic neuronal cell

death (Aarts *et al.* 2003).

DNA damage, including double strand breaks and pyrimidine dimerization, can

manifest as a result of oxidative stress (Ye *et al.* 2016) or increased exposure to UV

radiation (Zhang *et al.* 2000; Greinert *et al.* 2012), both of which increase at high

altitudes. Two DNA damage repair genes are in our set of genes identified in positively

selected regions. The ring finger protein RNF138 is in our list of top ten genomic

windows (Table 3) and has been shown to promote DNA double-strand break repair

(Ismail *et al.* 2015). Furthermore, the DNA polymerase POLH also appears in a

putatively adaptive region (Table S2) and is known to be able to efficiently bypass

pyrimidine dimer lesions (Zhang *et al.* 2000).

Previous studies in rhesus macaques have shown there may be drastic

differences in diets between high-and low-altitude populations (Zhao 2018). Indeed,

genes related to lipid and fat metabolism (DOCK7, ST6GALNAC5, ANGPTL3, and

ACACA) were also found in putative adaptive regions. Across human populations, these

genes are all responsible for varying blood levels of fatty acids (Guo *et al.* 2016; Dewey

et al. 2017; Hebbar *et al.* 2018). Furthermore, ACACA has been shown to vary fatty acid

blood concentrations and be differentially expressed in highland versus lowland swine

populations (Shang *et al.* 2019).

Discussion

When populations split and migrate, they may adapt in different ways in response

When populations split and migrate, they may adapt in different ways in response

to their local environments. Genetic adaptations that arise and sweep through the

population leave a characteristic genomic pattern of long haplotypes of low diversity and

high frequency. We develop a two-population haplotype-based statistic, XP-nSL, to

capture these patterns. With good power to detect hard and soft sweeps that occur in

one population but not another, XP-nSL can identify positively selected regions of the

genome likely the result of local adaptation. We apply this statistic to genomes sampled

from a pair of wild-born populations of Chinese rhesus macaques, inferred to have

diverged approximately 9500 generations ago, one of which lives at high altitude in far

western Sichuan province, and the other that lives close to sea level.

Life at high altitude presents extreme biological challenges, including low

atmospheric oxygen, increased UV exposure, harsh winters, and reduced nutrition

availability, which create strong selection pressure. Organisms that survive and persist

are likely to be carrying genetic mutations that confer an advantage for living in such

harsh environments. Common targets for adaptation to such an environment include

genes related to hypoxia, regulation of ROS, DNA damage repair, and metabolism.

Indeed, in the high-altitude macaque population, we identify a strong signal of positive

selection at the EGLN1 locus (Fig. 4), a classic target for adaptation to low-oxygen

environments, in addition to 302 other genes, many of which are related to the myriad

environmental selection pressures expected in high altitude environments. These results

suggest that, rather than a single adaptive mutation at a single locus, adaptation to this

extreme environment by rhesus macaques is polygenic and acts through multiple

biological systems.

Acknowledgments

407 The authors would like to thank members of the Stevison Lab for helpful discussions

408 and Lawrence Uricchio for helpful comments on early versions of the manuscript. This

409 work was supported by research start-up funds from the Department of Biological

410 Sciences at Auburn University (LSS).

Author Contributions

412 ZAS and LSS conceived of the study. ZAS performed all simulations and genomic

413 analyses and implemented novel statistics. ZAS, TEN, and NPB characterized gene

414 functions and ontologies with contributions LSS. ZAS wrote the manuscript with

415 contributions from NPB, TEN, and LSS. All authors read and approved of the

416 manuscript.

Data Accessibility

418 Macaque whole genome VCFs are available at http://dx.doi.org/10.5524/100484. Selection scan

419 data available at Dryad XXXX [Upon publication].

Tables

Table 1. Demographic parameters used for simulations. T values are given in number of generations before present. N values represent diploid effective population size.

Parameter	Value
T_1	4660
T_2	5605
T_3	9468
N_1	16188
N_2	3730
N_3	4506
N_4	20768
N_5	66210

Table 2. Gene ontology enrichment analysis results based on maximum XP-nSL scores per gene. Significant GO terms are enriched for high XP-nSL scores.

PANTHER GO-Slim Biological Process	Gene Ontology ID	p-value	FDR
anterograde trans-synaptic signaling	GO:0098916	4.43×10^{-5}	1.52×10^{-2}
chemical synaptic transmission	GO:0007268	4.43×10^{-5}	1.83×10^{-2}
regulation of transport	GO:0051049	3.60×10^{-5}	1.85×10^{-2}
trans-synaptic signaling	GO:0099537	6.32×10^{-5}	1.86×10^{-2}
synaptic signaling	GO:0099536	1.03×10^{-4}	2.66×10^{-2}
ion transport	GO:0006811	1.33×10^{-4}	2.75×10^{-2}
transmembrane transport	GO:0055085	1.57×10^{-4}	2.94×10^{-2}
regulation of ion transport	GO:0043269	1.30×10^{-4}	2.97×10^{-2}
metal ion transport	GO:0030001	2.21×10^{-4}	3.04×10^{-2}
regulation of localization	GO:0032879	2.13×10^{-4}	3.14×10^{-2}
regulation of ion transmembrane transport	GO:0034765	2.66×10^{-4}	3.23×10^{-2}
inorganic ion transmembrane transport	GO:0098660	3.01×10^{-4}	3.45×10^{-2}
sodium ion transport	GO:0006814	3.36×10^{-4}	3.65×10^{-2}
ion transmembrane transport	GO:0034220	4.50×10^{-4}	4.42×10^{-2}
regulation of transmembrane transport	GO:0034762	4.40×10^{-4}	4.54×10^{-2}
cation transport	GO:0006812	4.87×10^{-4}	4.57×10^{-2}

444 **Table 3.** The top ten 100 kb genomic regions as ranked by percentage of scores greater
445 than 2. Concatenated region represents the genomic region merged with adjacent top
446 1% regions. Max score represents the max XP-nSL score in the concatenated region.
447 Genes gives all genes overlapping the concatenated region.
448

Genomic Region	% > 2	Flanking Regions	Concatenated Region	Max Score	Genes
*chr2:42300001-42400000	81.62%	3	chr2:42200001-42600000	5.42043	STXBP5L
chr3:99100001-99200000	78.18%	3	chr3:99100001-99500000	5.54282	-
chr1:207600001-207700000	*77.39%*	*1*	*chr1:207600001-207800000*	*6.54809†*	*EGLN1, TSNAX*
chr10:69700001-69800000	76.60%	4	chr10:69700001-70200000	4.89388	PITPNB, ENSMMUT00000079195.1, TTC28
chr13:64300001-64400000	74.01%	1	chr13:64300001-64500000	4.17728	WDPCP, MDH1, ENSMMUT00000070474.1
chr7:96400001-96500000	73.86%	1	chr7:96400001-96600000	4.58886	FAM177A1, PPP2R3C, ENSMMUT00000039911.2, ENSMMUT00000027928.3
chr10:70400001-70500000	70.24%	0	chr10:70400001-70500000	3.69192	TTC28
*chr2:42400001-42500000	65.86%	3	chr2:42200001-42600000	5.42043	STXBP5L
chr7:120400001-120500000	62.54%	0	chr7:120400001-120500000	5.86114	KIAA0586, ENSMMUT00000059161.1
chr18:20600001-20700000	62.42%	0	chr18:20600001-20700000	5.37386	RNF138

449 ***These regions are contained in the same concatenated region.**
450 **†Top genome-wide score. This region contains 6 of the top 10 genome-wide scores.**

451
452 **Table S1.** Population classification by individual ID.
453 *See TableS1.xlsx*
454
455 **Table S2.** List of genomic windows with the top 1% highest fraction of extreme XP-nSL
456 scores.
457 *See TableS2.xlsx*

468 **Figures**

High Altitude Low Altitude

472 **Figure 1.** A representation of the demographic history for our high- and low-altitude
473 populations as inferred by Liu *et al.* (2018).

Figure 2. The distribution of maximum XP-nSL scores from simulations across various parameters, represented by medians and intervals containing 95% of the mass of the distribution. Neutral simulations are represented by the black solid horizontal line (median) and the black dashed horizontal line (95% interval). Non-neutral simulations represented by a colored box (median) and error bars (95% interval). The parameters are e (frequency at which selection begins, e > 0 indicates soft sweep), f (frequency of selected mutation at sampling), g (number of generations since fixation), and s (selection coefficient).

Figure 3. Power curves for (A) the max-score approach and (B) the window-based approach to identifying sweeps. The parameters are e (frequency at which selection begins, e > 0 indicates soft sweep), f (frequency of selected mutation at sampling), g (number of generations since fixation), and s (selection coefficient).

Figure 4. XP-nSL scores in the vicinity of the EGLN1 locus. This locus contains the genome wide top score (star) and six of the top ten genome wide scores (dark red).

Normal Q–Q Plot

510
511 **Figure S1.** A normal quantile-quantile plot of neutral XP-nSL scores showing generally
512 good adherence to a standard normal distribution. Due to autocorrelation along the
513 genome, only every 1000th score is plotted.

References

Aarts, M., Iihara, K., Wei, W.L., Xiong, Z.G., Arundine, M., Cerwinski, W. *et al.* (2003). A key role for TRPM7 channels in anoxic neuronal death. *Cell* 115**:** 863-877.

Ahmad, K.S., Hameed, M., Fatima, S., Ashraf, M., Ahmad, F., Naseer, M. *et al.* (2016). Morpho-anatomical and physiological adaptations to high altitude in some Aveneae grasses from Neelum Valley, Western Himalayan Kashmir. *Acta Physiologiae Plantarum* 38**:** 93.

Alston, C.L., Rocha, M.C., Lax, N.Z., Turnbull, D.M. & Taylor, R.W. (2017). The genetics and pathology of mitochondrial disease. *J Pathol* 241**:** 236-250.

BCM-HGSC. (2020) Baylor College of Medicine Human Genome Sequencing Center Rhemac8 Recombination Map. Available at: [ftp://ftp.hgsc.bcm.edu/ucscHub/rhesusSNVs/rheMac8/all.rate.bw].

Bigham, A., Bauchet, M., Pinto, D., Mao, X., Akey, J.M., Mei, R. *et al.* (2010). Identifying signatures of natural selection in Tibetan and Andean populations using dense genome scan data. *PLoS Genet* 6**:** e1001116.

Bigham, A.W., Mao, X., Mei, R., Brutsaert, T., Wilson, M.J., Julian, C.G. *et al.* (2009). Identifying positive selection candidate loci for high-altitude adaptation in Andean populations. *Hum Genomics* 4**:** 79-90.

Cai, Q., Qian, X., Lang, Y., Luo, Y., Xu, J., Pan, S. *et al.* (2013). Genome sequence of ground tit Pseudopodoces humilis and its adaptation to high altitude. *Genome Biol* 14**:** R29.

Campbell, K.L., Storz, J.F., Signore, A.V., Moriyama, H., Catania, K.C., Payson, A.P. *et al.* (2010). Molecular basis of a novel adaptation to hypoxic-hypercapnia in a strictly fossorial mole. *BMC Evol Biol* 10**:** 214.

Cardenas, A., Villalba, A., de Juan Romero, C., Pico, E., Kyrousi, C., Tzika, A.C. *et al.* (2018). Evolution of Cortical Neurogenesis in Amniotes Controlled by Robo Signaling Levels. *Cell* 174**:** 590-606 e521.

Cheviron, Z.A., Bachman, G.C., Connaty, A.D., McClelland, G.B. & Storz, J.F. (2012). Regulatory changes contribute to the adaptive enhancement of thermogenic capacity in high-altitude deer mice. *Proceedings of the National Academy of Sciences* 109**:** 8635-8640.

Corcoran, A. & O'Connor, J.J. (2013). Hypoxia-inducible factor signalling mechanisms in the central nervous system. *Acta Physiol (Oxf)* 208**:** 298-310.

Delaneau, O., Zagury, J.F., Robinson, M.R., Marchini, J.L. & Dermitzakis, E.T. (2019). Accurate, scalable and integrative haplotype estimation. *Nat Commun* 10**:** 5436.

Dewey, F.E., Gusarova, V., Dunbar, R.L., O'Dushlaine, C., Schurmann, C., Gottesman, O. *et al.* (2017). Genetic and Pharmacologic Inactivation of ANGPTL3 and Cardiovascular Disease. *N Engl J Med* 377**:** 211-221.

Fan, Z., Zhou, A., Osada, N., Yu, J., Jiang, J., Li, P. *et al.* (2018). Ancient hybridization and admixture in macaques (genus Macaca) inferred from whole genome sequences. *Mol Phylogenet Evol* 127**:** 376-386.

Ferrer-Admetlla, A., Liang, M., Korneliussen, T. & Nielsen, R. (2014). On detecting incomplete soft or hard selective sweeps using haplotype structure. *Mol Biol Evol* 31**:** 1275-1291.

Fooden, J. (2000). *Systematic review of the rhesus macaque, Macaca mulatta (Zimmermann,*
*1780)*Field Museum of Natural History, Chicago, Ill. :.
Foti, A., Dorendorf, F. & Leimkuhler, S. (2017). A single nucleotide polymorphism causes
enhanced radical oxygen species production by human aldehyde oxidase. *PLoS One* 12**:**
e0182061.
Garud, N.R., Messer, P.W., Buzbas, E.O. & Petrov, D.A. (2015). Recent selective sweeps in North
American Drosophila melanogaster show signatures of soft sweeps. *PLoS Genet* 11**:**
e1005004.
Ge, R.-L., Cai, Q., Shen, Y.-Y., San, A., Ma, L., Zhang, Y. *et al.* (2013). Draft genome sequence of
the Tibetan antelope. *Nature Communications* 4**:** 1858.
Gonzalo-Turpin, H. & Hazard, L. (2009). Local adaptation occurs along altitudinal gradient
despite the existence of gene flow in the alpine plant species Festuca eskia. *Journal of*
Ecology 97**:** 742-751.
Graham, A.M. & McCracken, K.G. (2019). Convergent evolution on the hypoxia-inducible factor
(HIF) pathway genes EGLN1 and EPAS1 in high-altitude ducks. *Heredity (Edinb)* 122**:** 819-
832.
Greinert, R., Volkmer, B., Henning, S., Breitbart, E.W., Greulich, K.O., Cardoso, M.C. *et al.* (2012).
UVA-induced DNA double-strand breaks result from the repair of clustered oxidative
DNA damages. *Nucleic Acids Res* 40**:** 10263-10273.
Guo, T., Yin, R.X., Huang, F., Yao, L.M., Lin, W.X. & Pan, S.L. (2016). Association between the
DOCK7, PCSK9 and GALNT2 Gene Polymorphisms and Serum Lipid levels. *Sci Rep* 6**:**
19079.
Guo, X., Hu, Q., Hao, G., Wang, X., Zhang, D., Ma, T. *et al.* (2018). The genomes of two Eutrema
species provide insight into plant adaptation to high altitudes. *DNA Research* 25**:** 307-
315.
Hebbar, P., Nizam, R., Melhem, M., Alkayal, F., Elkum, N., John, S.E. *et al.* (2018). Genome-wide
association study identifies novel recessive genetic variants for high TGs in an Arab
population. *J Lipid Res* 59**:** 1951-1966.
Hendrickson, S.L. (2013). A genome wide study of genetic adaptation to high altitude in feral
Andean Horses of the paramo. *BMC Evol Biol* 13**:** 273.
Hermisson, J. & Pennings, P.S. (2005). Soft sweeps: molecular population genetics of adaptation
from standing genetic variation. *Genetics* 169**:** 2335-2352.
Hernandez, R.D., Hubisz, M.J., Wheeler, D.A., Smith, D.G., Ferguson, B., Rogers, J. *et al.* (2007).
Demographic histories and patterns of linkage disequilibrium in Chinese and Indian
rhesus macaques. *Science* 316**:** 240-243.
Huerta-Sanchez, E., Degiorgio, M., Pagani, L., Tarekegn, A., Ekong, R., Antao, T. *et al.* (2013).
Genetic signatures reveal high-altitude adaptation in a set of ethiopian populations. *Mol*
Biol Evol 30**:** 1877-1888.
Huerta-Sanchez, E., Jin, X., Asan, Bianba, Z., Peter, B.M., Vinckenbosch, N. *et al.* (2014). Altitude
adaptation in Tibetans caused by introgression of Denisovan-like DNA. *Nature* 512**:** 194-
197.

Ismail, I.H., Gagne, J.P., Genois, M.M., Strickfaden, H., McDonald, D., Xu, Z. *et al.* (2015). The
RNF138 E3 ligase displaces Ku to promote DNA end resection and regulate DNA repair
pathway choice. *Nat Cell Biol* 17**:** 1446-1457.
Jandl, K., Thekkekara Puthenparampil, H., Marsh, L.M., Hoffmann, J., Wilhelm, J., Veith, C. *et al.*
(2019). Long non-coding RNAs influence the transcriptome in pulmonary arterial
hypertension: the role of PAXIP1-AS1. *J Pathol* 247**:** 357-370.
Janocha, A.J., Comhair, S.A.A., Basnyat, B., Neupane, M., Gebremedhin, A., Khan, A. *et al.*
(2017). Antioxidant defense and oxidative damage vary widely among high-altitude
residents. *Am J Hum Biol* 29.
Jeong, C., Alkorta-Aranburu, G., Basnyat, B., Neupane, M., Witonsky, D.B., Pritchard, J.K. *et al.*
(2014). Admixture facilitates genetic adaptations to high altitude in Tibet. *Nat Commun*
5**:** 3281.
Karle, C., Gehrig, T., Wodopia, R., Hoschele, S., Kreye, V.A., Katus, H.A. *et al.* (2004). Hypoxia-
induced inhibition of whole cell membrane currents and ion transport of A549 cells. *Am J*
Physiol Lung Cell Mol Physiol 286**:** L1154-1160.
Kern, A.D. & Schrider, D.R. (2016). Discoal: flexible coalescent simulations with selection.
Bioinformatics 32**:** 3839-3841.
Kim, E. & Donohue, K. (2013). Local adaptation and plasticity of Erysimum capitatum to altitude:
its implications for responses to climate change. *Journal of Ecology* 101**:** 796-805.
Kim, Y. & Nielsen, R. (2004). Linkage Disequilibrium as a Signature of Selective Sweeps. *Genetics*
167**:** 1513-1524.
Krueger, S.K. & Williams, D.E. (2005). Mammalian flavin-containing monooxygenases:
structure/function, genetic polymorphisms and role in drug metabolism. *Pharmacol Ther*
106**:** 357-387.
Kumar, R., Corbett, M.A., Smith, N.J., Jolly, L.A., Tan, C., Keating, D.J. *et al.* (2015). Homozygous
mutation of STXBP5L explains an autosomal recessive infantile-onset neurodegenerative
disorder. *Hum Mol Genet* 24**:** 2000-2010.
Li, M., Tian, S., Jin, L., Zhou, G., Li, Y., Zhang, Y. *et al.* (2013). Genomic analyses identify distinct
patterns of selection in domesticated pigs and Tibetan wild boars. *Nat Genet* 45**:** 1431-
1438.
Li, Y., Wu, D.D., Boyko, A.R., Wang, G.D., Wu, S.F., Irwin, D.M. *et al.* (2014). Population variation
revealed high-altitude adaptation of Tibetan mastiffs. *Mol Biol Evol* 31**:** 1200-1205.
Liu, J.-Q., Duan, Y.-W., Hao, G., Ge, X.-J. & Sun, H. (2014). Evolutionary history and underlying
adaptation of alpine plants on the Qinghai–Tibet Plateau. *Journal of Systematics and*
Evolution 52**:** 241-249.
Liu, Z., Tan, X., Orozco-terWengel, P., Zhou, X., Zhang, L., Tian, S. *et al.* (2018). Population
genomics of wild Chinese rhesus macaques reveals a dynamic demographic history and
local adaptation, with implications for biomedical research. *Gigascience* 7.
Madrid, J.E., Mandalaywala, T.M., Coyne, S.P., Ahloy-Dallaire, J., Garner, J.P., Barr, C.S. *et al.*
(2018). Adaptive developmental plasticity in rhesus macaques: the serotonin transporter
gene interacts with maternal care to affect juvenile social behaviour. *Proc Biol Sci* 285.

Mi, H., Muruganujan, A., Ebert, D., Huang, X. & Thomas, P.D. (2019). PANTHER version 14: more
genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools.
Nucleic Acids Res 47**:** D419-D426.
Munne-Bosch, S., Cotado, A., Morales, M., Fleta-Soriano, E., Villellas, J. & Garcia, M.B. (2016).
Adaptation of the Long-Lived Monocarpic Perennial Saxifraga longifolia to High Altitude.
Plant Physiol 172**:** 765-775.
O'Reilly, P.F., Birney, E. & Balding, D.J. (2008). Confounding between recombination and
selection, and the Ped/Pop method for detecting selection. *Genome Res* 18**:** 1304-1313.
Osada, N. & Akashi, H. (2012). Mitochondrial-nuclear interactions and accelerated
compensatory evolution: evidence from the primate cytochrome C oxidase complex. *Mol*
Biol Evol 29**:** 337-346.
Peng, Y., Yang, Z., Zhang, H., Cui, C., Qi, X., Luo, X. *et al.* (2011). Genetic variations in Tibetan
populations and high-altitude adaptation at the Himalayas. *Mol Biol Evol* 28**:** 1075-1081.
Pennings, P.S. & Hermisson, J. (2006). Soft sweeps II--molecular population genetics of
adaptation from recurrent mutation or migration. *Mol Biol Evol* 23**:** 1076-1084.
Przeworski, M. (2002). The Signature of Positive Selection at Randomly Chosen Loci. *Genetics*
160**:** 1179-1189.
Qiu, Q., Zhang, G., Ma, T., Qian, W., Wang, J., Ye, Z. *et al.* (2012). The yak genome and
adaptation to life at high altitude. *Nat Genet* 44**:** 946-949.
Qu, Y., Zhao, H., Han, N., Zhou, G., Song, G., Gao, B. *et al.* (2013). Ground tit genome reveals
avian adaptation to living at high altitudes in the Tibetan plateau. *Nat Commun* 4**:** 2071.
Ramasamy, S.K., Kusumbe, A.P., Wang, L. & Adams, R.H. (2014). Endothelial Notch activity
promotes angiogenesis and osteogenesis in bone. *Nature* 507**:** 376-380.
Richard, A.F., Goldstein, S.J. & Dewar, R.E. (1989). Weed macaques: The evolutionary
implications of macaque feeding ecology. *International Journal of Primatology* 10**:** 569.
Rodriguez, J., Pilkington, R., Garcia Munoz, A., Nguyen, L.K., Rauch, N., Kennedy, S. *et al.* (2016).
Substrate-Trapped Interactors of PHD3 and FIH Cluster in Distinct Signaling Pathways.
Cell Rep 14**:** 2745-2760.
Sabeti, P.C., Reich, D.E., Higgins, J.M., Levine, H.Z., Richter, D.J., Schaffner, S.F. *et al.* (2002).
Detecting recent positive selection in the human genome from haplotype structure.
Nature 419**:** 832-837.
Sabeti, P.C., Varilly, P., Fry, B., Lohmueller, J., Hostetter, E., Cotsapas, C. *et al.* (2007). Genome-
wide detection and characterization of positive selection in human populations. *Nature*
449**:** 913-918.
Schweizer, R.M., Velotta, J.P., Ivy, C.M., Jones, M.R., Muir, S.M., Bradburd, G.S. *et al.* (2019).
Physiological and genomic evidence that selection on the transcription factor Epas1 has
altered cardiovascular function in high-altitude deer mice. *PLoS Genet* 15**:** e1008420.
Shang, P., Li, W., Liu, G., Zhang, J., Li, M., Wu, L. *et al.* (2019). Identification of lncRNAs and
Genes Responsible for Fatness and Fatty Acid Composition Traits between the Tibetan
and Yorkshire Pigs. *Int J Genomics* 2019**:** 5070975.
Stewart, C.B. & Disotell, T.R. (1998). Primate evolution - in and out of Africa. *Curr Biol* 8**:** R582-
588.

Storz, J.F., Cheviron, Z.A., McClelland, G.B. & Scott, G.R. (2019). Evolution of physiological
performance capacities and environmental adaptation: insights from high-elevation deer
mice (Peromyscus maniculatus). *J Mammal* 100**:** 910-922.
Storz, J.F., Sabatino, S.J., Hoffmann, F.G., Gering, E.J., Moriyama, H., Ferrand, N. *et al.* (2007).
The molecular basis of high-altitude adaptation in deer mice. *PLoS Genet* 3**:** e45.
Szpiech, Z.A. & Hernandez, R.D. (2014). selscan: an efficient multithreaded program to perform
EHH-based scans for positive selection. *Mol Biol Evol* 31**:** 2824-2827.
Tanaka, T., Inazawa, J. & Nakamura, Y. (1996). Molecular cloning and mapping of a human cDNA
for cytosolic malate dehydrogenase (MDH1). *Genomics* 32**:** 128-130.
Velotta, J.P., Robertson, C.E., Schweizer, R.M., McClelland, G.B. & Cheviron, Z.A. (2020).
Adaptive shifts in gene regulation underlie a developmental delay in thermogenesis in
high-altitude deer mice. *Molecular Biology and Evolution*.
Voight, B.F., Kudaravalli, S., Wen, X. & Pritchard, J.K. (2006). A map of recent positive selection
in the human genome. *Plos Biol* 4**:** e72.
Wang, G.D., Fan, R.X., Zhai, W., Liu, F., Wang, L., Zhong, L. *et al.* (2014). Genetic convergence in
the adaptation of dogs and humans to the high-altitude environment of the tibetan
plateau. *Genome Biol Evol* 6**:** 2122-2128.
Wang, M.-S., Li, Y., Peng, M.-S., Zhong, L., Wang, Z.-J., Li, Q.-Y. *et al.* (2015). Genomic Analyses
Reveal Potential Independent Adaptation to High Altitude in Tibetan Chickens. *Molecular*
Biology and Evolution 32**:** 1880-1889.
Wang, M.-S., Wang, S., Li, Y., Jhala, Y., Thakur, M., Otecko, N.O. *et al.* (2020). Ancient
hybridization with an unknown population facilitated high altitude adaptation of canids.
Molecular Biology and Evolution.
Xing, J., Wang, M., Hong, J., Gao, Y., Liu, Y., Gu, H. *et al.* (2019). TRPM7 channel inhibition
exacerbates pulmonary arterial hypertension through MEK/ERK pathway. *Aging (Albany*
NY) 11**:** 4050-4065.
Xu, S., Li, S., Yang, Y., Tan, J., Lou, H., Jin, W. *et al.* (2010). A Genome-Wide Search for Signals of
High-Altitude Adaptation in Tibetans. *Molecular Biology and Evolution* 28**:** 1003-1011.
Xu, Z., Jin, X., Cai, W., Zhou, M., Shao, P., Yang, Z. *et al.* (2018). Proteomics Analysis Reveals
Abnormal Electron Transport and Excessive Oxidative Stress Cause Mitochondrial
Dysfunction in Placental Tissues of Early-Onset Preeclampsia. *Proteomics Clin Appl* 12**:**
e1700165.
Yang, X., Wang, Y., Zhang, Y., Lee, W.H. & Zhang, Y. (2016). Rich diversity and potency of skin
antioxidant peptides revealed a novel molecular basis for high-altitude adaptation of
amphibians. *Sci Rep* 6**:** 19866.
Ye, B., Hou, N., Xiao, L., Xu, Y., Xu, H. & Li, F. (2016). Dynamic monitoring of oxidative DNA
double-strand break and repair in cardiomyocytes. *Cardiovasc Pathol* 25**:** 93-100.
Yi, X., Liang, Y., Huerta-Sanchez, E., Jin, X., Cuo, Z.X., Pool, J.E. *et al.* (2010). Sequencing of 50
human exomes reveals adaptation to high altitude. *Science* 329**:** 75-78.
Zhang, Y., Yuan, F., Wu, X., Rechkoblit, O., Taylor, J.S., Geacintov, N.E. *et al.* (2000). Error-prone
lesion bypass by human DNA polymerase eta. *Nucleic Acids Res* 28**:** 4717-4724.
728