Transient intestinal colonization by a live-attenuated oral cholera vaccine induces

protective immune responses in streptomycin-treated mice

Bolutife Fakoya,[a,b,] Brandon Sit,[a,b] Matthew K. Waldor[a,b,c,#]

[a]Division of Infectious Diseases, Brigham & Women's Hospital, Boston, USA

[b]Department of Microbiology, Harvard Medical School, Boston, USA

[c]Howard Hughes Medical Institute, Bethesda, USA

[#]Address correspondence to Matthew K Waldor, mwaldor@research.bwh.harvard.edu

Abstract

Current mouse models for evaluating the efficacy of live oral cholera vaccines

(OCVs) have important limitations. Conventionally raised adult mice are resistant to

intestinal colonization by *Vibrio cholerae*, but germ-free mice can be colonized and have

been used to study OCV immunogenicity. However, germ free animals have impaired

immune systems and intestinal physiology; also, live OCVs colonize germ free mice for

many months, which does not mimic the clearance kinetics of live OCVs in humans.

Here, we leverage antibiotic-treated, conventionally raised adult mice to study the

effects of transient intestinal colonization by a live OCV *V. cholerae* strain. In a single

dose vaccination regimen, we found that HaitiV, a live-attenuated OCV candidate, was

cleared by streptomycin treated adult mice within a week after oral inoculation. This

transient colonization elicited far stronger adaptive immune correlates of protection

against cholera than did inactivated whole-cell HaitiV. Infant mice from HaitiV

vaccinated dams were also significantly protected from choleric disease than pups from

inactivated-HaitiV dams. Our findings establish the benefits of antibiotic treated mice for

live OCV studies as well as its limitations and underscore the immunogenicity of HaitiV.

Importance

Oral cholera vaccines (OCVs) are being deployed to combat cholera but current

killed OCVs require multiple doses and show little efficacy in young children. Live OCVs

have the potential to overcome these limitations but small animal models for testing

OCVs have shortcomings. We used an antibiotic treatment protocol for conventional

adult mice to study the effects of short-term colonization by a single dose of HaitiV, a

live OCV candidate. Vaccinated mice developed vibriocidal antibodies against *V.*

cholerae and delivered pups that were resistant to cholera, whereas mice vaccinated

with inactivated HaitiV did not. These findings demonstrate HaitiV's immunogenicity and

suggest that this antibiotic treatment protocol will be useful for evaluating the efficacy of

live OCVs.

Introduction

Vibrio cholerae is the cause of cholera and, following ingestion of water or food contaminated with this Gram-negative rod, humans can develop the severe and sometimes lethal dehydrating diarrhea that characterizes cholera. Cholera remains a major threat to global public health, with approximately 2.9 million cases and 95,000 deaths reported annually (1). Serologic classification of V. cholerae is based on the structure and chemistry of the abundant LPS O-antigen and the O1 serogroup of V. cholerae has given rise to all cholera pandemics. The O1 serogroup is further subdivided into Ogawa and Inaba serotypes that differ by the presence or absence of methylation of the terminal perosamine on their respective O-antigens (2). Current pandemic cholera is predominantly caused by an O1 'variant' El Tor biotype strain, such as the strain responsible for the Haitian epidemic in 2010 (3).

Prior exposure to V. cholerae can elicit long-lived protective O-antigen-specific responses against V. cholerae (4, 5), suggesting that vaccination has the potential to elicit protective immunity if vaccines can safely mimic elements of natural infection. As such, several vaccination strategies for cholera have been developed over the decades. Among these, oral cholera vaccines (OCV) are attractive options as they stimulate immunity at the intestinal mucosal surface, the site of infection, and because of their ease of administration. Killed whole-cell OCVs, such as Shancol, are being increasingly adopted as frontline public health tools both in endemic regions (6) as well as to limit outbreaks (7).Live-attenuated OCVs have also been developed and have theoretical advantages over killed OCVs including in vivo replication which enables continuous presentation of in vivo-induced antigens at the intestinal mucosal surface (8). In contrast

to killed OCVs, live vaccines will likely offer single dose efficacy, a particularly important

feature for reactive vaccination campaigns during epidemics. Furthermore, live OCVs

appear to be more effective in children less than 5 years of age (9, 10), a group that is

highly susceptible to death from cholera and that is not adequately protected by killed

OCVs. In volunteer studies, these vaccines have shown great promise (11, 12), but

none are approved for use in cholera endemic regions.

The development of both inactivated and live OCVs has been hampered by the

lack of a small animal model that closely recapitulates cholera pathogenesis in the

setting of normal immune reactivity. *V. cholerae* readily colonizes the intestines of infant

mice and infant rabbits, reviewed in (13), where cholera-like disease can be observed,

but these models lack mature immune systems. Conventionally raised adult mice

cannot be orally colonized by *V. cholerae,* likely due to their resident gut microbiota

(14). Germ free (GF) adult mice, which lack a microbiota, can be colonized by *V.*

cholerae and have been used to profile OCV immunogenicity, but immune and intestinal

physiological development is aberrant in these animals (15). Nonetheless, vaccinated

GF mice develop immune correlates of clinical protection against toxigenic *V. cholerae,*

including circulating vibriocidal antibodies and antigen specific antibody responses (14,

16, 17). Another limitation of the GF adult mouse model is that following oral

administration of a live OCV, they remain consistently colonized by *V. cholerae* for

periods exceeding 3 months (14, 17), making single-dose live OCV regimens difficult to

interpret; moreover, the prolonged colonization in this model precludes challenging

vaccinated animals with virulent *V. cholerae.* The consequence and significance of long-

term monocolonization with a live OCV in GF mice also remains unknown.

Many studies have shown that oral administration of broad-spectrum antibiotics

to mice depletes the gut microbiota and enables intestinal colonization by diverse

bacteria (18). This is beneficial as antibiotic treated mice are conventionally raised and

do not display the immunological and developmental defects that characterize GF mice.

Antibiotic treatments can enable wild-type *V. cholerae* intestinal colonization for similar

durations, typically 5-7 days, that humans are colonized by live OCVs (19, 20). We

sought to leverage this model to profile HaitiV, a live-attenuated OCV derived from

HaitiWT, a virulent *V. cholerae* O1 Ogawa clinical strain isolated during the Haitian

cholera outbreak (21). HaitiV is non-toxigenic and highly engineered for biosafety and in

infant rabbits HaitiV provides unprecedented rapid protection against virulent *V.*

cholerae within 24 hours of administration (21); furthermore, we showed that this

vaccine is immunogenic in GF mice (17).

Here, we adopted a streptomycin treated adult mouse model of *V. cholerae* (22)

to profile HaitiV's immunogenicity. Furthermore, we modified this model to assess the

protective efficacy of immune responses in immunized female mice, by challenging their

pups with HaitiWT. Our findings demonstrate the ease and utility of this approach for

studies of live-attenuated OCVs.

Results

Vaccine inoculation protocol and colonization kinetics.

We modified the protocol presented in Bueno et al (22) and orally treated 4-week-old C57BL/6 female mice with streptomycin (Sm) to deplete their intestinal microbiota to enable a longitudinal study of HaitV intestinal colonization and immunogenicity (Fig. 1A). Three days after initiating Sm treatment, mice were orally gavaged with either a single dose of 10^9 CFU of HaitiV, an equivalent dose of formalin-inactivated HaitiV (FI-HaitiV) to mimic oral vaccination with an inactivated OCV, or sodium bicarbonate as a vehicle control. There were no apparent untoward effects of any of these regimens and over the course of the experiment, all mice gained weight (Fig. 1B).

Plating of fresh fecal pellets (FP) from all animals inoculated with HaitiV revealed ~10^8 CFU/g FP for 5-7 days post inoculation (dpi), suggesting that initially the vaccine robustly colonized the intestines of Sm treated adult mice. However, HaitiV was no longer detectable in FPs by 8-12 dpi (Fig. 1C, Fig. S1B), indicating clearance of the vaccine strain. No CFUs of HaitiV were recovered from Sm treated mice that had been orally inoculated with FI-HaitiV or buffer control. The clearance kinetics of HaitiV from these mice resembles that previously charted by Nygren et al (19) for wild type cholera toxin-producing *V. cholerae* isolates, suggesting that cholera toxin does not play a substantial role in intestinal colonization in this model.

Transient HaitiV colonization elicits antibodies targeting multiple *V. cholerae*

serotypes*.*

Sera from all mice were individually assayed to quantify the circulating vibriocidal

antibody titers (VATs) targeting Ogawa and Inaba *V. cholerae* strains. By 7dpi most

mice did not have detectable VATs but by 14dpi when HaitiV was no longer detectable

in FPs, most (3/5 mice) seroconverted and developed high circulating VATs against

both serotyped matched (Ogawa) and serotype mismatched (Inaba) *V. cholerae* (Fig.

2A, Fig. S1C, Fig. S1D), though highest geometric mean VATs were directed against

serotype matched (Ogawa) isolates (Fig. 2A, Fig. S1C). Only one mouse in the FI-

HaitiV group developed circulating VATs against Ogawa serotype *V. cholerae,* and no

mice in this group developed vibriocidal antibodies against Inaba *V. cholerae* in this

group (Fig. 2B). None of the Sm treated mice that received sodium bicarbonate

developed detectable vibriocidal antibodies. Thus, transient colonization by HaitiV is

sufficient to elicit the generation of *V. cholerae* specific circulating markers of immunity.

Streptomycin treated adult mice resist recolonization by *V. cholerae.*

Administration of live-OCVs to GF mice leads to long term intestinal colonization

(14, 17), precluding the possibility of challenge studies with wild-type *V. cholerae.* Since

the Sm-treated mice cleared HaitiV, we investigated whether animals immunized with

this OCV could be successfully challenged with HaitiWT. At 28dpi, when all mice had

stopped shedding HaitiV, they (live HaitiV, FI-HaitiV, and vehicle control) were orally

treated with sulfamethoxazole and trimethoprim (SXT) to reduce the intestinal

microbiota (Fig. 3A); unlike HaitiV, HaitiWT is resistant to SXT. All three groups of mice

from above and another group, specific pathogen free (SPF) mice that had not been

Sm-treated or exposed to HaitiV was also included in this experiment, to test whether

SXT treatments modify susceptibility to HaitiWT colonization. All 4 groups of mice were

then orally gavaged with 10^9 CFU of HaitiWT, and their FPs monitored for HaitiWT

colonization. Unexpectedly, the three groups of mice (HaitiV, FI-HaitiV, and sodium

bicarbonate) that had been previously treated with Sm failed to be colonized by HaitiWT

(Fig. 3B). In contrast, the SXT-treated SPF mice were robustly colonized by HaitiWT

($\sim 10^8$ CFU/g FP) (Fig. 3B), indicating that although SXT treatment does susceptibilize

mice to *V. cholerae* colonization, the prior Sm treatment of the vaccinated animals

rendered them resistant to recolonization with *V. cholerae*.

Although the FPs of the previously Sm treated and HaitiWT challenged mice

lacked detectable HaitiWT on the Sm agar plates used to detect *V. cholerae*, these

plates contained high numbers of colonies of non-*V. cholerae* bacteria. 16s rRNA

sequencing of Sm resistant small colonies showed that these colonies corresponded to

either *Escherichia coli* or SXT resistant *Lactobacillus murinus*, which together were

present at $\sim 10^8$ CFU/g FP. These observations suggest that a bloom of Sm-resistant

organisms and other changes in the gut microbiota associated with prior oral Sm

treatment render mice resistant to *V. cholerae* colonization.

Offspring of vaccinated dams are protected from virulent *V. cholerae*.

As challenge studies could not be performed in Sm treated adult mice, we turned

to the suckling mouse model of cholera to assess the protective efficacy of the immune

response that was elicited by the transient HaitiV intestinal colonization. We recently

found that the survival of suckling mice in this lethal challenge model can be used to

gauge OCV efficacy (17). As such, animals from HaitiV, FI-HaitiV, and sodium

bicarbonate groups were mated with SPF males and their neonatal pups infected with a

lethal dose of HaitiWT (Fig. 4A).

Pups from HaitiV immunized dams survived significantly longer (median 30hpi)

than pups from dams treated with FI-HaitiV or sodium bicarbonate control (median

~25hpi) (Fig. 4B). Despite the increased survival in the offspring of HaitiV immunized

dams, at time of death, there were no differences in the burden of HaitiWT CFU

recovered from the small intestines of the 3 groups (Fig. 4C). Thus, oral immunization

and transient colonization of Sm-treated mice with HaitiV induces immune responses

that significantly protect against choleric disease in mice.

Discussion

The wide availability of diverse mutant mice and reagents for their study make

mice a preferred model mammal for studies of human disease and therapeutics.

However, adult mice are not susceptible to intestinal colonization with *V. cholerae*,

confounding evaluation of OCVs. Here, we found that Sm-treated adult mice can be

used to investigate the immunogenicity as well as the protective efficacy of live-

attenuated OCVs. Sm-treated adult mice orally inoculated with HaitiV were colonized by

this live-attenuated vaccine for 5-7days and this transient colonization was sufficient to

elicit vibriocidal antibodies against both *V. cholerae* serotypes. Furthermore, pups born

to HaitiV immunized Sm-treated mice exhibited prolonged survival following lethal

challenge with HaitiWT compared to pups born to dams immunized with FI-HaitiV or

dams treated with vehicle control. Together, these findings suggest that this model

should be valuable for further studies of the efficacy and protective mechanisms of live

OCVs.

Antibiotic treatment of adult mice limits the roles that the intestinal microbiota

plays in inhibiting live OCV colonization. Unexpectedly, we found that oral Sm treatment

enables the expansion of Sm and SXT-resistant microbes that also inhibit *V. cholerae*

colonization (Fig. 3). Detailed analyses of the composition of the microbiota that are

initially killed by oral Sm administration, and those that bloom after the antibiotic is

withdrawn would offer valuable clues into which bacteria inhibit *V. cholerae* colonization.

Though SXT treatment did not enable HaitiWT colonization of previously Sm treated

animals (Fig. 3), it is possible that different antibiotic cocktails would facilitate re-

challenge studies; for example, clindamycin was recently found to enable *V. cholerae*

intestinal colonization of adult mice (23). Establishing conditions for re-challenge studies

would simplify the model, bypassing the requirement for challenge studies in the

offspring of immunized mice; however, only intestinal colonization resistance could be

assayed in adult re-challenge studies since unlike suckling mice, adult mice are

resistant to choleric diarrhea.

Although pups from HaitiV immunized Sm-treated dams survived longer than

control animals, there was no difference in the CFU burden of HaitiWT in pups from

immunized or control dams at the time they became moribund (Fig. 4). It is possible that

vaccination partially delayed the replication of HaitiWT or, in addition, vaccination may

elicit antibodies that antagonize toxic factors, such as cholera toxin, that promote

disease, but do not directly modulate bacterial colonization. HaitiV ectopically expresses

the GM1-binding B subunit of cholera toxin and immune responses to this non-toxic

component of cholera toxin are linked to short term protection against cholera and

enterotoxigenic *E. coli* (24).

In contrast to GF mice, which were colonized HaitiV for many months in our

previous study (17),Sm-treated mice were only colonized by HaitiV for several days

(Fig. 1). This is similar to the duration that live OCVs colonize the human intestine (20,

25, 26) meaning that Sm treated mice may provide a more physiologically relevant

model to gauge OCV immunogenicity. However, it is important to note that although

Sm-treated mice display HaitiV clearance kinetics that mimic human OCV clearance,

mice are not natural hosts for *V. cholerae. V. cholerae* intestinal colonization in

susceptible adult mice occurs primarily in the colon and does not rely on TCP, a critical

factor for colonization of the small intestine in humans as well in infant mice and rabbits

(13, 19). Nonetheless, since adaptive protective immune responses are stimulated by

HaitiV in both Sm treated and GF mice, these models have value for testing vaccine

immunogenicity.

We adopted the strategy that Sit et al (17) used for studying the protective

efficacy of OCVs in immunized GF mice and coupled the suckling mouse model of

cholera with the Sm-treated adult model of OCV immunogenicity. However, in contrast

to the GF model, when the offspring of HaitiV immunized Sm-treated mice were

inoculated with HaitiWT, their dams had undetectable HaitiV CFUs in their feces and

VATs in their sera (Fig. 1, Fig. 4); these conditions more closely mimic those that will be

present when immunized humans are exposed to *V. cholerae*. Despite the absence of

circulating VATs, which are known to be relatively short-lived (27) , the offspring of

HaitiV-immunized dams exhibited significantly delayed death due to cholera-like

disease, but all succumbed (Fig. 4). In contrast, the offspring of HaitiV-immunized GF

dams were completely protected from challenge with HaitiWT (17), demonstrating the

increased potency of HaitiV immunization in GF mice. The greater effectiveness of

HaitiV vaccination in GF mice is likely attributable to the constant stimulation of the

intestinal mucosa by HaitiV in the persistently mono-colonized GF animals and could be

consistent with multiple-dose live OCV regimens. Thus, even though GF mice have

immune defects, they may overestimate the potency of HaitiV and other live OCVs.

Killed whole cell vaccines like Shancol have ~50% efficacy in single dose trials

(29). Notably, in marked contrast to HaitiV, we found in the Sm-treated mice model that

a single dose of FI-HaitiV, which is similar to current killed whole cell vaccines, did not

elicit protective immunity, consistent with the idea that live-attenuated HaitiV is far more

immunogenic than killed vaccines, at least when administered as a single dose.

In summary, we have described an adult mouse model for the investigation of the

protective efficacy of live OCVs. Using this model, we found that a single oral dose of

HaitiV and transient colonization by the vaccine can elicit vibriocidal antibody titers and

protective immune responses. Future work should enable refinement of this model to

allow re-challenge of immunized animals and further characterization of these protective

immune responses. Furthermore, given the availability of genetically defined mutant

C57BL/6 mice, the model described here should be a valuable approach for unraveling

the molecular determinants of live vaccine-mediated protection against cholera and

other mucosal pathogens.

Materials and Methods

Bacterial Strains and culture conditions

Vibrio cholerae strains were grown in Luria-Bertani (LB) broth supplemented with

relevant antibiotics: streptomycin (Sm) at 200μg/ml and sulfamethoxazole trimethoprim

(SXT) at 80μg/mL and 16μg/mL respectively at 37°C with continuous shaking at

220rpm. For LB agar plates 1.5% agar was used and was supplemented with 5-bromo-

4-chloro-3-indolyl-β-d-galactopyranoside (X-Gal) at 60μg/mL. Bacteria were kept as -

80°C stocks in LB with 40% glycerol.

Oral immunization scheme

4-week-old C57BL/6 female mice were purchased (Charles River) and housed in a BL-2

facility under conventional rearing conditions for the duration of the studies. On day 0,

all mice were briefly anesthetized with isoflurane and orally gavaged with 20mg of

streptomycin in 100uL of sterilized water. Mice were then provided drinking water

supplemented with 5mg/mL Sm for 72 hours (22). After this, mice were then gavaged

with 10^9 CFU of an overnight culture of either HaitiV, or HaitiV inactivated by 10%

formalin (FI-HaitiV) for 15 minutes, and then resuspended in 2.5% Na_2CO_3. Control

mice were also gavaged with 100uL of 2.5% Na_2CO_3 alone. Mice were then provided

drinking water supplemented with 200µg/ml Sm for 14 days before being returned to

unsupplemented water. All mice were weighed weekly and blood samples were

retrieved from each mouse by tail vein incision at each weighing. Blood samples were

clotted for 45 minutes at room temperature, centrifuged at 20000xg for 10 minutes, and

the serum stored at $-20^{o}C$ for future analysis.

Fresh fecal pellets were collected daily from each mouse, weighed, and plated in serial

dilutions on LB+Sm+X-Gal agar to determine the colonization of each mouse by HaitiV.

HaitiV colonizes are white due to a disrupted *lacZ*. The limit of detection of this assay

represents the lowest CFU count that could be detected for a fecal pellet of that weight.

Quantification of vibriocidal antibody titers

Circulating titers of vibriocidal antibodies were quantified by determining the minimal

serum dilution required to lyse PIC158 (Ogawa) or PIC018 (Inaba) *V. cholerae* as

described previously (30, 31) with minor modifications. Briefly, serial dilutions of serum

were incubated with guinea-pig complement (Sigma) and the target strain, and then

allowed to grow in BHI media. Reported titers are the dilution of serum that caused

more than 50% reduction in target strain optical density when compared to normal

saline control wells. A mouse monoclonal antibody 432A.1G8.G1.H12 targeting *V.*

cholerae O1 OSP was a positive control for the assay. The limit of detection of this

assay represents the lowest serum dilution at which no inhibition of growth could be

detected.

Colonization of immunized mice with toxigenic *V. cholerae*

Mice from all three groups (HaitiV, FI-HaitiV, and vehicle control) as well as 6 week old female SPF mice (Charles River) were anesthetized with isoflurane and gavaged with 16mg of sulfamethoxazole and 3.2mg trimethoprim (SXT) in 100uL of water and provided drinking water supplemented with 4mg/mL SX and 0.8mg/mL T for 72 hours. All mice were then gavaged with 10^9 CFU of an overnight culture of HaitiWT and switched to 0.16mg/mL SX and 0.032mg/mL T in their drinking water. Fecal pellets were collected daily and plated on both SXT agar and Sm agar. Non-*V. cholerae* colonies were isolated by streaking on fresh Sm plates, and 16s V1/V2 DNA sequences were amplified by colony PCR using primers F341 (TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG CCT ACG GGN GGC WGC AG) and R805 (GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA GGA CTA CHV GGG TAT CTA ATC C) and sequenced. Identification of the bacteria was performed using BLAST (NCBI)

Infant mouse survival assay

The infant mouse survival assay was adapted from a previous report (17). Female mice were mated with age matched SPF male mice (Charles River) and singly housed at E18 for delivery. At the third day of life (P3) pups were orally gavaged with 10^7 CFU of HaitiWT and returned to their dams. Pups were monitored every 6 hours until the first signs, typically including diarrhea and dehydration, were evident. At this point, monitoring was performed every 30 minutes until pups were moribund. Pups were then removed from the nest, euthanized with isoflurane, and decapitated and dissected. The small intestines of each pup were excised, homogenized, and plated on LB agar with Sm and X-Gal.

Statistical analysis

All statistical analyses were performed with Prism 8 (Graphpad). Infant mouse survival

curves were analyzed with a log rank (Mantel Cox) test and CFU data were analyzed

with a Mann Whitney U test.

Animal use statement

All experiments in this study were performed was approved by the Brigham and

Women's Hospital IACUC (protocol 2016N000416) and in compliance with the NIH

Guide for Use and Care of Laboratory animals.

Acknowledgements:

This work was supported by NIH grant (R01-AI-042347-24). BS was supported by an

NSERC PGS-D fellowship (PGSD3-487259-2016). B.F., B.S., and M.K.W. designed the

experiments. B.F. and B.S. performed the experiments. B.F. and M.K.W. analyzed the

data. B.F., B.S, and M.K.W. wrote the manuscript. All authors reviewed and approved

the manuscript.

We are grateful to Edward Ryan for providing the monoclonal antibody used for the

serum assays. We thank the MKW laboratory members for helpful discussions.

M.K.W. receives funding from the HHMI and the NIH.

References:

1. Ali M, Nelson AR, Lopez AL, Sack DA. 2015. Updated global burden of cholera in endemic countries. PLoS Negl Trop Dis 9:e0003832.

2. Hisatsune K, Kondo S, Isshiki Y, Iguchi T, Haishima Y. 1993. Occurrence of 2-O-Methyl-N-(3-Deoxy-L-glycero-tetronyl)-D-perosamine (4-amino-4,6-dideoxy-D-manno-pyranose) in Lipopolysaccharide from Ogawa but Not from Inaba O Forms of O1 Vibrio cholerae. Biochem Biophys Res Commun 190:302–307.

3. Chin CS, Sorenson J, Harris JB, Robins WP, Charles RC, Jean-Charles RR, Bullard J, Webster DR, Kasarskis A, Peluso P, Paxinos EE, Yamaichi Y, Calderwood SB, Mekalanos JJ, Schadt EE, Waldor MK. 2011. The origin of the Haitian cholera outbreak strain. N Engl J Med 364:33–42.

4. Levine MM, Black RE, Clements ML, Cisneros L, Nalin DR, Young CR. 1981. Duration Of Infection-Derived Immunity To Cholera. J Infect Dis 143:818–820.

5. Albert MJ, Alam K, Ansaruzzaman M, Qadri F, Sack RB. 1994. Lack of cross-protection against diarrhea due to vibrio cholerae o139 (bengal strain) after oral immunization of rabbits with v. cholerae o1 vaccine strain cvd103-hgr. J Infect Dis 169:230–231.

6. Bi Q, Ferreras E, Pezzoli L, Legros D, Ivers LC, Date K, Qadri F, Digilio L, Sack DA, Ali M, Lessler J, Luquero FJ, Azman AS, Cavailler P, Date K, Sreenivasan N, Matzger H, Luquero F, Grais R, Wiesner L, Ko M, Rouzier V, Peak C, Qadri F, Landegger J, Lynch J, Azman A, Sack D, Henkens M, Ciglenecki I, Ivers L, Diggle E, Weiss M, Hinman A, Maina K, Mirza I, Gimeno G, Levine M. 2017. Protection

against cholera from killed whole-cell oral cholera vaccines: a systematic review

and meta-analysis. Lancet Infect Dis 17:1080–1088.

7.   Parker LA, Rumunu J, Jamet C, Kenyi Y, Lino RL, Wamala JF, Mpairwe AM,

Muller V, Llosa AE, Uzzeni F, Luquero FJ, Ciglenecki I, Azman AS. 2017.

Neighborhood-targeted and case-triggered use of a single dose of oral cholera

vaccine in an urban setting: Feasibility and vaccine coverage. PLoS Negl Trop Dis

11:e0005652.

8.   Lycke N. 2012. Recent progress in mucosal vaccine development: Potential and

limitations. Nat Rev Immunol.

9.   Qadri F, Chowdhury MI, Faruque SM, Salam MA, Ahmed T, Begum YA, Saha A,

Al Tarique A, Seidlein L V., Park E, Killeen KP, Mekalanos JJ, Clemens JD, Sack

DA. 2007. Peru-15, a live attenuated oral cholera vaccine, is safe and

immunogenic in Bangladeshi toddlers and infants. Vaccine 25:231–238.

10.   Sur D, Lopez AL, Kanungo S, Paisley A, Manna B, Ali M, Niyogi SK, Park JK,

Sarkar B, Puri MK, Kim DR, Deen JL, Holmgren J, Carbis R, Rao R, Van NT,

Donner A, Ganguly NK, Nair GB, Bhattacharya SK, Clemens JD. 2009. Efficacy

and safety of a modified killed-whole-cell oral cholera vaccine in India: an interim

analysis of a cluster-randomised, double-blind, placebo-controlled trial. Lancet

374:1694–1702.

11.   Islam K, Hossain M, Kelly M, Mayo Smith LM, Charles RC, Bhuiyan TR, Kováč P,

Xu P, LaRocque RC, Calderwood SB, Simon JK, Chen WH, Haney D, Lock M,

Lyon CE, Kirkpatrick BD, Cohen M, Levine MM, Gurwith M, Harris JB, Qadri F,

403  Ryan ET. 2018. Anti-O-specific polysaccharide (OSP) immune responses

404  following vaccination with oral cholera vaccine CVD 103-HgR correlate with

405  protection against cholera after infection with wild-type Vibrio cholerae O1 El Tor

406  Inaba in North American volunteers. PLoS Negl Trop Dis 12:e0006376.

407 12. Chen WH, Garza J, Choquette M, Hawkins J, Hoeper A, Bernstein DI, Cohenb

408  MB. 2015. Safety and immunogenicity of escalating dosages of a single oral

409  administration of peru-15 pCTB, a candidate live, attenuated vaccine against

410  enterotoxigenic Escherichia coli and Vibrio cholerae. Clin Vaccine Immunol

411  22:129–135.

412 13. Ritchie JM, Waldor MK. 2009. Vibrio cholerae Interactions with the

413  Gastrointestinal Tract: Lessons from Animal Studies, p. 37–59. *In* Current Topics

414  in Microbiology and Immunology.

415 14. Butterton JR, Ryan ET, Shahin RA, Calderwood SB. 1996. Development of a

416  germfree mouse model of Vibrio cholerae infection. Infect Immun 64:4373–4377.

417 15. Round JL, Mazmanian SK. 2009. The gut microbiota shapes intestinal immune

418  responses during health and disease. Nat Rev Immunol.

419 16. Crean TI, John M, Calderwood SB, Ryan ET. 2000. Optimizing the germfree

420  mouse model for in vivo evaluation of oral Vibrio cholerae vaccine and vector

421  strains. Infect Immun 68:977–981.

422 17. 2019. Oral immunization with a probiotic cholera vaccine induces broad

423  protective immunity against Vibrio cholerae colonization and disease in mice.

424  PLoS Negl Trop Dis 13:e0007417.

18.   Kennedy EA, King KY, Baldridge MT. 2018. Mouse microbiota models:

Comparing germ-free mice and antibiotics treatment as tools for modifying gut

bacteria. Front Physiol 9.

19.   Nygren E, Li BL, Holmgren J, Attridge SR. 2009. Establishment of an adult mouse

model for direct evaluation of the efficacy of vaccines against Vibrio cholerae.

Infect Immun 77:3475–3484.

20.   Gotuzzo E, Butron B, Seas C, Penny M, Ruiz R, Losonsky G, Lanata CF,

Wasserman SS, Salazar E, Kaper JB, Cryz S, Levine MM. 1993. Safety,

immunogenicity, and excretion pattern of single-dose live oral cholera vaccine

CVD 103-HgR in Peruvian adults of high and low socioeconomic levels. Infect

Immun.

21.   Hubbard TP, Billings G, Dörr T, Sit B, Warr AR, Kuehl CJ, Kim M, Delgado F,

Mekalanos JJ, Lewnard JA, Waldor MK. 2018. A live vaccine rapidly protects

against cholera in an infant rabbit model. Sci Transl Med 10:eaap8423.

22.   Bueno E, Sit B, Waldor MK, Cava F. 2018. Anaerobic nitrate reduction divergently

governs population expansion of the enteropathogen Vibrio cholerae. Nat

Microbiol.

23.   You JS, Yong JH, Kim GH, Moon S, Nam KT, Ryu JH, Yoon MY, Yoon SS. 2019.

Commensal-derived metabolites govern Vibrio cholerae pathogenesis in host

intestine. Microbiome 7:132.

24.   Kauffman RC, Bhuiyan TR, Nakajima R, Mayo-Smith LM, Rashu R, Hoq MR,

Chowdhury F, Khan AI, Rahman A, Bhaumik SK, Harris L, O'Neal JT, Trost JF,

Alam NH, Jasinskas A, Dotsey E, Kelly M, Charles RC, Xu P, Kováč P, Calderwood SB, Ryan ET, Felgner PL, Qadri F, Wrammert J, Harris JB. 2016. Single-cell analysis of the plasmablast response to Vibrio cholerae demonstrates expansion of cross-reactive memory B cells. MBio 7.

25. Lagos R, San Martin O, Wasserman SS, Prado V, Losonsky GA, Bustamante C, Levine MM. 1999. Palatability, reactogenicity and immunogenicity of engineered live oral cholera vaccine CVD 103-HgR in Chilean infants and toddlers. Pediatr Infect Dis J 18:624–630.

26. Chen WH, Cohen MB, Kirkpatrick BD, Brady RC, Galloway D, Gurwith M, Hall RH, Kessler RA, Lock M, Haney D, Lyon CE, Pasetti MF, Simon JK, Szabo F, Tennant S, Levine MM. 2016. Single-dose live oral cholera vaccine CVD 103-HgR protects against human experimental infection with vibrio cholerae O1 El Tor. Clin Infect Dis 62:1329–1335.

27. Mosley WH, Ahmad S, Benenson AS, Ahmed A. 1968. The relationship of vibriocidal antibody titre to susceptibility to cholera in family contacts of cholera patients. Bull World Health Organ 38:777.

28. Sit B, Zhang T, Fakoya B, Akter A, Biswas R, Ryan ET, Waldor MK. 2019. Oral immunization with a probiotic cholera vaccine induces broad protective immunity against vibrio cholerae colonization and disease in mice. PLoS Negl Trop Dis 13:e0007417.

29. Qadri F, Ali M, Lynch J, Chowdhury F, Khan AI, Wierzba TF, Excler JL, Saha A, Islam MT, Begum YA, Bhuiyan TR, Khanam F, Chowdhury MI, Khan IA, Kabir A,

Riaz BK, Akter A, Khan A, Asaduzzaman M, Kim DR, Siddik AU, Saha NC,

Cravioto A, Singh AP, Clemens JD. 2018. Efficacy of a single-dose regimen of

inactivated whole-cell oral cholera vaccine: results from 2 years of follow-up of a

randomised trial. Lancet Infect Dis 18:666–674.

30.   Son MS, Taylor RK. 2011. Vibriocidal assays to determine the antibody titer of

patient sera samples. Curr Protoc Microbiol CHAPTER:Unit6A.3.

31.   Sayeed MA, Bufano MK, Xu P, Eckhoff G, Charles RC, Alam MM, Sultana T,

Rashu MR, Berger A, Escobedo GG, Mandlik A, Bhuiyan TR, Leung DT,

LaRocque RC, Harris JB, Calderwood SB, Qadri F, Vann WF, Kováč P, Ryan ET.

2015. A cholera conjugate vaccine containing ospecific polysaccharide (OSP) of

V. cholera o1 inaba and recombinant fragment of tetanus toxin heavy chain

(OSP:rTTHC) induces serum, memory and lamina proprial responses against

OSP and is protective in mice. PLoS Negl Trop Dis 9:e0003881.

Figure legends:

Figure 1. Transient intestinal colonization of HaitiV in Sm-treated adult mice following treatment with streptomycin. A) Schematic of streptomycin treatment and single oral inoculation with HaitiV. Black arrowheads indicate peroral treatment with either streptomycin in sodium bicarbonate, or oral gavage with bacteria. B) Bodyweight of all mice over the course of this study. C) Fecal shedding of HaitiV from mice inoculated with HaitiV; open symbols depict CFU levels below the limit of detection.

Figure 2. Serum vibriocidal antibody titers in Sm-treated mice immunized with HaitiV. Circles indicate the lowest dilutions at which specific vibriocidal activity was detected, and the height of the bars represent geometric mean titers in each group. Ogawa *V. cholerae* PIC158 was used to measure Ogawa serotype-specific antibodies (black) and Inaba *V. cholerae* PIC018 was used to measure Inaba serotype-specific antibodies (red). Titers below the limit of detection are indicated by open symbols.

Figure 3. SXT treatment and re-colonization of Sm-treated adult mice by *V. cholerae.* A) All mice were treated with sulfamethoxazole and trimethoprim (SXT) and dosed with 10^9 CFU of HaitiWT on day 0. B) Fecal shedding of HaitiWT from all mice; open symbols represent fecal pellets from which no CFUs of HaitiWT could be detected. An asterisk denotes differences with a P value of <0.05 as determined by a Mann Whitney U test.

Figure 4. Transient colonization by HaitiV is protective in an infant mouse model

of cholera. A) Survival curves of pups born to dams inoculated with HaitiV (n=15),

formalin inactivated-HaitiV (n=24) or vehicle buffer (n=16) after HaitiWT challenge.

Differences in the survival curves were determined by a log rank (Mantel Cox) test

(p=0.0013 for HaitiV vs FI-HaitiV). B) HaitiWT CFUs recovered from the small intestines

of the neonatal mice at the time they were moribund. Differences in CFU burdens were

determined by a Mann Whitney U test.

.

Figure 1. Transient intestinal colonization of HaitiV in Sm-treated adult mice following treatment with streptomycin. A) Schematic of streptomycin treatment and single oral inoculation with HaitiV. Black arrowheads indicate peroral treatment with either streptomycin in sodium bicarbonate, or oral gavage with bacteria. B) Bodyweight of all mice over the course of this study. C) Fecal shedding of HaitiV from mice inoculated with HaitiV; open symbols depict CFU levels below the limit of detection.

Figure 2. Serum vibriocidal antibody titers in Sm-treated mice immunized with HaitiV.
Circles indicate the lowest dilutions at which specific vibriocidal activity was detected, and
the height of the bars represent geometric mean titers in each group. Ogawa *V. cholerae*
PIC158 was used to measure Ogawa serotype-specific antibodies (black) and Inaba *V.
cholerae* PIC018 was used to measure Inaba serotype-specific antibodies (red). Titers below
the limit of detection are indicated by open symbols.

Figure 3. SXT treatment and re-colonization of Sm-treated adult mice by *V. cholerae*. A) All mice were treated with sulfamethoxazole and trimethoprim (SXT) and dosed with 10^9 CFU of HaitiWT on day 0. B) Fecal shedding of HaitiWT from all mice; open symbols represent fecal pellets from which no CFUs of HaitiWT could be detected. An asterisk denotes differences with a P value of <0.05 as determined by a Mann Whitney U test.

Figure 4. Transient colonization by HaitiV is protective in an infant mouse model of cholera. A) Survival curves of pups born to dams inoculated with HaitiV (n=15), formalin inactivated-HaitiV (n=24) or vehicle buffer (n=16) after HaitiWT challenge. Differences in the survival curves were determined by a log rank (Mantel Cox) test (p = 0.0013 for HaitiV vs FI-HaitiV). B) HaitiWT CFUs recovered from the small intestines of the neonatal mice at the time they were moribund. Differences in CFU burdens were determined by a Mann Whitney U test.