Hypothetical control of postural sway

Madhur Mangalam[1,*] and Damian G. Kelty-Stephen[2,†]

[1]Department of Physical Therapy, Movement and Rehabilitation Sciences, Northeastern University, Boston, MA 02115, USA

[2]Department of Psychology, Grinnell College, Grinnell, IA 50112, USA

Authors for correspondence:

Madhur Mangalam

email: m.mangalam@northeastern.edu

Damian G. Kelty-Stephen

keltysda@grinnell.edu

Abstract

Quiet standing exhibits strongly intermittent variability that has inspired at least two interpretations. First, variability can be intermittent through the alternating engagement and disengagement of complementary control processes at distinct scales. A second and perhaps deeper way to interpret this intermittency is through the possibility that postural control depends on cascade-like interactions across many timescales at once, suggesting specific non-Gaussian distributional properties at different timescales. Multiscale probability density function (PDF) analysis shows that quiet standing on a stable surface exhibits a crossover from low, increasing non-Gaussianity (consistent with exponential distributions) at shorter timescales, reflecting inertial control, towards higher non-Gaussianity. Feedback-based control at medium to longer timescales yields a linear decrease characteristic of cascade dynamics. Destabilizing quiet standing with unstable surface or closed eyes serves to attenuate inertial control and to elicit more of the feedback-based control over progressively shorter timescales. The result was to strengthen the appearance of the linear decay indicating cascade dynamics. Finally, both linear and nonlinear indices of postural sway also govern the relative strength of crossover or of linear decay, suggesting that tempering of non-Gaussianity across log-timescale is a function of both extrinsic constraints and endogenous postural control. These results provide new evidence that cascading interactions across longer-timescales supporting postural corrections can even recruit shorter-timescale processes with novel task constraints that can destabilize posture.

Keywords: cascades; center of pressure; fluctuations; postural control; quiet stance; scale invariance

1. Introduction

1.1. Postural sway in quiet standing

Quiet standing—a fundamental motor act supporting other movement tasks—exhibits intermittent variability. This intermittency can be interpreted in various ways. First, variability can be intermittent due to the engagement and disengagement of control processes. For instance, during quiet standing, the body continually sways about a constant average position of the global ground reaction force (center of pressure, CoP) trajectory. Within the center of the base of upright support, it is traditionally thought that the postural system relies on postural corrections based on sensory feedback at longer timescales, towards the margins of the base of support [1,2]. A second and perhaps a deeper way to interpret this intermittency is through the possibility that feedback-based corrections depend on the cascade-like interactions across timescales. For instance, more modern interpretations of postural control recognize that upright standing is not always about reining the body within a margin. Instead, postural control appears to respect the maintenance of equilibrium points around which the body can 'tremble,' but postural control can also 'ramble,' sometimes retaining equilibrium points in place but sometimes moving those about [3,4]. Quiet standing accomplishes its rambling equilibrium through the integration of information from visual, vestibular, and somatosensory receptors, in conjunction with the passive properties of the musculoskeletal system. The neurological and physiological processes that have been suggested to dictate intermittency in postural sway include active mechanical processes [5–7], passive reflex mechanisms [8,9] anticipatory [10–12] and feedback processes [13,14], and exploratory processes used by the central nervous system (CNS) to generate sensory information [15,16].

The sources of intermittency noted above should each entail characteristic signatures on the distributional properties of postural sway. Different modes of postural control enlist different kinds of relationships between consecutive fluctuations. For instance, feedback-based correction will entail a temporal structure that feedforward fluctuations may not, depending on how much independence postural control might leave between fluctuations. The change in expected independence will entail a change in expected distributions. An important theme in what follows is aligning the transition in control with a transition in distribution of measured postural fluctuations.

1.2. Aligning feedback-based control with multiplicative random processes and non-Gaussianity

The transition from more to less independence among postural fluctuations may prompt the transition from more Gaussian to less Gaussian (i.e., to more non-Gaussian) probability density functions (PDFs). As independence gives ways to interactivity, the statistical processes that these postural fluctuations resemble move from additive to multiplicative. Independence aligns with additivity, and interactivity aligns with multiplicative processes. First, postural sway remaining within the base of support is most compatible with independence among postural fluctuations. Under these conditions, feedback-based correction becomes a lower priority for maintaining an upright stance, especially as a precision suprapostural task begins to take higher priority [17]. Specifically, shorter-timescale sway roams 'unchecked' resembling Brownian motion [1,2], exhibiting inertia-driven temporal correlations across fluctuations whose nonzero size is drawn from distributions with thin tails (e.g., exponential) [18,19]. The independence of postural fluctuations from feedback in this inertial mode ensures an additive random process for which Gaussian variance effectively describes sway. Nearer to the boundaries of the support surface, the postural system draws on feedback to maintain stability, correcting for and so reversing the inertial tendencies. Feedback is fundamentally contingent and interactive, leading postural fluctuations to exhibit lognormal distributions characteristic of multiplicative random processes [20,21]. The lognormal PDFs' heavy tails entail that a non-Gaussian variance better describe sway. That is, stable quiet standing should exhibit exponential and lognormal distributions at shorter and longer timescales devoted to inertial and correction-based control modes, respectively. Variance requires different definitions for lognormal and Gaussian distributions, and the non-Gaussianity of multiplicative processes manifests as an excess of lognormal variance beyond Gaussian variance.

An important step beyond the simple additive-vs.-multiplicative dichotomy is our proposal that feedback-based corrections are explicitly cascade-like—multiplicative random processes with hierarchical interactivity [22–24]. Feedback is necessarily a consequence of outward movements of the organism into the

surrounding task context, and hence it is always sequential—organism activity always prompts later feedback.
The question of the hierarchical structure only enters into this sequence as a potential nuance.

The sparsest series of feedback may show no cascade structure. That is, feedback could
straightforwardly be a chain of sensorimotor links (sensory events only following motor events, and vice versa).
Time-series modeling often refers to sequences with feedback only on previous events as having 'short memory'
[25]. The popularity and controversy around heavy-tailed distributions warrants emphasis that this extremely
short memory is a valid interpretation of interactivity from non-Gaussian distributions [26–29]. Many heavy-
tailed distributions result from little more than outliers and so have not even the least meaningful sequential
relationship [30]. Lognormal and other heavy-tailed distributions are a signature of what could be feedback and
more than short-memory feedback, but we cannot take these interpretations for granted without further test. We
can conceive of longer-memory feedback following from longer links in this chain, e.g., a brief sensory event
(e.g., an action potential) that could follow short and long motor events (e.g., a muscle fiber contracting and the
extension of a joint, respectively). Longer-term, less-motor, and more-cognitive factors like intention or
instructions might contribute as well to the sensory event. So, we can conceive at least three lengths of links: the
shortest muscle-fiber link, the slightly longer joint-extension link, and the longest intention link. Feedback can
follow from a mix of long-memory and short-memory without entailing cascade organization [23,24].

1.3. Empirical test of cascades in feedback-based postural control

Multiplicative processes are cascade-like when longer events can reshape shorter events. It would be
wrong to think of cascades and short-memory as exclusive opposites; cascades can exhibit short-memory chain
structure [31,32]. What distinguishes cascades is a hierarchical structure affording not just short-memory but
extensions including long-memory and more fluid, flexible organizations of feedback.

The cascades reflect multiplicative random processes in which the memories at different timescales
intermix. They allow mathematically addressing situations in which feedback is deeply context-sensitive. For
instance, during quiet stance, the value of feedback may change depending upon how intention, joint extension,
and muscle-fiber activity influence each other. In these situations, the cascade formalism reflects an accrual of
possible feedback regimes, not sloughing the other simpler kinds. So, the preceding examples of feedback are
available as limiting cases of more generic cascade mechanisms [32]. The 'chain' metaphor above eventually
becomes too rigid to describe cascades respecting the fact that dynamics have been used to model fluid dynamics
(e.g., actual, hydrological cascades [33]). The interactions amongst links at different scales entail that
nonlinearity is the rule than an exception to a linear sequence.

Nonlinear interactions across scales unleash a deep potential of cascades to invite a wider class of
mathematical models of cause-effect relationships. As nonlinearity brings pliability to once-rigid links in our
chain, cascades begin to look much less like chains and more like tapestries [34–36], branching structures [37–
41], nestings [42–44], or webs [45–47]. This metaphoric diversity mainly illustrates how nonlinear interactions
across scales open up the scientific discourse to various proposals of how hierarchical geometry may (or may
not) constrain the shape of cause-effect relationships. In the context of human postural control, cascade modeling
allows testing whether preceding longer-memory links can all reshape one another as well as that brief sensory
event called feedback. But then, we might note that feedback might be no less multi-layered, including an action
potential in a sensory cell and longer-term perceptual impressions that the postural control system can use in the
future.

Cascade-like multiplicative processes offer a generic, common analytical thread to coordinate this
exploding range of possible forms for cascading feedback processes: 'multifractal geometry.' While cascades
may fire the scientific imagination and allow us to tailor our modeling to different model systems, there is a
promise that the widely diverse manifestations of nonlinear interactions across scales might operate along with
similar multifractal principles [22,48,49]. The present work aims to develop the topic of human postural control
through a multifractal lens to determine the sensitivity of cascade-formalisms to human postural control.

The present work applied multiscale PDF analysis rather than simply examining the ambiguous evidence
of heavy tails. Multiscale PDF elaborates the multifractal formalism for quantifying the strength and form of
cascades via quantifying non-Gaussianity. We tested for a crossover from Gaussianity (and exponential
distribution) at shorter timescales to non-Gaussianity (and lognormal distribution) at longer timescales. To test

whether the crossover was specific to the timescale-dependence of postural corrections, we examined multiscale
non-Gaussianity under manipulations intended to destabilize posture. We hypothesized that destabilizing posture
would elicit non-Gaussian postural corrections into the short timescales not traditionally associated with
corrections in response to feedback. Multiscale PDF results indicate cascade dynamics when the full distribution
of measured fluctuations exhibits a linear decay of non-Gaussianity with log-timescale. We compared original
data to surrogate data to ensure that these cascade-like results depended on nonlinear interactions across scales
[50–52].

1.4. Multiscale probability density function (PDF) analysis of postural sway in quiet standing

A central feature of intermittent, cascading fluctuations is the heterogeneity of variance beyond the bounds of a
Gaussian probability distribution function (PDF). An important question then is about the expected PDF shape of
a system whose intermittency reflects the cascading of fluctuation across many scales. Research on intermittency
in the turbulent fluid flow [53] has inspired a method for quantifying this heterogeneity of variance through a
metric λ_s^2 [54,55]. This metric speaks directly to a point raised above: multiplicative random processes will
generate a lognormal PDF with larger variance than the Gaussian PDF resulting from additive random processes.
As we had noted, variance calculated for Gaussian PDFs underestimates lognormal variance for multiplicative
random processes. λ_s^2 expresses the excess of lognormal variance compared to Gaussian variance. Hence, λ_s^2 is
zero only for Gaussian PDFs and increases as for progressively non-Gaussian PDFs.

Multiscale PDF analysis aims to distill deep insights about the random process producing measured
behavior via empirical calculations of variance. Empirical variance can be unstable [56], depending on
contingent aspects of measurement (e.g., sample size). Multiscale PDF analysis quantifies λ_s^2 at a variety of
timescales, coarse-graining measured fluctuations for progressively longer timescales. Controlling for how
coarse-graining might depress variance, multiscale PDF reveals crucial features of cascades based on how
quickly or slowly λ_s^2 decays with timescale [57]. At shorter and medium scales, higher λ_s^2 indicates stronger
evidence of cascades. Theoretical simulations with idealized cascades show monotonic decay of λ_s^2 with longer
timescales. Additionally, distinct kinds of cascades show distinct types of decay. For scale-invariant cascades, λ_s^2
decreases linearly with the natural logarithm of timescale (log-timescale), but cascades unfolding across a
limited range of timescales exhibit quadratic decay with log-timescale [52].

This issue of scale-dependency motivates our attempt to manipulate the transition between inertial and
feedback-based postural control by destabilizing the support surface of closing the eyes. This transition is scale-
dependent—that is, it depends on the width of the base of support. We expected that the short-scale postural
sway roaming inertially unchecked through the base of support would show low λ_s^2, and higher λ_s^2 only at
medium timescales. Thus, we expected that the scale of the base of support would produce a quadratic shape,
depressing the λ_s^2-vs.-log-timescale curve over the short and long timescales. Finally, we expect that
destabilizing the support surface—which effectively narrows the base of support and widens the range of
timescales—or closing the eyes would reduce the scale-dependence of λ_s^2 and resemble more the linear decay
associated with scale-invariant cascades.

While the manipulations of support surface or vision might make the λ_s^2-vs.-log-timescale curve less
quadratic and more linear, their effects must depend on endogenous postural control. Multiscale PDF analysis of
both heart-rate variability (HRV) [54,55,58–60] and neuronal interspike intervals (ISI) [61] suggests that
endogenous control processes account for the variability in λ_s^2-vs.-log-timescale curve for physiological
dynamics. For instance, endogenous sinoatrial dynamics could explain the heavier-tailed PDFs for pathological
cases than for healthy cases in HRV [62], and non-Gaussianity showed relationships with the magnitude
autocorrelation of the ISI [61]. Hence, in addition to testing whether postural corrections at different timescales
in quiet standing change the non-Gaussianity of CoP dynamics, we expected that endogenous variability in
postural control would moderate any observed effect of these manipulations on the non-Gaussianity of postural
fluctuations.

Here, we applied multiscale PDF method to study postural control and test three specific hypotheses in a
reanalysis of a public dataset of kinematics and ground reaction forces [63]. Hypothesis-1 was that quiet
standing on a stable surface would show a crossover between low non-Gaussianity in postural sway (consistent
with exponential distributions) at shorter timescales reflecting lack of sensory corrections, increasing towards
higher and subsequently decreasing (consistent with cascade-driven lognormal distributions) at longer
timescales. Specifically, we tested for this crossover as a negative quadratic polynomial with log-timescale.
Hypothesis-2 was that destabilizing quiet standing would require postural corrections at more of the shorter
timescales, eliciting non-Gaussianity that begins high and decreases across timescales (consistent with cascade-
driven lognormal distribution). Specifically, we expected that quiet standing on an unstable surface or with eyes
closed would cancel the negative quadratic polynomial. Hypothesis-3 was that known indices of postural sway
would significantly moderate the quadratic polynomial, effectively governing the appearance or disappearance
of the crossover.
189
2. Methods

The present work is a reanalysis of a public data set with ground reaction forces of human balance [63].

2.1. Participants

Forty nine participants grouped by age: young (15 males and 12 females, 18–40 years old) and old (11 males and
11 females, > 60 years old) took part in the present study after providing institutionally-approved informed
consent.

2.2. Experimental setup and procedure

Each participant stood still for 60 s under all four possible combinations of vision (eyes open/eyes closed) and
standing surface (stable/unstable). Each participant completed three 60-s trials per condition. Stable condition
involved standing on two 40×60 cm force platforms (100 Hz, OPT400600-1000; AMTI Inc., Watertown, MA)
under each foot, and unstable condition involved standing on two 6-cm high foam blocks (Balance Pad, Airex
AG Inc., Sins, Switzerland), each placed on each force platform.

2.3. Data processing

All data processing was performed in MATLAB 2019b (Matlab Inc., Natick, MA). Trial-by-trial ground reaction
forces yielded a 2D center of pressure (CoP) series, each dimension describing CoP position along anterior-
posterior (*AP*) and medial-lateral (*ML*) axes. Each 60-s trial yielded 6000-sample 2D CoP series and 5999-
sample 2D CoP displacement series. Finally, a 1D CoP planar Euclidean displacement (PED) series described
postural sway along the transverse plane of the body. Iterated Amplitude Adjusted Fourier Transformation
(IAAFT) provided phase-randomized surrogates using original series' spectral amplitudes to preserve only linear
temporal correlations [23,64]. We used the IAAFT surrogate data for two purposes. Primarily, we computed
multiscale PDF analysis on the IAAFT surrogates because any difference in the λ_s^2-vs.-log-timescale curves
between the original and IAAFT series would reveal differences in non-Gaussianity attributable to nonlinear
temporal correlations. Secondarily, as noted below (Section 2.4), we wanted to test whether the difference in
multifractal spectrum width due to nonlinear temporal correlations would moderate the non-Gaussianity in
original series.

2.4. Canonical indices of postural control

Three linear indices were computed for each CoP PED series: (i) Mean of all fluctuations (CoP_*Mean*), (ii)
Standard deviation (CoP_*SD*), and (iii) Root mean square estimate (CoP_*RMSE*). These linear indices quantify
the extent of and variation in postural sway with no reference to how the magnitude of sway at a given instance
is related to the magnitude of away at other instances. Postural control under unstable conditions is associated
with a higher magnitude of and variation in sway [65–67].

Four nonlinear indices were computed for each CoP PED series: (i) Sample entropy (CoP_*SampEn*),
indexing the extent of complexity in the signal [68]. We used $m = 2$ and $r = 0.2$ for computing the *SampEn* [cf.
34]. (ii) Multiscale sample entropy (CoP_*MSE*), indexing signal complexity over multiple timescales [70]. We
used $m = 2$, $r = 0.2$, and $\tau = 20$ for computing the *MSE* [cf. 34]. *SampEn* and *MSE* measure the repeatability or

predictability (i.e., uncertainty) of the possible configurations in a time series. *MSE* extends *SampEn* to multiple
timescales and, providing an additional perspective when the timescale of relevance to the behavior of interest is
unknown. Structural changes in postural sway revealed by *SampEn* and *MSE* have lent insights into the effects of
balance training, vision, and support surface on postural control [71,72]. (iii) We used detrended fluctuation
analysis [73] to compute Hurst's exponent, H_{fGn}, indexing monofractal scaling or the strength of temporal
correlations in original CoP PED series and shuffled versions of each series (CoP_H_{fGn}) using the scaling region:
4, 8, 12,… 1024 [cf. 37]. Shuffling destroys the temporal structure of a signal. (iv) Multifractal spectrum width,
$\Delta\alpha$, indexing multifractality or the extent of multifractal temporal correlations in the signal. Chhabra and
Jensen's direct method [74] was used to compute $\Delta\alpha$ for the original CoP series and the IAAFT surrogate
(CoP_dAlpha) [cf. 38]. Monofractal scaling and multifractal spectrum width have been traditionally implicated
in quantifying the complexity in postural sway and its decay with aging and disease [76–78].

2.5. Multiscale probability density function (PDF) analysis

238 Multiscale PDF analysis characterizes the distribution of abrupt changes in CoP PED series $\{b(t)\}$ using the
239 PDF tail. The first step is to generate $\{B(t)\}$ by integrating $\{b(t)\}$ after centering by the mean b_{ave} (figure 1a):

$$B(t) = \sum_{i=1}^{t/\Delta t} b(i\,\Delta t) - b_{ave}. \qquad (1)$$

A 3rd order polynomial detrends $\{B(t)\}$ within k overlapping windows of length $2s$, s being the timescale
(figure 1b). Intermittent deviation $\Delta_s B(t)$ in k^{th} window from $1+s(k-1)$ to sk in the detrended time series $\{$
$B^d(t) = B(t) - f_{fit}(t)\}$ is computed as $\Delta_s B^d(t) = B^d(t+s) - B^d(t)$, where $1+s(k-1) \le t \le sk$ and $f_{fit}(t)$ is the
polynomial representing the local trend of $\{B(t)\}$, of which the elimination assures the zero-mean probability
density function in the next step (figure 1c). Finally, $\Delta_s B$ is normalized by the SD (i.e., variance is set to one) to
quantify the PDF.

To quantify the non-Gaussianity of $\Delta_s B$ at timescale s, the standardized PDF constructed from all the
$\Delta_s B(t)$ values is approximated by the Castaing model [53], with λ_s as a single parameter characterizing the non-
Gaussianity of the PDF. λ_s is estimated as

$$\lambda_s^2 = \frac{2}{q(q-2)}\left[\ln\left(\sqrt{\pi}\left\langle\left|\Delta_s B\right|^q\right\rangle\right) + \ln\Gamma\left(\frac{q+1}{2}\right) - \frac{q}{2}\ln 2\right], \qquad (2)$$

where $\left\langle\left|\Delta_s B\right|^q\right\rangle$ denotes an estimated value of q^{th} order absolute moment of $\{\Delta_s B\}$. As λ_s^2 increases, the PDF
becomes increasingly peaked and fat-tailed (figure 2a). λ_s^2 can be estimated by Eq. (2) based on q^{th} order
absolute moment of a time series independent of q. Estimating λ_s^2 based on 0.2th moment ($q = 0.2$) emphasizes
the center part of the PDF, reducing the effects of extreme deviations due to heavy-tails and kurtosis. We used
0.2th moment because estimates of λ_s^2 for a time series of ~ 6000 samples are more accurate at lower values of q
[79].

Cascade-type multiplicative process yield the inverse relationship $\lambda_s^2 \sim -\log s$ (figure 2b) [55]. For the
present purposes, we quantified λ_s^2 for each original CoP PED series and corresponding IAAFT surrogate at
timescales 5 to 1000 samples (i.e., 50 ms to 10 s) at steps of 5 samples (50 ms).

To test explicitly for lognormality in postural fluctuations, we used Akaike Information Criterion (AIC)
weights obtained via the more common maximum likelihood estimation (MLE) to determine whether the PDF at
each timescale was a power law, lognormal, exponential, or gamma distribution. We include this analysis also to
complement multiscale PDF analysis to portray the analytical effect of sensitivity to different portions of the
distribution.

2.6. Statistical analysis

To test for crossovers in non-Gaussianity, a linear mixed-effect (LME) model using lmer() function in package *lme4* [80] for *R* [81] tested λ_s^2 vs. log-timescale curves for orthogonal linear and quadratic polynomials, for interactions with grouping variables (Surface × Vision × Age-group × Original, where Original encoded differences in original series from surrogates) and with indices of postural control (Section 2.4). Statistical significance was assumed at the alpha level of 0.05 using package *lmerTest* [82]. To test how lognormality changed with log-timescale, a generalized linear mixed-effect (GLME) fit changes in Lognormality as a dichotomous variable using orthogonal linear, quadratic, and cubic polynomials and tested interaction effects of grouping variables (Surface × Vision × Age × Original) with those polynomials using glmer() function in package *lme4* [80].

3. Results

Figures 3 and 4 provide a schematic of multiscale PDF characterization of postural sway in representative young and old participants standing quietly for 60 s on stable and unstable surfaces with eyes closed (figure 3) and eyes open (figure 4).

3.1. Hypothesis-1

3.1.1. λ_s^2-vs.-log-timescale curves for stable postural conditions

Surface stability elicited nonlinearity in non-Gaussianity with timescale, that is, low short-scale λ_s^2, increasing up to a maximum roughly at 75 samples (750 ms), and decreasing with subsequently longer timescales (table S1; figure 5a). λ_s^2 showed stronger linear decrease ($b = -13.37$, $p < 0.001$) and downward-facing parabolic shape ($b = -2.57$, $p < 0.001$). This profile of λ_s^2-vs.-log-timescale curves held for original series and not for surrogates without original nonlinear interactions across timescales.

3.1.2. The growth of lognormality in postural sway with log-timescale for stable postural conditions

For the most stable postural conditions (i.e., stable surface, eyes open), heaviness of PDF tails aligned with results from multiscale PDF analysis of non-Gaussianity (table S2; figure 6a). The original CoP PED series exhibited less lognormality overall than the respective surrogates ($b = -1.75$, $p < 0.001$). The stable-surface and eyes-open conditions yielded a cubic-polynomial increase in lognormality across log-timescale. Specifically, the CoP PED series showed positive linear ($b = 296.15$, $p < 0.001$) and quadratic ($b = 161.98$, $p < 0.001$) but negative cubic ($b = -209.16$, $p = < 0.001$) growth with log-timescale. However, a striking result is notably heavier PDF tails (i.e., more likely to be lognormal) for surrogates without original nonlinear interactions across timescales.

3.2. Hypothesis-2

3.2.1. λ_s^2-vs.-log-timescale curves for less stable postural conditions

Less stable postural conditions consistently showed increased non-Gaussianity and more cascade-like linear decay of with log-timescale in original series as compared to linear surrogates (table S1; figures 5b and 5d). First, young participants showed a significantly weaker linear decrease in stable-surface with longer timescales ($b = 3.74$, $p < 0.001$), indicating that stronger linear descent of λ_s^2 is attributable to weaker stability with age. However, changes in the nonlinearity of λ_s^2-vs.-log-timescale curves did not depend on age alone or differ between the eyes-open and eyes-closed conditions ($p > 0.05$).

Standing on the unstable surface with eyes open showed significantly less of the downward-facing parabolic shape (figure 5a) as when standing on the stable surface with eyes open ($b = 8.64$, $p < 0.001$; figure 5c). λ_s^2 showed stronger linear decay with log-timescale for the older participants with eyes open ($b = -5.37$, $p < 0.001$). Combining surface instability with eyes closed elicited characteristic evidence of scale-invariant cascades, that is, stronger linear decreases ($b = -1.24$, $p < 0.001$) and weaker quadratic forms ($b = 3.26$, $p <$

309 0.001) of λ_s^2-vs.-log-timescale irrespective of age (figure 5*d*). Young participants exhibited even stronger linear

310 decay ($b = -2.26$, $p = 0.001$) but slightly less reversal of the downward-facing parabolic decay in λ_s^2 ($b = -1.67$, p

311 $= 0.018$). Young participants also showed an overall increase in λ_s^2 with eyes closed ($b = 0.0069$, $p < 0.001$).

3.2.2. The growth of lognormality in postural sway with log-timescale for less stable postural conditions

The relationship between feedback-based control and PDF tails began to unravel, as results about non-

Gaussianity and lognormality of postural fluctuations began to diverge, although some changes in lognormality

aligned with the results about non-Gaussianity. For instance, young participants showed less lognormality

overall ($b = -0.42$, $b < 0.001$; figure 6), indicating that greater lognormality is attributable to weaker stability

with age. However, factors that should have destabilized posture did not increase lognormality—for example,

closing eyes also reduced the likelihood of lognormality ($b = -0.41$, $p < 0.001$; figures 6*b* and 6*d*), signifying a

departure of PDF tail heaviness from non-Gaussian evidence of cascade strength.

Surface instability and its combination with closed eyes and age produced progressively stronger

signatures of cascade structure, but the effects of these factors on PDF tail heaviness were much smaller and

mixed in direction. Surface instability increased lognormality for young participants to a greater degree than for

old participants ($b = 0.37$, $p = 0.002$; figures 6*c* and 6*d*) as well as increased lognormality even more for all

participants with eyes closed ($b = 0.62$, $p < 0.001$). Surface instability weakened quadratic growth with eyes

open ($b = -248.63$, $p = 0.042$; figure 6c) but strengthened it with eyes closed ($b = 417.63$, $p = 0.041$; figure 6d).

Effects of surface instability depended on age, with closed eyes accentuating the cubic form for the old

participants ($b = -231.37$, $p = 0.027$) but reversing it for young participants ($b = 353.95$, $p = 0.040$).

3.3. Hypothesis-3: Effects of indices of endogenous postural control on λ_s^2-vs.-log-timescale curves

All linear and nonlinear indices of postural control showed predictive effects on λ_s^2 (table S1). λ_s^2 typically

decreased with increases in CoP_*Mean* ($b = -41.67$, $p < 0.001$), CoP_*SD* ($b = 31.01$, $p < 0.001$), CoP_*SampEn* (b

$= -0.01$, $p < 0.001$), and CoP_*MSE* ($b = -0.08$, $p < 0.001$), and increased with increases in CoP_*RMSE* ($b =$

51.76, $p < 0.001$). λ_s^2 also increased with both the strength of temporal correlations or fractality (i.e., CoP_H_{fGn}; b

$= 0.08$, $p < 0.00$) and more variation in these temporal correlations or multifractality (i.e., CoP_dAlpha; $b = -$

0.05, $p < 0.00$). Nonetheless, in the tradeoff between temporal correlations, greater 'original-minus-surrogate'

differences in both fractality and multifractality diminished (CoP_H_{fGn}_diff_OS: $b = 2.72$, $p < 0.001$;

CoP_dAlpha_diff_OS: $b = -2.72$, $p < 0.001$). This result confirms the fact that evidence of multifractality with a

relatively weak difference from the surrogates suggests heavy-tailed distributions and thus non-Gaussianity

[59,60]. However, this model, including only the main effects of these indices, speaks only to average

differences in intercept and not to the trajectory of across log-timescale.

Including the interactions of all indices of postural sway with linear and quadratic terms in the prior

model significantly improved the model fit ($\chi_{2423} = 32699$, $p < 0.001$; table S3). Table S3 details the full model,

but in short, it illustrated two major points: first, most of the variability in the quadratic form of the λ_s^2-vs.-log-

timescale curves depended on these indices. This newer model with interactions of these indices with log-

timescale had 69 significant quadratic effects ($p < 0.05$), and all but one significant quadratic effects were

interactions of these indices with log-timescale. That is, the parabolic tempering of non-Gaussianity across log-

timescale in stable-surface condition was mostly a function of endogenous postural control.

4. Discussion

Multiscale PDF analysis was used to test whether unperturbed postural sway exhibits weaker and stronger

evidence of cascade dynamics at relatively short and relatively long timescales. Our major finding is that of a

crossover in non-Gaussianity profiles of postural sway with log-timescale. This crossover likely reflects a

transition between distinct control processes enlisting feedback on short timescales but not long timescales. The

present work joins a growing enterprise in attempting to characterize diverse control processes through nonlinear

methods. Perturbing postural sway with unstable support surface or with closing eyes reveals the capacity of

non-Gaussian cascade dynamics to spread from the longer timescales into shorter timescales in both old and
young participants.

The present findings offer new grounds for doubting whether the traditional open-/closed-loop
interpretation of control is fundamental [1,2,83,84]. Sensory processes and postural corrections do not
necessarily observe a firm divide between fast and slow processes [85]. Instead, they reflect contingent scale-
dependencies of postural corrections emerging from constraints subject to task manipulation. Task manipulation
specifically working to destabilize posture—whether by standing on an unstable surface or closing eyes—reveal
rich cascade-like dynamics in postural control [50–52]. Cascade dynamics afford richly interactive capacity for
movement organization to emerge through the coordination in accordance with constraints at multiple scales
[86,87].

Non-Gaussianity complements the existing understanding that healthy human postural sway exhibits
complexity quantifiable through various means (entropy [69], fractality [88], and multifractality [78]). The
present work shows that the cascade structure of healthy human postural sway varies not simply with the
extrinsic manipulation of task constraints but also with the postural system's endogenous complexity. Indices of
postural sway governed the appearance or disappearance of crossovers, suggesting that tempering of non-
Gaussianity across log-timescale in stable condition is mostly a function of endogenous postural control. This
pliability of a crossover between inertial and feedback-based control could have life-and-death consequences for
healthy functioning (e.g., preventing falls). The present work suggests that an important direction for future
research would be using multiscale PDF as a lens on the healthy maintenance of upright standing, particularly as
a balance between endogenous control and exogenous constraints.

Supplementary materials

Supplementary Table S1. Coefficients of linear mixed-effect (LME) model examining the effects of Surface × Vision × Age-group on λ_s^2 vs. log-timescale curves.

Supplementary Table S2. Coefficients of generalized linear mixed-effect (GLME) model examining the effects of Surface × Vision × Age-group × Type on the growth of lognormality in postural sway with log-timescale.

Supplementary Table S3. Coefficients of LME model examining the effects of Surface × Vision × Age-group on λ_s^2 vs. log-timescale curves for orthogonal linear and quadratic polynomials, for interactions of grouping variables (Surface × Vision × Age-group × Type) and with indices of postural control.

Author contributions

M.M. and D.G.K-S. conceived and designed research; M.M. and D.G.K-S. analyzed data; M.M. and D.G.K-S. interpreted results of experiments; M.M. prepared figures; M.M. and D.G.K-S. drafted manuscript; M.M. and D.G.K-S. edited and revised manuscript; M.M. and D.G.K-S. approved final version of manuscript.

References

1. Collins JJ, De Luca CJ. 1993 Open-loop and closed-loop control of posture: A random-walk analysis of center-of-pressure trajectories. *Exp. Brain Res.* **95**, 308–318. (doi:10.1007/BF00229788)

2. Collins JJ, De Luca CJ. 1995 Upright, correlated random walks: A statistical-biomechanics approach to the human postural control system. *Chaos An Interdiscip. J. Nonlinear Sci.* **5**, 57–63. (doi:10.1063/1.166086)

3. Zatsiorsky VM, Duarte M. 2000 Rambling and trembling in quiet standing. *Motor Control* **4**, 185–200. (doi:10.1123/mcj.4.2.185)

4. Zatsiorsky VM, Duarte M. 1999 Instant equilibrium point and its migration in standing tasks: Rambling and trembling components of the stabilogram. *Motor Control* **3**, 28–38. (doi:10.1123/mcj.3.1.28)

5. Winter DA. 1995 Human balance and posture control during standing and walking. *Gait Posture* **3**, 193–214. (doi:10.1016/0966-6362(96)82849-9)

6. Winter DA, Patla AE, Ishac M, Gage WH. 2003 Motor mechanisms of balance during quiet standing. *J. Electromyogr. Kinesiol.* **13**, 49–56. (doi:10.1016/S1050-6411(02)00085-8)

7. Winter DA, Patla AE, Prince F, Ishac M, Gielo-Perczak K. 1998 Stiffness control of balance in quiet standing. *J. Neurophysiol.* **80**, 1211–1221. (doi:10.1152/jn.1998.80.3.1211)

8. Fitzpatrick R, Burke D, Gandevia SC. 1996 Loop gain of reflexes controlling human standing measured with the use of postural and vestibular disturbances. *J. Neurophysiol.* **76**, 3994–4008. (doi:10.1152/jn.1996.76.6.3994)

9. Bacsi AM, Colebatch JG. 2005 Evidence for reflex and perceptual vestibular contributions to postural control. *Exp. Brain Res.* **160**, 22–28. (doi:10.1007/s00221-004-1982-2)

10. Gantchev GN, Dimitrova DM. 1996 Anticipatory postural adjustments associated with arm movements during balancing on unstable support surface. *Int. J. Psychophysiol.* **22**, 117–122. (doi:10.1016/0167-8760(96)00016-5)

11.   Klous M, Mikulic P, Latash ML. 2011 Two aspects of feedforward postural control: Anticipatory postural
adjustments and anticipatory synergy adjustments. *J. Neurophysiol.* **105**, 2275–2288.
(doi:10.1152/jn.00665.2010)

12.   Klous M, Mikulic P, Latash ML. 2012 Early postural adjustments in preparation to whole-body voluntary
sway. *J. Electromyogr. Kinesiol.* **22**, 110–116. (doi:10.1016/j.jelekin.2011.11.005)

13.   Peterka RJ, Loughlin PJ. 2004 Dynamic regulation of sensorimotor integration in human postural control.
J. Neurophysiol. **91**, 410–423. (doi:10.1152/jn.00516.2003)

14.   Peterka RJ. 2002 Sensorimotor integration in human postural control. *J. Neurophysiol.* **88**, 1097–1118.
(doi:10.1152/jn.2002.88.3.1097)

15.   Carpenter MG, Murnaghan CD, Inglis JT. 2010 Shifting the balance: Evidence of an exploratory role for
postural sway. *Neuroscience* **171**, 196–204. (doi:10.1016/j.neuroscience.2010.08.030)

16.   Murnaghan CD, Horslen BC, Inglis JT, Carpenter MG. 2011 Exploratory behavior during stance persists
with visual feedback. *Neuroscience* **195**, 54–59. (doi:10.1016/j.neuroscience.2011.08.020)

17.   Haddad JM, Ryu JH, Seaman JM, Ponto KC. 2010 Time-to-contact measures capture modulations in
posture based on the precision demands of a manual task. *Gait Posture* **32**, 592–596.
(doi:10.1016/j.gaitpost.2010.08.008)

18.   Codling EA, Plank MJ, Benhamou S. 2008 Random walk models in biology. *J. R. Soc. Interface* **5**, 813–
834. (doi:10.1098/rsif.2008.0014)

19.   Liebovitch LS, Yang W. 1997 Transition from persistent to antipersistent correlation in biological
systems. *Phys. Rev. E* **56**, 4557–4566. (doi:10.1103/PhysRevE.56.4557)

20.   Holden JG, Van Orden GC, Turvey MT. 2009 Dispersion of response times reveals cognitive dynamics.
Psychol. Rev. **116**, 318–342. (doi:10.1037/a0014849)

21.   Stephen DG, Mirman D. 2010 Interactions dominate the dynamics of visual cognition. *Cognition* **115**,
154–165. (doi:10.1016/j.cognition.2009.12.010)

22.   Mandelbrot B. 1976 Intermittent turbulence and fractal dimension: Kurtosis and the spectral exponent
5/3+B. In *Turbulence and Navier Stokes Equations* (ed R Temam), pp. 121–145. Berlin, Heidelberg:
Springer Berlin Heidelberg. (doi:10.1007/BFb0091452)

23.   Ihlen EAF, Vereijken B. 2010 Interaction-dominant dynamics in human cognition: Beyond $1/f$
fluctuation. *J. Exp. Psychol. Gen.* **139**, 436–463. (doi:10.1037/a0019098)

24.   Kelty-Stephen DG, Wallot S. 2017 Multifractality versus (mono-) fractality as evidence of nonlinear
interactions across timescales: Disentangling the belief in nonlinearity from the diagnosis of nonlinearity
in empirical data. *Ecol. Psychol.* **29**, 259–299. (doi:10.1080/10407413.2017.1368355)

25.   Box GEP, Jenkins GM, Reinsel GC, Ljung GM. 2015 *Time Series Analysis: Forecasting and Control*.
Hoboken, NJ: John Wiley & Sons.

26.   Wagenmakers E-J, Farrell S, Ratcliff R. 2004 Estimation and interpretation of $1/f^{\alpha}$ noise in human
cognition. *Psychon. Bull. Rev.* **11**, 579–615. (doi:10.3758/BF03196615)

27.   Reynolds AM. 2018 Current status and future directions of Lévy walk research. *Biol. Open* **7**, bio030106.
(doi:10.1242/bio.030106)

28.   Broido AD, Clauset A. 2019 Scale-free networks are rare. *Nat. Commun.* **10**, 1017. (doi:10.1038/s41467-
019-08746-5)

29.   Virkar Y, Clauset A. 2014 Power-law distributions in binned empirical data. *Ann. Appl. Stat.* **8**, 89–119.

30.   Gheorghiu S, Coppens M-O. 2004 Heterogeneity explains features of "anomalous" thermodynamics and
statistics. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 15852–15856. (doi:10.1073/pnas.0407191101)

31.   Mangalam M, Chen R, McHugh TR, Singh T, Kelty-Stephen DG. 2020 Bodywide fluctuations support
manual exploration: Fractal fluctuations in posture predict perception of heaviness and length via
effortful touch by the hand. *Hum. Mov. Sci.* **69**, 102543. (doi:10.1016/j.humov.2019.102543)

32.   Schertzer D, Lovejoy S. 1993 Multifractal generation of self-organized criticality. In *Fractals in the
Natural and Applied Sciences*, pp. 325–339.

33.   Lovejoy S, Schertzer D. 2006 Multifractals, cloud radiances and rain. *J. Hydrol.* **322**, 59–88.
(doi:10.1016/j.jhydrol.2005.02.042)

34.   AghaKouchak A, Huning LS, Chiang F, Sadegh M, Vahedifard F, Mazdiyasni O, Moftakhari H,
Mallakpour I. 2018 How do natural hazards cascade to cause disasters? *Nature* **561**, 458–460.
(doi:10.1038/d41586-018-06783-6)

35.   Yaswen P, Campisi J. 2007 Oncogene-induced senescence pathways weave an intricate tapestry. *Cell* **128**,
233–234. (doi:10.1016/j.cell.2007.01.005)

36.   Van Breusegem F, Bailey-Serres J, Mittler R. 2008 Unraveling the tapestry of networks Involving
reactive oxygen species in plants. *Plant Physiol.* **147**, 978–984. (doi:10.1104/pp.108.122325)

37.   Garcia-Castro M, Kremer L, Reinkemeier CD, Unkelbach C, Strohmann C, Ziegler S, Ostermann C,
Kumar K. 2015 *De novo* branching cascades for structural and functional diversity in small molecules.
Nat. Commun. **6**, 6516. (doi:10.1038/ncomms7516)

38.   Liu W, Khedkar V, Baskar B, Schürmann M, Kumar K. 2011 Branching cascades: A concise synthetic
strategy targeting diverse and complex molecular frameworks. *Angew. Chemie Int. Ed.* **50**, 6900–6905.
(doi:10.1002/anie.201102440)

39.   Blaizot J-P, Iancu E, Mehtar-Tani Y. 2013 Medium-induced QCD cascade: Democratic branching and
wave turbulence. *Phys. Rev. Lett.* **111**, 52001. (doi:10.1103/PhysRevLett.111.052001)

40.   Jo E-K, Yang C-S, Choi CH, Harding C V. 2007 Intracellular signalling cascades regulating innate
immune responses to *Mycobacteria*: Branching out from toll-like receptors. *Cell. Microbiol.* **9**, 1087–
1098. (doi:10.1111/j.1462-5822.2007.00914.x)

41.   Patil NT, Shinde VS, Sridhar B. 2013 Relay catalytic branching cascade: A technique to access diverse
molecular scaffolds. *Angew. Chemie Int. Ed.* **52**, 2251–2255. (doi:10.1002/anie.201208738)

42.   Dunne JA, Williams RJ. 2009 Cascading extinctions and community collapse in model food webs.
Philos. Trans. R. Soc. B Biol. Sci. **364**, 1711–1723. (doi:10.1098/rstb.2008.0219)

43. Chauhan VK, Perera S, Brintrup A. 2020 The relationship between nested patterns and the ripple effect in
complex supply networks. *Int. J. Prod. Res.* , 1–17. (doi:10.1080/00207543.2020.1831096)

44. Bernier M, Fortin V, El-Amraoui M, Messaddeq Y, Vallée R. 2014 3.77 μm fiber laser based on cascaded
Raman gain in a chalcogenide glass fiber. *Opt. Lett.* **39**, 2052–2055. (doi:10.1364/OL.39.002052)

45. Li K. 2009 Pathology and radiology beyond looking at pictures. *Arch. Pathol. Lab. Med.* **133**, 587–590.
(doi:10.1043/1543-2165-133.4.587)

46. James DG. 1990 The many faces of sarcoidosis. The Thomé Villar memorial lecture. *Sarcoidosis* **7**, 1—8.

47. Schillemans M, Karampini E, Kat M, Bierings R. 2019 Exocytosis of Weibel–Palade bodies: How to
unpack a vascular emergency kit. *J. Thromb. Haemost.* **17**, 6–18. (doi:10.1111/jth.14322)

48. Schertzer D, Lovejoy S. 2004 Uncertainty and predictability in geophysics: Chaos and multifractal
insights. *State Planet Front. Challenges Geophys.* , 317–334. (doi:10.1029/150GM25)

49. Lovejoy S, del Rio Amador L, Hébert R. 2015 The Scaling Linear Macroweather Model (SLIM): Using
scaling to forecast global scale macroweather from months to decades. *Earth Syst. Dyn Disc* **6**, 489–545.

50. Mangalam M, Lee I-C, Newell KM, Kelty-Stephen DG. 2020 Visual effort moderates postural cascade
dynamics. *Neurosci. Lett.* **742**, 135511. (doi:10.1016/j.neulet.2020.135511)

51. Furmanek MP, Mangalam M, Kelty-Stephen DG, Juras G. 2020 Postural constraints recruit shorter-
timescale processes into the non-Gaussian cascade processes. *Neurosci. Lett.* **741**, 135508.
(doi:10.1016/j.neulet.2020.135508)

52. Kelty-Stephen DG, Furmanek MP, Mangalam M. 2020 Multifractality distinguishes reactive from
proactive cascades in postural control. *Chaos, Solitons & Fractals* **145**, 110471.
(doi:10.1016/j.chaos.2020.110471)

53. Castaing B, Gagne Y, Hopfinger EJ. 1990 Velocity probability density functions of high Reynolds
number turbulence. *Phys. D Nonlinear Phenom.* **46**, 177–200. (doi:10.1016/0167-2789(90)90035-N)

54. Kiyono K, Struzik ZR, Aoyagi N, Sakata S, Hayano J, Yamamoto Y. 2004 Critical scale invariance in a
healthy human heart rate. *Phys. Rev. Lett.* **93**, 178103. (doi:10.1103/PhysRevLett.93.178103)

55. Kiyono K, Hayano J, Watanabe E, Struzik ZR, Yamamoto Y. 2008 Non-Gaussian heart rate as an
independent predictor of mortality in patients with chronic heart failure. *Hear. Rhythm* **5**, 261–268.
(doi:10.1016/j.hrthm.2007.10.030)

56. Shlesinger MF, West BJ, Klafter J. 1987 Lévy dynamics of enhanced diffusion: Application to turbulence.
Phys. Rev. Lett. **58**, 1100–1103. (doi:10.1103/PhysRevLett.58.1100)

57. Kiyono K. 2009 Log-amplitude statistics of intermittent and non-Gaussian time series. *Phys. Rev. E* **79**,
31129. (doi:10.1103/PhysRevE.79.031129)

58. Kiyono K, Struzik ZR, Aoyagi N, Yamamoto Y. 2006 Multiscale probability density function analysis:
Non-Gaussian and scale-Invariant fluctuations of healthy human heart rate. *IEEE Trans. Biomed. Eng.*
53, 95–102. (doi:10.1109/TBME.2005.859804)

59. Kiyono K, Hayano J, Kwak S, Watanabe E, Yamamoto Y. 2012 Non-gaussianity of low frequency heart
rate variability and sympathetic activation: Lack of increases in multiple system atrophy and parkinson
disease. *Front. Physiol.* **3**, 34. (doi:10.3389/fphys.2012.00034)

60. Hayano J, Kiyono K, Struzik Z, Yamamoto Y, Watanabe E, Stein P, Watkins L, Blumenthal J, Carney R.
2011 Increased non-Gaussianity of heart rate variability predicts cardiac mortality after an acute
myocardial infarction. *Front. Physiol.* **2**, 65. (doi:10.3389/fphys.2011.00065)

61. Safonov LA, Isomura Y, Kang S, Struzik ZR, Fukai T, Câteau H. 2010 Near scale-free dynamics in neural
population activity of waking/sleeping rats revealed by multiscale analysis. *PLoS One* **5**, e12869.
(doi:10.1371/journal.pone.0012869)

62. West BJ, Turalska M. 2019 Hypothetical control of heart rate variability. *Front. Physiol.* **10**, 1078.
(doi:10.3389/fphys.2019.01078)

63. dos Santos DA, Fukuchi CA, Fukuchi RK, Duarte M. 2017 A data set with kinematic and ground reaction
forces of human balance. *PeerJ* **5**, e3626. (doi:10.7717/peerj.3626)

64. Schreiber T, Schmitz A. 1996 Improved surrogate data for nonlinearity tests. *Phys. Rev. Lett.* **77**, 635–
638. (doi:10.1103/PhysRevLett.77.635)

65. Hansson EE, Beckman A, Håkansson A. 2010 Effect of vision, proprioception, and the position of the
vestibular organ on postural sway. *Acta Otolaryngol.* **130**, 1358–1363.
(doi:10.3109/00016489.2010.498024)

66. Teasdale N, Stelmach GE, Breunig A. 1991 Postural sway characteristics of the elderly under normal and
altered visual and support surface conditions. *J. Gerontol.* **46**, B238–B244.
(doi:10.1093/geronj/46.6.B238)

67. Le Clair K, Riach C. 1996 Postural stability measures: What to measure and for how long. *Clin. Biomech.*
11, 176–178. (doi:10.1016/0268-0033(95)00027-5)

68. Richman JS, Moorman JR. 2000 Physiological time-series analysis using approximate entropy and
sample entropy. *Am. J. Physiol. Circ. Physiol.* **278**, H2039–H2049.
(doi:10.1152/ajpheart.2000.278.6.H2039)

69. Ko J-H, Newell KM. 2016 Aging and the complexity of center of pressure in static and dynamic postural
tasks. *Neurosci. Lett.* **610**, 104–109. (doi:10.1016/j.neulet.2015.10.069)

70. Costa M, Goldberger AL, Peng C-K. 2002 Multiscale entropy analysis of complex physiologic time
series. *Phys. Rev. Lett.* **89**, 68102. (doi:10.1103/PhysRevLett.89.068102)

71. Riley MA, Mitra S, Stoffregen TA, Turvey MT. 1997 Influences of body lean and vision on unperturbed
postural sway. *Motor Control* **1**, 229–246. (doi:10.1123/mcj.1.3.229)

72. Vaillancourt DE, Sosnoff JJ, Newell KM. 2004 Age-related changes in complexity depend on task
dynamics. *J. Appl. Physiol.* **97**, 454–455. (doi:10.1152/japplphysiol.00244.2004)

73. Peng C-K, Buldyrev S V, Havlin S, Simons M, Stanley HE, Goldberger AL. 1994 Mosaic organization of
DNA nucleotides. *Phys. Rev. E* **49**, 1685–1689. (doi:10.1103/PhysRevE.49.1685)

74. Chhabra A, Jensen R V. 1989 Direct determination of the $f(\alpha)$ singularity spectrum. *Phys. Rev. Lett.* **62**,
1327–1330. (doi:10.1103/PhysRevLett.62.1327)

75. Mangalam M, Kelty-Stephen DG. 2020 Multiplicative-cascade dynamics supports whole-body
coordination for perception via effortful touch. *Hum. Mov. Sci.* **70**, 102595.
(doi:10.1016/j.humov.2020.102595)

76. Amoud H, Abadi M, Hewson DJ, Michel-Pellegrino V, Doussot M, Duchêne J. 2007 Fractal time series
analysis of postural stability in elderly and control subjects. *J. Neuroeng. Rehabil.* **4**, 12.
(doi:10.1186/1743-0003-4-12)

77. Duarte M, Sternad D. 2008 Complexity of human postural control in young and older adults during
prolonged standing. *Exp. Brain Res.* **191**, 265–276. (doi:10.1007/s00221-008-1521-7)

78. Shimizu YU, Thurner S, Ehrenberger K. 2002 Multifractal spectra as a measure of complexity in human
posture. *Fractals* **10**, 103–116. (doi:10.1142/S0218348X02001130)

79. Kiyono K, Struzik ZR, Yamamoto Y. 2007 Estimator of a non-Gaussian parameter in multiplicative log-
normal models. *Phys. Rev. E* **76**, 41113. (doi:10.1103/PhysRevE.76.041113)

80. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting linear mixed-effects models using lme4. *J. Stat.*
Softw. **1**, 1–48. (doi:10.18637/jss.v067.i01)

81. Team RC. 2013 R: A language and environment for statistical computing. (doi:URL https://www.R-
project.org/.)

82. Kuznetsova A, Brockhoff PB, Christensen RHB. 2017 lmerTest package: Tests in linear mixed effects
models. *J. Stat. Softw.* **82**, 1548–7660. (doi:/10.18637/jss.v082.i13)

83. Federolf P, Zandiyeh P, von Tscharner V. 2015 Time scale dependence of the center of pressure entropy:
What characteristics of the neuromuscular postural control system influence stabilographic entropic half-
life? *Exp. Brain Res.* **233**, 3507–3515. (doi:10.1007/s00221-015-4420-8)

84. Delignières D, Torre K, Bernard P-L. 2011 Transition from persistent to anti-persistent correlations in
postural sway indicates velocity-based control. *PLOS Comput. Biol.* **7**, e1001089.
(doi:10.1371/journal.pcbi.1001089)

85. Chagdes JR, Rietdyk S, Haddad JM, Zelaznik HN, Raman A, Rhea CK, Silver TA. 2009 Multiple
timescales in postural dynamics associated with vision and a secondary task are revealed by wavelet
analysis. *Exp. Brain Res.* **197**, 297–310. (doi:10.1007/s00221-009-1915-1)

86. Lee I-C, Pacheco MM, Newell KM. 2018 Constraints specific influences of vision, touch and surface
compliance in postural dynamics. *Gait Posture* **59**, 117–121. (doi:10.1016/j.gaitpost.2017.09.014)

87. Newell KM, Slobounov SM, Slobounova ES, Molenaar PCM. 1997 Stochastic processes in postural
center-of-pressure profiles. *Exp. Brain Res.* **113**, 158–164. (doi:10.1007/BF02454152)

88. Duarte M, Zatsiorsky VM. 2000 On the fractal properties of natural human standing. *Neurosci. Lett.* **283**,
173–176. (doi:10.1016/S0304-3940(00)00960-5)

Figure 1. Schematic illustration of multiscale probability density function (PDF) analysis. (*a*) The first step is to
generate $\{B(t)\}$ by integrating $\{b(t)\}$ after centering by the mean b_{ave}: $B(t)=\sum_{i=1}^{t/\Delta t} b(i\,\Delta t)-b_{ave}$. (*b*) A a 3$^{\text{rd}}$ order
polynomial detrends $\{B(t)\}$ within k overlapping windows of length $2\,s$, s being the timescale. (*c*) Intermittent
deviation $\Delta_s B(t)$ in k^{th} window from $1+s(k-1)$ to sk in the detrended time series $\{B^d(t)=B(t)-f_{fit}(t)\}$ is
computed as $\Delta_s B^{\square}(t)=B^{\square}(t+s)-B^{\square}(t)$, where $1+s(k-1)\leq t\leq sk$ and $f_{fit}(t)$ is the polynomial representing the
local trend of $\{B(t)\}$, of which the elimination assures the zero-mean probability density function in the next
step.

Figure 2. Schematic illustration of non-Gaussianity index λ_s^2. (*a*) The relationship between λ_s^2 and shapes of PDF
plotted in linear-log coordinates for $\lambda_s^2 = 0.9$, 0.33, and 0.13 (from top to bottom). For clarity, the PDFs have
been shifted vertically. As λ_s^2 increases, the PDF becomes increasingly peaked and fat-tailed. As λ_s decreases,
the PDF increasingly resembles the Gaussian (dashed line), assuming a perfect Gaussian as λ_s^2 approaches 0. (*b*)
In a cascade-type multiplicative process yield the inverse relationship $\lambda_s^2 \sim -\log s$.

Figure 3. Multiscale PDF characterization of postural sway in representative young and old participants standing quietly for 60 s on stable and unstable surfaces with eyes open. From top to bottom: CoP trajectories along the anterior-posterior (*AP*) and medial-lateral (*ML*) axes. CoP PED series. $\{\Delta_s B(i)\}$ for s = 0.5, 2, and 8 s. Standardized PDFs (in logarithmic scale) of $\{\Delta_s B(i)\}$ for s = 0.5, 2, and 8 s (from top to bottom), where σ_s denotes the *SD* of $\{\Delta_s B(i)\}$.

Figure 4. Multiscale PDF characterization of postural sway in representative young and old participants standing quietly for 60 s on stable and unstable surfaces with eyes closed. From top to bottom: CoP trajectories along the anterior-posterior (*AP*) and medial-lateral (*ML*) axes. CoP PED series. $\{\Delta_s B(i)\}$ for $s = 0.5$, 2, and 8 s. Standardized PDFs (in logarithmic scale) of $\{\Delta_s B(i)\}$ for $s = 0.5$, 2, and 8 s (from top to bottom), where σ_s denotes the *SD* of $\{\Delta_s B(i)\}$.

Figure 5. Log-timescale dependence of the non-Gaussianity index λ_s^2 for the young and old participants standing quietly for 60 s. (*a*) Stable surface, eyes open. (*b*) Stable surface, eyes closed. (*c*) Unstable surface, eyes open. (*d*) Unstable surface, eyes closed. Vertical bars indicate $\pm 1 s.e.m.$ of the group averages (young, $n = 27$; old, $n = 22$).

Figure 6. Log-timescale dependence of the mean proportion of trials with a lognormal distribution for the young and old participants standing quietly for 60 s. (*a*) Stable surface, eyes open. (*b*) Stable surface, eyes closed. (*c*) Unstable surface, eyes open. (*d*) Unstable surface, eyes closed. Vertical bars indicate ±1*s.e.m.* of the group averages (young, $n = 27$; old, $n = 22$).