Time course of alterations in adult spinal motoneuron properties in the SOD1(G93A) mouse model of ALS

Seoan Huh[1], Charles J. Heckman[1,2,3], and Marin Manuel[1,4,§]

[1]Department of Physiology, [2]Department of Physical Medicine and Rehabilitation, and [3]Department of Physical Therapy and Human Movement Science, Northwestern University, Feinberg School of Medicine, Chicago, Illinois, USA. [4]Université de Paris, SPPIN - Saints-Pères Paris Institute for the Neurosciences, CNRS, Paris, France.

§ Corresponding author:

Marin MANUEL
SPPIN - Saints-Pères Paris Institute for the Neurosciences
Université de Paris, 45, rue des Saints-Pères, 75006 Paris, France
email: marin.manuel@neurobio.org

Author contribution

Conceptualization: S.H., C.J.H., and M.M.; Methodology: S.H., C.J.H., and M.M.; Investigation: S.H.; Formal analysis: S.H. and M.M.; Software: M.M.; Visualization: S.H. and M.M.; Writing - Original Draft: S.H.; Writing - Review & Editing: S.H., C.J.H., and M.M.; Funding Acquisition: C.J.H and M.M.; S.H., C.J.H., and M.M. approved the final version of this manuscript.

Number of pages: 33

Number of figures: 7 + 1 table

Number of words for abstract: 149

Number of words for Introduction: 514

Number of words for Discussion: 1860

Conflict of Interest

The authors have no conflict of interest to declare

Acknowledgements

This work was supported by NIH NINDS R01NS077863 and NINDS R01NS110953. S.H. was supported by a National Science Foundation Graduate Research Fellowship (NSF GRFP).

Abstract

Although ALS is an adult-onset neurodegenerative disease, motoneuron electrical

properties are already altered during embryonic development. Motoneurons must

therefore exhibit remarkable capacity for homeostatic regulation to maintain a

normal motor output for most of the life of the patient. In the present paper, we

demonstrate how maintaining homeostasis could come at a very high cost. We

studied the excitability of spinal motoneurons from young adult SOD1(G93A) mice

to end-stage. Initially homeostasis is highly successful in maintaining their overall

excitability. This initial success, however, is achieved by pushing some cells far

above the normal range of passive and active conductances. As the disease

progresses, both passive and active conductances shrink below normal values in the

surviving cells. This shrinkage may thus promote survival, implying the previously

large values contribute to degeneration. These results support the hypothesis that

motoneuronal homeostasis may be "hyper-vigilant" in ALS, and a source of

accumulating stress.

Impact statement

51   During ALS, motoneurons attempt to maintain a normal motor output by altering

52   their electrical properties, sometimes beyond the normal range, which may be

53   causing long-lasting harm.

Introduction

ALS is a fatal neurodegenerative disorder characterized by progressive loss of cortical and spinal motoneurons. Five to ten percent of the ALS cases are familial (FALS), and ~20% of the FALS cases are due to mutations in the Superoxide Dismutase1 (SOD1) gene (Rosen et al., 1993). Studies in the mutant SOD1 (mSOD1) mouse and other animal models have revealed that multiple cellular functions become impaired as the disease progresses (Ilieva et al., 2009; Kanning et al., 2009; Johnson and Heckman, 2010; Brownstone and Lancelin, 2018), implying that homeostatic mechanisms are failing. Yet, the intrinsic excitability of neonatal and young adult mSOD1 motoneurons undergoes a very potent and effective homeostatic response to compensate an earlier surge in excitability in the embryonic state (Kuo et al., 2004; van Zundert et al., 2008; Pieri et al., 2009; Martin et al., 2013). The mechanisms of this successful homeostasis are however highly unusual.

The standard measure of the net excitability of motoneurons is the relation between frequency of firing of spikes and amplitude of injected current. This frequency-current (F-I) function is determined by multiple ionic currents. Remarkably, the successful F-I homeostasis in the neonatal state is achieved by abnormal but compensatory increases in these currents. Persistent inward currents (PICs), which are mediated by voltage gated Na^+ and Ca^{2+} channels, become aberrantly large. On its own, this change would increase F-I gain, but it is compensated by large increases in the leak currents that set the input conductance

of the cell. These abnormal changes show that F-I homeostasis during the neonatal

period is achieved via a compensatory mechanism. If, however, these compensatory

increases persist into the young adult state, then the continued distortions in input

conductances and PIC amplitudes could induce substantial stresses within the

motoneurons. In this scenario, the initial success in homeostatic regulation of net

excitability would be coupled to a severe cost, likely accelerating the onset of

degeneration. On the other hand, if homeostatic processes instead successfully

return input conductance and PIC values to normal ranges before the onset of

denervation, then homeostasis for the F-I function would reduce stress on the cell

and effects on subsequent degeneration would likely be small.

To investigate how homeostasis for excitability develops as the disease

progresses, we undertook the first *in vivo* voltage clamp studies of motoneurons in

the mouse and pursued these measurements across a wide range of ages, from P30

to P120. This age range spans the young adult period, the onset of denervation

period (∼P50, Pun et al., 2006) and the development of overt symptoms (∼P90).

We hypothesized that input conductance and PIC values would continue to grow in

the young adult state. Our results supported this hypothesis, revealing in fact a

continual increase in amplitude of these parameters, followed by a collapse, so that

motoneurons surviving beyond onset of overt symptoms (∼P90) had aberrantly

small values of each. These results are consistent with the possibility that

homeostasis for excitability is not in fact weak but excessively strong and that this

overreaction contributes to subsequent degeneration (Mitchell and Lee, 2012).

Methods

Animals

This study was performed in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. All of the animals were handled according to protocols approved by Northwestern University's institutional animal care and use committee (IACUC). All surgery was performed under sodium pentobarbital anesthesia, and every effort was made to minimize suffering. Because of the reproducible and stereotyped progression of symptoms, we chose to use mice overexpressing the human SOD1(G93A) gene, in which glycine has been substituted by alanine at residue 93 as a mouse model of ALS. All mice had a B6SJL background (Jackson Lab, Bar Harbor, ME). The transgene copy number was compared to a housekeeping gene at Transnetyx (Cordova, TN), and only mice with relative copy number >45 were used in this study. Control group consisted of non-transgenic littermates, with the same B6SJL background (WT). 33 animals of either sex were used in this study, and they were divided into different age groups for analysis: P30–60 (N=9 mice), P60–90 (N=12 mice), and P90–120 (N=12 mice).

In vivo preparation

Procedures were similar to those in our previous studies (Manuel and Heckman, 2011). Initially, atropine (0.2mg/kg) was administered subcutaneously to prevent salivation. Ten minutes later, anesthesia was initiated with an intraperitoneal injection of pentobarbital sodium (70 mg/kg) and maintained by intravenous infusion of supplemental doses of pentobarbital (6 mg/kg) mixed in perfusion

solution containing 4% glucose, 1% $NaHCO_3$, and 14% Plasmion. The trachea was

cannulated, allowing the mouse to be artificially ventilated with 100% O_2. The end-

tidal PCO_2 was continuously monitored and maintained around 4% by adjusting

ventilator parameters including respiratory rate around 120–150 bpm and tidal

volume around 0.09–0.23 mL. The hind limb muscles were then dissected; biceps

femoris was removed to expose the sciatic nerve, which was thereafter placed over

a stimulating electrode. A laminectomy was performed at T13–L1 level, and L3–L4

spinal segments were exposed. To prevent the spinal cord from dehydration, a

custom made bath was affixed using silicone elastomer and covered with mineral

oil. To locate the motoneurons of interest, the sciatic nerve was stimulated at

intensity 1.8× threshold for the apparition of an afferent volley.

Electrophysiology

Intracellular recordings of spinal motoneurons were performed by impaling them

with glass micropipette electrodes filled with 3M KCl with resistance of 8–15 MΩ.

Motoneurons were identified by the presence of antidromic action potential from

stimulation of the sciatic nerve. Cells with unstable resting membrane potential

(RMP) or with RMP more depolarized than −50mV were excluded from analysis.

PICs were recorded in discontinuous voltage clamp mode, with switching rates of

6–8 kHz. Clamp feedback gain was between 0.3 and 1.5. In addition to the feedback

gain from Axoclamp amplifier, an additional low-frequency feedback loop with a

gain of 11 and a cutoff of −3 dB at 0.3 kHz was used to improve voltage control in

such large cells as motoneurons (Lee and Heckman, 1998). Outputs were observed

at all times to assure reasonable settling of the electrode. To record PICs, a slow

triangular voltage ramp (−80 to −40 mV) was applied (Lee and Heckman, 1998).

Leak current was determined by fitting a regression line through the subthreshold

region of the I-V function. Then, this leak was subtracted from the total function to

determine the PIC amplitude (measured both on the ascending and descending part

of the ramp). In addition, the voltage at which the PIC was maximal on each part of

the ramp was recorded (PIC peak voltage), as well as the voltage at which the

current trace deviated from the leak current (PIC onset voltage) on the ascending

ramp, and joined back the leak trace on the descending ramp (PIC end voltage).

Input conductance was estimated as the slope of the leak current.

To assess intrinsic properties of the motoneuron in current clamp mode, we

measured the frequency-current (F-I) relationship of action potential firing. The F-I

relationship was determined based on the firing produced by a triangular current

injection. The interspike frequency was then plotted against the intensity of the

injected current. All these measurements were carried out in the discontinuous

current clamp mode of the Axoclamp 2A amplifier, with switching rates of 6–8 kHz.

Mouse motoneurons possess two regimes of firing: a sub-primary range of firing,

followed by a linear primary range (Manuel et al., 2009; Manuel and Heckman,

2011). The "gain" of the F-I relationship was determined by fitting a regression line

over the linear primary range (separately on the ascending and descending phase of

the ramp). The other parameters measured in the F-I relationship were: the current

at which the first action potential (AP) fires on the ascending ramp (recruitment

current). the current at which the last AP fires on the descending ramp (current at

de-recruitment); the voltage threshold for spiking, which was determined as the

voltage where the slope of the membrane voltage reaches 10mV/s before the first

AP (Sekerli et al., 2004); the current and firing frequency at the transition between

the sub-primary and primary ranges (Manuel and Heckman, 2011).

Data analysis

171   Recordings were acquired and analyzed using Spike2 v.7 (CED, Cambridge, UK).

172   Data were collected and analyzed using Pandas v.1.0.0 (McKinney, 2011), SciPy

173   Python v.1.4.1 (Virtanen et al., 2020) and DABEST v.0.3.0 (Ho et al., 2019). Figures

174   were generated using matplotlib v.3.1.3 (Hunter, 2007) and seaborn v.0.10.0

175   (Waskom et al., 2020).

Statistical analysis

All data are reported as mean ± standard deviation with their respective sample

size. No test was performed to detect outliers, and no data points were excluded

from analysis. When comparing between WT and mSOD1 samples, we focus on

estimation statistics that rely on effect sizes and confidence intervals (95%CI),

rather than null hypothesis significance testing, as recommended by several

scientific societies and editorial boards (Bernard, 2019; Makin and Orban de Xivry,

2019; Wasserstein et al., 2019; Michel et al., 2020). Unless otherwise specified, effect

sizes are reported as difference of means and Hedges' g (Hedges, 2016), as

calculated using DABEST v.0.3.0. (Ho et al., 2019). Where appropriate, data are

presented as Cumming plots (Cumming, 2012) generated using DABEST. In these

plots, the raw data are plotted as swarms of points. In addition, the mean ± standard

deviation of each group is plotted as a notched line immediately to the right of each

group. The effect size and bootstrapped 95% CIs are plotted on a separate axes

beneath the raw data. Confidence intervals were bias-corrected and accelerated, and

are displayed with the bootstrap distribution of the mean; resampling was

performed 5000 times (Ho et al., 2019). Welch's t-test (Welch, 1947) results are

provided for information only.

Results

Motoneurons from young adult mutant mice have overly large PICs

196
Figure 1: PIC amplitude is larger is young adult mSOD1 mice
A. Example of a PIC recording from a P43 WT mouse. Top trace: recorded current, bottom trace: voltage command. The red dashed line shows the leak current estimated by fitting a straight line around the point of zero current, and which is used to measure the input conductance of the cell. Below is a plot of the leak-subtracted current vs. the voltage command color coded for the ascending phase (thick red line) and the descending phase (thin blue line). The amplitude of the PIC is measured at the point indicated by the double arrow. Note that, in this example, a breakaway spike was filtered out. B. Example of a PIC recording from a P62 mSOD1 mouse. Same organization as in A. C. Plot of the amplitude of the PICs (in nA) vs. the age of the mouse in WT (blue square) and mSOD1 mice (red diamonds). The solid

We set out to measure persistent inward currents (PICs) in 103 motoneurons

from SOD1(G93A) mice (mSOD1), and their non-transgenic (WT) littermate, aged

between 31 and 123 days old. PICs were measured in response to a slow (5 mV/s)

ramp in voltage-clamp mode. The amplitude of the PIC was measured on the leak-

subtracted trace at the point where the downward deflection of the trace was

maximal (see Figure 1). In WT animals, the PIC amplitude remained roughly steady

during the age period studied (Figure 1C). However, in mSOD1 motoneurons, PIC

amplitudes decreased strikingly over time (slope=−0.45 nA/week, r=−0.54

95%CI[−0.70−−0.31], p=3.4e-5). When broken down by age groups (Figure 1D), our

results shows that, in the young adult stage (P30–60), when animals are pre-

symptomatic and neuromuscular-junction denervation is just starting, SOD1

mutation lead to an almost 3× increase in PIC amplitudes compared to WT

motoneurons. This increase appears to be a continuation of the trend observed in

neonates, where PICs were already increased ～2-fold compared to controls

(Quinlan et al., 2011). In the presymptomatic group (P60–90), however, this effect

became negligible despite the fact that substantial denervation is already underway.

At symptomatic stages (P90–120), the trend seen in young animals is reversed: PICs

are much smaller in mSOD1 motoneurons compared to WT motoneurons. Similar

results could be observed on the PIC amplitudes measured on the descending ramp

(Table 1).

In addition to its effect on PIC amplitude, SOD1 mutation also affected the voltage

at which PICs are recruited. The "PIC onset voltage", measured as the voltage at the

point where the leak-subtracted motoneuron I-V curve initially begins to curve

downward, was hyperpolarized by 10 mV in young mSOD1 mice compared to WT

controls (WT: −58.03±6.35mV, N=19 vs. mSOD1: −69.41±5.71mV, N=15; g=−1.83

95%CI[−2.64−−0.78]; t(32)=5.49, p=5.1e−06). The SOD1 mutation had a similarly

large effect on the voltage at which the PIC reached its maximum in young animals

(WT: −43.83±4.79mV, N=19 vs. SOD1: −52.32±4.20mV, N=15; g=−1.83

95%CI[−2.63−−1.04]; t(32)=5.50, p=4.8e−06). Similarly to the situation with PIC

amplitude, the effect of the SOD1 mutation on its activation voltage was reduced in

older animals, at both presymptomatic and symptomatic ages (Table 1).

These profound changes in PIC amplitude and activation voltage would be

expected to have an impact on the firing properties of motoneurons. We therefore

performed current-clamp recordings in the same motoneurons as above. We

injected a triangular ramp of current to elicit the repetitive firing of the motoneuron

(Figure 2A−B), and measured the voltage threshold for spiking on the first spike

triggered by the ramp. In concordance with the hyperpolarization of the PICs,

spiking threshold was also hyperpolarized in young animals (WT: −45.35±5.48mV,

N=19 vs. mSOD1: −54.86±5.15mV, N=14; g=−1.74 95%CI[−2.53−−0.81]; t(31)=5.10,

p=1.9e−05).

Young adult motoneurons are nonetheless not hyperexcitable

258
Figure 2. Mutant motoneurons are not hyperexcitable
A. Example of the response of a P101 WT mouse to a triangular ramp of current.
From bottom to top, traces are: injected current, membrane potential and
instantaneous firing frequency. The dash-dotted line shows the measure of the
recruitment current, the current intensity at which the motoneuron starts to fire.
The red dashed line shows the linear fit of the firing frequency in the primary range,
used to estimate the gain of the F-I relationship. B. Example of the response of a P60
mSOD1 mouse. Same organization as in A. C. Current intensity required for eliciting

the first spike on a triangular ramp of current in WT and mSOD1 motoneurons in each of the age groups. In young adult and presymptomatic mice, mutant motoneurons require the same amount of current than WT motoneuron to fire, P30–60 WT: 4.73±3.14nA, N=19 vs. mSOD1: 6.21±2.51nA, N=14; g=0.50 95%CI[−0.24–1.24]; t(31)=−1.51, p=0.14. P60–90 WT: 5.28±2.64nA, N=16 vs. mSOD1: 4.53±3.38nA, N=20; g=−0.24 95%CI[−0.93–0.41]; t(34)=0.75, p=0.46. At the symptomatic stages (P90–120), mutant motoneurons exhibit a lower current threshold for firing (WT: 5.43±3.07nA, N=15 vs. mSOD1: 2.19±1.56nA, N=13; g=−1.26 95%CI[−1.95−−0.47]; t(26)=3.59, p=0.0017), compatible with the loss of the least excitable cells. D. The slope of the F-I relationship, measured over the primary range, is not affected by the mutation. P30–60 WT: 9.3±3.6Hz/nA, N=10 vs. mSOD1: 8.3±3.4Hz/nA, N=14; g=−0.28 95%CI[−1.10–0.64]; t(22)=0.69, p=0.5. P60–90 WT: 12.2±7.3Hz/nA, N=8 vs. mSOD1: 10.3±6.4Hz/nA, N=16; g=−0.27 95%CI[−1.22–0.54]; t(22)=0.62, p=0.54. P90–120 WT: 8.0±3.2Hz/nA, N=10 vs. mSOD1: 10.8±9.1Hz/nA, N=13; g=0.38 95%CI[−0.43–0.93]; t(21)=−1.04, p=0.31.

Despite the large alterations in PIC amplitude and activation voltage, and despite the change in the voltage threshold for spiking, the excitability of the motoneurons was remarkably unaffected by the SOD1 mutation, regardless of age. Figure 2 shows the response of WT and mSOD1 young adult motoneurons to a slow triangular ramps of current (F-I relationship). We measured the intensity of the current required to elicit the first spike on the ascending ramp ("recruitment current"). Young adult (P30–60) mutant motoneurons, whose neuromuscular junctions are just starting to be denervated, and pre-symptomatic motoneurons (P60–90), which experience substantial denervation, did not require, on average, less current to reach firing threshold than WT controls (Figure 2C). At symptomatic stages (P90–120), the SOD1 mutation does lead to a decrease in the recruitment current compared to WT animals (Figure 2C), which is probably caused by the degeneration of the high threshold motoneurons at this stage.

As shown previously (Manuel et al., 2009), most F-I curves, regardless of age and

genotype, showed a distinct sub-primary range (SPR) with a steep slope followed by

a linear phase called primary range (PR). We used the "gain" of the motoneuron, i.e.

the slope of the F-I curve in the PR, as another measure of excitability of

motoneurons. Despite the alterations in PICs in mutant motoneurons, the gain of the

motoneuron F-I curves were unaffected by both age and mutation (Figure 2D).

Compensatory changes responsible to maintain excitability

302
Figure 3: Young mutant motoneurons have an aberrantly large input
conductance
A. Plot of the motoneuron input conductance vs. age in WT (blue squares) and
mSOD1 (red diamonds) animals. The solid lines correspond to the linear regression
lines with 95% confidence intervals (shaded areas). Pearson correlation coefficients
& 95%CI, WT: r=0.40 95%CI[0.14–0.61], p=0.0035, mSOD1: r=−0.73 95%CI[−0.83–
−0.57], p=6.1e−10. B. Breakdown of the difference in input conductance between
WT and mSOD1 animals by age groups. P30–60 WT: 0.42±0.16μS, N=19 vs. mSOD1:
0.72±0.18μS, N=15; g=1.73 95%CI[0.92–2.41]; t(32)=−5.06, p=2.2e−05. P60–90 WT:
0.46±0.16μS, N=16 vs. mSOD1: 0.42±0.15μS, N=24; g=−0.28 95%CI[−0.96–0.37];
t(38)=0.88, p=0.39. P90–120 WT: 0.60±0.15μS, N=15 vs. mSOD1: 0.33±0.13μS,
N=14; g=−1.85 95%CI[−2.57−−1.16]; t(27)=5.16, p=2.1e−05.

The fact that young adult motoneurons are not hyperexcitable despite substantial

alteration in the PICs suggests that other compensatory mechanisms are acting to

preserve the functional output of the cells. Motoneuron input conductance is an

important factor controlling neuron excitability. We estimated the input

conductance of mSOD1 and WT motoneurons from the slopes of their I-V relations

around the level of the resting membrane potential (see Figure 1 and Methods).

Input conductances of mSOD1 and WT motoneurons were plotted against the age of

the animal in Figure 3A. Consistent with the continuous increase in size of the

animals over the age span studied, WT motoneurons exhibit a moderate increase in

input conductance in WT motoneurons over time (slope=0.02 µS/week). On the

other hand, mSOD1 motoneurons showed the opposite trend (slope=−0.04

µS/week). Young adult mutant motoneurons had an input conductance almost twice

as high as WT controls at P30–P60 (Figure 3B). This increase is relatively greater (

～1.7 fold) than that observed in neonatal motoneurons at P0–7 (～1.25 fold)

(Quinlan et al., 2011), suggesting that the trend for increased conductance has

become stronger as the animal matures into the young adult state. However, by the

late pre-symptomatic stage (P60–90), the mutation had no longer any effect on

input conductance. Finally, symptomatic (P90–120) mSOD1 motoneurons had a

smaller input conductance than WT controls, but this difference could be due to the

degeneration of the largest, high-threshold units at this stage.

335
Figure 4. The resting membrane potential of young adult mutant mice is
hyperpolarized
Breakdown of the difference in input conductance between WT and mSOD1 animals
by age groups. P30–60 WT: −62.38±5.52mV, N=19 vs. mSOD1: −71.45±8.78mV,
N=15; g=−1.24 95%CI[−2.13−−0.23]; $t(32)$=3.49, p=0.002. P60–90 WT:
−60.98±5.35mV, N=16 vs. mSOD1: −66.18±5.63mV, N=22; g=−0.92 95%CI[−1.54−
−0.18]; $t(36)$=2.89, p=0.0067. P90–120 WT: −64.71±7.42mV, N=15 vs. mSOD1:
−61.99±5.43mV, N=13; g=0.40 95%CI[−0.35−1.10]; $t(26)$=−1.12, p=0.27.

The increase in input conductance is not the only compensatory change

happening in young adult mutant motoneurons. In particular, one would have

expected the leftward shift in the activation voltage of PICs seen in these

motoneurons to have a profound impact on their firing behavior. This discrepancy

can be explained by a parallel almost 10 mV hyperpolarization of the resting

membrane potential in young adult mSOD1 motoneurons (Figure 4, P30–60). This

effect is still observable at pre-symptomatic stages, but to a lesser degree (Figure 4,

P60–90). At symptomatic ages, however, mutant motoneurons had similar resting

membrane potentials as WT controls (Figure 4, P90–120).

When taking into account this hyperpolarization of the resting membrane

potential, the relative values of the voltage threshold for spiking (ΔV_{th}, difference

between the voltage threshold and the resting membrane potential), as well as the

relative activation voltage of the PICs (ΔV_{PIC}) and the relative voltage at the peak of

the PIC (ΔV_{peak}) were all similar between WT and mSOD1 motoneuron regardless of

age (Table 1), which suggest that these are the quantities that are homeostatically

regulated.

Arms race between PICs and input conductance

361
Figure 5. Some motoneurons exhibit properties outside of the normal range
A. Breakdown of the difference in normalized PIC amplitude (PIC amplitude divided
by input conductance) between WT and mSOD1 animals by age groups. P30–60 WT:
5.35±4.11nA/μS, N=19 vs. mSOD1: 9.32±3.94nA/μS, N=15; g=0.96 95%CI[0.16–
1.76]; t(32)=−2.86, p=0.0075. P60–90 WT: 5.65±3.35nA/μS, N=16 vs. mSOD1:
6.46±5.39nA/μS, N=24; g=0.17 95%CI[−0.50–0.66]; t(38)=−0.59, p=0.56. P90–120

WT: 6.80±5.01nA/μS, N=15 vs. mSOD1: 5.93±4.53nA/μS, N=14; g=−0.18 95%CI[−0.88–0.63]; t(27)=0.49, p=0.63. B. Plot of the PIC amplitude vs. the input conductance of young adult motoneurons (P30–60) in WT (blue squares) and mSOD1 (red diamonds) animals. The marginal plots indicate the kernel density estimation of the distributions of the values in the two populations. The asterisks indicate point to fraction of the mSOD1 population that is outside the range of the WT population. C. Same as A for the presymptomatic age range P60–90. D. Same as A for the symptomatic age range P90–120.

Our results show that, during the disease progression, motoneurons are actively engaged in a homeostatic process to maintain their firing output. It is plausible that this process would, by itself, cause undue stress to the cell. For example, the ratio of PIC to conductance is a major determinant of net excitability (Huh et al., 2017). If these two parameters grow in proportion, then net excitability is likely to stay about the same. We thus analyzed PIC amplitudes normalized by input conductances (Figure 5A). Although pre-symptomatic and symptomatic cells manage to maintain a normal normalized-PIC amplitudes, young adult mSOD1 motoneurons have a larger normalized PIC amplitude than their WT controls (Figure 5A). This implies that, in these animals, the cells that are abnormally large (large input conductance) also have PICs that are larger than expected for their size. It is likely that these cells are under considerable stress to respond to the high metabolic demand imposed by their large size, as well as the potential massive influx of calcium caused by their abnormally large PICs.

This effect is also visible in the plots of Figure 5B–D showing the relationship between PIC amplitude and input conductance. In the young adult animals (Figure 5B), the majority of the mutant motoneurons are clustered in the upper right hand corner, with values of both conductances and PIC amplitudes that are outside of the

range of WT controls. At the pre-symptomatic stages, the WT and mSOD1

populations roughly overlap (Figure 5C), and at the symptomatic ages, the situation

is reversed, with the appearance of a cluster (N=6) of very small cells (asterisk in

Figure 5D), most of which also have very small PICs. These results show that the

overall initial increase in PIC and conductance amplitudes driven by the SOD1

mutation generates a sub-population of motoneurons with conductance and PIC

amplitudes that are actually above those of the normal, WT range (P30–60). The

steady reduction in these amplitudes is so strong that by P90–120, a sub-population

develops that falls below the normal range.

Hypoexcitable cells

404
405   **Figure 6: Some cells are hypoexcitable and cannot fire repetitively**

A. Example of a mSOD1 motoneuron (from a P88 mouse) that is unable to fire repetitively in response to a triangular ramp of current. Top trace: membrane potential, bottom trace: injected current. B. This same motoneuron was able to generate a single full-height action potential in response to a square pulse of current. Same organization as in A. C. Non-firing motoneurons had a similar input conductance compared to motoneurons that were able to fire repetitively. Firing: 0.49±0.22μS, N=47 vs. non-firing: 0.38±0.20μS, N=6; $g=−0.54$ 95%CI[−1.11–0.50]; $t(51)=1.36$, $p=0.22$ D. Non-firing motoneurons had much smaller PICs than motoneurons able to fire repetitively. Firing: Firing: 4.04±3.18nA, N=47 vs. non-firing: 0.60±0.49nA, N=6; $g=−1.12$ 95%CI[−1.42−−0.86]; $t(51)=6.81$, $p=1.3e−08$ E. Non-firing motoneurons appear earlier in mSOD1 animals compared to WT animals. WT: 100±16 days old, N=7 vs. SOD1: 75±23 days old, N=6; $g=−1.20$ 95%CI[−2.06–0.01]; $t(11)=2.24$, $p=0.053$

As previously shown (Delestrée et al., 2014; Martínez-Silva et al., 2018), mSOD1 motoneuron tend to lose their ability to fire repetitively as the disease progresses. In the present study, out of the 53 mutant motoneurons recorded, 6 motoneurons could not fire repetitively to current ramps despite still being able to fire a single or a few action potentials to current steps. Figure 8 shows an example of such a motoneuron. Although those cells were not, on average, larger than those that fired repetitively (Figure 6B), non-firing cells were characterized by a very PIC amplitude (Figure 6C, see also (Huh et al., 2017)). Even more striking were the difference in PIC amplitude normalized by the input conductance of the cells. Non-firing motoneurons had a much smaller normalized PIC amplitude (1.91±1.82nA/μS, N=6) compared to repetitively firing motoneurons (7.80±4.79nA/μS, N=47; $g=−1.27$ 95%CI[−1.65−−0.80]; $t(51)=5.77$, $p=2.6e−05$). Finally, although some non-firing motoneurons could be found in WT mice, non-firing motoneurons appear on average 25 days earlier in mSOD1 mice (Figure 6D).

Table 1: summary of the properties of the motoneurons, broken down by age range

The table shows the various properties that were measured in each motoneuron. The rows are sorted in descending order of effect size at P30–60. WT and mSOD1 columns show mean values ± standard deviation, with the 95%CI around the mean below. The "Mean diff." column shows the mean difference between mSOD1 and WT population, with 95% CI below. The "Hedges' g" column shows a graphical representation of the effect size. ΔF: difference between the instantaneous firing frequency at de-recruitment and at recruitment. ΔI: difference between the de-recruitment and recruitment currents. ΔV_{th}: voltage threshold for spiking relative to the resting membrane potential. ΔV_{PIC}: PIC onset voltage relative to the resting membrane potential. ΔV_{peak}: voltage at the peak of the PICs relative to the resting membrane potential

Discussion

In this paper, we studied how motoneuron electrical properties evolve over the time course of ALS, focusing particularly on the persistent inward currents (PICs). We show that, in young adult mutant mice, before and up to the time where denervations are just starting, PICs are abnormally large compared to controls. This increase in PIC amplitude is accompanied by a parallel increase in motoneuron input conductance with the net effect that the excitability of the cells remains normal. Later, while the animals remain pre-symptomatic but denervation has begun, motoneuron properties return to normal levels. Finally, in symptomatic animals, mutant motoneurons tend to have smaller input conductance and smaller recruitment current.

Homeostatic regulation of motoneuron output

Although ALS is classically considered an adult-onset disease, we know from earlier studies of motoneurons in animal models of ALS that the change in motoneurons' intrinsic excitability is the first sign to be seen in the pathogenesis, long before any overt motor deficits manifest. Hyperexcitability is observed in

cultured E13 spinal motoneurons (Kuo et al., 2004), E15 cortical motoneurons (Pieri

et al., 2009), E17.5 spinal motoneurons (Martin et al., 2013 p.20), as well as early

postnatal hypoglossal motoneurons (van Zundert et al., 2008). Consequently, the

motor system in general, and motoneurons in particular, must exhibit remarkable

capability for homeostatic regulation to maintain a quasi-normal motor output until

overt symptoms appear, but there are many combinations of intrinsic properties

that can produce the same firing pattern (Marder and Goaillard, 2006). In neonatal

mSOD1 mice, the majority of motoneurons exhibited normal excitability (based on

recruitment current and F-I gain), although the most resistant population of

motoneuron seem to retain some hyperexcitability (Quinlan et al., 2011; Leroy et al.,

2014). Yet, PICs were almost twice as large in mutant compared to WT neonatal

motoneurons. This increased PIC amplitude was seemingly compensated by a

parallel increase in input conductance (Quinlan et al., 2011). In young adult animals,

this trend seems to continue, as the present study demonstrated an almost 3×

increase in PIC amplitude, accompanied by an almost 2× increase in input

conductance. Similarly, we observed large changes in PIC onset and peak voltages,

compensated by a commensurate hyperpolarization of the resting membrane

potential. These trends with disease progression from the neonatal state through

development of severe symptoms are summarized in Figure 7. An overall pattern of

oscillations is evident, with the PIC oscillations tending to increase excitability but

the conductance and rest potential oscillations tending to reduce it.

483
Figure 7: summary of the changes in motoneuron properties over time
Schematic representation of the changes in four key electrophysiological properties
over time. The dots represent the effect size (Hedges' g) and the vertical bars show
the 95%CI around g. The thin lines are cubic splines interpolation of the data over
time. The points have been slightly staggered so that the vertical bars do not occlude
each other. Data from neonates (P0–5 and P6–12) are from Quinlan et al. (2011).

The properties of the motoneurons seem to normalize at the late pre-

symptomatic stage (P60–90), but it is unclear whether this is due to changes in the

homeostatic pathways involved, or whether it is caused by the start of the

degeneration process in the most vulnerable motoneurons. At symptomatic stages,

motoneurons appear to be hypoexcitable. They have a smaller input conductance

(Figure 3), and smaller recruitment current (Figure 2), but that most likely reflect

the fact that the largest cells have degenerated by this age, and that the remaining

cells may have shrunk (see below, and Dukkipati et al., 2018).

The hidden cost of homeostatic regulation

Although the degeneration of motoneurons in ALS constitutes a failure in cell

homeostasis, our results show that, throughout the "silent", presymptomatic phase

of ALS, motoneurons are actively engaged in a highly successful homeostatic process

to maintain their firing output. However, this process is not perfect, and, by itself,

could be a major source of stress for the cells. We do not know whether the cells

increase their conductance to compensate for the fast increase in PICs or *vice versa*,

but the end result is the appearance of cells with both input conductance and PIC

amplitudes outside of the normal range (Figure 5B). The increase in input

conductance could be due to the insertion of more channels in the membrane.

However, direct measurements of soma sizes have revealed that pre-symptomatic

mutant cells are physically larger than controls (Shoenfeld et al., 2014; Dukkipati et

al., 2018). This increase in size, coupled with the larger calcium entry in the cells due

to the larger PICs (which are in part calcium-medicated (Li and Bennett, 2003)), is

bound to cause undue metabolic stress on those cells that are outside of the normal

range, despite the fact that cells manage to maintain a normal output. These cells are

likely to be Fast Fatigable (FF) motoneurons, as they are the cells with the largest

input conductance, and the largest PICs (Lee and Heckman, 1998; Huh et al., 2017)

in normal conditions, and they are the cells that are the most vulnerable to ALS (Pun

et al., 2006; Hegedus et al., 2007, 2008).

As the disease progresses, those large, vulnerable cells become the first

motoneurons that will embark on a degeneration pathway (Saxena et al., 2009,

2013; Martínez-Silva et al., 2018), and will lose their ability to fire repetitively

(Martínez-Silva et al., 2018). This early loss of the large cells is consistent with stress

induced by aberrantly large conductances and PICs. This loss is also likely the

primary reason why motoneurons appear to recover normal properties at late pre-

symptomatic stages. Nevertheless, it is probable that all motoneurons, regardless of

type, experience this form of stress. Interestingly, we observed that some of the

smallest motoneurons, which are the most resistant to the disease, appear to shrink

in the oldest animals (Figure 5D), consistent with anatomical observations (Kiernan

and Hudson, 1991; Dukkipati et al., 2018). At this point, it is unclear whether this

shrinkage is a pathological feature or a strong homeostatic effort by the surviving

motoneurons to counteract the stress generated by their initial hypertrophy.

Hyper-vigilant homeostasis

Mitchell and colleagues have recently proposed that cellular homeostasis is in

ALS motoneurons is in fact too strong (Mitchell and Lee, 2012; Irvin et al., 2015).

Several aspects of our results are consistent with this hyper-vigilant homeostasis

hypothesis. A major impetus for this hypothesis was that meta-analyses of data

obtained from many studies on multiple cellular properties revealed patterns of

oscillations as the disease progressed, suggesting the existence of high feedback

gains for homeostatic processes. Similar oscillations are clearly present in our

results, as illustrated in Figure 7. The highly successful homeostasis for net

excitability, achieved at the cost of excessively high amplitudes for conductances

and PICs, clearly shows homeostasis is strong. The development of subpopulations

that are initially well above and then well below the normal range supports the

possibility that this strong homeostasis is excessive. Direct tests of the hyper-

vigilant homeostasis hypothesis will require quantitative comparisons of the

responses of ALS and WT motoneurons to controlled homeostatic challenges.

Comparison with previous studies

The present study displays some differences with other studies, including by our

own. Others have described signs of hyperexcitability in spinal motoneurons

(Meehan et al., 2010; Jensen et al., 2020), and in the motor cortex (Kim et al., 2017),

accompanied with changes in local circuitry (Hedegaard et al., 2015; Kim et al.,

2017; Bączyk et al., 2020; Jara et al., 2020), although that aspect was not

investigated in the present work. We have previously studied the properties of

motoneurons in the same SOD1(G93A) mouse model of ALS group (Delestrée et al.,

2014; Martínez-Silva et al., 2018) but have mainly focused on a short period of time

just at the onset of denervations (P45–55). Although that time point partly overlaps

with the present young adult stage, we had not detected differences in input

conductances (Martínez-Silva et al., 2018). This might be due to the fact that the

difference in input conductance is largest at earlier time points (Figure 3A). Indeed,

the difference is gone at P60–90, and we did report an increase in input conductance

in an earlier series of experiments, which was more pronounced at younger time

points (Delestrée et al., 2014). Similarly, we did not detect a difference in resting

membrane potential in the Martínez-Silva et al. (2018) study. Again, the discrepancy

could be due to the difference in the distributions of the ages of the animals studied.

It is also possible that, despite our best efforts, the disease is not following the exact

same time course in the two colonies used (the present paper is based on animals

bred at Northwestern University in Chicago, while our previous studies were

conducted on animals bred in Paris). It is therefore conceivable that even though the

ages overlap, those ages correspond to slightly different stages of the disease.

In the present study, we recorded only 13 motoneurons out of 103 (7/50 WT and

6/53 mSOD1) that were unable to fire repetitively, contrary to our previous study

where we showed that a large proportion of vulnerable motoneurons became

hypoexcitable before the onset of denervation (P45–55, Martínez-Silva et al., 2018).

This discrepancy could be explained by the experimental constraints of the present

study. Indeed, obtaining stable voltage clamp recordings in vivo in mice is quite a

challenge, requiring electrodes able to pass very large amounts of current and

motoneuron able to withstand the protocol. Since we have only included for analysis

cells in which the voltage clamp protocol was completed successfully, we

hypothesize that we may have biased our sample towards the healthiest cells that

have yet to take the path of degeneration.

Limitations

The results presented here have been collected on the SOD1(G93A) mouse model

of ALS, a model that has attracted criticisms due to it being an overexpression

model, and its failure to bring seemingly promising therapies to the clinic (Philips

and Rothstein, 2015). Nevertheless, some parallel with other studies suggest that

our results could be extended to other models. First, Meehan et al. (2010) have

studied an unrelated mSOD1(G127X) model and have shown that although the net

excitability of the motoneurons was not affected by the mutation, there were

nonetheless signs of increased PICs in these motoneurons. Second, we observed

that, in late symptomatic animals, some cells exhibited a very small input

conductance, well below the normal range at this age, suggesting that these cells

actually shrank. This observation matches morphological measurements in mSOD1

mice (Dukkipati et al., 2018), but also observations from sporadic human patients

(Kiernan and Hudson, 1991).

We demonstrated a remarkably successful homeostatic control of motoneuron

firing, that we can attribute to commensurate changes in PICs and input

conductance. However, it is likely that other currents are also implicated in this

process (Marder and Goaillard, 2006). Limitations inherent to in vivo

electrophysiology prevent us from isolating many different currents and further in

vitro investigations in adults (Jiang and Heckman, 2006; Mitra and Brownstone,

2011; Jiang et al., 2017; Bhumbra and Beato, 2018) are warranted to shed more light

on the panoply of channels involved in this process.

Conclusion

Overall, our results show homeostasis for net excitability in mutant SOD1

motoneurons is remarkably strong in the presymptomatic state. This success

however comes at a cost of large compensatory changes in basic electrical

properties. These results support the hypothesis that ALS is not due to a single

specific root cause but instead caused by an inherent instability at the system-level

(Mitchell and Lee, 2012), possibly caused by an "hyper-vigilant" homeostatic system

in motoneurons. Initial perturbations in the electrical properties of motoneurons

would be overcompensated for, leading to new sources of stress, which, in turn,

would be overcorrected, and so on and so forth until the metabolic burden becomes

too high to sustain for the motoneuron. FF motoneurons have higher metabolic

needs (Le Masson et al., 2014) and lower calcium-buffering capabilities

(Grosskreutz et al., 2010) in the first place, which make them particularly vulnerable

to this vicious cycle.

Data availability

617   Data files and a computational notebook allowing reproducing the analysis and

618   figures of this article are provided at the URL:

619   http://doi.org/10.5281/zenodo.3831947

References

Bączyk M, Alami NO, Delestrée N, Martinot C, Tang L, Commisso B, Bayer D, Doisne
N, Frankel W, Manuel M, Roselli F, Zytnicki D (2020) Synaptic restoration by
cAMP/PKA drives activity-dependent neuroprotection to motoneurons in ALS. J Exp
Med 217 Available at: https://rupress-
org.insb.bib.cnrs.fr/jem/article/217/8/e20191734/151829/Synaptic-restoration-
by-cAMP-PKA-drives-activity [Accessed June 2, 2020].

Bernard C (2019) Changing the Way We Report, Interpret, and Discuss Our Results
to Rebuild Trust in Our Research. eNeuro 6 Available at:
https://www.eneuro.org/content/6/4/ENEURO.0259-19.2019 [Accessed April 13,
2020].

Bhumbra GS, Beato M (2018) Recurrent excitation between motoneurones
propagates across segments and is purely glutamatergic. PLOS Biol 16:e2003586
Available at:
https://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.2003586
[Accessed May 7, 2020].

Brownstone RM, Lancelin C (2018) Escape from homeostasis: spinal microcircuits
and progression of amyotrophic lateral sclerosis. J Neurophysiol 119:1782–1794
Available at: https://www.physiology.org/doi/full/10.1152/jn.00331.2017
[Accessed October 25, 2018].

Cumming G (2012) Understanding the new statistics: effect sizes, confidence

intervals, and meta-analysis. New York: Routledge, Taylor & Francis Group.

Delestrée N, Manuel M, Iglesias C, Elbasiouny SM, Heckman CJ, Zytnicki D (2014)
Adult spinal motoneurones are not hyperexcitable in a mouse model of inherited
amyotrophic lateral sclerosis. J Physiol 592:1687–703 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/24445319.

Dukkipati SS, Garrett TL, Elbasiouny SM (2018) The vulnerability of spinal
motoneurons and soma size plasticity in a mouse model of amyotrophic lateral
sclerosis. J Physiol 596:1723–1745.

Grosskreutz J, van den Bosch L, Keller BU (2010) Calcium dysregulation in
amyotrophic lateral sclerosis. Cell Calcium 47:165–174 Available at:
https://doi.org/10.1016/j.ceca.2009.12.002.

Hedegaard A, Lehnhoff J, Moldovan M, Grøndahl L, Petersen NC, Meehan CF (2015)
Postactivation depression of the Ia EPSP in motoneurons is reduced in both the
G127X SOD1 model of amyotrophic lateral sclerosis and in aged mice. J
Neurophysiol 114:1196–210.

Hedges LV (2016) Distribution Theory for Glass's Estimator of Effect size and
Related Estimators: J Educ Stat Available at:
https://journals.sagepub.com/doi/10.3102/10769986006002107 [Accessed May 7,
2020].

Hegedus J, Putman CT, Gordon T (2007) Time course of preferential motor unit loss
in the SOD1 G93A mouse model of amyotrophic lateral sclerosis. Neurobiol Dis
28:154–164 Available at: https://doi.org/10.1016/j.nbd.2007.07.003.

Hegedus J, Putman CT, Tyreman N, Gordon T (2008) Preferential motor unit loss in
the SOD1 G93A transgenic mouse model of amyotrophic lateral sclerosis. J Physiol
586:3337–3351 Available at: https://doi.org/10.1113/jphysiol.2007.149286.

Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A (2019) Moving beyond P values:
data analysis with estimation graphics. Nat Methods 16:565–566 Available at:
https://www-nature-com.insb.bib.cnrs.fr/articles/s41592-019-0470-3 [Accessed
December 30, 2019].

Huh S, Siripuram R, Lee RH, Turkin VV, O'Neill D, Hamm TM, Heckman CJ, Manuel M
(2017) PICs in motoneurons do not scale with the size of the animal: a possible
mechanism for faster speed of muscle contraction in smaller species. J Neurophysiol
118:93–102 Available at: https://doi.org/10.1152/jn.00045.2017.

Hunter JD (2007) Matplotlib: A 2D Graphics Environment. Comput Sci Eng 9:90–95.

Ilieva H, Polymenidou M, Cleveland DW (2009) Non-cell autonomous toxicity in
neurodegenerative disorders: ALS and beyond. J Cell Biol 187:761–772 Available at:
https://doi.org/10.1083/jcb.200908164.

Irvin CW, Kim RB, Mitchell CS (2015) Seeking homeostasis: temporal trends in
respiration, oxidation, and calcium in SOD1 G93A Amyotrophic Lateral Sclerosis
mice. Front Cell Neurosci 9:248.

Jara JH, Sheets PL, Nigro MJ, Perić M, Brooks C, Heller DB, Martina M, Andjus PR,
Ozdinler PH (2020) The Electrophysiological Determinants of Corticospinal Motor
Neuron Vulnerability in ALS. Front Mol Neurosci 13:73.

Jensen DB, Kadlecova M, Allodi I, Meehan CF (2020) Spinal motoneurones are
intrinsically more responsive in the adult G93A SOD1 mouse model of Amyotrophic
Lateral Sclerosis. J Physiol n/a Available at:
https://physoc.onlinelibrary.wiley.com/doi/abs/10.1113/JP280097 [Accessed July
28, 2020].

Jiang MC, Adimula A, Birch D, Heckman CJ (2017) Hyperexcitability in synaptic and
firing activities of spinal motoneurons in an adult mouse model of amyotrophic
lateral sclerosis. Neuroscience 362:33–46.

Jiang MC, Heckman CJ (2006) In vitro sacral cord preparation and motoneuron
recording from adult mice. J Neurosci Off J Soc Neurosci 156:31–36 Available at:
https://doi.org/10.1016/j.jneumeth.2006.02.002.

Johnson MD, Heckman CJ (2010) Interactions between focused synaptic inputs and
diffuse neuromodulation in the spinal cord. Ann N Y Acad Sci 1198:35–41 Available
at: https://doi.org/10.1111/j.1749-6632.2010.05430.x.

Kanning KC, Kaplan A, Henderson CE (2009) Motor neuron diversity in development
and disease. Annu Rev Neurosci 33:409–440 Available at:
https://doi.org/10.1146/annurev.neuro.051508.135722.

Kiernan JA, Hudson AJ (1991) Changes in Sizes of Cortical and Lower Motor Neurons
in Amyotrophic Lateral Sclerosis. Brain 114:843–853 Available at:
https://academic-oup-com.insb.bib.cnrs.fr/brain/article/114/2/843/263744
[Accessed April 13, 2020].

Kim J, Hughes EG, Shetty AS, Arlotta P, Goff LA, Bergles DE, Brown SP (2017)
Changes in the Excitability of Neocortical Neurons in a Mouse Model of Amyotrophic
Lateral Sclerosis Are Not Specific to Corticospinal Neurons and Are Modulated by
Advancing Disease. J Neurosci Off J Soc Neurosci 37:9037–9053.

Kuo JJ, Schonewille M, Siddique T, Schults ANA, Fu R, Bär PR, Anelli R, Heckman CJ,
Kroese ABA (2004) Hyperexcitability of cultured spinal motoneurons from
presymptomatic ALS mice. J Neurophysiol 91:571–575 Available at:
https://doi.org/10.1152/jn.00665.2003.

Le Masson G, Przedborski S, Abbott LF (2014) A Computational Model of Motor
Neuron Degeneration. Neuron 83:758–760 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/25088365.

Lee RH, Heckman CJ (1998) Bistability in spinal motoneurons in vivo: systematic
variations in persistent inward currents. J Neurophysiol 80:583–593.

Leroy F, Lamotte d'Incamps B, Imhoff-Manuel RD, Zytnicki D (2014) Early intrinsic
hyperexcitability does not contribute to motoneuron degeneration in amyotrophic
lateral sclerosis. eLife 3 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/25313866.

Li Y, Bennett DJ (2003) Persistent sodium and calcium currents cause plateau
potentials in motoneurons of chronic spinal rats. J Neurophysiol 90:857–869
Available at: https://doi.org/10.1152/jn.00236.2003.

Makin TR, Orban de Xivry J-J (2019) Ten common statistical mistakes to watch out
for when writing or reviewing a manuscript Rodgers P, Parsons N, Holmes N, eds.
eLife 8:e48175 Available at: https://doi.org/10.7554/eLife.48175 [Accessed
October 28, 2019].

Manuel M, Heckman CJ (2011) Adult mouse motor units develop almost all of their
force in the subprimary range: a new all-or-none strategy for force recruitment? J
Neurosci 31:15188–15194 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/22016552.

Manuel M, Iglesias C, Donnet M, Leroy F, Heckman CJ, Zytnicki D (2009) Fast
kinetics, high-frequency oscillations, and subprimary firing range in adult mouse
spinal motoneurons. J Neurosci 29:11246–11256 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/19741131.

Marder E, Goaillard J-M (2006) Variability, compensation and homeostasis in
neuron and network function. Nat Rev Neurosci 7:563–574.

Martin E, Cazenave W, Cattaert D, Branchereau P (2013) Embryonic alteration of
motoneuronal morphology induces hyperexcitability in the mouse model of
amyotrophic lateral sclerosis. Neurobiol Dis 54:116–26.

Martínez-Silva M de L, Imhoff-Manuel RD, Sharma A, Heckman C, Shneider NA,
Roselli F, Zytnicki D, Manuel M (2018) Hypoexcitability precedes denervation in the
large fast-contracting motor units in two unrelated mouse models of ALS Rubin LL,
ed. eLife 7:e30955 Available at: https://doi.org/10.7554/eLife.30955 [Accessed
February 7, 2019].

McKinney W (2011) pandas: a foundational Python library for data analysis and
statistics. Python High Perform Sci Comput 14.

Meehan CF, Moldovan M, Marklund SL, Graffmo KS, Nielsen JB, Hultborn H (2010)
Intrinsic properties of lumbar motor neurones in the adult G127insTGGG
superoxide dismutase-1 mutant mouse in vivo: evidence for increased persistent
inward currents. Acta Physiol Oxf Engl 200:361–76.

Michel MC, Murphy TJ, Motulsky HJ (2020) New Author Guidelines for Displaying
Data and Reporting Data Analysis and Statistical Methods in Experimental Biology.
Mol Pharmacol 97:49–60 Available at:
http://molpharm.aspetjournals.org/lookup/doi/10.1124/mol.119.118927
[Accessed January 13, 2020].

Mitchell CS, Lee RH (2012) Dynamic Meta-Analysis as a Therapeutic Prediction Tool
for Amyotrophic Lateral Sclerosis. In: Amyotrophic Lateral Sclerosis (Maurer M, ed).
InTech. Available at: http://www.intechopen.com/books/amyotrophic-lateral-
sclerosis/dynamic-meta-analysis-as-a-therapeutic-prediction-tool-for-amyotrophic-
lateral-sclerosis [Accessed January 5, 2020].

Mitra P, Brownstone RM (2011) An in vitro spinal cord slice preparation for
recording from lumbar motoneurons of the adult mouse. J Neurophysiol 107:728–
741 Available at: https://www.physiology.org/doi/full/10.1152/jn.00558.2011
[Accessed December 6, 2018].

Philips T, Rothstein JD (2015) Rodent Models of Amyotrophic Lateral Sclerosis. Curr
Protoc Pharmacol Editor Board SJ Enna Ed--Chief Al 69:5.67.1-5.67.21 Available at:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4562058/ [Accessed April 13,
2020].

Pieri M, Carunchio I, Curcio L, Mercuri NB, Zona C (2009) Increased persistent
sodium current determines cortical hyperexcitability in a genetic model of
amyotrophic lateral sclerosis. Exp Neurol 215:368–379 Available at:
https://doi.org/10.1016/j.expneurol.2008.11.002.

Pun S, Santos AF, Saxena S, Xu L, Caroni P (2006) Selective vulnerability and pruning
of phasic motoneuron axons in motoneuron disease alleviated by CNTF. Nat
Neurosci 9:408–19 Available at: http://www.ncbi.nlm.nih.gov/pubmed/16474388.

Quinlan KA, Schuster JE, Fu R, Siddique T, Heckman CJ (2011) Altered postnatal
maturation of electrical properties in spinal motoneurons in a mouse model of
amyotrophic lateral sclerosis. J Physiol 589:2245–2260 Available at:
http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=21486
770&retmode=ref&cmd=prlinks.

Rosen DR, Siddique T, Patterson D, Figlewicz DA, Sapp P, Hentati A, Donaldson D,
Goto J, O'Regan JP, Deng H-X (1993) Mutations in Cu/Zn superoxide dismutase gene
are associated with familial amyotrophic lateral sclerosis. Nature 362:59–62
Available at: https://doi.org/10.1038/362059a0.

Saxena S, Cabuy E, Caroni P (2009) A role for motoneuron subtype-selective ER
stress in disease manifestations of FALS mice. Nat Neurosci 12:627–636 Available
at: https://doi.org/10.1038/nn.2297.

Saxena S, Roselli F, Singh K, Leptien K, Julien J, Gros-louis F, Caroni P (2013)
Neuroprotection through Excitability and mTOR Required in ALS Motoneurons to
Delay Disease and Extend Survival. Neuron 80:80–96.

Sekerli M, Del Negro CA, Lee RH, Butera RJ (2004) Estimating action potential
thresholds from neuronal time-series: new metrics and evaluation of
methodologies. IEEE Trans Biomed Eng 51:1665–1672.

Shoenfeld L, Westenbroek RE, Fisher E, Quinlan K a, Tysseling VM, Powers RK,
Heckman CJ, Binder MD (2014) Soma size and Cav1.3 channel expression in
vulnerable and resistant motoneuron populations of the SOD1G93A mouse model of
ALS. Physiol Rep 2:e12113 Available at:
http://physreports.physiology.org/content/2/8/e12113.abstract.

van Zundert B, Peuscher MH, Hynynen M, Chen A, Neve RL, Brown RH, Constantine-
Paton M, Bellingham MC (2008) Neonatal neuronal circuitry shows hyperexcitable
disturbance in a mouse model of the adult-onset neurodegenerative disease

amyotrophic lateral sclerosis. J Neurosci Off J Soc Neurosci 28:10864–10874
Available at: https://doi.org/10.1523/JNEUROSCI.1340-08.2008.

Virtanen P et al. (2020) SciPy 1.0: fundamental algorithms for scientific computing
in Python. Nat Methods:1–12 Available at:
https://www.nature.com/articles/s41592-019-0686-2 [Accessed March 1, 2020].

Waskom M et al. (2020) mwaskom/seaborn: v0.10.0 (January 2020). Zenodo.
Available at: https://doi.org/10.5281/zenodo.3629446.

Wasserstein RL, Schirm AL, Lazar NA (2019) Moving to a World Beyond "p < 0.05."
Am Stat 73:1–19 Available at:
https://amstat.tandfonline.com/doi/full/10.1080/00031305.2019.1583913
[Accessed April 13, 2020].

Welch BL (1947) The generalisation of student's problems when several different
population variances are involved. Biometrika 34:28–35.