Seasonal variation in brain mu-opioid receptor availability

Lihua Sun[1*], Jing Tang[2], Heidi Liljenbäck[1,3], Aake Honkaniemi[1,4], Jenni Virta[1,3], Janne Isojärvi[1],

Tomi Karjalainen[1], Tatu Kantonen[1,5], Pirjo Nuutila[1,6], Jarmo Hietala[1,7], Valtteri Kaasinen[1,5], Kari

Kalliokoski[1], Jussi Hirvonen[8], Harry Scheinin[1], Semi Helin[1], Kim Eerola[9,10], Eriika

Savontaus[9,10], Emrah Yatkin[11], Juha O. Rinne[1,5], Anne Roivainen[1,3], Lauri Nummenmaa[1,12]

1. Turku PET Centre, University of Turku, Turku, Finland

2. Research Program in Systems Oncology, Faculty of Medicine, University of Helsinki, Helsinki, Finland

3. Turku Center for Disease Modelling, University of Turku, Turku, Finland

4. Turku University Hospital, Turku, Finland

5. Clinical Neuroscience, Turku University Hospital, Finland

6. Department of Endocrinology, Turku University Hospital, Turku, Finland

7. Department of Psychiatry, Turku University Hospital, Turku, Finland.

8. Department of Radiology, University of Turku, Turku, Finland

9. Research Unit for Integrative Physiology and Pharmacology, Institute of Biomedicine, University of Turku, Turku, Finland

10. Unit of Clinical Pharmacology, Turku University Hospital, Turku, Finland

11. Central Animal Laboratory, University of Turku, Turku, Finland

12. Department of Psychology, University of Turku, Turku, Finland

*Correspondence to: lihua.sun@utu.fi

Turku PET Centre, University of Turku, Turku, Finland

24   **Abstract**

Seasonal rhythms influence mood and sociability. The brain μ-opioid receptor (MOR) system

modulates a multitude of seasonally varying socioemotional functions, but its seasonal variation

remains elusive with no previously reported *in vivo* evidence. Here, we studied the seasonal effects

on brain MOR availability via analysing a dataset (n=204) of [^{11}C]carfentanil positron emission

tomography (PET) scans of healthy volunteers. We found that seasonally varying daylength had

an inverted U-shaped functional relationship with brain MOR availability. Brain regions sensitive

to daylength spanned the socio-emotional brain circuits, where MOR availability formed a spring-

like peak. Causal effect of daylength on brain MOR availability was further verified by a *post hoc*

experiment with repeated PET imaging of rats (n=9) under seasonal photoperiodic simulation.

Therefore, the *in vivo* brain MOR availability in normal humans shows significant seasonal

variation, which aligns with expected seasonal variation in mood and suicidality.

Introduction

Seasonal rhythms profoundly impact mood. Negative affect including depression, anger, and hostility is at lowest during the summer [1], whereas seasonal affective disorder rates peak during the winter months [2]. These changes are mediated by slow phasic changes in different neuroreceptor systems. For instance, long daylength increases brain serotonin and norepinephrine levels in mice and reduces depression and anxiety behaviour, compared to short daylength [3]. Duration of daylight exposure is similarly associated with brain serotonin turnover in humans [4]. The intimate link between seasonal fluctuations in mood and the contribution of opioidergic neurotransmission in human emotions [5] suggests potential seasonal variation of *in vivo* μ-opioid receptor (MOR) signalling.

Several lines of evidence suggest that MOR availability could vary seasonally in humans. Opioids are among the most commonly used illicit drugs in the US, where 2 % of the population have had opioid use disorder during their lifetime [6]. Suicidal behaviour with prescription opioids follows a clear seasonal pattern, with suicide attempts peaking in the spring and fall [7]. Further, post-mortem studies have established that suicide victims have increased MOR densities [8,9], and serious self-injurious behaviour is associated with increased plasma opioid growth factor levels [10]. Although these studies have not directly assessed seasonal effects, it is well established that suicide rate peaks often in the spring regardless of the geographical location of the country [11]. Finally, MORs are potent modulators of feeding [12–14] due to their contribution of hedonic or "liking" responses in the brain [15], and human feeding patterns show seasonal variation with caloric intake of fats typically peaking in the fall or winter [16,17]. However, direct *in vivo* evidence on the causal seasonal effects on opioidergic neurotransmission is currently lacking.

Animal studies also suggest that there is seasonal variation in MOR-dependent signalling. The

endogenous opioid signalling is crucial for the photoperiodic control of the seasonal reproductive

cycle in mammals [18]. Endogenous opioids inhibit the release of gonadotropin and sex hormones,

and this inhibitory effect is enhanced under short daylength compared to long daylength [19]. Also,

MOR expression in hamster testes is increased during short days, with increased inhibitory control

for testosterone secretion [20]. In Siberian hamsters, brain expression of dynorphin A, an endogenous

peptide agonist for opioid receptors, is increased under relatively longer daylength compared to

shorter daylength [21], suggesting a potential impact of daylength on opioidergic signalling in the

brain. Furthermore, effects of morphine on feeding behaviour of ground squirrels vary in accord

to the hibernation state [22], indirectly reflecting a seasonal variation of MOR signalling.

Here, we tested whether seasonal variation in daylength, a noise-free cue for seasonal rhythms,

influences MOR availability in the brain. MOR availability was quantified using *in vivo* positron

emission tomography (PET) with MOR-sensitive agonist ligand [^{11}C]carfentanil. We first

conducted a cross-sectional study with previously acquired human PET imaging data to test

whether there was seasonal difference in MOR availability. We then investigated experimentally

whether seasonal variation in daylength causally influences brain MOR availability in rats. While

rats underwent daylength cycle simulating seasonal changes, repeated [^{11}C]carfentanil PET

imaging was conducted (Supplementary Fig. 1). Our data suggest that there is seasonal variation

in brain MOR availability in both humans and rats, with MOR availability peaking in seasons with

intermediate daylengths.

Results

Daylength associated brain MOR availability in Humans

Figure 1A shows the mean MOR distribution in the human brain. We examined the association

between natural variation in daylength (Fig. 1B) and MOR availability in humans. Brain full

volume analysis revealed a statistically significant quadratic polynomial effects in a large brain cluster (39000 voxels; Fig. 1C-D), spanning the cingulate cortex, the superior frontal, medial frontal, middle temporal and superior temporal gyrus, insula and orbitofrontal cortex. Brain MOR availability in the brain cluster peaked at 15-19 hours (Fig. 1D).

Figure 1. Human brain MOR availability and natural variation in daylength. **A.** Mean distribution of MOR availability in the subjects. **B.** Local seasonal distribution of daylength (day + civil twilight). **C.** The brain cluster sensitive to the seasonal daylength changes in quadratic polynomial regression model. $p < 0.05$, FDR-corrected. **D.** [^{11}C]carfentanil BP_{ND} in the brain cluster as a function of daylength. Red line shows polynomial LS regression curve ($y \sim x + x^2$) predicting BP_{ND}, and shaded area shows 95% CI.

In addition, we did brain region of interest (ROI) analysis by involving thirty structurally

determined brain ROIs (see Method). While involving all ROIs in the statistical analysis, both

daylength ($\beta = 0.044$, 95% CI [0.008, 0.08]) and squared daylength ($\beta = -0.0014$, 95% CI [-0.0026,

-0.00021]) were significant predictors for [^{11}C]carfentanil BP_{ND}, indicating an inverted-U shape

functional relationship. No interaction effects between daylength and squared daylength with brain

hemisphere (left versus right) or ROIs were found. Effect of daylength on individual ROIs is shown

in Supplementary Table 1.

Meteorological seasons and brain MOR availability in Humans

Traditionally defined seasons, according to the northern meteorological definition, including

spring (March, April, May), summer (June, July, August), fall (September, October, November),

and winter (December, January, February) were used as predictors for MOR BP_{ND} in the brain

cluster (Fig. 1C-D). Each season is dummy-coded in a linear regression model, and we found that

spring predicted higher brain MOR availability ($\beta = 0.08$, 95% CI [0.02, 1.14]), Fig. 2A. No other

seasons had significant effects on MOR availability. In the local region, spring is with daylength

of 16.39 ± 2.78 h (Fig. 2B). Therefore, MOR availability peaking at daylength 15-19 hours (Fig.

1D) suggests a spring-like peak.

Figure 2. Spring is associated with highest brain MOR BP_{ND} in humans. **A.** MOR availability in

the daylength activated brain cluster is plotted to seasons. Blue dots stand for means and error bars

for standard deviations. **B.** Local daylength distribution in different seasons. * = p < 0.05.

Causal effect of daylength on brain MOR availability in Rats

In addition to analysis of the human database, we set up a rat model PET imaging experiment to

verify potential causal effect of daylength on brain MOR availability. This is also a crucial

complement considering the quasi-experimental design nature of human data analysis which could

not differentiate the effect of daylength from other highly correlated seasonal factors such as

temperature. The rat model study consisted of experimental groups (n = 9) where daylength

simulated the local daylength changes (no twilight) and control groups (n = 3) where daylength

were kept consistent (12h light ON). When both experimental and control groups were included

in the statistical model, both daylength (β = 0.16, 95% CI [0.035, 0.28]) and squared daylength (β

= -0.0067, 95% CI [-0.012, -0.0017]) were significant predictors; the MOR availability and

daylength had an inverted U-shaped relationship. Age, sex or group did not influence MOR

availability, and no interaction effects were found. Analysis excluding the control group gave

similar results (see Supplementary data).

We also modelled regional BP_{ND} separately for the experimental group rats (within-animal design),

using fixed-effect factors daylength and squared daylength and varying intercepts for each rat.

Daylength and squared daylength were significant predictors for [^{11}C]carfentanil BP_{ND} in all

regions (Fig 3; Supplementary Table 2). Including age and sex in the models did not improve the

models, as evidenced by higher Akaike Information Criterion (AIC) values (Supplementary results

& Supplementary Table 3). In addition, rats in the experimental versus control group showed

slower weight gain (Supplementary Fig. 3) and had a higher level of blood corticosterone (β =

44.93, 95% CI [12.37, 77.48]).

Figure 3. Within-animal changes in brain MOR BP_{ND} as a function of daylength. Individual

animals are shown as separate lines. Black dashed line shows the LS-regression curve for the

polynomial model (y ~ x + x^2) predicting BP_{ND} across the whole sample. Shaded area shows 95%

CI for the LS curve.

<div align="center">**Discussion**</div>

The current study reveals that seasonally varying daylength influences in vivo brain MOR

availability in healthy humans. Increasing daylength modulated brain MOR availability following

an inverted-U shape, with highest MOR bindings observed in intermediate daylengths. In rats,

daylength causally influences MOR availability paralleling the quasi-experimental study in

humans using a large database of historical PET imaging data. Most profound seasonal variation

of MOR availability was found in large brain clusters including the amygdala, striatum, thalamus,

cingulate cortex, orbitofrontal cortex, superior frontal and temporal cortex. Previous studies have

established the role of seasonal variation in serotonin signalling on brain affective functions and

pointed towards the role of daylight duration on its seasonal rhythm [3,4]. The present data show that

also MOR system undergoes significant seasonal variation, which might be linked with seasonal

variation in mood and sociability.

The human data revealed salient seasonal variation of MOR availability in brain regions linked

with emotional and social processing, such as amygdala, cingulate cortex and posterior superior

temporal cortices [5]. To our knowledge, this is the first *in vivo* evidence indicative of seasonal

variation in MOR system in humans, and it fits with other types of evidence implying seasonal

effects on MOR. In high-latitude regions daylight changes influence sleep quality, mood, and

social behaviour [23–26]. Prevalence of seasonal affective disorders or subsyndromes (time-locked to

short daylength) in the local region is remarkably high [27]. The spring-like peak of MOR availability

(Fig. 2A) parallels with seasonal variation in opioid-modulated emotional changes such as mood

[1], suicidality [11], particularly with prescription opioids [7] that peak particularly in the spring.

Considering the impact of photoperiod on seasonal affective disorders [28] as well as the intimate

linkage between MOR signalling and social and affective functions [5,29], the current study suggests

that seasonal variation in MOR availability could be a potential brain mechanism under seasonal

affective changes.

The rat experiment finding demonstrates that daylength causally influences brain MOR availability,

verifying the human data findings, and goes in line with previously reported impact of photoperiod

on MOR dependent signalling in mammals [19–22]. Exact mechanism on the daylength-mediated

alterations in brain MORs is however unclear. In mammals, neural responses to daylength are

largely mediated by the superchiasmatic nucleus (SCN). The SCN generates endogenous neural

signal, and adjusts the duration of this signal in accord to changes of the solar day length as is

detected via the retinohypothalamic tract [30,31]. While MORs are most widely expressed in the

subcortical and limbic brain regions, it is possible that daylength affects MOR signalling via the

retinohypothalamic tract, calling for in-depth studies.

In rats, we also found that seasonally alternating daylength slowed weight gain and increased blood

corticosterone levels in adult rats (in comparison with rats kept in a fixed 12-hour light cycle), in

line with previous findings [32,33]. Daylength-dependent changes in corticosterone levels may be due

to elevated stress caused by extremely long daylength, as supported by a positive effect of

daylength on blood corticosterone levels in rats (Supplementary results). In humans, exposure to

extremely long daylength in summer also increases human blood cortisol levels [34]. Blood

corticosterone levels have been linked with social stress and activity of MOR signalling in

mammals [35], where increased MOR avidity is along with lower cortisol level under stress. This is

similarly supported by our finding that brain MOR availability in rats is negatively associated with

blood corticosterone levels (Supplementary Fig. 4).

While the current study cannot disentangle the relation between corticosterone levels and MOR

signalling in humans, the findings suggest that social and physiological stress could be linked with

MOR availability. Regardless of the euphoric effect of longer daylength [1,28], cerebral MOR

availability did not increase linearly with daylength but demonstrated an inverted-U shape,

potentially due to the increased physiological or psychological stress under extremely long

daylength. This link between MOR signalling and stress is corroborated by the fact that opioid

agonists and partial agonist buprenorphine reduce symptoms of anxiety and depression in rats [36]

as well as alleviate the effects of psychological stressors such as separation distress [37]. In line with

this, human molecular and functional imaging studies have found that high opioidergic tone

suppresses hemodynamic brain responses to viewing distressing videos [38,39].

While PET being the optimal non-invasive approach to quantify *in vivo* brain MOR availability,

drawbacks exist in the current study. The human study was based on historical data, where each

subject was imaged only once, and quasi-experimental design (natural changes in daylight), as

longitudinal multi-scan studies would yield a significant radiation load. Contribution of other

seasonal factors (such as temperature) cannot however be ruled out in the human data, whereas we

verified the causal effect of daylength on brain MOR availability using animal experiments.

Furthermore, daylength was estimated using astronomical data without considering day-to-day

(e.g. cloud cover) and individual variation (e.g. time spent outside) in light exposure. Nevertheless,

we had a large sample and the results paralleled those from the experiment done in rats, albeit with

smaller effect size. The human data were also sampled from different projects and scanners, which

were corrected for potential scanner-related biases in the analyses. Technically, while

$[^{11}C]$carfentanil BP_{ND} in baseline condition is proportional to MOR density, the exact contributions

of MOR density, receptor affinity, and baseline occupancy by endogenous opioids cannot be

assessed in a single measurement [40,41], and these components cannot be differentiated in a single

scan. $[^{11}C]$carfentanil is an agonist tracer preferably binding MORs in the high-affinity state [41] and

has low test-retest variability in PET imaging [42]. Although endogenous opioids compete for

binding sites with $[^{11}C]$carfentanil [43,44], the basal opioid concentrations are often very low at least

in rats [45]. Accordingly, [^{11}C]carfentanil BP_{ND} in baseline likely reflects density of MORs. Finally,

current findings based on human data should be interpreted while keeping in mind the large

magnitude of local photoperiodic variation, which should be cautiously interpolated to regions

with lower latitudes.

Taken together, we conclude that seasonal variation in daylength influences brain MOR

availability, following an inverted U-shaped curve and peaking at 15-19 hours in humans. Given

the intimate links between MOR signalling and socioemotional behaviour [29,46], these results

suggest that the MOR system might underlie seasonal variation in human mood and social

behaviour [1,2], and imply that MOR might be a feasible target for treating seasonal affective

disorders.

Methods

Study with human PET image database

Subjects

Human study was performed using retrospective analysis of historical subjects scanned with PET

using radioligand [^{11}C]carfentanil at different times of the year (Supplementary Fig. 1). The data

were retrieved from the AIVO database (http://aivo.utu.fi) of *in vivo* molecular images hosted by

Turku PET Centre. We identified all the baseline scans from individuals with no neurologic or

psychiatric disorders who had been scanned between 2003 and 2018. The resulting data frame of

204 individuals (132 males, 72 females; age 32.4 ± 10.8 y) consists of scans from 11 research

projects and five PET scanners, with details of the subjects described in [47]. No subjects abused

alcohol or illicit drugs or used medications affecting the central nervous system.

PET image analysis

PET data were analysed and modelled using Magia toolbox (https://github.com/tkkarjal/magia)

[48], an automated processing pipeline developed at the Turku PET Centre running on MATLAB

(The MathWorks, Inc., Natick, Massachusetts, United States). Pre-processing consisted of

framewise realignment and co-registration of the PET and magnetic resonance images. Tracer

binding was quantified using BP_{ND} [49], which is the ratio of specific binding to non-displaceable

binding in tissue. BP_{ND} was estimated using simplified reference tissue model with occipital cortex

as the reference region [50,51]. Parametric BP_{ND} images were also calculated for full-volume analysis.

They were spatially normalized to MNI-space via segmentation of T1-weighted MRIs and

smoothed with an 8-mm Gaussian kernel. Regions of interest, including the reference region, were

parcellated for each subject using FreeSurfer (http://surfer.nmr.mgh.harvard.edu/).

Full-volume human data analysis

For each subject, daylength was calculated as the daytime plus civil twilight on the day when the

PET image was acquired. Civil twilight involves morning civil twilight that begins when the

geometric centre of the sun is 6° below the horizon and ends at sunrise and evening civil twilight

that begins at sunset and ends when the geometric centre of the sun reaches 6° below the horizon.

Calculation was done using R package "suncalc", where calculations were based on geographic

location of the Turku PET Centre (Turku, Finland; latitude = 60.4518, longitude = -22.2666) and

formulas provided by Astronomy Answers articles about position of the sun and the planets

(https://www.aa.quae.nl/en/reken/zonpositie.html).

We first modelled the effect of daylength on MOR availability using multiple regression, as

implemented in SPM12 (http://www.fil.ion.ucl.ac.uk/spm/). Polynomial expansion of daylength

to the second order and linear component of daylength were used as regressors. Subject age, sex,

scanner and body mass index (BMI) were used as nuisance covariates. MOR BP_{ND} values in the

daylength-sensitive brain cluster were also estimated using the MarsBaR toolbox [52], and plotted as

a function of daylength and seasons for visualization.

Region of interest analysis

We also analysed the regional MOR availability in 15 brain ROIs (based on subject-wise

FreeSurfer delineations) for both hemispheres (in total 30 ROIs), as listed in Supplementary Table

1. Pooled ROI values were analysed using linear mixed-effects model with varying intercepts for

each subject. Fixed effect factors included daylength, squared daylength, age, sex, BMI, scanner

type, and their interaction effects with ROI and brain hemisphere (right vs. right). In the single

ROI data analysis, linear regression model was used with factors including daylength, squared

daylength, age, sex, BMI, and scanner type. Regional MOR BP_{ND} was log-transformed in the

statistical analysis as previously [47]. R statistical software (version 3.6.0) using lme4 package were

used in analysing the human ROI data.

Rat model PET imaging study

Animal handling and seasonal simulation

The National Animal Experiment Board of Finland approved all the procedures and protocols

(license number ESAVI/8355/2019) in accordance with the EU Directive 2010/63/EU on the

protection of animals used for scientific purposes. Eighteen adult (age > 90 days) Sprague-Dawley

rats (twelve of them were PET imaged as described in the next section; 11 males, 7 females;

Central Animal Laboratory, University of Turku, Turku, Finland) were housed under controlled

laboratory conditions in open top cages with free access to CRM-E diet (SDS, UK) and water.

Rats were caged in groups of two or three same-sex rats. No rats were removed before finishing

the last PET imaging to keep the social environment stable and to avoid additional stress [53]. For

the rats in the experimental group (9 males, 5 females), in-house light with variable ON/OFF

duration was programmed, simulating the local seasonal change of daylength with a speeded cycle

completed in three months (Supplementary Fig. 1). Rats in the control group (2 males, 2 females)

were kept in a different room with constant daylength cycle (12h ON/ 12h OFF, no twilight), with

all other conditions the same with the experimental group. The same type of LED lighting was

used for control groups and the experimental groups.

PET imaging and processing

Twelve out of eighteen rats (experimental group: 6 males, 3 females; control group: 2 males, 1

female) were studied with dynamic [^{11}C]carfentanil PET imaging for 3-4 times under isoflurane

anaesthesia. Radiotracers were divided by three rats at each scanning day (2 males, 1 female), and

via using two different scanners to maximize the data collection. Due to larger body size, Inveon

Multimodality PET/CT (Siemens Medical Solutions, Knoxville, TN, USA) was used for imaging

the male rats (two rats each time), whereas Molecubes PET/CT (Molecubes, Gent, Belgium) was

used for the female rats. In total, there were 42 PET/CT scans. Rats were weighed on the scanning

day. For the Inveon Multimodality PET/CT scanner, with relatively lower resolution and

sensitivity, the aimed dosage of radiotracer was 5 MBq, while for Molecubes PET/CT scanner rats

were scanned with aimed dosage of 1 MBq. Accordingly, sex effects cannot be separated from

scanner type and dosage dependent effects.

Dynamic PET images were analysed using Carimas software (version 2.10.3.0) developed at the

Turku PET Centre, Finland. The PET data sets were reconstructed into 20 time frames using

OSEM3D algorithm: 6 × 0.5 minutes, 3 × 1 minutes, 4 × 3 minutes, and 7 × 6 minutes. We first

created a template of structural CT image and brain atlas. In the template, a rat's skull CT image

was aligned with the Waxholm Space Atlas of the Sprague Dawley rat brain [54] to be used as the

reference for the normalization of the brain scans. We first visually checked that the PET and CT

images were in alignment. Next, PET and CT images were locked regarding their positions using

Carimas. Individual skull images were then manually realigned and resized to the template CT

image. Once the skull image was scaled with the CT image, the CT and the PET images were

rigidly co-registered with the CT template (Supplementary Fig. 2).

Simplified reference tissue model was used for estimating of non-displaceable MOR binding

potential (BP_{ND}) in the neocortex, striatum and thalamus [49]. Cerebellum was used as the reference

region because it is devoid of MOR in rats (see Supplementary Fig. 5 for *ex vivo* biodistribution

results). We excluded the first five minutes from model fitting to rule out confounding perfusion

effects. Time-activity curves (TACs) from brain regions of interest (thalamus, striatum and

neocortex) defined by the Waxholm atlas were then extracted for data analysis.

Serum sampling and measurement of corticosterone levels

Blood samples were collected by lateral tail vein puncture at 9:00-11:00 am. For experimental

group rats, samples included collections conducted one day prior to the PET imaging (30 samples

out of 33). For the control group, sample collections were randomly conducted. Altogether, 51

blood samples were obtained (18 for the control group). The samples were centrifuged at 7000

rpm for 1.5 minutes and the serum was stored at -20°C. Levels of serum corticosterone were

measured using an ELISA kit (Enzo Life Sciences, Ann Arbor, MI, USA).

Statistical analysis

Data were analysed with R (version 3.6.0) using lme4 package [55]. To estimate the effects of

daylength on MOR availability, we used a linear mixed-effects model with varying intercepts for

each rat. Fixed effect factors included daylength, squared daylength, age, sex, group, and the

interaction effects between regions of interests (ROI) and all fixed effect factors were included.

Reference

1.   Harmatz, M. G. *et al.* Seasonal variation of depression and other moods: A longitudinal

approach. *J. Biol. Rhythms* (2000) doi:10.1177/074873000129001350.

2.   Lam, R. W. & Levitan, R. D. Pathophysiology of seasonal affective disorder: a review. *J*

Psychiatry Neurosci **25**, 469–480 (2000).

3.   Green, N. H., Jackson, C. R., Iwamoto, H., Tackenberg, M. C. & McMahon, D. G.

Photoperiod programs dorsal raphe serotonergic neurons and affective behaviors. *Curr. Biol.*

(2015) doi:10.1016/j.cub.2015.03.050.

4.   Lambert, G. W., Reid, C., Kaye, D. M., Jennings, G. L. & Esler, M. D. Effect of sunlight

and season on serotonin turnover in the brain. *Lancet* (2002) doi:10.1016/S0140-

6736(02)11737-5.

5.   Nummenmaa, L. & Tuominen, L. Opioid system and human emotions. *British Journal of*

Pharmacology (2018) doi:10.1111/bph.13812.

6.   Tung, A., Wong, C. & Fairbairn, N. Perioperative Management of Opioid Use Disorder

Patients on Buprenorphine/Naloxone. (2018).

7.   Davis, J. M., Searles, V. B., Severtson, S. G., Dart, R. C. & Bucher-Bartelson, B. Seasonal

variation in suicidal behavior with prescription opioid medication. *J. Affect. Disord.* (2014)

doi:10.1016/j.jad.2014.01.010.

8.   Gross-Isseroff, R., Dillon, K. A., Israeli, M. & Biegon, A. Regionally selective increases in

μ opioid receptor density in the brains of suicide victims. *Brain Res.* (1990)

doi:10.1016/0006-8993(90)91301-V.

9.   Gabilondo, A. M., Javier Meana, J. & García-Sevilla, J. A. Increased density of μ-opioid

receptors in the postmortem brain of suicide victims. *Brain Res.* (1995) doi:10.1016/0006-

8993(95)00333-L.

10.   Coid, J., Allolio, B. & Rees, L. H. RAISED PLASMA METENKEPHALIN IN PATIENTS

352   WHO HABITUALLY MUTILATE THEMSELVES. *Lancet* (1983) doi:10.1016/S0140-

353   6736(83)90572-X.

354   11.   Woo, J. M., Okusaga, O. & Postolache, T. T. Seasonality of suicidal behavior. *International*

355   *Journal of Environmental Research and Public Health* (2012) doi:10.3390/ijerph9020531.

356   12.   Tuulari, J. J. *et al.* Feeding releases endogenous opioids in humans. *J. Neurosci.* (2017)

357   doi:10.1523/JNEUROSCI.0976-17.2017.

358   13.   Nummenmaa, L. *et al.* μ-opioid receptor system mediates reward processing in humans.

359   *Nat. Commun.* **9**, 1500 (2018).

360   14.   Karlsson, H. K. *et al.* Obesity Is Associated with Decreased -Opioid But Unaltered

361   Dopamine D2 Receptor Availability in the Brain. *J. Neurosci.* (2015)

362   doi:10.1523/JNEUROSCI.4744-14.2015.

363   15.   Berridge, K. C. 'Liking' and 'wanting' food rewards: Brain substrates and roles in eating

364   disorders. *Physiol. Behav.* (2009) doi:10.1016/j.physbeh.2009.02.044.

365   16.   Shahar, D. R. *et al.* Changes in dietary intake account for seasonal changes in cardiovascular

366   disease risk factors. *Eur. J. Clin. Nutr.* (1999) doi:10.1038/sj.ejcn.1600761.

367   17.   Ma, Y. *et al.* Seasonal variation in food intake, physical activity, and body weight in a

368   predominantly overweight population. *Eur. J. Clin. Nutr.* (2006)

369   doi:10.1038/sj.ejcn.1602346.

370   18.   Tortonese, D. J. Interaction between Hypothalamic Dopaminergic and Opioidergic Systems

371   in the Photoperiodic Regulation of Pulsatile Luteinizing Hormone Secretion in Sheep.

372   *Endocrinology* (1999) doi:10.1210/en.140.2.750.

19.   Lincoln, G. A., Ebling, F. J. P. & Martin, G. B. Endogenous opioid control of pulsatile LH

secretion in rams: Modulation by photoperiod and gonadal steroids. *J. Endocrinol.* (1987)

doi:10.1677/joe.0.1150425.

20. Mukherjee, A. & Haldar, C. Effect of Naltrexone on photoperiodic regulation of testicular

steroidogenesis in adult golden hamster, Mesocricetus auratus. *Gen. Comp. Endocrinol.*

(2016) doi:10.1016/j.ygcen.2015.10.001.

21. Meyza, K. Z. & Sotowska-Brochocka, J. Photoperiod affects distribution of dynorphin A in

the brain of Siberian hamster. *Acta Neurobiol. Exp. (Wars).* (2006).

22. Nizielski, S. E., Levine, A. S., Morley, J. E., Hall, K. A. & Gosnell, B. A. Seasonal variation

in opioid modulation of feeding in the 13-lined ground squirrel. *Physiol. Behav.* (1986)

doi:10.1016/0031-9384(86)90375-6.

23. Aronen, E. T., Fjällberg, M., Paavonen, E. J. & Soininen, M. Day length associates with

activity level in children living at 60 degrees north. *Child Psychiatry Hum. Dev.* (2002)

doi:10.1023/A:1017956706208.

24. Hiltunen, L., Suominen, K., Lönnqvist, J. & Partonen, T. Relationship between daylength

and suicide in Finland. *J. Circadian Rhythms* (2011) doi:10.1186/1740-3391-9-10.

25. Lindblom, N., Heiskala, H., Kaski, M., Leinonen, L. & Laakso, M. L. Sleep fragmentation

in mentally retarded people decreases with increasing daylength in spring. *Chronobiol. Int.*

(2002) doi:10.1081/CBI-120002880.

26. Friborg, O., Bjorvatn, B., Amponsah, B. & Pallesen, S. Associations between seasonal

variations in day length (photoperiod), sleep timing, sleep quality and mood: A comparison

between Ghana (5°) and Norway (69°). *J. Sleep Res.* (2012) doi:10.1111/j.1365-

2869.2011.00982.x.

27. Saarijärvi, S., Lauerma, H., Helenius, H. & Saarilehto, S. Seasonal affective disorders

among rural Finns and Lapps. *Acta Psychiatr. Scand.* (1999) doi:10.1111/j.1600-0447.1999.tb07206.x.

28. Magnusson, A. & Boivin, D. Seasonal Affective Disorder: An overview. *Chronobiology International* (2003) doi:10.1081/CBI-120019310.

29. Lutz, P. E., Courtet, P. & Calati, R. The opioid system and the social brain: implications for depression and suicide. *Journal of Neuroscience Research* (2020) doi:10.1002/jnr.24269.

30. Schwartz, W. J., De La Iglesia, H. O., Zlomanczuk, P. & Illnerová, H. Encoding Le Quattro Stagioni within the mammalian brain: Photoperiodic orchestration through the suprachiasmatic nucleus. *Journal of Biological Rhythms* (2001) doi:10.1177/074873001129002024.

31. Jacob, N., Vuillez, P., Lakdhar-Ghazal, N. & Pévet, P. Does the intergeniculate leaflet play a role in the integration of the photoperiod by the suprachiasmatic nucleus? *Brain Res.* (1999) doi:10.1016/S0006-8993(99)01324-4.

32. Mariné-Casadó, R. *et al.* The exposure to different photoperiods strongly modulates the glucose and lipid metabolisms of normoweight fischer 344 rats. *Front. Physiol.* (2018) doi:10.3389/fphys.2018.00416.

33. Deibel, S. H., Hong, N. S., Himmler, S. M. & McDonald, R. J. The effects of chronic photoperiod shifting on the physiology of female Long-Evans rats. *Brain Res. Bull.* (2014) doi:10.1016/j.brainresbull.2014.03.001.

34. Vondrašová, D., Hájek, I. & Illnerová, H. Exposure to long summer days affects the human melatonin and cortisol rhythms. *Brain Res.* (1997) doi:10.1016/S0006-8993(97)00358-2.

35. Chong, R. Y. *et al.* The mu-opioid receptor polymorphism A118G predicts cortisol responses to naloxone and stress. *Neuropsychopharmacology* (2006)

doi:10.1038/sj.npp.1300856.

36.   Falcon, E. *et al.* Antidepressant-like effects of buprenorphine are mediated by kappa opioid

receptors. *Neuropsychopharmacology* (2016) doi:10.1038/npp.2016.38.

37.   Herman, B. H. & Panksepp, J. Effects of morphine and naloxone on separation distress and

approach attachment: Evidence for opiate mediation of social affect. *Pharmacol. Biochem.*

Behav. (1978) doi:10.1016/0091-3057(78)90167-3.

38.   Karjalainen, T. *et al.* Dissociable roles of cerebral μ-opioid and type 2 dopamine receptors

in vicarious pain: A combined PET-fMRI study. *Cereb. Cortex* **27**, 4257–4266 (2017).

39.   Karjalainen, T. *et al.* Opioidergic Regulation of Emotional Arousal: A Combined PET–

fMRI Study. *Cereb. Cortex* (2018) doi:10.1093/cercor/bhy281.

40.   Mintun, M. A., Raichle, M. E., Kilbourn, M. R., Wooten, G. F. & Welch, M. J. A

quantitative model for the in vivo assessment of drug binding sites with positron emission

tomography. *Ann. Neurol.* (1984) doi:10.1002/ana.410150302.

41.   Henriksen, G. & Willoch, F. Imaging of opioid receptors in the central nervous system.

Brain (2008) doi:10.1093/brain/awm255.

42.   Hirvonen, J. *et al.* Measurement of central μ-opioid receptor binding in vivo with PET and

[11C]carfentanil: A test-retest study in healthy subjects. *Eur. J. Nucl. Med. Mol. Imaging*

(2009) doi:10.1007/s00259-008-0935-6.

43.   Quelch, D. R., Katsouri, L., Nutt, D. J., Parker, C. A. & Tyacke, R. J. Imaging endogenous

opioid peptide release with [^{11}C]carfentanil and [^{3}H]diprenorphine: Influence of agonist-

induced internalization. *J. Cereb. Blood Flow Metab.* (2014) doi:10.1038/jcbfm.2014.117.

44.   Saanijoki, T. *et al.* Opioid release after high-intensity interval training in healthy human

subjects. *Neuropsychopharmacology* (2018) doi:10.1038/npp.2017.148.

45.   Maidment, N. T., Brumbaugh, D. R., Rudolph, V. D., Erdelyi, E. & Evans, C. J.

Microdialysis of extracellular endogenous opioid peptides from rat brain in vivo.

Neuroscience (1989) doi:10.1016/0306-4522(89)90407-7.

46.   Becker, J. A. *et al.* Autistic-Like Syndrome in Mu Opioid Receptor Null Mice is Relieved

by Facilitated mGluR4 Activity. *Neuropsychopharmacology* (2014)

doi:10.1038/npp.2014.59.

47.   Kantonen, T. *et al.* Interindividual variability and lateralization of μ-opioid receptors in the

human brain. *Neuroimage* (2020) doi:10.1016/j.neuroimage.2020.116922.

48.   Karjalainen, T. *et al.* Magia: Robust Automated Image Processing and Kinetic Modeling

Toolbox for PET Neuroinformatics. *Front. Neuroinform.* (2020)

doi:https://doi.org/10.3389/fninf.2020.00003.

49.   Innis, R. B. *et al.* Consensus nomenclature for in vivo imaging of reversibly binding

radioligands. *Journal of Cerebral Blood Flow and Metabolism* (2007)

doi:10.1038/sj.jcbfm.9600493.

50.   Lammertsma, A. A. & Hume, S. P. Simplified reference tissue model for PET receptor

studies. *Neuroimage* (1996) doi:10.1006/nimg.1996.0066.

51.   Frost, J. J. *et al.* Multicompartmental analysis of [11C]-carfentanil binding to opiate

receptors in humans measured by positron emission tomography. *J. Cereb. Blood Flow*

Metab. (1989) doi:10.1038/jcbfm.1989.59.

52.   Brett, M., Anton, J.-L. L., Valabregue, R. & Poline, J.-B. Region of interest analysis using

an SPM toolbox - Abstract Presented at the 8th International Conference on Functional

Mapping of the Human Brain, June 2-6, 2002, Sendai, Japan. *Neuroimage* (2002)

doi:10.1016/S1053-8119(02)90010-8.

53.   Beery, A. K. & Kaufer, D. Stress, social behavior, and resilience: Insights from rodents.

467   *Neurobiology of Stress* (2015) doi:10.1016/j.ynstr.2014.10.004.

468   54.   Papp, E. A., Leergaard, T. B., Calabrese, E., Johnson, G. A. & Bjaalie, J. G. Waxholm Space

469   atlas   of   the   Sprague   Dawley   rat   brain.   *Neuroimage*   (2014)

470   doi:10.1016/j.neuroimage.2014.04.001.

471   55.   Bates, D., Mächler, M., Bolker, B. M. & Walker, S. C. Fitting linear mixed-effects models

472   using lme4. *J. Stat. Softw.* (2015) doi:10.18637/jss.v067.i01.

Conflict of interest

The authors declare no conflict of interest.

Author contribution

LS, JT, EY, AR designed the rat imaging experiment. LS analysed the data under supervision of JT, AR, and LN. All authors wrote the manuscript. HL, AH, JV, SH conducted the rat imaging experiments under supervision of EY and AR. KE and ES measured the corticosterone levels. JI, TKarjalainen, TKantonen, PN, JH, VK, KK, JH, HS, JOR, LN contributed to construction of the human brain database. All the authors discussed the results and commented on the manuscript.

Acknowledgements

The study was supported by the Academy of Finland (grant #317680 to JT, #294897 and #304385 to LN, and #310962 to JOR), Sigrid Juselius Foundation, and Jane and Aatos Erkko Foundation (to AR). LS is supported by Turku Collegium for Science and Medicine, University of Turku. TKantonen is supported by Finnish Cultural Foundation (Southwest Finland Fund) and Emil Aaltonen Foundation. JOR is supported also by the Sigrid Juselius Foundation and Finnish State Research Funding. We thank Sauli Piirola, for help with Carimas scripting, and Elina Kahra for

489   her assistance in laboratory analysis of blood corticosterone levels. We thank Paulina Chrusciel,

490   Ella Kujala, Malgorzata Major and Joonas Khabbal from UTUCAL for assisting lighting planning

491   and blood sampling.