Genome-wide CRISPR-Cas9 screen reveals the importance of the heparane sulphate

pathway and the conserved oligomeric golgi complex for synthetic dsRNA uptake and

Sindbis virus infection

Olivier Petitjean[1,][*], Erika Girardi[1,][*], Richard Patryk Ngondo[2], Vladimir Lupashin[3], Sébastien

Pfeffer[1,][‡]

1 Université de Strasbourg, Architecture et Réactivité de l'ARN, Institut de biologie moléculaire

et cellulaire du CNRS, 2 allée Konrad Roentgen, 67084 Strasbourg France

2 Université de Strasbourg, Institut de Biologie Moléculaire des Plantes du CNRS, 12 rue du

Général Zimmer, 67084 Strasbourg France

3 University of Arkansas for Medical Sciences, 4301 West Markham Street, Little Rock,

Arkansas 72205, United States of America

* These authors contributed equally

‡ To whom correspondence should be addressed: spfeffer@unistra.fr

Abstract

Double stranded RNA (dsRNA) is the hallmark of many viral infections. dsRNA is produced either by RNA viruses during replication or by DNA viruses upon convergent transcription. Synthetic dsRNA is also able to mimic viral-induced activation of innate immune response and cell death. In this study, we employed a genome-wide CRISPR-Cas9 loss of function screen based on cell survival in order to identify genes implicated in the host response to dsRNA. By challenging HCT116 human cells with either synthetic dsRNA or Sindbis virus (SINV), we identified the heparan sulfate (HS) pathway as a crucial factor for dsRNA entry and we validated SINV dependency on HS. Interestingly, we uncovered a novel role for COG4, a component of the Conserved Oligomeric Golgi (COG) complex, as a factor involved in cell survival to both dsRNA and SINV in human cells. We showed that COG4 knock-out led to a decrease of extracellular HS, specifically affected dsRNA transfection efficiency and reduced viral production, explaining the increased cell survival of these mutants.

Importance

When facing a viral infection, the organism has to put in place a number of defense mechanisms in order to clear the pathogen from the cell. At the early phase of this preparation for fighting against the invader, the innate immune response is triggered by the sensing of danger signals. Among those molecular cues, double-stranded (dsRNA) is a very potent inducer of different reactions at the cellular level that can ultimately lead to cell death. Using a genome-wide screening approach, we set to identify genes involved in dsRNA entry, sensing and apoptosis induction in human cells. This allowed us to determine that the heparan sulfate pathway and the Conserved Oligomeric Golgi complex are key determinants allowing entry of both dsRNA and viral nucleic acid leading to cell death.

Introduction

Upon infection by a virus, numerous mechanisms are put in place at the cellular level to raise the alarm and get rid of, or at least limit, the invader. One of the first barrier that the virus has to overcome to is to enter the cell by taking advantage of wide diversity of ubiquitous or cell-specific cellular receptors. In addition to protein receptors, glycosaminoglycans present at the cell surface also represent crucial factors for efficient viral attachment and entry (1). Glycosaminoglycans, and more precisely heparan sulfates are ubiquitously expressed in human cells. They possess a global negative charge that is able to interact electrostatically with the basic residues that are exposed by viral surface glycoproteins. This allows viruses to increase their concentration at cell surface and so the possibility to interact with their specific entry receptor (2). For instance, alphaviruses such as Semliki Forest virus (SFV) and Sindbis virus (SINV) are enveloped positive-strand RNA viruses that contain two glycoproteins at the envelope, the proteins E1 and E2. E2 is involved in the interaction of the virus particle to the cell surface (3, 4), while E1 serves in the fusion process (5).

Once inside the cell, the replication of the viral genome represents another critical step to trigger the antiviral immune response. Double-stranded (ds) RNA is a ubiquitous pathogen-associated molecular pattern (PAMP) recognized by the cellular machinery, which can arise as a replication intermediate for viruses with an RNA genome, or from convergent transcription for DNA viruses (6). In mammals, dsRNA recognition is driven by specific receptors including the cytoplasmic RIG-Like Receptors (RLRs) and endosomal Toll Like Receptors (TLRs) (7). Sensing of dsRNA by these receptors results in the activation of a complex signaling cascade leading to the production of type I interferon (IFN), which in turn triggers the expression of IFN stimulated genes (ISG) and the establishment of the antiviral state (8). The ultimate outcome of this vertebrate-specific antiviral response is translation arrest and cell death by apoptosis (9).

The revolution brought by the discovery of the CRISPR-Cas9 technology has provided biologists with an invaluable tool to edit the genome at will and easily perform individual gene knock-out (10). This technique is perfectly suited to perform genome-wide screens in a relatively fast and easy to implement manner, especially when the readout is based on cell survival. For this reason, numerous CRISPR-Cas9 loss of function screens have been performed based on cell survival after infection with different viruses (11–13). These approaches allowed the identification of novel virus specific as well as common factors involved in antiviral defense mechanisms or in cellular permissivity to virus infection.

Here, we chose to take advantage of the fact that dsRNA is almost always detected in virus-infected cells (6) and is a potent inducer of apoptosis to design a genome-wide screen aiming at identifying host genes that when edited resulted in increased cell survival to dsRNA and viral challenge. To this aim, we performed a CRISPR-Cas9 screen based on cell survival in HCT116 cells after either cationic lipid-based transfection of an *in vitro* transcribed long dsRNA or infection with the model alphavirus SINV, which replicates *via* a dsRNA intermediate.

Our results indicate that genes involved in limiting attachment and therefore entry, be it of the synthetic dsRNA or SINV, are vastly over-represented after selection. We validated two genes of the heparan sulfate pathway (namely *SLC35B2* and *B4GALT7*) as required for dsRNA transfectability and SINV infectivity. We also identified and characterized COG4, a component of the Conserved Oligomeric Golgi (COG) complex, as a novel factor involved in susceptibility to dsRNA and viral induced cell death linked to the heparan sulfate biogenesis pathway.

Results

Genome-wide CRISPR/Cas9 screen based on cell survival upon dsRNA transfection identify factors of the heparan sulfate pathway

In order to identify cellular genes that are involved in the cellular response to dsRNA which culminates with cell death, we performed a CRISPR/Cas9 genome-wide loss-of-function screen

in the human colon carcinoma cell line HCT116. This cell line is highly suitable for

CRISPR/Cas9 genetic screening procedures (14) and can be easily infected with SINV with

visible cytopathic effects at 24 and 48 hours post infection (hpi) (Fig. S1A). Moreover,

transfection of an *in vitro* transcribed 231 bp-long dsRNA by a cationic lipid-based transfection

reagent in HCT116 cells led to strong cell death at 24 and 48 hours post treatment (hpt) (Fig.

S1B).

We generated a Cas9-expressing HCT116 monoclonal cell line (Fig. S1C) that we stably

transduced with the human genome-wide lentiviral Brunello library composed of 76 441

sgRNAs targeting 19 114 genes, as well as about 1000 non-targeting sgRNAs as controls (15).

We then applied a positive selection by lipofecting 30 million transduced cells per replicate

with the synthetic long dsRNA and we collected the surviving cells 48h later. In parallel, the

same initial amount of stably transduced cells was left untreated as control for (Input) for each

replicate (Fig. 1A). DNA libraries from the input samples were generated, sequenced and

quality checked. In particular, we verified the sgRNA coverage by observing the presence of

the 4 guides per genes for 18 960 genes (99.2% of the genes) and 3 sgRNAs per gene for the

remaining 154 genes (0.2% of the genes) (Dataset S1).

Using the MAGeCK software (16), we assessed the normalized read count distribution of the

control and dsRNA-treated biological triplicates, which, despite a quite homogenous sgRNA

distribution, showed the presence of few outliers upon selection (Fig. 1B). We identified eight

genes that were significantly enriched with a false discovery rate lower than 1% (FDR1%).

Among those, four genes belonged to the heparan sulphate biosynthesis pathway (namely,

SLC35B2, *B4GALT7*, *EXT1* and *EXT2*) and three were components of the conserved oligomeric

golgi complex (namely *COG3*, *COG4* and *COG8*) (Fig. 1C, Dataset S2). In particular, all four

sgRNAs targeting each of *SLC35B2*, *B4GALT7, COG4* genes were enriched upon dsRNA

selection (Fig. S1D).

Heparan sulphate (HS) is a linear polysaccharide that is covalently attached to core proteins in

proteoglycans (PG) on the cell surface (for review, see Li and Kusche-Gullberg, 2016). Among

many properties, HS play a role in binding of protein ligands and as a carrier for lipases,

chemokines and growth factor (Bishop et al., 2007; Li and Kusche-Gullberg, 2016), but also as

a viral receptor (19). HS biosynthesis takes place in the Golgi, where most of the biosynthetic

enzymes are anchored to the Golgi membrane (20).

We first validated the resistance phenotype to dsRNA of *SLC35B2* and *B4GALT7*, the two top

hits identified in the screen (Fig. 1C, Fig. S1D), by generating two individual knock-out clones

for each gene by CRISPR-Cas9 editing in HCT116cas9 (Fig. S2). Knock-out of either *SLC35B2*

or *B4GALT7* genes abolished cell death induced by dsRNA lipofection compared to parental

HCT116cas9 cells, as assessed by the measurement of cell viability 48h post transfection (Fig.

1D, left part of the graph). These results demonstrated the involvement of SLC35B2 and

B4GALT7 in dsRNA-induced cell death.

The observed resistance to dsRNA in the mutants could occur at many different steps: dsRNA

liposome attachment and entry, recognition, induction of the IFN pathway or apoptosis. To test

whether the first step was affected, we employed a nucleic acid delivery method that was not

based on cationic lipid transfection. In particular, we used nucleofection (an electroporation-

based transfection method) to introduce long dsRNAs into HCT116 cells and we showed that

this approach totally restored cell death in *SLC35B2* and *B4GALT7* knock-out cells (Fig. 1D,

right part of the graph). In addition, we also performed liposome-based transfection of an *in*

vitro transcribed Cy5-labelled dsRNA in *SLC35B2* and *B4GALT7* KO cells and assessed the

Cy5 fluorescence at 48 h post-transfection by FACS analysis (Fig. 1E-F). Although the number

of Cy5 positives (Cy5+) cells was not significantly different in the *B4GALT7* KO clones and

only slightly lower in the *SLC35B2* KO cells compared to WT cells (Fig. 1E), we observed a

significant reduction of at least 80% of the median Cy5 fluoresence in both *B4GALT7* and

SLC35B2 KO cells relative to control (Fig. 1F), thereby indicating a significant drop in the

number of transfected Cy5-labeled RNA molecules per cells.

We also confirmed that liposome-based transfection of nucleic acid such as plasmidic DNA

was impaired in *SLC35B2* and *B4GALT7* KO cells, by transfecting a GFP-expressing plasmid

using Lipofectamine 2000 in wild type or knock-out cells (Fig. S3A, left panels and Fig. S3B).

Nonetheless, GFP expression could be restored in all cell lines upon nucleofection (Fig. S3A,

right panels).

To establish whether impairment of the HS synthesis is directly linked to a defect in dsRNA

entry and increased cell survival, we measured the extracellular HS levels in *SLC35B2* and

B4GALT7 KO cells. We measured a substantial reduction of the extracellular HS staining as

assessed by FACS measurement of two independent *SLC35B2* and *B4GALT7* KO clones

compared to HCT116 wild type cells (Fig. 1G). To confirm the importance of HS at the cell

surface for liposome-based transfection, we mimicked the HS-defective phenotype by

removing extracellular HS in parental HCT116cas9 cells either enzymatically (with heparinase)

or chemically (with sodium chlorate NaClO3) (Fig. S3C-D). We tested the transfectability of a

GFP-expressing plasmid by measuring either the relative number of GFP positive cells or the

relative median of GFP intensity of fluorescence by FACS analysis. Although the relative

number of GFP positive cells was not significantly reduced by heparinase treatment (Fig. S3C),

it caused a reduction in GFP intensity in HCT116cas9 treated cells, thereby recapitulating the

GFP plasmid lipofection defect observed in *SLC35B2* and *B4GALT7* KO (Fig. S3D). Moreover,

this effect correlates with the reduction of extracellular HS by enzymatic treatment quantified

by FACS (Fig. S3E), which demonstrates that extracellular HS are crucial for transfection by

lipofection. In the case of NaClO3 treatment, despite the reduction in both the relative number

of GFP positive cells (Fig. S3C) and the relative median of GFP intensity of fluorescence(Fig.

S3D) compared to control, we could not observe a correlation with a decrease in overall

extracellular HS (Fig. S3E). This could be due to the fact that while a mix of heparinase I & III

remove every kind of extracellular heparan sulfates, NaClO3 only impairs the O-sulfation (21).

Taken together, our results show that knocking out *SLC35B2* and *B4GALT7* results in reduced

levels of extracellular HS, which in turn impairs liposome-based transfectability of HCT116

cells. Moreover, the validation of these two top hits indicates that other candidates might be

suitable for further analysis and may also have an impact on the dsRNA resistance phenotype.

COG4 is involved in dsRNA-induced cell death partly via the heparan sulphate pathway

Among the significant hits of our genome-wide screen were proteins related to COG complex,

namely COG4, COG3 and COG8. The COG complex is a hetero-octameric complex containing

8 subunits (COG1-8) interacting with numerous proteins mainly involved in intra-Golgi

membrane trafficking such as vesicular coats, Rab proteins and proteins involved in SNARE

complex (22, 23). This interaction with the trafficking machinery is crucial for the proper

functionality of the Golgi apparatus and mutations in the COG complex result in severe cellular

problems such as glycosylation defect (24–27), which are due to mislocalization of recycling

Golgi enzymes (28, 29).

Since we retrieved three out of the eight COG family members in our CRISPR/Cas9 screen

suggesting their importance in dsRNA-induced cell death, we tested the effect of their

inactivation by CRISPR-Cas9. *COG4* being the most enriched COG gene in our screen, we

generated a polyclonal *COG4* KO HCT116cas9 cell line and validated their dsRNA resistance

phenotype resulting in an increase survival in response to synthetic dsRNA transfection (Fig.

S4A). We also observed a reduction in the relative number of Cy5 positives (Cy5+) cells and

the median Cy5 fluorescence in *COG4* KO cells relative to control by FACS analysis (Fig.

S4B-C).

To further confirm the involvement of the COG complex, we also tested the effect of dsRNA

transfection in previously generated HEK293T KO *COG3*, 4 and 8 cells (Fig. 2A) (30).

Interestingly, while *COG8* mutants mutants did not display a significant survival phenotype in

response to dsRNA lipofection, *COG3* and *COG4* KO HEK293 cells did. In addition, the

survival phenotype could be complemented by stable expression of a COG4-GFP construct

compared to *COG4* KO cells (Fig. 2A). Moreover, although we could not detect a decrease in

the relative number of Cy5 positives (Cy5+) cells in *COG4* KO cells relative to the controls

(Fig. 2B), the median Cy5 fluorescence in *COG4* KO cells was significantly reduced compared

to both HEK293T and COG4 rescued cells (Fig. 2C), thereby indicating a significant decrease

in the number of transfected Cy5-labeled RNA molecules per cells.

In agreement, dsRNA accumulation appeared to be significantly reduced, but still present, in

HEK293T *COG4* KO cells compared to control cells as determined by RT-qPCR analysis of

dsRNA isolated from cells 24h after transfection (Fig 2D) and this correlated with reduced IFN-

beta accumulation in HEK293T KO *COG4* cells compared to control cells (Fig. 2E).

These results indicated that dsRNA transfectability was strongly reduced but not totally

impaired in the absence of COG4 and that dsRNA could be still detected in *COG4* KO cells in

order to activate type-I IFN response.

We confirmed the reduced internalization of dsRNAs in COG4 mutant cells by transfecting

rhodamine-labeled poly(I:C), a synthetic dsRNA analog, in HEK293T *COG4* KO or WT cells

(Fig. 2F) and by counting the number of poly I:C foci per cell at 6, 12 and 24 h post transfection

(Fig. 2G). We could observe a significant reduction in rhodamine positive foci in HEK293T

COG4 KO during the time course, suggesting a defect in dsRNA internalization, which could

explain the increased survival phenotype.

In order to assess whether the *COG4* KO survival phenotype was associated with the heparan

sulphate pathway defect, we stained extracellular HS and measured the HS expression by FACS

analysis. We observed a decrease of extracellular HS in KO *COG4* cells compared to control

cells (WT and rescued), which demonstrated that the COG complex is related to the HS

biosynthesis pathway (Fig. S4D)

The reduction in extracellular HS could correlate with a decrease in transfectability and explain

the survival phenotype in KO *COG4* cells. Surprisingly however, lipofection of a GFP

expressing plasmid indicated that HEK293T *COG4* KO cells are still transfectable with a

plasmid DNA compared to control cells as observed by FACS analysis (Fig. S4E-F).

Altogether these findings indicate that the COG complex is involved in HS biosynthesis and

that removal of *COG4* results in a lower accumulation of HS at the cell surface, which most

likely translate in a reduced transfectability of dsRNA. However, as opposed to the observations

in *SLC35B2* or *B4GALT7* KO cells, the cells are still transfectable with a plasmid DNA and,

although to a lower extent, with dsRNA. Interestingly, the increased cell survival phenotype of

COG4 KO cells upon dsRNA transfection does correlate with a reduced, but still measurable,

IFN-B production.

Cell survival based genome-wide CRISPR/Cas9 screen identifies *COG4* as a permissivity

factor to SINV

SINV is a small enveloped virus with a single stranded RNA genome of positive polarity. The

virus belongs to the *Togaviridae* family, alphavirus genus and is considered as the model for

other medically important viruses such as chikungunya virus (CHIKV) and Semliki forest virus

(SFV). During its infectious cycle, SINV produces dsRNA as a replication intermediate and

induces cytopathic effects in mammalian cells, leading to cell death within 24 to 48 h post-

infection (31).

In order to identify host genes that are related to SINV-induced cell death and infection, we

performed a CRISPR/Cas9 knock-out screen in HCT116cas9 cells, which are susceptible to

this virus (Fig. 3A, Fig. S1). After transduction with the CRISPR lentiviral genome-wide

knockout library, puromycin-resistant HCT116 cells were infected with SINV-GFP at a MOI

of 0.1 and selected for cell survival. Using the MAGeCK software (16), we assessed the

normalized read count distribution of the control and SINV-infected biological triplicates,

which, despite a quite homogenous sgRNA distribution, showed the presence of few outliers

upon selection (Fig. 3B). We identified two genes that were significantly enriched with a false

discovery rate lower than 25% (FDR25%), notably *SLC35B2* and *B4GALT7* (Dataset S3, Fig.

3C). Genes of the heparan sulphate pathway have been previously found in genome-wide

CRISPR-Cas9 loss of function studies looking for factors involved in the accumulation of

viruses such as Influenza, Zika and chikungunya virus (Han et al., 2018; Li et al., 2019; Tanaka

et al., 2017). Interestingly, among the top-ranking hits, we retrieved *COG4*, which was not

previously associated with SINV infection (Fig. 3C).

To validate the involvement of *COG4* during SINV infection, we infected HEK293T, *COG4*

KO or COG4 KO rescued HEK293T cells with SINV and measured cell viability at 24, 48 and

72 hpi. The cell viability assay revealed that the *COG4* KO cells were less sensitive at early

times points of SINV infection (24 & 48 hpi), but this tendency disappeared at 72 hpi (Fig. 3D.

In agreement, the determination of viral titer by plaque assay showed that *COG4* KO HEK293T

cells produced significantly less infectious viral particles than HEK293T or *COG4*-rescued

cells at 24 hpi but not at 48 hpi, underlining a possible delay in the infection and virus-induced

cell death (Fig. 3E). We also observed that GFP accumulated to lower levels in *COG4* KO than

in WT cells both at 24 and 48 h post-infection (hpi) (Fig. S5A). Finally, we noticed that the

reduced viral production in KO *COG4* cells was associated with a reduced accumulation of

viral dsRNA in the cytoplasm, when we infected WT or *COG4* KO cells with SINV-GFP and

performed immunostaining with the anti-dsRNA J2 antibody 24 hpi (Fig. S5B). Overall, our

results indicate that COG4 expression is needed for an efficient SINV infection and that its

absence can delay the infection thereby increasing cell survival in COG4 KO cells.

Discussion

Several CRISPR/Cas9 screens aiming at identifying factors required for infection by specific

viruses have been described in the literature, but to our knowledge, none has been designed to

look at the effect of the only common factor between all those viruses, *i.e.* dsRNA. Here, we

used the Brunello sgRNA lentiviral library to screen for genes involved in HCT116 cells

survival to synthetic dsRNA transfection and to SINV infection. This allowed us to identify

components of the heparan sulphate biosynthesis pathway and of the COG complex as critical

host factors in the cellular response to above mentioned challenges. It has been already reported

that cell-survival-based CRISPR screens for viral host factors are biased toward genes linked

to the initial steps of the infection, and even more so to viral entry (11, 35). Thus, in our case,

HS is a well-known factor required for SINV entry due to the virus adaptation to cell culture

(36). We also retrieved genes of the HS pathway in our dsRNA-based screen, and we confirmed

the importance of extracellular HS for dsRNA-induced toxicity. This is mostly due to a decrease

of transfectability of the cells when HS are missing, which is linked to the fact that the

polyplexes used for transfection are positively charged and can interact electrostatically with

glycosaminoglycans (37, 38). Our work also points out to the limitations of survival-based

CRISPR-Cas9 screens, *i.e.* the selection pressure is too strong to allow the identification of

genes intervening after the entry step, thereby making the screen less sensitive. New strategies

should be designed to overcome this problem, such as using fluorescent-based cell sorting in

order to be less stringent in the selection.

In addition to the HS pathway, we identified members of the COG complex, and more

specifically *COG4*, as factors involved in dsRNA transfection and SINV infection. Loss of

function COG4 mutant cells show a dsRNA-resistant phenotype as well as a reduction in

extracellular HS expression, similar to previously published reports for other COG proteins (34,

39). Surprisingly, even if the removal of COG4 expression results in a defect in the HS pathway,

we were still able to transfect the KO *COG4* cell line either with a plasmid encoding GFP or,

although to a lesser extent, dsRNA. In addition, the dsRNA molecules that are able to enter into

HEK293T KO *COG4* are still sufficient to induce the IFN-β mRNA production indicating that

the innate immune response is still functional in these mutant background. Nonetheless, cell

death induced by dsRNA appears to be lower in *COG4* KO cells, most likely due to a delay in

the infection.

Future work will be needed in order to assess whether this phenotype upon SINV infection is

only correlated with a defect in HS biogenesis in *COG4* mutants or to other function of COG4.

The fact that *COG4* KO cells seem to better survive to both synthetic dsRNA transfection and

viral infection opens several interesting perspectives. Indeed, since the COG complex is related

to glycosylation and membrane trafficking (25, 40–42), deficiency in one or more of its

components could potentially lead to a glycosylation and/or subcellular localization defect of

components of the innate immune response or of the apoptosis pathway, although this

possibility remains to be formally proven. The difference of transfectability of plasmid DNA

and dsRNA in *COG4* KO cells is also intriguing and could indicate that different kinds of

nucleic acids do not necessarily use the exact same routes to enter the cells upon liposome-

based transfection. Finally, there could be other defects linked to COG deficiencies(40, 43) that

could account for our observations and elucidating those will require further work. It is

particularly interesting that *COG3* and *COG4* knock-out cells display a dsRNA-induced cell

death resistance phenotype, while *COG8* mutants do not. This implies that only part of the COG

complex is involved in dsRNA uptake.

In conclusion, our work uncovered *COG4* as a new player in the HS production, which is

required for both SINV infection and dsRNA transfection. These results also highlight that

synthetic dsRNA is a powerful tool to identify novel key pathways of the cellular response to

RNA viruses.

Materials and Methods

Cell culture and virus

HCT116, HEK293T, HEK293T COG KOs, BHK21 and Vero R cells were maintained in

Dulbecco's Modified Eagle Medium (DMEM) 4.5 g/L glucose (Gibco, Thermo Fisher

Scientific Inc.) supplemented with 10% fetal bovine serum (FBS) (Takara) in a humidified

atmosphere of 5% CO_2 at 37°C. HEK293T *COG3*, *COG8* and *COG4* KO and *COG4* KO stably

rescued with COG4-GFP were described previously (44, 45). HCT116cas9 and HEK293T

COG4rescued were maintained in the same medium with addition of 10 μg/mL blasticidine.

HCT116 KO clones (*COG4*, *SLC35B2*#1 and #2, *B4GALT7*#1 and #2 were maintained in the

same medium with addition of 10 μg/mL blasticdine and 1 μg/mL puromycine.

SINV wild type (SINV) or expressing the green fluorescent protein (SINV-GFP) were produced

as previously decribed (46) in BHK21 cells. In the later, the promoter of SINV subgenomic

RNA was duplicated and inserted at the 3' extremity of the viral genome, the GFP sequence

was then inserted after this new promoter. Cells were infected with either SINV WT or SINV

GFP at a MOI of 10^{-1} and samples were harvested at 24 or 48 hours post-infection (hpi).

Standard plaque assay

10-fold dilutions of the viral supernatant were prepared. 50 μl aliquots were inoculated onto

Vero R cell monolayers in 96-well plates for 1 hour. Afterwards, the inoculum was removed

and cells were cultured in 2.5% carboxymethyl cellulose for 72 h at 37°C in a humidified

atmosphere of 5% CO_2. Plaques were counted manually under the microscope. For plaque

visualization, the medium was removed, cells were fixed with 4% formaldehyde for 20 min and

stained with 1x crystal violet solution (2% crystal violet (Sigma-Aldrich), 20% ethanol, 4%

formaldehyde).

J2 immunostaining

HEK293T or KO *COG4* HEK293T cells were plated on millicell EZ slide (Millipore) and were

infected with SINV at an MOI of 0.1 for 24h. Cells were fixed with 4% formaldehyde diluted

in PBS 1x for 10 min at room temperature, followed by incubation in blocking buffer (0.2%

Tween X-100; PBS 1x; 5% normal goat serum) for 1 h. J2 antibody (Scicons) diluted in

blocking buffer at 1:1000 was incubated over night at 4°C. Between each steps, cells were

washed with PBS 1X-Tween 0.2%. Secondary antibody goat anti-mouse Alexa 594

(ThermoFisher) diluted at 1:1000 in PBS 1x-Tween 0.2% were incubated for 1 h at room

temperature. After DAPI staining (1:5000 dilution in PBS 1x for 5 min), slides were mounted

with a coverslip over anti-fading medium and observed by epifluorescence microscopy using

the BX51 (Olympus) microscope with a X 40 objective.

Generation of HCT116cas9 line

The HCT116cas9 cells, expressing human codon-optimized S. pyogenes Cas9 protein, were

obtained by transducing wild type HCT116 colorectal carcinoma cell line (ATCC® CCL-

247™) with lentiCas9-Blast lentiviral vector (Addgene #52962). Briefly, wild-type HCT116

cells were cultured in standard DMEM (GIBCO) medium supplemented with 10% Fetal bovine

serum (FBS, Gibco) and 100 U/mL of penicillin-Streptomycin (Gibco) at 37°C in 5% CO_2. The

cells were transduced at 80% confluency in a 10 cm tissue culture plate, using 6 mL lentiviral

supernatant supplemented with 4μg/mL of polybrene (H9268, sigma) for 6 hours. The

transduction medium was replaced with fresh growing medium for 24h before starting the

selection. Transduced HCT116cas9 cells were selected for 10 days and maintained in growing

medium supplemented with 10μL/mL of Blasticidin (Invivogen).

High titer lentiviral sgRNA library production

The production of high titer Human sgRNA Brunello lentiviral library which contains 4 sgRNA

per gene (Doench et al 2018) (Addgene #73178), was performed by transfecting HEK293T

cells in five 15 cm tissue culture plates using PEI (Polyethylenimin Linear, MW 25,000, 23966-

1-A, Polysciences) transfection method (Boussif et al 1995). Briefly, for each 15 cm plate

containing 20 mL of medium, 10μg of sgRNA library, 8μg of psPAX2 and 2μg of pVSV diluted

in 500μL of NaCl 150 mM were combined with 40μL of PEI (1.25 mg/mL) dissolved in 500μL

of NaCl 150 mM. The mix was incubated 30 minutes at room temperature and the formed

complexes were added dropwise on the cells. After 6 hours, the medium was replaced and the

viral supernatant was collected after 48 hours and after 72 hours. The supernatant was filtered

through a 0.45μm PES filter and the viral particles concentrated 100 times using Lenti-X™

Concentrator (Takara) before storage at -80. The viral titer was established by counting

puromycin resistant colonies formed after transducing HCT116 cells with serial dilutions of

viral stock. HCT116cas9 cells were transduced with lentivirus-packaged Brunello sgRNA

library at a MOI of 0.3. The lentiviral library has been sequenced to verify that all the lenti-

sgRNA are represented.

Genome-wide CRISPR/Cas9 knock-out screens

For each replicate (n=3), 5×10^6 stably transduced cells/dish were seeded in 6x15cm^2 plate in

order to keep a 300X representativity of the sgRNA library. Untreated samples (Input) were

collected as controls. One day later, cells were either lipofected with $1\,\mu$g/mL dsRNA-Citrine

or infected with SINV at MOI of 0.1 and cultured at 37°C 5% CO_2. Cells were washed with

PBS 1X 48 hours post treatment, to remove dead cells and fresh media was added to surviving

clones. Cells were expanded and all cells were collected 6 days post dsRNA transfection and

18 days post SINV infection.

Genomic DNA was isolated by resuspending the cell pellet in 5 mL of resuspension buffer (50

mM Tris-HCl pH 8.0, 10 mM EDTA, 100 µg/ml RNaseA), 0,25 mL of 10% SDS was added

and incubated 10 mins at RT after mix. After incubation, the sample was sonicated and

incubated 30 mins at RT with 10 uL of proteinase K (10mg/mL). 5 mL of

Phenol/Chloroform/Isoamyl Alcohol solution was added and followed by a spin down 60 mins

at 12000g/20°C. Upper phase was transferred into a new tube and 500uL of NaAc 3M and 5

mL of isopropanol was added then incubation over night at RT and followed by a centrifuge 30

mins/20°C/12000g. Pellet was washed using EtOH and dissolve in H2O.

Illumina P5 and P7-barcoded adaptors were added by PCR on gDNA samples according to the

GoTaq protocol (Promega). PCR amplicons were gel purified and sequenced on a HiSeq4000

(Illumina) to obtain about 30 million reads for each samples. Enrichment of sgRNAs was

analysed using MaGeCK with default parameters (16). Primers used to generate the PCR

products are listed in Table S1. The results of the dsRNA and SINV screen are available in

Dataset S1 and S2, respectively

Generation of monoclonal SLC35B2 and B4GALT7 and polyclonal COG4 knock-out

HCT116 cell lines

The sgRNA expression vectors targeting either SLC35B2,B4GALT7 or COG4 (sgRNA

sequences selected were the 2 most enriched sgRNA from the Brunello library in the dsRNA

screen) genes were produced by annealing the "sense" and "antisense" oligonucleotides (Table

S1) at a concentration of 10 µM in 10 mM Tris-HCl (pH 8.0), 50 mM MgCl2 in 100 μL. The

mixture was incubated at 95 ° C for 5 minutes and then allowed to cool down to room

temperature. The oligonucleotide duplex thus formed was cloned into the BbsI restriction site

of the plasmid pKLV-U6gRNA (BbsI)-pGKpuro2ABFP (Addgene#62348). The lentiviral

supernatant from single transfer vector was produced by transfecting HEK293T cells (ATCC®

CRL-3216™) with the transfer vector, psPAX2 packaging plasmid (Addgene #12260) and the

pVSV envelope plasmid (Addgene #8454) in proportion 5:4:1 using Lipofectamine™ 2000

(Themofisher) reagent according to manufacturer protocol. Standard DMEM (GIBCO) medium

supplemented with 10% Fetal bovine serum (FBS, Gibco) and 100 U/mL of penicillin-

Streptomycin (Gibco) was used for growing HEK293T cells and for lentivirus production. One

10 cm plate of HEK293T cells at 70-80% confluency was used for the transfection. The medium

was replaced 8 hours post-transfection. After 48 hours the medium containing viral particles

was collected and filtered through a 0.45μm PES filter. The supernatant was directly used for

transfection or stored at -80°C. A 6 well plate of HCT116cas9 cells at 80% confluency was

transduced using 600 μL of lentiviral supernatant (300 μL of each lentivirus produced for each

duplexes) supplemented with 4 μg/mL polybrene (Sigma) for 6 hours. The transduction media

was then changed with fresh DMEM for 24 hours then the transduced cells were selected using

DMEM containing 10 μg/mL Blasticidin (Invivogen) and 1 μg/mL Puromycin (Invivogen).

Genomic DNA was isolated from individual colonies and KO clones were screened by PCR

(primers in Table S1). The expected WT band for SLC35B2 is 469 bp and the mutant band 132

bp. For B4GALT7, the WT band is 341 bp and mutant band 180 bp. For laboratory purposes,

the SLC35B2 clones have been generated into HCT116cas9 cells that are expressing mCherry

and Citrine due to integration of miReporter-PGK (Addgene#82477).

Nucleic acids delivery

Transfection using lipofectamine2000 (InvitroGen – 11668019) were performed following

manufacturer's instructions. For nucleofection, cells were nucleofected using Nucleofector SE

solution and reagent into Nucleocuvette following manufacturer's instructions using 4D-

Nucleofector System (Lonza). The cell number and nucleic acid amounts are indicated in each

figure legends. P-EGFP-N1 (Addgene plasmid#2491) was used in transfection and

nucleofection experiments as a control.

Viability assay

PrestoBlue (PB) reagent (ThermoFisher – A13261) was used for viability assay according to

the manufacturer's protocol. After 24 to 48 hours post treatment (SINV, dsRNA

transfection/nucleofection), cells were incubated with PB reagent and cell viability was

assessed by measuring the fluorescence (excitation 570 nm; emission 590 nm) after 20 mins

incubation using SAFAS spectrofluorometer (Xenius XC).Cell viability was expressed as a

percentage relative to untreated cells.

Heparinase & sodium chlorate treatment and heparan sulfate staining

Heparinase: $1*10^6$ cells were treated with 2U of Heparinase I and III blend from

Flavobacterium heparinum (Merck – H3917) for 1 h at 37°C 5% CO_2 in DMEM and then cells

were reverse transfected with 2 µg of GFP using lipofectamine2000 (Invitrogen) in 6 well plate.

Sodium chlorate: HCT116cas9 cells grow in 50 mM sodium chlorate (Merck – 1.06420)

DMEM 10 % FBS for at least 48 h then 150 000 cells were reverse transfected with 500 ng of

GFP using lipofectamine 2000 (Invitrogen) in 24 well plate.

24 (heparinase) or 48 (sodium chlorate) hours post treatment, cells were detach using PBS

0.02% EDTA then heparan sulfate of the cells were stained using 1:30 of F58-10E4 as primary

antibody (AMSBIO, cat#370255-S) in PBS 3% BSA for 30 to 40 minutes on ice then washed

with PBS 1% FBS and incubated with 1:30 anti-mouse Alexa Fluor 594 (Thermo, A-11032) in

PBS 3% BSA, washed twice using PBS 1% FBS then and analysed at FACS using

FACSCalibur.

dsRNA preparation

PCR fragments corresponding to 231 nts of the Citrine coding sequence were amplified from

ES-FUCCI plasmid (Addgene plasmid#62451) using primers containing T7 promoter sequence

with 2 distinct PCR fragment allowing the positive-sense or negative-sense RNA. Primers used

to generate the PCR products are listed in Table S1. The PCR fragments were produced using

DyNAzyme EXT DNA Polymerase (F-505S - Thermo Scientific) and purified using Monarch

DNA extraction (T1020L - New England Biolab). In vitro transcription (IVT) with homemade

T7 RNA polymerase was performed for 4 hours at 37°C. To label the IVT RNA, 1/10th of Cy5-

CTP (Amersham CyDye Fluorescent Nucleotides Cy5-CTP, GE Healthcare Life sciences) was

included in the IVT reaction. IVT RNA was digested with DNase I (EN0525 - Thermo

Scientific) for 30 min at 37°C and IVT product was purified, unincorporated nucleotides

removed and size checked using UV shadow (8% acrylamide-urea gel) followed by phenol-

chloroform extraction the nanodrop quantification for each strand. We then mix an equal

quantity of positive-strand and negative-strand RNA, heat for 5' at 95°C followed by slow cool

down to RT. The integrity of dsRNA is then checked by RNases T1 (EN0541 - Thermo

Scientific) and V1 (AM2275, Ambion) digestion.

Microscopy

Imaging of cells treated with dsRNA/GFP plasmid was carried out on the Observer A1 (Zeiss)

microscope and analyzed using Fiji (47). Images of cells transfected with Poly(I:C) (LMW)

Rhodamine (tlrl-piwr – Invivogen)(1,8 μg/mL, 24h post plating of 76 000 cells) into Lab-Tek

on glass coverslip (155411- Thermo Scientific) were acquired using a 100x Plan Apochromat

oil immersion NA1.4 objective on a spinning disk system Axio Observer Z1 (Zeiss) every 20

minutes for 72 hours. All pictures were acquired under the same conditions (laser power and

amplification gain) then processed with Fiji. Images of cells infected with SINV stained with

J2 antibody were carried out on the BX51 (Olympus).

FACS analysis

The cells intended for analysis by flow cytometry are recovered mechanically (PBS 0,5 mM

EDTA) or using trypsin, washed in PBS and then suspended in PBS 1 % FBS. Each acquisition

includes at least 10,000 events and is performed on the FACScalibur (BD Bioscience) device.

The data produced is processed using FlowJo software (FlowJo LLC).

RT-qPCR analysis

Total RNA was isolated using TRIzol (Invitrogen - 15596026) following manufacturer's

instructions. 1 µg of RNA was reverse transcribed using SuperScript IV Vilo (Invitrogen –

11756050) according to manufacturer's instructions. Real-time PCR was performed using

SYBR Green (Applied Biosystem – 4309155) and primers listed in Table S1 at an annealing

temperature of 60°C on a CFX96 thermal cycler (Biorad). Generated data were analysed using

the CFX Manager Software (Biorad).

Western blot analysis

Proteins were extracted using RIPA lysis buffer. Proteins were quantified by the Bradford

method and 20 to 30 µg of total protein extract was loaded on 4-20% Mini-PROTEAN®

TGX™ Precast Gels (Biorad). After transfer onto nitrocellulose membrane, equal loading was

verified by Ponceau staining. Membranes were blocked in 5% milk and probed with the

following antibodies: anti-Flag M2 (SIGMA, #F1804) and anti-GAPDH (clone 6C5, BioRad #

MCA4739P). Detection was performed using Chemiluminescent Substrate (Thermo Fisher).

Data availability

523 The CRISPR-Cas9 screen sequencing data discussed in this manuscript has been deposited on

524 NCBI's Sequence Read Archive (SRA) and has been attributed the submission ID

525 SUB8109948. It can be accessed at the following URL:

526 http://www.ncbi.nlm.nih.gov/bioproject/662202 (release pending).

Acknowledgments

The authors would like to thank members of the Pfeffer laboratory for discussion, Delphine

Richer for technical help, Dr. Jean-Daniel Fauny for help with the spinning disk microscope

and Dr. Frédéric Gros for help with FACS analysis, Dr Carla Saleh for providing us the SINV

WT and GFP clones.

This work was funded by the European Research Council (ERC-CoG-647455 RegulRNA) and

was performed under the framework of the LABEX: ANR-10-LABX-0036_NETRNA, which

benefits from a funding from the state managed by the French National Research Agency as

part of the Investments for the future program. This work has also received funding from the

People Programme (Marie Curie Actions) of the European Union's Seventh Framework

Program (FP7/2007-2013) under REA grant agreement n° PCOFUND-GA-2013-609102,

through the PRESTIGE program coordinated by Campus France (to EG), and from the French

Minister for Higher Education, Research and Innovation (PhD contract to OP). VL was

supported by the National Institutes of Health (R01GM083144).

Authors contribution

SP and EG conceived the project. SP, EG and OP designed the work and analysed the results.

OP, EG, and RPN performed the experiments. EG and OP set up the CRISPR/Cas9 screens,

RPN generated the lentivirus library and the HCT16-Cas9 cell line. OP and RPN performed the

bioinformatics analysis of the screens. OP generated the SLC35B2 and B4GALT7 KO clones.

OP generated IVT dsRNA and perform validation of cell survival. EG produced SINV-GFP

viral stock, performed the infections and analysed viral titers. OP performed FACS analyses.

EG and OP performed the immunofluorescence assays. OP analysed the live-imaging

microscopy data. VL provided the COG KO cells and antibodies. OP, EG and SP wrote the

manuscript with input from the other authors. SP and EG coordinated the work. SP assured

funding. All authors reviewed the final manuscript.

References

1. Bomsel M, Alfsen A. 2003. Entry of viruses through the epithelial barrier: pathogenic
trickery. Nat Rev Mol Cell Biol 4:57–68.

2. Rusnati M, Vicenzi E, Donalisio M, Oreste P, Landolfo S, Lembo D. 2009. Sulfated K5
Escherichia coli polysaccharide derivatives: A novel class of candidate antiviral
microbicides. Pharmacology & Therapeutics 123:310–322.

3. Byrnes AP, Griffin DE. 1998. Binding of Sindbis Virus to Cell Surface Heparan Sulfate. J
VIROL 72:8.

4. Smith TJ, Cheng RH, Olson NH, Peterson P, Chase E, Kuhn RJ, Baker TS. 1995. Putative
receptor binding sites on alphaviruses as visualized by cryoelectron microscopy.
Proceedings of the National Academy of Sciences 92:10648–10652.

5. Wahlberg JM, Bron R, Wilschut J, Garoff H. 1992. Membrane fusion of Semliki Forest
virus involves homotrimers of the fusion protein. J Virol 66:7309–7318.

6. Weber F, Wagner V, Rasmussen SB, Hartmann R, Paludan SR. 2006. Double-Stranded
RNA Is Produced by Positive-Strand RNA Viruses and DNA Viruses but Not in Detectable
Amounts by Negative-Strand RNA Viruses. Journal of Virology 80:5059–5064.

7. Schlee M, Hartmann G. 2016. Discriminating self from non-self in nucleic acid sensing.
Nat Rev Immunol 16:566–580.

8. Stetson DB, Medzhitov R. 2006. Type I Interferons in Host Defense. Immunity 25:373–
381.

9. García MA, Meurs EF, Esteban M. 2007. The dsRNA protein kinase PKR: Virus and cell
control. Biochimie 89:799–811.

10. Shalem O, Sanjana NE, Zhang F. 2015. High-throughput functional genomics using
CRISPR-Cas9. Nat Rev Genet 16:299–311.

11. Han J, Perez JT, Chen C, Li Y, Benitez A, Kandasamy M, Lee Y, Andrade J, tenOever B,
Manicassamy B. 2018. Genome-wide CRISPR/Cas9 Screen Identifies Host Factors
Essential for Influenza Virus Replication. Cell Reports 23:596–607.

12. Park RJ, Wang T, Koundakjian D, Hultquist JF, Lamothe-Molina P, Monel B, Schumann
K, Yu H, Krupzcak KM, Garcia-Beltran W, Piechocka-Trocha A, Krogan NJ, Marson A,
Sabatini DM, Lander ES, Hacohen N, Walker BD. 2017. A genome-wide CRISPR screen
identifies a restricted set of HIV host dependency factors. Nat Genet 49:193–203.

13. Zhang R, Miner JJ, Gorman MJ, Rausch K, Ramage H, White JP, Zuiani A, Zhang P,
Fernandez E, Zhang Q, Dowd KA, Pierson TC, Cherry S, Diamond MS. 2016. A CRISPR

screen defines a signal peptide processing pathway required by flaviviruses. Nature
535:164–168.

14. Golden RJ, Chen B, Li T, Braun J, Manjunath H, Chen X, Wu J, Schmid V, Chang T-C,
Kopp F, Ramirez-Martinez A, Tagliabracci VS, Chen ZJ, Xie Y, Mendell JT. 2017. An
Argonaute phosphorylation cycle promotes microRNA-mediated silencing. Nature
542:197–202.

15. Doench JG, Fusi N, Sullender M, Hegde M, Vaimberg EW, Donovan KF, Smith I, Tothova
Z, Wilen C, Orchard R, Virgin HW, Listgarten J, Root DE. 2016. Optimized sgRNA design
to maximize activity and minimize off-target effects of CRISPR-Cas9. Nat Biotechnol
34:184–191.

16. Li W, Xu H, Xiao T, Cong L, Love MI, Zhang F, Irizarry RA, Liu JS, Brown M, Liu XS.
2014. MAGeCK enables robust identification of essential genes from genome-scale
CRISPR/Cas9 knockout screens. Genome Biol 15:554.

17. Li J-P, Kusche-Gullberg M. 2016. Heparan Sulfate: Biosynthesis, Structure, and Function,
p. 215–273. *In* International Review of Cell and Molecular Biology. Elsevier.

18. Bishop JR, Schuksz M, Esko JD. 2007. Heparan sulphate proteoglycans fine-tune
mammalian physiology. Nature 446:1030–1037.

19. Cagno V, Tseligka ED, Jones ST, Tapparel C. 2019. Heparan Sulfate Proteoglycans and
Viral Attachment: True Receptors or Adaptation Bias? Viruses 11.

20. Lindahl U, Couchman J, Kimata K, Esko JD. 2015. Proteoglycans and Sulfated
Glycosaminoglycans, p. . *In* Varki, A, Cummings, RD, Esko, JD, Stanley, P, Hart, GW,
Aebi, M, Darvill, AG, Kinoshita, T, Packer, NH, Prestegard, JH, Schnaar, RL, Seeberger,
PH (eds.), Essentials of Glycobiology, 3rd ed. Cold Spring Harbor Laboratory Press, Cold
Spring Harbor (NY).

21. Safaiyan F, Kolset SO, Prydz K, Gottfridsson E, Lindahl U, Salmivirta M. 1999. Selective
Effects of Sodium Chlorate Treatment on the Sulfation of Heparan Sulfate. J Biol Chem
274:36267–36273.

22. Blackburn JB, D'Souza Z, Lupashin VV. 2019. Maintaining order: COG complex controls
Golgi trafficking, processing, and sorting. FEBS Lett 593:2466–2487.

23. Willett R, Ungar D, Lupashin V. 2013. The Golgi puppet master: COG complex at center
stage of membrane trafficking interactions. Histochem Cell Biol 140:271–283.

24. Kranz C, Ng BG, Sun L, Sharma V, Eklund EA, Miura Y, Ungar D, Lupashin V, Winkel
RD, Cipollo JF, Costello CE, Loh E, Hong W, Freeze HH. 2007. COG8 deficiency causes
new congenital disorder of glycosylation type IIh. Human Molecular Genetics 16:731–741.

25. Reynders E, Foulquier F, Leão Teles E, Quelhas D, Morelle W, Rabouille C, Annaert W,
Matthijs G. 2009. Golgi function and dysfunction in the first COG4-deficient CDG type II
patient. Human Molecular Genetics 18:3244–3256.

26. Spaapen LJM, Bakker JA, van der Meer SB, Sijstermans HJ, Steet RA, Wevers RA, Jaeken
J. 2005. Clinical and biochemical presentation of siblings with COG-7 deficiency, a lethal
multiple O- and N-glycosylation disorder. J Inherit Metab Dis 28:707–714.

27. Zeevaert R, Foulquier F, Jaeken J, Matthijs G. 2008. Deficiencies in subunits of the
Conserved Oligomeric Golgi (COG) complex define a novel group of Congenital
Disorders of Glycosylation. Molecular Genetics and Metabolism 93:15–21.

28. Shestakova A, Zolov S, Lupashin V. 2006. COG Complex-Mediated Recycling of Golgi
Glycosyltransferases is Essential for Normal Protein Glycosylation: COG Complex and
Golgi Glycosylation. Traffic 7:191–204.

29. Zolov SN, Lupashin VV. 2005. Cog3p depletion blocks vesicle-mediated Golgi retrograde
trafficking in HeLa cells. J Cell Biol 168:747–759.

30. Bailey Blackburn J, Pokrovskaya I, Fisher P, Ungar D, Lupashin VV. 2016. COG Complex
Complexities: Detailed Characterization of a Complete Set of HEK293T Cells Lacking
Individual COG Subunits. Front Cell Dev Biol 4.

31. Strauss JH, Strauss EG. 1994. The alphaviruses: gene expression, replication, and
evolution. Microbiol Rev 58:491–562.

32. Li Y, Muffat J, Omer Javed A, Keys HR, Lungjangwa T, Bosch I, Khan M, Virgilio MC,
Gehrke L, Sabatini DM, Jaenisch R. 2019. Genome-wide CRISPR screen for Zika virus
resistance in human neural cells. Proc Natl Acad Sci USA 201900867.

33. Murakami S, Takenaka-Uema A, Kobayashi T, Kato K, Shimojima M, Palmarini M,
Horimoto T. 2017. Heparan Sulfate Proteoglycan Is an Important Attachment Factor for
Cell Entry of Akabane and Schmallenberg Viruses. J Virol 91:e00503-17,
/jvi/91/15/e00503-17.atom.

34. Tanaka A, Tumkosit U, Nakamura S, Motooka D, Kishishita N, Priengprom T, Sa-ngasang
A, Kinoshita T, Takeda N, Maeda Y. 2017. Genome-Wide Screening Uncovers the
Significance of N-Sulfation of Heparan Sulfate as a Host Cell Factor for Chikungunya
Virus Infection. J Virol 91:e00432-17, /jvi/91/13/e00432-17.atom.

35. Savidis G, McDougall WM, Meraner P, Perreira JM, Portmann JM, Trincucci G, John SP,
Aker AM, Renzette N, Robbins DR, Guo Z, Green S, Kowalik TF, Brass AL. 2016.
Identification of Zika Virus and Dengue Virus Dependency Factors using Functional
Genomics. Cell Reports 16:232–246.

36. Klimstra WB, Ryman KD, Johnston RE. 1998. Adaptation of Sindbis Virus to BHK Cells
Selects for Use of Heparan Sulfate as an Attachment Receptor. Journal of Virology
72:7357–7366.

37. Mislick KA, Baldeschwieler JD. 1996. Evidence for the role of proteoglycans in cation-
mediated gene transfer. Proceedings of the National Academy of Sciences 93:12349–
12354.

38. Payne CK, Jones SA, Chen C, Zhuang X. 2007. Internalization and Trafficking of Cell
Surface Proteoglycans and Proteoglycan-Binding Ligands. Traffic 8:389–401.

39. Jae LT, Raaben M, Riemersma M, van Beusekom E, Blomen VA, Velds A, Kerkhoven
RM, Carette JE, Topaloglu H, Meinecke P, Wessels MW, Lefeber DJ, Whelan SP, van
Bokhoven H, Brummelkamp TR. 2013. Deciphering the Glycosylome of
Dystroglycanopathies Using Haploid Screens for Lassa Virus Entry. Science 340:479–483.

40. Blackburn JB, Kudlyk T, Pokrovskaya I, Lupashin VV. 2018. More than just sugars:
Conserved oligomeric Golgi complex deficiency causes glycosylation-independent
cellular defects. Traffic 19:463–480.

41. Ungar D, Oka T, Brittle EE, Vasile E, Lupashin VV, Chatterton JE, Heuser JE, Krieger M,
Waters MG. 2002. Characterization of a mammalian Golgi-localized protein complex,
COG, that is required for normal Golgi morphology and function. The Journal of Cell
Biology 157:405–415.

42. Willett R, Ungar D, Lupashin V. 2013. The Golgi puppet master: COG complex at center
stage of membrane trafficking interactions. Histochem Cell Biol 140:271–283.

43. D'Souza Z, Blackburn JB, Kudlyk T, Pokrovskaya ID, Lupashin VV. 2019. Defects in
COG-Mediated Golgi Trafficking Alter Endo-Lysosomal System in Human Cells. Front
Cell Dev Biol 7:118.

44. Climer LK, Pokrovskaya ID, Blackburn JB, Lupashin VV. 2018. Membrane detachment
is not essential for COG complex function. MBoC 29:964–974.

45. Blackburn JB, Lupashin VV. 2016. Creating Knockouts of Conserved Oligomeric Golgi
Complex Subunits Using CRISPR-Mediated Gene Editing Paired with a Selection Strategy
Based on Glycosylation Defects Associated with Impaired COG Complex Function, p.
145–161. *In* Brown, WJ (ed.), The Golgi Complex. Springer New York, New York, NY.

46. López P, Girardi E, Mounce BC, Weiss A, Chane-Woon-Ming B, Messmer M, Kaukinen
P, Kopp A, Bortolamiol-Becet D, Fendri A, Vignuzzi M, Brino L, Pfeffer S. 2020. High-
throughput fluorescence-based screen identifies the neuronal microRNA miR-124 as a
positive regulator of alphavirus infection. J Virol JVI.02145-19, jvi;JVI.02145-19v1.

47. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image
analysis. Nat Methods 9:676–682.
48. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-
quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol 7:539.
697

 Figures legends

Figure 1. CRISPR-Cas9 survival screen to long dsRNAs identifies the extracellular

heparan-sulfates as necessary for nucleic acids internalization and cell death induction.

A. Schematic representation of the CRISPR-Cas9 approach. HCT116 cells stably expressing a

human codon-optimized *S. pyogenes* Cas9 protein were transduced with the lentiviral sgRNA

library Brunello (MOI 0.3). 60*10^6 cells transduced cells were selected with 1 μg/mL

puromycin to obtain a mutant cell population to cover at least 300X the library. Selective

pressure via synthetic long dsRNA (1 μg/mL) was applied to induce cell death (in red). DNA

libraries from input cells and cells surviving the dsRNA treatment as three independent

biological replicates were sequenced on an Illumina HiSeq4000. Comparison of the relative

sgRNA boundance in the Input and dsRNA condition were done using the MAGeCK standard

pipeline. **B.** Median normalized read count distribution of all sgRNAs for the Input (in black)

and dsRNA (in red) replicates. **C.** Bubble plot of the candidate genes. Significance of RRA

score was calculated for each gene in the dsRNA condition compared to INPUT using the

MAGeCK software. The number of enriched sgRNAs for each gene is represented by the

bubble size. The gene ontology pathways associated to the significant top hits are indicated in

orange and green. **D.** Viability assay. Cells were transfected (80 000 cells; 1 μg/mL) or

nucleofected (200 000 cells; 400 ng) with synthetic long dsRNA and cell viability was

quantified 24 h (nucleofection) or 48 h (transfection) post treatment using PrestoBlue reagent.

Average of at least three independent biological experiments +/- SD is shown. One-way

ANOVA analysis, * $p < 0.05$. **E-F.** Cy5-labeled dsRNA (80 000 cells; 1 μg/mL) was transfected

into HCT116cas9, B4GALT7#1 and 2, SLC35B2#1 and #2 cells and Cy5 fluorescence was

quantified using FACS (10 000 events). The relative number of the Cy5 positive (Cy5+) cells

(**E**) and the relative median of Cy5 intensity of fluorescence (**F**) compared to HCT116cas9 cells

are shown. Average of three independent biological experiments +/- SD are shown. Paired t.test

analysis, * p < 0.05. **G.** Quantification of extracellular heparan-sulfates. FACS analysis of

HCT116 control or KO clones stained with the HS-specific antibody 10E4 (in red) compared

to unstained samples (in blue) (10 000 events). One representative experiment out of three is

shown.

Figure 2. *COG4* is a novel host susceptibility factor to long dsRNA induced cell death.

A. Viability assay. Cells (80 000 cells; 1 µg/mL) were transfected with dsRNA then the

viability of the cells was quantified 48 h post transfection using PrestoBlue reagent. Data from

at least three independent biological experiments are shown. One-way ANOVA analysis, * p <

0.05. **B-C.** Cy5-labeled dsRNA transfection (80 000 cells; 1 µg/mL) in HEK293T, KO COG4

and rescued cells. Cy5 fluorescence was quantified using FACSCalibur (10 000 events). The

percentage of Cy5+ cells (**B**) and the relative median of Cy5 intensity of fluorescence (**C**)

compared to parental HEK293T cells is shown. Average of three experiments +/- SD are shown.

Paired t.test analysi analysis, * p < 0.05. **D-E.** qPCR quantification of dsRNA and IFN-beta.

Cells (300 000 cells; 1 µg/mL) were transfected with synthetic long dsRNA. Total RNA was

extracted 24 h post transfection and quantified by RT-qPCR. The histogram represents the

expression fold change of synthetic dsRNA (**D**) and IFN-B mRNA (**E**) relative to GAPDH

mRNA in dsRNA transfected HEK293T KO COG4 rescued compared to HEK293T KO

COG4. Average of three independent biological experiments +/- SD are shown. Paired t.test

analysis, * p < 0.05. **F-G.** poly I:C rhodamine transfection and immunofluorescence in

HEK293T and HEK293T KO COG4. Cells were transfected with rhodamine-labeled poly I:C

(in red) and with a Rab5-GFP plasmid (in green). Images were acquired using a spinning disk

microscope at different time post transfection. Representative pictures (**E**) and the

approximative number of rhodamine-positive foci per cells quantified by counting 7 fields per

conditions (**F**) are shown. Two-way ANOVA analysis, * p < 0.05.

Figure 3. CRISPR-Cas9 screen identifies *COG4* as a permissivity factor to SINV

A. Schematic representation of the CRISPR-Cas9 approach. HCT116 cells stably expressing a

human codon-optimized *S. pyogenes* Cas9 protein were transduced with the lentiviral sgRNA

library Brunello (MOI 0.3). 60*10^6 cells transduced cells were selected with 1 μg/mL

puromycin to obtain a mutant cell population to cover at least 300X the library. Selective

pressure via SINV infection (MOI 0,1) was applied to induce cell death (in red). DNA libraries

from input cells and cells surviving the dsRNA treatment as three independent biological

replicates were sequenced on an Illumina HiSeq4000. Comparison of the relative sgRNA

boundance in the Input and dsRNA condition were done using the MAGeCK standard pipeline

B. Median normalized read count distribution of all sgRNAs for the Input (in black) and SINV

(in red) replicates**. C.** Bubble plot of the candidate genes. Significance of RRA score was

calculated for each gene in the dsRNA condition compared to INPUT using the MAGeCK

software. The number of enriched sgRNAs for each gene is represented by the bubble size. The

gene ontology pathways associated to the significant top hits are indicated in orange and green.

D. Viability of cells upon SINV infection. Cells were infected with SINV at MOI of 0.1 then

the viability of the cells was quantified 24, 48 and 72 h post infection using PrestoBlue reagent.

One-way ANOVA analysis, * $p < 0.05$. **D.** SINV-GFP plaque assay. WT, COG4KO and

rescued HEK293T cells were infected with SINV-GFP for 24 and 48h at MOI of 1 and

supernatant was collected in order to measure viral production. The fold change in titer relative

to HEK293T arbitrarily set to 1 is shown. Average of three independent biological experiments

+/- SD is shown. Paired t.test analysis * $p < 0.05$.

 Supplemental information

Supplemental figures legends

Figure S1. Set up of the HCT116 cells as model for SINV infection and dsRNA-induced

cell death.

A. Representative pictures of HCT116cas9 cells uninfected or infected with SINV-GFP MOI

of 1 at 24 and 48h post infection (hpi). 5X optical microscopy**. B.** Representative pictures of

HCT116cas9 cells at 24 and 48h post dsRNA (80 000 cells; 1 μg/mL) compared to non

transfected ones. 5X optical microscopy. **C.** Western blot of FLAG-Cas9 expression in

HCT116cas9 cells. Antibody against FLAG and GAPDH (normalizer) were used. **D-E**.

Distribution of sgRNA normalized read counts of selected genes. The read counts for the four

individual sgRNAs targeting enriched (*SLC35B2, BGAL4T7, COG4*) and not enriched genes

(*DDX58*, negative control) in the dsRNA samples (**D**) and in the SINV samples (**E**) over the

Input are shown.

Figure S2. Generation of SLC35B2 and B4GALT7 HCT116 CRISPR/Cas9 knock-out

monoclonal cell lines.

A. PCR screen of *SLC35B2* and *B4GALT7* knock-out clones obtained by CRISPR/Cas9. The

gels show the amplicons corresponding to wild type and deleted alleles that were subsequently

sequenced. **B.** Clustal Omega (48) alignment of the wild type and mutated peptide sequences

corresponding to the genomic deletions identified in the different clones. The reference

aminoacidic sequence is represented in yellow (unknown domains - InterPro) and orange

(known domains – InterPro). The peptidic sequences resulting in shorter protein due to

deletions or formation of premature stop codon in the knock-out clones are represented with

black bars or red rectangles, respectively.

Figure S3: **GFP plasmid transfectability in SLC35B2 and B4GALT7 HCT116**

CRISPR/Cas9 KO cells.

A. Representative pictures of GFP plasmid transfectability assay. Cells were transfected

(80 000 cells; 2 µg/mL) or nucleofected (200 000 cells; 500 ng) with a plasmid coding for GFP

and the GFP positive cells were observed by fluorescence microscopy at 24h (nucleofection)

or 48h (transfection) post treatment. Pictures were taken at 10× magnification, scale bar

represents 50 µm. **B.** GFP transfectability by FACS analysis. The different cell types were

transfected (80 000 cells; 2 µg/mL) with plasmid coding for GFP for 48h. The percentage of

GFP+ cells were determined by FACS analysis using a FACSCalibur. **C-D-E.** DNA plasmid

transfectability assay upon heparan-sulfates depletion. Cells were treated with 50 mM sodium

chlorate or with 2 units of heparinase then transfected with GFP plasmid. The relative number

of GFP positive (GFP+) cells (**C**), the GFP intensity of fluorescence (**D**) and the relative median

intensity of fluorescence of extracellular heparan-sulfates (HS) staining (**E**) were quantified

48h post transfection using FACS (10 000 events). Data from at least three independent

biological experiments +/- SD are shown. Paired t.test analysis, * $p < 0.05$.

Figure S4. Characterization of the survival phenotype in HCT116 KO COG4 and

HEK293 KO COG4 cells.

A. Viability assay. HCT116cas9 and HCT116cas9 KO COG4 cells (80 000 cells; 1 µg/mL)

were transfected with dsRNA then the cell viability was quantified 48 h post transfection using

PrestoBlue reagent. Data from at least three independent biological experiments are shown.

Paired t.test analysis, * $p < 0.05$. **B-C.** Cy5-labeled dsRNA transfection (80 000 cells; 1 µg/mL)

in HCT116cas9 and HCT116cas9 KO COG4 cells. Cy5 fluorescence was quantified using

FACSCalibur (10 000 events). The relative number of Cy5+ cells (**B**) and the relative median

of Cy5 intensity of fluorescence (**C**) compared to parental HCT116cas9 cells is shown. Average

of three experiments +/- SD are shown. Paired t.test analysis, * p < 0.05. **D.** Quantification of

extracellular heparan-sulfates. Upper panel correspond to FACS analysis of HEK293T control

(WT & rescued – respectively in red and orange) or HEK293T KO *COG4* (in blue) cells stained

with the HS-specific antibody 10E4 and lower panel show the unstained sample and HEK293T

KO *COG4*. One representative experiment out of three is shown. (10 000 events) **E-F.** GFP

transfectability by FACS analysis. HEK293T and HEK293T KO *COG4* cells were transfected

(80 000 cells; 2 µg/mL) with plasmid coding for GFP for 48h. The percentage of GFP+ cells

(**E**) and the relative median of GFP intensity of fluorescence (**F**) compared to parental

HEK293T cells was determined by FACS analysis using a FACSCalibur. Average of three

experiments +/- SD are shown Paired t.test analysis, * p < 0.05.

Figure S5. Accumulation of GFP and dsRNA upon SINV infection in presence or absence

of COG4. A. Representative pictures of HEK293T and HEK293T KO COG4 cells infected

with SINV-GFP MOI of 0.1 at 24 and 48h post infection (hpi). Pictures were taken at 20x

magnification. One representative experiment out of three is shown. **B.** DsRNA

immunofluorescence assay. Cells were infected with SINV at MOI of 0.1 for 24 h, then fixed

and stained with J2 antibody which recognizes dsRNA longer than 40bp and DAPI to stain the

nuclei. Pictures were taken at 40× magnification with BX51 (Olympus) microscope.

Table S1

List of primers used in the study.

Dataset S1

Count of sequenced sgRNA per genes in every replicates

Dataset S2

MAGeCK comparison report enriched sgRNAs in dsRNA versus Input samples

Dataset S3

MAGeCK comparison report enriched sgRNAs in SINV versus Input samples

Figure 1

Figure 2

Figure 3

Figure S1

Figure S2

Figure S3

Figure S4

A.

	24hpi SINV-GFP			48hpi SINV-GFP	
	brightfield	GFP		brightfield	GFP

HEK293T

HEK293T
KO COG4

B.

	DAPI	J2 (dsRNAs)	Merge

HEK293T

HEK293T
KO COG4

Figure S5