Development of thalamus mediates paternal age effect

on offspring reading: A preliminary investigation

Zhichao Xia [a, b, c], Cheng Wang [a], Roeland Hancock [d], Maaike Vandermosten [e],

Fumiko Hoeft [a, d, f, g, h] [†]

[a] Department of Psychiatry and Weill Institute for Neurosciences, University of

California, San Francisco, CA USA

[b] State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern

Institute for Brain Research, Beijing Normal University, China

[c] School of Systems Science, Beijing Normal University, China

[d] Department of Psychological Sciences and Brain Imaging Research Center,

University of Connecticut, Storrs, CT USA

[e] Experimental ORL, Department of Neuroscience, KU Leuven, Belgium

[f] Dyslexia Center, University of California, San Francisco, CA USA

[g] Haskins Laboratories, New Haven, CT USA

[h] Department of Neuropsychiatry, Keio University School of Medicine, Japan

[†] **Corresponding Author**: Fumiko Hoeft, Department of Psychological Sciences

and Brain Imaging Research Center, University of Connecticut, 850 Bolton Rd

#1271, Storrs, CT 06269; E-mails: fumiko.hoeft@uconn.edu.

Abbreviated Title: Thalamus mediates PatAGE effect on reading

Authors' Contributions: F. Hoeft designed the study and collected data with her

students. F. Hoeft, Z.C. Xia and C. Wang conceived the particular idea of the

manuscript. Z.C. Xia, C. Wang and M. Vandermosten analyzed the data. Z.C. Xia,

F. Hoeft, C. Wang, R. Hancock, and M. Vandermosten wrote the manuscript.

Acknowledgments: The authors thank all the families for their participation in

this longitudinal study. They also thank Albert Galaburda and Tuong-Vi Nguyen

for their thoughtful suggestions during manuscript preparation.

Funding: This study was supported by the Eunice Kennedy Shriver National

Institute of Child Health and Human Development (NICHD) K23HD054720 (PI: F.

Hoeft), Child Health Research Program (aka Lucile Packard Foundation for

Children's Health, Spectrum Child Health & Clinical and Translational Science

Award) (PI: F. Hoeft). F. Hoeft was supported by NIH R01HD078351 (PI: F. Hoeft),

R01HD086168 (PIs: F. Hoeft & K. Pugh), P50HD52120 (PI: R. Wagner),

R01HD044073 (PI: Cutting), R01HD096261 (PI : Hoeft), R01HD094834 (PIs: Hoeft

& Hancock), NSF BCS-2029373 (PI: F. Hoeft), UCSF Dyslexia Center, Oak

Foundation ORIO-16-012 & OCAY-19-215 (PI: F. Hoeft), and Lori and Ray DePole.

Z.C. Xia was supported by China Postdoctoral Science Foundation 2019T120062

(PI: Z.C. Xia), 2018M641235 (PI: Z.C. Xia) and China Scholarship Council (CSC)

No.201406040106.

Conflict of Interest Statement: The authors declare no competing financial

interests.

Data Availability Statement: Data that support the findings of this study are

available from the corresponding author on request.

Abstract

The importance of (inherited) genetic impact in reading development is well-

established. *De novo* mutation is another important contributor that is recently

gathering interest as a major liability of neurodevelopmental disorders, but has

been neglected in reading research to date. Paternal age at childbirth (PatAGE) is

known as the most prominent risk factor for *de novo* mutation, which has been

shown repeatedly by molecular genetic studies. As one of the first effort, we

performed a preliminary investigation of the relationship between PatAGE,

offspring's reading, and brain structure in a longitudinal neuroimaging study

following 51 children from kindergarten through third grade. The results showed

that greater PatAGE was associated significantly with worse reading, explaining an

additional 9.5% of the variations after controlling for a number of confounds —

including familial factors and cognitive-linguistic reading precursors. Moreover, this

effect was mediated by volumetric maturation of the left posterior thalamus from

ages 5 to 8. Complementary analyses indicated the PatAGE-related thalamic region

was most likely located in the pulvinar nuclei and related to the dorsal attention

network, by using offspring's diffusion MRI data, brain atlases, and public datasets.

Altogether, these findings provide novel insights into neurocognitive mechanisms

underlying the PatAGE effect on reading acquisition during its earliest phase and

suggest promising areas of future research.

Keywords

Paternal age, dorsal attention network, longitudinal design, pulvinar nuclei,

reading, dyslexia

Highlights

67   • Paternal age at childbirth (PatAGE) is negatively correlated with reading in

68   offspring.

69   • PatAGE is related to volumetric maturation of the thalamus.

70   • Brain maturation mediates the PatAGE effect on reading.

71   • PatAGE-related thalamic area is connected to the dorsal attention network.

Abbreviations

ARHQ, Adult Reading History Questionnaire; DAN, dorsal attention network;

DNA, deoxyribonucleic acid; FDR, false discovery rate; FWE, family wise error;

MatAGE, maternal age at childbirth; MNI, Montreal Neurological Institute; MRI,

magnetic resonance imaging; PA, phonological awareness; PatAGE, paternal age at

childbirth; pIQ, performance intelligence quotient; RAN, rapid automatic naming;

RD, reading disorder; READ, reading composite score; ROI, region-of-interest;

RSFC, resting-state functional connectivity; SES, socioeconomical status; $t1$, time-

point 1; $t2$, time-point 2; TIV, total intracranial volume; V5/MT, middle temporal

visual area; VAN, ventral attention network; VBM, voxel-based morphometry

Introduction

There has been a global trend of postponed childbearing, especially in

developed countries (Kohler, Billari, & Ortega, 2002). This so-called "postponement

transition" is primarily owing to changing patterns of education, employment and

marriage (Khandwala, Zhang, Lu, & Eisenberg, 2017; Sobotka, 2010). Although the

research field is still in its infancy, increasing evidence reveals that advanced

paternal age at childbirth (PatAGE) increases risk for a wide range of

neuropsychiatric conditions, such as schizophrenia and autism spectrum disorders

(D'Onofrio et al., 2014).

In contrast to the area of mental health, few studies investigated effects of

PatAGE on offspring's cognition such as reading, which is essential to success in the

modern society. The pioneering study in 1978 reported a negatively skewed

distribution of PatAGE in 48 boys with reading disorders (RD; a.k.a. and referred

here to dyslexia) (Jayasekara & Street, 1978). Four decades later, the topic remains

controversial. In a broader sample of 7-year-old children, Saha and colleagues

revealed a significantly negative effect of PatAGE on offspring's reading after

controlling for maternal age at childbirth (MatAGE), gestational age, sex and race

(Saha et al., 2009). However, when parental education and number of siblings were

added to the same dataset, the effect of PatAGE on reading was no longer

significant (Edwards & Roff, 2010). Such inconsistency underlies the importance of

more research that controls for possible confounds examining the PatAGE effect on

reading.

In addition to the controversy over the PatAGE effect on reading, no studies

have yet examined its underlying mechanisms. Nascent research in molecular

genetics however, show that PatAGE explains nearly all the variance in the amount

of *de novo* mutation, which is an alteration in a gene as the result of a mutation in a

germ cell (egg or sperm) that increases by cell divisions of the gametes

(approximately 38-fold in males at the age of 50 compared to females) (Breuss et al.,

2019; Jónsson et al., 2017; Kong et al., 2012). Hence, *de novo* mutation is the most

likely molecular mechanism underlying the PatAGE effect. In a separate line of

research, *de novo* mutations is known to increase risk by up to 20-fold in

neurodevelopmental disorders (De Rubeis et al., 2014; Deciphering Developmental

Disorders Study, 2017). Taken together, it is conceivable that *de novo* mutations are

at least partially responsible for the negative effect of PatAGE on offspring's mental

health, offering a plausible explanation of the PatAGE effect on children's reading

abilities.

At the neurocognitive level, whether and how cognitive-linguistic factors

mediate the PatAGE effect on reading development is unknown. Studies to date

have focused on genetic and environmental factors that contribute to the

multifactorial liability of dyslexia (Pennington, 2006; Petrill, Deater-Deckard,

Thompson, DeThorne, & Schatschneider, 2006). One such example, phonological

processing, is thought to exert its effect more dominantly through inherited genetic

impact, often estimated by family reading history (van Bergen, Bishop, van Zuijen,

& de Jong, 2015). Under the same framework, whether PatAGE serves as a

contributor to the multifactorial liability, and if so, what is the neural and cognitive

mediators (that would likely be heritable but not inherited traits if caused by *de*

novo mutation), have not been examined. Brain measures derived from

neuroimaging techniques including magnetic resonance imaging (MRI) are

particularly informative as they have been suggested as useful mediators between

genetic etiology and behavioral outcome, acting as endophenotypes (Grasby et al.,

2020; Shaw et al., 2012). Further, longitudinal investigation, in combination with

cross-sectional analysis, can provide comprehensive insights into the neural basis

underlying typical reading acquisition and its impairment (Clark et al., 2014;

Yeatman, Dougherty, Ben-Shachar, & Wandell, 2012).

Therefore, we conducted a preliminary study examining behavioral and

multimodal neuroimaging data both cross-sectionally and longitudinally in a cohort

of 51 children followed from kindergarten through third grade in conjunction with

analyses of publicly available datasets. The aim of the study was threefold: (1) to

examine the relationship between PatAGE and offspring's reading while

systematically controlling for potential contributing/confounding factors; (2) to

examine the role of previously known cognitive-linguistic precursors,

neuroanatomy, and its maturational process in the relation to PatAGE and reading;

and (3) to understand the functional role of the neuroanatomical findings in this

study by identifying convergent evidence through the use of brain atlases, public

datasets and our multimodal imaging data.

Materials and Methods

Participants

Participants in this study were drawn from a longitudinal NIH-funded

project (K23HD054720) focusing on children's reading development and followed

from kindergarten (time-point 1 [$t1$], mean age = 5.58 years, SD = 0.43) to third

grade (time-point 2 [$t2$], mean age = 8.30 years, SD = 0.46). All children were

healthy native English speakers without neurological/psychiatric disorders (e.g.,

attention deficit/hyperactivity disorder) or contraindications to MRI based on

parental report. Among the participating children, 76% were White, 6% were Asian,

and 18% were of multiradical heritage. In addition, 8% identified as Hispanic or

Latino. Based on the household annual income, parental educational levels and

occupation, the participants in this study was of relatively high socioeconomic

status (also see Black et al., 2012). The initial sample consisted of 51 children

recruited from local newspapers, school mailings, flyers, and mothers' clubs. In the

behavior analyses, eight children were excluded because of attrition (n = 5), no

record of PatAGE (n = 1), or being siblings (n = 2). In the latter case, the child with

poorer T1 image quality was excluded. The final sample for behavioral analyses

included 43 unrelated children (17 girls). In the neuroanatomical analysis, another

7 children (2 girls) were excluded because of incomplete T1 data collection or poor

image quality at either time-point. In the diffusion imaging analysis, a sub-group of

23 children (8 girls) with the same acquisition sequence were included. The

differences in either familial or behavioral measures between the entire cohort and

sub-groups were non-significant (all p's > 0.1). The Institutional Review Boards of

Stanford University where data were collected and principal investigator was at the

time of the study, and the University of California San Francisco where data were

analyzed due to transition of the principal investigator, approved the present study.

Both informed assent and consent were obtained from children and their guardians.

Behavioral measurements

Demographics, family information and behavioral measures are summarized

in Table 1. Family information collected at t1 include: PatAGE; MatAGE; Adult

Reading History Questionnaire from both parents (PatARHQ, MatARHQ) that were

used to estimate familial history of reading difficulty (Lefly & Pennington, 2000);

numbers of older and younger siblings; parental education (PatEDU, MatEDU);

socioeconomic status (SES), a composite index computed from family annual

income, parental education and occupation with principal component analysis

(Noble, Wolmetz, Ochs, Farah, & McCandliss, 2006); Home Observation

Measurement of the Environment (HOME), an index for home environment

including home literacy environment (Segers, Damhuis, van de Sande, &

Verhoeven, 2016). A battery of behavioral tests measuring intelligence, language

and reading related skills was administered. Verbal Comprehension, Concept

Formation, and Visual Matching sub-tests of the Woodcock-Johnson III Tests of

Cognitive Abilities (McGrew & Schrank, 2007) were used to estimate general

cognitive abilities. These tests have reliabilities of at least 0.80 and have been used

as a proxy for intelligence quotient (IQ) (Shaw, 2010). Vocabulary was measured

with Peabody Picture Vocabulary Test (4th edition) (Dunn & Dunn, 2007). Blending,

Elision, Memory for Digit, Nonword Repetition sub-tests from the Comprehensive

Test of Phonological Processing (CTOPP 1st Edition) (Wagner, Torgesen, &

Rashotte, 1999) were used to measure phonological skills. Rapid Automatized

Naming (RAN; Objects and Colors sub-tests) (Wolf & Denkla, 2005) and Letter

Identification sub-test of Woodcock Reading Mastery Test R/NU (WRMT-R/NU)

(Mather, 1998) were also administered.

The same set of tests were used at *t*2 (Table 1). Numbers and Letters sub-

tests of RAN were further included to measure print-sound mapping efficiency.

Additionally, we administered tests measuring different aspects of reading,

including Sight Word Efficiency and Phonemic Decoding Efficiency sub-tests from

the Test of Word Reading Efficiency (TOWRE 1st Edition) (Torgesen, Wagner, &

Rashotte, 1999), Word Identification, Word Attack, and Passage Comprehension

sub-tests from WRMT-R/NU, and Reading Fluency and Spelling sub-tests from WJ-

III Tests of Achievement.

Image acquisition

High-resolution T1-weighted images were collected at both time-points with

the following parameters: 128 slices; thickness = 1.2 mm; NEX = 1; repetition time =

8.5 ms; echo time = 3.4 ms; inversion time = 400 ms; in-plane resolution = 256 ×

256; voxel size = 0.9 × 0.9 × 1.2 mm^3; flip angle = 15 °; field of view = 22 cm. High-

angular resolution diffusion-imaging data were collected at $t2$ with the following

parameters: 46 axial slices; slice thickness = 3 mm; repetition time = 5000 ms; echo

time = 81.7 ms; in-plane resolution = 128 × 128; voxel size = 2.0 × 2.0 × 3.0 mm^3;

150 directions with b = 2500 s/mm^2; 6 volumes with b = 0 s/mm^2. All images were

acquired using a GE Healthcare 3.0 T 750 scanner with eight-channel phased-array

head coil at Richard M. Lucas Center for Imaging at Stanford University. The

quality of images was qualitatively evaluated by an investigator who was blinded to

the behavioral and demographic information prior to any analyses.

Behavior analyses

To reduce dimensionality of behavioral metrics, factor analyses were

conducted on reading-related tests for each time-point; $t1$: Blending, Elision,

Memory for Digits, Nonword Repetition sub-tests of CTOPP, Objects and Colors

sub-tests of RAN, Letter Identification sub-test of WRMT; $t2$: Blending, Elision,

Memory for Digits, Nonword Repetition sub-tests of CTOPP, Numbers, Letters,

Objects and Colors sub-tests of RAN, Sight Word Efficiency and Phonemic Decoding

Efficiency sub-tests of TOWRE, Word Identification, Word Attack, Passage

Comprehension sub-tests of WRMT-R/NU, Reading Fluency and Spelling sub-tests

of WJ-III Tests of Achievement. Maximum Likelihood, Varimax, and Bartlett

methods were used for extraction, rotation, and factor score calculation,

respectively. Criteria of eigenvalue greater than 1 was used to identify factors.

From $t1$ behavioral metrics, we obtained two factors, explaining 53.8% of the total

variance. Since phonological awareness (PA) and RAN loaded heavily on each

factor, we named these factors as $t1$PA and $t1$RAN (Fig. 1A). Given PA and RAN are

the most reliable predictors for reading development in alphabetic languages

(Caravolas et al., 2012), we used these two composite scores as cognitive-linguistic

precursors of reading in subsequent analyses. Using the same approach, we

extracted three factors from $t2$ behavioral metrics, explaining 67.2% of the total

variance. The scores were labeled as $t2$READ, $t2$PA, and $t2$RAN according to the

corresponding factor loading (Fig. 1B).

Since a consensus on the definition of advanced paternal age remained

lacking (Couture, Delisle, Mercier, & Pennings, 2020), in this study we treated

PatAGE as a continuous variable rather than splitting children into different

groups. To examine the relationship between PatAGE and reading, we first

calculated zero-order correlation. Once the correlation was significant, hierarchical

linear regressions were conducted to answer four questions in the following order:

(1) whether the PatAGE effect on reading remains significant after controlling for

demographic variables; (2) whether the PatAGE effect on reading exists after

regressing out MatAGE, which is known to correlate highly with PatAGE and is a

possible confound; (3) whether the PatAGE effect on reading is present above and

beyond familial risk (representing inherited risk; Swagerman et al., 2017), and

environmental factors to address the issue of multifactorial liability; (4) whether the

PatAGE effect on reading is explained by $t1$ cognitive-linguistic precursors of

reading to examine whether the most common predictors were the mediating factor.

254   Specifically, in the first model we entered $t2$ age, sex, handedness and

average performance IQ (pIQ) across $t1$ and $t2$ in the first step and PatAGE in the

second step (Table 2). In the second model, besides the aforementioned nuisance

variables, we regressed out MatAGE, which was correlated with both PatAGE and

$t2$READ. In the third model, familial risk measured by ARHQ of both parents (van

Bergen et al., 2015), and environmental factors including educational level of both

parents (Edwards & Roff, 2010), number of older and younger siblings (Price, 2008),

SES (Pan et al., 2016), HOME (Segers et al., 2016), which are known to be

associated with reading were additionally controlled. In the final model, $t1$PA and

$t1$RAN were entered in the fourth step, just before PatAGE, to examine whether the

PatAGE effect was present beyond $t1$ cognitive-linguistic skills. Since $t1$RAN and

$t1$PA did not correlate with PatAGE, these factors were not examined further in the

mediation analyses. All statistics were done with SPSS 24.0 (IBM, Inc.), and p-

values were two-tailed while statistical significance was set at 0.05.

Structural image preprocessing

269   Both cross-sectional and longitudinal analyses were conducted by using

voxel-based morphometry toolbox (v435; http://www.neuro.uni-jena.de/vbm/) with

SPM8 (v4667; http://www.fil.ion.ucl.ac.uk/spm8/) implemented in Matlab R2014a.

In the cross-sectional data preprocessing for $t1$ and $t2$, individual T1 volumes were

segmented into gray matter, white matter and cerebrospinal fluid with a

resampling at 1.5 mm³. Then, the gray matter segments were registered to a T1

template in Montreal Neurological Institute (MNI) space by using both affine

normalization and Diffeomorphic Anatomical Registration Through Expoentiated

Lie Algebra (Ashburner, 2007), and subsequently modulated by the 'affine and non-

linear' modulation (http://www.neuro.uni-jena.de/vbm/segmentation/modulation/).

The modulated images containing regional tissue volume of gray matter for each

voxel were smoothed with an 8-mm full-with half-maximum isotropic Gaussian

kernel. Voxels with gray matter values < 0.1 were excluded to avoid edge effects.

In the longitudinal analysis, 'Preprocessing of Longitudinal Data' module

that contains specific preprocessing steps for processing longitudinal structural MRI

data was used. Intra-subject realignment, bias correction, segmentation, and

normalization were conducted sequentially as described elsewhere (Ridgway et al.,

2007). After applying spatial smoothing with an 8-mm full-width half-maximum

Gaussian kernel, we obtained maps of gray matter volume for both time-points. We

generated GMV maps reflecting change from $t1$ to $t2$ for further analyses (such that

a positive value indicates enlarging from $t1$ to $t2$).

Whole-brain analyses

Prior to voxel-wise analyses, we examined relationships between PatAGE

and global measurement (i.e., the total intracranial volume that defined as the sum

of total gray matter, white matter and cerebrospinal fluid) at each time-point ($t1$TIV

and $t2$TIV). Then, we examined whether PatAGE correlated with the change of TIV

(ΔTIV) from $t1$ to $t2$ while controlling for the baseline measure ($t1$TIV). To explore

relationships between regional gray matter volume (GMV) at each time-point (i.e.,

cross-sectional analyses), as well as the change of regional GMV (ΔGMV) across

time-points ($t2$GMV-$t1$GMV) with PatAGE (i.e., longitudinal analysis), voxel-wise

whole brain regressions were conducted while controlling for global measurements.

Specifically, $t1$TIV or $t2$TIV was controlled in cross-sectional analyses for $t1$ and $t2$,

respectively. In the longitudinal analysis, $t1$TIV and ΔTIV were controlled to

exclude the effects from initial gross volume and its development. Since correlations

between $t1$TIV, ΔTIV, and PatAGE were not significant (all p's > 0.1), the models

were free from multicollinearity. Topological Family Wise Error (FWE) correction

was used to determine the corrected thresholds of statistical significance. All

clusters significant at a threshold of corrected $p < 0.05$ corrected for the whole brain

(p-voxel < 0.005 for height) were reported in MNI space. Since no significant

clusters were found for cross-sectional analyses at either time-point, the subsequent

analyses focused on longitudinal data. In particular, region-of-interest (ROI)

analyses were conducted in the significant clusters to examine the robustness and

specificity of the effect. For this, values of each voxel in the cluster were extracted

and averaged, then included in hierarchical multiple regression analyses as the

dependent variable. Demographic variables ($t1$ age, time interval between $t1$ and $t2$,

sex, handedness, average pIQ across $t1$ and $t2$), $t1$TIV and ΔTIV were entered in the

first step, while PatAGE was entered in the second step. Then, we further

controlled for MatAGE and MatARHQ since they showed significant correlation

with PatAGE as in previous analyses of this study.

Next, we examined the relationship between ΔGMV and children's t2READ

in the cluster that was significantly associated with PatAGE (hereafter PatAGE-

cluster) by using small volume correction (p-voxel < 0.005, p-cluster < 0.05,

topological FWE correction) while t1TIV and ΔTIV were statistically controlled.

Average ΔGMV of the significant cluster was extracted for subsequent ROI

analyses. Hierarchical multiple regression analyses were conducted to test for the

robustness and specificity of the effect. In the first model, t2READ was the

dependent variable and demographic variables (t2 age, sex, handedness, average

pIQ across t1 and t2), t1TIV and ΔTIV were entered in the first step. In the second

model, t1PA and t1RAN were additionally entered in the second step since they

were significantly associated with t2READ in previous analysis of this study. The

average ΔGMV was entered in the final step.

Mediation

In the region where the volumetric change significantly correlated with

PatAGE and t2READ, we examined whether the negative PatAGE effect on reading

was mediated by brain maturation. We first examined a model without controlling

for covariates. Bootstrapping (10,000 samples) was used to obtain 95% confidence

intervals of the indirect effect. If a significant indirect effect was observed, we

further adjusted the model for demographic variables (t2 age, sex, handedness,

average pIQ across $t1$ and $t2$), $t1$TIV, ΔTIV, and $t1$ cognitive-linguistic precursors

($t1$RAN and $t1$PA; i.e., significant predictors on $t2$READ in addition to PatAGE as

shown in the regression analysis) to confirm the uniqueness and robustness of the

effect.

Existing evidence suggests that dyslexia is largely genetically transmitted

from parent (often assayed by parental-report of reading difficulty) to offspring

(Soden et al., 2015; Swagerman et al., 2017). Further, Twin studies find a

dissociation between sources of variance in phonological and orthographic processes,

with variance in phonological skills being primarily genetic compared to

orthographic skills (Olson, Wise, Conners, Rack, & Fulker, 1989; Olson et al., 2011).

These findings are consistent with the idea that PA mediates the negative effect of

parental reading difficulty (a proxy for inherited genetic transmission) on reading in

offspring (van Bergen et al., 2015). In line with the previous literature, we observed

significant correlations between MatARHQ and $t1$PA, MatARHQ and $t2$READ,

$t1$PA and $t2$READ. We therefore examined the role of poor PA on the relationship

between history of maternal reading difficulty and lower reading performance in

offspring. Age at $t2$, sex, handedness, average pIQ across $t1$ and $t2$, and $t1$RAN were

controlled statistically.

If both a PatAGE effect (a proxy for non-inherited genetic risk) and

MatARHQ effect via PA processing (a proxy for inherited genetic risk) are to be

observed in the current sample, the result will provide support of the multifactorial

liability model.

PROCESS procedure (release 2.16.1) implemented in SPSS was used to

conduct these mediation analyses (Hayes, 2013).

Complementary analyses

We adopted multiple complementary analytical approaches to depict fine-

grained spatial localization and connectivity patterns of the PatAGE-cluster,

capitalizing on the fact that these have been shown to inform possible functional

roles of a particular brain region (in this case, the left posterior thalamus) in the

absence of a comprehensive set of cognitive measures. First, we spatially localized

the PatAGE-cluster with two brain atlases. (1) Since the thalamus consists of

multiple nuclei that have different functions, we calculated the percentage of

overlapped voxels between the PatAGE-cluster and each thalamic nucleus from the

Morel Atlas, a histological atlas that is optimal for thalamic targets in MNI space

(Jakab, Blanc, Berényi, & Székely, 2012; Krauth et al., 2010); (2) Since the

structural connectivity provides information about the function of a given brain

region (Barron, Eickhoff, Clos, & Fox, 2015), we used Oxford Thalamic Connectivity

Probability Atlas with the atlasquery tool implemented in FSL

(https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/Atlases) to obtain the probability that the

PatAGE-cluster is structurally connected to different cortical areas (de Moura et al.,

2016).

To further identify the functional role of the PatAGE-cluster and

complementary to the results from analyses using the histological and connectivity

probability atlases, we examined PatAGE-cluster-associated cortical patterns by

using an online database, Neurosynth v0.5 (Yarkoni, Poldrack, Nichols, Van Essen,

& Wager, 2011). In particular, we generated a co-activation map by including all

fMRI studies in the database (N > 10,900), with the PatAGE-cluster as ROI. False

Discovery Rate (FDR) corrected $q < 0.01$ was used as the threshold to obtain

significant regions reported in fMRI studies when the PatAGE-cluster is also

reported (i.e., forward inference). In addition, we generated a map of whole-brain

resting-state functional connectivity (RSFC) by using the 1,000 Functional

Connectome dataset (Biswal et al., 2010). The center of gravity of the PatAGE-

cluster (MNI: x = -19, y = -28, z = 6) was used as the seed, and its connectivity to the

rest of the brain was calculated. The resultant brain map was thresholded with a

liberal cutoff value of $r = 0.01$, same as that in the previous literature (Yang,

Rosenblau, Keifer, & Pelphrey, 2015). To be conservative, we only considered the

overlapping regions between the co-activation map and the RSFC map. Dice

coefficients between the conjunction map and the 7 large-scale intrinsic connectivity

networks (i.e., visual, somatomotor, dorsal attention, ventral attention, limbic,

frontoparietal, and default networks) from Yeo et al. (2011) were then calculated to

examine which functional network most overlapped with the PatAGE-cluster-

associated cortical pattern.

In the final step, we analyzed white matter connectivity, where fibers passing

the PatAGE-cluster were reconstructed using deterministic tractography. Diffusion-

weighted imaging preprocessing was performed by using ExploreDTI (Leemans,

Jeurissen, Sijbers, & Jones, 2009). Next, to visually inspect for possible artefacts,

rigorous motion correction with CATNAP and eddy current correction were

conducted by using the required reorientation of the b-matrix (Leemans & Jones,

2009). The diffusion tensors were calculated using a non-linear regression procedure

(Pierpaoli & Basser, 1996). The individual datasets were non-rigidly normalized to

MNI space. Next, whole brain tractography was performed for each individual

dataset using a deterministic approach (Basser, Pajevic, Pierpaoli, Duda, &

Aldroubi, 2000). Fibers (streamlines) were reconstructed by defining seed points

distributed uniformly throughout the data at $2.0 \times 2.0 \times 2.0$ mm^3 resolution,

following the main direction with step size set at 1.0 mm. Fiber tracking was

discontinued when the fiber entered a voxel with fractional anisotropy < 0.2 or

made a high angular turn (angle > 40°), or when the fiber was outside the fiber

length range of 50-500 mm. Two analyses were then conducted: (1) To localize fibers

and get a general view, the cluster related to PatAGE was used as ROI and all

fibers passing through this cluster were delineated. The delineated fibers and its

projection points were visually inspected, after which individual maps were

binarized and summed to acquire probabilistic map across participants. (2) To

complement the Neurosynth analysis above and to identify the functional network

most relevant to the PatAGE-cluster, the numbers of streamlines passing through

the PatAGE-cluster and each of the 7 functional networks (Yeo et al., 2011) were

calculated and normalized by dividing this number by size of the target network.

The results were treated as the connectivity strength and compared between

networks with one-way ANOVA (as well as post-hoc analyses). Furthermore, we

examined the correlations between the connectivity strength with PatAGE and

t2READ.

Results

PatAGE is negatively associated with offspring's reading above and

beyond commonly known predictors

PatAGE (M = 36.12 years, SD = 4.91, $Range$ = 25-47; Table 1; distribution

presented in Fig. S1A) was positively correlated with MatAGE (r = 0.63, p = 5 × 10-

6) but not with other potentially confounding demographic variables reported in the

past such as SES, number of siblings and parental education (all p's > 0.1; Table

S1). Importantly, greater PatAGE was significantly correlated with lower reading

composite scores in offspring (t2READ; r = -0.39, p = 0.011). Similar to PatAGE and

not surprisingly, MatAGE was negatively correlated with t2READ (r = -0.33, p =

0.031). No significant correlations were found between PatAGE and cognitive-

linguistic skills typically found to be predictors of later reading ability at either

time-point (p's > 0.1, for t1PA, t1RAN, t2PA, and t2RAN). In accordance with prior

literature on factors that predict reading outcomes (Segers et al., 2016; Thompson et

al., 2015; van Bergen et al., 2015), lower t2READ was predicted by poorer reading

reported by mothers (MatARHQ; r = -0.46, p = 0.002), poorer home literacy

environment measured by HOME (r = 0.31, p = 0.047), and poorer cognitive-

linguistic skills at time-point 1 (t1PA: $r = 0.46$, $p = 0.002$; t1RAN: $r = 0.31$; $p =$

0.041).

To examine whether the PatAGE effect on reading existed above and beyond

commonly identified confounds and additional variables known to influence reading

development, hierarchical linear regressions were conducted with t2READ as the

dependent variable in a systematic and hypothesis-driven fashion. In the first

model, before PatAGE was entered, confounds related to children were entered as

predictors in the first step, including t2 age, sex, handedness and average pIQ

across two time-points. The effect of PatAGE remained significant, explaining an

additional 14.9% of the variance ($t = -3.12$, $p = 0.004$; Model 1 in Table 2; Fig. S1B).

In the second model, MatAGE was further added in the second step, since it was

significantly correlated with PatAGE and t2READ. As shown in Table 2 (Model 2),

PatAGE explained an additional 9.7% of the variance in reading outcomes ($t = -2.48$,

$p = 0.018$). Then, in the third model, familial risk (PatARHQ, MatARHQ) and

environmental factors (number of siblings, parental education, SES and HOME) in

relation to reading development were added. We still observed a significant PatAGE

effect, explaining an additional 10.7% of the variance ($t = -2.45$, $p = 0.023$; Model 3

in Table 2). Thus far, we demonstrated that the PatAGE effect on reading was not

accounted for by factors that predict child's reading outcome and known to be either

inherited or environmental. In the final model, we investigated its relationship with

early cognitive-linguistic predictors of reading outcomes by entering t1PA and

t1RAN in the fourth step. The PatAGE effect on offspring's reading was above and

beyond that of cognitive-linguistic variables, explaining an additional 9.5% of the

variance (t = -2.71, p = 0.014; Model 4 in Table 2). In accord with the prior

literature, t1PA and t1RAN also significantly predicted t2READ in the final model

and accounted for 13.8% of the variation (t1PA: t = 2.87, p = 0.010; t1RAN: t = 2.19,

p = 0.042). That is, contributions from PatAGE and cognitive-linguistic precursors

were relatively independent and jointly predicted children's reading outcomes.

PatAGE is associated with thalamic maturation

In the brain analyses, we found no significant correlations between PatAGE

and TIV at t1, t2, or ΔTIV from t1 to t2 (all p's > 0.1). Second, whole brain analyses

on regional GMV at each time-point did not show any significant clusters at the

FWE corrected threshold of p < 0.05. Finally, we examined the PatAGE effect on

regional ΔGMV while controlling for t1TIV and ΔTIV. Results revealed a

significantly positive correlation between PatAGE and ΔGMV in a cluster covering

the left posterior thalamus (i.e., PatAGE-cluster; p = 0.017, FWE corrected, 819

voxels, 2,764 mm^3, peak MNI coordinate [-27, -30, 6]; Fig. 2A). Specifically, greater

PatAGE was associated with less GMV decrease (Fig. 2B). To verify that this effect

was not due to confounds, hierarchical linear regression analyses were performed.

In the first model, after regressing out nuisance variables commonly controlled in

longitudinal VBM analysis (t1 age, time interval between t1 and t2, sex,

handedness, average pIQ across t1 and t2, t1TIV and ΔTIV), PatAGE explained

34.6% of the ΔGMV variance of the PatAGE-cluster (t = 4.59, p < 0.001). Since

MatARHQ and MatAGE were significantly correlated with PatAGE, we additionally

included them as covariates in the second model, and found PatAGE still explained

19.0% of the ΔGMV variance of the PatAGE-cluster ($t = 3.28$, $p = 0.003$).

The PatAGE effect on offspring's reading is mediated by ΔGMV in the left

posterior thalamus

To examine whether ΔGMV in the PatAGE-cluster was correlated with

reading, we performed voxel-wise regression analysis within the cluster while

controlling for $t1$TIV and ΔTIV. As shown in Fig. 2A, there was a negative

correlation between ΔGMV and $t2$READ in a sub-region of the PatAGE-cluster (p-

cluster = 0.037, small volume FWE corrected, 86 voxels, 290 mm³, peak MNI

coordinate [-27, -31, 10]), indicating the less the thalamic GMV decrease between

the two time-points, the poorer the reading (Fig. 2C). Hierarchical linear regression

analyses were performed to verify this effect. ΔGMV of the significant cluster

explained 19.7% of variance in $t2$READ after regressing out nuisance variables ($t2$

age, sex, handedness, average pIQ across $t1$ and $t2$, $t1$TIV and ΔTIV) in the first

step ($t = -3.20$, $p = 0.003$). Since $t2$READ was significantly correlated with $t1$PA and

$t1$RAN, we further examined whether the observed effect was above and beyond

these two cognitive-linguistic skills by entering them in the second step. The results

showed that ΔGMV additionally explained 19.9% of the variance in $t2$READ ($t = -$

3.66, $p = 0.001$).

Given that ΔGMV in the left thalamic sub-region was associated with both

PatAGE and $t2$READ, we conducted mediation analyses and observed ΔGMV

significantly mediated the PatAGE effect on offspring's reading; 95% confidence

interval was [-0.522 -0.041] when age at $t2$, sex, handedness, average pIQ across $t1$

and $t2$, TIV at $t1$, ΔTIV, $t1$PA, and $t1$RAN were not controlled for, and [-0.552 -

0.043] when these covariates were controlled (Fig. 2D). These results are in contrast

to the commonly found results in the literature that we also find in the present

study, i.e., $t1$PA mediates the negative effect of family history on offspring's reading

(95% confidential interval was [-0.303 -0.022] when not controlling for age at $t2$, sex,

handedness, average pIQ across $t1$ and $t2$, and $t1$RAN, and changed to [-0.249 -

0.001] when all these covariates were controlled; Fig. S2).

PatAGE-cluster is localized in the pulvinar nuclei and linked to the dorsal

attention network

520   To understand the neurostructural profile of the PatAGE-cluster in the left

thalamus, we compared it with a histological atlas and a connectivity atlas. 548 out

of 819 voxels in the cluster overlapped with the human thalamus of the Morel

histological atlas (Jakab et al., 2012; Krauth et al., 2010), while the remaining 271

voxels were unlabeled, possibly because the cluster also contained white matter. As

presented in Fig. 3A, within the overlapping region, 380 voxels (69.5%) were in the

subdivision labeled as pulvinar nuclei, especially the medial portion, which is

known to have widespread connections with the inferior parietal lobule (Arcaro,

Pinsk, & Kastner, 2015). These results were corroborated by examining the

Thalamic Connectivity Probability Atlas

(http://fsl.fmrib.ox.ac.uk/fsl/fslview/atlas.html): the PatAGE-cluster was most likely

localized in the subdivision that was connected to the posterior parietal cortex, with

a probability of 38.9% (Fig. 3B).

Next, we examined functional connectivity of the PatAGE-cluster by utilizing

two approaches available in Neurosynth v0.5 (Yarkoni et al., 2011). These included

generation of a meta-analytic map of regions that co-activate with the PatAGE-

cluster across more than 10,900 fMRI studies, as well as a RSFC map from the

PatAGE-cluster to the rest of the brain by using the 1000 Functional Connectome

dataset (Biswal et al., 2010). The co-activated areas included subcortical structures

and cortical regions such as bilateral intraparietal sulci, inferior temporal gyrus,

and frontal eye fields in the frontal cortex (Fig. S3A). The RSFC map showed

similar but more widespread pattern than the co-activation map (Fig. S3B). A

conjunction analysis revealed that bilateral frontal eye fields, intraparietal sulci,

middle temporal visual area (V5/MT), and cerebellum were among the overlapped

regions across the two approaches, in addition to subcortical structures.

Sørensen-Dice coefficients (s) between the overlapping areas and the

previously identified networks deriving from resting-state functional MRI data (Yeo

et al., 2011) were calculated. The derived pattern of overlapping areas showed the

greatest resemblance to the dorsal attention network (DAN; $s = 0.360$; Fig. 4B) and

to the ventral attention network (VAN; $s = 0.261$), much higher than its

resemblance to other networks (visual network: $s = 0.086$; somatomotor network: $s =$

0.071; limbic network: $s < 0.001$; frontoparietal network: $s = 0.041$; and default

network: $s = 0.005$). Together with the aforementioned findings utilizing structural

atlases, these results using large-scale fMRI databases from functional

neuroimaging studies point to the attention network, in particular the DAN, to be

the candidate brain functional system associated with the PatAGE-cluster in the

left thalamus.

Finally, we used diffusion imaging data available in a sub-group of 23

participants to confirm DAN was more likely the candidate system associated with

the PatAGE effect on reading. Using deterministic tractography, we reconstructed

white matter fibers through the PatAGE-cluster, covering inferior fronto-occipital

fasciculus, corticospinal tract, forceps major, superior corona radiata, as well as

anterior and posterior limbs of the internal capsule. Fig. 5A shows reconstructed

fibers in a representative child and Fig. 5B shows intersection across participants,

for demonstrative purposes. More importantly, the PatAGE-cluster showed

significantly stronger connectivity (defined as the total number of streamlines

divided by the size of the target network) with DAN than with VAN ($t = 6.61$, $p <$

0.001; Fig. 5C). Finally, consistent with the aforementioned results, significant

positive correlation between PatAGE-cluster-DAN streamlines and PatAGE ($r =$

0.49, $p = 0.018$; Fig. 5D), and significant negative correlation between PatAGE-

cluster-DAN streamlines and t2READ ($r = -0.45$, $p = 0.030$; Fig. 5E) were observed.

No significant correlations were found between PatAGE-cluster-VAN streamlines

and PatAGE or t2READ (both $p's > 0.1$).

Discussion

In this study, we observed a significant effect of PatAGE on offspring's

reading at the earliest stages of formal schooling from ages 5 to 8, independent of

confounds (e.g., maternal age) and factors that play key roles in learning to read

(i.e., family reading history, environmental factors, and cognitive-linguistic

precursors of reading), explaining an additional 9.5% of the variance. Furthermore,

we revealed volumetric maturation of the left thalamus as a potential neural

endophenotype mediating this effect. With multimodal neuroimaging and public

datasets, we identified that this area is most relevant to the dorsal attention

network. These findings are in contrast to and complement the current literature

linking phonological and orthographic processing in reading to the left temporo-

parietal and occipito-temporal regions. The mediation revealed here was distinct

from the mediating role of phonological processing on the relationship between

reading and familial risk, which has been attributed to hereditary effects. Taken

together, this study provides novel and converging evidence suggesting PatAGE as

a significant factor that associated with offspring's reading, independent of

phonological processing, possibly through the maturational process of the left

posterior thalamus.

The PatAGE effect on offspring's reading

Jayasekara and Street (1978) for the first time reported that advanced

PatAGE was associated with greater incidence of dyslexia, independent of SES and

birth order. While their analysis was restricted to dyslexic boys, Saha and

colleagues extended the finding to a broader population of 7-year-old children with

varying reading abilities measured using Wide Range Achievement Test (Saha et

al., 2009). Negative PatAGE effects on several cognitive measures including reading

were observed after controlling for confounds that included MatAGE, SES, and

parental psychiatric illness. A follow-up study re-analyzed the same dataset and

found that the PatAGE effect was no longer significant after further adjusting

parental education and the number of siblings (Edwards & Roff, 2010). Therefore,

the PatAGE effect on reading was equivocal, and the inconsistency was related to

covariates controlled in the model, especially parental characteristics such as

educational level.

In the present study, with the range of PatAGE restricted to 25-47 years, we

found PatAGE was negatively associated with reading performance measured using

variety of tests, even after additionally controlling for strong predictors of reading

that were not included in previous studies (Edwards & Roff, 2010; Saha et al.,

2009). These predictors included familial risk and cognitive-linguistic skills (e.g.,

phonological processing) that shown to be more genetically than environmentally

mediated, as well as home literacy environment (Hulme, Snowling, Caravolas, &

Carroll, 2005; Swagerman et al., 2017; van Bergen et al., 2015). These findings

support an adverse PatAGE effect on reading and suggest such effect may occur

through a mechanism different to factors such as inherited genetic and

environmental risks.

While the number of studies examining the PatAGE effect on reading is too

few to infer potential mechanisms, studies on other PatAGE-linked

neurodevelopmental disorders offer insights. One predominant explanation is that

advanced PatAGE exerts its effect on the risk of disorders such as autism and

schizophrenia through accumulated *de novo* genetic mutations and epigenetic

modifications (e.g., DNA methylation and repressive histone modification) in

paternal gametes (Deciphering Developmental Disorders Study, 2017; Girard et al.,

2016; Saha et al., 2009).

At a more macroscopic scale, understanding of the mechanisms can be

deepened by identifying intermediate (endo)phenotypes through behavioral and

neuroimaging measures such as we did in the current study. That is, advanced

PatAGE may impact precursors of neurodevelopmental disorders, which in turn

leads to higher occurrence of such disorders (Cannon, 2009). For example, the

likelihood of having impaired social functioning in offspring, a core symptom of

psychiatric disorders, increases with PatAGE (Weiser et al., 2008). While the

underlying mechanisms are yet to be fully understood, multifactorial liability

confers risk for neurodevelopmental disorders and may involve liability such as *de*

novo mutations in addition to inherited and environmental risks. Adding to prior

research, the current study offers insights into potential mechanisms at the

macroscopic level.

The intermediary role of the left posterior thalamus

The thalamus is an important relay center in the human brain, receiving

information from sensory cortices and relaying it to higher-level association cortex.

Previous studies paint a mixed picture on thalamic development: gross volume

relative to its brain size is smaller (Sussman, Leung, Chakravarty, Lerch, & Taylor,

2016) or larger (Brain Development Cooperative, 2012) in older compared to

younger children of ages 4 to 18, and the pulvinar compared to other thalamic

nuclei show no apparent change with age (Raznahan et al., 2014). Despite

controversial evidence on typical thalamic maturation, its anomalous development

undoubtedly affects the maturation of other cortical and subcortical brain regions

(Ball et al., 2012), which in turn could impact higher level cognitive processes that

underlie typical reading acquisition. In support of this, anomalies in thalamic

structure (Giraldo-Chica, Hegarty, & Schneider, 2015), activation (Diaz, Hintz,

Kiebel, & von Kriegstein, 2012; Koyama, Molfese, Milham, Mencl, & Pugh, 2020),

and connectivity (Müller-Axt, Anwander, & von Kriegstein, 2017; Paz-Alonso et al.,

2018; Tschentscher, Ruisinger, Blank, Diaz, & von Kriegstein, 2018) are associated

with dyslexia. While most of these studies adopt a cross-sectional design with adult

participants, here we conducted a longitudinal investigation and found volumetric

change in the posterior thalamus from ages 5 to 8 was significantly associated with

PatAGE; children with younger fathers showed GMV decrease, whereas those with

older fathers showed less decrease or an increase. This pattern suggests that

PatAGE is associated with the development of this subcortical structure.

Noteworthy is that although no PatAGE effect were observed when examining a

single time-point (cross-sectional), it does not indicate this effect cannot manifest at

a specific age (i.e., this may be due to a particular developmental stage examined).

Rather, it underscores the importance of considering developmental dynamics when

examining brain-behavior relationships. A similar pattern has been revealed in

white matter development in dyslexia (Yeatman et al., 2012).

Moreover, examination of the cluster location with the Morel Atlas suggested

the foci in the left pulvinar, an integral region supporting visuo-spatial attention

(Amso & Scerif, 2015; Fischer & Whitney, 2012), and attentional control (Barron et

al., 2015; Xuan et al., 2016). Analysis with connectivity-based thalamic atlas

showed that this region was most likely to overlap with the subdivision connected

with posterior parietal areas. Furthermore, RSFC and co-activation maps produced

by Neurosynth revealed connectivity patterns were suggestive of the attention

networks, especially the DAN. The diffusion imaging-based connectivity were also

suggestive of the DAN. At the behavioral level, studies have repeatedly

demonstrated the relationship between visuo-spatial attention and reading

(Facoetti, Franceschini, & Gori, 2019; Vidyasagar & Pammer, 2010). For example,

selective visual attention is associated with acquisition of orthographic knowledge

(Stevens & Bavelier, 2012) and decoding skill (Matthews & Martin, 2015). Besides,

both dyslexic adults and children manifest visuo-spatial attention deficits such as

lower visuo-spatial span capacities, and slower response in a visuo-spatial

attention-orienting task (Bosse, Tainturier, & Valdois, 2007; Buchholz & Davies,

2008). Longitudinal research also demonstrated impaired visuo-spatial processing

in pre-reading kindergarteners as an important risk factor for developing RD (Gori

& Facoetti, 2015). Together, these findings indicate that maturation of the pulvinar

and brain networks underlying visuo-spatial attention are parsimonious

neurocognitive mechanisms impacted by advanced PatAGE, impeding on reading

acquisition.

To date, research investigating the PatAGE effect on brain networks and

corresponding cognitive processes is scarce. As the first step, Shaw et al. (2012)

revealed PatAGE effects on cortical morphometry. However, the authors did not

examine the relationship with cognitive functions, making the study somewhat

inconclusive as to the role of PatAGE on neurocognitive processes. Taking one step

further, the current study revealed thalamic maturation as an intermediary

between PatAGE and reading – a specific behavioral phenotype, offering insights

into the complex mechanisms underlying PatAGE effects.

Limitations and future directions

In the present study, we observed a negative PatAGE effect on offspring's

reading and provided possible brain intermediator. Given the preliminary nature of

this investigation and the small sample size, the findings should be interpreted with

caution until they are replicated in large samples. Second, PatAGE in this study

was restricted to 25-47 years, with which we observed a negative linear

relationship. On the other hand, the findings do not necessarily extend to children

with extreme young and old fathers. For example, Saha et al. (2009) observed a non-

linear relationship with a range of PatAGE between 14 to 66 years, while the

correlation appears to be linear in the age range as ours. Of relevance, young

fatherhood is also associated with adverse outcomes in offspring but possibly due to

other factors including immature sperm and economic disadvantages (Chen et al.,

2008). Third, the complementary analyses implied DAN as the candidate functional

system associated with PatAGE. But it should be noted that the brain atlases and

public datasets implemented in Neurosynth are primarily from research on adults.

Since pediatric samples may have specific characteristics in terms of brain

organization, the neurostructural profile and function of the PatAGE-cluster need to

be re-visited as pediatric-specific atlases and tools become available. Fourth, while

we found that the left posterior thalamus mediated the PatAGE effect on reading, it

remains unknown why this subcortical structure is susceptible to advanced PatAGE

(and related to *de novo* mutations). Given that typical thalamus maturation is also

affected by prenatal and postnatal factors such as preterm birth (Ball et al., 2012),

questions including how PatAGE influences maturation of the thalamus and

relevant functional systems, together with other factors, require further

elaboration.

To advance understanding of the PatAGE effect on reading, future studies

are warranted in which a more comprehensive battery of behavioral tests (e.g.,

measuring visuo-spatial attention, executive function, etc.), neural measures (e.g.,

task-driven activation), and molecular approaches measuring the number and

origins of *de novo* mutations (e.g., trio-based whole-genome/exome sequencing; Jin

et al., 2017) are included. Fusing such neural, cognitive and molecular genetic

approaches at multiple levels will provide the much-needed vertical and multi-level

explanatory models that will further our understanding of risk factors associated

with poor reading. In particular, future research aiming at disentangling different

sources of genetic variations associated with reading development and their

interplays will greatly further our understanding. In addition, advanced research

designs such as intergenerational neuroimaging approach can be adopted to gain in-

depth knowledge on how multiple factors including PatAGE affect the development

of offspring's reading and the corresponding networks interactively from preliteracy

to mature stages of reading (Ho, Sanders, Gotlib, & Hoeft, 2016; Hoeft & Hancock,

2017).

Conclusion

The current study examined the PatAGE effect on offspring's reading at both

behavioral and neurobiological levels. The results provide initial evidence that

advanced PatAGE is an independent factor associated with poor reading in

beginning readers, above and beyond previously identified familial and cognitive-

linguistic precursors. This effect was mediated by maturation of the posterior

thalamus, suggesting a neurobiological pathway to intergenerational influence on

reading acquisition, complementing prior findings that offspring's reading is

influenced by parental reading via offspring's phonological skills (van Bergen et al.,

2015). Based on these evidences we argue that PatAGE should be regarded as an

important factor influencing literacy development, and included in a cumulative

risk (and protection) model (Hayiou-Thomas, Smith-Woolley, & Dale, 2020;

Menghini et al., 2010; Pennington, 2006; van Bergen, van der Leij, & de Jong,

2014).

References

Amso, D., & Scerif, G. (2015). The attentive brain: insights from developmental

cognitive neuroscience. *Nat Rev Neurosci, 16*(10), 606-619.

Arcaro, M. J., Pinsk, M. A., & Kastner, S. (2015). The Anatomical and Functional

Organization of the Human Visual Pulvinar. *J Neurosci, 35*(27), 9848-9871.

Ashburner, J. (2007). A fast diffeomorphic image registration algorithm.

Neuroimage, 38(1), 95-113.

Ball, G., Boardman, J. P., Rueckert, D., Aljabar, P., Arichi, T., Merchant, N., . . .

Counsell, S. J. (2012). The effect of preterm birth on thalamic and cortical

development. *Cereb Cortex, 22*(5), 1016-1024.

Barron, D. S., Eickhoff, S. B., Clos, M., & Fox, P. T. (2015). Human pulvinar

functional organization and connectivity. *Human Brain Mapping, 36*(7), 2417-

2431.

Basser, P. J., Pajevic, S., Pierpaoli, C., Duda, J., & Aldroubi, A. (2000). In vivo fiber

tractography using DT‐MRI data. *Magnetic Resonance in Medicine, 44*(4), 625-

632.

Biswal, B. B., Mennes, M., Zuo, X.-N., Gohel, S., Kelly, C., Smith, S. M., . . . Milham,

M. P. (2010). Toward discovery science of human brain function. *Proceedings of*

the National Academy of Sciences, 107(10), 4734-4739.

Black, J. M., Tanaka, H., Stanley, L., Nagamine, M., Zakerani, N., Thurston, A., . . .

Hoeft, F. (2012). Maternal history of reading difficulty is associated with

reduced language-related gray matter in beginning readers. *Neuroimage, 59*(3),

3021-3032.

Bosse, M.-L., Tainturier, M. J., & Valdois, S. (2007). Developmental dyslexia: The

visual attention span deficit hypothesis. *Cognition, 104*(2), 198-230.

Brain Development Cooperative, G. (2012). Total and Regional Brain Volumes in a

Population-Based Normative Sample from 4 to 18 Years: The NIH MRI Study of

Normal Brain Development. *Cereb Cortex, 22*(1), 1-12.

Breuss, M. W., Antaki, D., George, R. D., Kleiber, M., James, K. N., Ball, L. L., . . .

Gleeson, J. G. (2019). Autism risk in offspring can be assessed through

quantification of male sperm mosaicism. *Nature Medicine, 26*(1), 143-150.

Buchholz, J., & Davies, A. A. (2008). Adults with Dyslexia Demonstrate Attentional

Orienting Deficits. *Dyslexia, 14*(4), 247-270.

Cannon, M. (2009). Contrasting effects of maternal and paternal age on offspring

intelligence: the clock ticks for men too. *PLoS Med, 6*(3), e42.

Caravolas, M., Lervag, A., Mousikou, P., Efrim, C., Litavsky, M., Onochie-

Quintanilla, E., . . . Hulme, C. (2012). Common patterns of prediction of literacy

development in different alphabetic orthographies. *Psychol Sci, 23*(6), 678-686.

Chen, X. K., Wen, S. W., Krewski, D., Fleming, N., Yang, Q. Y., & Walker, M. C.

(2008). Paternal age and adverse birth outcomes: teenager or 40+, who is at

risk? *Human Reproduction, 23*(6), 1290-1296.

Clark, K. A., Helland, T., Specht, K., Narr, K. L., Manis, F. R., Toga, A. W., &

Hugdahl, K. (2014). Neuroanatomical precursors of dyslexia identified from pre-

reading through to age 11. *Brain, 137*(Pt 12), 3136-3141.

Couture, V., Delisle, S., Mercier, A., & Pennings, G. (2020). The other face of

advanced paternal age: a scoping review of its terminological, social, public

health, psychological, ethical and regulatory aspects. *Human reproduction*

update.

D'Onofrio, B. M., Rickert, M. E., Frans, E., Kuja-Halkola, R., Almqvist, C.,

Sjölander, A., . . . Lichtenstein, P. (2014). Paternal age at childbearing and

offspring psychiatric and academic morbidity. *JAMA Psychiatry, 71*(4), 432-438.

de Moura, L. M., Cogo-Moreira, H., Brandão de Ávila, C. R., Pan, P. M., Gadelha,

A., Moriyama, T., . . . Picon, F. A. (2016). Children with Poor Reading Skills at

the Word Level Show Reduced Fractional Anisotropy in White Matter Tracts of

Both Hemispheres. *Brain Connect, 6*(7), 519-523.

De Rubeis, S., He, X., Goldberg, A. P., Poultney, C. S., Samocha, K., Cicek, A. E., . . .

Buxbaum, J. D. (2014). Synaptic, transcriptional and chromatin genes

disrupted in autism. *Nature, 515*(7526), 209-215.

Deciphering Developmental Disorders Study. (2017). Prevalence and architecture of

de novo mutations in developmental disorders. *Nature, 542*(7642), 433-438.

Diaz, B., Hintz, F., Kiebel, S. J., & von Kriegstein, K. (2012). Dysfunction of the

auditory thalamus in developmental dyslexia. *Proc Natl Acad Sci U S A,*

109(34), 13841-13846.

Dunn, D. M., & Dunn, L. M. (2007). *Peabody picture vocabulary test: Manual*:

Pearson.

Edwards, R. D., & Roff, J. (2010). Negative effects of paternal age on children's

neurocognitive outcomes can be explained by maternal education and number of

siblings. *PLoS One, 5*(9), e12157.

Facoetti, A., Franceschini, S., & Gori, S. (2019). Role of Visual Attention in

Developmental Dyslexia. In C. Perfetti, K. Pugh, & L. Verhoeven (Eds.),

Developmental Dyslexia across Languages and Writing Systems (pp. 307-326).

Cambridge: Cambridge University Press.

Fischer, J., & Whitney, D. (2012). Attention gates visual coding in the human

pulvinar. *Nature Communications, 3*, 1051.

Giraldo-Chica, M., Hegarty, J. P., 2nd, & Schneider, K. A. (2015). Morphological

differences in the lateral geniculate nucleus associated with dyslexia.

Neuroimage Clin, 7, 830-836.

Girard, S. L., Bourassa, C. V., Lemieux Perreault, L. P., Legault, M. A., Barhdadi,

A., Ambalavanan, A., . . . Rouleau, G. A. (2016). Paternal Age Explains a Major

Portion of De Novo Germline Mutation Rate Variability in Healthy Individuals.

PLoS One, 11(10), e0164212.

Gori, S., & Facoetti, A. (2015). How the visual aspects can be crucial in reading

acquisition: The intriguing case of crowding and developmental dyslexia.

Journal of vision, 15(1), 8-8.

Grasby, K. L., Jahanshad, N., Painter, J. N., Colodro-Conde, L., Bralten, J., Hibar,

D. P., . . . Enhancing NeuroImaging Genetics through Meta-Analysis

Consortium -Genetics working, g. (2020). The genetic architecture of the human

cerebral cortex. *Science, 367*(6484).

Hayes, A. F. (2013). *Introduction to mediation, moderation, and conditional process*

analysis: A regression-based approach: Guilford Press.

Hayiou-Thomas, M. E., Smith-Woolley, E., & Dale, P. S. (2020). Breadth versus

depth: Cumulative risk model and continuous measure prediction of poor

language and reading outcomes at 12. *Dev Sci,* e12998.

Ho, T. C., Sanders, S. J., Gotlib, I. H., & Hoeft, F. (2016). Intergenerational

Neuroimaging of Human Brain Circuitry. *Trends Neurosci, 39*(10), 644-648.

Hoeft, F., & Hancock, R. (2017). Intergenerational Transmission of Reading and

Reading Brain Networks. In A. M. Galaburda, N. Gaab, & F. Hoeft (Eds.),

Dyslexia and Neuroscience: The Geschwind-Galaburda Hypothesis 30 Years

Later. Baltimore: Paul H. Brookes Publishing Co.

Hulme, C., Snowling, M., Caravolas, M., & Carroll, J. (2005). Phonological skills are

(probably) one cause of success in learning to read: A comment on Castles and

Coltheart. *Scientific Studies of Reading, 9*(4), 351-365.

Jakab, A., Blanc, R., Berényi, E., & Székely, G. (2012). Generation of individualized

thalamus target maps by using statistical shape models and thalamocortical

tractography. *American Journal of Neuroradiology, 33*(11), 2110-2116.

Jayasekara, R., & Street, J. (1978). Parental age and parity in dyslexic boys.

Journal of biosocial science, 10(03), 255-261.

Jin, Z. B., Li, Z., Liu, Z., Jiang, Y., Cai, X. B., & Wu, J. (2017). Identification of de

novo germline mutations and causal genes for sporadic diseases using trio -

based whole - exome/genome sequencing. *Biological Reviews, 93*(2), 1014-1031.

Jónsson, H., Sulem, P., Kehr, B., Kristmundsdottir, S., Zink, F., Hjartarson, E., . . .

Stefansson, K. (2017). Parental influence on human germline de novo mutations

in 1,548 trios from Iceland. *Nature,549*(7673), 519-522.

Khandwala, Y. S., Zhang, C. A., Lu, Y., & Eisenberg, M. L. (2017). The age of

fathers in the USA is rising: an analysis of 168 867 480 births from 1972 to

2015. *Human Reproduction*, 1-7.

Kohler, H.-P., Billari, F. C., & Ortega, J. A. (2002). The Emergence of Lowest-Low

Fertility in Europe During the 1990s. *Population and Development Review,*

28(4), 641-680.

Kong, A., Frigge, M. L., Masson, G., Besenbacher, S., Sulem, P., Magnusson, G., . . .

Stefansson, K. (2012). Rate of de novo mutations and the importance of father's

age to disease risk. *Nature, 488*(7412), 471-475.

Koyama, M. S., Molfese, P. J., Milham, M. P., Mencl, W. E., & Pugh, K. R. (2020).

Thalamus is a common locus of reading, arithmetic, and IQ: Analysis of local

intrinsic functional properties. *bioRxiv*, 2020.2005.2005.076232.

Krauth, A., Blanc, R., Poveda, A., Jeanmonod, D., Morel, A., & Szekely, G. (2010). A

mean three-dimensional atlas of the human thalamus: generation from multiple

histological data. *Neuroimage, 49*(3), 2053-2062.

Leemans, A., Jeurissen, B., Sijbers, J., & Jones, D. (2009). *ExploreDTI: a graphical*

toolbox for processing, analyzing, and visualizing diffusion MR data. Paper

presented at the 17th Annual Meeting of Intl Soc Mag Reson Med.

Leemans, A., & Jones, D. K. (2009). The B-matrix must be rotated when correcting

for subject motion in DTI data. *Magnetic Resonance in Medicine, 61*(6), 1336-

1349.

Lefly, D. L., & Pennington, B. F. (2000). Reliability and validity of the adult reading

history questionnaire. *J Learn Disabil, 33*(3), 286-296.

Mather, M. (1998). Woodcock Reading Mastery Tests—Revised/Normative Update.

Circle Pines, MN: American Guidance Service.

Matthews, A. J., & Martin, F. H. (2015). Spatial attention and reading ability: ERP

correlates of flanker and cue-size effects in good and poor adult phonological

decoders. *Brain and Language, 151*, 1-11.

McGrew, K. S., & Schrank, F. A. (2007). Technical manual. Woodcock-Johnson III

normative update. *Rolling Meadows, IL: Riverside Publishing.*

Menghini, D., Finzi, A., Benassi, M., Bolzani, R., Facoetti, A., Giovagnoli, S., . . .

Vicari, S. (2010). Different underlying neurocognitive deficits in developmental

dyslexia: a comparative study. *Neuropsychologia, 48*(4), 863-872.

Müller-Axt, C., Anwander, A., & von Kriegstein, K. (2017). Altered Structural

Connectivity of the Left Visual Thalamus in Developmental Dyslexia. *Current*

Biology, 27(23), 3692-3698.

Noble, K. G., Wolmetz, M. E., Ochs, L. G., Farah, M. J., & McCandliss, B. D. (2006).

Brain–behavior relationships in reading acquisition are modulated by

socioeconomic factors. *Dev Sci, 9*(6), 642-654.

Olson, R., Wise, B., Conners, F., Rack, J., & Fulker, D. (1989). Specific deficits in

component reading and language skills: genetic and environmental influences.

J Learn Disabil, 22(6), 339-348.

Olson, R. K., Keenan, J. M., Byrne, B., Samuelsson, S., Coventry, W. L., Corley, R., .

. . Hulslander, J. (2011). Genetic and Environmental Influences on Vocabulary

and Reading Development. *Sci Stud Read, 15*(1), 26-46.

Pan, J., Kong, Y., Song, S., McBride, C., Liu, H., & Shu, H. (2016). Socioeconomic

status, parent report of children's early language skills, and late literacy skills:

a long term follow-up study among Chinese children. *Reading and Writing*, 1-

16.

Paz-Alonso, P. M., Oliver, M., Lerma-Usabiaga, G., Caballero-Gaudes, C., Quiñones,

I., Suárez-Coalla, P., . . . Carreiras, M. (2018). Neural correlates of phonological,

orthographic and semantic reading processing in dyslexia. *NeuroImage:*

Clinical, 20, 433-447.

Pennington, B. F. (2006). From single to multiple deficit models of developmental

disorders. *Cognition, 101*(2), 385-413.

Petrill, S. A., Deater-Deckard, K., Thompson, L. A., DeThorne, L. S., &

Schatschneider, C. (2006). Genetic and environmental effects of serial naming

and phonological awareness on early reading outcomes. *Journal of Educational*

Psychology, 98(1), 112-121.

Pierpaoli, C., & Basser, P. J. (1996). Toward a quantitative assessment of diffusion

anisotropy. *Magnetic Resonance in Medicine, 36*(6), 893-906.

Price, J. (2008). Parent-Child Quality Time Does Birth Order Matter? *Journal of*

Human Resources, 43(1), 240-265.

Raznahan, A., Shaw, P. W., Lerch, J. P., Clasen, L. S., Greenstein, D., Berman, R., .

. . Giedd, J. N. (2014). Longitudinal four-dimensional mapping of subcortical

anatomy in human development. *Proc Natl Acad Sci U S A, 111*(4), 1592-1597.

Ridgway, G., Camara, O., Scahill, R., Crum, W., Whitcher, B., Fox, N., & Hill, D.

(2007). Longitudinal Voxel-based morphometry with unified segmentation:

evaluation on simulated Alzheimer's disease, British Machine Vision

Association.

Saha, S., Barnett, A. G., Foldi, C., Burne, T. H., Eyles, D. W., Buka, S. L., &

McGrath, J. J. (2009). Advanced paternal age is associated with impaired

neurocognitive outcomes during infancy and childhood. *PLoS Med, 6*(3),

e1000040.

Segers, E., Damhuis, C. M., van de Sande, E., & Verhoeven, L. (2016). Role of

executive functioning and home environment in early reading development.

Learning and Individual Differences, 49, 251-259.

Shaw, L. A. (2010). *The relationship between the Wechsler Intelligence Scale for*

Children-and the Woodcock-Johnson III Tests of Cognitive Abilities in a

clinically referred pediatric population: Nova Southeastern University.

Shaw, P., Gilliam, M., Malek, M., Rodriguez, N., Greenstein, D., Clasen, L., . . .

Giedd, J. (2012). Parental age effects on cortical morphology in offspring. *Cereb*

Cortex, 22(6), 1256-1262.

Sobotka, T. (2010). Shifting Parenthood to Advanced Reproductive Ages: Trends,

Causes and Consequences. In *A young generation under pressure?* (pp. 129-154).

Berlin, Heidelberg: Springer Berlin Heidelberg.

Soden, B., Christopher, M. E., Hulslander, J., Olson, R. K., Cutting, L., Keenan, J.

M., . . . Petrill, S. A. (2015). Longitudinal stability in reading comprehension is

largely heritable from grades 1 to 6. *PLoS One, 10*(1), e0113807.

Stevens, C., & Bavelier, D. (2012). The role of selective attention on academic

foundations: a cognitive neuroscience perspective. *Dev Cogn Neurosci, 2 Suppl*

1, S30-48.

Sussman, D., Leung, R. C., Chakravarty, M. M., Lerch, J. P., & Taylor, M. J. (2016).

Developing human brain: age‐related changes in cortical, subcortical, and

cerebellar anatomy. *Brain and Behavior, 6*(4), e00457.

Swagerman, S. C., van Bergen, E., Dolan, C., de Geus, E. J. C., Koenis, M. M. G.,

Hulshoff Pol, H. E., & Boomsma, D. I. (2017). Genetic transmission of reading

ability. *Brain and Language, 172*, 3-8.

Thompson, P. A., Hulme, C., Nash, H. M., Gooch, D., Hayiou-Thomas, E., &

Snowling, M. J. (2015). Developmental dyslexia: predicting individual risk. *J*

Child Psychol Psychiatry, 56(9), 976-987.

Torgesen, J. K., Wagner, R., & Rashotte, C. (1999). TOWRE–2 Test of Word

Reading Efficiency. *Austin, TX: Pro-Ed*.

Tschentscher, N., Ruisinger, A., Blank, H., Diaz, B., & von Kriegstein, K. (2018).

Reduced structural connectivity between left auditory thalamus and the

motion-sensitive planum temporale in developmental dyslexia. *arXiv preprint*

arXiv:1811.11658.

van Bergen, E., Bishop, D., van Zuijen, T., & de Jong, P. F. (2015). How does

parental reading influence children's reading? A study of cognitive mediation.

Scientific Studies of Reading, 19(5), 325-339.

van Bergen, E., van der Leij, A., & de Jong, P. F. (2014). The intergenerational

multiple deficit model and the case of dyslexia. *Front Hum Neurosci, 8*(346),

346.

Vidyasagar, T. R., & Pammer, K. (2010). Dyslexia: a deficit in visuo-spatial

attention, not in phonological processing. *Trends Cogn Sci, 14*(2), 57-63.

Wagner, R. K., Torgesen, J. K., & Rashotte, C. A. (1999). *Comprehensive test of*

phonological processing: CTOPP: Pro-ed.

Weiser, M., Reichenberg, A., Werbeloff, N., Kleinhaus, K., Lubin, G.,

Shmushkevitch, M., . . . Davidson, M. (2008). Advanced Parental Age at Birth Is

Associated With Poorer Social Functioning in Adolescent Males: Shedding Light

on a Core Symptom of Schizophrenia and Autism. *Schizophr Bull, 34*(6), 1042-

984   1046.

985   Wolf, M., & Denkla, M. (2005). Rapid Automatized Naming and Rapid Alternating

986   Stimulus Tests. Austin, TX: PROED. In: Inc.

987   Xuan, B., Mackie, M. A., Spagna, A., Wu, T., Tian, Y., Hof, P. R., & Fan, J. (2016).

988   The activation of interactive attentional networks. *Neuroimage, 129*, 308-319.

989   Yang, D. Y., Rosenblau, G., Keifer, C., & Pelphrey, K. A. (2015). An integrative

990   neural model of social perception, action observation, and theory of mind.

991   *Neurosci Biobehav Rev, 51*, 263-275.

992   Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., & Wager, T. D. (2011).

993   Large-scale automated synthesis of human functional neuroimaging data.

994   *Nature methods, 8*(8), 665-670.

995   Yeatman, J. D., Dougherty, R. F., Ben-Shachar, M., & Wandell, B. A. (2012).

996   Development of white matter and reading skills. *Proc Natl Acad Sci U S A,*

997   *109*(44), E3045-3053.

998   Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead,

999   M., . . . Polimeni, J. R. (2011). The organization of the human cerebral cortex

1000   estimated by intrinsic functional connectivity. *Journal of Neurophysiology,*

1001   *106*(3), 1125-1165.

1002

Table Legend

Table 1 Demographic profiles, familial variables and performance on reading-

related tests (n = 43). *Acronyms: ARHQ, Adult Reading History Questionnaire; CS,*

composite score; CTOPP BW, Comprehensive Test of Phonological Processing,

Blending sub-test; CTOPP EL, Comprehensive Test of Phonological Processing,

Elision sub-test; CTOPP MD, Comprehensive Test of Phonological Processing,

Memory for Digit sub-test; CTOPP NR, Comprehensive Test of Phonological

Processing, Nonword Repetition sub-test; HOME, Home Observation Measurement of

the Environment; Mat, maternal; Pat, paternal; PPVT, Peabody Picture Vocabulary

Test; RAN COL, Rapid Naming, Colors sub-test; RAN LTR, Rapid Naming, Letters

sub-test; RAN NUM, Rapid Naming, Numbers sub-test; RAN OBJ, Rapid Naming,

Objects sub-test; RS, raw score; SES, socioeconomic status; SS, standard score;

TOWRE PDE, Test of Word Reading, Phonemic Decoding Efficiency sub-test;

TOWRE SWE, Test of Word Reading, Sight Word Efficiency sub-test; WJA RF,

Woodcock-Johnson III Tests of Achievement, Reading Fluency sub-test; WJA SP,

Woodcock-Johnson III Tests of Achievement, Spelling sub-test; WJC CF, Woodcock-

Johnson III Tests of Cognitive Abilities, Concept Formation sub-test; WJC NR,

Woodcock-Johnson III Tests of Cognitive Abilities, Numbers Reversed sub-test; WJC

VC, Woodcock-Johnson III Tests of Cognitive Abilities, Verbal Comprehension sub-

test; WJC VM, Woodcock-Johnson III Tests of Cognitive Abilities, Visual Matching

sub-test; WRMT LID, Woodcock Reading Mastery Test, Letter Identification sub-test;

WRMT PC, Woodcock Reading Mastery Test, Passage Comprehension sub-test;

WRMT WA, Woodcock Reading Mastery Test, Word Attack sub-test; WRMT WID,

Woodcock Reading Mastery Test, Word Identification sub-test.

Table 2 Results of multiple linear regression analyses examining the unique

contribution of paternal age on offspring's reading at time-point 2. *Acronyms:*

ARHQ, Adult Reading History Questionnaire; EDU, educational level; HOME,

Home Observation Measurement of the Environment; Mat, maternal; PA,

phonological awareness; Pat, paternal; pIQ, performance intelligence quotient; RAN,

rapid naming; SES, socioeconomic status; t1, time-point 1; t2, time-point 2.

Figure Legends

Fig. 1 Principal components extracted from reading-related tests. **A**. Component

loadings for each factor at time-point 1. **B**. Component loadings for each factor at

time-point 2. *Acronyms: CTOPP BW, Comprehensive Test of Phonological*

Processing, Blending sub-test; CTOPP EL, Comprehensive Test of Phonological

Processing, Elision sub-test; CTOPP MD, Comprehensive Test of Phonological

Processing, Memory for Digit sub-test; CTOPP NR, Comprehensive Test of

Phonological Processing, Nonword Repetition sub-test; RAN COL, Rapid Naming,

Colors sub-test; RAN LTR, Rapid Naming, Letters sub-test; RAN NUM, Rapid

Naming, Numbers sub-test; RAN OBJ, Rapid Naming, Objects sub-test; t1, time-

point 1; t2, time-point 2; TOWRE PDE, Test of Word Reading, Phonemic Decoding

Efficiency sub-test; TOWRE SWE, Test of Word Reading, Sight Word Efficiency sub-

test; WJA RF, Woodcock-Johnson III Tests of Achievement, Reading Fluency sub-

test; WJA SP, Woodcock-Johnson III Tests of Achievement, Spelling sub-test; WRMT

LID, Woodcock Reading Mastery Test, Letter Identification sub-test; WRMT PC,

Woodcock Reading Mastery Test, Passage Comprehension sub-test; WRMT WA,

Woodcock Reading Mastery Test, Word Attack sub-test; WRMT WID, Woodcock

Reading Mastery Test, Word Identification sub-test.

Fig. 2 Results of whole-brain longitudinal voxel-based morphometry and mediation

analysis. **A.** Brain regions significantly correlated with paternal age (yellow) and

composite score of reading at time-point 2 (cyan). **B**. Scatter plot of the relationship

between gray matter volume change in the yellow cluster and paternal age. The

linear regression line is presented. **C**. Scatter plot of the relationship between gray

matter volume change in the cyan cluster and reading composite score at time-point

2. The linear regression line is presented. **D.** The effect of paternal age on

offspring's reading is mediated by gray matter volume change in the thalamus.

Confounds were controlled statistically. The bias corrected 95% confidence interval

for indirect effect was [-0.052 -0.043], indicating a significant mediation

relationship.

Fig. 3 Localization and structural connectivity pattern of the PatAGE-cluster (i.e.,

the left posterior thalamus) according to the Morel and Thalamic Connectivity

atlases. **A.** Bar plot displaying the percentage of total voxels in the PatAGE-cluster

overlaps with divisions of the Morel atlas. **B.** Bar plot showing the probability of the

cluster belonging to different subdivisions of the Thalamic connectivity atlas,

calculated by 'autoaq' function implemented in FSL

(https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/Atlases). *Acronyms: CL, central lateral nucleus;*

CM, central median nucleus; LP, lateral posterior nucleus; VLpv, ventral lateral

posterior nucleus, ventral; VPI, ventral posterior inferior nucleus; VPL, ventral

posterior lateral nucleus; VPM, ventral posterior medial nucleus.

Fig. 4 Functional co-activation and connectivity of the PatAGE-cluster produced by

Neurosynth. **A.** Brain map representing overlapping regions between co-activation

and resting-state functional connectivity maps of the PatAGE-cluster. **B.** Bar plot

showing the degree of overlap between the overlapping areas and Yeo's 7 functional

networks represented by Dice coefficients. Dice coefficient measures the similarity

between the overlapping areas and a given function network, ranging from 0 to 1.

While 0 indicates the two networks are disjoint, 1 indicates the two networks are

identical. *Acronyms: LH, left hemisphere; RH, right hemisphere.*

Fig. 5 White matter tractography of the PatAGE-related thalamic region using

subject-specific diffusion imaging data. **A.** Example of reconstructed fibers in a

representative child with the seed being the PatAGE-cluster. **B.** Intersection across

23 children with diffusion imaging data is shown for demonstrative purposes. In

particular, only fibers (streamlines) observed in more than 25% of the subjects (6

children) are displayed. The color bar represents the number of subjects where the

streamline is observed in a given voxel. **C.** The DAN compared to the VAN derived

from Yeo's 7 networks showed significantly greater (normalized) number of

streamlines to go through the PatAGE-cluster. **D.** Scatter plot of the positive

correlation between paternal age and number of streamlines labeled as DAN

passing through the cluster. The linear regression line and R^2 are shown. **E.** Scatter

plot of the negative correlation between reading composite scores at time-point 2

and the number of streamlines passing through the PatAGE-cluster and DAN. The

linear regression line and R^2 are presented. *Acronyms: DAN, dorsal attention*

network; LH, left hemisphere; RH, right hemisphere; VAN, ventral attention

network.

Tables

Table 1 Demographic profiles, familial variables and performance on reading-related tests (n = 43).

	Mean	Std.Dev	Min	Max
Time-point 1				
Age (years)	5.58	0.43	5.03	6.99
Gender, Male (%)	60.50	--	--	--
Handedness, Right (%)	88.40	--	--	--
WJC VC (SS)	121.53	13.43	86	145
WJC CF (SS)	118.16	11.43	94	137
WJC VM (SS)	105.65	11.83	72	127
WJC NR (SS)	112.05	12.06	83	138
PPVT (SS)	121.23	9.94	97	148
# Older Siblings (RS)	0.65	0.81	0	3
# Younger Siblings (RS)	0.63	0.62	0	2
PatAGE (years)	36.12	4.91	24.78	46.71
MatAGE (years)	33.01	4.09	23.04	41.08
PatARHQ (RS)	0.35	0.14	0.09	0.66
MatARHQ (RS)	0.31	0.15	0.07	0.67
PatEDU (years)	16.95	2.05	13	22
MatEDU (years)	16.97	2.04	12	22
SES (CS) [a]	0.04	1.00	-2.87	2.38
HOME (RS) [b]	51.39	2.25	44	55
CTOPP BW (SS) [c]	12.28	1.80	8	17
CTOPP EL (SS) [c]	11.93	2.80	7	19
CTOPP MD (SS) [c]	10.79	2.22	7	16
CTOPP NR (SS) [c]	11.23	2.85	6	19
RAN OBJ (SS)	100.56	17.57	55	135

RAN COL (SS)	97.98	16.51	55	137
WRMT LID (SS)	109.84	10.84	80	138
Time-point 2				
Age (years)	8.30	0.46	7.51	9.76
WJC VC (SS)	116.07	10.18	92	144
WJC CF (SS)	118.21	12.29	97	150
WJC VM (SS)	99.23	15.04	77	138
WJC NR (SS)	109.26	15.13	80	140
PPVT (SS)	120.02	14.56	81	160
TOWRE SWE (SS)	111.49	12.29	86	138
TOWRE PDE (SS)	106.35	14.85	77	144
WRMT WID (SS)	116.23	11.66	94	139
WRMT WA (SS)	113.84	14.40	90	146
WRMT PC (SS)	114.91	9.95	99	141
WJA RF (SS)	112.49	16.31	84	162
WJA SP (SS)	105.28	18.33	74	148
CTOPP BW (SS) [c]	12.67	2.24	6	16
CTOPP EL (SS) [c]	13.00	3.11	4	17
CTOPP MD (SS) [c]	10.47	2.60	5	15
CTOPP NR (SS) [c]	10.09	2.26	6	16
RAN NUM (SS)	100.35	12.47	76	129
RAN LTR (SS)	102.98	11.63	78	134
RAN OBJ (SS)	96.53	17.07	62	132
RAN COL (SS)	97.02	15.29	60	121

Notes: [a] SES: n = 38; [b] HOME: n = 41; [c] T Scores are presented for CTOPP sub-tests where mean is 10 and SD is 3. All other test scores are in standard scores where mean is 100 and SD is 15.

Table 2 Results of multiple linear regression analyses examining the unique

contribution of paternal age on offspring's reading at time-point 2

Model	Step	Predictor	ΔR^2	β
1	1	Age (*t*2)	0.284 *	-0.351 *
		Sex		-0.022
		Handedness		0.081
		Average pIQ		0.302 *
	2	PatAGE	0.149 **	-0.393 **
2	1	Age (*t*2)	0.284 *	-0.351 *
		Sex		-0.022
		Handedness		0.079
		Average pIQ		0.309 *
	2	MatAGE	0.052 †	0.026
	3	PatAGE	0.097 *	-0.408 *
3	1	Age (*t*2)	0.279 *	-0.315 †
		Sex		-0.002
		Handedness		0.119
		Average pIQ		0.203
	2	MatAGE	0.071 †	0.269
	3	PatARHQ	0.170	-0.129
		MatARHQ		-0.162
		# Older Siblings		-0.298
		# Younger Siblings		-0.171
		PatEDU		0.031
		MatEDU		-0.075
		SES		-0.214
		HOME		0.233
	4	PatAGE	0.107 *	-0.592 *

4	1	Age (*t2*)	0.279 *	-0.261
		Sex		-0.051
		Handedness		0.136
		Average pIQ		-0.013
	2	MatAGE	0.071 †	0.200
	3	PatARHQ		-0.105
		MatARHQ		-0.039
		# Older Siblings		-0.174
		# Younger Siblings		-0.144
		PatEDU		0.086
		MatEDU		-0.211
		SES		-0.146
		HOME		0.201
	4	Time 1 PA	0.138 *	0.423 *
		Time 1 RAN		0.350 *
	5	PatAGE	0.095 *	-0.567 *

Note: β is value at the final step (all predictors included). ** $p < 0.01$; * $p < 0.05$, † $p < 0.1$

Figures

Fig. 1

Fig. 2

Fig. 3

Fig. 4

A Intersection of Neurosynth-Derived Maps

RH

Frontal eye field

Inferior parietal sulci

Z = 54

Middle temporal visual area

X = -47 (LH)

Overlapping areas between co-activation and RSFC maps

B Similarity with the Intrinsic Functional Networks

Fig. 5