Body size and tree species composition determine variation in prey consumption in a forest-inhabiting generalist predator

Irene M. van Schrojenstein Lantman[a], Eero J. Vesterinen[b,c,d], Lionel R. Hertzog[a,e], An Martel[e], Kris Verheyen[f], Luc Lens[a], Dries Bonte[a]

Affiliations

[a]Terrestrial Ecology Unit, Department of Biology, Ghent University, K. L. Ledeganckstraat 35, 9000 Gent, Belgium

[b]Spatial Foodweb Ecology Group, Department of Agricultural Sciences, PO Box 27, (Latokartanonkaari 5), FI-00014 University of Helsinki, Helsinki, Finland

[c]Department of Ecology, Swedish University of Agricultural Sciences, Box 7044, 750 07 Uppsala, Sweden

[d]Department Biology, University of Turku, 20014 Turku, Finland

e Thünen Institut for Biodiversity, Bundesallee 68, 38116 Braunschweig, Germany

[e]Department Pathology, Bacteriology and Avian Diseases, Ghent University, Salisburylaan 133, 9820 Merelbeke, Belgium

[f]Forest & Nature Lab, Department of Environment, Ghent University, Geraardsbergse steenweg 267, 9090 Gontrode, Belgium

Corresponding author: Dries Bonte (dries.bonte@ugent.be)

<u>Short running title</u>: body size, forest structure and spider prey

Abstract

Trophic interactions may strongly depend on body size and environmental variation, but this prediction has been seldom tested in nature. Many spiders are generalist predators that use webs to intercept flying prey. The size and mesh of orb webs increases with spider size, allowing a more efficient predation on larger prey. We studied to this extent the orb-weaving spider *Araneus diadematus* inhabiting forest fragments differing in edge distance, tree diversity and tree species. These environmental variables are known to correlate to insect composition, richness and abundance. We anticipated these forest characteristics to be a principle driver of prey consumption. We additionally hypothesised them to impact spider size at maturity and expect shifts towards larger prey-size distributions in larger individuals independently from the environmental context.

We quantified spider diet by means of metabarcoding of nearly 1000 *A. diadematus* from a total of 53 forest plots. This approach allowed a massive screening of consumption dynamics in nature, though at the cost of identifying the exact prey identity, as well as their abundance and putative intraspecific variation. Our study confirmed *A. diadematus* as a generalist predator, with more than 300 prey ZOTUs detected in total. At the individual level, we found large spiders to consume fewer different species, but adding larger species to their diet. Tree species composition affected both prey species richness and size in the spider's diet, although tree diversity per se had no influence on the consumed prey. Edges had an indirect effect on the spider diet as spiders closer to the forest edge were larger and therefore consumed larger prey. We conclude that both intraspecific size variation and tree species composition shape the consumed prey of this generalist predator.

Keywords: Metabarcoding, predator-prey interaction, tree diversity, edge effects, prey-size spectrumμ

Short title: Body size, forest structure and spider prey

Introduction

Trophic interactions are a key component of ecological networks (Thébault & Loreau, 2005; Landi *et al.*, 2018). Food webs are strongly impacted by the level of diet specialisation of the involved consumers. With higher per capita consumption rates, specialist predators are more effective in prey control than generalists (Diehl, Sereda, Wolters & Birckhofer, 2013), yet opportunistic and generalist predators provide stronger stabilizing effect within a food web (Gross, Rudolg, Levin & Dieckmann, 2009). Generalist species may, however, be a collection of individuals that specialise on specific components of the full prey spectrum (Bolnick *et al.*, 2003; Araújo, Bolnick & Layman, 2011). This intraspecific variation is known to arise from behavioural or physiological niche divergence, but equally from environmental variation constraining prey availability and the species spectrum (Bolnick et al. 2013).

Spiders are known to be the principal consumers of insects (Nyffeler & Birkhofer, 2017), and many species are opportunistic and generalist predators that rely on active hunting or trapping for prey capture (Eitzinger *et al.*, 2019). Orb-web spiders build vertical webs from silk to intercept flying prey, from small insects to smaller vertebrates (e.g. Brooks, 2012). As interception-feeders, their prey spectrum, is by definition strongly determined by the environment. The diet of generalist and opportunistic predators is therefore expected to reflect the species richness of the prey community (Bison *et al.*, 2015; Schmidt, Mosbacher, Eitzinger, Vesterinen & Roslin, T., 2018; Etzinger, Roslin, Vesterinen, Robinson & O'Gorman 2021).

Insect, and thus prey, diversity is largely structured by the plant community composition and plant diversity (Price, 2002; Scherber *et al.*, 2010; Rzanny *et al.*, 2013). By providing a larger variation in resources, increases in plant diversity are known to promote coexistence of herbivore and predator species (Haddad *et al.*, 2009; Hertzog, Meyer, Weisser & Ebeling, 2016; O'Brien *et al.*, 2017).

Additionally, as plant diversity is linked to an increase in productivity (Loreau *et al.*, 2001; Hooper *et*

al., 2005), prey abundance and mean prey size should increase as well (Allen *et al.*, 2006). Despite

the general prediction of plant diversity increasing insect diversity (Scherber *et al.*, 2010), plant

species identity may have larger effects on the community composition of arthropods than its

diversity per se (Vehviläinen, Koricheva & Ruohomäki, 2008; Scherber, Vockenhuber, Stark, Meyer &

Tscharntke 2014; van Schrojenstein Lantman et al. 2019). Equally, and independent from these

primary producer dynamics, insect and thus spider prey variation can be impacted by the spatial

dimensions of their habitat, especially by habitat size and isolation (Debinski & Holt, 2000; Krauss *et*

al., 2010). According to the trophic rank hypothesis, higher trophic levels are generally more affected

by habitat fragmentation than lower trophic levels (Holt, 2002; Cagnolo *et al.*, 2009; Martinson &

Fagan, 2014, Hillaert, Vandegehuchte, Hovestadt & Bonte, 2018). Edge densities increase with

fragmentation (Haddad *et al.*, 2015) and are prominent drivers of insect composition (Murcia, 1995;

Schmidt, Jochheim, Kersebaum & Lischeid, 2017) and abundance (Debinski & Holt, 2000; Rand,

Tylianakis& Tscharntke, 2006). In forests fragments, arthropod diversity has been shown to be

strongly determined by tree species diversity, identify and patch size (Hertzog et al. 2019, 2020;

Perring et al. 2021). The patch-size effects are directly translated to edge effects which allow allowing

coexistence of both forest and matrix-species (e.g., Rand et al. 2006). The warmer microclimate at

forest edges also favours smaller arthropods (Atkinson & Sibly, 1997; Kingsolver & Huey, 2008).

In general, and despite potential variation due to weather conditions and prey availability (Schneider

& Vollrath, 1998; Bonte, Lanckacker, Wiersma & Lens, 2008; Sensenig, Agnarsson, Gondek &

Blackledge, 2010; Tew, Adamson & Hesselberg, 2015), larger orb web spider species (Dahirel,

Dierick, Cock & Bonte, 2017) but also larger individuals of the same species (Bonte *et al.*, 2008;

Dahirel, De Cock, Vantieghem & Bonte 2019) build larger webs with larger mesh sizes. This allows

predation on larger prey that provide essential resources for reproduction (Venner & Casas, 2005).

Since predator body size determines the maximum prey size that can be caught (as shown for spiders

by Nentwig & Wissel, 1986), large predators can be expected to expand or shift their prey

composition towards larger prey (Woodward & Hildrew, 2002). This size-dependency of web-building

has been demonstrated in the cross spider *Araneaus diadematus*. As other in species (Dahirel et al.

2017), *A. diadematus* may show individual-level resource specialization in relation to the prey-

availability by adapting web building behaviour (Schnieder & Vollrath 1998). More-over, since prey

capture does not automatically imply prey consumption (Janetos, 1982), the relationship between

spider size and prey (size) consumption remains untested.

Describing and understanding trophic interactions in complex habitats, such as forests, rather than in

experiments or simple agricultural systems is a challenging endeavour. We lack a profound

understanding on the relative importance of environmental relative to intraspecific variation for

realised trophic interactions in nature. We therefore engaged in an unprecedented barcoding

(O'Rorke, Lavery & Jeffs, 2012; Pompanon *et al.*, 2012) of the species' gut content to understand to

what extent these components of environmental variation as well as predator size determine tropic

interactions of *A. diadematus* in forest fragments dominated by different proportions of *Quercus*

robur L. (pedunculate oak), *Quercus rubra* L. (red oak) and *F. sylvatica* L. (common beech). We

specifically tested the hypothesis that prey richness in the diet increases with both tree diversity, the

relative availability of pedunculated oak and forest edge proximity, due to already documented

increases in prey species availability (Hertzog et al. 2020). Independent of these tree composition

and spatial effects, we tested the prediction that large spiders consume larger prey, and therefore

have a wider (size) range of prey species in their diet. Since more diverse forests are expected to

contain larger insects, an similar shift in prey consumption is expected under the assumption of orb

web spider diet to be primarily determined by prey availability rather than prey selection.

Materials and methods

Study site

This study was conducted within the TREEWEB research platform (www.treedivbelgium.ugent.be/pl_treeweb.html) situated in the fragmented landscape of northern Belgium (50.899°N, 3.946°E – 50.998°N, 3.584°E). This platform consists of 53 research plots of 30 x 30 m². All have a similar land-use history (forest since at least 1850), management (no forest management in the last decade) and soil (dry sandy-loam). *Quercus robur* L. (pedunculate oak), *Quercus rubra* L. (red oak) and *Fagus sylvatica* L. (common beech) are the focal tree species in these forests. Plots of the three monocultures and all possible species mixtures (7 different stand compositions in total) were replicated 6 to 8 times along a fragmentation gradient. Edge distance (ranging from 7.0 to 215.5 m) was used as a proxy for edge effects. Edge distance was not correlated to tree diversity, neither did it differ between tree species combinations. Tree diversity was calculated by taking the exponent of the Shannon diversity index. The Shannon diversity index was calculated using the basal stem area of the tree species per plot. For more information on the setup of the study plots see (De Groote *et al.*, 2017).

Study species

We sampled common orb-weaver spiders (*Araneus diadematus* Clerk, 1757) for this study, as they are abundant in the study area and present in all of our study plots. We collected, if possible, 20 adult female *A. diadematus* in each plot from the 29[th] of August till 8[th] of September 2016. The spiders were taken from their webs, which were located at breast height. Collected spiders were immediately killed and stored in 99.6% alcohol. In some plots we could not collect 20 spiders, even after multiple visits. Spider size was taken by measuring the maximum width of the cephalothorax (i.e. carapace) under a stereomicroscope using a calibrated eyepiece graticule. Cephalothorax or

carapace width has been a common used proxy for body size in mature spiders (Hagstrum, 1971;

Greenstone, Morgan & Hultsch, 1985).

Molecular analysis

To establish the diet of the spiders, we used a proven metabarcoding protocol for spiders and other

invertebrate predators (Vesterinen, Lilley, Laine & Wahlberg 2013; Kaunisto, Roslin, Sääksjärvi &

Vesterinen, 2017; Eitzinger et al., 2019). Shortly, we extracted DNA from the spiders' abdomen using

NucleoSpin ® Tissue kit (cat. nr. 740952.250, Germany). To amplify mitochondrial COI gene, we used

primers ZBJ-ArtF1c and ZBJ-ArtR2c from (Zeale, Butlin, Barker, Lees & Jones, 2011). As these primers

also amplify the spiders themselves, we designed a blocking primer to decrease predator

amplification in favour of prey amplification (Vestheim & Jarman, 2008). To prepare the blocking

primer, we first downloaded all unique *A. diadematus* sequences from BOLD and GenBank and

aligned them with multiple potential prey sequences using Geneious (Kearse et al., 2012). Then we

designed three primer sequences that overlapped the reverse primer ZBJ-ArtR2c and that were

specific for *A. diadematus* (zero mismatches) but that did not match to any potential prey. The

blocking primers were tested using primer BLAST (Koressaar & Remm, 2007; Untergasser et al., 2012;

Ye, Coulouris, Zaretskaya, Cutcutache, Rozen & Madden, 2012). The best candidate (that did not bind

to anything in the database except *A. diadematus*) was chosen. This primer sequence was ordered

with C3 spacer modification at the 3' end (Aradia-R-blk-C3: 5'- CCA AAT CCC CCA ATT AAA ATA GGT

ATA-C3 spacer -3'). PCR conditions and library preparation followed (Kaunisto et al., 2017) and

(Vesterinen, Lilley, Puisto & Blomberg, 2018). Shortly, the first PCR included locus-specific linker-

tagged primers, and in this stage the target gene was amplified. In the second PCR, we introduced

linker-tagged indexed (different index in forward and reverse primers) adapters that were

compatible with Illumina platforms and perfectly matched the linker-tags in the first initial PCR. To

minimize the risk of contamination, all the extraction steps were carried out in carefully cleaned lab

space, using purified pipettes with filter tips. All the extraction batches included negative controls to

account for contamination issues. Washing the spiders several times in 99.6% ethanol during the

collection, storage and preparation for extraction process was deemed to be appropriate for

sterilization. Negative controls containing all but template DNA were included in each PCR assay. PCR

products were never introduced to the pre-PCR space. All the uniquely dual-indexed reactions were

pooled and purified using SPRI beads as in (Vesterinen *et al.*, 2016). The pool in this study was

combined with a similarly prepared pool (from a vertebrate dietary analysis) to increase nucleotide

diversity in the sequencing and to lower costs per project . Sequencing was performed by Macrogen

Korea (Macrogen Inc., Seoul, Rep. of Korea) using HiSeq4000 with TruSeq 3000 4000 SBS Kit v3

chemistry and 151 bp paired-end read length following HiSeq 3000 4000 System User Guide

(Document #15066496 v04 HCS 3.3.52).

After sequencing, the reads separated by each original sample were uploaded on CSC servers

(IT Center for Science, www.csc.fi) for trimming and further analysis. Trimming and quality control of

the sequences were carried out as in (Vesterinen *et al.*, 2018). Briefly, paired-end reads were

merged, trimmed, and collapsed using 64-bit software VSEARCH (Rognes ,Flouri, Nichols, Quince &

Mahé, 2016). For chimera-filtering, denoising, and clustering into ZOTUs ('zero-radius OTU'), we used

32-bit USEARCH (Edgar, 2010; Edgar & Flyvbjerg, 2015). Before collapsing, primers were removed

using software Cutadapt (Martin, 2011). Then, ZOTUs were mapped back to the original trimmed

reads using VSEARCH, and finally ZOTUs were assigned to prey taxa as explained below.

Data preparation

We summed the presence or absence of each prey taxon in each sample to end up with a frequency

of occurrence (FOO) for each prey taxon. Additionally, all the frequencies were scaled to per cent of

occurrence as explained in (Deagle *et al.*, 2019), creating a modified frequency of occurrence (MFO).

We identified prey to the species level, where possible. The ZOTUs were initially identified using local

BLAST against all COI sequences downloaded from BOLD and GenBank (Altschul *et al.*, 1990;

Ratnasingham & Hebert, 2007). When species name was not available but match to the database

was high, we used BIN codes from BOLD (Ratnasingham & Hebert, 2013). For details of the ZOTUs

see Supplementary materials 1. To visualize the trophic interactions structures resolved by the

molecular data, we used package BIPARTITE (Dormann, Fruend, Bluetghen & Gruber, 2008; Dormann,

Grubver & Fruend 2009) implemented in program R (R Core Team, 2018). Semi-quantitative

foodwebs were constructed using per cent of occurrence as explained above.

For further analysis, the cut-off threshold per ZOTU for the number of reads was set 0.05% of

the total number of reads per spider, with a minimum cut-off threshold of 10 reads. A first

multivariate analysis was performed to explore the variation in prey composition within the spider

diet. The variation in the prey ZOTU composition within the diet of individual spiders was analysed as

a function of tree species combination, edge distance and spider size. A distance-based redundancy

analysis (Euclidean distance) using the CAPSCALE function from the RDA package (Guo et al., 2018) was

applied. We performed an analysis of variance on the distance-based redundancy analysis with 1000

permutations (permanova) to quantify the variation in prey species composition explained by the

different variables. Taxonomic units (prey species) were treated as binary data (absence or presence)

as the data collection and chosen set-up using the metabarcoding technique does not allow a more

quantitative approach. We detected DNA from potential parasitoids (e.g. Tachinid flies, Ichneumonid

wasps) as well. These records were rare, and can be equally part of the diet. The metabarcoding

does, in the same vein, not allow separating potential DNA from species consumed by predators that

are here predated by spiders (e.g., Vespidae, Staphylidae). If some bias is created by these, we don't

expect this to be associated with specific environments or size classes.

Response variables

Per individual spider with known size, four diet-related response variables were calculated.

Prey richness was taken to be the number of ZOTUs in the diet of each spider. For every single

assigned prey ZOTU, its size (body length) was taken from literature (Supplementary materials 1).

Prey size was taken to be the average prey size of taxonomic units preyed by each spider. These two

response variables allow us to answer the four specific hypothesis. In addition, analyses were

performed with response variables to unearth possible explanations for the found patterns.

Taxonomic units with a body length over 1 cm are considered to be of highest gain (Venner & Casas,

2005), and could help us understand patterns arising. Prey richness of large prey was the number of

taxonomic units with a body length of over 1 cm present in the diet of each spider. Prey size of large

prey was the average prey size of taxonomic units larger than 1 cm present in the diet of each spider.

Additional to the four individual-level response variables, we additionally calculated the Sørensen

index (within-plot turnover of species composition in the diet), coefficient of variation (CV) for prey

richness and CV for prey size were calculated at plot level.

Statistics and model selection

Four models were applied to the response variables to explore different aspects of the data.

M_{div} was the first model in which we tested for the effects of spider size, edge distance, tree diversity

and the interacting effects of edge distance and tree diversity. The three other models were a set of

models to compare to each other in order to understand the effects of spider size, edge distance and

tree species composition (Kirwan *et al.*, 2009). The null model (M_{null}) includes only spider size and

edge distance and assumes no effect of tree species composition. The additive model (M_{add}) includes,

besides spider size and edge distance the relative basal area of each of the three focal tree species,

and the intercept was forced through zero. This model assumes that tree species exert only additive

identity effects. The pair-wise interaction model (M_{pair}) includes additionally the pair-wise

interactions between the relative basal areas of the focal tree species. This model assumes not only

additive effects, but also interacting effects between the tree species. The three composition models

(M_{null}, M_{add} and M_{pair}) were compared to each other to understand in which way the tree species

composition impacted on the diet-related response variables. The model with the lowest AICc

(obtained using the AICcMODAVG package by (Mazerolle, 2017) was considered the best. Spider size,

edge distance and tree diversity were scaled around their mean in all models (Schielzeth, 2010).

Spider size was excluded as an explanatory variable from the models applied to spider size itself. For

the plot-level mean response variables, an average plot-level spider size was included as a variable,

instead of individual-level spider size.

All analyses were performed in R, version 3.5.1 (R Core Team, 2018). All models, except the

models with Sørensen index and coefficients of variation as a response variables, included plot ID as

a random factor. This accounts for our data structure and the potential effect of plot ID. Models with

overall prey size as response variable had a negative binomial distribution (log-link) with a variance

increasing quadratically to the mean, applied through the GLMMTMB package (Brooks *et al.*, 2017).

Models with the overall prey richness as response variable had a negative binomial distribution with

constant variance using glm. Models with the richness of prey larger than 1 cm as response variable

had a Poisson distribution. All other models had Gaussian distributions. All models are two-tailed and

have no effect of the explanatory variables as H_0.

Results

General results

A total of 340 distinct prey species from 85 families in 8 orders and two classes were detected (Figure 1). The HiSeq 4000 sequencing yielded 265 871 470 paired-end reads. After assigning these reads to unique dual-indexes used in this study, and after trimming and filtering, we ended up with 5765096 prey reads that could be mapped to the original samples. A total of 983 spiders were included in the molecular analysis, and 857 (87.2%) of these provided prey data after bioinformatic filtering including 298 prey species and were included in the subsequent analysis. The highest observed prey species richness within a single spider sample was 15 prey ZOTUs. The most frequent prey detected was *Phaonia pallida* (N = 357), a forest-living muscid fly. The average prey size was 7.5 mm (SD ± 4.1), with only 39 prey species or genera larger than 1 cm. For a full list of prey taxa, and size data, see Supplementary materials 1.

Spider cephalothorax width ranged from 2.6 to 5.31 mm (average of 3.57 mm). Spider size decreased further from the edge; tree species identity had additive effects on spider size (Table 1-3, Figure 2). Spider size was not impacted by tree diversity or the interaction between tree diversity and edge distance, but size was largest in monocultures of *Q. robur* and smallest in monocultures of *F. sylvatica* (Table 1-3, Figure 2).

Diet composition

The composition of prey species in spider diets is highly variable (Figure 3). Although the constrained components of the ordination only explained 2,2% of the variation in composition, spider size (PERMANOVA, F_{pseudo} = 4.41, p = 0.001) and tree species combination (PERMANOVA, F_{pseudo} = 2.65, p = 0.001) strongly influenced prey species composition. However, all tree species compositions overlapped and showed large variation (Figure 3). Edge distance had no effect (PERMANOVA, F_{pseudo} =

1.28, p = 0.127). Compositional similarity of spider diet as measured by the Sørensen Index was not

correlated to edge distance, tree diversity or tree species composition (Table 1, 3).

Prey richness & size

The models including tree diversity (M_{div}) revealed that edge distance, tree diversity and the

interaction between them had no impact on the four diet-related response variables: prey richness,

prey size, richness of prey > 1cm and size of prey > 1cm (Table 1). In the best fitting composition

models (M_{null}, M_{add}, M_{pair}) edge distance had no impact on the diet-related response variables either

(Table 3). Spider size had an impact on the overall prey richness and prey size, only marginally on the

size of prey larger than 1 cm, and no impact on the richness of prey larger than 1 cm (Table 1). A

lower prey richness was detected in large spiders (Figure 4), although prey were larger in these

spiders. The effect of identity of the tree species and their relative contribution to prey richness was

absent, as M_{null} was the best fitting model (Table 2). For prey size there were additive effects of tree

identity (Table 2). Spiders in monocultures of *F. sylvatica* consumed the largest prey species, whilst

spiders from *Q. robur* and *Q. rubra* monocultures had very similar sized prey (Table 3, Figure 5).

Spiders inhabiting mixtures consumed prey which were the average size of the prey in the

monoculture values of tree species included in the tree species composition.

Richness of prey with the highest gain (prey larger than 1 cm) showed no effects of tree species

identity; size of prey with the highest gain (prey larger than 1 cm) did show effects of tree species

identity (Table 1-3). These large prey were proportionally most abundant in the diet of spiders from

F. sylvatica monocultures (Figure 6A). Prey body size was lowest in *F. sylvatica* - *Q. robur* mixtures,

relative to their anticipated size in the respective monocultures (Table 3, Figure 6B).

Patch-level individual variation in prey consumption.

The coefficient of variation (CV) for prey richness revealed that across tree diversity, tree species

composition, edge distance and spider size, the level of specialization was the same, as none of the

explanatory variables were significant (Table 1), and M_{null} was the best fitting composition model

(Table 2). The CV for prey size revealed that plots with on average larger spiders were more

consistent in their consumed prey size, independent of tree diversity or edge distance (Table 1).

Spiders in monocultures of *Q. robur* were least consistent in the consumed prey size, while those in

F. sylvatica monoculture were most consistent (Table 3, Figure 7). Spiders inhabiting mixtures

showed a level of consistency in prey size that was the average of the monoculture values of tree

species included in the tree species composition (Table 1; Figure 7).

Discussion

Tree diversity is an important driver of biodiversity in smaller forest fragments in Flanders (Hertzog et al. 2020), but different taxa react in different ways to the specific environmental gradients (Perring et al. 2021). We here use gut-content metabarcoding to show how changes in tree diversity, identity and edge effects affect an important functional biodiversity component, namely trophic interactions between an orb-web spider and its prey. Tree species composition, rather than tree diversity, impacted diet. More specifically, spiders consistently consumed larger prey in monocultures of *F. sylvatica* than in other tree species compositions. Contrary to expectations, we found prey species richness in the spider diet to decrease with spider size, but the proportion of large prey items to increase. Since spiders were largest in monocultures of *Q. robur* and smallest in monocultures of *F. sylvatica,* but also larger closer to the forest edge, size-mediated interactions are prevalent. Plots inhabited by on average larger spiders showed lower individual-variation in prey (size) consumption

Unlike our expectation that plant diversity impacts insect diversity and trophic interactions (Price, 2002; Scherber *et al.*, 2010; Rzanny *et al.*, 2013), we did not find any effect of tree diversity on the richness or size of prey in the diet. The does not imply that plant diversity cannot have an impact on the diet of other predators. In one of the rare studies that focussed on trophic interactions of a generalist predator in relation to plant diversity, it was shown that the richness of consumed prey in carabid beetles did increase along an experimental grassland diversity gradient (Tiede *et al.*, 2016). The general expectation that the diet of a predator contains more prey in prey rich habitat, is based on ecological opportunism (Bison *et al.*, 2015). Essentially, a generalist predator's diet would reflect the diversity of prey available. However, this is contradicted by the idea that the availability of more prey species allows intraspecific specialization on species (Staudacher *et al.*, 2018). We found no support for increased specialization in more diverse forest, as the coefficient of variation in prey richness did not depend on tree diversity. It should, however, be made clear that the occurrence of

ecological opportunism can neither be confirmed nor dismissed as both processes act at the

individual level, and may be levelled out at the population-level. In line with earlier research finding

mixtures between *Q. robur* and *F. sylvatica* to harbour the highest (arthropod) diversity (Hertzog et

al. 2020), we found this two-species mixture to held a different composition of species in the spiders'

diet, but not richness, compared to the other tree species composition. However, if spider diet did

indeed reflect the prey availability, we would expect spider diet in forests with *Q. robur* to have the

more species and being composed of larger prey, since *Q. robur* is known to harbour a more diverse

arthropod community (Southwood, Wint, Kennedy & Greenwood, 2004). This could not be

demonstrated, but we need to emphasise that it remains difficult to confidently conclude whether

our findings are pure reflections of the prey availability without its effective quantification, or

whether some prey selection is prevalent. Indeed, larger spiders tended to be more selective in

their foraging by the consumption of fewer, but larger prey species. Consumption of larger prey is

likely needed to achieve their higher energy requirement (Brown *et al.*, 2004). We can, however, not

attribute the general prey size increases to the selection of the largest most energy efficient prey, as

prey larger than one cm did not show a clear relationship with spider size. More-over, our approach

does neither allow more quantitative analyses on the relative abundances, consumed biomasses or

intraspecific variation in prey size. Prey sizes could only be derived from literature, and while caught

prey are adult active flying stages, intraspecific variation may be substantial. Irrespective of spider

size, spiders also consumed consistently large prey, and more and larger prey of high gain (> 1cm) in

monocultures of *F. sylvatica* than in other tree species composition. Interestingly, monocultures of *F.*

sylvatica also hold the smallest spiders.

We found no changes in prey richness or composition in relation to edge proximity. A possible

pattern could, however, be masked by the occurrence of larger spiders in proximity of the forest

edge. Neither larger spiders close to the edge, nor the absence of larger prey species consumed close

to the edge fit the expectation that the warmer forest edges could favour smaller arthropods

(Atkinson & Sibly, 1997; Kingsolver & Huey, 2008). Wind is stronger in the forest edges (Schmidt *et*

al., 2017). Wind damages webs, which reduces the foraging efficiency and enforces costly web

repairs (Tew, Adamson & Hesselberg, 2015). This may select for larger spiders with higher silk

production (Vollrath, 1999). It is possible that within this study a stronger effect is overlooked, as

edge effects for both biotic and abiotic gradients are generally observed in first few meters from the

edge (Murcia, 1995; Schmidt, Jochheim, Kersebaum & Lischeid, 2017; De Smedt *et al.*, 2018).

In conclusion, we show that an interception predator like the orb web spider *A. diadematus* is not

only generalist in prey capture but also prey consumption. At the individual level, however, the prey

spectrum of this generalist predator is linked to body size, with larger spiders foraging on a smaller

number of consistently larger prey species. Contrary to expectations from earlier research on insect

diversity in forest fragments differing in tree species composition, prey species richness and size are

only marginally explained by tree species composition. Edges effects on trophic relations were only

indirectly prevalent through differences in spider size.

Acknowledgements

We thank the research group of Tomas Roslin for providing the workspace for performing the

molecular analysis, all the forest owners for their permission to conduct our sampling in their forest,

Daan Dekeukeleire for many discussions and Pieter Vantieghem for his field work assistance. This

study was financially supported by Concerted Research Actions – Special Research Fund – Ghent

University for the UGent GOA project Scaling up Functional Biodiversity Research: from Individuals to

Landscapes and Back (TREEWEB) and FWO project G080221N.

Data accessibility Statement

All data on spider size, plot composition and DNA sequence assemblies are available on DRYAD:

https://doi.org/10.5061/dryad.7sqv9s4s3

Author contributions

IMSL, AM, KV, LL, DB designed the study, EJV assisted IMSL with the molecular analysis, LRH, DB

assisted IMSL with the data-analysis. All authors contributed to the writing of the manuscript.

Conflict of interests

The authors declare no competing interests.

Table 1 Summary of results for all M_{div} models. Except for the Sørensen index, Plot ID was added as a random factor in the models, but is not shown here. CV stands for coefficient of variation. (. p < 0.100; * p < 0.050, ** p < 0.010, *** p < 0.001)

		estimate	s.e.	z-value	p-value	
Spider size	Edge distance	-0.08	0.03	-2.74	0.009	**
	Tree diversity	0.01	0.03	0.38	0.705	
	Interaction	-0.003	0.03	-0.08	0.937	
Sørensen index	Edge distance	0.01	0.02	0.65	0.516	
	Tree diversity	0.01	0.02	0.48	0.635	
	Interaction	-0.01	0.02	-0.27	0.791	
Prey richness	Edge distance	-0.03	0.05	-0.60	0.550	
	Tree diversity	0.04	0.05	0.85	0.393	
	Spider size	-0.07	0.03	-2.48	0.013	*
	Interaction	-0.01	0.05	-0.12	0.908	
Prey size	Edge distance	-0.01	0.03	-0.21	0.808	
	Tree diversity	-0.02	0.03	-0.56	0.574	
	Spider size	0.06	0.02	3.37	<0.001	***
	Interaction	-0.03	0.03	-1.01	0.311	

		estimate	s.e.	z-value	p-value	
Prey richness > 1cm	Edge distance	-0.02	0.08	-0.28	0.777	
	Tree diversity	0.06	0.08	0.70	0.487	
	Spider size	0.06	0.05	1.27	0.204	
	Interaction	-0.01	0.09	-0.14	0.891	
Prey size > 1 cm	Edge distance	0.19	0.31	0.61	0.546	
	Tree diversity	0.03	0.31	0.08	0.933	
	Spider size	-0.33	0.19	-1.76	0.080	.
	Interaction	0.001	0.34	0.01	0.996	
CV of prey richness	Edge distance	-0.02	0.03	-0.51	0.610	
	Tree diversity	-0.01	0.03	-0.26	0.797	
	Spider size	-0.01	0.03	-0.17	0.869	
	Interaction	-0.01	0.04	-0.38	0.705	
CV of prey size	Edge distance	-0.02	0.02	-1.01	0.316	
	Tree diversity	0.02	0.02	0.80	0.429	
	Spider size	-0.02	0.02	-0.84	0.406	
	Interaction	-0.01	0.02	-0.29	0.773	

404   **Table 2** Comparison of the tree species composition models for all response variables. Best fitting models based on AICc and parsimony are bold. CV stands for coefficient of

405   variation.

	AICc		
	M_{null}	M_{add}	M_{pair}
Spider size	1169.26	**1164.76**	**1164.70**
Sørensen index	**-72.46**	-68.54	-62.71
Prey richness	**4169.50**	4173.21	4175.28
Prey size	4521.33	**4512.74**	4517.46
CV of prey richness	**-0.94**	3.38	6.37
CV op prey size	-58.96	**-61.93**	-56.76
Prey richness > 1cm	**1866.22**	**1864.35**	**1865.08**
Prey size > 1cm	2173.30	2173.76	**2161.18**

Table 3 Summary of results for the best fitting models which test for tree species composition effects. CV stands for coefficient of variation. (. p < 0.100; * p < 0.050, ** p < 0.010, *** p < 0.001)

		estimate	s.e.	z-value/t-value	p-value	
Spider size	Edge distance	-0.06	0.03	-2.27	0.028	*
	Rel. *F. sylvatica*	3.41	0.05	65.58	<0.001	***
	Rel. *Q. robur*	3.75	0.05	70.35	<0.001	***
	Rel. *Q. rubra*	3.55	0.05	69.38	<0.001	***
Sørensen index	Edge distance	0.01	0.02	0.72	0.475	
Prey richness	Edge distance	-0.03	0.05	-0.58	0.566	
	Spider size	-0.06	0.03	-2.47	0.014	*
Prey size	Edge distance	-0.01	0.03	-0.34	0.737	
	Spider size	0.07	0.02	3.80	<0.001	***
	Rel. *F. sylvatica*	2.17	0.05	41.68	<0.001	***
	Rel. *Q. robur*	1.91	0.05	34.92	<0.001	***
	Rel. *Q. rubra*	1.90	0.05	36.63	<0.001	***

		estimate	s.e.	z-value/t-value	p-value	
Prey richness > 1cm	Edge distance	-0.02	0.08	-0.26	0.795	
	Spider size	0.06	0.05	1.28	0.199	
Prey size > 1cm	Edge distance	0.17	0.31	0.57	0.570	
	Spider size	-0.29	0.19	-1.52	0.130	
	Rel. *F. sylvatica*	19.75	0.70	28.08	<0.001	***
	Rel. *Q. robur*	19.08	0.75	25.56	<0.001	***
	Rel. *Q. rubra*	18.56	0.80	23.07	<0.001	***
	Rel. *F. sylvatica* x Rel. *Q. robur*	-6.75	3.54	-1.91	0.065	.
	Rel. *F. sylvatica* x Rel. *Q. rubra*	3.87	3.79	1.02	0.313	
	Rel. *Q. robur* x Rel. *Q. rubra*	3.40	3.59	0.95	0.350	
CV of prey richness	Edge distance	-0.02	0.03	-0.49	0.630	
	Spider size	-0.01	0.03	-0.18	0.858	
CV of prey size	Edge distance	-0.02	0.02	-1.25	0.266	
	Spider size	-0.04	0.02	-2.04	0.047	*
	Rel. *F. sylvatica*	0.20	0.04	5.02	<0.001	***
	Rel. *Q. robur*	0.38	0.04	8.89	<0.001	***
	Rel. *Q. rubra*	0.32	0.04	8.80	<0.001	***

411   **Figures**

412   **Figure 1** Visual representation of the taxonomic distribution and quantified strength of trophic links from *Araneus diadematus* to their prey. The blocks in the lower row

413   represent prey species. A line connecting the predator with a prey represents detected predation events, and the thickness of the line represents the modified frequency of

414   occurrences   (MFO)   of   each   predation   record.   See   the   "Data   analysis"   in   the   main   text   for   details   on   the   MFO.

Figure 2 Relationship between the spider's cephalothorax size (mm) and distance to the forest edge (m) per tree species composition. Data points are the individual spiders (N = 983). Lines with 95% CI are the estimated slope based on model M_{add}. Colours refer to the tree species composition. There is a negative relationship between spider size and edge distance, and additive tree species composition effects, where mixtures are averages of each individual monoculture contributing to the mixture.

421
Figure 3 Distance-based redundancy analysis on the prey composition of individual spiders. Colours refer to the

different tree species compositions in which the spiders were sampled. Ellipses correspond with the standard

deviation of each tree species composition centroid. Size refers to the spider size and Edge to edge distance.

The constrained components of the ordination (spider size, edge distance and tree species composition)

explained 2.2% of all variation in diet composition.

Figure 4 Relationship between the number of prey species in the diet of a spider and its cephalothorax size in
mm. Data points are the number of prey species per individual spider (n = 983). Red line with 95% CI is the
estimated slope based on model M_{div}

Figure 5 Relationship between the average prey size (mm) of the species caught and the spider's cephalothorax

size (mm) per tree species composition. Data points are the individual spiders (N = 983). Lines with 95% CI are

the estimated slopes based on model M_{add}. For estimation, edge distance was taken to be the overall average.

Colours refer to the tree species composition. There is a positive relationship between average prey size and

spider, and additive tree species composition effects, in which monocultures of *F. sylvatica* catch relatively

larger prey species than the other monocultures. In mixtures, the prey size are averages of each individual

monoculture contributing to the mixture. The lower part of the graph shows density plots of the spider size

distribution within each tree species composition.

Figure 6 Tree species composition effect on (A) the number of prey species larger than 1 cm and (B) the average prey size of prey larger than 1 cm. Open data points are the averages per plot. Red point ranges are the estimated values based on the best parsimonious model. Edge distance and spider size were taken to be the overall average. For (A) there are additive effects of the monocultures in mixtures, averaging out the number of prey species. For (B) there are pairwise interaction effects in combining *F. sylvatica* with *Q. robur* results in (marginally) lower prey size than expected based on the monocultures.

Figure 7. Relationship between the coefficient of variation (CV) for prey size per plot and the mean spider's cephalothorax size (mm) per plot. Data points are the individual plots (N = 53). Lines with 95% CI are the estimated slopes based on model M_{add}. For estimation, edge distance was taken to be the overall average. Colours refer to the tree species composition. In plots with larger spiders, there is a stronger stability in prey size (lower CV). Additive tree species composition effects are present. The strongest stability in prey size is found in monocultures of *F. sylvatica*, and the lowest stability in monoculture of *Q. robur*. In mixtures, the CV is an average of the CV of each individual monoculture contributing to the mixture.

References

Allen, C.R., Garmestani, A.S., Havlicek, T.D., Marquet, P.A., Peterson, G.D., Restrepo, C., *et al.* (2006) Patterns in body mass distributions: sifting among alternative hypotheses. *Ecology Letters*, 9, 630–643.

Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. (1990) Basic local alignment search

tool. *Journal of Molecular Biology*, 215, 403–410.

Araújo, M.S., Bolnick, D.I. & Layman, C.A. (2011) The ecological causes of individual specialisation.

Ecology Letters, 14, 948–958.

Atkinson, D. & Sibly, R.M. (1997) Why are organisms usually bigger in colder environments? Making

sense of a life history puzzle. *Trends in ecology & evolution*, 12, 235–239.

Bison, M., Ibanez, S., Redjadj, C., Boyer, F., Coissac, E., Miquel, C., *et al.* (2015) Upscaling the niche

variation hypothesis from the intra - to the inter - specific level. *Oecologia*, 179, 835–842.

Bolnick, D.I., Amarasekare, P., Araújo, M.S., Bürger, R., Levine, J.M., Novak, M., *et al.* (2011) Why

intraspecific trait variation matters in community ecology. *Trends in ecology & evolution*, 26, 183–

192.

Bolnick, D.I., Svanbäck, R., Fordyce, J.A., Yang, L.H., Davis, J.M., Hulsey, C.D., *et al.* (2003) The Ecology

of Individuals: Incidence and Implications of Individual Specialization. *The American naturalist*, 161,

1–28.

Bonte, D., Lanckacker, K., Wiersma, E. & Lens, L. (2008) Web building flexibility of an orb-web spider

in a heterogeneous agricultural landscape. *Ecography*.

Brooks, D.M. (2012) Birds caught in spider webs: a synthesis of patterns. *The Wilson Journal of

Ornithology*, 124, 345–353.

Brooks, M.E., Kristensen, K., Benthem, K.J. van, Magnusson, A., Berg, C.W., Nielse, A., *et al.* (2017)

glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed

Modeling. *The R Journal*, 9, 378–400.

Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. (2004) Toward a metabolic theory of

ecology. *Ecology*, 85, 1771–1789.

Cagnolo, L., Valladares, G., Salvo, A., Cabido, M. & Zak, M. (2009) Habitat fragmentation and species

loss across three interacting trophic levels: effects of life-history and food-web traits. *Conservation*

biology, 23, 1167–1175.

Dahirel, M., Dierick, J., Cock, M. De & Bonte, D. (2017) Intraspecific variation shapes community-level

behavioral responses to urbanization in spiders. *Ecology*, 98, 2379–2390.

Dahirel, M., De Cock, M. , Vantieghem, P. & Bonte, D. (2019). Urbanization-driven changes in web-

building and body size in an orb-web spider. *Journal of Animal Ecology,* 88: 79-91.

Deagle, B.E., Thomas, A.C., McInnes, J.C., Clarke, L.J., Vesterinen, E.J., Chare, E.L., *et al.* (2019)

Counting with DNA in metabarcoding studies: How should we convert sequence reads to dietary

data? *Molecular Ecology*, 28, 391–406.Debinski, D.M. & Holt, R.D. (2000) A survey and overview of

habitat fragmentation experiments. *Conservation Biology*, 14, 342–355.

Diehl, E., Sereda, E., Wolters, V. & Birkhofer, K. (2013) Effects of predator specialization, host plant

and climate on biological control of aphids by natural enemies: a meta-analysis. *Journal of Applied*

Ecology, 50, 262–270.

Dormann, C.F., Fruend, J., Bluethgen, N. & Gruber, B. (2009) Indices, graphs and null models:

analyzing bipartite ecological networks. *The Open Ecology Journal*, 2, 7–24.

Dormann, C.F., Gruber, B. & Fruend, J. (2008) Introducing the bipartite Package: Analysing Ecological

Networks. *R news*, 8, 8–11.

Edgar, R.C. (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*, 26,

2460–2461.

Edgar, R.C. & Flyvbjerg, H. (2015) Sequence analysis Error filtering, pair assembly and error correction

for next-generation sequencing reads. *Bioinf*, 31, 3476–3482.

Eitzinger, B., Abrego, N., Gravel, D., Huotari, T., Vesterinen, E.J. & Roslin, T. (2019) Assessing changes

in arthropod predator – prey interactions through DNA - based gut content analysis — variable

environment , stable diet. *Molecular Ecology*, 28, 266–280.

Eitzinger, B., Roslin, T., Vesterinen, E. J., Robinson, S. I., & O'Gorman, E. J. (2021). Temperature

affects both the Grinnellian and Eltonian dimensions of ecological niches – A tale of two Arctic wolf

spiders. *Basic and Applied Ecology*, *50*, 132–143.

Greenstone, M.H., Morgan, C.E. & Hultsch, A.L. (1985) Ballooning methodology: equations for

estimating masses of sticky-trapped spiders. *Journal of Arachnology*, 13.

Groote, S.R.E. De, Schrojenstein Lantman, I.M. van, Sercu, B.K., Dekeukeleire, D., Boonyarittichaikij,

R., Smith, H.K., *et al.* (2017) Tree species identity outweighs the effects of tree species diversity and

forest fragmentation on understorey diversity and composition. *Plant Ecology and Evolution*, 150,

229–239.

Gross, T., Rudolf, L., Levin, S.A. & Dieckmann, U. (2009) Generalized models reveal stabilizing factors

in food webs. *Science*, 235, 747–750.

Guo, Y., Hastie, T. & Tibshirani, R. (2018) rda: Shrunken Centroids Regularized Discriminant Analysis.

Haddad, N.M., Brudvig, L.A., Clobert, J., Davies, K.F., Gonzalez, A., Holt, R.D., *et al.* (2015) Habitat

fragmentation and its lasting impact on Earth's ecosystems. *Applied Ecology*, 1, e1500052.

Haddad, N.M., Crutsinger, G.M., Gross, K., Haarstad, J., Knops, J.M.H. & Tilman, D. (2009) Plant

species loss decreases arthropod diversity and shifts trophic structure. *Ecology Letters*, 12, 1029–

1039.

Hagstrum, D.W. (1971) Carapace width as a tool for evaluating the rate of development of spiders in

the laboratory and the field. *Annals of the Entomological Society of America*, 64, 757–760.

Hertzog, L.R., Meyer, S.T., Weisser, W.W. & Ebeling, A. (2016) Experimental Manipulation of

Grassland Plant Diversity Induces Complex Shifts in Aboveground Arthropod Diversity. *PLoS ONE*, 11,

e0148768.

Hertzog, L. R., Boonyarittichaikij, R., Dekeukeleire, D., de Groote, S. R. E., van Schrojenstein Lantman,

I. M., Sercu, B. K., Smith, H. K., de la Peña, E., Vandegehuchte, M. L., Bonte, D., Martel, A., Verheyen,

K., Lens, L., and Baeten, L. (2019). Forest fragmentation modulates effects of tree species richness

and composition on ecosystem multifunctionality. *Ecology* 100(4):e02653. [10.1002/ecy.2653](10.1002/ecy.2653)

Hillaert, J., Hovestadt, T., Vandegehuchte, M.L. & Bonte, D. (2018) Size-dependent movement

explains why bigger is better in fragmented landscapes. *Ecology and Evolution*, 8, 10754–10767.

Holt, R.D. (2002) Food webs in space: On the interplay of dynamic instability and spatial processes.

Ecological Research, 17, 261–273.

Hooper, D.U., Chapin, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., *et al.* (2005) Effects of

Biodiversity on Ecosystem Functioning: A Consensus of Current Knowledge. *Ecological Monographs*,

75, 3–35.

Janetos, A.C. (1982) Foraging Tactics of Two Guilds of Web-Spinning Spiders. *Behavioral Ecology and*

Sociobiology, 10, 19–27.

Kaunisto, K.M., Roslin, T., Sääksjärvi, I.E. & Vesterinen, E.J. (2017) Pellets of proof: First glimpse of the

dietary composition of adult odonates as revealed by metabarcoding of feces. *Ecology and Evolution*,

7, 8588–8598.

Kearse, M., Moir, R., Wilson, A., Stones-havas, S., Sturrock, S., Buxton, S., *et al.* (2012) Geneious

Basic: An integrated and extendable desktop software platform for the organization and analysis of

sequence data. *Bioinformatics*, 28, 1647–1649.

Keiser, C.N., Slyder, J.B., Carson, W.P. & Pruitt, J.N. (2015) Individual differences in predators but not

producers mediate the magnitude of a trophic cascade. *Arthropod-Plant Interactions*, 9, 225–232.

Kingsolver, J.G. & Huey, R.B. (2008) Size, temperature , and fitness: three rules. *Evolutionary Ecology*

Research, 10, 251–268.

Kirwan, L., Connolly, J., Finn, J.A., Brophy, C., Lüscher, A., Nyfeler, D., *et al.* (2009) Diversity–

interaction modeling: estimating contributions of species identities and interactions to ecosystem

function. *Ecology*, 90, 2032–2038.

Koressaar, T. & Remm, M. (2007) Enhancements and modifications of primer design program

Primer3. *Bioi*, 23, 1289–1291.

Krauss, J., Bommarco, R., Guardiola, M., Heikkinen, R.K., Helm, A., Kuussaari, M., *et al.* (2010) Habitat

fragmentation causes immediate and time- delayed biodiversity loss at different trophic levels.

Ecology Letters, 13, 597–605.

Landi, P., Minoarivelo, H.O., Brännström, A., Hui, C. & Dieckmann, U. (2018) Complexity and stability

of ecological networks: a review of the theory. *Population Ecology*, 60, 319–345.

Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J.P., Hector, A., *et al.* (2001) Biodiversity

and Ecosystem Functioning: Current Knowledge and Future Challenges. *Science*, 294, 804–808.

Martin, M. (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal, 17, 10–12.

Martinson, H.M. & Fagan, W. (2014) Trophic disruption: a meta-analysis of how habitat

fragmentation affects resource consumption in terrestrial arthropod systems. *Ecology Letters*, 17,

1178–1189.

Mazerolle, M.J. (2017) Model selection and multimodel inference based on (Q)AIC(c). R package

version 2.1-1 [WWW Document]. URL http://cran.r-project.org/package=AICcmodavg [accessed on

2017].

Morse, D.R., Lawton, J.H., Dodson, M.M. & Williamson, M.H. (1985) Fractal dimension of vegetation

and the distribution of arthropod body lengths. *Nature*, 314, 731–733.

Murcia, C. (1995) Edge effects in fragmented forests: implications for conservation. *Trends in ecology*

& evolution, 10, 58–62.

Nentwig, W. & Wissel, C. (1986) A Comparison of Prey Lengths among Spiders. *Oecologia*, 68, 595–

600.

Nyffeler, M. & Birkhofer, K. (2017) An estimated 400 – 800 million tons of prey are annually killed by

the global spider community. *The Science of Nature*, 140, 30.

O'Brien, M.J., Brezzi, M., Schuldt, A., Zhang, J.-Y., Ma, K., Schmid, B., *et al.* (2017) Tree diversity drives

diversity of arthropod herbivores , but successional stage mediates detritivores. *Ecology and*

Evolution, 7, 8753–8760.

O'Rorke, R., Lavery, S. & Jeffs, A. (2012) PCR enrichment techniques to identify the diet of predators.

Molecular Ecology Resources, 12, 5–17.

Perring, MP, Hertzog, LR, De Groote, SRE, et al. (2021). Overstorey composition shapes across-trophic

level community relationships in deciduous forest regardless of fragmentation context. *J Ecol.* 2021;

109: 1591– 1606. https://doi.org/10.1111/1365-2745.13580

Pompanon, F., Deagle, B.E., Symondson, W.O.C., Brown, D.S., Jarman, S.N. & Taberlet, P. (2012) Who

is eating what: diet assessment using next generation sequencing. *Molecular Ecology*, 21, 1931–

1950.

Post, D.M., Palkovacs, E.P., Schielke, E.G. & Dodson, S.I. (2008) Intraspecific Variation in a Predator

Affects Community Structure and Cascading Trophic Interactions. *Ecology*, 89, 2019–2032.

Price, P.W. (2002) Resource-driven terrestrial interaction webs. *Ecological Research*, 17, 241–247.

R Core Team. (2018) R: A language and environment for statistical comunting. *R Foundation for*

Statistical Computing, Vienna, Austria.

Rand, T.A., Tylianakis, J.M. & Tscharntke, T. (2006) Spillover edge effects: the dispersal of

agriculturally subsidized insect natural enemies into adjacent natural habitats. *Ecology Letters*, 9,

603–614.

Ratnasingham, S. & Hebert, P. (2007) BOLD: The Barcode of Life Data System

(www.barcodinglife.org). *Molecular Ecology Notes*, 7, 355–364.

Ratnasingham, S. & Hebert, P. (2013) A DNA-Based Registry for All Animal Species: The Barcode Index

Number (BIN) System. *PLoS ONE*, 8.

Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. (2016) VSEARCH⬚: a versatile open source

tool for metagenomics. *PeerJ*, 4, e2584.

Rzanny, M., Kuu, A. & Voigt, W. (2013) Bottom – up and top – down forces structuring consumer

communities in an experimental grassland. *Oikos*, 122, 967–976.

Scherber, C., Eisenhauer, N., Weisser, W.W., Schmid, B., Voigt, W., Fischer, M., *et al.* (2010) Bottom-

up effects of plant diversity on multitrophic interactions in a biodiversity experiment. *Nature*, 468,

553–556.

Scherber, C., Vockenhuber, E.A., Stark, A., Meyer, H. & Tscharntke, T. (2014) Effects of tree and herb

biodiversity on Diptera, a hyperdiverse insect order. *Oecologia*, 174, 1387–1400.

Schielzeth, H. (2010) Simple means to improve the interpretability of regression coefficients.

Methods in Ecology and Evolution, 1, 103–113.

Schmidt, M., Jochheim, H., Kersebaum, K. & Lischeid, G. (2017) Gradients of microclimate, carbon

and nitrogen in transition zones of fragmented landscapes – a review. *Agricultural and Forest*

Meteorology, 232, 659–671.

Schmidt, N.M., Mosbacher, J.B., Eitzinger, B., Vesterinen, E.J., Roslin, T. & Schmidt, N.M. (2018) High

resistance towards herbivore-induced habitat change in a high Arctic arthropod community. *Biology*

L, 14, 20180054.

Schneider, J.M. & Vollrath, F. (1998) The effect of prey type on the geometry of the capture web of

Araneus diadematus. *The Science of Nature*, 85, 391–394.

Sensenig, A., Agnarsson, I., Gondek, T.M. & Blackledge, T.A. (2010) Webs in vitro and in vivo: spiders

alter their orb-web spinning behavior in the laboratory. *Journal of Arachnology*, 38, 183–191.

Smedt, P. De, Baeten, L., Berg, M.P., Gallet-moron, E., Brunet, J., Cousins, S.A.O., *et al.* (2018)

Desiccation resistance determines distribution of woodlice along forest edge- to-interior gradients.

European Journal of Forest Research, 85, 1–3.

Southwood, T.R.E., Wint, G.R.W., Kennedy, C.E.J. & Greenwood, S.R. (2004) Seasonality, abundance,

species richness and specificity of the phytophagous guild of insects on oak (Quercus) canopies.

European Journal of Entomology, 101, 43–50.

Staudacher, K., Rubbmark, O.R., Birkhofer, K., Malsher, G., Sint, D., Jonsson, M., *et al.* (2018) Habitat

heterogeneity induces rapid changes in the feeding behaviour of generalist arthropod predators.

Functional Ecology, 23, 809–819.

Tew, E.R., Adamson, A. & Hesselberg, T. (2015) The web repair behaviour of an orb spider. *Animal*

Behaviour, 103, 137–146.

Thébault, E. & Loreau, M. (2005) Trophic Interactions and the Relationship between Species. *The*

American naturalist, 166, E95–E114.

Tiede, J., Wemheuer, B., Traugott, M., Daniel, R., Tscharntke, T., Ebeling, A., *et al.* (2016) Trophic and

Non-Trophic Interactions in a Biodiversity Experiment Assessed by Next- Generation Sequencing. *PloS*

ONE, 11, e0148781.

Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B.C., Remm, M., *et al.* (2012) Primer3 —

new capabilities and interfaces. *Nucleic Acids Research*, 40, 1–12.

Vehviläinen, H., Koricheva, J. & Ruohomäki, K. (2008) Effects of stand tree species composition and

diversity on abundance of predatory arthropods. *Oikos*, 117, 935–943.

Venner, S. & Casas, J. (2005) Spider webs designed for rare but life-saving catches. *Proceedings of the*

Royal Society Biological sciences, 272, 1587–1592.

Vesterinen, E.J., Lilley, T., Laine, V.N. & Wahlberg, N. (2013) Next Generation Sequencing of Fecal

DNA Reveals the Dietary Diversity of the Widespread Insectivorous Predator Daubenton' s Bat

(Myotis daubentonii) in Southwestern Finland. *PLoS ONE*, 8, 1–14.

Vesterinen, E.J., Lilley, T.M., Puisto, A.I.E. & Blomberg, A.S. (2018) Table for five, please: Dietary

partitioning in boreal bats. *Ecology and Evolution*, 8, 10914–10937.

Vesterinen, E.J., Ruokolainen, L., Wahlberg, N., Peña, C., Roslin, T., Laine, V.N., *et al.* (2016) What you

need is what you eat? Prey selection by the bat Myotis daubentonii. *Molecular Ecology*, 25, 1581–

1594.

Vestheim, H. & Jarman, S.N. (2008) Blocking primers to enhance PCR amplification of rare sequences

in mixed samples – a case study on prey DNA in Antarctic krill stomachs. *Frontiers in Zoology*, 11, 1–

11.

Vollrath, F. (1999) Biology of spider silk. *Internation Journal of Biological Macromolecules*, 24, 81–88.

Woodward, G. & Hildrew, A.G. (2002) Body-size determinants of niche overlap and intraguild

predation within a complex food web. *Journal of Animal Ecology*, 71, 1063–1074.

Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S. & Madden, T.L. (2012) Primer-BLAST: A

tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics*, 13, 134.

Zeale, M.R.K., Butlin, R.K., Barker, G.L.A., Lees, D.C. & Jones, G. (2011) Taxon-specific PCR for DNA

barcoding arthropod prey in bat faeces. *Molecular Ecology Resources*, 11, 236–244.