The impact of neural attentional filters on listening behaviour in a large cohort of ageing individuals

Sarah Tune[1,2]*,
Mohsen Alavash[1,2],
Lorenz Fiedler[1,2] §,
& Jonas Obleser[1,2] *

1 – Department of Psychology, University of Lübeck, 23562 Lübeck, Germany
2 – Center for Brain, Behavior, and Metabolism, University of Lübeck, 23562 Lübeck, Germany

Author correspondence:
Sarah Tune, Jonas Obleser
Department of Psychology
University of Lübeck
Maria-Goeppert-Str. 9a
23562 Lübeck
Email: sarah.tune@uni-luebeck.de; jonas.obleser@uni-luebeck.de

§ current address: Eriksholm Research Centre, Snekkersten, Denmark

Running title: Modelling neural filters and behavioural outcome in attentive listening

Number of words in Abstract: 159
Number of words in Introduction: 868
Number of words in Discussion: 2048

31 pages, 6 Figures, 0 Tables, Includes Supplemental Information

Keywords: EEG, alpha power lateralization, neural speech tracking, speech processing, auditory selective attention

Abstract

Successful listening crucially depends on intact attentional filters that separate relevant from irrelevant information. Research into their neurobiological implementation has focused on two potential auditory filter strategies: the lateralization of alpha power and selective neural speech tracking. However, the functional interplay of the two neural filter strategies and their potency to index listening success in an ageing population remains unclear. Using electroencephalography and a dual-talker task in a representative sample of listeners (N=155; age=39–80 years), we here demonstrate an often-missed link from single-trial behavioural outcomes back to trial-by-trial changes in neural attentional filtering. First, we observed preserved attentional–cue-driven modulation of both neural filters across chronological age and hearing levels. Second, neural filter states varied independently of one another, demonstrating complementary neurobiological solutions of spatial selective attention. Stronger neural speech tracking but not alpha lateralization boosted trial-to-trial behavioural performance. Our results highlight the translational potential of neural speech tracking as an individualized neural marker of adaptive listening behaviour.

Significance statement

Successful listening requires attentional filtering into behaviourally relevant and irrelevant sounds. Most previous studies have focused on one of two candidate neural filter strategies: the lateralization of alpha power or selective neural speech tracking. Closing the gap between hitherto separate lines of research, we used electroencephalography and a dual-talker task in a large sample of age-varying listeners. We directly probe the functional relevance of state- and trait-level changes in these neural filter strategies to listening success. We demonstrate the co-existence of largely independent neural filters that pose complementary neurobiological implementations of selective attention. Additionally, our results emphasize the utility of neural speech tracking over alpha lateralization as a potential neural marker of an individual's adaptive listening behaviour.

Introduction

Real-life listening is characterized by the concurrence of sound sources that compete for our attention[1]. Successful speech comprehension thus relies on the differentiation of relevant and irrelevant inputs. Here, the concept of neural attentional 'filters' serves as an important and pervasive algorithmic metaphor of how auditory attention is implemented at the neural level[2-4]. Neural attentional filters can be instantiated by different mechanistic principles and recent studies have predominantly focused on two potential but nonexclusive neural filter strategies originating from distinct research traditions:

From the visual domain stems an influential line of research that supports a role of alpha-band (~8–12 Hz) oscillatory activity in the implementation of controlled, top-down suppression of behaviourally-irrelevant information[5-8]. Importantly, across modalities, it was shown that spatial-attention tasks are neurally supported by a hemispheric lateralization of alpha power over occipital, parietal but also the respective sensory cortices[9-18]. This suggests that asymmetric alpha modulation could act as a filter mechanism by modulating sensory gain already in early processing stages.

In addition, a prominent line of research focuses on the role of low-frequency (1–8 Hz) neural activity in auditory and, broadly speaking, perisylvian cortex in the selective representation of speech input ("neural speech tracking"). Slow cortical dynamics temporally align with (or "track") auditory input signals to prioritize the neural representation of behaviourally-relevant sensory information[19-22] (see also refs.[23,24] for the neural tracking of contextual semantic information). In human speech comprehension, a key finding is the preferential neural tracking of attended compared to ignored speech in superior temporal brain areas close to auditory cortex[25-29].

However, with few exceptions[9], these two proposed neural auditory filter strategies have been studied independently of one another (but see refs.[30,31] for recent results on visual attention). Also, they have often been studied using tasks that are difficult to relate to natural, conversation-related listening situations[32,33].

We thus lack understanding whether or how modulations in lateralized alpha power and the neural tracking of attended versus ignored speech in wider auditory cortex interact in the service of successful listening behaviour. Moreover, few studies using more real-life listening and speech-tracking measures were able to explicitly address the functional relevance of the discussed neural filter strategies, that is, their potency to explain behavioural listening success[27,28].

As part of an ongoing large-scale project on the neural and cognitive mechanisms supporting adaptive listening behaviour in healthy ageing, the current study aims at closing these gaps by leveraging the statistical power and representativeness of our large, age-varying participant sample. We use a novel dichotic listening paradigm to enable a synergistic look at concurrent single-trial changes in lateralized alpha power and neural speech tracking.

More specifically, our linguistic variant of a classic Posner paradigm[34] emulates a challenging dual-talker listening situation in which speech comprehension is supported by two different listening cues[35,36]. Of particular interest for the present

scientific endeavour is the spatial-attention cue that guides auditory attention in space.
We additionally manipulated the semantic predictability of upcoming speech via a
semantic category cue. While the effects of the semantic cue are of secondary
importance for the present research questions, its manipulation still allows insights into
whether semantic predictability modulates the engagement of neural attentional filter
mechanisms, and how it affects listening success in a large cohort of middle-aged and
older adults. Previous research has shown that the sensory analysis of speech and, to a
lesser degree, the modulation of alpha power are influenced by the availability of
higher-order linguistic information[37-42].

Varying from trial to trial, both cues were presented either in an informative or
uninformative version. This manipulation allowed us to understand how concurrent
changes in the neural dynamics of selective attention and the resulting listening
behaviour are connected.

We focus on four main research questions (see Fig. 1). Note that in addressing
these, we model additional known influences on listening success: age, hearing loss, as
well as hemispheric asymmetries in speech processing due to the well-known right-ear
advantage[43,44].

Figure 1. Schematic illustration of the research questions addressed in the present study. The dichotic listening task manipulated the attentional focus and semantic predictability of upcoming input using two separate visual cues. We investigated whether informative cues would enhance behavioural performance (Q1). In line with (Q2), we also examined the degree to which a spatial (and semantic) cue modulated the two auditory neural measures of interest: neural speech tracking and lateralization of auditory alpha power. Finally, we assessed (Q3) the co-variation of neural measures, and (Q4) their potency in explaining behavioural performance. Furthermore, we investigated the impact of age, hearing loss, and probed ear on listening success and its underlying neural strategies.

First, informative listening cues should increase listening success: These cues allow the
listener to deploy auditory selective attention (compared to divided attention), and to
generate more specific (compared to only general) semantic predictions, respectively.

Second, we asked how the different cue–cue combinations would modulate the
two key neurobiological measures of selective attention—alpha lateralization and
neural speech tracking. We aimed to replicate previous findings of increased alpha
lateralization and a preferential tracking of the target compared to the distractor speech
signal under selective (compared to divided) spatial attention. At the same time, we
capitalized on the large, age-varying sample to quantify the hitherto contested
dependence of these neurobiological filters on participants' chronological age and
hearing loss[14,45-47].

Third, an important and often neglected research question pertains to a direct,
trial-by-trial relationship of these two candidate neural measures: Do changes in alpha

lateralization impact the degree to which attended and ignored speech signals are
neurally tracked by low-frequency cortical responses?
Our final research question is arguably the most relevant one for all translational
aspects of auditory attention; it has thus far only been answered indirectly when
deeming these neurobiological filter mechanisms "attentional": To what extent do
alpha lateralization and neural speech tracking allow us to explain behavioural
outcome—that is, listening success—at the individual level and in a single trial?

Results

We recorded and source-localized electroencephalography (EEG) signals in a large, age-
varying sample of healthy middle-aged and older adults (N=155; age=39–80 years, see
Supplementary Fig. 1) who performed a challenging dichotic listening task. In this
linguistic variant of a classic Posner paradigm[35], participants listened to two concurrent
five-word sentences spoken by the same female talker and were asked to identify the
final word in one of the two sentences. Sentence pairs were temporally aligned to the
onset of these task-relevant final words which led to slightly asynchronous sentence
onsets.
Importantly, sentence presentation was preceded by two visual cues. First, a
spatial-attention cue encouraged the use of either selective or divided attention by
providing informative or uninformative instructions about the to-be-attended, and thus
later probed, ear. The second cue indicated the semantic category that applied to both
final target words. The provided category could represent a general or specific level,
thus allowing for more or less precise prediction of the upcoming speech signal (Fig. 2a,
b). While this listening task does not tap into the most naturalistic forms of speech
comprehension, it still approximates a dual-talker listening situation to probe the neural
underpinnings of successful selective listening[35].
Using generalized linear mixed-effects models on single-trial data, we focus on
two key neurobiological instantiations of auditory attention: the lateralization of 8–12
Hz alpha power, emerging from auditory as well as parietal cortex, and the differential
neural tracking of attended versus ignored speech by slow (1–8 Hz) auditory cortical
responses. We investigate the how spatial cues, age, and hearing status modulate
behaviour and neural filters, whether neural filters operate independently, and to which
extent they influence selective listening success.

Figure 2. Experimental design and behavioural benefit from informative cues.
(a) Visualization of used 2×2 design[35]. Levels of spatial and semantic cue differed on a trial-by-trial basis. Note that the effects of the semantic cue were of secondary importance to the current analyses. Top row shows the informative [+] cue levels, bottom row the uninformative [−] cue levels.
(b) Schematic representation of the trial structure. Successive display of the two visual cues precedes the dichotic presentation of two sentences spoken by the same female talker. After sentence presentation, participants had to select the final word from four alternative words.
(c) Grand average and individual results of accuracy shown per cue-cue combination. Coloured dots are individual (N=155) trial-averages, black dots and vertical lines show group means ± bootstrapped 95 % confidence intervals (CI; left side). Box plots show median, 25th to 75th percentile data, and 1.5 interquartile range. Individual cue benefits displayed separately for the two cues (top: spatial cue, bottom: semantic cue). Black dots indicate individual trial-averages ± bootstrapped 95 % CI. Histograms show the distribution of the difference for informative vs. uninformative levels. OR: Odds ratio parameter estimate from generalized linear mixed-effects models (right side); *** = p < .001.
(d) As in (c) but shown for response speed; β: slope parameter estimate from general linear mixed-effects models.

Informative spatial cues improve listening success

For behavioural performance, we tested the impact of informative versus uninformative cues on listening success. Overall, participants achieved a mean accuracy of 87.8 % ± sd 9.1 % with a mean reaction time of 1742 ms ± sd 525 ms; as response speed: 0.62 s^{-1} ± sd 0.17 s^{-1}.

As expected, behaviour depended on the different combinations of listening cues (Fig. 2c,d). Informative compared to uninformative spatial-attention cues yielded a strong behavioural benefit. In selective-attention trials, participants responded more accurately and faster (accuracy: generalized linear mixed-effects model (GLMM); *odds*

*ratio (OR)=3.5, std. error (SE) =.12, p<.001; response speed: linear mixed-effects model
(LMM); β=.57, SE=.04, p<.001; see Supplementary Tables 1 and 2).* That is, when cued to
one of the two sides, participants responded on average 261 ms faster and their
probability of giving a correct answer increased by 6 %.

Also, participants responded generally faster in trials in which they were given
a specific, more informative semantic cue (LMM; *β=.2, SE=.03, p<.001*), most likely
reflecting a semantic priming effect that led to faster word recognition. Contrary to our
expectations, a more informative semantic cue did not lead to more accurate responses
(GLMM; *OR=1.1, SE=.11, p=.69*).

As in a previous fMRI implementation of this task[35], we did not find evidence for
any interactive effects of the two listening cues on either accuracy (GLMM; *OR=1.3,
SE=.21, p=.36*) or response speed (LMM; *β=.09, SE=.06, p=.31*). Moreover, the breakdown
of error trials revealed a significantly higher proportion of spatial stream confusions (6
% ± sd 8.3 %) compared to random errors (3 % ± sd 3.4 %; paired t-test on logit-
transformed proportions: t_{155} = 6.53, *p*<.001; see Supplementary Fig. 2). The increased
rate of spatial stream confusions (i.e., responses in which the last word of the to-be-
ignored sentence was chosen) attests to the distracting nature of dichotic sentence
presentation and thus heightened task difficulty.

Spatial attention modulates both alpha lateralization and neural speech tracking in
auditory cortex

In line with our second research question, following source projection of EEG data, we
probed whether the presence of an informative spatial attention cue would lead to
reliable modulation of both alpha power and neural speech tracking within an *a priori*
defined auditory region of interest (ROI; see Supplementary Fig. 3 and Methods for
details).

For alpha power, we expected an attention-induced lateralization due to a
decrease in power contralateral and concomitant increase in power ipsilateral to the
focus of attention. For neural speech tracking, we expected stronger neural tracking of
attended compared to ignored speech under selective attention but no such systematic
difference in the neural tracking of probed and unprobed sentences in divided-
attention trials. Accordingly, our analyses of alpha power and neural speech tracking
focused on attentional modulation index measures that contrast the relative strength
of neural responses to target versus distractor stimuli. In line with previous results, we
expected alpha lateralization to be present throughout the auditory sentence
presentation but to potentially increase around the task-relevant final word[12,14,48].

We compared alpha power changes ipsi- and contralateral to the probed ear to
derive a temporally-resolved single-trial measure of alpha power lateralization [alpha
lateralization index (ALI) = (α-power$_{ipsi}$ − α-power$_{contra}$) / (α-power$_{ipsi}$ + α-power$_{contra}$)][15].

As shown in Figure 3a, an informative spatial cue—that is, the instruction to pay
attention to a given side—elicited a pronounced lateralization of 8–12 Hz alpha power
within the auditory ROI. Lateralization of alpha power was evident following the spatial

cue itself and during dichotic sentence presentation with its strongest peak around final
word presentation.

As expected, the statistical analysis of alpha lateralization during sentence
presentation (time window: 3.5–6.5 s; see control analysis section below for results on
the final word period) revealed a significant modulation by attention that was
additionally influenced by the probed ear (LMM; spatial cue x probed ear: β=.13, SE=.02,
p<.001; Fig. 3b). Follow-up analysis showed a significant difference in alpha
lateralization between selective- and divided-attention trials when the right ear but not
when the left was probed (LMM, right ear probed: β=.12, SE=.01, p<.001; LMM, left ear
probed: β=−.016, SE=.013, p=.55; see Supplementary Table 3). This pattern suggests that
when given an uninformative spatial cue, participants presumably payed overall more
attention to the left-ear stimulus leading to an increase in alpha lateralization for
probed-left compared to probed-right trials.

Notably, we did not find any evidence for a modulation by the semantic cue nor
any joint influence of the spatial and semantic cue on alpha lateralization during
sentence presentation (LMM; semantic cue main effect: β=−.01, SE=.01, p=.53, spatial x
semantic cue: β=−.02, SE=.02, p=.53).

Not least, the extent of overall as well as attention-specific alpha lateralization
was unaffected by participants' chronological age and hearing loss (p-values > .27 for
main effects of age, PTA, and their respective interactions with the spatial-attention cue;
see also Supplementary Table 6 for a corresponding analysis of alpha power during the
interval of the final word).
272

Figure 3. Informative spatial cue elicits increased alpha power lateralization before and during speech presentation.

(a) Attentional modulation of 8–12 Hz auditory alpha power throughout the trial for all N=155 participants. Brain models indicate the spatial extent of the auditory region of interest (shown in red). Purple traces show the grand-average alpha lateralization index (ALI) for the informative (solid dark purple line) and uninformative spatial cue (dashed light purple line), each collapsed across semantic cue levels. Error bands indicate ± 1 SE. Positive values indicate relatively higher alpha power in the hemisphere ipsilateral to the attended/-probed sentence compared to the contralateral hemisphere. Shaded grey area shows the time window of sentence presentation.

(b) ALI during sentence presentation (3.5–6.5 s) shown separately per spatial cue condition and probed ear (left plot). Coloured dots show trial-averaged individual results, black dots and error bars indicate the grand-average and bootstrapped 95% confidence intervals. Box plots show median, 25th to 75th percentile data, and 1.5 interquartile range. For probed-right trials there was a significant difference in ALI between selective- and divided-attention trials (right plot). Black dots represent individual mean ALI values ± bootstrapped 95 % confidence intervals. Histogram shows distribution of differences in ALI for informative vs. uninformative spatial-cue levels. β: slope parameter estimate from the corresponding general linear mixed-effects model; *** = p<.001.

In close correspondence to the alpha-power analysis, we investigated whether changes in attention or semantic predictability would modulate the neural tracking of attended versus ignored speech. We used linear backward ('decoding') models to reconstruct the onset envelopes of the to-be-attended and ignored sentences (for simplicity hereafter referred to as *attended* and *ignored*) from neural activity in the auditory ROI. Reconstruction models were trained on selective-attention trials only, but then utilized to reconstruct attended (probed) and ignored (unprobed) envelopes for both attention conditions (see Methods, Fig. 4a and Supplementary Fig. 4 for details).

In line with previous studies[26,28,49], the forward transformed temporal response functions (TRFs) show increased encoding of attended compared to ignored speech in the time window covering the $N1_{TRF}$ and $P2_{TRF}$ component (see Fig. 4b, left panel). Here, however, this was observed particularly for right-ear inputs processed in the left auditory ROI.

Further attesting to the validity of our reconstruction models, reconstructed attended envelopes were overall more similar to the envelope of the to-be-attended sentence than to that of the to-be-ignored sentence, and vice versa for the reconstructed ignored envelopes (see Fig. 4b, right panel).

As shown in Figure 4c, the differential neural tracking of attended and ignored envelopes (probed and unprobed envelopes under divided attention) was modulated

by attention. Following an informative spatial cue, the neural tracking index becomes
increasingly positive during the second half of sentence presentation with its highest
peaks around final word onset.

The statistical analysis of single-trial index values averaged for the time interval
of final word presentation confirmed this pattern: The difference in the neural tracking
of attended and ignored sentence was generally more pronounced under selective
compared to divided attention (see control analysis section below for results on the
entire sentence presentation). However, this effect was also modulated by differences
in sentence onset: The difference in neural speech tracking between the two attention
conditions was reduced when the attended/probed sentence started ahead of the
distractor sentence. This effect was driven by an increase in differential neural speech
tracking for divided attention in such trials: in absence of an informative spatial cue,
participants' attention was captured by the sentence with the earlier onset.
Consequently, we observed overall more positive index values when the earlier
sentence was probed compared to when it was not probed (LMM, earlier onset x spatial
cue: β=−.05, SE=.02, p=.049, see Fig. 4d and Supplementary Table 4 for full model
details).

We also found a neural correlate of the known right-ear advantage for verbal
materials, that is, an overall stronger tracking of left-ear inputs. This effect was
independent of spatial-attention cueing (LMM; probed ear main effect: β=−.03, SE=.01,
p=.023; spatial cue x probed ear: β=.02, SE=.02, p=.54). As for alpha power, we did not
observe any modulation of neural tracking by the semantic cue, nor any joint influence
of the spatial and semantic cue (LMM; semantic cue main effect: β=−.01, SE=.01, p=.53,
interaction spatial x semantic cue: β=−.02, SE=.02, p=.53).

Again, participants' age and hearing status did not prove significant predictors
of neural speech tracking (p-values > .54 for main effects of age, PTA, and their
respective interactions with the spatial-attention cue, see also Supplementary Table 7
for a corresponding analysis of neural tracking during the entire sentence presentation).

Figure 4. Neural speech tracking of attended and ignored sentences.

(a) Schematic representation of linear backward model approach used to reconstruct onset envelopes from recorded EEG responses. Following the estimation of linear backward models using only selective-attention trials (see Supplementary Fig. 4 for details), envelopes of attended and ignored sentences were reconstructed via convolution of EEG responses with the estimated linear backward models within the auditory ROI. Reconstructed onset envelopes were compared to the onset envelopes of presented sentences to assess neural tracking strength and decoding accuracy (see Supplements).

(b) Left: forward-transformed temporal response functions (TRFs) for attended (green) and ignored (red) speech trained on selective-attention trials. Models are averaged across all N=155 participants, and shown separately for the left and right auditory ROI. Error bands indicate 95 % CI. Right: Single-subject average Pearson correlation of reconstructed envelopes with those of presented sentences. Top panel shows results for reconstruction of the attended envelope, bottom panel for the ignored envelope along with 95 % CI. Shown for all N=155 participants.

(c) Top: Grand average peri-stimulus time course (see Supplements for details) of neural tracking index shown separately for selective (solid dark green curve) and divided attention (dashed light green curve) ± 1 SE error band. Histograms indicate the distribution of sentence and final word onsets, respectively. Grey shaded area shows time window of final word presentation used for statistical analysis. Bottom: 45° plot contrasting the neural tracking of attended (probed) and ignored (distractor) envelope during final word presentation for selective and divided attention, respectively.

(d) Left: Neural tracking index shown per spatial-attention condition and for trials in which cued/probed sentences started ahead of ('probed first') or after the distractor ('probed second'). Coloured dots represent trial-averaged individual results, black dots and error bars indicate the grand-average and bootstrapped 95 % CI. Box plots show median, 25th to 75th percentile data, and 1.5 interquartile range. Right: Significant difference in neural tracking between selective- and divided-attention trials in probed-second trials (top plot), and stronger neural tracking in probed-left trials. Black dots in 45° plot represent individual mean neural tracking index ± bootstrapped 95 % CI. Histogram shows distribution of differences in neural tracking in informative vs. uninformative spatial-cue trials, and probed-left vs. probed-right trials,

354   respectively. β: slope parameter estimate from the corresponding general linear mixed-effects model; *** =
355   p<.001, *= p<.5.
356

Trial-to-trial neural speech tracking is independent of synchronous alpha lateralization

Our third major goal was to investigate whether neural speech tracking versus the
modulation of alpha power reflect two dependent neural mechanisms of auditory
attention at all. We asked whether neural speech tracking could be explained by
auditory alpha lateralization either at the state level (i.e., within an individual from trial
to trial) or at the trait level (i.e., between individual mean levels; see Statistical analysis
for details). If modulations of alpha power over auditory cortices indeed act as a neural
filter mechanism to selectively gate processing during early stages of sensory analysis
then heightened levels of alpha lateralization should lead to a more differentiated
neural tracking of the attended vs. ignored speech and thus more a positive neural
tracking index (cf. Fig. 5a).

However, in the analysis of the task-relevant final word period, we did not find
evidence for an effect of alpha lateralization on neural speech tracking at neither the
state nor the trait level (Fig. 5b, LMM; ALI within-subject effect: β=−.008, *SE*=.005, *p*=.35;
ALI between-subject effect: β=−.0007, *SE*=.007, *p*=.98; see Supplementary Table 5). This
notable absence of an alpha lateralization–neural speech tracking relationship held
irrespective of spatial-attention condition or probed ear (all *p-values*>.35).

To complement our fine-grained single-trial level investigation into the brain-
brain relationship with a coarser, yet time-resolved analysis, we related the temporal
dynamics of both neural measures in an exploratory between-subjects cross-correlation
analysis. As shown in Figure 5c, under selective attention, neural speech tracking and
alpha lateralization follow different temporal trajectories with neural speech tracking
peaking earlier than alpha lateralization around final word presentation. The average
cross-correlation of the two neural time courses during sentence presentation confirms
a systematic temporal delay with fluctuations in neural speech tracking leading those in
alpha power by about 520 ms (see Fig. 5d).

Figure 5. Relationship of alpha lateralization and neural speech tracking.
(a) Hypothesized relationship of alpha power and neural speech tracking within the auditory region of interest. Changes in alpha lateralization are assumed to drive changes in neural tracking. Schematic representation for an attend-left trial.
(b) Independence of neural speech tracking and alpha lateralization during the final word presentation as shown by the predictions from the same linear mixed effect model. Plots show the predicted, non-significant effect of within- and between-subject variations in alpha lateralization on selective neural tracking, respectively. Blue lines indicates the respective predicted fixed effects ± 95 % CI, grey thin lines in the left plot show N=155 subject-specific random slopes (included for illustrative purposes only), and grey dots show average predictions per subject. β: slope parameter estimate from the corresponding general linear mixed-effects model; *** = p<.001.
(c) Grand-average time courses of alpha lateralization and neural speech tracking during sentence presentation mapped to the same peri-stimulus time axis. Shown separately for selective-attention (darker, solid curves) and divided-attention trials (lighter, dashed curves). Error bands reflect ± 1 SE. Note how the peak in neural speech tracking under selective attention precedes the peak in alpha lateralization.
(d) Mean normalized cross-correlation of trial-averaged neural speech tracking and alpha lateralization time courses. Upper and lower bound of shaded area reflect the 97.5th and 2.5th percentile of surrogate data derived from 5000 independently permuted time courses of alpha power and neural speech tracking.

Neural speech tracking but not alpha lateralization explains listening behaviour

Having established the functional independence of alpha lateralization and neural speech tracking at the single-trial level, the final piece of our investigation was to probe their relative functional importance for behavioural outcome.

Using the same (generalized) linear mixed-effects models as in testing our first research question (Q1 in Fig.1), we investigated whether changes in task performance could be explained by the independent (i.e., as main effects) or joint influence (i.e., as an interaction) of neural measures. Again, we modelled the influence of the two neural filter strategies on behaviour at the state and trait level[50].

For response accuracy, our most important indicator of listening success, we
observed an effect of trial-by-trial variation in neural speech tracking both during the
presentation of the final word and across the entire sentence: participants had a higher
chance of responding correctly in trials in which they neurally tracked the cued/probed
sentence more strongly than the distractor sentence (see Fig. 6a, left panel). For changes
in neural speech tracking extracted from the entire sentence presentation, this effect
occurred independently of other modelled influences (GLMM; main effect neural
tracking (within-subject effect): *OR*=1.06, *SE*=.02, *p*=.03; see Supplementary Table 12)
while it was generally less pronounced and additionally modulated by the probed ear
for the period of the task-relevant final word (GLMM; probed ear x neural tracking
(within-subject effect): *OR*=1.1, *SE*=.04, *p*=.03; see Supplementary Table 1).

The data held no evidence for any direct effects of trial-to-trial or participant-to-
participant variation in alpha lateralization during sentence or final word presentation
on accuracy (all *p*-values > .18; see Supplementary Tables 1 and 12). We also did not find
evidence for any joint effects of alpha power and neural speech tracking extracted from
either of the two time windows (all *p*-values >.33). Importantly, the absence of an effect
did not hinge on differences in neural measures across spatial-cue, or probed-ear levels
as relevant interactions of neural measures with these predictors were included in the
model (all *p*-values > .55).

The observed effects of neural filters on response speed depended on the
analysed time window: while participants with relatively higher average levels of neural
speech tracking during sentence presentation responded overall faster (LMM, neural
tracking (between-subject effect): *β*=0.08, SE=.03, *p*=.01; see Fig 6a, right panel and
Supplementary Table 13), we found a combined effect of neural dynamics during final
word presentation. Under selective but not divided attention, response speed
depended on a combination of trial-to-trial variation in both alpha lateralization and
neural speech tracking (LMM; spatial cue x ALI (within-subject effect) x neural tracking
index (within-subject effect): *β*=0.08, SE=.03, *p*=.01; see Supplementary Table 2). In
short, responses were fastest in trials where relatively elevated levels in either neural
speech tracking or alpha lateralization were paired with relatively reduced levels in the
respective other neural measure thus highlighting the influence of two independent
complementary filter solutions (see also Supplementary Fig. 5).

In line with the literature on listening behaviour in ageing adults[e.g. 51,52], the
behavioural outcome was further reliably predicted by age, hearing loss, and probed
ear. We observed that participants' performance varied in line with the well-attested
right-ear advantage (REA, also referred to as *left-ear disadvantage*) in the processing of
linguistic materials[44]. More specifically, participants responded both faster and more
accurately when they were probed on the last word presented to the right compared to
the left ear (response speed: LMM; *β*=.08, *SE*=.013, *p*<.001; accuracy: GLMM*; OR*=1.25,
SE=.07, *p*=.006; see also Supplementary Fig. 6).

Increased age led to less accurate and slower response (accuracy: GLMM; *OR*=.80, *SE*=.08, *p*=.025; response speed: LMM; *β*=−.15, *SE*=.03, *p*<.001). In contrast, increased hearing loss led to less accurate (GMM; *OR*=.75, *SE*=.08, *p*=.002) but not slower responses (LMM; *β*=−.05, *SE*=.03, *p*=.21, see Supplementary Tables 1–2, and Supplementary Fig. 7).

Figure 6. Neural speech tracking predicts listening behavior.

(a) Model predictions for the effect of neural tracking on behavior for N=155 participants. Left panel shows the predicted group-level fixed effect (green line ± 95 % CI) of trial-to-trial variation in neural tracking on accuracy. Grey thin lines indicate estimated subject-specific slopes. Right panel shows the predicted group-level fixed effect of neural tracking at the between-subject level on response speed (green line ± 95 % CI). Grey dots indicate subject-level model predictions. OR = odds ratio, *β* = β: slope parameter estimate from the corresponding general linear mixed-effects model; ** = p<.01, * = p<.05.

(b) Summary of results. Black arrows highlight statistically significant effects from (generalized) single-trial linear mixed-effects modelling. Grey arrow connecting alpha power dynamics with neural tracking reflects their systematic temporal delay observed in the exploratory cross-correlation analysis. Notably, changes in age and hearing loss did not modulate the fidelity of the two key neural measures.

Control analyses

We ran additional control analyses to validate our main set of results. First, we asked whether the observed independence of alpha lateralization and neural speech tracking hinged on the precise time window and cortical site from neural measures were extracted. One set of control models thus included alpha lateralization during spatial-cue rather than sentence presentation as a predictor of neural speech tracking during sentence and final word presentation. Next, we related the two neural filters during the entire sentence presentation rather than only during the final word.

Second, we tested the hypothesis that neural speech tracking might be driven not primarily by alpha-power modulations emerging in auditory cortices, but rather by those generated in domain-general attention networks in parietal cortex[53]. We therefore ran control models including alpha lateralization within the inferior parietal lobule. However, none of these additional analyses found evidence for an effect of alpha lateralization on neural speech tracking (see Supplementary Tables 8–11 and Supplementary Fig. 8).

Third, we asked whether our neural speech tracking results were impacted by
the range of time lags used for reconstruction, or by the specific decoder model
underlying the neural tracking index. Reconstructing envelopes using a shorter time
window (50–250 ms) did not significantly change the resulting neural tracking index
values (LMM, β = .002, *SE* = .007, *p* = .84; see also Supplementary Fig. 9). In a separate
analysis, we calculated the neural tracking index using only the attended decoder
model, and probed its influence on behaviour. The results are overall in line with our
main conclusions and particularly underscore the impact of neural speech tracking on
response accuracy (see Supplementary Tables 14–17 for details).
Finally, we tested whether changes in age or hearing loss would modulate the
relationship of neural tracking and alpha lateralization with listening behaviour.
However, the inclusion of the respective interaction terms did not further improve the
statistical models of accuracy and response speed (all *p*-values > .44).
493

Discussion

We have utilized the power of a large, representative sample of middle-aged and older listeners to explicitly address the question of how the two most eminent neurobiological implementations of attentional filtering, typically studied in isolation, relate to one another, and how they jointly shape listening success. In addition, we leveraged our large age-varying sample to ask how chronological age and hearing loss affect the fidelity of neural filter strategies and their influence on behaviour.

In our dichotic listening task, we source-localized the electroencephalogram, and primarily focused on systematic spatial-cue-driven changes within auditory cortex in alpha lateralization and the neural tracking of attended versus ignored speech. These results provide unprecedented large-sample support for their suggested roles as neural instantiations of selective attention.

First, an informative spatial-attention cue not only boosted both neural measures but also consistently boosted behavioural performance. Listening behaviour was additionally influenced by both trial-to-trial and individual-to-individual variation in neural speech tracking, with relatively stronger tracking of the target sentence leading to better performance. An informative semantic cue led to faster responses but did not affect the two neural measures, thus most likely reflecting a priming effect speeding up the analysis of response alternatives rather than the differential processing of the sentences themselves.

Second, when related at the single-trial, single-subject level, the two neural attentional filter mechanisms were found to operate statistically independently of each other. This underlines their functional segregation and speaking to two distinct neurobiological implementations. Yet, when related in a coarser, between-subjects analysis across time, peaks in selective neural tracking systematically preceded those in alpha lateralization.

Importantly, while chronological age and hearing loss reliably decreased behavioural performance they did not systematically affect the fidelity of neural filter strategies nor their influence on behaviour.

Neural speech tracking but not alpha lateralization predicts listening success

The present study explicitly addressed the often overlooked question of how neural filter states (i.e., fluctuations from trial-to-trial) impact behaviour, here single-trial listening success[54-57]. Using a sophisticated linear-model approach that probed the impact of both state- and trait-level modulation of neural filters on behaviour, we only found evidence for a direct influence of neural speech tracking but not alpha lateralization on behavioural performance even though all three measures were robustly modulated by the presence of a spatial cue (see Fig. 6). What could be the reason for this differential impact of neural measures on behaviour?

To date, the behavioural relevance of selective neural speech tracking is still poorly supported given the emphasis on more naturalistic, yet more complex language stimuli[58,59]. While these stimuli provide a window onto the most natural forms of speech

comprehension, they are not easily paired with fine-grained measures of listening
behaviour. This makes it particularly challenging to establish a direct link between
differential neural speech tracking and listening success[23,25,26,49,60]. Nevertheless, there is
preliminary evidence linking stronger neural tracking to improved comprehension
when it is tested at a comparably high level[28] (i.e., content questions on longer speech
segments). Our current results thus provide important additional fine-grained and
temporally resolved support to the functional relevance of selective neural speech
tracking for moment-to-moment listening behaviour[61-63].

Despite a vast number of studies investigating the role of (lateralized) alpha
oscillations in attentional tasks, the circumstances under which their top-down
modulation may affect the behavioural outcome are still insufficiently understood[31].
Rather, the presence of a stable brain–behaviour relationship hinges on several factors.

First, the link of neural filter state to behaviour seems to be impacted by age:
most evidence linking increased alpha lateralization to better task performance in
spatial-attention tasks stems from smaller samples of young adults[12,15,64,65]. By contrast,
the presence of such a systematic relationship in middle-age and older adults is
obscured by considerable variability in age-related changes at the neural (and to some
extent also behavioural) level[14,66-69] (see discussion below).

Second, previous findings differ along (at least) two dimensions: (i) whether the
functional role of alpha lateralization is studied during attention cueing, stimulus
anticipation, or stimulus presentation[9,66,70], and (ii) whether behaviour is related to the
overall strength of alpha lateralization or its stimulus-driven rhythmic modulation[12,14].
Depending on these factors, the observed brain–behaviour relations may relate to
different top-down and bottom-up processes of selective auditory attention.

Third, as shown in a recent study by Wöstmann et al.[70], the neural processing of
target and distractor are supported by two uncorrelated lateralized alpha responses
emerging from different neural networks. Notably, their results provide initial evidence
for the differential behavioural relevance of neural responses related to target selection
and distractor suppression, respectively.

In summary, it is still a matter of debate by which mechanistic pathway, and at
which processing stage the modulation of alpha power will impact behaviour. While it
is (often implicitly) assumed that alpha oscillations impact behaviour via a modulation
of neural excitability and thus early sensory processing, there is little evidence that
shows a direct influence of alpha oscillation on changes in neural excitability and on
subsequent behaviour[31,71].

Lastly, the increase in alpha lateralization around final word presentation could
at least partially reflect post-perceptual processes associated with response selection
rather than the perceptual analysis itself [72]. The observed combined influence of neural
tracking and alpha lateralization on response speed but not accuracy would seem
compatible with such an interpretation (but see also ref.[73] for the combined influence of
non-lateralized alpha power and neural speech tracking on intelligibility in a non-spatial
listening task).

Taken together, our results underscore the impact of prioritized sensory
encoding of relevant sounds via selective neural speech tracking on listening
performance and highlight the difficulty in establishing a comparable link for a neural
signature as multifaceted as alpha oscillations[74-76].

Are fluctuations in lateralized alpha power and neural speech tracking functionally
connected?

We investigated attention-related changes in two neural filter strategies that (i) involve
neurophysiological signals operating at different frequency regimes, (ii) are assumed to
support auditory attention by different neural mechanisms, and (iii) are typically studied
in isolation[6,22]. Here, we found both neural filter strategies to be impacted by the same
spatial-attention cue which afforded new insights into their neurobiological
dependence.

There is preliminary evidence, mostly from between-subjects analyses,
suggesting that the two neural filter strategies may exhibit a systematic
relationship[9,14,32,33,77]. How the two neural filter strategies may be connected
mechanistically is thus still an open question. We here asked whether concurrent
changes in neural filter states would imply a neural hierarchy in which alpha-driven
controlled inhibition modulates the amplification of behaviourally relevant sensory
information via selective neural speech tracking[78-80].

Our in-depth trial-by-trial analysis revealed independent modulation of alpha
power and neural speech tracking. At the same time, in our exploratory between-
subjects cross-correlation analysis we observed a systematic temporal delay with peaks
in neural speech tracking leading those in alpha lateralization. While the direction and
duration of this delay were closely in line with previous findings[12,14], at this coarser level
of analysis, they speak against a hierarchy of neural processing in which lateralized alpha
responses govern the differential neural tracking of attended versus ignored speech[81].

Our single-trial results are well in line with recent reports of independent
variation in alpha band activity and steady-state (SSR) or frequency following responses
(FFR) in studies of visual spatial attention[30,31,82]. Additionally, the inclusion of single-trial
alpha lateralization as an additional training feature in a recent speech-tracking study
failed to improve the decoding of attention[83]. The results from our most fine-grained
single-trial level of analysis thus speak against a consistent, linear relationship of
momentary neural filter states. We observed instead the co-existence of two
complementary but seemingly independent neurobiological solutions to the
implementation of auditory selective attention.

How can this finding be reconciled with findings from previous electrophysiological
studies[9,32,33] pointing towards a functional trade-off between neurobiological
attentional filter mechanisms? And what could be an advantage to independent neural
solutions for selective auditory attention?

Our between-subjects cross-correlation analysis appears to provide at least
tentative support for a systematic relationship in which peaks in neural speech tracking
precede those in alpha lateralization. A closer inspection of the group-level temporal
modulation of neural measures throughout sentence presentation, however, reveals
some important differences to previous results. Whereas earlier studies reported an
acyclic waxing and waning of neural entrainment and alpha power in response to
rhythmic auditory stimulation[12,14,33], in the current study, the two neural measures show
different temporal dynamics: neural speech tracking gradually increases leading up to
the final word, while alpha lateralization peaks at sentence and final word onset. The
temporal dynamics of alpha lateralization, in particular, may point to strategic
intermittent engagement of spatial attention in line with task demands[48].

Do these differences in temporal dynamics of the two neural filters challenge
the existence of a systematic single-trial brain–brain relationship? Yes, but they also
point to a potential benefit of independent neural filter solutions. If the two neural
measures of auditory attention were indeed functionally unconnected as suggested by
the current results, they would allow for a wider range of neural filter state
configurations to flexibly adapt to the current task demands and behavioural goals. The
co-existence of two independent but complementary filter mechanisms operating
either via the selective amplification of relevant or via the controlled inhibition of
irrelevant sounds, enables different modes of auditory attention to serve a listener's
goal in the face of complex real-life listening situations[20,84,85].

640
641

Do age and hearing loss affect neural filter strategies?

The detrimental effects of increasing age and associated hearing loss on speech
comprehension in noisy listening situations are well attested[86] and borne out by the
current results. However, the extent to which the neural implementations of attentional
filtering are affected by age and hearing loss, and in how far they may constitute neural
markers of age-related speech comprehension problems, remains poorly understood[51].

As in a previous study on a subset of the current sample, we found the fidelity
of alpha lateralization unchanged with age[14]. Other studies on auditory attention,
however, have observed diminished and less sustained alpha lateralization for older
compared to younger adults that were to some extend predictive of behaviour[66,87,88].

Our observation of preserved neural speech tracking across age and hearing
levels only partially agree with earlier findings. They are in line with previous reports of
differential neural tracking of attended and ignored speech for hearing impaired older
adults that mirrored the attentional modulation observed for younger or older normal-
hearing adults[45-47,89]. As revealed by follow-up analysis (data not shown), however, our
data do not provide evidence for a differential impact of hearing loss on the neural
tracking of attended or ignored speech as found in some of these studies. We also did
not find evidence for overall increased levels of cortical neural tracking with age as
observed in earlier studies[90,91].

The discrepancy in results may be explained by differences in (i) the studied
populations (i.e., whether groups of younger and older participants were contrasted
compared to the modelling of continuous changes in age and hearing loss), (ii) whether
natural stories or short Matrix sentence speech materials were used[92], or (iii) by
differences in task details. In sum, the results suggest that the commonly observed
adverse effects of age and hearing loss on speech-in-noise processing are not readily
paired with concomitant changes at the neural level.
668

Conclusion

In a large, representative, age-varying sample of listeners, we underscore the functional
significance of lateralized alpha power and neural speech tracking to spatial attention.
Our results point to the co-existence of two independent yet complementary neural
filter mechanisms to be flexibly engaged depending on a listener's attentional goals.
However, we see no direct, behaviourally relevant impact of alpha power modulation
on early sensory gain processes.

Only for neural speech tracking, we establish a mechanistic link from trial-to-trial
neural filtering during the concurrent sound input to the ensuing behavioural outcome.
This link exists irrespective of age and hearing status, which points to the potency of
neural speech tracking to serve as an individualized marker of comprehension problems
in clinical settings and as a basis for translational neurotechnological advances.

This key advance notwithstanding, the notable absence of an association
between alpha lateralization and listening behaviour also highlights the level of
complexity associated with establishing statistically robust relationships of complex
neural signatures and behaviour in the deployment of auditory attention. To
understand how the brain enables successful selective listening it is necessary that
studies go beyond characterization of neurobiological filter mechanisms alone, and
further jointly account for the variability in both neural states and behavioural
outcomes[93].
689
690
691

Materials and Methods

Data collection

The analysed data are part of an ongoing large-scale study on the neural and cognitive mechanisms supporting adaptive listening behaviour in healthy middle-aged and older adults ("The listening challenge: How ageing brains adapt (AUDADAPT)"; https://cordis.europa.eu/project/rcn/197855_en.html). This project encompasses the collection of different demographic, behavioural, and neurophysiological measures across two time points. The analyses carried out on the data aim at relating adaptive listening behaviour to changes in different neural dynamics[35,36].

Participants and procedure

A total of $N = 155$ right-handed German native speakers (median age 61 years; age range 39–80 years; 62 males; see Supplementary Fig. 1 for age distribution) were included in the analysis. Handedness was assessed using a translated version of the Edinburgh Handedness Inventory[94]. All participants had normal or corrected-to-normal vision, did not report any neurological, psychiatric, or other disorders and were screened for mild cognitive impairment using the German version of the 6-Item Cognitive Impairment Test (6CIT[95]).

During the EEG measurement, participants performed six blocks of a demanding dichotic listening task (see Fig. 2 and Supplements for details on task and sentence materials).

As part of our large-scale study on adaptive listening behaviour in healthy aging adults, prior to the EEG session, participants also underwent a session consisting of a general screening procedure, detailed audiometric measurements, and a battery of cognitive tests and personality profiling (see ref.[14] for details). Only participants with normal hearing or age-adequate mild-to-moderate hearing loss were included (see Supplementary Fig. 1 for individual audiograms). As part of this screening procedure an additional 17 participants were excluded prior to EEG recording due to non-age-related hearing loss or a medical history. Three participants dropped out of the study prior to EEG recording and an additional 9 participants were excluded from analyses after EEG recording: three due to incidental findings after structural MR acquisition, and six due to technical problems during EEG recording or overall poor EEG data quality. Participants gave written informed consent and received financial compensation (8€ per hour). Procedures were approved by the ethics committee of the University of Lübeck and were in accordance with the Declaration of Helsinki.

Behavioural data analysis

We evaluated participants' behavioural performance in the listening task with respect to accuracy and response speed. For the binary measure of accuracy, we excluded trials in which participants failed to answer within the given 4-s response window ('timeouts'). Spatial stream confusions, that is trials in which the sentence-final word of the to-be-ignored speech stream were selected, and random errors were jointly classified as incorrect answers. The analysis of response speed, defined as the inverse of reaction time, was based on correct trials only. Single-trial behavioural measures were subjected to (generalized) linear mixed-effects analysis and regularized regression (see Statistical analysis).

EEG data analysis

The preprocessed continuous EEG data (see Supplements for details on data collection and preprocessing) were high-pass-filtered at 0.3 Hz (finite impulse response (FIR) filter, zero-phase lag, order 5574, Hann window) and low-pass-filtered at 180 Hz (FIR filter, zero-phase lag, order 100, Hamming window). The EEG was cut into epochs of –2 to 8 s relative to the onset of the spatial-attention cue to capture cue presentation as well as the entire auditory stimulation interval.

For the analysis of changes in alpha power, EEG data were down-sampled to f_s=250 Hz. Spectro-temporal estimates of single-trial data were then obtained for a time window of –0.5 to 6.5 s (relative to the onset of the spatial-attention cue) at frequencies ranging from 8 to 12 Hz (Morlet's wavelets; number of cycles = 6).

For the analysis of the neural encoding of speech by low-frequency activity, the continuous preprocessed EEG were down-sampled to f_s=125 Hz and filtered between f_c=1 and 8 Hz (FIR filters, zero-phase lag, order: $8f_s/f_c$ and $2f_s/f_c$, Hamming window). The EEG was cut to yield individual epochs covering the presentation of auditory stimuli, beginning at noise onset until the end of auditory presentation.

Following EEG source and forward model construction (see Supplements for details), sensor-level single-trial data in each of our two analysis routines were projected to source space by matrix-multiplication of the spatial filter weights. To increase signal-to-noise ratio in source estimates and computationally facilitate source-level analyses, source-projected data were averaged across grid-points per cortical area defined according to the HCP functional parcellation template[96,97]. This parcellation provides a symmetrical delineation of each hemisphere into 180 parcels for a total of 360 parcels. We constrained the analysis of neural measures to an a priori defined, source-localized auditory region of interest (ROI) as well as one control ROI in the inferior parietal lobule (see Supplements for details).

Attentional modulation of alpha power

Absolute source power was calculated as the square-amplitude of the spectro-temporal estimates. Since oscillatory power values typically follow a highly skewed, non-normal distribution, we applied a nonlinear transformation of the Box-Cox family (power$_{trans}$ =

772 (power^p −1)/p with p=0.5) to minimize skewness and to satisfy the assumption of
773 normality for parametric statistical tests involving oscillatory power values[98]. To quantify
774 attention-related changes in 8–12 Hz alpha power, per ROI, we calculated the single-
775 trial, temporally resolved alpha lateralization index as follows[12,14,15].

776
777 ALI = (α-power$_{ipsi}$ − α-power$_{contra}$)/(α-power$_{ipsi}$ + α-power$_{contra}$)   (1)
778

To account for overall hemispheric power differences that were independent of
attention modulation, we first normalized single-trial power by calculating per parcel
and frequency the whole-trial (−0.5–6.5 s) power averaged across all trials and
subtracted it from single trials. We then used a robust variant of the index that applies
the inverse logit transform [(1 / (1+ exp(−x))] to both inputs to scale them into a
common, positive-only [0;1]-bound space prior to index calculation.
For visualization and statistical analysis of cue-driven neural modulation, we
then averaged the ALI across all parcels within the auditory ROI and extracted single-
trial mean values for the time window of sentence presentation (3.5–6.5 s), and treated
them as the dependent measure in linear mixed-effects analysis (see Statistical Analysis
below). They also served as continuous predictors in the statistical analysis of brain–
behaviour and brain–brain relationships. We performed additional analyses that
focused on the ALI in auditory cortex during presentation of the sentence-final word
and spatial-attention cue, respectively. Further control analyses included single-trial ALI
during sentence and final-word presentation that were extracted from the inferior
parietal ROI.

795
796
Estimation of envelope reconstruction models

To investigate how low-frequency (i.e., < 8 Hz) fluctuations in EEG activity relate to the
encoding of attended and ignored speech, we trained stimulus reconstruction models
(also termed decoding or backward models) to predict the onset envelope (see
Supplements for details) of the attended and ignored speech stream from EEG[99,100]. In
this analysis framework, a linear reconstruction model *g* is assumed to represent the
linear mapping from the recorded EEG signal, *r(t,n)*, to the stimulus features, *s(t)*:
804
805 $\hat{s}(t) = \sum_n \sum_\tau g(\tau, n) r(t + \tau, n)$ (2)
806

807 where *s^(t)* is the reconstructed onset envelope at time point *t*. We used all parcels within
808 the bilateral auditory ROI and time lags τ in the range of −100 ms to 500 ms to compute
809 envelope reconstruction models using ridge regression[101]:
810
811 $g = (R^T R + \lambda m I)^{-1} R^T s$ (3)
812

where R is a matrix containing the sample-wise time-lagged replication of the neural
response matrix r, λ is the ridge parameter for regularization, I is the identity matrix , and
m is a subject-specific scalar representing the mean of the trace of R^{T}R[102,103]. The same

grid of ridge parameters ($\lambda = 10^{-5}, 10^{-4}, \ldots 10^{10}$) was used across subject, and *m* proved to
be relatively stable across subjects (387.2 ± 0.18, mean ± sd). The optimal ridge value of
$\lambda = 1$ was determined based on the average Pearson's correlation coefficient and mean
squared error of the reconstructed and actually presented envelope across all trials and
subjects.

Compared to linear forward ('encoding') models that derive temporal response
functions (TRFs) independently for each EEG channel or source, stimulus reconstruction
models represent multivariate impulse response functions that exploit information
from all time lags and EEG channels/sources simultaneously. To allow for a
neurophysiological interpretation of backward model coefficients, we additionally
transformed them into linear forward model coefficients following the inversion
procedure described in ref.[104]. All analyses were performed using the multivariate
temporal response function (mTRF) toolbox[99] for Matlab.
Prior to model estimation, we split the data based on the two spatial attention
conditions (selective vs. divided), resulting in 120 trials per condition. Envelope
reconstruction models were trained on concatenated data from selective-attention
trials, only. Prior to concatenation, single trials were zero-padded for 600ms to reduce
discontinuity artefacts, and one trial was left out for subsequent testing. On each
iteration, two different backward models were estimated, an envelope reconstruction
model for the-be-attended speech stream (short: attended reconstruction model), and
one for the to-be-ignored speech stream (short: ignored reconstruction model.
Reconstruction models for attended and ignored speech signals were trained separately
for attend-left and attend-right trials which yielded 120 decoders (60 attended, 60
ignored) per attentional setting. For illustrative purposes, we averaged the forward
transformed models of attended and ignored speech per hemisphere across all
participants (Fig. 4B).

Evaluation of neural tracking strength

We analysed how strongly the attended compared to ignored sentences were tracked
by slow cortical dynamics by quantifying the envelope reconstruction accuracy for
individual trials. To this end, we reconstructed the attended and ignored envelope of a
given trial using a leave-one-out cross-validation procedure. The two envelopes of a
given trial were reconstructed using the models trained on all but the current trial from
the same attention condition. The reconstructed onset envelope obtained from each
model was then compared to onset envelopes of the actually presented speech signals
using a 248-ms sliding window (rectangular window, step size of 1 (8 ms) sample). The
resulting time courses of Pearson-correlation coefficients, $r_{attended}$ and $r_{ignored}$, reflect a
temporally resolved measure of single-trial neural tracking strength or reconstruction
accuracy[28] (see Fig. 4 and Supplementary Fig. 4).
We proceeded in a similar fashion for divided-attention trials. Since these trials
could not be categorized based on the to-be-attended and -ignored side, we split them
based on the ear that was probed at the end the trial. Given that even in the absence of

a valid attention cue, participants might still (randomly) focus their attention to one of
the two streams, we wanted to quantify how strongly the probed and unprobed
envelopes were tracked neurally. We used the reconstruction models trained on
selective-attention trials to reconstruct the onset envelopes of divided-attention trials.
Sentences presented in probed-left/unprobed-right trials were reconstructed using the
attend-left/ignore-right reconstruction models while probed-right/unprobed-left trials
used the reconstruction models trained on attend-right/ignore-left trials.
866
867 *Attentional modulation of neural tracking*

868 In close correspondence to the alpha lateralization index, we calculated a neural
869 tracking index throughout sentence presentation. The index expresses the difference in
870 neural tracking of the to-be-attended and ignored sentence (in divided attention:
871 probed and unprobed, respectively)[27]:
872
873 Neural tracking index $= (r_{attended} - r_{ignored})/(r_{attended} + r_{ignored})$ (4)
874
Positive values of the resulting index indicate that the attended envelope was tracked
more strongly than the ignored envelope, and vice versa for negative values. Since
individual sentences differed in length, for visualization and statistical analysis, we
mapped their resulting neural tracking time courses onto a common time axis
expressed in relative (percent) increments between the start and end of a given
stimulus. We first assigned each sample to one of 100 bins covering the length of the
original sentence in 1% increments. We then averaged across neighbouring bins using
a centred rectangular 3% sliding window (1% overlap). The same procedure was applied
to the time course of alpha power lateralization following up-sampling to 125 Hz.
Single-trial measures for the interval of final word presentation were averaged across
the final 35% of sentence presentation as this interval covered final word onset across
all 240 sentence pairs. We used the single-trial neural tracking index as (in-)dependent
variables in our linear mixed-effects analyses (see below).
888

Statistical analysis

We used (generalized) linear mixed-effect models to answer the research questions
outlined in Figure 1. This approach allowed us to jointly model the impact of listening
cues, neural filter strategies and various additional covariates known to influence
behaviour. These included the probed ear (left/right), whether the later-probed
sentence had the earlier onset (yes/no), as well as participants' age and hearing acuity
(pure-tone average across both ears).
To arbitrate between state-level (i.e., within-subject) and trait-level (i.e.,
between-subject) effects, our models included two separate regressors for each of the
key neural measures. Between-subject effect regressors consisted of neural measures
that were averaged across all trials at the single-subject level, whereas the within-

subject effect was modelled by the trial-by-trial deviation from the subject-level mean
[50].

Deviation coding was used for categorical predictors. All continuous variables
were z-scored. For the dependent measure of accuracy, we used generalized linear
mixed-effects model (binomial distribution, logit link function). For response speed, we
used general linear mixed-effects model (gaussian distribution, identity link function).
Given the large sample size, *p*-values for individual model terms are based on Wald *t-as-*
z-values for linear models[105] and on z-values and asymptotic Wald tests in generalized
linear models. All reported p-values are corrected to control for the false discovery
rate[106].

In lieu of a standardized measure of effect size for mixed-effects models, we
report odds ratios (OR) for generalized linear models and standardized regression
coefficients (β) for linear models along with their respective standard errors (SE).

All analyses were performed in R (version 3.6.1)[107] using the packages lme4[108],
and sjPlot[109].

Model selection

To avoid known problems associated with a largely data-driven stepwise model
selection that include the overestimation of coefficients[110] or the selection of irrelevant
predictors[111], the inclusion of fixed effects was largely constrained by our *a priori* defined
hypotheses. The influence of visual cues and of neural measures were tested in same
brain–behaviour model. The brain–behaviour model of accuracy and response speed
included random intercepts by subject and item. In a data-driven manner, we then
tested whether model fit could be further improved by the inclusion of subject-specific
random slopes for the effects of the spatial-attention cue, semantic cue, or probed ear.
The change in model fit was assessed using likelihood ratio tests on nested models.

Data availability

The complete neural and behavioral data required to reproduce the main analyses
supporting this work, as well as the auditory stimuli used in this study are publicly
available in the study's Open Science Framework repository (https://osf.io/28r57/).
Source data are provided with this paper.

Code availability

Code for the main analyses supporting this work is publicly available in the study's Open
Science Framework repository (https://osf.io/28r57/).

References

1.   Cherry, E. C. Some experiments on the recognition of speech, with one and with two ears. *J. Acoust. Soc. Am.* **25,** 975–979 (1953).

2.   Obleser, J. & Erb, J. in *The Cognitive Neurosciences* (eds. Poeppel, D., Mangun, G. S. & Gazzaniga, M. S.) (MIT Press, 2020).

3.   Broadbent, D. E. *Perception and communication*. (Pergamon Press, 1958). doi:10.1186/s12889-020-10047-2

4.   Fernandez-Duque, D. & Johnson, M. L. Attention metaphors: How metaphors guide the cognitive psychology of attention. *Cognitive Science* **23,** 83–116 (1999).

5.   Jensen, O. & Mazaheri, A. Shaping functional architecture by oscillatory alpha activity: gating by inhibition. *Front. Hum. Neurosci.* **4,** 186 (2010).

6.   Foxe, J. J. & Snyder, A. C. The Role of Alpha-Band Brain Oscillations as a Sensory Suppression Mechanism during Selective Attention. *Front. Psychology* **2,** 154 (2011).

7.   Händel, B. F., Haarmeier, T. & Jensen, O. Alpha oscillations correlate with the successful inhibition of unattended stimuli. *J. Cogn. Neurosci.* **23,** 2494–2502 (2011).

8.   Rihs, T. A., Michel, C. M. & Thut, G. Mechanisms of selective inhibition in visual spatial attention are indexed by α-band EEG synchronization. *Eur. J. Neurosci.* **25,** 603–610 (2007).

9.   Kerlin, J. R., Shahin, A. J. & Miller, L. M. Attentional gain control of ongoing cortical speech representations in a 'cocktail party'. *J. Neurosci.* **30,** 620–628 (2010).

10.   Ahveninen, J., Huang, S., Belliveau, J. W., Chang, W.-T. & Hämäläinen, M. Dynamic oscillatory processes governing cued orienting and allocation of auditory attention. *J. Cogn. Neurosci.* **25,** 1926–1943 (2013).

11.   Müller, N. & Weisz, N. Lateralized auditory cortical alpha band activity and interregional connectivity pattern reflect anticipation of target sounds. *Cereb. Cortex* **22,** 1604–1613 (2011).

12.   Wöstmann, M., Herrmann, B., Maess, B. & Obleser, J. Spatiotemporal dynamics of auditory attention synchronize with speech. *Proceedings of the National Academy of Sciences* **113,** 3873–3878 (2016).

13.   Wöstmann, M., Vosskuhl, J., Obleser, J. & Herrmann, C. S. Opposite effects of lateralised transcranial alpha versus gamma stimulation on auditory spatial attention. *Brain Stimulation* **11,** 752–758 (2018).

14.   Tune, S., Wöstmann, M. & Obleser, J. Probing the limits of alpha power lateralisation as a neural marker of selective attention in middle-aged and older listeners. *Eur. J. Neurosci.* **48,** 2537–2550 (2018).

15.   Haegens, S., Handel, B. F. & Jensen, O. Top-Down Controlled Alpha Band Activity in Somatosensory Areas Determines Behavioral Performance in a Discrimination Task. *J. Neurosci.* **31,** 5197–5204 (2011).

16.   Bauer, M., Kennett, S. & Driver, J. Attentional selection of location and modality in vision and touch modulates low-frequency activity in associated sensory cortices. *J. Neurophysiol.* **107,** 2342–2351 (2012).

17.   Worden, M. S., Foxe, J. J., Wang, N. & Simpson, G. V. Anticipatory biasing of visuospatial attention indexed by retinotopically specific alpha-band electroencephalography increases over occipital cortex. *J. Neurosci.* 1–6 (2000). doi:https://doi.org/10.1523/JNEUROSCI.20-06-j0002.2000

18.   Kelly, S. P., Lalor, E. C., Reilly, R. B. & Foxe, J. J. Increases in alpha oscillatory power reflect an active retinotopic mechanism for distracter suppression during sustained visuospatial attention. *J. Neurophysiol.* **95,** 3844–3851 (2006).

19. Schroeder, C. E. & Lakatos, P. Low-frequency neuronal oscillations as
instruments of sensory selection. *Trends Neurosci.* **32,** 9–18 (2009).
20. Schroeder, C. E., Wilson, D. A., Radman, T., Scharfman, H. & Lakatos, P.
Dynamics of Active Sensing and perceptual selection. *Current Opinion in*
Neurobiology **20,** 172–176 (2010).
21. Henry, M. J. & Obleser, J. Frequency modulation entrains slow neural
oscillations and optimizes human listening behavior. *Proc. Natl. Acad. Sci.*
U.S.A. **109,** 20095–20100 (2012).
22. Obleser, J. & Kayser, C. Neural Entrainment and Attentional Selection in the
Listening Brain. *Trends Cogn. Sci.* **23,** 913–926 (2019).
23. Brodbeck, C., Hong, L. E. & Simon, J. Z. Rapid Transformation from Auditory to
Linguistic Representations of Continuous Speech. *Curr. Biol.* **28,** 3976–3982.e6
(2018).
24. Broderick, M. P., Anderson, A. J., Di Liberto, G. M., Crosse, M. J. & Lalor, E. C.
Electrophysiological Correlates of Semantic Dissimilarity Reflect the
Comprehension of Natural, Narrative Speech. *Curr. Biol.* **28,** 1–11 (2018).
25. Zion Golumbic, E. M. Z. *et al.* Mechanisms Underlying Selective Neuronal
Tracking of Attended Speech at a '"Cocktail Party"'. *Neuron* **77,** 980–991
(2013).
26. Ding, N. & Simon, J. Z. Emergence of neural encoding of auditory objects
while listening to competing speakers. *Proc. Natl. Acad. Sci. U.S.A.* **109,** 11854–
11859 (2012).
27. Mesgarani, N. & Chang, E. F. Selective cortical representation of attended
speaker in multi-talker speech perception. *Nature* **485,** 233–236 (2012).
28. O'Sullivan, J. A. *et al.* Attentional Selection in a Cocktail Party Environment Can
Be Decoded from Single-Trial EEG. *Cereb. Cortex* **25,** 1697–1706 (2014).
29. Horton, C., D'Zmura, M. & Srinivasan, R. Suppression of competing speech
through entrainment of cortical oscillations. *J. Neurophysiol.* **109,** 3082–3093
(2013).
30. Keitel, C. *et al.* Stimulus-Driven Brain Rhythms within the Alpha Band: The
Attentional-Modulation Conundrum. *J. Neurosci.* **39,** 3119–3129 (2019).
31. Gundlach, C., Moratti, S., Forschack, N. & Müller, M. M. Spatial Attentional
Selection Modulates Early Visual Stimulus Processing Independently of Visual
Alpha Modulations. *Cereb. Cortex* **16,** 637–18 (2020).
32. Henry, M. J., Herrmann, B., Kunke, D. & Obleser, J. Aging affects the balance of
neural entrainment and top-down neural modulation in the listening brain.
Nat Commun **8,** 15801 (2017).
33. Lakatos, P. *et al.* Global dynamics of selective attention and its lapses in
primary auditory cortex. *Nat. Neurosci.* **19,** 1707–1717 (2016).
34. Posner, M. I. Orienting of attention. *Quarterly Journal of Experimental*
Psychology **32,** 3–25 (1980).
35. Alavash, M., Tune, S. & Obleser, J. Modular reconfiguration of an auditory
control brain network supports adaptive listening behavior. *Proc. Natl. Acad.*
Sci. U.S.A. **116,** 660–669 (2019).
36. Alavash, M., Tune, S. & Obleser, J. Large-scale cortical connectivity dynamics of
intrinsic neural oscillations support adaptive listening behavior. *bioRxiv*
2021.02.22.432315 (2021). doi:10.1101/2021.02.22.432315
37. Presacco, A., Simon, J. Z. & Anderson, S. Effect of informational content of
noise on speech representation in the aging midbrain and cortex. *J.*
Neurophysiol. **116,** 2356–2367 (2016).
38. Sohoglu, E., Peelle, J. E., Carlyon, R. P. & Davis, M. H. Predictive Top-Down
Integration of Prior Knowledge during Speech Perception. *J. Neurosci.* **32,**
8443–8453 (2012).

39. Peelle, J. E., Gross, J. & Davis, M. H. Phase-Locked Responses to Speech in
Human Auditory Cortex are Enhanced During Comprehension. *Cereb. Cortex*
23, 1378–1387 (2013).

40. Obleser, J. & Weisz, N. Suppressed Alpha Oscillations Predict Intelligibility of
Speech and its Acoustic Details. *Cereb. Cortex* **22,** 2466–2477 (2012).

41. Wöstmann, M., Lim, S.-J. & Obleser, J. The human neural alpha response to
speech is a proxy of attentional control. *Cereb. Cortex* **27,** 3307–3317 (2017).

42. Broderick, M. P., Anderson, A. J. & Lalor, E. C. Semantic Context Enhances the
Early Auditory Encoding of Natural Speech. *J. Neurosci.* **39,** 7564–7575 (2019).

43. Broadbent, D. E. & Gregory, M. Accuracy of recognition for speech presented
to the right and left ears. *Quarterly Journal of Experimental Psychology* **16,** 359–
360 (1964).

44. Kimura, D. Cerebral dominance and the perception of verbal stimuli. *Canadian
Journal of Psychology/Revue canadienne de psychologie* **15,** 166–171 (1961).

45. Decruy, L., Vanthornhout, J. & Francart, T. Hearing impairment is associated
with enhanced neural tracking of the speech envelope. *Hearing Research* **393,**
107961 (2020).

46. Presacco, A., Simon, J. Z. & Anderson, S. Speech-in-noise representation in the
aging midbrain and cortex: Effects of hearing loss. *PLoS ONE* **14,** e0213899–26
(2019).

47. Petersen, E. B., Wöstmann, M., Obleser, J. & Lunner, T. Neural tracking of
attended versus ignored speech is differentially affected by hearing loss. *J.
Neurophysiol.* **117,** 18–27 (2017).

48. Bonacci, L. M., Bressler, S. & Shinn-Cunningham, B. G. Nonspatial Features
Reduce the Reliance on Sustained Spatial Auditory Attention. *Ear and Hearing*
41, 1635–1647 (2020).

49. Fiedler, L., Wöstmann, M., Herbst, S. K. & Obleser, J. Late cortical tracking of
ignored speech facilitates neural selectivity in acoustically challenging
conditions. *NeuroImage* **186,** 33–42 (2019).

50. Bell, A., Fairbrother, M. & Jones, K. Fixed and random effects models: making
an informed choice. *Qual Quant* **53,** 1051–1074 (2018).

51. Passow, S. *et al.* Human aging compromises attentional control of auditory
perception. *Psychol Aging* **27,** 99–105 (2012).

52. Anderson, S., White-Schwoch, T., Parbery-Clark, A. & Kraus, N. A dynamic
auditory-cognitive system supports speech-in-noise perception in older
adults. *Hearing Research* **300,** 18–32 (2013).

53. Banerjee, S., Snyder, A. C., Molholm, S. & Foxe, J. J. Oscillatory alpha-band
mechanisms and the deployment of spatial attention to anticipated auditory
and visual target locations: supramodal or sensory-specific control
mechanisms? *J. Neurosci.* **31,** 9923–9932 (2011).

54. Waschke, L., Tune, S. & Obleser, J. Local cortical desynchronization and pupil-
linked arousal differentially shape brain states for optimal sensory
performance. *eLife Sciences* **8,** 1868–27 (2019).

55. Krakauer, J. W., Ghazanfar, A. A., Gomez-Marin, A., MacIver, M. A. & Poeppel, D.
Neuroscience needs behavior: correcting a reductionist bias. *Neuron* **93,** 480–
490 (2017).

56. van Ede, F., Köster, M. & Maris, E. Beyond establishing involvement:
quantifying the contribution of anticipatory α- and β-band suppression to
perceptual improvement with attention. *J. Neurophysiol.* **108,** 2352–2362
(2012).

57. Ding, N. & Simon, J. Z. Cortical entrainment to continuous speech: functional
roles and interpretations. *Front. Hum. Neurosci.* **8,** 13367 (2014).

58.   Hamilton, L. S. & Huth, A. G. The revolution will not be controlled: natural
stimuli in speech neuroscience. *Language, Cognition and Neuroscience* **27,** 1–
10 (2018).

59.   Sassenhagen, J. How to analyse electrophysiological responses to naturalistic
language with time-resolved multiple regression. *Language, Cognition and*
Neuroscience **0,** 1–17 (2018).

60.   Ding, N. & Simon, J. Z. Neural coding of continuous speech in auditory cortex
during monaural and dichotic listening. *J. Neurophysiol.* **107,** 78–89 (2012).

61.   Pernet, C. R., Sajda, P. & Rousselet, G. A. Single-trial analyses: why bother?
Front. Psychology **2,** 322 (2011).

62.   Riecke, L., Formisano, E., Sorger, B., Başkent, D. & Gaudrain, E. Neural
Entrainment to Speech Modulates Speech Intelligibility. *Curr. Biol.* **28,** 1–9
(2018).

63.   Zoefel, B., Archer-Boyd, A. & Davis, M. H. Phase Entrainment of Brain
Oscillations Causally Modulates Neural Responses to Intelligible Speech. *Curr.*
Biol. **28,** 401–408.e5 (2018).

64.   Thut, G. α-band electroencephalographic activity over occipital cortex
indexes visuospatial attention bias and predicts visual target detection. *J.*
Neurosci. **26,** 9494–9502 (2006).

65.   Bengson, J. J., Mangun, G. R. & Mazaheri, A. The neural markers of an
imminent failure of response inhibition. *NeuroImage* **59,** 1534–1539 (2012).

66.   Dahl, M. J., Ilg, L., Li, S.-C., Passow, S. & Werkle-Bergner, M. Diminished pre-
stimulus alpha-lateralization suggests compromised self-initiated attentional
control of auditory processing in old age. *NeuroImage* **197,** 414–424 (2019).

67.   Hong, X., Sun, J., Bengson, J. J., Mangun, G. R. & Tong, S. Normal aging
selectively diminishes alpha lateralization in visual spatial attention.
NeuroImage **106,** 353–363 (2015).

68.   Leenders, M. P., Lozano-Soldevilla, D., Roberts, M. J., Jensen, O. & De Weerd, P.
Diminished Alpha Lateralization During Working Memory but Not During
Attentional Cueing in Older Adults. *Cereb. Cortex* **28,** 21–32 (2018).

69.   Mok, R. M., Myers, N. E., Wallis, G. & Nobre, A. C. Behavioral and neural markers
of flexible attention over working memory in aging. *Cereb. Cortex* **26,** 1831–
1842 (2016).

70.   Wöstmann, M., Alavash, M. & Obleser, J. Alpha Oscillations in the Human Brain
Implement Distractor Suppression Independent of Target Selection. *J.*
Neurosci. **39,** 9797–9805 (2019).

71.   Iemi, L. *et al.* Spontaneous neural oscillations influence behavior and sensory
representations by suppressing neuronal excitability. *bioRxiv*
2021.03.01.433450 (2021). doi:10.1101/2021.03.01.433450

72.   Kloosterman, N. A. *et al.* Humans strategically shift decision bias by flexibly
adjusting sensory evidence accumulation. *eLife Sciences* **8,** (2019).

73.   Hauswald, A., Keitel, A., Chen, Y.-P., Rösch, S. & Weisz, N. Degradation levels of
continuous speech affect neural speech tracking and alpha power differently.
in (John Wiley & Sons, Ltd, 2020). doi:10.1111/ejn.14912

74.   Clayton, M. S., Yeung, N. & Cohen Kadosh, R. The many characters of visual
alpha oscillations. *Eur. J. Neurosci.* **48,** 2498–2508 (2018).

75.   Womelsdorf, T., Valiante, T. A., Sahin, N. T., Miller, K. J. & Tiesinga, P. Dynamic
circuit motifs underlying rhythmic gain control, gating and integration. *Nat.*
Neurosci. **17,** 1031–1039 (2014).

76.   Sadaghiani, S. & Kleinschmidt, A. Brain Networks and α-Oscillations: Structural
and Functional Foundations of Cognitive Control. *Trends Cogn. Sci.* **20,** 805–
817 (2016).

77.   Wöstmann, M. & Obleser, J. Acoustic detail but not predictability of task-
1147   irrelevant speech disrupts working memory. *Front. Hum. Neurosci.* **10,** 201–9
1148   (2016).

1149   78.   van Kerkoerle, T. *et al.* Alpha and gamma oscillations characterize feedback
1150   and feedforward processing in monkey visual cortex. *Proceedings of the*
1151   *National Academy of Sciences* **111,** 14332–14341 (2014).

1152   79.   Spaak, E., Bonnefond, M., Maier, A., Leopold, D. A. & Jensen, O. Layer-Specific
1153   Entrainment of Gamma-Band Neural Activity by the Alpha Rhythm in Monkey
1154   Visual Cortex. *Curr. Biol.* **22,** 2313–2318 (2012).

1155   80.   Chapeton, J. I., Haque, R., Wittig, J. H., Jr, Inati, S. K. & Zaghloul, K. A. Large-
1156   Scale Communication in the Human Brain Is Rhythmically Modulated through
1157   Alpha Coherence. *Curr. Biol.* **29,** 2801–2811.e5 (2019).

1158   81.   Antonov, P. A., Chakravarthi, R. & Andersen, S. K. Too little, too late, and in the
1159   wrong place: Alpha band activity does not reflect an active mechanism of
1160   selective attention. *NeuroImage* **219,** 117006 (2020).

1161   82.   Zhigalov, A., Herring, J. D., Herpers, J., Bergmann, T. O. & Jensen, O. Probing
1162   cortical excitability using rapid frequency tagging. *NeuroImage* **195,** 59–66
1163   (2019).

1164   83.   Teoh, E. S. & Lalor, E. C. EEG decoding of the target speaker in a cocktail party
1165   scenario: considerations regarding dynamic switching of talker location. *J*
1166   *Neural Eng* **16,** 036017–30 (2019).

1167   84.   Schroeder, C. E. & Lakatos, P. Low-frequency neuronal oscillations as
1168   instruments of sensory selection. *Trends Neurosci.* **32,** 9–18 (2009).

1169   85.   Herrmann, B., Henry, M. J., Haegens, S. & Obleser, J. Temporal expectations
1170   and neural amplitude fluctuations in auditory cortex interactively influence
1171   perception. *NeuroImage* **124,** 487–497 (2016).

1172   86.   Pichora-Fuller, M. K. Using the Brain when the Ears are Challenged Helps
1173   Healthy Older Listeners Compensate and Preserve Communication Function.
1174   1–14 (2017).

1175   87.   Rogers, C. S., Payne, L., Maharjan, S., Wingfield, A. & Sekuler, R. Older adults
1176   show impaired modulation of attentional alpha oscillations: Evidence from
1177   dichotic listening. *Psychol Aging* **33,** 246–258 (2018).

1178   88.   Getzmann, S., Klatt, L.-I., Schneider, D., Begau, A. & Wascher, E. EEG correlates
1179   of spatial shifts of attention in a dynamic multi-talker speech perception
1180   scenario in younger and older adults. *Hearing Research* **398,** 108077–46
1181   (2020).

1182   89.   Fuglsang, S. A., Märcher-Rørsted, J., Dau, T. & Hjortkjær, J. Effects of
1183   Sensorineural Hearing Loss on Cortical Synchronization to Competing Speech
1184   during Selective Attention. *J. Neurosci.* **40,** 2562–2572 (2020).

1185   90.   Decruy, L., Vanthornhout, J. & Francart, T. Evidence for enhanced neural
1186   tracking of the speech envelope underlying age-related speech-in-noise
1187   difficulties. *J. Neurophysiol.* **122,** 601–615 (2019).

1188   91.   Presacco, A., Simon, J. Z. & Anderson, S. Evidence of degraded representation
1189   of speech in noise, in the aging midbrain and cortex. *J. Neurophysiol.* **116,**
1190   2346–2355 (2016).

1191   92.   Verschueren, E., Vanthornhout, J. & Francart, T. The Effect of Stimulus Choice
1192   on an EEG-Based Objective Measure of Speech Intelligibility. *Ear and Hearing*
1193   **41,** 1586–1597 (2020).

1194   93.   Waschke, L., Kloosterman, N. A., Obleser, J. & Garrett, D. D. Behavior needs
1195   neural variability. *Neuron* **109,** 751–766 (2021).

1196   94.   Oldfield, R. C. The assessment and analysis of handedness: The Edinburgh
1197   inventory. **9,** 97–113 (1971).

95. Jefferies, K. & Gale, T. M. in *Cognitive Screening Instruments* 209–218 (Springer,
London, 2013). doi:10.1007/978-1-4471-2452-8_11
96. Glasser, M. F. *et al.* A multi-modal parcellation of human cerebral cortex.
Nature **536,** 171–178 (2016).
97. Keitel, A. & Gross, J. Individual Human Brain Areas Can Be Identified from Their
Characteristic Spectral Activation Fingerprints. *PLoS Biol* **14,** e1002498–22
(2016).
98. Smulders, F. T. Y., Oever, ten, S., Donkers, F. C. L., Quaedflieg, C. W. E. M. & van
de Ven, V. Single-trial log transformation is optimal in frequency analysis of
resting EEG alpha. *Eur. J. Neurosci.* **44,** 94–14 (2018).
99. Crosse, M. J., Di Liberto, G. M., Bednar, A. & Lalor, E. C. The Multivariate
Temporal Response Function (mTRF) Toolbox: A MATLAB Toolbox for Relating
Neural Signals to Continuous Stimuli. *Front. Hum. Neurosci.* **10,** 604 (2016).
100. Lalor, E. C. & Foxe, J. J. Neural responses to uninterrupted natural speech can
be extracted with precise temporal resolution. *Eur. J. Neurosci.* **31,** 189–193
(2010).
101. Hoerl, A. E. & Kennard, R. W. Ridge Regression: Biased Estimation for
Nonorthogonal Problems. *Technometrics* **12,** 55–67 (1970).
102. Biesmans, W., Das, N., Francart, T. & Bertrand, A. Auditory-Inspired Speech
Envelope Extraction Methods for Improved EEG-Based Auditory Attention
Detection in a Cocktail Party Scenario. *IEEE Trans Neural Syst Rehabil Eng* **25,**
402–412 (2017).
103. Fiedler, L. *et al.* Single-channel in-ear-EEG detects the focus of auditory
attention to concurrent tone streams and mixed speech. *J Neural Eng* **14,**
036020 (2017).
104. Haufe, S. *et al.* On the interpretation of weight vectors of linear models in
multivariate neuroimaging. *NeuroImage* **87,** 96–110 (2014).
105. Luke, S. G. Evaluating significance in linear mixed-effects models in R. *Behav*
Res **49,** 1494–1502 (2017).
106. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical
and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol*
57, 289–300 (1995).
107. R Core Team. R: A language and environment for statistical computing. (2019).
doi:https://www.R-project.org/
108. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects
Models Using lme4. *J. Stat. Soft.* **67,** 1–48 (2015).
109. Lüdecke, D. & Lüdecke, D. Data Visualization for Statistics in Social Science [R
package sjPlot version 2.6.1]. (2018). doi:10.5281/zenodo.1308157
110. Tibshirani, R. Regression shrinkage and selection via the lasso. *J R Stat Soc*
Series B Stat Methodol **56,** 267–288 (1996).
111. Derksen, S. & Keselman, H. J. Backward, forward and stepwise automated
subset selection algorithms: Frequency of obtaining authentic and noise
variables. *British Journal of Mathematical and Statistical Psychology* **45,** 265–
282 (1992).

1242

1243 **Acknowledgments:** Research was funded by the European Research Council (grant
1244 no. ERC-CoG-2014-646696 "Audadapt" awarded to J.O.). The authors are grateful for
1245 the help of Franziska Scharata in acquiring the data.

1246

1247 **Author contributions:** S.T., M.A., and J.O. designed the experiment; S.T. and M.A.
1248 oversaw data collection and preprocessing of the data; S.T., M.A. and L.F. analysed the
1249 data under supervision by J.O.; S.T. wrote the first draft; all authors contributed to the
1250 final version of the manuscript.

1251

1252 **Competing interests:** The authors declare no competing interests.