High-resolution population-specific recombination rates and their effect on phasing and genotype imputation

Running Title: Population-specific recombination maps in phasing & imputation

Shabbeer Hassan, *Institute for Molecular Medicine Finland, FIMM, HiLIFE, University of Helsinki, Helsinki, Finland*

Ida Surakka, *Institute for Molecular Medicine Finland, FIMM, HiLIFE, University of Helsinki, Helsinki, Finland*

Marja-Riitta Taskinen, *Clinical and molecular metabolism, Research program unit, University of Helsinki, Helsinki, Finland*

Veikko Salomaa, *Finnish Institute for Health and Welfare, Helsinki, Finland*

Aarno Palotie, *Institute for Molecular Medicine Finland, FIMM, HiLIFE, University of Helsinki, Helsinki, Finland, Psychiatric & Neurodevelopmental Genetics Unit, Department of Psychiatry, Analytic and Translational Genetics Unit, Department of Medicine, Department of Neurology, Massachusetts General Hospital, Boston, MA, USA*

Maija Wessman, *Institute for Molecular Medicine Finland, FIMM, HiLIFE, University of Helsinki, Helsinki, Finland*

Taru Tukiainen, *Institute for Molecular Medicine Finland, FIMM, HiLIFE, University of Helsinki, Helsinki, Finland*

Matti Pirinen, *Institute for Molecular Medicine Finland, FIMM, HiLIFE, University of Helsinki, Helsinki, Finland, Public Health, Clinicum, University of Helsinki, Helsinki, Finland, Department of Mathematics and Statistics, University of Helsinki, Helsinki, Finland*

Priit Palta, *Institute for Molecular Medicine Finland, FIMM, HiLIFE, University of Helsinki, Helsinki, Finland , Estonian Genome Center, Institute of Genomics, University of Tartu, Tartu, Estonia*

Samuli Ripatti, *Institute for Molecular Medicine Finland, FIMM, HiLIFE, University of Helsinki, Helsinki, Finland, Broad Institute of the Massachusetts Institute of Technology and Harvard University, Cambridge, MA, USA, Public Health, Clinicum, University of Helsinki, Helsinki, Finland*

CORRESPONDING AUTHOR

Samuli Ripatti, **Email**: samuli.ripatti@helsinki.fi

Abstract:

Founder population size, demographic changes (eg. population bottlenecks or rapid

expansion) can lead to variation in recombination rates across different populations.

Previous research has shown that using population-specific reference panels has a

significant effect on downstream population genomic analysis like haplotype phasing,

genotype imputation and association, especially in the context of population isolates.

Here, we developed a high-resolution recombination rate mapping at 10kb and 50kb

scale using high-coverage (20-30x) whole-genome sequenced 55 family trios from

Finland and compared it to recombination rates of non-Finnish Europeans (NFE). We

tested the downstream effects of the population-specific recombination rates in

statistical phasing and genotype imputation in Finns as compared to the same analyses

performed by using the NFE-based recombination rates. We found that Finnish

recombination rates have a moderately high correlation (Spearman's ρ =0.67-0.79) with

NFE, although on average (across all autosomal chromosomes), Finnish rates

(2.268±0.4209 cM/Mb) are 12-14% lower than NFE (2.641±0.5032 cM/Mb). Finnish

recombination map was found to have no significant effect in haplotype phasing

accuracy (switch error rates ~ 2%) and average imputation concordance rates (97-98%

for common, 92-96% for low frequency and 78-90% for rare variants). Our results

suggest that downstream population genomic analyses like haplotype phasing and

genotype imputation mostly depend on population-specific contexts like appropriate

reference panels and their sample size, but not on population-specific recombination

maps or effective population sizes. Currently, available HapMap recombination maps

seem robust for population-specific phasing and imputation pipelines, even in the

context of relatively isolated populations like Finland.

Keywords: recombination, phasing, imputation, Finland, population genomics

1. Introduction:

Recombination is not uniform across the human genome with large areas having lower

recombination rates, so-called 'coldspots', which are then interspersed by shorter

regions marked by a high recombinational activity called 'hotspots' [1]. With long

chunks of human genome existing in high linkage disequilibrium, LD [2], and organised

in the form of 'haplotype blocks', the 'coldspots' tend to coincide with such regions of

high LD [3].

Direct estimation methods of recombination are quite time-consuming, and evidence

has suggested that they do not easily scale up to genome-wide, fine-scale

recombinational variation estimation [4]. A less time-consuming but computationally

intensive alternative is to use the LD patterns surrounding the SNPs [5]. Such methods

have been used in the past decade or so, to create fine-scale recombination maps [6].

Besides the International HapMap project that focused on capturing common variants

and haplotypes in diverse populations, international WGS-based collaborations like

1000 Genomes Project, provided genetic variation data for 20 worldwide populations

[7]. This led to further refinement of the recombination maps coupled with

methodological advances of using coalescent methods for recombination rate [8, 9].

With the rise of international collaborative projects, it was realised that founder

populations can often have very unique LD patterns [10], subsequently also displaying

unique increased genetics-driven health risks [11], suggesting that population-specific

reference datasets should be used to leverage the LD patterns to better detect disease

variants in downstream genetic analysis   [12].   Genomic analysis methods like

haplotype phasing and imputing genotypes require recombination maps and other

population genetic parameters as input to obtain optimal results [13, 14, 15, 16]

In theis study, we set to test this by 1) estimating recombination rates along the genome

in Finnish population using ~55 families of whole-genome sequenced (20-30x) Finns,

2) comparing these rates to some other European populations, and 3) comparing the

effect of using Finnish recombination rate estimates and cosmopolitan estimates in

phasing and imputation errors in Finnish samples.

2. Materials & Methods:

2.1 Datasets used:

Finnish Migraine Families Collection

Whole-genome sequenced trios (n = 55) consisting of the parent-offspring combination

were drawn from a large Finnish migraine families collection consisting of 1,589

families totalling 8,319 individuals [17]. The trios were used for the recombination map

construction using LDHAT version 2. The families were collected over 25 years from

various headache clinics in Finland (Helsinki, Turku, Jyväskylä, Tampere, Kemi, and

Kuopio) and via advertisements in the national migraine patient organisation web page

(https://migreeni.org/). The families consist of different pedigree sizes from small to

large (1-5+ individuals). Of the 8319 individuals, 5317 have a confirmed migraine

diagnosis based on the third edition of the established International Classification for

Headache Disorders (ICHD-3) criteria [18].

EUFAM cohort

To check the phasing accuracy of our Finnish recombination map, we used an

independently sourced 49 trios from the European Multicenter Study on Familial

Dyslipidemias in Patients with Premature Coronary Heart Disease (EUFAM). Finnish

familial combined hyperlipidemia (FCH) families were identified from patients initially

admitted to hospitals with premature cardiovascular heart disease (CHD) diagnosis who

also had elevated levels of total cholesterol (TC), triglycerides (TG) or both in the $\geq$

90th Finnish population percentile. Those families who had at least one additional first-

degree relative also affected with hyperlipidemia were also included in the study apart

from individuals with elevated levels of TG. [19, 20, 21].

FINRISK cohort

The imputation accuracy of the Finnish and previously published HapMap based

recombination maps   [8, 9] was subsequently tested on an independent FINRISK

CoreExome chip dataset consisting of 10,481 individuals derived from the national-

level FINRISK cohort. Primarily, it comprises of respondents of representative, cross-

sectional population surveys that are conducted once every 5 years since 1972 to get a

national assessment of various risk factors of chronic diseases and other health

behaviours among the working-age population drawn from 3 to 4 major cities in

Finland [22].

FINNISH reference panel cohort

The whole-genome sequenced samples used were obtained from PCR-free methods and

PCR-amplified methods, which was followed by sequencing on a Illumina HiSeq X

platform with a mean depth of ~30×. The obtained reads were then aligned to the

GRCh37 (hg19) human reference genome assembly using BWA-MEM. Best practice

guidelines from Genome Analysis Toolkit (GATK) were used to process the BAM files

and variant calling. Several criteria were used in this stage for sample exclusion:

relatedness (identity-by-descent (IBD) > 0.1), sex mismatches, among several others.

Furthermore, samples were filtered based on other criteria such as: non-reference

variants, singletons, heterozygous/homozygous variants ratio, insertion/deletion ratio

for novel indels, insertion/deletion ratio for indels observed in dbSNP, and

transition/transversion ratio.

After this stage, some exclusion criteria were applied to set some variants as missing:

GQ < 20, phred-scaled genotype likelihood of reference allele < 20 for heterozygous

and homozygous variant calls, and allele balance <0.2 or >0.8 for heterozygous calls. A

truth sensitivity percentage threshold of 99.8% for SNVs and of 99.9% for indels was

used based on the GATK Variant Quality Score Recalibration (VQSR) to filter variants

with, quality by depth (QoD) < 2 for SNVs and < 3 for indels, call rate < 90%, and

Hardy-Weinberg equilibrium (HWE) p-value < 1×10-9. Some other variants like

monomorphic, multi-allelic and low-complexity regions [23] were further excluded.

The final reference dataset used in this study for imputation consisted of high coverage

(20-30x) whole-genome sequence-based reference panel of 2690 individuals from the

SISu project (Sequencing Initiative Suomi, http://www.sisuproject.fi/, [24]).

2.2 Recombination map construction:

Coalescent-based fine-scale recombination map construction [8] is greatly eased by

using trios which provide more accurate haplotype phasing resolution [25]. Hence, we

used trio data (n=55, 110 independent parents) from the Finnish Migraine Families

Cohort described above. These were filtered primarily using VCFtools [26] and custom

R scripts. Firstly, sites were thinned with within 15bp of each other such that only one

site remained followed by a filtering step of removing variants with a minor allele

frequency of <5%   [27]. The resultant data were then phased using family-aware

method of SHAPEIT [28] using the standard HapMap recombination map [8, 9],

which was then split into segments of ~10000 SNPs with a 1000 SNP overhang on each

side of the segments. LDhat version 2 was run for 10^7 iterations with a block penalty of

5, every 5000 iterations of them of which the first 10% observations were discarded [8,

29]. The CEU based maps, used here for comparison, were obtained similarly using

LDhat [29].

However, LDHat is computationally intensive, and calculations suggest that the 1000

Genomes OMNI data set [30] would be too much computationally intensive to

complete [31], hence limiting the maximum number of haplotypes which could be

used.

To overcome this and make the recombination map independent of the underlying

methodology, we used a machine learning method implemented in FastEPRR [31, 32].

It supports the use of larger sample sizes, than LDHat and the recombination estimates

for sample sizes > 50, yields smaller variance than LDHat based estimates [31]. The

method was then applied to each autosome with overlapping sliding windows (*i.e.*,

window size, 50 kb and step length, 25 kb) under default settings for diploid organisms.

As seen in [31] both methods produce similar estimates, with only variance of the

estimate of mean being different.

The output of LDHat and FastEPRR is in terms of population recombination rate (p)

and to convert them into per-generational rate (r) used in phasing/imputation algorithms

we used optimal effective population size values derived from our testing (as explained

in the Supplementary Text). The estimates from LDHat and FastEPRR were then

averaged, to obtain a new combined estimate with the lowest variance amongst all the

three [31].

2.3 Phasing and imputation accuracy

To test whether the usage of different recombination maps affects the efficiency of

haplotype phasing and imputation , we used the aforesaid Finnish genotype data to

evaluate: (i) switch error rates across all chromosomes and (ii) imputation concordance

rates for chromosome 20.

2.3.1 Phasing Accuracy

The gold standard method to estimate haplotype phasing accuracy is to count the

number of switches (or recombination events) needed between the computationally

phased dataset and the true haplotypes [33].The number of such switches divided by

the number of all possible switches is called switch error rate (SER).

For testing the influence of recombination maps on phasing accuracy, we used three

different recombination maps: HapMap, fine-scale Finnish recombination map and a

constant background recombination rate (1cM/Mb), to phase the 55 offspring

haplotypes without using any reference dataset. To check whether reference panels used

during haplotype phasing made any impact on the switch error rates, we used the

Finnish SISU based reference (n=2690), to check whether the size of the reference

panel made any impact on the results in phasing the offspring's haplotypes (Figure 1).

The SER in the offspring's phased haplotypes were then calculated by determining the

true offspring haplotypes using data from the parents (98 individuals) with a custom

script [34].

2.3.2 Imputation Accuracy

Imputation concordance was used as the metric for calculating the imputation accuracy.

For this, we randomly masked FINRISK CoreExome chip data consisting of 10,480

individuals [22] from chromosome 20. To test the role of reference panel size in

influencing the imputation accuracy in conjunction with varying the population genetics

parameters, we imputed the masked dataset with BEAGLE (Browning *et al.*, 2016)

using the Finnish reference panel (n = 2690). The concordance was then calculated

between the imputed genotypes and the original masked variants. Masking was done by

randomly removing ~10% of variants from the chip dataset.

The influence of recombination maps on imputation accuracy was checked by

calculating the concordance values between imputed and original variants, using the

Finnish reference panel in various combinations of recombination maps (constant rate,

HapMap, Finnish map) during the imputation (Figure 1).

3. Results:

3.1 Finnish recombination map and its comparison to the HapMap recombination

map:

The primary aim of our study was to derive a high-resolution genetic recombination

map for Finland and use it for comparative tests in commonly used analyses like

haplotype phasing and imputation. To derive a population-specific Finnish

recombination map, we used the high-coverage WGS data and an average of different

estimation methods (LDHat and FastEPRR). We used the Ne value of 10,000 derived

from our extensive testing of different Ne values (See supplementary text) to get the

per-generation recombination rates. The average recombination rates of Finnish

population isolate depicted 12-14% lower values (autosome-wide average 2.268±0.4209

cM/Mb) for all chromosomes compared to CEU based maps (2.641±0.5032 cM/Mb)

(Figure 2).

These differences in average recombination rates are reflected in the correlation values

across all chromosomes (Spearman's ρ ~ 0.67 - 0.79) between the developed Finnish

map and HapMap based one (Figure 2). We also present a direct comparison between

the two maps, of the recombination rates at 5Mb scales, which presents a similar visual

pattern of rates across the genome (Supplementary Figure 1).

3.2 Effects of the population-specific recombinations map on haplotype phasing

Variation in population-specific recombination maps (and effective population sizes)

can affect the downstream genomic analyses like haplotype phasing and imputation.

We tested the Finnish map, HapMap map and a constant recombination rate map

(1cM/Mb) to understand the effects of population-specific maps on downstream

genomic analyses. The phasing accuracy was tested under two different conditions:

using no additional reference panel and using an population-specific . SISu v2 reference

panel (n= 2690) in phasing. We observed that, on average, SER ranged between 1.8-

3.7% (Supplementary Figure 2) across the different chromosomes and recombination

maps. We found statistically significant differences within both no-reference panel and

the Finnish reference panel results (Kruskal Wallis, p-value = 5.3e-10 and 4.7e-10,

respectively; Figure 3). The constant recombination map (1cM/Mb) had significantly

higher SER values when compared to the Finnish map or the HapMap map (Figure 3)

both when no reference panels were used (p-value = 2.9e-11 and 2.6e-09, respectively)

and when the Finnish reference panel was used (p-value = 2.9e-11 and 9.5e-13,

respectively). The choice of recombination maps mattered more when no reference

panel was used (p-value = 0.0046), however when using the Finnish reference panel, the

difference in SER was statistically insignificant (p-value = 0.25).

3.3 Effects of the population-specific recombinations map on genotype imputation

Imputation accuracy was similarly tested using the reference panel under three different

recombination map settings. We observed that when the imputation target dataset was

phased and imputed using the Finnish reference panel (n=2690) irrespective of the

population-specific recombination maps, it had a high imputation accuracy (overall

concordance rate ~98%, Figure 4) across MAF bins (>0.1%). Though some differences

in concordance rates are seen in for rare variants (MAF <0.1%). The concordance rate

was lower when the test dataset was phased without reference panels (concordance rate

72~77%, Figure 5).

4. Discussion:

Population isolates like Finland, have had a divergent demographic history as compared

to the outbred European populations, with a less historic migration, more fluctuating

population sizes and higher incidences of bottleneck events and founder effects [35, 36]

This unique demographic history then affects different population genetic parameters,

like recombination rates [37]. It has been shown previously that using population-

specific genomic reference panels augmented the accuracy of imputation accuracy

leading to better mapping of diseases specific variants in GWAS [12]. Since

recombination rates (in the form of recombination maps), features in much of the

downstream genomic analyses' methods like imputation and haplotype phasing [15,

34], we wanted to study their effect on downstream analyses.

Firstly, we characterised the Finnish recombination map using high-coverage (~30x)

whole-genome sequencing (WGS) samples from large SISu v2 reference panel

(n=2690). Previously used recombination maps hail from the HapMap and

1000Genomes projects which used sparse genotypic datasets or low-depth sequencing

samples. This is a first attempt in creating a recombination map for Finland using

population-specificWGS samples. We used two different methods in estimating the

recombination rates, to achieve accurate estimates with lower variance [29,31]. In

addition, we estimated effective population sizes using identity-by-descent (IBD) based

methods [15] for both Finnish and CEU based datasets. The obtained recombination

map was then used to test their role and importance in two selected downstream

genomic analyses – haplotype phasing and imputation concordance. Since the

recombination rate determination requires effective population size estimates, we also

tested the role of varying effective population size on these two analyses (See

Supplementary Text). The extensive testing of Ne yielded the estimate of 10,000

originally derived theoretically [38] and most used commonly for humans fits quite

rightly for the recombination map.

The Finnish recombinational landscape when compared to the HapMap based map,

showed, on average, a high degree of correlation across scales (10, 50kb and 5Mb),

however, on average, Finnish recombination rates across chromosomes were found to

be lower. Such moderate to high correlations (Figure 2) and similar recombinational

landscape (Supplementary Figure 1) could be due to high sharing of recombinations in

individuals from closely-related populations. The degree of dissimilarity in the

population-level differences between Finnish and mainland Europeans in terms of

recombination rates could be due to population-specific demographic processes like

founder effects, bottleneck events and migration [39], or chromatin structure PRDM9

binding locations, for example [40]. And the broad similarity in terms of correlational

structure seen here, reflects a shared ancestral origin of Finns and other mainland

Europeans [41]. Other studies on population isolates like Iceland [9] have previously

found a high degree of correlation with CEU based maps, albeit with substantial

differences as seen here. Previous studies [42] have additionally explored the

relationship between recombination rate differences between populations and allele

frequency differences, with evidence suggesting that the differences between rates show

the selection impact in the past 100,000 years since the out-of-Africa movement of

humans.

As seen in previous studies, much of the downstream genomic analyses like getting

more refined GWAS hits or, accurate copy number variants (CNV) imputation, can be

highly improved with the addition/use of population-specific datasets [12]. To test this

in the context of population-specific recombination maps, we used them to test the

haplotype phasing and imputation accuracy and observed that despite large differences

in the effective population sizes between populations, it did not affect the tested metrics.

One possible explanation for the insignificant effect seen here is that the role of

parameters like effective population size and recombination maps is to scale over the

haplotypes for efficient coverage of the whole genome. However, when sufficiently

large, population-specific genomic reference panels are available with tens of thousands

of haplotypic combinations, such scaling over for specific populations, does not yield

in substantial improvements. As we showed here, reference panel size could play an

important role in the downstream genomic analyses and in most cases, the current

practice of using the standard HapMap recombination map can be reasonably used.

Another point of interest here is that the use of different Ne parameters during

phasing/imputation might be redundant as we observed no change in the accuracy of our

estimates on varying the Ne parameters. Similarly, when using population-specific

recombination maps, we did not find any tangible benefits in using them over the

current standard maps based on the HapMap data.

Our study suggests a couple of important points for future studies: (a) varying effective

population size for downstream genomic analyses, such as phasing and imputation,

might have a relatively small impact, and it might be better to use the default option of

the particular software; (b) when available, it is beneficial to use a population-specific

genomic reference panel as they increase the accuracy; (c) HapMap can be used for

current downstream genomic analyses like haplotype phasing or genotype imputation in

European-based populations. And, if need be, can be substituted for using population-

specific maps, as the accuracy rates are quite similar to the population-based maps.

Though the sample used here is from a disease cohort but is nevertheless representative

of Finland's population and hence provides a reasonable recombination rate estimates.

On the other hand, our reliance on disease cohorts could lead to minor variation in the

resultant recombination. Though as we share a similar out-of-Africa origin, much of our

history is shared and though biological differences in the recombinational landscape do

exist between different populations, much of the downstream genomic analyses

(haplotyping, imputation or, GWAS), might not be affected by recombination map or

values of effective population size.

Funding

This work was financially supported by the Academy of Finland (251217 and 255847 to

S.R.). S.R. was further supported by the Academy of Finland, Center of Excellence for

Complex Disease Genetics, the Finnish Foundation for Cardiovascular Research,

Biocentrum Helsinki, and the Sigrid Jusélius Foundation. S.H. was supported by

FIMM-EMBL PhD program doctoral funding and I.S. by Academy of Finland

Postdoctoral Fellowship (298149). V.S. was supported by the Finnish Foundation for

Cardiovascular Research. T.T. was supported by Academy of Finland grant number

315589.

Acknowledgements

We would like to thank Sari Kivikko and Huei-Yi Shen for management assistance. The FINRISK analyses were conducted using the THL biobank permission for project BB2015_55.1. We thank all study participants for their generous participation in the FINRISK study.

Conflict of Interest: VS has received honoraria from Novo Nordisk and Sanofi for consulting and has ongoing research collaboration with Bayer ltd (all unrelated to the present study).

References

1. Baudat F, Buard J, Grey C, Fledel-Alon A, Ober C, Przeworski M et al. PRDM9 is a major determinant of meiotic recombination hotspots in humans and mice. Science 2009; 327: 836–840

2. Daly MJ, Rioux JD, Schaffner SF, Hudson TJ, Lander ES. High-resolution haplotype structure in the human genome. Nature Genetics 2001; 29: 229–232

3. Hudson RR, Kaplan NL. Statistical properties of the number of recombination events in the history of a sample of DNA sequences. Genetics 1985; 111: 147-164

4. Chan AH, Jenkins PA, Song YS. Genome-wide fine-scale recombination rate variation in Drosophila melanogaster. PLoS Genet 2012; 8: e1003090

5. McVean GA, Myers SR, Hunt S, Deloukas P, Bentley DR, Donnelly P. The fine-scale structure of recombination rate variation in the human genome. Science 2004; 304: 581-584

6. Myers S, Bottolo L, Freeman C, McVean G, Donnelly P. A fine-scale map of

recombination rates and hotspots across the human genome. Science 2005; 310:

321-324.

7. Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO et al. A

global reference for human genetic variation. Nature 2015; 526: 68-74

8. Auton A, McVean G. Recombination rate estimation in the presence of hotspots.

Genome Res 2007; 17: 1219-1227.

9. Kong A, Thorleifsson G, Gudbjartsson DF, Masson G, Sigurdsson A,

Jonasdottir A et al. Fine-scale recombination rate differences between sexes,

populations and individuals. Nature 2010; 467: 1099-1103.

10. Service S, DeYoung J, Karayiorgou M, Roos JL, Pretorious H, Bedoya G et al.

Magnitude and distribution of linkage disequilibrium in population isolates and

implications for genome-wide association studies. Nat Genet 2006; 38: 556-560.

11. Peltonen L, Jalanko A, Varilo T. Molecular genetics of the Finnish disease

heritage. Hum Mol Genet 1999; 8: 1913-1923.

12. Surakka I, Kristiansson K, Anttila V, Inouye M, Barnes C, Moutsianas L et al.

Founder population-specific HapMap panel increases power in GWA studies

through improved imputation accuracy and CNV tagging. Genome Res 2010;

20: 1344-1351.

13. Tewhey R, Bansal V, Torkamani A, Topol EJ, Schork NJ. The importance of

phase information for human genomics. Nat Rev Genet 2011; 12: 215-223.

14. Browning SR, Browning BL. Haplotype phasing: existing methods and new

developments. Nat Rev Genet 2011; 12: 703-714.

15. Browning BL, Browning SR. Genotype Imputation with Millions of Reference
Samples. Am J Hum Genet 2016; 98: 116-126.

16. Delaneau O, Zagury JF, Marchini J. Improved whole-chromosome phasing for
disease and population genetic studies. Nat Methods 2013; 10: 5-6.

17. Gormley P, Kurki MI, Hiekkala ME, Veerapen K, Häppölä P, Mitchell AA et al.
Common Variant Burden Contributes to the Familial Aggregation of Migraine
in 1,589 Families. Neuron 2018; 98: 743-753.e4.

18. The International Classification of Headache Disorders, 3rd edition (beta
version). Cephalalgia 2013; 33: 629-808.

19. Borodulin K, Vartiainen E, Peltonen M, Jousilahti P, Juolevi A, Laatikainen T et
al. Forty-year trends in cardiovascular risk factors in Finland. Eur J Public
Health 2015; 25: 539-546.

20. Porkka KV, Nuotio I, Pajukanta P, Ehnholm C, Suurinkeroinen L, Syvänne M et
al. Phenotype expression in familial combined hyperlipidemia. Atherosclerosis
1997; 133: 245-253.

21. Ripatti P, Rämö JT, Söderlund S, Surakka I, Matikainen N, Pirinen M et al. The
Contribution of GWAS Loci in Familial Dyslipidemias. PLOS Genetics 2016;
12: e1006078.

22. Vartiainen E, Laatikainen T, Peltonen M, Juolevi A, Mannisto S, Sundvall J et
al. Thirty-five-year trends in cardiovascular risk factors in Finland. International
Journal of Epidemiology 2009; 39: 504–518.

23. Li H. Toward better understanding of artifacts in variant calling from high-
coverage samples. Bioinformatics 2014; 30: 2843–2851.

24. Mart Kals, Tiit Nikopensius, Kristi Läll, Kalle Pärn, Timo Tõnis Sikka, Jaana

Suvisaari, Veikko Salomaa, Samuli Ripatti, Aarno Palotie, Andres Metspalu,

Tõnu Esko, Priit Palta, Reedik Mägi Advantages of genotype imputation with

ethnically matched reference panel for rare variant association analyses bioRxiv

579201; doi: https://doi.org/10.1101/579201

25. Roach JC, Glusman G, Hubley R, Montsaroff SZ, Holloway AK, Mauldin DE et

al. Chromosomal haplotypes by genetic phasing of human families. Am J Hum

Genet 2011; 89: 382-397.

26. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA et al. The

variant call format and VCFtools. Bioinformatics 2011; 27: 2156-2158.

27. Stevison LS, Woerner AE, Kidd JM, Kelley JL, Veeramah KR, McManus KF et

al. The Time Scale of Recombination Rate Evolution in Great Apes. Mol Biol

Evol 2016; 33: 928-945.

28. O'Connell J, Gurdasani D, Delaneau O, Pirastu N, Ulivi S, Cocca M et al. A

general approach for haplotype phasing across the full spectrum of relatedness.

PLoS Genet 2014; 10: e1004234.

29. Auton A, Fledel-Alon A, Pfeifer S, Venn O, Ségurel L, Street T et al. A fine-

scale chimpanzee genetic map from population sequencing. Science 2012; 336:

193-198.

30. Abecasis GR, Auton A, Brooks LD, DePristo MA, Durbin RM, Handsaker RE

et al. An integrated map of genetic variation from 1,092 human genomes. Nature

2012; 491: 56-65.

31. Gao F, Ming C, Hu W, Li H. New Software for the Fast Estimation of Population Recombination Rates (FastEPRR) in the Genomic Era. G3 (Bethesda) 2016; 6: 1563-1571.

32. Lin K, Futschik A, Li H. A fast estimate for the population recombination rate based on regression. Genetics 2013; 194: 473-484.

33. Bansal V. Integrating read-based and population-based phasing for dense and accurate haplotyping of individual genomes. Bioinformatics 2019; 35: i242-i248.

34. Loh PR, Danecek P, Palamara PF, Fuchsberger C, A Reshef Y, K Finucane H et al. Reference-based phasing using the Haplotype Reference Consortium panel. Nat Genet 2016; 48: 1443-1448.

35. Martin AR, Karczewski KJ, Kerminen S, Kurki MI, Sarin AP, Artomov M et al. Haplotype Sharing Provides Insights into Fine-Scale Population History and Disease in Finland. Am J Hum Genet 2018; 102: 760-775.

36. Kerminen S, Havulinna AS, Hellenthal G, Martin AR, Sarin AP, Perola M et al. Fine-Scale Genetic Structure in Finland. G3 (Bethesda) 2017; 7: 3459-3468.

37. Wang J, Santiago E, Caballero A. Prediction and estimation of effective population size. Heredity (Edinb) 2016; 117: 193-206.

38. Takahata N, Satta Y, Klein J. Divergence time and population size in the lineage leading to modern humans. Theor Popul Biol 1995; 48: 198-221.

39. Novembre J, Johnson T, Bryc K, Kutalik Z, Boyko AR, Auton A et al. Genes mirror geography within Europe. Nature 2008; 456: 98-101.

40. Ségurel L. The complex binding of PRDM9. Genome Biol 2013; 14: 112.

41. Mallick S, Li H, Lipson M, Mathieson I, Gymrek M, Racimo F et al. The

Simons Genome Diversity Project: 300 genomes from 142 diverse populations.

Nature 2016; 538: 201-206.

42. Keinan A, Reich D. Human population differentiation is strongly correlated with

local recombination rate. PLoS Genet 2010; 6: e1000886.

Figure 1: Flowchart overview of the analyses and comparisons performed.

Figure 2: Average (± standard deviation) recombination rates of Finnish v/s CEU per

autosome measured in cM/Mb and Correlation between Finnish and CEU

recombination rates across all chromosomes. The comparisons are made for similar

physical positions.

Figure 3: Statistical comparison of Switch Error Rates across all autosomes calculated

for all children in the trios using different recombination maps with respect to different

reference panel conditions (absent or present). The p-values are shown at the top of each

panel from Kruskal Wallis ANOVA testing between panel groups and ones between

boxplots for within-group comparisons.

Figure 4: Comparison of Imputation Concordance across different Minor Allele

Frequency (MAF) groups for a range of different recombination map combinations

phased with NO reference panels

Figure 5: Comparison of Imputation Concordance across different Minor Allele

Frequency (MAF) groups for a range of different recombination map combinations

phased with reference panels.

With RefPanels(Phasing) + SISU_2700(Imputation)

No RefPanels(Phasing) + SISU_2700(Imputation)