No evidence for dominance-discovery tradeoffs in *Pheidole* (Hymenoptera: Formicidae) assemblages

CRISTIAN L. KLUNK[1] and MARCIO R. PIE[1,2]

[1]Graduate Program in Ecology and Conservation, Universidade Federal do Paraná, Curitiba-PR, Brazil. [2]Department of Zoology, Universidade Federal do Paraná, Curitiba-PR, Brazil.

Corresponding author: klunkcristian@gmail.com

Abstract

Understanding the mechanisms that allow species coexistence across spatial scales is of great interest to

ecologists. Many such proposed mechanisms involve tradeoffs between species in different life-history traits, with

distinct tradeoffs being expected to be prevalent at varying temporal and spatial scales. The dominance-discovery

tradeoff posits that species differ in their ability to find and use resources quickly, in contraposition to their ability to

monopolize those resources, a mechanism analogous to the competition-colonization tradeoff. We investigated the

occurrence of this structuring mechanism in *Pheidole* (Hymenoptera: Formicidae) assemblages in Atlantic Forest

remnants. According to the dominance-discovery tradeoff, we should observe a consistent interspecific variation along

the axis of discovery and dominance. We established 55 sampling units across two sites, with each unit consisting of

a sardine bait monitored for three hours. There was no distinction among *Pheidole* species in their ability to find or

dominate food sources, suggesting that the dominance-discovery tradeoff does not explain their coexistence. The low

levels of aggression between *Pheidole* species could prevent the establishment of dominance hierarchies, whereas the

species order of arrival at food sources could allow for resource partitioning through priority effects.

Keywords: Aggressiveness; Ants; Dominance-discovery tradeoff; Priority effects; Species interactions.

Introduction

Species coexistence is a fundamental theme in community ecology. How some regions are able to harbor high levels of diversity is still far from being understood, despite a long history of investigation (Hutchinson 1959). Also challenging is to understand how closely related species, which share many aspects of their niches, are able to coexist at fine spatial scales (Hutchinson 1961). Many ecological mechanisms proposed to date to explain species coexistence consider the occurrence of tradeoffs in the performance of species in some life-history traits, which affect their spatial distribution and interspecific interactions (Kneitel and Chase 2004). Species can differ in their competitive and colonization abilities, the levels of abiotic stresses and predation pressures they tolerate, their degree of habitat specialization, among other characteristics, and those differences could explain species coexistence across distinct spatial scales (Kneitel and Chase 2004). Given that diversity translates to ecosystem functioning, the study of species coexistence is essential to understand and manage ecosystem services.

The competition-colonization tradeoff is one of the most classical tradeoff mechanisms proposed to explain species coexistence by suggesting that some species are better at dispersing and colonizing new habitat patches, whereas others are better at displacing incumbent species once they finally colonize the patch (Kneitel and Chase 2004). Although the spatial scale considered under this mechanism can vary, that tradeoff purports to explain coexistence in long temporal scales. Likewise, the competition-colonization tradeoff inspired the proposition of an analogous mechanism that operates in a reduced time range, the dominance-discovery tradeoff (Parr and Gibb 2012). This mechanism proposes that species can differ in their ability to find and dominate resources, and was suggested to explain the coexistence of bees (Hubbell and Johnson 1978; Nagamitsu and Inoue 1997), rodents (Brown et al. 1994) and ants (Fellers 1987; Lebrun and Feener 2007; Feener et al. 2008; Perfecto and Vandermeer 2001; Bertelsmeier et al. 2015*a*).

Ants are important components of terrestrial ecosystems worldwide, particularly in the tropics, where they account for a substantial portion of the animal biomass (Fittkau and Klinge 1973), show considerable species diversity (Economo et al. 2018) and are essential to sustain many ecosystem services (Del Toro et al. 2012). Most ant species are food generalists (Kaspari 2001), because of the marked differences in brood and adult nutritional requirements (Blüthgen and Feldaar 2010). Therefore, if several ant species co-occur and share common resources, one would

expect that they might show some level of competition, which suggests the existence of mechanisms of coexistence

to sustain high levels of diversity. However, the role of competition in ant community structure is far from settled

(Parr and Gibb 2010), despite having traditionally been considered as a "hallmark" of ant communities (Hölldobler

and Wilson 1990). Competition in ant communities can be manifested in two ways. Through interference, some ant

species exclude competitors from resources by direct aggression or territoriality, whereas by exploitative behaviors

some species are specialized in finding and removing resources before other species arrive (Davidson 1998). This

leads to dominance hierarchies, in which some species (dominants) are capable of monopolizing resources, whereas

others are better at finding resources quickly and removing them before the arrival of the dominants (subordinates)

(Fellers 1987; Feener et al. 2008; Arnan et al. 2012; Yitbarek and Philpott 2019).

Ant species are considered as dominant, in the context of those hierarchies, either by the outcomes of pairwise

interactions with other species, where the winners are considered dominants (behavioral dominance), or by a large

recruitment of workers on food sources (numerical dominance), preventing the use of the resource by other species

(Davidson 1998). The dominance-discovery tradeoff is an interesting mechanism to explain coexistence between

subordinate and dominant ant species. To find resources at faster rates, an ant colony needs to optimize the number of

scout workers (Pearce-Duvet and Feener 2010) that search for food, as opposed to dominant species that invest more

in recruiters (Fellers 1987). This tradeoff was tested in several situations, including in intraspecific interactions (Jordan

and Blüthgen 2007), between invasive species (Bertelsmeier et al. 2015*a*) and within guilds (Fellers 1987). Although

some studies demonstrated the importance of this mechanism in warranting ant species coexistence (Fellers 1987;

Lebrun and Feener 2007; Feener et al. 2008; Bertelsmeier et al. 2015*a*), others questioned its ubiquity (Adler et al.

2007; Jordan and Blüthgen 2007; Parr and Gibb 2012; Stuble et al. 2013; Castracani et al. 2014; Camarota et al. 2018;

Antoniazzi et al. 2021; Dáttilo and MacGregor-Fors 2021). Some authors also stressed factors that can modify the

outcome of dominance-discovery tradeoffs (Davidson 1998; Feener 2000; Adler et al. 2007; Parr and Gibb 2012).

The genus *Pheidole* is one of the most diverse ant lineages, with more than 1000 species described to date

(Bolton 2021), showing high levels of diversity in the subtropical and tropical regions (Economo et al. 2015; 2019).

The combination of high levels of local diversity and the fact that most *Pheidole* species are food generalists and

consequently share many resources (Wilson 2003), raise the question of what mechanisms could explain such high

levels of species co-occurrence. Here we test, through bait experiments, the possibility that *Pheidole* species co-exist

at local spatial scales by sharing resources according to the predictions of the dominance-discovery tradeoff. If the

dominance-discovery tradeoff explains the coexistence of *Pheidole* species, we expect to observe a consistent

interspecific variation along the axis of discovery (order of arrival) and dominance (number of workers recruited)

(Fellers 1987). Alternatively, a lack of evidence for the dominance-discovery tradeoff could be the result of *Pheidole*

species being more spatially segregated than the expected by chance (Connor and Simberloff 1979; Gotelli 2000), so

they would rarely interact directly to dispute resources. We therefore test whether *Pheidole* species are spatially

segregated, and expect that they are not more segregated than expected by chance.

Material and Methods

We carried out field experiments in two neighboring urban forest fragments located in the campus of the

Universidade Federal do Paraná in Curitiba, Paraná, southern Brazil (Site A - approximately 5.5 ha; 25º26'45.9"S;

49º13'55.5"W and site B - approximately 15 ha; 25°26'52.06"S; 49°14'19.24"W). The vegetation in these sites is

classified as mixed ombrophylous forest (Reginato et al. 2008), which is part of the Atlantic Forest biome (IBGE

2012). The climate in the study region is subtropical humid with temperate summers, categorized as *Cfb* according to

Köppen's system (Alvares et al. 2013). Experiments were carried out between the end of February and the beginning

of April of 2018, during the warm season, always in the morning, to minimize the effects of climatic differences

between sampling days. We established 55 sampling units along three transects in each of the two sampling sites (25

and 30 in sites A and B, respectively). In each transect, sampling units were set 10 meters apart from each other to

avoid resampling individuals from the same colony in different sampling units (Bestelmeyer et al. 2000). To properly

follow the behavior of ant workers at each sampling unit during each experiment, only five sampling units were

established each day, for a total of 11 sampling dates over the course of the study. Each sampling unit consisted of a

bait including 2-mL mixture of sardine and wheat flour, placed on a Petri dish (9 cm in diameter) that prevented ants

from having access to the bait from below, which could hamper direct observation and potentially underestimate their

presence on the baits. We offered a large amount of sardine to avoid quick removal of the resource before the end of

the experimentation time. There was no evidence that Petri dishes prevented workers to access the sardine. *Pheidole*

species are heavily attracted to sardine baits, which represent a rich source of proteins, sodium and lipids, and it has

been demonstrated that protein resources are more attractive to ants in forest remains at earlier successional stages

(Bihn et al. 2008). Each experiment lasted for three hours, such that each sampling unit was checked every 20 min,

for a total of nine time intervals, and the same person led all experiments.

We estimated the number of major and minor workers of *Pheidole* species at the sampling units with the aid

of a manual counter, and recorded the occurrence of negative interactions with other *Pheidole* species and with non-

Pheidole species (defined as interactions where a direct contact between workers leads them to get away from each

other, without necessarily a physical injury). In the case of interactions between *Pheidole* species, the same interaction

was counted twice, once for each species, given that we were not always able to define which species led the attack

against the other. For later species identification, we collected one individual per subcaste and morphospecies at each

sampling unit and deposited them in micro tubes filled with 95% ethanol. Voucher specimens were deposited at the

"Coleção Entomológica Padre Jesus Santiago Moure", Zoology Department, UFPR. Identifications followed the

Pheidole taxonomic key available in Wilson (2003). No permits were required for fieldwork, and collections of

specimens followed Brazilian legal provisions.

Analysis

If species were spatially segregated due to competitive interactions (Connor and Simberloff 1979; Gotelli

2000), we would rarely see more than one *Pheidole* species in the same sampling unit, which would necessarily lead

to a pattern in which the species that discovered the bait will remain alone until the end of the experiment. To test if

species are more spatially segregated than expected by chance, we calculated the C-score (Stone and Roberts 1990)

based on presence-absence matrices considering all sampling units. The observed C-score was then tested for

significance against an estimated C-score obtained from null models. We generated 5000 random matrices with fixed

row and fixed column sums (Gotelli 2000). We estimate the C-score for the combined data as well as the data for each

sampling site independently and for the common species (defined as the species that occurred in more than one

sampling unit). The C-score analysis were carried out in EcoSim7.0 (Gotelli and Entsminger 2001).

To assess species discovery ability, we defined a discovery ability index (DAI), which ranges from 0 to 1 and

is defined as the number of sampling units a species arrived first, divided by the total number of sampling units that

the species occurred at. If two or more *Pheidole* species occurred together in the same sampling unit at the first time

interval, both were treated as discoverers. Although observations started after 20 minutes, which prevent us from

ascertaining the exact order of arrival in a few instances in which two *Pheidole* species were presented in the first time

interval (four sampling units), there was no evidence that this time was enough for a species to discover the bait and

be completely removed by another *Pheidole* species, a process that tends to demand a greater amount of time (Perfecto

and Vandermeer 2011). To determine dominance ability as a proxy for numerical dominance (Davidson 1998), we

defined a dominance index (DI), which also ranges from 0 to 1 as the number of sampling units that a *Pheidole* species

dominated, divided by the total number of sampling units in which that species occurred. A species dominated the

sampling unit if it recruited more minor workers at the final time interval than the median number of minors recruited

across all baits and time intervals in which it occurred (Table 1), which resembles the monopolization index proposed

in Castracani et al. (2014) and Antoniazzi et al. (2021), based on the original proposition of Santini et al. (2007). In

considering the median number of minors recruited by each species to the baits as a threshold to define numerical

dominance, instead of a pre-defined specific number of workers for all species, we avoid the issues associated with

interspecific differences in recruitment behavior.

We calculated species occurrence, the frequency of negative interactions among *Pheidole* species and

between *Pheidole* and non-*Pheidole* species. The details of these calculations are provided in Fig. 2, Tables 1 and S3.

To test if the occurrence of negative interactions influences the number of workers being recruited to the sampling

units, we estimated the median number of minor workers being recruited at sampling units where negative interactions

occurred among *Pheidole* species and between *Pheidole* and non-*Pheidole* species. We applied a paired Wilcoxon test

(Zar 2010*a*) to compare if these numbers are different from the median number of minors recruited at all sampling

units, which is the number considered to define the DI. To test if the dominance-discovery tradeoff influences the

coexistence of *Pheidole* species, we applied a Spearman rank correlation test (Zar 2010*b*) between the DI and DAI,

with a negative correlation being considered as support for the dominance-discovery tradeoff (Fellers 1987). We

combined the data of both study areas but also considered them independently, as well as the data from only the

common species. Analyses were carried out in R 4.0-2 (R Core Team 2020).

Results

We recorded 12 *Pheidole* species across our two sampling sites (Table 1). Two morphospecies were

indistinguishable in the field, so to avoid any under or overestimation of each morphospecies worker numbers we

combined their data under a unique species cluster, the *P. flavens* complex. *Pheidole* species visited 48 out of 55

sampling units during our field experiments. Of all sampling units, 47% were discovered by only one species, 24%

by two species and 16% by three species. The most frequent species was *P. flavens* complex (45.45%), followed by

P. hetschkoi Emery, 1896 (32.73%) and *P. risii* Forel, 1892 (20%) (Table 1). *Pheidole* aff. *susannae*, *P.* cf. *lucretii*

and *P.* sp. 1 recruited the highest number of minor workers, whereas *P.* aff. *bucculenta* and *P.* sp. 2 recruited the

lowest numbers (Table 1). Four species never recruited major workers (*P.* aff. *bucculenta*, *P.* aff. *rosae*, *P. risii* and

P. sp. 2), and only *P.* aff. *susannae* and *P.* cf. *lucretii* recruited more than five majors at any sampling unit and time

interval (Table 1).

Regarding negative interactions, *Pheidole* sp.2 did not exhibit negative interactions against other *Pheidole*

species and *P.* aff. *susannae* never co-occurred with another *Pheidole* species (Table 1). The remaining ten species

showed aggressive behaviors against other *Pheidole* species (Table S1), with *P.* aff. *rosae* showing negative

interactions in 52.63% of the time intervals in which it occurred, followed by *P. nesiota* Wilson, 2003 (40%) and *P.*

hetschkoi (36.84%). However, aggressive behaviors between *Pheidole* species were observed in only 26.07% of time

intervals in which at least two *Pheidole* species co-occurred. During the experiments, some non-*Pheidole* species

(*Brachymyrmex* sp., *Camponotus rufipes* (Fabricius, 1775)*, Gnamptogenys* sp., *Linepithema* sp., *Nylanderia* sp.,

Odontomachus chelifer (Latreille, 1802)*, Pachycondyla striata* Smith, 1858, and *Solenopsis* sp.) were recorded and

showed agonistic interactions with *Pheidole* species (Table 1). *Pheidole* sp. 2 interacted negatively against a non-

Pheidole species in 28.57% of time intervals where it was recorded, followed by *P.* cf. *lucretii* (20%) and *P.* aff.

longiseta (15.38%). Negative interactions had no effect on the number of minor workers being recruited at sampling

units, given that the median number of minors of each *Pheidole* species along the entire experimentation time was not

different from the median number of minors being recruited at baits that registered negative interactions among

Pheidole species (V = 4; p = 0.42) and where negative interactions happened between *Pheidole* and non-*Pheidole*

species (V = 7.5; p-value = 0.46). Consequently, there was no evidence that negative interactions influenced the DI.

By comparing the spatial distribution of *Pheidole* species with random distributions generated by null models,

we did not find any evidence that species are more aggregated than expected by chance (Table S2; observed ≤

expected, p = 0.314) or less aggregated than expected by chance (Table S2; observed ≥ expected, p = 0.694) for the

combined data. When looking at site A individually the same patterns were found (Table S2; observed ≤ expected, p

= 0.106; observed ≥ expected, p = 0.919), but the results from site B suggest otherwise that *Pheidole* species are

spatially segregated (Table S2; observed ≤ expected, p = 0.967; observed ≥ expected, p = 0.038). However, when we

considered only the most common species, the co-occurrence patterns of all sites are not different from what is

expected by chance (Table S2). We did not find any negative correlation between DI and DAI (Figure 1), considering

both the combined data (Figure 1a) and the independent data from sites A (Figure 1b) and B (Figure 1c). The

conclusions remain the same when we considered just the common species (Figure S1). In general, most *Pheidole*

species stay on the sampling units throughout the entire duration of the experiment (Table S3; Figure 2). Even rarely

recorded species, like *P*. aff. *tijucana* and *P*. aff. *susannae*, which occurred just at one sampling unit, were never

replaced for those baits (Table S3).

Discussion

Despite the relatively high number of co-occurring *Pheidole* species at our sampling sites, we did not find

any evidence for dominance-discovery tradeoffs promoting species coexistence, suggesting that *Pheidole* species

show similar strategies to invest in resource discovery and dominance. Specifically, we found no negative correlation

between the discovery and dominance indices, contrary to the expectation for the occurrence of the tradeoff (Fellers

1987). By reviewing the literature on dominance-discovery tradeoff, some authors suggested that the tradeoff might

be context dependent (Davidson 1998; Parr and Gibb 2012), as also proposed under a simulation approach (Adler et

al. 2007). The levels of aggressiveness between pairwise species interactions can contribute to establish dominance

hierarchies (Savolainen and Vepsäläinen 1988). However, here we recorded negative interactions between *Pheidole*

species in only 26.07% of the encounters between at least two *Pheidole* species, and we found no evidence that

Pheidole species changed their recruitment behavior due to the occurrence of any kind of negative interaction.

Although commonly recorded in ant interactions, aggressiveness need not be the norm in interspecific encounters,

even among dominant and highly invasive ants (Bertelsmeier et al. 2015*b*). Neutral interactions between coexisting

species at bait experiments represent a great portion of the observed outcomes (Blüthgen et al. 2004; Stuble et al.

2017; Gray et al. 2018), yet they are mainly neglected, with emphasis in studies granted preferentially to true agonistic

behaviors (Stuble et al. 2017). In *Pheidole* species, the increased proportion of major workers recruited to food sources

could be an important trait to enhance the probability of resource dominance (Mertl et al. 2010). Although the species

considered here were markedly distinct with respect to the number of majors recruited at baits, this difference was not

enough to be translated into a dominance hierarchy, and majors were rarely observed engaging in negative interactions.

Reacting aggressively or not against a potential competitor is context-dependent (Tanner and Adler 2009;

Barbieri et al. 2013). The "dear enemy" phenomenon posits that animals can recognize potential competitors that live

close to their own territory from others that are strangers, and react more aggressively against strangers (Langen et al.

2000; Tanner and Adler 2009). If the *Pheidole* species that approach the sampling units are neighbors that constantly

interact, this could explain the relative low levels of aggression in interspecific encounters. However, the opposite

trend was also recorded for ants (Gordon 1989), and to evaluate the importance of this phenomenon in the present

context would require a more detailed investigation of the nature of the interspecific encounters between the *Pheidole*

species. The type of resource being disputed and the nutritional demands of the colony can also influence the foraging

behavior of ant workers (Cornelius and Grace 1997; Kay 2004; Silberman et al. 2016). In general, proteins and lipids

are the main nutrients to feed larvae and the queens, whereas for adult workers carbohydrates are the main energy

sources to sustain their activities (Blüthgen and Feldhaar 2010). Although we used here a unique source of food to

attract *Pheidole* workers to baits, sardine baits are sources of both proteins and lipids (Lasmar et al. 2021), which are

commonly searched by *Pheidole* species in the field along with carbohydrate (Rosumek 2017). Further studies can

investigate if other food types can elicit more aggressive behaviors between *Pheidole* workers, which occasionally

can result in interactions that follow the predictions of the dominance-discovery tradeoff.

Coupled with the low number of agonistic interactions between ant species at baits, in nearly half of our

sampling units we recorded only one *Pheidole* species throughout the entire experimentation time. In some situations,

it was clear that the first species that arrived at baits had nests close to the baits (author's pers. obs.), which represents

a clear advantage to find the resource and recruit workers to dominate it. The implication is that the crucial stage for

resource assurance could be the discovery event, a fact recently suggested for Neotropical arboreal ant communities

(Camarota et al. 2018; Antoniazzi et al. 2021) and a Neotropical urban gradient (Dátillo and MacGregor-Fors), which

was defined as a discovery-defense strategy (Camarota et al. 2018). Although we did not find a positive relationship

between discovery and dominance abilities, there is a trend towards it in our results from both the combined data and

individual data from sites A and B, which was not confirmed due to the contrasting behavior of three species (*P*. sp.1,

P. sp.2 and *P*. aff. *tijucana*). This pattern differs from the two step view whereby the resource used by ants is

partitioned between a food search phase, followed by a latter moment of competition displacement and species

turnover where other ants have greater success, the so called dominance-discovery tradeoff (Fellers 1987). This

suggests that other mechanisms can be involved in structuring this community, such as priority effects (Andersen

2008). Priority effects predict that the order of species arrival at empty patches determines the subsequent structuring

of the community (Fukami 2015; De Meester et al. 2016). The order of arrival into a resource patch can determine the

level of aggressiveness of dominant ant species (Barbieri et al. 2013), in a way that influences resource partitioning.

Order of arrival can also change the colony level of activity and survival (Barbieri et al. 2013), with important

consequences for species coexistence and community structure.

By arriving first at baits, the *Pheidole* species in our study have more time to recruit nestmates and dominate

the resource before the arrival of other *Pheidole* species, and in the absence of successful takeovers in resource patches,

coexistence can be improved (Adler et al. 2007). If the *Pheidole* species investigated here are spatially segregated to

some degree, this could explain why so many sampling units were visited by only one species. Although our analysis

of spatial structure from site B suggests that those *Pheidole* species are spatially segregated, it seems that this result is

an effect of the rare species. When just common species are considered, the patterns of species co-occurrence agree

with what is expected by chance (Stone and Roberts 1990; Gotelli 2000). Another possibility is that the complexity

of the habitat delays the time to discover and recruit nestmates to the baits. More complex habitats affect the way that

ants sense the environment and influence its locomotion capacity (Kaspari and Weiser 1999), which consequently

affects the time to discover and recruit nestmates to the baits (Parr and Gibb 2012). The time to discover food sources

can vary because of several biological aspects of ant species, but there are cues that in more simple vegetation types

ants are able to find resources at faster rates (Gibb and Hochuli 2004) than in more complex habitats (Holway 1999).

Although our sampling sites conserve vegetation elements of secondary stages of succession (Reginato et al. 2008),

both sites are composed of arboreal species that provide a complex leaf-litter in which ants forage, which can hinder

ant locomotion and navigation.

We show here that a classical mechanism of species coexistence, the dominance-discovery tradeoff (Fellers

1987), was not able to explain *Pheidole* species co-occurrence in two assemblages of Atlantic Forest remains in south

Brazil. Similarly, coexistence of *Pheidole* species seems not to be related to tradeoffs between dominance and

environmental resistance (Tschá and Pie 2019), which suggests that evolutionary tradeoffs can have a secondary role

to structuring the *Pheidole* assemblages investigated. Although there are cues for the importance of priority effects

(Andersen 2008) to the resource partitioning between *Pheidole* species, this hypothesis remains to be tested in the

present community. Also important to advance our knowledge about *Pheidole* species coexistence is a deeper

understanding of their dietary preferences (Rosumek 2017; Rosumek et al. 2018) and the role of stochastic

mechanisms in promoting species coexistence (Andersen et al. 2013; Stuble et al. 2017). The importance of the

dominance-discovery tradeoff in structuring ant communities is highly context-dependent (Davidson 1998; Parr and

Gibb 2012), and the prevalence of ecological tradeoffs in general depend on the spatial scales being considered

(Kneitel and Chase 2004). Therefore, a lack of evidence for the occurrence of the dominance-discovery tradeoff in

many ant assemblages (Parr and Gibb 2012) should not be viewed as a definitive proof against the importance of such

mechanisms in the structuring of ant communities. The dominance-discovery tradeoff should be more broadly

explored, along different spatial scales and with different organisms, to provide a deeper understanding about the

range of their influence in ecological communities to ensure species coexistence.

Acknowledgments

The authors thank Alexandre Casadei Ferreira for assistance in species identification and Félix Rosumek for

comments on the manuscript.

Conflicts of interest/Competing interests

The authors declare no competing interests.

Authors' contributions

CLK and MRP designed the study, CLK carried out field and lab work and data analysis, CLK and MRP

wrote the manuscript.

Funding

The Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Brazil, provided a graduate

fellowship to CLK (Financial Code 001). MRP was partially funded by a grant from CNPq/MCT (302904/2020-4).

References

Adler, F.R., LeBrun, E.G., and Feener Jr., D.H. 2007. Maintaining Diversity in an Ant Community: Modeling,

Extending, and Testing the Dominance-Discovery Trade-Off. The American Naturalist **169**(3): 323–333.

doi:10.1086/510759.

Alvares, C.A., Stape, J.L., Sentelhas, P.C., De Moraes Gonçalves, J.L., and Sparovek, G. 2013. Köppen's climate

classification map for Brazil. Meteorol. Zeitschrift **22**(6): 711–728. doi:10.1127/0941-2948/2013/0507.

Andersen, A.N. 2008. Not enough niches: Non-equilibrial processes promoting species coexistence in diverse ant

communities. Austral Ecol. **33**(2): 211–220. doi:10.1111/j.1442-9993.2007.01810.x.

Andersen, A.N., Arnan, X., and Sparks, K. 2013. Limited niche differentiation within remarkable co-occurrences of

congeneric species: Monomorium ants in the Australian seasonal tropics. Austral Ecol. **38**(5): 557–567.

doi:10.1111/aec.12000.

Antoniazzi, R., Camarota, F., Leponce, M., and Dáttilo, W. 2021. Discovery-defense strategy as a mechanism of social

foraging of ants in tropical rainforest canopies. Behavioral Ecology: arab054. doi:10.1093/beheco/arab054.

Arnan, X., Cerdá, X., and Retana, J. 2012. Distinctive life traits and distribution along environmental gradients of

dominant and subordinate Mediterranean ant species. Oecologia **170**(2): 489–500. doi:10.1007/s00442-012-

2315-y.

Barbieri, R.F., Grangier, J., and Lester, P.J. 2013. Arrival sequence and diet mediate interspecific competition in an

ant community. Insectes Soc. **60**(4): 463–473. doi:10.1007/s00040-013-0312-8.

Bertelsmeier, C., Avril, A., Blight, O., Confais, A., Diez, L., Jourdan, H., Orivel, J., Saint Germès, N., and Courchamp,

F. 2015*b*. Different behavioural strategies among seven highly invasive ant species. Biol. Invasions **17**(8): 2491–

2503. doi:10.1007/s10530-015-0892-5.

Bertelsmeier, C., Avril, A., Blight, O., Jourdan, H., and Courchamp, F. 2015*a*. Discovery-dominance trade-off among

widespread invasive ant species. Ecol. Evol. **5**(13): 2673–2683. doi:10.1002/ece3.1542.

Bestelmeyer, B.T., Agosti, D., Alonso, L.E., Brandão, R.F., Brown Jr., W.L., Delabie, J.H.C., and Silvestre, R. 2000.

Field techniques for the study of ground-dwelling ants: an overview, description and evaluation. *In* Ants:

standard methods for measuring and monitoring biodiversity, 1st edition. Smithsonian Institution Press.,
Washington, USA. pp. 122–144.

Bihn, J.H., Verhaagh, M., and Brandl, R. 2008. Ecological Stoichiometry along a Gradient of Forest Succession: Bait
Preferences of Litter Ants. Biotropica **40**(5): 597–599. doi:10.1111/j.1744-7429.2008.00423.x.

Blüthgen, N., and Feldhaar, H. 2010. Food and shelter: how resources influence ant ecology. *In* Ant Ecology. Oxford
University Press, New York. pp. 115–136.

Bluthgen, N., Stork, N.E., and Fiedler, K. 2004. Bottom-Up Control and Co-Occurrence in Complex Communities :
Honeydew and Nectar Determine a Rainforest Ant Mosaic. Oikos **106**(2): 344–358.

Bolton, B. 2021. An online catalog of the ants of the world. Available from antcat.org [accessed 31 March 2021].

Brown, J.S., Kotler, B.P., and Mitchell, W.A. 1994. Foraging Theory, Patch Use, and the Structure of a Negev Desert
Granivore Community. Ecology **75**(8): 2286. doi:10.2307/1940884.

Camarota, F., Vasconcelos, H.L., Koch, E.B.A., and Powell, S. 2018. Discovery and defense define the social foraging
strategy of Neotropical arboreal ants. Behav. Ecol. Sociobiol. **72**(7): 1–11. Behavioral Ecology and
Sociobiology. doi:10.1007/s00265-018-2519-1.

Castracani, C., Spotti, F.A., Grasso, D.A., and Fanfani, A. 2014. A new exception to the dominance-discovery trade-
off rule in ant communities. Redia **97**: 171–175.

Connor, E.F., and Simberloff, D. 1979. The Assembly of Species Communities: Chance or Competition? Ecology
60(6): 1132. doi:10.2307/1936961.

Cornelius, M.L., and Grace, J.K. 1997. Influence of Brood on the Nutritional Preferences of the Tropical Ant Species,
Pheidole megacephala (F.) and *Ochetellus glaber* (Mayr). Journal of Entomological Science **32**(4): 421–429.
doi:10.18474/0749-8004-32.4.421.

Dáttilo, W., and MacGregor-Fors, I. 2021. Ant social foraging strategies along a Neotropical gradient of urbanization.
Sci Rep **11**(1): 6119. doi:10.1038/s41598-021-85538-2.

Davidson. 1998. Resource discovery versus resource domination in ants: a functional mechanism for breaking the
trade-off. Ecol. Entomol. **23**: 484–490.

Del Toro, I., Ribbons, R.R., and Pelini, S.L. 2012. The little things that run the world revisited: a review of ant-
mediated ecosystem services and disservices (Hymenoptera: Formicidae). Myrmecol. News **17**: 133–146.

De Meester, L., Vanoverbeke, J., Kilsdonk, L.J., and Urban, M.C. 2016. Evolving Perspectives on Monopolization
and Priority Effects. Trends Ecol. Evol. **31**(2): 136–146. doi:10.1016/j.tree.2015.12.009.

Economo, E.P., Huang, J.P., Fischer, G., Sarnat, E.M., Narula, N., Janda, M., Guénard, B., Longino, J.T., and
Knowles, L.L. 2019. Evolution of the latitudinal diversity gradient in the hyperdiverse ant genus *Pheidole*.
Global Ecol Biogeogr **28**(4): 456–470. doi:10.1111/geb.12867.

Economo, E.P., Klimov, P., Sarnat, E.M., Guénard, B., Weiser, M.D., Lecroq, B., and Knowles, L.L. 2015. Global
phylogenetic structure of the hyperdiverse ant genus *Pheidole* reveals the repeated evolution of macroecological
patterns. Proc. R. Soc. B **282**(1798): 20141416. doi:10.1098/rspb.2014.1416.

Economo, E.P., Narula, N., Friedman, N.R., Weiser, M.D., and Guénard, B. 2018. Macroecology and macroevolution
of the latitudinal diversity gradient in ants. Nat. Commun. **9**(1): 1–8. Springer US. doi:10.1038/s41467-018-
04218-4.

Feener, D.H. 2000. Is the Assembly of Ant Communities Mediated by Parasitoids? Oikos **90**(1): 79–88.

Feener, D.H., Orr, M.R., Wackford, K.M., Longo, J.M., Benson, W.W., Gilbert, L.E., Feener, D.H., Orr, M.R.,
Wackford, K.M., Longo, J.M., Benson, W.W., and Gilbert, L.E. 2008. Geographic Variation in Resource
Dominance-Discovery in Brazilian Ant Communities Ecology **89**(7): 1824–1836.

Fellers, J.H. 1987. Interference and Exploitation in a Guild of Woodland Ants. Ecol. Soc. Am. **68**(5): 1466–1478.

Fittkau, E.J., and Klinge, H. 1973. On Biomass and Trophic Structure of the Central Amazonian Rain Forest
Ecosystem. Biotropica **5**(1): 2. doi:10.2307/2989676.

Fukami, T. 2015. Historical Contingency in Community Assembly: Integrating Niches, Species Pools, and Priority
Effects. Annu. Rev. Ecol. Evol. Syst. **46**: 1–23. doi:10.1146/annurev-ecolsys-110411-160340.

Garnier, S. 2018. viridis: Default Color Maps from "matplotlib." Available from https://CRAN.R-
project.org/package=viridis.

Gibb, H., and Hochuli, D.F. 2004. REMOVAL EXPERIMENT REVEALS LIMITED EFFECTS OF A

BEHAVIORALLY DOMINANT SPECIES ON ANT ASSEMBLAGES. Ecology **85**(3): 648–657. doi:10.1890/03-0007.

Gordon, D.M. 1989. Ants distinguish neighbors from strangers. Oecologia **81**(2): 198–200. doi:10.1007/BF00379806.

Gotelli, N.J., and Entsminger, G.L. 2001. EcoSim: Null models software for ecology. Acquired Intelligence Inc. & Kesey-Bear. Available from http://homepages.together.net/~gentsmin/ecosim.htm.

Gotelli, N.J. 2000. NULL MODEL ANALYSIS OF SPECIES CO-OCCURRENCE PATTERNS. Ecology **81**(9): 2606–2621. doi:10.1890/0012-9658(2000)081[2606:NMAOSC]2.0.CO;2.

Gray, R.E.J., Ewers, R.M., Boyle, M.J.W., Chung, A.Y.C., and Gill, R.J. 2018. Effect of tropical forest disturbance on the competitive interactions within a diverse ant community. Sci. Rep. **8**(1): 1–12. Springer US. doi:10.1038/s41598-018-23272-y.

Hölldobler, B.., and Wilson, E.O. 1990. The Ants. Harvard University Press.

Holway, D.A. 1999. COMPETITIVE MECHANISMS UNDERLYING THE DISPLACEMENT OF NATIVE ANTS BY THE INVASIVE ARGENTINE ANT. Ecology **80**(1): 238–251. doi:10.1890/0012-9658(1999)080[0238:CMUTDO]2.0.CO;2.

Hubbell, S.P., and Johnson, L.K. 1978. Comparative Foraging Behavior of Six Stingless Bee Species Exploiting a Standardized Resource. Ecology **59**(6): 1123–1136. doi:10.2307/1938227.

Hutchinson, G.E. 1959. Homage to Santa Rosalia or Why Are There So Many Kinds of Animals? The American Naturalist **93**(870): 145–159. doi:10.1086/282070.

Hutchinson, G.E. 1961. The Paradox of the Plankton. The American Naturalist **95**(882): 137–145. doi:10.1086/282171.

IBGE. 2012. Manual Técnico da Vegetação Brasileira. IBGE, Rio de Janeiro.

Jordan, R., and Blüthgen, N. 2007. No evidence for a dominance-discovery trade-off among Formica pratensis colonies (Hymenoptera: Formicidae). Myrmecol. News **10**: 7–10.

Kaspari, M. 2001. Taxonomic Level, Trophic Biology and the Regulation of Local Abundance.Glob. Ecol. Biogeogr. **10**(3): 229–244.

Kaspari, M., and Weiser, M.D. 1999. The size–grain hypothesis and interspecific scaling in ants. Functional Ecology
13(4): 530–538. doi:10.1046/j.1365-2435.1999.00343.x.

Kay, A. 2004. The relative availabilities of complementary resources affect the feeding preferences of ant colonies.
Behavioral Ecology **15**(1): 63–70. doi:10.1093/beheco/arg106.

Kneitel, J.M., and Chase, J.M. 2004. Trade-offs in community ecology: linking spatial scales and species coexistence.
Ecol Letters **7**(1): 69–80. doi:10.1046/j.1461-0248.2003.00551.x.

Langen, T.A., Tripet, F., and Nonacs, P. 2000. The red and the black: habituation and the dear-enemy phenomenon in
two desert *Pheidole* ants. Behavioral Ecology and Sociobiology **48**(4): 285–292. doi:10.1007/s002650000223.

Lasmar, C.J., Bishop, T.R., Parr, C.L., Queiroz, A.C.M., Schmidt, F.A., and Ribas, C.R. 2021. Geographical variation
in ant foraging activity and resource use is driven by climate and net primary productivity. J Biogeogr **48**(6):
1448–1459. doi:10.1111/jbi.14089.

Lebrun, E.G., and Feener, D.H. 2007. When trade-offs interact: Balance of terror enforces dominance discovery trade-
off in a local ant assemblage. J. Anim. Ecol. **76**(1): 58–64. doi:10.1111/j.1365-2656.2006.01173.x.

Leite, P.F.., and Klein, R.M. 1990. Leite Klein 1990- Vegetação do Sul do Brasil.pdf. IBGE, Rio de Janeiro.

Mertl, A.L., Sorenson, M.D., and Traniello, J.F.A. 2010. Community-level interactions and functional ecology of
major workers in the hyperdiverse ground-foraging *Pheidole* (Hymenoptera, Formicidae) of Amazonian
Ecuador. Insectes Soc. **57**(4): 441–452. doi:10.1007/s00040-010-0102-5.

Nagamitsu, T., and Inoue, T. 1997. Aggressive foraging of social bees as a mechanism of floral resource partitioning
in an Asian tropical rainforest. Oecologia **110**(3): 432–439. doi:10.1007/s004420050178.

Parr, C.L., and Gibb, H. 2010. Competition and the role of dominant ants. *In* Ant Ecology. Oxford University Press,
Oxford. pp. 77–96.

Parr, C.L., and Gibb, H. 2012. The discovery-dominance trade-off is the exception, rather than the rule. J. Anim. Ecol.
81(1): 233–241. doi:10.1111/j.1365-2656.2011.01899.x.

Pearce-Duvet, J.M.C., and Feener, D.H. 2010. Resource discovery in ant communities: Do food type and quantity
matter? Ecol. Entomol. **35**(5): 549–556. doi:10.1111/j.1365-2311.2010.01214.x.

Perfecto, I., and Vandermeer, J. 2011. Discovery dominance tradeoff: the case of Pheidole subarmata and Solenopsis
geminata (Hymenoptera: Formicidae) in neotropical pastures. Environ. Entomol. **40**(5): 999–1006.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing,
Vienna, Austria. Available from www.R-project.org.

Reginato, M., Matos, F.B., Lindoso, G. da S., Souza, C.M.F. de, Prevedello, J.A., Morais, J.W., and Evangelista,
P.H.L. 2008. A vegetação na Reserva Mata Viva, Curitiba, Paraná, Brasil. Acta Biológica Parana. **37**(1989):
229–252. doi:10.5380/abpr.v37i0.13211.

Rosumek, F.B., Blüthgen, N., Brückner, A., Menzel, F., Gebauer, G., and Heethoff, M. 2018. Unveiling community
patterns and trophic niches of tropical and temperate ants using an integrative framework of field data, stable
isotopes and fatty acids. PeerJ **6**: e5467. doi:10.7717/peerj.5467.

Rosumek, F.B. 2017. Natural history of ants: What we (do not) know about trophic and temporal niches of neotropical
species. Sociobiology **64**(3): 244–255. doi:10.13102/sociobiology.v64i3.1623.

Santini, G., Tucci, L., Ottonetti, L., and Frizzi, F. 2007. Competition trade-offs in the organisation of a Mediterranean
ant assemblage. Ecol Entomol **32**(3): 319–326. doi:10.1111/j.1365-2311.2007.00882.x.

Savolainen, R., and Vepsäläinen, K. 1988. A Competition Hierarchy among Boreal Ants: Impact on Resource
Partitioning and Community Structure. Oikos **51**(2): 135. doi:10.2307/3565636.

Silberman, R.E., Gordon, D., and Ingram, K.K. 2016. Nutrient stores predict task behaviors in diverse ant species.
Insect. Soc. **63**(2): 299–307. doi:10.1007/s00040-016-0469-z.

Stone, L., and Roberts, A. 1990. The checkerboard score and species distributions. Oecologia **85**(1): 74–79.
doi:10.1007/BF00317345.

Stuble, K.L., Juric, I., Cerdá, X., and Sanders, N.J. 2017. Dominance hierarchies are a dominant paradigm in ant
ecology (Hymenoptera: Formicidae), but should they be? and what is a dominance hierarchy anyways?
Myrmecological News **24**: 71–81.

Stuble, K.L., Rodriguez-Cabal, M.A., McCormick, G.L., Jurić, I., Dunn, R.R., and Sanders, N.J. 2013. Tradeoffs,
competition, and coexistence in eastern deciduous forest ant communities. Oecologia **171**(4): 981–992.
doi:10.1007/s00442-012-2459-9.

Tanner, C.J., and Adler, F.R. 2009. To fight or not to fight: context-dependent interspecific aggression in competing

ants. Animal Behaviour **77**(2): 297–305. doi:10.1016/j.anbehav.2008.10.016.

Tschá, M.K., and Pie, M.R. 2019. Correlates of ecological dominance within *Pheidole* ants (Hymenoptera:

Formicidae). Ecol. Entomol. **44**(2): 163–171. doi:10.1111/een.12685.

Wickham, H. 2009. Ggplot2: elegant graphics for data analysis. Springer, New York.

Wilson, E.O. 2003. *Pheidole* in the New World: a dominant, hyperdiverse ant genus. Harvard University Press,

Cambridge.

Yitbarek, S., and Philpott, S.M. 2019. Arboreal twig-nesting ants form dominance hierarchies over nesting resources.

PeerJ **2019**(11): 1–17. doi:10.7717/peerj.8124.

Zar, J.H. 2010*a*. Paired-Sample Hypothesis. *In* Biostatistical analysis, 5th edition. Prentice-Hall/Pearson, Upper

Saddle River, New Jersey.

Zar, J.H. 2010*b*. Simple Linear Correlation. *In* Biostatistical analysis, 5th edition. Prentice-Hall/Pearson, Upper Saddle

River, New Jersey.

Table 1. Species sampled, approximate number (median) of major and minor workers registered, occurrence at baits, number

(frequency) of negative interactions among *Pheidole* species and between *Pheidole* species (Neg. int. *Pheidole*) and non-*Pheidole*

species (Neg. int. non-*Pheidole*). Species occurrence represents the number of sampling units that each species occurred at out of

the total number of sampling units (55). The frequency of negative interactions among *Pheidole* species is defined as the number

of time intervals that each species engages in a negative interaction with another *Pheidole* species out of the total number of time

intervals that it co-occurred with any other *Pheidole* species. The frequency of negative interactions between *Pheidole* species and

non-*Pheidole* species is defined as the number of time intervals that a *Pheidole* species engages in a negative interaction with a

non-*Pheidole* species out of the total number of time intervals that it occurred.

Species	Nº major records[a]	Nº minor records[a]	Occurrence (%)	Neg. int. *Pheidole*	Neg. int. non-*Pheidole*
P. aff. *bucculenta*	0 (0)	4 (1)	1.82	1(25)	0(0)
P. aff. *longiseta*	24 (2)	345 (30)	3.64	2(25)	2(15.38)
P. aff. *rosae*	0 (0)	209 (4.5)	7.27	10(52.63)	2(10)
P. aff. *susannae*	211 (25)	900 (100)	1.82	0(0)	0(0)
P. aff. *tijucana*	3 (0)	109 (10)	1.82	1(20)	0(0)
P. cf. *lucretii*	218 (2)	3833 (60)	18.18	6(28.57)	14(20)
P. flavens complex	15 (0.10)	1544 (5)	45.45	19(20.43)	9(5.81)
P. hetschkoi	1 (0)	699 (4)	32.73	14(36.84)	15(14.29)
P. nesiota	35 (1)	197 (6)	7.27	2(40)	1(5)
P. risii	0 (0)	831 (5)	20	5(17.24)	0(0)
P. sp.1	4 (0)	335 (40)	1.82	1(14.29)	0(0)
P. sp.2	0 (0)	16 (2)	1.82	0(0)	2(28.57)

450 [a]Approximate sum (median) of major and minor workers at each sampling unit at each time interval. This sum does not correspond

451 to abundance, since it is likely that the same individuals were present in the same Petri dish over the entire experiment time. If

452 more than 100 workers were counted, we stopped to count and considered this as the final number of workers.

Fig. 1 Correlations between discovery and dominance indices from the combined (a), site A (b) and site B (c) data.

Insets display the results of the Spearman correlation tests. Plots were generated with the R package GGPLOT2

(Wickham 2009).

Fig.2 Relative frequency of *Pheidole* species over the experiment time (a). Relative frequency considers the number

of records of each *Pheidole* species out of the total number of time intervals that a *Pheidole* species occurred (483).

461   In (b), the same relationship is visualized with the relative frequencies of each species standardized to a constant

462   height. Plots were generated with the R package "GGPLOT2" (Wickham 2009) and "viridis" (Garnier 2018).