Title: TET1 dioxygenase is required for FOXA2-associated chromatin remodeling in

pancreatic beta-cell differentiation

Authors: Xinwei Wu[1#], Jianfang Li[2#], Minjung Lee[3], Qingping Lan[1], Jia Li[3], Yun Huang[3], De-Qiang

Sun[2,3*], Ruiyu Xie[1*]

Affiliations:

[1] Centre of Reproduction, Development, and Aging, Faculty of Health Sciences, University of Macau,

Taipa, Macau SAR.

[2] Department of Center Laboratory, The Fifth Affiliated Hospital of Guangzhou Medical University,

Guangzhou 510700, China.

[3] Center for Epigenetics & Disease Prevention, Institute of Biosciences and Technology, College of

Medicine, Texas A&M University, Houston, TX 77030, USA

# These authors contributed equally to the work.

* Correspondence: ruiyuxie@um.edu.mo; dsun@ibt.tamhsc.edu

Abstract: Existing knowledge of the role of epigenetic modifiers in pancreas development has exponentially increased. However, the function of TET dioxygenases in pancreatic endocrine specification remains obscure. We set out to tackle this issue using a human embryonic stem cell (hESC) differentiation system, in which *TET1/TET2/TET3* triple knock-out cells displayed severe defects in pancreatic β-cell specification. Integrative whole-genome analysis identified unique cell-type-specific hypermethylated regions (hyper-DMRs) displaying reduced chromatin activity and remarkable enrichment for the binding of FOXA2, a pioneer transcription factor essential for pancreatic endoderm specification. Intriguingly, hundreds of hyper-DMRs recently identified in type-2 diabetes pathogenesis overlapped with the hyper-DMRs we found in TET-deficient cells. Furthermore, transduction of *TET1* in *TET*-deficient cells effectively rescued β-cell differentiation and reversed hypermethylation and suppression of the β-cell determinant *PAX4*. Genome-wide mapping of TET1 showed that TET1 co-localized at a subset of FOXA2 targets featuring high levels of active chromatin modifications in pancreatic progenitors. Taking these findings together with the defective generation of functional β-cells upon TET1-inactivation, our study not only unveils an essential role of TET1-dependent epigenetic regulation in establishing β-cell identity but also provides a new mechanistic clue regarding the complex crosstalk between TET dioxygenases and pioneer transcription factors in lineage specification.

Introduction

During embryonic development, pluripotent human embryonic stem cells (hESCs) differentiate into many diverse lineages that make up the complex tissues and organs of the human body. Pancreatic lineage specification relies on cross-talk between the genome and environmental cues in the progenitor niche. This cross-talk is mediated by cis-regulatory elements that play a prominent role in spatiotemporal gene regulation during embryogenesis. In particular, distal regulatory elements, such as enhancers, serve as information integration hubs that allow binding of multiple regulators, including lineage-specific transcription factors (TFs) as well as epigenetic readers, writers, and erasers to ensure integration of intrinsic and extrinsic environmental cues at these loci [1,2]. It was recently demonstrated that pioneer TFs, such as FOXA1 and FOXA2, are required for proper chromatin opening and establishment of enhancer marks H3K4me1 and H3K27ac during pancreatic fate specification [3,4]. These pioneer TFs bind to nucleosomal DNA to initiate chromatin remodeling associated with DNA demethylation at newly accessible enhancers [5-9].

DNA demethylation is mediated by ten-eleven-translocation methylcytosine dioxygenases (TETs), which catalyze sequential oxidation of 5-methylcytosine (5mC) to 5-hydroxymethylcytosine (5hmC), 5-formylcytosine, and 5-carboxylcytosine [10-14]. Distribution of 5hmC, a novel epigenetic modification, is dynamically changed by and positively correlated with active gene transcription during early lineage specification [15-17]. Consequently, inhibition of TET family enzymes (TET1, TET2, and TET3) impairs cell fate commitment into neural, hematopoietic, cardiac, and several other lineages [18-20]. Although TETs have been shown to regulate 5mC homeostasis during early embryonic

development, the functional relevance and mechanisms by which TETs regulate pancreatic endocrine

specification remain mostly unknown.

Results

TET deficiency impairs pancreatic endoderm differentiation. We previously demonstrated that

5hmC positively correlates with 'open' chromatin at poised and active enhancers in multiple

endodermal lineage intermediates using a stepwise hESC differentiation system toward pancreatic

progenitors [21]. To determine the biological significance of TET-dependent regulation during pancreas

specification, we generated *TET1*, *TET2*, and *TET3* single knock-out (KO), double knock-out (DKO),

and triple knock-out (TKO) H1 hESC lines using CRISPR/Cas9 technology (Supplementary Fig. 1a).

Mutations resulted in premature termination codons, which were confirmed by Sanger sequencing.

Global DNA hydroxymethylation levels from positive clones were assayed by 5hmC dot blot. A

significant reduction of 5hmC signals was shown in TKO, TET1KO, TET1/2DKO, and TET1/3DKO

cells, whereas TET2KO, TET3KO, and TET2/3DKO cells showed minimal alterations of 5hmC

levels (Supplementary Fig. 1b). To avoid functional redundancy, we focused our analysis on TKO

cells devoid of any TET-mediated active demethylation. Consistent with other *TET* knock-out mESC

and hESC lines [18,22], TKO H1-hESCs exhibited no apparent defects in stem cell self-renewal capacity

or expression of pluripotent factors (Supplementary Fig. 1e).

To determine whether TET proteins affect hESC differentiation toward pancreatic endocrine fate,

we used an established stepwise differentiation platform [21] to induce efficient differentiation of

hESCs to definitive endoderm (DE), primitive gut tube (GT), pancreatic progenitor (PP), and

pancreatic endocrine (PE) (Fig. 1a). Both TET triple-deficient lines (clones 2 and 6) displayed similar

differentiation efficiency toward endoderm germ layer as the wild-type (WT) hESC line, in which

over 90% of cells were SOX17$^+$ by day 3 of differentiation (Supplementary Fig. 1c). Moreover,

expression levels of other endoderm markers, such as *FOXA2*, *FOXA1*, and *CXCR4*, were unchanged

in TET-depleted cells compared with control cells (Supplementary Fig. 1d, e). This analysis suggests

that TET dioxygenases are dispensable for endoderm specification in the context of *in vitro* hESC

differentiation. To examine the effects of TET ablation on pancreatic commitment, we subsequently

examined the expression of critical pancreatic markers at the PP and PE stages. Using flow cytometry

to quantitate the expression of *PDX1* at the PP stage and co-expression of *PDX1* and *NKX6.1* at the

PE stage, we observed a substantial decrease in TET knock-out cells (Fig. 1b). These data were

confirmed by immunofluorescence staining and RT-qPCR analysis (Fig. 1c, Supplementary Fig. 1e).

Consistent with these findings, expression of the pancreatic progenitor markers *SOX9* and *PTF1A*

and endocrine hormones insulin, and glucagon were significantly downregulated in TKO lines (Fig.

1c, Supplementary Fig. 1e). Interestingly, *PAX4*, a key determinant for β-cell specification, failed to

induce in TET-inactivated cells, whereas expression of the α-cell determinant, *ARX*, was not affected

(Supplementary Fig. 1e). These results demonstrate that TETs are required for pancreatic β-cell

lineage specification.

To investigate TET-dependent global transcriptional changes during pancreatic endocrine cell

fate commitment, we performed transcriptome analysis of WT and TKO cells at the ES, DE, and PP

stages. Hierarchical clustering of genes specifically expressed at each stage revealed a remarkable

downregulation of PP-specific genes in TKO_PP cells, whereas ES-specific and DE-specific genes

showed no obvious changes in TET-deficient cells compared with WT cells (Supplementary Fig. 1f).

In particular, 1,173 differentially expressed genes (DEGs) were identified in TKO_PP cells (Fig. 1d,

Supplementary Table 1). Among them, there were 921 downregulated DEGs in TET-deficient cells,

including many important developmental determinants for pancreatic endocrine specification, such

as *PDX1*, *NKX2.2*, *NKX6.1*, *NKX6.2*, *NEUROG3*, and *PAX4*. We also observed 252 genes,

particularly HOX family members, which were significantly upregulated upon TET deletion,

suggesting that failure to differentiate into pancreatic β-cells is tightly linked to an aberrant TF

network. In addition, KEGG pathway analysis of the DEGs showed enrichment for terms associated

with maturity onset diabetes of the young (MODY) and insulin secretion (Supplementary Fig. 1g),

suggesting that loss of TETs impairs pancreatic endocrine formation at later stages of endoderm

specification.

FOXA motifs are enriched at distal regulatory elements displaying decreased accessibility upon

TET deletion. Given that our previous study demonstrated that TET-mediated cytosine oxidation is

strongly associated with open chromatin regions during pancreatic differentiation [21], we performed

ATAC-seq to evaluate chromatin accessibility landscapes in WT and TET-deficient cells. We

assessed differential accessible regions (DARs) between WT_PP and TKO_PP cells from a total of

124,322 identified ATAC peaks and found that 4.3% of ATAC peaks significantly changed upon

TET deletion (Supplementary Table 2). We found that loss of TET led to an overall reduction in

chromatin accessibility, with 5,060 DARs showing reduced accessibility and only 290 DARs showing

increased accessibility in TKO_PP cells (Supplementary Fig. 1h). Moreover, DARs with decreased

accessibility were located primarily at distal regions (> 1 kb transcription start site (TSS)) (Fig. 1e).

Decomposition of chromatin accessibility and DEGs in TKO_PP cells revealed that downregulated

genes became less accessible (Fig. 1f). To enhance the biological insights of TET depletion-mediated

changes in chromatin accessibility, we determined DNA binding motifs within DARs and found that

regions with reduced accessibility were mostly enriched for motifs of pioneer TFs essential for

pancreas organogenesis, such as FOXAs and GATAs [23,24] (Fig. 1g).

TET deficiency induces a differentiation-associated loss of 5hmC at pioneer TF binding sites.

To gain insight into the TET-mediated DNA methylation network governing pancreatic

differentiation, we used cytosine-5-methylenesulfonate immunoprecipitation (CMS-IP) coupled with

high-throughput sequencing to profile DNA hydroxymethylation landscapes in WT_PP and TKO_PP

cells. As a consequence of TET deletion, we found genome-wide loss of DNA hydroxymethylation

in TET-deficient cells (Supplementary Fig. 2a, b). 94% of CMS-positive regions showed a significant

reduction in TKO_PP cells relative to WT_PP cells (Supplementary Fig. 2c). These differentially

hydroxymethylated regions showing reduced 5hmC (hypo-DHMRs) were primarily located at non-

promoter regions (> 1 kb from TSS) (Supplementary Fig. 2d).

Reduced DNA hydroxymethylation in TET-deficient cells could be a result of initial loss of

5hmC at the ES stage or unsuccessful oxidation of 5mC during lineage progression. To provide

biological insight into development-specific changes of hydroxymethylation upon TET inactivation,

we systematically compared hypo-DHMRs found in TKO_PP cells to previously identified regions

showing increased 5hmC during differentiation of hESCs toward pancreatic endoderm [21].

Specifically, 27,113 hypo-DHMRs with higher 5hmC levels in WT_PP cells than in WT_ES cells

(named 'differentiation-specific hypo-DHMRs') were first isolated from a total of 70,857 TKO hypo-

DHMRs (Fig. 2a). We subsequently clustered them into four categories based on the dynamic changes

of 5hmC across stages (Fig. 2b). In particular, one group gained 5hmC specifically at the PP stage

(PP-specific), whereas others gained 5hmC at the earlier DE (DE-to-PP) or GT (GT-to-PP) stages

and sustained hydroxymethylation until the PP stage. We found that GATA motifs were mostly

associated with the DE-to-PP cluster, in which gain of 5hmC began at the DE stage (Fig. 2c, purple).

By contrast, the PP-specific cluster displayed a prominent presence of binding sites for HNF6,

which is critical for pancreatic endocrine differentiation [25] (Fig. 2c, red). Most notably, FOXA motifs

were predominantly associated with the GT-to-PP(h) cluster in which 5hmC levels increased at the

GT stage and continued to be elevated until the PP stage (Fig. 2c, blue), whereas both FOXA and

GATA motifs were highly associated with the GT(h)-to-PP cluster in which 5hmC peaked at the GT

stage and subsequently decreased at the PP stage (Fig. 2c, brown). These results suggest a unique

temporal binding pattern of GATAs, FOXAs, and HNF6s associated with the dynamic distribution

of 5hmC during pancreatic differentiation, which is supported by a recent study demonstrating

sequential requirements for these TFs during transitions in pancreas development [4].

It was previously demonstrated that the pioneer TF FOXA2 is essential for enhancer priming and

chromatin remodeling during pancreatic differentiation [3,4]. By analyzing FOXA2-binding datasets

previously generated [4] from a similar stepwise pancreatic differentiation system, we found that

binding of FOXA2 corresponded well with the dynamic changes of 5hmC across differentiation

stages in the four 'differentiation-specific hypo-DHMRs' clusters (Fig. 2d). These results suggest that

TET-mediated hydroxymethylation strongly correlates with FOXA2 deposition during development.

Given our previous finding that 5hmC is positively associated with chromatin accessibility and

enhancer activity [21], we hypothesized that loss of TET inhibits chromatin opening and enhancer

activation at FOXA2 targets during pancreatic differentiation. Thus, we conducted ChIP-seq for

enhancer signatures H3K4me1 and H3K27ac in WT_PP and TKO_PP cells and performed integrated

analysis. We found remarkable decreases in ATAC and H3K27ac signals on FOXA2 targets

overlapping with 'differentiation-specific hypo-DHMRs' upon TET deletion (Fig. 2e), particularly at

distal regions (Supplementary Fig. 2e). Moreover, ATAC, H3K4me1, and H3K27ac signals were

substantially higher at FOXA2/hypo-DHMR-coexisting sites in pancreatic progenitors (Fig. 2e),

suggesting that 5hmC-associated FOXA2 targets are predominantly enriched for active chromatin.

Nevertheless, FOXA and GATA family members were expressed at similar levels in TKO and WT

cells (Supplementary Fig. 2f), suggesting that inhibition of TET did not alter the expression levels of

pioneer TFs. Collectively, these data reveal that TET deficiency results in a progressive failure of

5mC oxidation at a subset of FOXA2 targets essential for the establishment of active chromatin

during differentiation.

Pancreas-specific DMRs feature reduced chromatin activity during differentiation. To

investigate TET deficiency-mediated alterations in global DNA methylation, we performed whole-

genome bisulfite sequencing (WGBS) in WT_PP and TKO_PP cells. Over 86% of sequencing reads

were uniquely aligned to hg38 with a high sequencing depth of $24 \times$ on CpG dinucleotides and typical

bimodal distribution of methylation ratio in each sample (Supplementary Fig. 3a, b). The methylation

ratio (5mC/C) was depleted at TSS but enriched across gene coding regions, especially within

TKO_PP cells (Supplementary Fig. 3c). Among 26.6 million CpG sites detected in both WT_PP and

TKO_PP cells (depth $\geq$ 5), a total of 251,658 differentially methylated cytosines (DMCs; credible

difference of methylation ratio > 20%) were identified (Fig. 3a). Strikingly, 97.5% of DMCs were

hypermethylated (hyper-DMCs), suggesting that TET inhibition results in a pronounced gain of

methylation during pancreas differentiation. We observed enrichment of DMCs primarily at

intergenic regions and introns (Supplementary Fig. 3d), whereas substantial enrichment of hypo-

DMCs was also found in repeat elements, such as long interspersed elements, consistent with a

suggested connection between hypomethylation and activation of transposable elements [26,27]. Based

on an established link between DNA methylation and transcription regulation, we further calculated

changes in methylation levels on cis-regulatory elements previously identified in pancreatic

progenitors [3,28]. We found increased methylation at bivalent promoters, active enhancers, and, to a

lesser extent, poised enhancers (Fig. 3b). Notably, hypermethylation was preferentially enriched at

open chromatin (ATAC) as well as FOXA2, GATA4, GATA6, and PDX1 binding regions located

distally (Fig. 3b, Supplementary Fig. 3e). This analysis demonstrates that active regulatory elements

are hypermethylated upon TET depletion during pancreatic differentiation.

Our previous studies reveal that DNA demethylation is correlated with pancreatic endocrine

patterning [21]. To gain better insight into the role of TET-mediated methylation changes in pancreatic

differentiation, we analyzed differentially methylated regions (DMRs) by connecting at least three

consecutive DMCs [29] between TKO_PP and WT_PP cells. We found a total of 16,490 hyper-DMRs

and classified them into two categories based on their 5mC levels in pancreatic progenitors versus

hESCs [30] (Fig. 3c). Notably, more than half (n = 10,254) of hyper-DMRs exhibited less 5mC in

WT_PP cells than in hESCs (named pancreas-specific hyper-DMRs), whereas the others (n = 6,236)

showed similar methylation levels (non-pancreas hyper-DMRs) (Fig. 3c, Supplementary Fig. 3f).

Interestingly, we found that FOXA2, GATA4, and GATA6 were predominantly enriched at pancreas-

specific hyper-DMRs in pancreatic progenitors (Fig. 3d), suggesting that TET-dependent

demethylation is primarily associated with pioneer TF binding. We further compared ATAC,

H3K4me1, and H3K27ac signals at pancreas-specific hyper-DMRs with those at non-pancreatic

hyper-DMRs. We found that chromatin activity was markedly reduced in TET-deficient cells at

distally located hyper-DMRs in the pancreatic-specific group (Fig. 3e, Supplementary Fig. 3g).

Consistently, genomic loci displaying decreased accessibility showed a substantial increase in DNA

methylation upon depletion of TET (Fig. 3f). For example, we detected significant hypermethylation

at the distal regulatory elements of *PTF1A* and *NKX2.2* loci (Supplementary Fig. 3h). Altogether,

these analyses demonstrate that TET-dependent demethylation at enhancers and other distal

regulatory elements is essential for chromatin remodeling during pancreas development.

Additionally, we found that 774 type 2 diabetes (T2D)-associated islet hyper-DMRs, which were

recently identified [31], overlapped with hyper-DMRs found in TET-deficient cells (Supplementary

Table 3). Remarkably, four of the most significant T2D-associated islet hyper-DMRs annotated to

the *PDX1* and lncRNA *PDX1-AS1* loci were hypermethylated upon TET deletion during pancreatic

differentiation (Fig. 3g, green areas). We also uncovered an adjacent hyper-DMR (Fig. 3g, pink area)

that overlapped with the pioneer TF binding sites and showed decreased ATAC, H3K4me1, and

H3K27ac signals in a TET-dependent manner. Given that both *PDX1* and *PDX1-AS1* are

downregulated in islets from donors with T2D [32], these data suggest that TET inactivation leads to

remarkable DNA hypermethylation featuring loss of chromatin activity at regulatory elements, which

might be crucial for the development of diabetes.

TET1 is required in pancreatic β-cell specification. Our findings suggest that TET inactivation-induced aberrant methylation/hydroxymethylation at least in part contributes to defective β-cell specification. To determine the essential role of each TET family member in pancreatic differentiation, we further analyzed mutants with single (*TET1*, *TET2*, or *TET3*) or double (*TET1/2*, *TET2/3*, or *TET1/3*) *TET* knock-out. All mutant lines were able to induce the expression of *PDX1* and *NKX6.1* to levels comparable to WT (Supplementary Fig. 4a). However, only those retaining intact *TET1* expression (TET2KO, TET3KO, and TET2/3DKO) displayed proper induction of *PAX4* (Fig. 4a). In comparison with *TET2/TET3* double-deletion, inhibition of TET1 alone had significant effects on the formation of INS- and C-peptide (CPEP)-expressing β-cells but not GCG-expressing α-cells (Fig. 4b, Supplementary Fig. 4b). To further determine whether TET1 is responsible for β-cell induction, we restored *TET1* expression using lentivirus-mediated gene transduction to TET-depleted hESCs. *TET1*-repaired TKO cells showed a global increase in 5hmC levels, confirming the replenishment of hydroxymethylation (Supplementary Fig. 4c). Differentiation of the TKO-TET1 line toward pancreatic endocrine fate led to robust restoration of β-cell genes, including *NKX6.1*, *PAX4*, *INS*, and *CPEP* (Fig. 4c, d, Supplementary Fig. 4d), suggesting that TET1 is critical for β-cell specification.

To evaluate global transcriptome changes in response to *TET1* deletion, we performed RNA-seq of TET1KO cells at the PP stage. Integrated analysis of WT_PP, TKO_PP, and TET1KO_PP data revealed a total of 1,590 DEGs among the three lines (Supplementary Table 4). Specifically, 555 genes, including most pancreas developmental regulators such as *PDX1*, *SOX9*, *NKX2.2*, *NKX6.1*, *NEUROD1*, and *PAX6*, were downregulated in TKO_PP cells but not in TET1KO_PP cells (Fig. 4e), consistent with effective differentiation of TET1KO-hESCs into PDX1$^+$/NKX6.1$^+$ cells

(Supplementary Fig. 4a). However, 536 genes, including the β-cell fate determinants *PAX4*, *NKX6.2*,

and *FEV* [33-35], were significantly inhibited in TET1-deficient cells, implicating TET1 as responsible

for the activation of a subset of genes essential for β-cell identity. To further analyze the functional

consequences of TET1 inhibition, WT-, TET1KO-, and TKO-hESCs were differentiated to the PE

stage and subsequently engrafted into SCID-beige mice under the kidney capsules (Fig. 4f). Glucose-

stimulated human C-peptide secretion was determined 18 weeks post-transplantation. Notably, mice

engrafted with WT_PE cells produced substantial fasting C-peptide in serum (1605 ± 527 pg/ml) and

showed statistically significant glucose-stimulated C-peptide secretion (2,358 ± 839 pg/ml). By

contrast, mice transplanted with TET1KO_PE cells secreted extremely low amounts of basal C-

peptide (163 ± 122 pg/ml) and showed no response to glucose stimulation (149 ± 161 pg/ml). In

agreement with these functional results, excised WT_PE cell grafts were highly composed of insulin$^+$

β-cells (Supplementary Fig. 4e). The TET1KO_PE cell grafts displayed much less insulin content,

and only the δ-cell hormone somatostatin was detected in TKO_PE cell grafts. Together, these data

demonstrate that loss of TET1 impairs β-cell specification and maturation.

TET1 binding sites are distinct in pancreatic progenitors. To identify the direct targets of TET1,

we performed ChIP-seq for TET1 in pancreatic progenitors and identified a total of 8,476 peaks with

higher signals around TSSs (Supplementary Fig. 5a). Consistent with previous reports [18,36], TET1

was enriched at CpG islands (Supplementary Fig. 5b) and was primarily associated with

unmethylated genomic loci (Fig. 5a). To further examine the epigenetic features of TET1 target sites

in pancreatic progenitors, we centered all TET1 peaks and analyzed their accessibility, 5hmC, and

deposition of H3K4me1, H3K27ac, H3K4me3 [28], and H3K27me3 [28]. We found that TET1 binding

sites predominantly overlapped with active chromatin modifications at proximal (Fig. 5b, left panel) and distal (Fig. 5b, right panel) regions, suggesting that TET1 is primarily present at active regulatory elements in pancreatic progenitors.

Recent studies show that TET1 safeguards bivalent promoters in ESCs to ensure proper neuronal differentiation [18]. Therefore, it is interesting to examine whether TET1 occupies similar or different loci upon hESC differentiation toward pancreatic endoderm. We compared TET1 binding sites identified in pancreatic progenitors and pluripotent stem cells using TET1 ChIP-seq data previously generated from hESCs [18]. 4,463 TET1 peaks specific to pancreatic progenitors were referred to 'PP-specific', 22,177 binding sites specific to hESCs were called 'ES-specific', and 4,013 TET1 sites shared between the two groups were named 'common' (Fig. 5c). We then analyzed the annotations of their nearby genes with Genomic Regions Enrichment of Annotations Tool (GREAT) and found that only PP-specific peaks were enriched for terms of endocrine pancreas development (Supplementary Fig. 5c). In agreement with these findings, PP-specific sites prominently featured FOXA2 and CTCF motifs, whereas insulator CTCF and its paralogue BORIS motifs were mostly enriched at ES-specific sites (Fig. 5d). This analysis suggests that TET1 is involved in different biological functions in ESCs and lineage-committed cells. Interestingly, in contrast to the PP-specific sites, most of which were distally located (Fig. 5e), the 'common' TET1 binding sites were mainly located at proximal promoters and enriched for gene ontology annotations connected to essential biological processes, such as mRNA processing, biosynthesis, and translation (Supplementary Fig. 5c). We further examined chromatin accessibility and histone modifications of promoter and enhancer regions at proximal and distal TET1 binding sites. We found that TET1 bound more strongly

at both proximal and distal 'common' sites than at 'PP-specific' sites (Fig. 5f, TET1). Higher active

chromatin signals were also observed at the proximal (Fig. 5f, ATAC and H3K4me3) and distal (Fig.

5f, ATAC and H3K27ac) 'common' sites. The more active chromatin modifications at 'common'

TET1 targets may contribute to maintaining continually active transcripts essential for basic

biological functions in different types of cells. By contrast, cell-type-specific TET1 binding may

participate in chromatin remodeling during lineage specification.

Recent studies demonstrate that FOXA2 triggers chromatin opening on a subset of its potential

binding sites during endoderm differentiation [37]. In the present study, we found that FOXA2 target

sites acquiring 5hmC during pancreatic differentiation correlate with high levels of active chromatin

modifications (Fig. 2). We thus hypothesized that TET1 binds to a specific set of FOXA2 targets

associated with unique chromatin features. We isolated FOXA2 binding sites in pancreatic

progenitors [4] and analyzed DNA methylation, chromatin accessibility, and enhancer activity with or

without TET1 binding. The functional relevance of TET1 and FOXA2 co-binding is supported by

exceptionally low levels of methylation as well as more active chromatin modifications (ATAC,

H3K4me1, and H3K27ac), particularly at distal regions (Fig. 5g, Supplementary Fig. 5d). Moreover,

only the loci bound by both FOXA2 and TET1 were significantly enriched for terms associated with

pancreas development (Supplementary Fig. 5e), implicating a unique feature of TET1/FOXA2 co-

binding elements. Taken together, our data suggest that TET1 binds to a specific subset of FOXA2

targets that are confined mainly to active regulatory compartments during pancreatic differentiation.

TET1 is required for the *PAX4* enhancer to achieve a hypomethylated state. As TET1 depletion

inhibited the induction of genes essential for β-cell specification (Fig. 4), we reasoned that co-binding

of TET1 and FOXA2 at distal regulatory elements modulates focal DNA

methylation/hydroxymethylation and subsequent gene activation. To identify potential TET1

regulatory elements, we integrated gene expression, chromatin accessibility, and TET1/FOXA2

binding profiles into a comprehensive analysis. We found that a putative *PAX4* enhancer (~4.0 kb

upstream of the TSS), where TET1/FOXA2 co-bound, displayed hypermethylation and decreased

accessibility in TET-deficient cells (Fig. 6a, pink area). We next examined whether the deletion of

TET1 alone results in hypermethylation at this site. Locus-specific methylation was determined in

TET1KO_PP, TET2/3DKO_PP, TKO_PP, and WT_PP cells using a glucosylation and digestion-

based method followed by qPCR analysis with two independent pairs of primers (PAX4-P1 and

PAX4-P2) located within the TET1/FOXA2 co-binding site (Fig. 6a). Notably, the percentage of

5mC was increased to 35–50% for TET1KO and 60–65% for TKO compared with WT but remained

unchanged for TET2/3DKO (Fig. 6b, left panel). A corresponding decrease in unmethylated cytosine

content was observed, with values of 45–65% and ~35% of C in TET1KO and TKO samples,

respectively (Fig. 6b, right panel). As a housekeeping control, we examined methylation content at a

nearby region using the PAX4-P3 primer, and, as expected, no significant differences in 5mC and C

were observed in any samples.

Furthermore, we determined methylation contents at the same *PAX4* locus in TET1-repaired

TKO cells. Consistent with the results found in TET2/3DKO cells, no detectable differences in 5mC

and C were found among TKO-TET1 samples (Supplementary Fig. 6). Together with the increased

expression of *PAX4* in TKO-TET1 cells (Fig. 4d), we conclude that TET1 is responsible for the

establishment or maintenance of a hypomethylated state at the *PAX4* enhancer, which is essential for

transcription activation of *PAX4*.

Discussion

In the present work, we dissected the roles of TET proteins in pancreatic endocrine commitment

based on a stepwise hESC differentiation system. We found that the loss of all three TET family

members significantly impaired the differentiation of pancreatic β-cells. Furthermore, we discovered

that locus-specific hypermethylation was associated with genes essential for β-cell specification and

maturation, such as *PAX4*[33], *PDX1*[38], and *NKX2.2*[39]. The reintroduction of *TET1* in TET-deficient

cells effectively reversed hypermethylation and restored the expression of *PAX4*. We further

demonstrated that TET1 functions as an upstream epigenetic regulator of *PAX4* through direct

binding at a putative *PAX4* enhancer to preserve its unmethylated status, thereby potentiating *PAX4*

expression to adopt β-cell fate during endocrine lineage commitment. Consistently, we observed

striking increases in methylation at the *PAX4* enhancer in TET1 knock-out cells but not in

TET2/TET3 DKO cells, suggesting that TET1 epigenetically regulates induction of the β-cell

program in a locus-specific manner. Moreover, despite the successful induction of *PDX1* and *INS*

upon deletion of TET1 alone, mice receiving TET1KO cell grafts showed a persistent defective

insulin response to glucose, implying that TET1 is also essential for β-cell maturation.

Whereas the specification of β-cells was strongly influenced by depletion of *TET1*, we did not

observe a significant impact on the expression of *ARX*, a critical α-cell fate determinant[40]. In contrast

to *PAX4*, no hyper-DMRs were identified within the *ARX* locus, where methylation levels were nearly

undetectable in pancreatic progenitors. Previous studies demonstrate that several CpG-rich sites of

Arx, including one site close to TSS and another site 2 kb upstream of TSS, are heavily methylated

in adult β-cells but not α-cells [41]. Moreover, pharmacological inhibition of DNA methyltransferases,

Dnmts, in pancreatic progenitors promotes α-cell specification [42], whereas deletion of *Dnmt1* in β-

cells results in demethylation and depression of *Arx* [41]. We thus speculate that *ARX* is hypomethylated

in pancreatic progenitors in a TET-independent manner. The *ARX* locus maintains an unmethylated

state when progenitors differentiate into α-cells, whereas it becomes hypermethylated once cells

commit to β-cell fate in the presence of DNMTs and other β-cell-specific factors. In line with this

hypothesis are findings that Nkx2.2 recruits Dnmt3a to the *Arx* promoter to repress its expression in

β-cells [43]. In the future, it will be interesting to explore whether there are differences in the complete

epigenetic landscapes of endocrine progenitors, which subsequently differentiate into α- or β-cells.

TET dioxygenases are critical for lineage induction in a cell-type-specific manner [44]. How TETs

recognize lineage-specific regulatory elements and modulate chromatin remodeling during pancreas

development remains unknown. Here, we addressed these questions by performing analyses

integrating gene expression with multiple chromatin features, including DNA methylation,

hydroxymethylation, chromatin accessibility, and histone modifications of promoters and enhancers.

We found extensive hypermethylation in TET-deficient cells that differentiated to the pancreatic

progenitor stage. A significant portion of hyper-DMRs were distributed in a cell-type-specific manner,

in which they were enriched for the binding of FOXA2, a pioneer TF essential for pancreatic

endoderm differentiation, and showed remarkable decreases in chromatin activity upon TET

inactivation. It is noteworthy that we unveiled a unique TF binding pattern associated with a stepwise

increase of 5hmC during pancreatic differentiation. The presence of GATA, FOXA, and HNF6

binding motifs in a temporal manner suggests that TET proteins are recruited to chromatin by a set

of lineage-specific TFs sequentially expressed during pancreatic differentiation. More intriguingly,

nearly 800 hyper-DMRs recently identified in T2D pathogenesis [31] overlapped with hyper-DMRs

found in TET-deficient cells in the present study, implying that TET dioxygenases are crucial for β-

cell function.

During lineage specification, chromatin structure is dynamically changed between 'closed' and

'open' states. Open chromatin regions such as primed and active enhancers are accessible for TF and

epigenetic modulator binding to initiate gene transcription. Proper chromatin remodeling is believed

to be at least partly triggered by pioneer TFs, which can directly bind to nucleosomal DNA [6]. It was

previously suggested that pioneer TF FOXA2 initiates chromatin remodeling and enhancer priming

during pancreatic differentiation [3,4]. Interestingly, our analyses revealed that changes in 5hmC mirror

the dynamic binding of FOXA2 in cells differentiated from hESCs through defined lineage

intermediates toward pancreatic endocrine fate. We showed that chromatin activity was markedly

decreased at the FOXA2 binding sites associated with DHMRs in TET-deficient cells, suggesting

that lineage-specific hydroxymethylation is processed at a subset of FOXA2 target sites to influence

chromatin opening.

How pioneer TFs initiate chromatin opening is still mostly unknown. One suggestion is that

FOXA2 can recruit nucleosome remodeling complexes SWI/SNF to alter the surrounding chromatin

structure [45]. Our present study demonstrates that TET1 binds to two distinct groups of regulatory

elements: universal and lineage-specific TET1 sites. The fact that 1) pancreatic progenitor-specific

TET1 targets are enriched for the FOXA2 binding motif and 2) co-occupancy of TET1 and FOXA2

at distal regulatory elements is associated with more active chromatin suggests that TET1 residing in

specific genomic loci modulates FOXA2-mediated chromatin opening. Several studies suggest that TET can be directly recruited by pioneer TFs, such as FOXA1 and PU.1, to lineage-specific enhancers to facilitate local demethylation and gene induction [46,47]. However, a recent study found that chromatin opening requires cooperative binding of FOXA2 and additional TFs, such as GATA4, during endoderm differentiation [37]. This suggests complex cross-talk between pioneer TFs and chromatin remodelers in endodermal lineage specification. Further investigation is warranted to determine whether TET1 is recruited to distinct FOXA2-binding sites by lineage-specific TFs to synergistically regulate chromatin remodeling.

Methods

Cell lines. The human embryonic stem cell line H1 was obtained from WiCell Research Institute. H1 cells were maintained on Matrigel (Corning) in mTeSR1 (STEMCELL Technologies) and passaged every for 4-5 days using 0.5 mM EDTA and 3 mM NaCl. Mycoplasma detection was regularly performed by the Stem Cell Core Facility at the University of Macau.

Pancreatic differentiation. All differentiation experiments were repeated at least three times with three individual clones of the same phenotype. Pancreatic differentiation was performed as previously described with slight modifications [48]. In brief, hESCs were dissociated with Accutase (eBioscience) and seeded at a density of 19,000 cells/cm^2 on Matrigel in mTeSR1 supplemented with 10 μM Rho-associated protein kinase inhibitor Y-27632 (Miltenyi Biotec). Upon reaching 95% confluence, cells were exposed to differentiation medium with daily media feeding.

ES-to-DE (3 d): hESCs were exposed to 100 ng/ml Activin A (PeproTech) and 25 ng/ml Wnt-3a (PeproTech) in basal medium-I containing MCDB 131 medium (Gibco), 1.5 g/l NaHCO$_3$, 1×

GlutaMAX (Gibco), 10 mM glucose, and 0.5% BSA for 1 day. For 2 additional days, cells were

cultured in basal medium-I further supplemented with 100 ng/ml Activin A.

DE-to-GT (3 d): DE intermediates were incubated in basal medium-I supplemented with 50

ng/ml FGF7 (PeproTech) for 3 days.

GT-to-PP (5 d): GT intermediates were cultured for 3 days in basal medium-II containing MCDB

131 medium, 2.5 g/l $NaHCO_3$, 1× GlutaMAX, 10 mM glucose, 2% BSA, and 1:200 ITS-X (Gibco),

which was further supplemented with 50 ng/ml FGF7, 0.25 µM hedgehog inhibitor SANT-1 (Sigma),

1 µM retinoic acid (Sigma), 100 nM BMP inhibitor LDN193189 (Stemgent), and 200 nM PKC

activator TPB (Millipore). After 3 days of culture, cells were treated for 2 days with 2 ng/ml FGF7,

0.25 µM SANT-1, 0.1 µM retinoic acid, 200 nM LDN193189, and 100 nM TPB in basal medium-II.

PP-to-PE (3 d): PP intermediates were differentiated in basal medium-III containing MCDB 131

medium, 1.5 g/l $NaHCO_3$, 1× GlutaMAX, 20 mM glucose, 2% BSA, and 1:200 ITS-X, which was

further supplemented with 0.25 µM SANT-1, 0.05 µM retinoic acid, 100 nM LDN193189, 1 µM T3

(3,3',5-Triiodo-L-thyronine sodium salt, Sigma), 10 µM ALK5 inhibitor II (Enzo Life Sciences), 10

µM $ZnSO_4$, and 10 µg/ml heparin (Sigma).

Animal experiments. All animal experiments were approved by the University of Macau Animal

Ethics Committee. Immunocompromised SCID-beige mice were obtained from Charles River and

maintained under a 12-h light/dark cycle with free access to standard mouse diet. For transplantation,

~5 million hESC-derived d14 cells were enzymatically dissociated and aggregated using ultra-low

adherent plates (Corning) in basal medium-III further supplemented with 100 nM LDN193189, 1 µM

T3, 10 µM ALK5 inhibitor II, 10 µM $ZnSO_4$, 100 nM γ-secretase inhibitor XX (Millipore), and 10

µM Y-27632 for 1 day. Cell aggregates were then transplanted into 6-week-old SCID-beige mice

under the kidney capsule as previously described [49].

Glucose-stimulated human C-peptide secretion was assessed with mice 18 weeks post-

transplantation. Blood samples were collected after overnight fasting and 30 min following an

intraperitoneal injection of glucose (2 g/kg body weight). Human C-peptide levels in isolated plasma

were quantified using the STELLUX Chemi Human C-peptide ELISA kit (ALPCO Diagnostics)

according to the manufacturer's instructions.

Generation of *TET* knock-out lines. *TET* knock-out hESCs were generated using CRISPR/Cas9

technology. gRNAs (Supplementary Table 5) were designed to target the sequences encoding exon 7

of *TET1*, exon 3 of *TET2*, or exon 3 of *TET3* and cloned into pX330 vector (Addgene #42230) as

previously described [50]. Constructs containing validated gRNAs were electroporated together with a

vector expressing puromycin into hESCs using the P3 Primary Cell 4D-Nucleofector X kit (Lonza)

following the manufacturer's instructions. The electroporated cells were plated at ~2,000 cells/cm^2

and cultured on Matrigel in mTeSR1 supplemented with 10 μM Y-27632 for 2 days. Successfully

transfected cells were selected with 1 μg/ml puromycin in mTeSR™1 and allowed to expand to form

visible colonies from a single cell. Subsequently, clonal colonies were manually picked and reseeded

individually into 24-well plates. The amplified colonies were analyzed by Sanger sequencing at

targeted loci for the presence of Indel mutations.

TET1 expression cell lines. TET1 lentivirus expression plasmid was constructed by amplifying the

human *TET1* ORF from FH-TET1-pEF (Addgene #49792) and inserting it into *KpnI* and *BamHI* sites

of the lentiCRISPR v2 vector (Addgene #52961). Lentivirus particles were prepared as previously

described [51]. For viral transduction, TKO-hESCs were grown on Matrigel in mTeSR1 using 24-well

plates and treated with 6 µg/ml polybrene for 15 min at 37°C. Concentrated lentivirus particles (10

µl, 1×10^6 TU/ml) were added to cell culture and incubated overnight at 37°C. On the next day, viral

infection was repeated (30 µl, 1×10^6 TU/ml) to increase transduction efficiency. Infected cells were

cultured in mTeSR1 for 2 days and then exposed to 1 µg/ml puromycin for 10 days. The TKO line

TKO-TET1, in which an intact *TET1* cDNA was expressed, was amplified and frozen down. Control

cells (TKO-CRTL) carrying the lentiCRISPR v2 plasmid were established with the same

experimental procedure.

Locus-specific detection of 5mC. Detection of 5mC content at particular CCGG sites was performed

using the Epimark 5hmC and 5mC analysis kit (New England Biolabs) following the manufacturer's

instructions. Briefly, genomic DNA was extracted using the DNeasy Blood and Tissue kit (Qiagen)

followed by RNase treatment. RNase-treated DNA samples were incubated with T4 β-

glucosyltransferase at 37°C for 16 h. The glycosylated DNA was subsequently digested with *MspI* or

HpaII for 8 h at 37°C. Samples were treated with Proteinase K at 40°C for 30 min and then at 95°C

for 10 min to inactivate the enzymes. Site-specific methylation contents were examined by RT-qPCR

using the primers listed in Supplementary Table 6. The percentage of 5mC and unmodified cytosine

were calculated using the comparative C_t method.

5hmC dot blot assay. Genomic DNA was extracted using the DNeasy Blood and Tissue kit (Qiagen)

according to the manufacturer's instructions. DNA was denatured in 1 M NaOH supplemented with

25 mM EDTA at 95°C for 10 min and then neutralized with 2 M ice-cold ammonium acetate for 10

min. Two-fold serial dilutions of the DNA samples were spotted onto nitrocellulose membrane. The

air-dried membrane was fixed with UV irradiation (CL-1000 UV crosslinker, Ultra-Violet Products),

blocked with 5% non-fat milk, and incubated with a rabbit anti-5hmC antibody (1:10,000, Active

Motif) followed by an HRP-conjugated anti-rabbit antibody (1:5,000, Jackson ImmunoResearch).

Signal was visualized with SuperSignal West Pico PLUS chemiluminescent substrate (Thermo

Fisher Scientific). The same membrane was subsequently stained with 0.02% methylene blue in 0.3

M sodium acetate to ensure equal loading of input DNA.

RNA isolation for real-time quantitative PCR. Total RNA was extracted using the RNeasy Plus

Mini kit (Qiagen) according to the manufacturer's instructions. cDNA was synthesized using the

PrimeScript RT reagent kit (Takara). Real-time quantitative PCR was performed in triplicate using

the SYBR Premix Ex Taq (Tli RNase H Plus) kit (Takara). The expression of *TBP* was used for the

normalization of mRNA expression. All primers used for RT-qPCR were listed in Supplementary

Table 6.

Immunocytochemical analysis. Differentiated cells were fixed in 4% paraformaldehyde for 30 min

at room temperature, washed, and then permeabilized with 0.15% Triton X-100 at room temperature

for 1 h. Following blocking with 5% normal donkey serum (Jackson Immuno Research Laboratories),

samples were incubated with primary antibodies at 4°C overnight and then appropriate secondary

antibodies for 1 h at room temperature. Images were acquired using the Zeiss Axio Observer

microscope. For transplant grafts, tissues were fixed with 4% paraformaldehyde overnight at 4°C,

washed with PBS, and subsequently exposed to 30% sucrose overnight at 4°C. Samples were

mounted with Optimal Cutting Temperature Compound (Tissue-Tek) and sectioned at 10 μm.

Immunofluorescent staining was performed on cryosections as described above. Antibodies used for

immunofluorescent staining are listed in Supplementary Table 7.

Flow cytometry analysis. Cells were enzymatically dissociated into single cells and fixed with 4%

paraformaldehyde for 20 min at 4°C. Fixed cells were permeabilized with Perm/Wash Buffer (BD

Biosciences) and stained with primary antibodies (Supplementary Table 7) diluted in Perm/Wash

Buffer overnight at 4°C. Cells were subsequently washed, stained with appropriate secondary

antibodies for 1 h at room temperature, and assessed using an Accuri C6 flow cytometer (BD

Biosciences). Data were analyzed using FlowJo software (TreeStar).

RNA-seq library construction and data analysis. Total RNA was isolated using Trizol reagent

following the manufacturer's instructions. RNA integrity was determined using an Agilent 2100

Bioanalyzer (Agilent Technologies). Subsequently, polyA-tailed RNA was selected using Dynabeads

oligo(dT) (Thermo Fisher Scientific), and libraries were prepared using the NEBNext Ultra RNA

Library Prep kit for Illumina (New England Biolabs). Libraries were subjected to high-throughput

sequencing on an Illumina HiSeq 2500 system (150 bp, paired-end) at Novogene (Tianjin, China).

To process the sequencing data, low-quality bases and adaptor were trimmed using TrimGalore

v.0.5.0 (options: --quality 20 and --length 50, https://github.com/FelixKrueger/TrimGalore). Clean

reads were aligned to the hg38 reference genome using STAR v.2.5.3 [52] with default parameters, and

only uniquely mapped reads were used for downstream analysis. A count matrix for each gene was

generated using htseq-count (HTSeq package [53]). DESeq2 [54] was used to identify significant DEGs

in knock-out samples compared with WT samples at different differentiation stages (fold change (FC)

≥ 2; False Discovery Rate (FDR) < 0.05). Hierarchical cluster analysis [55] of the union DEGs was used

to determine stage-specific signature genes. The 'ClusterProfilter' package in R [56] was used for the

functional enrichment analysis of DEGs in KEGG pathways.

ATAC-seq library preparation and data analysis. ATAC-seq libraries were prepared as previously

described [57]. In brief, cells were enzymatically dissociated and lysed in lysis buffer (10 mM Tris-HCl,

pH 7.4, 10 mM NaCl, 3 mM MgCl2, 0.1% IGEPAL CA-630). Immediately after centrifugation,

transposition reactions were carried out by adding Tn5 transposes from the Illumina Nextera DNA

library preparation kit to the isolated nuclei and incubation at 37°C for 30 min. DNA fragments were

purified using the MinElute PCR Purification kit (Qiagen) and amplified using the KAPA real-time

library amplification kit (Roche). Libraries were purified using PCRClean DX beads (Aline

Biosciences) and subsequently subjected to high-throughput sequencing on an Illumina NextSeq 500

instrument (75 cycle, paired-end).

Adaptor trimming of raw reads was performed using TrimGalore v0.5.0 (options: --quality 20

and --length 20), and high-quality ($Q \geq 20$) reads were uniquely aligned to the hg38 reference genome

using Bowtie2 with the '--very-sensitive' option. Reads mapped to mitochondrial DNA and PCR

duplicate reads were removed, and uniquely mapped reads were extracted for downstream analysis.

Genrich v.0.5 (https://github.com/jsh58/Genrich) with ATAC-seq mode (option: -j,-q 0.01) was

applied for each sample (with two biological replicates) to call ATAC peaks. In total, 52,817 and

38,697 peaks were detected in WT_PP and TKO_PP cells, respectively. Bedtools intersect was used

to count the reads falling into 124,322 non-overlapping peak regions, and significant DARs were

detected using DESeq2 by the criteria $FC \geq 2$ and $FDR < 0.05$. Volcano plots were generated using

the R package ggplot2. Motif annotation of DARs was performed using HOMER software [58].

CMS-IP-seq library preparation and data analysis. CMS-IP-seq libraries were performed as

previously described with minor modifications (Huang et al., 2012). Purified genomic DNA (with 5%

mouse DNA and 0.5% lambda DNA spike-in) was sheared to 200–500 bp fragments using a M220

Focused-ultrasonicator (Covaris). Bisulfite-converted DNA libraries with methylated adapters were

enriched using an in-house anti-CMS antibody bound to protein A/G Dynabeads. Amplified libraries

were purified by AmpuXP beads (Beckman Coulter) and then sequenced using the Illumina NextSeq

(75 and 40 cycles, single-end) system.

Analysis of CMS-IP data was performed by in-house software 'CMSIP' v0.1.1

(https://github.com/lijinbio/cmsip). Briefly, raw reads were mapped to hg38 and mm10 genome

references using bsmap v.2.89 [59] (options: -n 1 -q 3 -r 0). After removing PCR duplicates and reads

mapped to both human and spike-in mouse genome, species-specific reads were used to perform

normalization for each sample according to the spike-in size factors. The mean wigsum for every

200-bp window of the whole human genome was used to call the hydroxymethylation-enriched peak

regions for each group against input. In total, 75,324 and 503 peaks were detected in WT_PP and

TKO_PP samples, respectively. Differentially DHMRs between WT_PP and TKO_PP were

identified (g-test; FDR < 0.05). GREAT analysis with single-nearest genes option was used to

perform functional annotation of hypo-DHMRs.

CMS-IP-seq datasets of hESCs and multiple pancreatic lineage intermediates (DE, GT, and PP)

were downloaded from GSE97992 [21] and mapped to the hg38 reference genome using bsmap with

the same parameters as in the previous analysis. Bam2wig.py in RSeQC [60] was used to transform the

bam file to normalized bigWig files (option: -t 2000000000). The average 5hmC signals within each

hypo-DHMR for all stages were calculated using deepTools [61]. Compared with hESCs, increased

5hmC peaks were defined as peaks in PP cells with a ≥ 1.5-fold increase.

WGBS library preparation and data analysis. Genomic DNA was isolated using the DNeasy

Blood & Tissue kit (Qiagen). Library preparation and high-throughput sequencing were conducted

by BGI (Shengzhen, China). In brief, purified genomic DNA (with 1% unmethylated lambda DNA

spike-in, Promega) was sheared to a fragment size of 100–700 bp (primary size 250 bp). Sheared

DNA was ligated with methylated adaptors (MGIEasy WGBS Adapters-16, MGI) and subjected to

bisulfite conversion using the EZ DNA Methylation-gold kit (Zymo Research). Bisulfite-converted

DNA was amplified with 13 PCR cycles and purified by AMPure XP beads (Beckman Coulter). All

libraries were sequenced on an MGISEQ-2000 system (100-cycle, paired-end).

For data analysis, paired-end 100 bp reads were mapped against hg38 using bsmap v.2.89 [59] with

paired mode (options: -n 1 -q 3 -r 0), and only uniquely mapped reads were retained. More than 26.6

million CpG sites with coverage of $\geq$ 5 reads were detected in both WT_PP and TKO_PP samples,

which were used for downstream analyses. BSeQC [62] and the mcall module in MOABS [29] was applied

to perform quality control and calculate the methylation ratio for each CpG site (options: --

trimWGBSEndRepairPE2Seq 40). Bisulfite conversion efficiencies were estimated using spike-in

unmethylated lambda phage DNA. The Mcomp module was used to call significant DMCs and DMRs

(absolute credible difference of DNA methylation ratio > 20% and adjusted $p < 0.05$). The output

bedGraph files from mcall included single-base resolution DNA methylation ratios, which were

transformed into bigwig file format. The bigwig files were uploaded to the UCSC genome browser

for visualization. Motif enrichment analysis of DMRs was performed using HOMER software,

functional annotation was performed using GREAT [20] with default settings, and many plots related

to WGBS was performed using MMINT.

ChIP-seq library preparation and data analysis. ChIP-seq was performed for TET1 (WT_PP cells),

H3K4me1 (WT_PP and TKO_PP cells), and H3K27ac (WT_PP and TKO_PP cells). For TET1 ChIP-

seq, ~5×10^7 cells were crosslinked, washed, and snap-frozen following the Cell Fixation protocol

from Active Motif. TET1 ChIP-seq library preparation and sequencing were conducted by Active

Motif. Chromatin immunoprecipitation of H3K4me1 and H3K27ac were performed as previously

described [28]. WT_PP and TKO_PP cells were fixed, washed, and lysed in nuclear lysis buffer (50

mM Trish HCl, pH 8.0, 5 mM EDTA, 1% SDS, and 1× protease inhibitor cocktail). Chromatin was

sheared to 200–500-bp fragments using Bioruptor (Diagenode), and DNA fragments were

precipitated with appropriate antibodies. ChIP-seq libraries were prepared using the NEBNext Ultra

II DNA Library Prep kit (New England Biolabs) following the manufacturer's instruction and

subjected to high-throughput sequencing on an Illumina HiSeq 2500 system (150 bp, paired-end) at

Novogene (Tianjin, China).

Quality control and alignment of raw reads of ChIP-seq data were performed similarly to the

ATAC-seq data analysis described above. TrimGalore (options: --quality 20 and --length 50) was

used to remove the adaptor, and only uniquely mapped reads were retained. Bam2wig.py was used

to transform the bam file to normalized bigWig files (option: -t 2000000000). For WT_PP TET1

ChIP-seq, Macs2 [63] was used to call ChIP-seq enriched peak regions with default parameters.

Integration of analyses. RNA-seq, WGBS, CMS-IP-seq, ATAC-seq and ChIP-seq library

preparations were performed as previously described. Detail experimental procedures can be found

in Supplementary Materials. The number of mapped reads and mapped ratios are listed in

Supplementary Table 8. To compare 5hmC and FOXA2 enriched signals, FOXA2 ChIP-seq datasets

of multiple pancreatic lineage intermediates were downloaded from GSE117136. Data were analyzed

as described above. Deeptools was used to calculate average signals at pancreas-specific hypo-

DHMRs (up- and down- 3 kb) in each group. The FOXA2 bound sites were detected using Macs2 [63].

To compare DNA methylation between TKO_PP and WT_PP cells at annotated genomic

features, ChIP-seq datasets of FOXA2, GATA4, GATA6, and PDX1 were downloaded from

GSE117136. Bivalent promoters and poised and active enhancers in hESC-derived pancreatic

progenitors were downloaded from EMTAB1086 and GSE54471, respectively. The analysis was

similar to that described above.

Quantification and statistical analysis. All statistical analyses were performed using GraphPad

Prism software. Student's unpaired two-tailed t-tests were used for qRT-PCR and flow cytometry

experiments. Student's paired one-tailed t-tests were used for the glucose-stimulated human C-

peptide secretion experiment. Quantification data are presented as mean ± SD. For all statistical

analyses, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$. Statistical analyses for RNA-seq,

WGBS, CMS-IP-seq, and ChIP-seq data are described in the corresponding sections.

Data availability

WGBS, CMS-IP, RNA-seq, ATAC-seq, and ChIP-seq data from this study were submitted to the NCBI Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE146486. All relevant data supporting the critical findings of this study are available within the article and its supplementary information files or from the corresponding author upon reasonable request.

Acknowledgments

We thank the University of Macau, Faculty of Health Sciences, Animal Research Facility for animal housing. This work was supported by the National Natural Science Foundation of China (NSFC 31701276) and the University of Macau Multi-Year Research Grant (MYRG2016-00065-FHS) to R.X.

Author contributions

R.X. conceived the project. R.X. and D.S. directed and oversaw the project. X.W. performed most experiments and collected data. J.F.L. performed computational analysis for WGBS, RNA-seq, CMS-IP-seq, ATAC-seq, and ChIP-seq. M.L. prepared the CMS-IP library and performed sequencing. Q.L. assisted with the animal studies. J.L. provided expert advice on bioinformatic analyses. Y.H. critically reviewed the manuscript. R.X., X.W., D.S., and J.F.L. wrote the manuscript; all other authors provided editorial advice.

634 **Competing interests**

635 The authors declare that no conflicts of interest exist.

References

1. Buecker, C. & Wysocka, J. Enhancers as information integration hubs in development: lessons from genomics. *Trends Genet.* **28**, 276-284 (2012).

2. Long, H. K., Prescott, S. L. & Wysocka, J. Ever-changing landscapes: transcriptional enhancers in development and evolution. *Cell* **167**, 1170-1187 (2016).

3. Wang, A. et al. Epigenetic priming of enhancers predicts developmental competence of hESC-derived endodermal lineage intermediates. *Cell Stem Cell* **16**, 386-399 (2015).

4. Lee, K. et al. FOXA2 is required for enhancer priming during pancreatic differentiation. *Cell Rep* **28**, 382-393 (2019).

5. Cirillo, L. A. & Zaret, K. S. An early developmental transcription factor complex that is more stable on nucleosome core particles than on free DNA. *Mol. Cell* **4**, 961-969 (1999).

6. Cirillo, L. A. et al. Opening of compacted chromatin by early developmental transcription factors HNF3 (FoxA) and GATA-4. *Mol. Cell* **9**, 279-289 (2002).

7. Iwafuchi-Doi, M. et al. The pioneer transcription factor foxA maintains an accessible nucleosome configuration at enhancers for tissue-specific gene activation. *Mol. Cell* **62**, 79-91 (2016).

8. Serandour, A. A. et al. Epigenetic switch involved in activation of pioneer factor FOXA1-dependent enhancers. *Genome Res.* **21**, 555-565 (2011).

9. Donaghey, J. et al. Genetic determinants and epigenetic effects of pioneer-factor occupancy. *Nat. Genet.* **50**, 250-258 (2018).

10. He, Y. F. et al. Tet-mediated formation of 5-carboxylcytosine and its excision by TDG in mammalian DNA. *Science* **333**, 1303-1307 (2011).

11. Ito, S. et al. Role of Tet proteins in 5mC to 5hmC conversion, ES-cell self-renewal and inner cell mass specification. *Nature* **466**, 1129-1133 (2010).

12. Ito, S. et al. Tet proteins can convert 5-methylcytosine to 5-formylcytosine and 5-carboxylcytosine. *Science* **333**, 1300-1303 (2011).

13. Tahiliani, M. et al. Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* **324**, 930-935 (2009).

14. Zhang, H. et al. TET1 is a DNA-binding protein that modulates DNA methylation and gene transcription via hydroxylation of 5-methylcytosine. *Cell Res.* **20**, 1390-1393 (2010).

15. Tsagaratou, A. et al. Dissecting the dynamic changes of 5-hydroxymethylcytosine in T-cell development and differentiation. *Proc. Natl. Acad. Sci. USA* **111**, E3306-3315 (2014).

16. Tan, L. et al. Genome-wide comparison of DNA hydroxymethylation in mouse embryonic stem cells and neural progenitor cells by a new comparative hMeDIP-seq method. *Nucleic Acids Res.* **41**, e84 (2013).

17. Kim, M. et al. Dynamic changes in DNA methylation and hydroxymethylation when hES cells undergo differentiation toward a neuronal lineage. *Hum. Mol. Genet.* **23**, 657-667 (2014).

18. Verma, N. et al. TET proteins safeguard bivalent promoters from de novo methylation in human embryonic stem cells. *Nat. Genet.* **50**, 83-95 (2018).

19. Ko, M. et al. Ten-Eleven-Translocation 2 (TET2) negatively regulates homeostasis and differentiation of hematopoietic stem cells in mice. *Proc. Natl. Acad. Sci. USA* **108**, 14566-

14571 (2011).

20.   Fang, S. et al. Tet inactivation disrupts YY1 binding and long-range chromatin interactions
during embryonic heart development. *Nat. Commun.* **10**, 4297 (2019).

21.   Li, J. et al. Decoding the dynamic DNA methylation and hydroxymethylation landscapes in
endodermal lineage intermediates during pancreatic differentiation of hESC. *Nucleic Acids*
Res. **46**, 2883-2900 (2018).

22.   Lu, F. L., Liu, Y. T., Jiang, L., Yamaguchi, S. & Zhang, Y. Role of Tet proteins in enhancer
activity and telomere elongation. *Genes Dev.* **28**, 2103-2119 (2014).

23.   Rodriguez-Segui, S., Akerman, I. & Ferrer, J. GATA believe it: new essential regulators of
pancreas development. *J. Clin. Invest.* **122**, 3469-3471 (2012).

24.   Kaestner, K. H. The FoxA factors in organogenesis and differentiation. *Curr. Opin. Genet.*
Dev. **20**, 527-532 (2010).

25.   Zhang, H. et al. Multiple, temporal-specific roles for HNF6 in pancreatic endocrine and ductal
differentiation. *Mech. Dev.* **126**, 958-973 (2009).

26.   Lopez-Moyado, I. F. et al. Paradoxical association of TET loss of function with genome-wide
DNA hypomethylation. *Proc. Natl. Acad. Sci. USA* **116**, 16933-16942 (2019).

27.   Walsh, C. P., Chaillet, J. R. & Bestor, T. H. Transcription of IAP endogenous retroviruses is
constrained by cytosine methylation. *Nat. Genet.* **20**, 116-117 (1998).

28.   Xie, R. et al. Dynamic chromatin remodeling mediated by polycomb proteins orchestrates
pancreatic differentiation of human embryonic stem cells. *Cell Stem Cell* **12**, 224-237 (2013).

29.   Sun, D. et al. MOABS: model based analysis of bisulfite sequencing data. *Genome Biol.* **15**,
R38 (2014).

30.   Consortium, E. P. An integrated encyclopedia of DNA elements in the human genome. *Nature*
489, 57-74 (2012).

31.   Volkov, P. et al. Whole-genome bisulfite sequencing of human pancreatic islets reveals novel
differentially methylated regions in type 2 diabetes pathogenesis. *Diabetes* **66**, 1074-1085
(2017).

32.   Akerman, I. et al. Human pancreatic beta cell lncRNAs control cell-specific regulatory
networks. *Cell Metab.* **25**, 400-411 (2017).

33.   Sosa-Pineda, B., Chowdhury, K., Torres, M., Oliver, G. & Gruss, P. The Pax4 gene is essential
for differentiation of insulin-producing beta cells in the mammalian pancreas. *Nature* **386**,
399-402 (1997).

34.   Nelson, S. B., Schaffer, A. E. & Sander, M. The transcription factors Nkx6.1 and Nkx6.2
possess equivalent activities in promoting beta-cell fate specification in Pdx1+ pancreatic
progenitor cells. *Development* **134**, 2491-2500 (2007).

35.   Ohta, Y. et al. Convergence of the insulin and serotonin programs in the pancreatic beta-cell.
Diabetes **60**, 3208-3216 (2011).

36.   Wu, H. et al. Dual functions of Tet1 in transcriptional regulation in mouse embryonic stem
cells. *Nature* **473**, 389-393 (2011).

37.   Cernilogar, F. M. et al. Pre-marked chromatin and transcription factor co-binding shape the
pioneering activity of Foxa2. *Nucleic Acids Res.* **47**, 9069-9086 (2019).

38.   Ahlgren, U., Jonsson, J., Jonsson, L., Simu, K. & Edlund, H. Beta-cell-specific inactivation

of the mouse Ipf1/Pdx1 gene results in loss of the beta-cell phenotype and maturity onset diabetes. *Genes Dev.* **12**, 1763-1768 (1998).

39. Sussel, L. et al. Mice lacking the homeodomain transcription factor Nkx2.2 have diabetes due to arrested differentiation of pancreatic beta cells. *Development* **125**, 2213-2221 (1998).

40. Collombat, P. et al. Opposing actions of Arx and Pax4 in endocrine pancreas development. *Genes Dev.* **17**, 2591-2603 (2003).

41. Dhawan, S., Georgia, S., Tschen, S. I., Fan, G. & Bhushan, A. Pancreatic beta cell identity is maintained by DNA methylation-mediated repression of Arx. *Dev. Cell* **20**, 419-429 (2011).

42. Liu, J. et al. Neurog3-independent methylation is the earliest detectable mark distinguishing pancreatic progenitor identity. *Dev. Cell* **48**, 49-63 (2019).

43. Papizan, J. B. et al. Nkx2.2 repressor complex regulates islet beta-cell specification and prevents beta-to-alpha-cell reprogramming. *Genes Dev.* **25**, 2291-2305 (2011).

44. Wu, X., Li, G. & Xie, R. Decoding the role of TET family dioxygenases in lineage specification. *Epigenetics Chromatin* **11**, 58 (2018).

45. Serandour, A. A. et al. Dynamic hydroxymethylation of deoxyribonucleic acid marks differentiation-associated enhancers. *Nucleic Acids Res* **40**, 8255-8265 (2012).

46. Yang, Y. A. et al. FOXA1 potentiates lineage-specific enhancer activation through modulating TET1 expression and function. *Nucleic Acids Res.* **44**, 8153-8164 (2016).

47. Lio, C. W. et al. Tet2 and Tet3 cooperate with B-lineage transcription factors to regulate DNA modification and chromatin accessibility. *Elife* **5**, e18290 (2016).

48. Rezania, A. et al. Reversal of diabetes with insulin-producing cells derived in vitro from human pluripotent stem cells. *Nat. Biotechnol.* **32**, 1121-1133 (2014).

49. Zmuda, E. J., Powell, C. A. & Hai, T. A method for murine islet isolation and subcapsular kidney transplantation. *J. Vis. Exp.* **50**, 2096 (2011).

50. Cong, L. et al. Multiplex genome engineering using CRISPR/Cas systems. *Science* **339**, 819-823 (2013).

51. Tu, S. et al. Takusan: a large gene family that regulates synaptic activity. *Neuron* **55**, 69-85 (2007).

52. Dobin, A. et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

53. Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**, 166-169 (2015).

54. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

55. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics* **32**, 2847-2849 (2016).

56. Yu, G., Wang, L. G., Han, Y. & He, Q. Y. ClusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* **16**, 284-287 (2012).

57. Buenrostro, J. D., Wu, B., Chang, H. Y. & Greenleaf, W. J. ATAC-seq: a method for assaying chromatin accessibility genome-wide. *Curr. Protoc. Mol. Biol.* **109**, 21.29.21-21.29.29 (2015).

58. Heinz, S. et al. Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576-589 (2010).

59. Cowley, M. et al. Effects of cadmium exposure on DNA methylation at imprinting control
regions and genome-wide in mothers and newborn children. *Environ. Health Perspect.* **126**,
037003 (2018).

60. Wang, L., Wang, S. & Li, W. RSeQC: quality control of RNA-seq experiments. *Bioinformatics*
28, 2184-2185 (2012).

61. Ramirez, F., Dundar, F., Diehl, S., Gruning, B. A. & Manke, T. DeepTools: a flexible platform
for exploring deep-sequencing data. *Nucleic Acids Res.* **42**, W187-191 (2014).

62. Lin, X. et al. BSeQC: quality control of bisulfite sequencing experiments. *Bioinformatics* **29**,
3227-3229 (2013).

63. Feng, J., Liu, T., Qin, B., Zhang, Y. & Liu, X. S. Identifying ChIP-seq enrichment using
MACS. *Nat. Protoc.* **7**, 1728-1740 (2012).

Fig. 1 TET hydroxylases are required for pancreatic endocrine lineage specification.

a Stepwise differentiation of hESCs to pancreatic endocrine cells. Wild-type (WT) and *TET1/TET2/TET3* triple knock-out (TKO) hESC lines were differentiated and analyzed by RNA-seq, WGBS, CMS-IP-seq, ATAC-seq, and ChIP-seq at the indicated stages (ES: embryonic stem cell; DE: definitive endoderm; GT: primitive gut tube; PP: pancreatic progenitor; PE: pancreatic endocrine). **b** Representative flow cytometry plots for PDX1 and NKX6.1 in WT or TKO cells at the PP and PE stages. Quantification of the percentage of PDX1$^+$ NKX6.1$^+$ cells is shown in the right panel. n = 3 independent differentiation. **c** Immunostaining of PDX1/NKX6.1 at the PP stage and insulin (INS)/glucagon (GCG) at the PE stage are shown in WT and TKO cells. Scale bar = 50 μm. **d** Genes with significant changes in expression in TKO cells relative to WT cells at the PP stage. Differentially expressed genes (DEGs) are classified into down- and upregulated groups. Each row corresponds to one individual gene and each column to a different biological repeat. The color scale from white to blue represents Z-score normalized gene expression levels from low to high (FDR < 0.05). **e** Number of regions with significant changes in chromatin accessibility upon TET depletion at proximal (≤ 1 kb from TSS) and distal (> 1 kb from TSS) regions ($p < 0.05$). **f** Dot plots depicting the ratios (WT_PP over TKO_PP) of ATAC signals (y-axis) at proximal (left) or distal (right) regions of DEGs that were downregulated (blue) or upregulated (red) in TKO_PP cells compared with WT_PP cells. Values indicate the fraction of increased or decreased chromatin accessibility regions in down- and upregulated genes, respectively. **g** Transcription factor (TF) motif enrichment analysis of genomic regions showed significantly decreased ATAC-seq signals in TKO_PP cells. The top 10 significant motifs are shown after removing redundant motifs.

Fig. 2 Loss of TET disrupts epigenetic dynamics at pancreas-specific hypo-DHMRs associated

with distinct pioneer TFs motifs.

a Scatterplot presenting 5hmC differences between pancreatic progenitors (WT_PP) and hESCs

(WT_ES) at hypo-DHMRs identified in TKO_PP cells. Red indicates increased 5hmC (fold change >

1.5), and blue shows decreased 5hmC in pancreatic progenitors relative to hESCs. **b** Clustering of

5hmC signals in WT cells at four differentiation stages (ES, DE, GT, and PP). Each row represents

one hypo-DHMR showing increased 5hmC in pancreatic progenitors compared with hESCs. Red

represents high 5hmC; blue indicates low 5hmC. **c** Motif enrichment analysis among the four clusters

defined in **b**. The top five known motifs are shown. **d** Heatmap and density plots showing dynamic

changes of FOXA2 in the DE, GT, and PP lineage intermediates across the four clusters defined in **b**.

The color scale from white to blue represents normalized FOXA2 signal from low to high. **e**

Enrichment profile of chromatin accessibility (ATAC), H3K4me1, and H3K27ac at regions of

FOXA2 binding alone (left column), FOXA2 binding together with 'differentiation-specific hypo-

DHMRs' (middle column), and 'differentiation-specific hypo-DHMRs' alone (right column). The

WT signal is marked in blue and the TKO signal in red.

Fig. 3 Pancreas-specific hyper-DMRs show reduced chromatin activity during pancreatic differentiation.

a Volcano plot of WGBS data illustrating differentially methylated CpGs (DMCs) identified in TKO_PP cells compared with WT_PP cells. Red and blue represent increased and decreased 5mC in TKO_PP cells, respectively (credible methylation difference > 0.2). **b** Heatmap illustrating methylation difference between TKO_PP and WT_PP cells at centers of annotated genomic features (± 5 kb) for chromatin accessibility (ATAC), hydroxylation (5hmC), TF binding (FOXA2, GATA4, GATA6, and PDX1), bivalent promoters, poised enhancers, and active enhancers. Average 5mC signals of every 100-bp bin were calculated. **c** Classification of TKO hyper-DMRs based on 5mC levels in hESCs (green), WT_PP cells (blue), and TKO_PP cells (red). **d** Average plots of FOXA2 (left column), GATA4 (middle column), and GATA6 (right column) signal at pancreas-specific hyper-DMRs or non-pancreatic hyper-DMRs in pancreatic progenitors. **e** Average plots of ATAC (left column), H3K27ac (middle column), and H3K4me1 (right column) at pancreas-specific hyper-DMRs or non-pancreatic hyper-DMRs in WT_PP (blue) and TKO_PP (red) cells. **f** Average plots of 5mC at proximal (≤ 1 kb from TSS) and distal (> 1 kb from TSS) decreased accessible regions in WT_PP (blue) and TKO_PP (red) cells. **g** Genome-browser view of the *PDX1/PDX1-AS1* locus. Four type 2 diabetes-associated islet hyper-DMRs [31] overlapping with TKO hyper-DMRs are highlighted in green. A specific TKO hyper-DMR showing decreased ATAC, H3K4me1, and H3K27ac signals is highlighted in pink.

Fig. 4 TET1 is required in pancreatic β-cell specification.

a Expression of *PAX4* in WT, TET1KO, TET2KO, TET3KO, TET1/2DKO, TET1/3DKO, TET2/3DKO, and TKO cells at the PP stage by RT-qPCR. RT-qPCR validation was performed with three independent batches of samples. **b** Representative plots of flow cytometry of human C-peptide

(CPEP) and glucagon (GCG) in WT, TET1KO, and TET2/3DKO cells at the PE stage.

Quantifications of the percentage of CPEP$^+$ or GCG$^+$ cells are shown in the right panel. n = 3

independent differentiation. **c** Immunostaining of NKX6.1, PDX1, INS, and GCG in WT, TKO-

CTRL, and TKO-TET1 cells at the PE stage of pancreatic differentiation. Scale bar = 50 μm. **d**

Expression of *PDX1*, *NKX6.1*, and *PAX4* in WT, TKO-CTRL, and TKO-TET1 cells at the PP stage

by RT-qPCR. RT-qPCR validation was performed with three independent batches of samples. **e**

Heatmap showing the hierarchical clustering of DEGs among WT_PP, TKO_PP, and TET1KO_PP

cells. Each row represents one DEG, and each column represents one biological replicate. The color

scale from white to blue represents normalized gene expression levels from low to high (FDR < 0.05).

f WT, TKO, and TET1KO cells were differentiated to the PE stage and transplanted under the kidney

capsule of nondiabetic female SCID-beige mice. Eighteen weeks post-implantation, human C-peptide

levels were measured after an overnight fast and 30 min following an i.p. glucose injection. C-peptide

levels from individual mice are shown in box and whisker plots.

Fig. 5 TET1 binding sites are distinct in pancreatic progenitors.

a Methylation levels at TET1 binding sites in WT_PP cells. The x-axis represents the average

methylation ratio of CpG sites located within each peak (at 5% methylation ratio intervals). The green

bar represents the percentage of TET1 peaks within the corresponding methylation ratio interval (left

y-axis). The red curve indicates TET1-enriched signals within each corresponding methylation ratio

interval (right y-axis). The numerical labels at the top represent the fraction of the four class peaks

with different average methylation ratios: UMR, unmethylated regions (0-0.1); LMR, low methylated

regions (0.1-0.5); MMR, medium methylated regions (0.5-0.9); and HMR, high methylated regions

(0.9-1). **b** Heatmaps demonstrating enrichment patterns of ATAC, 5hmC, and histone modification

(H3K4me1, H3K27me3, H3K4me1, and H3K27ac) relative to the center of TET1 peaks ($\pm$ 5 kb) at

proximal ($\leq$ 1 kb from TSS) and distal ($>$ 1 kb from TSS) regions. **c** Venn diagram illustrating distinct

and overlapping TET1 binding sites between hESCs and pancreatic progenitors ($p < 0.05$). **d** TF motif

enrichment analysis of TET1 targets identified specifically in pancreatic progenitors (top, PP-specific)

or hESCs (bottom, ES-specific). The top three motifs are shown. **e** Number of TET1 binding sites

identified specifically in pancreatic progenitors (PP-specific) or in both hESCs and pancreatic

progenitors (common) at proximal and distal regions ($p < 0.05$). **f** Average plots (top) and heatmaps

(bottom) of TET1, H3K4me3, H3K27me3, H3K4me1, H3K27ac, and ATAC at 'common' or 'PP-

specific' TET1 peaks ($\pm$ 5 kb) at proximal or distal regions. Heatmaps are ranked by decreased TET1-

binding signal of 'common' and 'PP-specific' regions, respectively. The color scale from white to

blue represents the normalized signal from low to high. **g** Distribution profiles of average methylation

(5mC), ATAC, H3K4me1, and H3K27ac located at FOXA2-bound sites with (green) or without

(orange) co-binding of TET1.

Fig. 6 TET1 is required for the PAX4 enhancer to achieve a hypomethylated state.

a Genome-browser view of the *PAX4* locus with increased methylation and decreased chromatin

associability upon TET depletion at a TET1/FOXA2 co-bound region featuring enhancer signatures

H3K4me1 and H3K27ac. **b** Locus-specific increase in 5mC at the *PAX4* enhancer in TKO or

TET1KO samples compared with TET2/3DKO samples. Percentages of unmethylated cytosine and

5mC at CCGG sites are shown. n = 3 independent differentiation.