Controlling for off-target genetic effects using polygenic scores improves the power of genome-wide association studies

Declan Bennett, Derek Morris and Cathal Seoighe[*]

[*]Correspondence:
cathal.seoighe@nuigalway.ie
National University of Ireland,
Galway, University Road, Galway,
Ireland
Full list of author information is
available at the end of the article

Abstract

Ongoing increases in the size of human genotype and phenotype collections offer the promise of improved understanding of the genetics of complex diseases. In addition to the biological insights that can be gained from the nature of the variants that contribute to the genetic component of complex trait variability, these data have brought forward the prospect of predicting complex traits and the risk of complex genetic diseases from genotype data. Optimal realization of these objectives requires ongoing methodological developments, designed to identify true trait-associated variants and accurately predict phenotype from genotype. In addition to optimal accuracy, these methods must also be computationally efficient, in order to remain tractable in the context of high variant densities and very large sample sizes. Here we show that the power of linear mixed models that are in widespread use for GWAS can be increased significantly by modeling off target genetic effects using polygenic scores derived from SNPs that are not on the same chromosome as the target SNP. Applying this method to simulated and real data we found that it can result in a substantial increase in the number of variants passing genome-wide significance thresholds. This increase in power to detect trait-associated variants also translates into an increase in the accuracy with which the resulting polygenic score predicts the phenotype from genotype data.

Keywords: GWAS; PGS; Off-target effects

Bachground

Linear mixed effects models (LMMs) are routinely applied to detect associations

between SNPs and phenotypes in genome-wide association studies (GWAS). The

contribution of a given SNP to the phenotype is modelled as a fixed effect, while

the contribution of all other SNPs is modelled as a random effect, with the co-

variance structure of this random effect corresponding to the genetic relationships

between the individuals in the study. In order to avoid including the component

of the genetic relatedness contributed by variants in linkage disequilibrium (LD)

with the tested variant, the genetic relationship matrix (GRM) is calculated using

an LD-pruned set of variants [1, 2]. Originally these methods assumed that a large

number of SNPs may each make small contributions to the phenotypic variation,

such that the phenotypic contribution of the genetic background could be mod-

elled as a multivariate normal (with covariance matrix proportional to the GRM)

[2, 3, 4, 5]. Newer implementations of these models have allowed for the possibility

of a combination of SNPs with effect sizes close to zero as well as some SNPs with

large effect sizes. This can be done by modelling the effect sizes of background SNPs

as a mixture of normal distributions, with a proportion, $p << 1$, of SNPs with large

effect sizes and the remainder with effect sizes close to zero. This results in a 'spike

and slab' mixture model for the SNP effect sizes [6] (the spike corresponds to a nor-

mal component with small variance and the slab corresponding to a second normal

component with large variance). Implementations of these normal mixture mod-

els have been developed that can be applied to biobank-scale data [7]. In addition

to controlling for the direct genetic effects of other SNPs on the phenotype LMM

methods have also been shown to account well for subtle population structure, and

cryptic relatedness between individuals in the sample population [1, 2, 3, 4, 8].

Various options have been explored for which SNPs to include in the calculation of

the GRM [9]. Including SNPs in LD with the target SNP results in loss of power, as

the effect of the target SNP is partially accounted for by the random effect through

the GRM. This has been referred to as proximal contamination [10]. On the other

hand, including all (or most) SNPs that are not in LD with the target SNP, e.g.

using a Leave One Chromosome Out (LOCO) approach, can result in dilution of

the extent to which the relevant part of the genetic background is captured by

the GRM. In the latter case, SNPs that are not relevant, in that they do not

capture direct genetic effects or tag relevant population structure effects, effectively

add noise to the GRM [10]. Alternatively, the GRM can be built from only the

SNPs that are found using a linear model to be associated with the phenotype.

Although this results in an increase in statistical power [5, 9, 11], it does not fully control for population structure and is not recommended if population structure is of substantial concern [6, 9]. Methods have been developed that incorporate principal components into the GRM calculation built from significant SNPs; however, most of these methods are not suited to large biobank-scale data, without access to cloud computing or large compute farms [12, 13, 14]. Off-target SNPs can also be included in the statistical model as fixed effects and this is the recommended approach when there are SNPs with large effect sizes [9]. A model fitting approach to determine the SNPs to include as fixed effects has been developed, and this also results in increased power in GWAS [10].

As the genomic architecture of complex diseases is uncovered with the help of large biobanks, the application of GWAS results in a clinical setting is increasing in importance [15], including through the use of polygenic scores (PGS). PGS are constructed from weighted sums of allele dosages, with the weights corresponding to the effects size of the variants. Risk variants (variants associated with the phenotype) are typically inferred from the largest available GWAS, generally a meta-analysis. The clinical potential of PGS has already been shown in complex diseases such as coronary artery disease (CAD), diabetes and cancer [16, 17, 18]. In CAD, the identification of individuals with similar risk to those with rare high-risk monogenic variants has been reported [16]. Similarly, in breast cancer, pathogenic variants in BRCA1/2 account for 25% of familial risk of the disease with genome wide variants accounting for a further 18% of the risk [19, 20]. European bias is a significant concern in clinical application of PGS, resulting from over-represented of individuals of European ancestry in GWAS and poor generalization of PGS when applied to individuals with genetic ancestry distinct to the population from which the score was derived [21, 22].

Here, we set out to investigate whether including an additive PGS, estimated using a LOCO scheme, as a fixed effect in a linear mixed model has the potential to improve the power of GWAS. Our hypothesis is that the contribution of background genetic variation may not be adequately captured by a random effect, with covariance proportional to the GRM. In particular, stochastic variation resulting from SNPs with relatively large contributions to phenotypic variation may be better captured by the PGS. We tested this hypothesis in two ways. Firstly, using

simulated data we tested for an improvement in power on the task of recovering

known causal variants as a function of study size, number of causal variants and

trait heritability. In addition, we applied the method to standing height data from

the UK Biobank and determined the number and characteristics of additional vari-

ants that were detected. For an objective assessment of performance on real data,

where the true associations are unknown, we divided the data into test and training

sets and predicted the phenotype in the test set. The improved performance on the

critical task of complex phenotype prediction illustrates the utility of the PGS as a

means of accounting for background genetic variance.

Results

We simulated data to evaluate the impact of including the PGS as a fixed effect

in GWAS. The simulations consisted initially of a normally-distributed continuous

trait in 100,000 individuals. The trait had a narrow-sense heritability (h^2) of 0.5

and there were 1,000 causal SNPs with normally-distributed effects on the trait

(see Methods for details). We incorporated the LOCO PGS as a fixed effect in a

linear mixed model using GCTA fastGWA [23] (a method we refer to as PGS-LMM)

and found a substantial improvement in power to detect the known causal SNPs

(Fig. 1). In 100 simulations we found that the PGS-LMM recovered, on average,

76 additional causal variants below the conventional P-value threshold of $5x10^{-8}$

compared to fastGWA (Fig. 1), corresponding to a relative increase in power of 17%.

The contribution to phenotype variance of off-target SNPs can also be modelled as

a random effect in a linear mixed model. This approach is applied by BOLT-LMM,

which uses a normal mixture random effect, with a component corresponding to

SNPs with large effects; however, PGS-LMM achieved higher power than BOLT-

LMM, suggesting that first calculating a PGS from an initial GWAS and including

this as a fixed effect in a mixed model may provide a better means to control for

off-target genetic effects (Fig. 1).

To investigate whether the increased number of causal variants recovered under

PGS-LMM reflected a reduction in P-values across the board or also an improvement

in the ordering of the variants, when the variants are ordered by the evidence of

an association with the phenotype, we calculated receiver operator characteristic

(ROC) curves for the fastGWA and PGS-LMM method (i.e. fastGWA with the

LOCO PGS included as a fixed effect covariate). Over 100 simulations we found

that the area under the ROC curve (AUC) was always higher for PGS-LMM than

for fastGWA. The difference in sensitivity as a function of specificity (Fig. 2) showed

that the ordering of SNPs is significantly improved under PGS-LMM. The PGS-

LMM method provided an improvement of up to 0.051 in the mean sensitivity

(corresponding to a relative improvement of 8.2% in sensitivity compared to the

fastGWA method at a specificity of 0.9992).

Dependence on trait heritability and number of causal variants

We simulated data over a range of values of h^2 and of the number of causal SNPs.

As we would expect, for very low values of h^2 PGS-LMM offers very little improve-

ment over fastGWA (although the proportion of causal variants detected below a

p-value threshold of $5x10^{-8}$ was always greater for PGS-LMM). From a h^2 value of

around 0.4 the proportion of the causal variants tended to be significantly greater

for PGS-LMM than for fastGWA (Fig. 3a). Similarly, when there was a small num-

ber of causal variants (with a fixed value of $h^2 = 0.5$), including the polygenic

score as a covariate provided little advantage; however, again, the number of causal

variants recovered when the PGS was included as a covariate was always greater

than or equal to the number recovered when the PGS covariate was omitted. For

simulations with at least 80 causal variants the proportion of causal variants cor-

rectly identified with PGS-LMM always exceeded that obtained with the fastGWA

and this difference in proportions became statistically significant as the number of

causal variants approached 1,000 (Fig. 3b).

Results of simulations with variable numbers of GWAS participants

Including the PGS as fixed effect resulted in a higher proportion of causal vari-

ants recovered across the full range of sample sizes investigated (Fig. 4). Even at

N=10,000 there was a significant difference between the proportion of causal vari-

ants recovered with PGS-LMM over fastGWA without PGS. This is somewhat

surprising given that it is assumed that large sample sizes are required for accurate

phenotype prediction from PGS [24]. As the sample size increased, the number of

variants recovered increased sharply for both models (with and without PGS), with

the rate of increase in power gradually coming down as sample numbers approach

100,000.

Application to UK Biobank height data

We assessed performance of PGS-LMM on real data using standing height of in-

dividuals of British ancestry (429,359 individuals) from the UK Biobank. The dis-

tribution of P-values obtained from the PGS-LMM was lower than that obtained

using fastGWA (See Supplementary Figure 1, Additional File 1). At a genome-wide

significance level of $5x10^{-8}$ there were 127,306 and 152,626 variants that achieved

significance using fastGWA and PGS-LMM, respectively. A total of 31,332 variants

were identified by PGS-LMM that did not pass the significance threshold using fast-

GWA, termed from here as PGS-LMM only variants (See Supplementary Figure 2,

Additional File 1). These variants were significantly enriched for low ($< 1\%$) minor

allele frequency (OR=2.44; CI=2.2-2.7; P $= 6x10^{-65}$, using Fisher's Exact test;

Fig. S3 Table S1). We performed a gene-property analysis for tissue specificity of

the PGS-LMM-only variants across 53 GTEx (v8) tissue types. Muscle and skeletal

tissue had the strongest, albeit modest, association (P $= 0.02$; Fig. S4, Table S2,

Additional File 1).

Application to phenotype prediction

One way to determine objectively whether PGS-LMM outperforms fastGWA on real

data is to apply both methods on the key task of phenotype prediction. We par-

titioned the UKB height data into an 80%/20% training and testing datasets and

calculated the PGS scores using a partitioning and thresholding (P+T) method

as well as a more powerful Bayesian method to estimate the phenotype in the

test dataset (see Methods for details). In both cases PGS-LMM consistently out-

performed fastGWA across all inclusion thresholds (Posterior inclusion probability

(PIP) and P-value). The maximum R^2 using P+T was achieved at a P value thresh-

old of 0.05 for both the fastGWA ($R^2 = 0.129$) and PGS-LMM ($R^2 = 0.138$), cor-

responding to a relative improvement in the proportion of the phenotype explained

of 7% (Fig. 5a). The Bayesian model provided substantially better prediction of the

phenotype than the P+T method when variants below 0.4 PIP were included in

the model and, again, this performance was further enhanced when the SNP effects

were estimated using PGS-LMM rather than fastGWA (Fig. 5b). The maximum

R^2 for PGS-LMM was 0.194, compared to 0.184 with fastGWA (corresponding to

a relative increase of 5.4%). Although smaller, the improvement in performance

on the prediction task remained consistent when we repeated this analysis with a 20%/80% training-test split (Fig. S5, Additional File 1). There are three possible reasons for the improvement in prediction obtained using PGS-LMM. It could result from a larger number of SNPs passing a given threshold being used in prediction, from improvement in the ordering of SNPs, so that a better set of SNPs are being used or, lastly, from better estimates of the SNP effect sizes. To distinguish between these possibilities we repeated the prediction using the same number of SNPs for PGS-LMM and fastGWA and then for the same set of SNPs for both (see Methods for details). The performance gain was maintained when we used the same number of SNPs, but was much smaller when we used exactly the same set of SNPs (Table S3, Additional File 1).

Discussion

Omitting covariates that are associated with a response and independent of an effect of interest can result in a reduction in the efficiency of the estimation of the effect of interest [25]. Complex traits are associated with the genotype of many loci across the genome, but the effects of the off-target SNPs are frequently not explicitly modelled in GWAS and, instead tests of association are performed for one target SNP at a time. We evaluated a simple two-stage approach to accounting for off-target genetic effects that consists of performing an initial GWAS and using the summary statistics to calculate a polygenic score and then including the polygenic score, derived from SNPs not on the same chromosome as the target SNP, as a fixed effect in a second round of association testing. We found that this led to a substantial improvement in GWAS power using simulated data. A popular approach to account for the effects of off-target SNPs is to make use of a mixture distribution for the random effect in a linear mixed model, with a component with large variance corresponding to off-target SNPs with large effect sizes. This approach is implemented in BOLT-LMM [6]. However, PGS-LMM had higher power in simulations than BOLT-LMM, suggesting that incorporating the PGS as a fixed effect in a linear model offers a better alternative for accounting for off-target genetic effects. It is also possible to include the LOCO PGS as a fixed effect with BOLT-LMM. In this case the effects of off-target SNPs are modelled both as a fixed and as a random effect. We found that this offered a small further improvement over fastGWA with the LOCO PGS

(Fig. S6). However, this improvement comes at a substantial cost, as fastGWA is

much more computationally efficient than BOLT-LMM [23].

The increase in power with PGS-LMM was largest for simulated phenotypes with

high heritability and a large number of causal variants (Fig. 3b). In these cases

the many off-target SNPs collectively explain a substantial proportion of the phe-

notypic variance and summarizing the contribution of these off-target SNPs to the

phenotype via the LOCO PGS is likely to result in a better estimate of the effect

of the target SNP and its standard error. The boost in performance derived from

including the LOCO PGS as a fixed effect was evident even for relatively small

study sizes. Across all the simulation parameters we investigated, the performance

of the PGS-LMM was never worse than fastGWA without the LOCO PGS.

We also applied the method to real data (standing height in individuals of British

ancestry in the UK Biobank). Height was chosen as there have been over 3,000 near

independent variants associated with height, explaining 0.483 of the phenotypic

variance. Heritability is often estimated to be as high as 80%. Although GWAS

can only give us insight into h^2, the additive component of heritability, estimates

of h^2 for height range from 0.3 for high quality common variants up to 0.88 in

monozygotic twins for all SNPs [26, 27, 28]. Consistent with the simulation results,

we found nearly 20% more associated variants using the PGS-LMM, including a

large number of variants (31,332) that were significant using PGS-LMM, but not

using fastGWA (a much smaller set of 6,012 SNPs were identified as significant by

fastGWA but not PGS-LMM; Fig. S2). The newly identified variants are enriched

for rare variants while common variants were depleted. The PGS-LMM identified

variants were associated with expression in skeletal and muscle tissue, consistent

with PGS-LMM identifying additional true causal variants over fastGWA.

The h^2 estimate in the simulations corresponds to the combined additive contribu-

tion to the variance of the causal SNPs; however, when we apply GWAS methods to

the simulated data we do not recover all of the associated SNPs and, consequently,

the estimated SNP heritabilities tends to be lower than the simulated values. For

example, the simulations that were the basis of Figure 1 used $h^2 = 0.5$, but the mean

estimated h^2 for these simulations was 0.41. This has implications for the interpre-

tation of Figure 3a, which shows the relationship between simulated h^2 and power,

rather than the estimated h^2. Within UK Biobank, standing height has the highest

estimated SNP h^2, at 0.52 [23]. PGS-LMM offers improvement over fastGWA at higher values of h^2, so we can expect it to yield more independent association signals for other traits such as weight, heel bone mineral density, basal metabolic rate and platelet count that all have estimated $h^2 > 0.2$ in UK Biobank [23, 29]. These are traits with high heritabilities that can be easily measured in large population-based cohorts, which has enabled GWAS to already detect many association signals. Some diseases have equally high heritability, e.g., type 1 diabetes, schizophrenia and Alzheimer's disease, but do not yet have sample sizes on the same scale as height. As this changes and statistical power of GWAS of these diseases increases, inclusion of the LOCO PGS as a fixed effect in GWAS may prove useful for identifying additional risk loci for them over and above conventional GWAS analysis.

The use of polygenic scores to contribute to phenotype prediction from genotype is an increasingly important application of the results of GWAS [30]. Recent work has shown that PGS can identify sizeable proportions of a population with substantially greater risk for CAD, type II diabetes (T2D), arterial fibrillation (AF), infammatory bowel disease (IBD) and breast cancer [16]. A separate study using the FinnGen dataset compared the overall lifetime risk in coronary heart disease (CHD), T2D, AF, breast cancer and prostate cancer between individuals having an average compared to a high PGS score. A high PGS score was associated with a 21% to 38% increase in overall lifetime risk with age of onset 4 to 9 years earlier than in the average group [31]. Prophylactic interventions can reduce both the physical and economic stress on health systems by entering individuals on measures to prevent and manage disease earlier. PGS when used to target individuals with high efficacy of treatment can also reduce the numbers of individuals to treat [32]. Although having a high PGS for a disease is not akin to a diagnosis, it can guide individuals to limit their exposure to other risk factors. Performing GWAS on a subset of samples and predicting on the remainder, we observed a consistent improvement in accuracy of the prediction of standing height in UK Biobank with PGS-LMM compared to fastGWA. Although there are many caveats surrounding the prediction of complex phenotypes, particularly clinical phenotypes [22], accurate prediction of disease susceptibility can have profound implications for disease management and prevention. Our results suggest that incorporating PGS into the

263 GWAS before recalculating PGS or otherwise predicting the phenotype or disease
264 risk leads to improved performance on this crucial task.

Conclusion

The tasks of detecting trait-associated variants and predicting the trait in a new
sample from the summary statistics of these variants are closely intertwined. Im-
proved performance on the trait-association task can result in more associated vari-
ants and better estimates of their effect sizes, resulting in improvement on the pre-
diction task. On the other hand, improved methods for phenotype prediction can
help to control for off-target effects in methods that identify the trait-associated
variants and their effects. The method that we have explored here consists of in-
corporating a LOCO PGS as a fixed-effect covariate to control for these off-target
effects; however, any method for phenotype prediction could play this role, once
its application is restricted to variants that are not linked to the target SNP. We
show here that incorporating the PGS as a fixed-effect covariate results in increased
power to detect trait-associated variants in GWAS. The resulting trait-associated
variants and effect size estimates lead to an improvement in the PGS, as illustrated
by improved performance in the task of predicting the phenotype in a test dataset.

Methods

Genotype data

The simulations and application to real data were based on autosomal genotype
data from the UK Biobank. To limit the effects of population stratification only
individuals reporting white British ancestry (data field 21000; code 1001) were
included in all analyses. The genotype data for the simulation analysis was based
on directly genotyped variants with minor allele frequency (MAF) greater than
0.005%. Variants with genotype missingness greater than 2% or that failed a test
for Hardy-Weinberg equilibrium (HWE) at $\alpha = 0.0001$ were excluded. To compute
the genetic relationship matrix (GRM), a set of variants was extracted with MAF
greater than 0.25%, a genotype missingness rate of less than 2% and passing a HWE
test at $\alpha = 0.0005$. The GRM set was further reduced by elimination of variants with
an R^2 greater than 0.1 in a 200 variant sliding window of size 500. All genotype QC is
implemented in plink2 [33]. Sparsity of the GRM was increased by removing sample
pairs with an estimated relatedness of 0.05. To account for population structure in

the association studies, principal component analysis (PCA) was performed on the
set of GRM variants using plink2.

Simulations

Based on the above genotype data, we simulated a continuous phenotype using the
GCTA software suite [34]. The initial simulation consisted of 100,000 individuals,
1,000 causal variants and $h^2 = 0.5$. This simulation was repeated 100 times. Power
was calculated as the proportion of the causal variants recovered. To calculate false
positive rates we first removed all SNPs within 1 Mb of the causal SNPs. Further
simulations were carried out to investigate the effects of varying the number of causal
SNPs, h^2 and the sample size on method performance. In each case all parameters
other than the one being varied were the same as the initial simulation, and one
simulation was performed per set of parameter values. Error bars on the proportion
of causal variants recovered were calculated based on twice the standard error of a
proportion. To assess the sensitivity and specificity of the simulations, the pROC R
package was used to generate receiver operating characteristic (ROC) curves [35].

Association tests

Association testing was performed using fastGWA and using BOLT-LMM with
10 PCs and assessment centre as fixed covariates. For the PGS-LMM method we
first performed GWAS (using fastGWA or BOLT-LMM) and calculated PGS scores
on a Leave One Chromosome Out (LOCO) basis. This resulted in 22 sets of PGS
values (one for each autosomal chromosome, calculated from the summary statistics
of variants on all other autosomal chromosomes). The PGS were calculated using
PRSice2 (version 2.2.12 (2020-02-20))[36]. To decrease computation time, resources
and reduce the likelihood of overfitting a p-value threshold of 5e-05 was chosen
a priori for the LOCO PGS calculation. Association testing was then performed
using fastGWA in a chromosome-wise manner with the corresponding LOCO PGS
included as a fixed effect.

Application to standing height from the UK biobank

Autosomal genotype data was cleaned using MAF of 0.1%, SNP-missingness of 2%
and sample-missingness cut-off of 10% and imputation quality thresholds of 0.8.
Variants that failed HWE testing at $\alpha = 5x10^{-7}$ were excluded from subsequent

analysis. The total number of variants remaining after filtering was 9,156,362. The

GRM was calculated using a set of LD-pruned variants with a MAF of at least 0.01,

window size of 1000, R^2 of 0.9 and window shift of 100 variants. This left a total of

1,326,927 to be used in the calculation of the sparse GRM using the same methods

as described for the simulation analysis. Batch number, sex, age, assessment centre

and 10 PCs were included as covariates in all association tests.

Gene-property analysis

To test whether the PGS-LMM only variants had an association with tissue spe-

cific expression data from GTEx, gene-property analysis was performed using the

SNP2GENE functionality of the FUMA web platform [37]. Briefly, FUMA using

MAGMA [38], maps variants to genes assigning a p-value based gene score. The

Z-transformed gene scores are then modelled along with the log2 RPKM average

expression in the tissue of interest, log2 RPKM of average expression across all tis-

sues and technical confounders. To test for a positive relationship a one-sided test

is then performed.

Prediction

To test the performance of PGS-LMM on the task of predicting standing height, the

UK Biobank data was partitioned into 80/20 and an independent 20/80 training and

test datasets. We used two PGS strategies, pruning and thresholding (P+T) and a

Bayesian method (SBayesR) [36, 39]. As the Bayesian method requires LD matrices

we used a pre-computed set of matrices built on 2.8 million common pruned UK

Biobank variants. The precomputed LD matrices for 2.8 million high-confidence UK

biobank variants were downloaded from the Zenodo public repository (10.5281/zen-

odo.3375373). Both PRSice and SBayesR predictions were performed using this 2.8

million UK biobank set of genotypes.

For the P+T method, the same PGS-LMM methodology was performed as in

the simulation study and PGS scores were calculated in the test set using p-value

thresholds (0.001, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 1). The Bayesian method replaces the

P+T method in the PGS-LOCO step with SBayesR. To limit the effect of over-

fitting, we excluded variants with a posterior inclusion probability (PIP) score of

less than 0.2 in the PGS-LOCO calculation. As SBayesR uses Bayesian inference

to estimate the true effect sizes, Plink2 was used to calculate the PGS-LOCO score

from the genotype and SBayesR effect size data before association testing. SBayesR

was then applied once more on the PGS-LMM summary statistics over a range of

PIP scores. The model fit was assessed for each method by fitting a linear model to

the values of the phenotype in the test set as a function of their predicted values.

We repeated the phenotype prediction using the same number of SNPs (74,000)

with the PGS-LMM and fastGWA methods. We chose 74,000 as it fell in the range of

the number of top SNPs that maximise the R^2 value for PGS-LMM and fastGWA.

We also performed the prediction using the 37,514 SNPs that were in common

among the top 74,000 SNPs from fastGWA and PGS-LMM.

Availability of data and materials

The PGS-LMM implementation described in this work is available under MIT license from the github repository,

https://github.com/declan93/PGS-LMM/.

Competing interests

The authors declare that they have no competing interests.

Author's contributions

CS conceived and supervised the project and performed analyses. DB implemented the pipeline and performed

analyses. DB and CS wrote the manuscript, with input from DM. DM advised on application of the method to

human phenotypes.

Acknowledgements

This research has been conducted using the UK Biobank Resource under Application Number 23739. This

publication has emanated from research conducted with the financial support of Science Foundation Ireland under

Grant number 16/IA/4612.

References

1. Yu, J., Pressoir, G., Briggs, W.H., Bi, I.V., Yamasaki, M., Doebley, J.F., McMullen, M.D., Gaut, B.S., Nielsen,

D.M., Holland, J.B., *et al.*: A unified mixed-model method for association mapping that accounts for multiple

levels of relatedness. Nature genetics **38**(2), 203–208 (2006)

2. Kang, H.M., Zaitlen, N.A., Wade, C.M., Kirby, A., Heckerman, D., Daly, M.J., Eskin, E.: Efficient control of

population structure in model organism association mapping. Genetics **178**(3), 1709–1723 (2008)

3. Kang, H.M., Sul, J.H., Service, S.K., Zaitlen, N.A., Kong, S.-y., Freimer, N.B., Sabatti, C., Eskin, E., *et al.*:

Variance component model to account for sample structure in genome-wide association studies. Nature

genetics **42**(4), 348 (2010)

4. Zhou, X., Stephens, M.: Genome-wide efficient mixed-model analysis for association studies. Nature genetics

44(7), 821–824 (2012)

5. Listgarten, J., Lippert, C., Heckerman, D.: Fast-lmm-select for addressing confounding from spatial structure

and rare variants. Nature genetics **45**(5), 470–471 (2013)

6. Loh, P.-R., Tucker, G., Bulik-Sullivan, B.K., Vilhjalmsson, B.J., Finucane, H.K., Salem, R.M., Chasman, D.I.,

Ridker, P.M., Neale, B.M., Berger, B., *et al.*: Efficient bayesian mixed-model analysis increases association

power in large cohorts. Nature genetics **47**(3), 284 (2015)

7. Loh, P.-R., Kichaev, G., Gazal, S., Schoech, A.P., Price, A.L.: Mixed-model association for biobank-scale

datasets. Nature genetics **50**(7), 906–908 (2018)

8. Price, A.L., Zaitlen, N.A., Reich, D., Patterson, N.: New approaches to population stratification in

genome-wide association studies. Nature Reviews Genetics **11**(7), 459–463 (2010)

9. Yang, J., Zaitlen, N.A., Goddard, M.E., Visscher, P.M., Price, A.L.: Advantages and pitfalls in the application of mixed-model association methods. Nature genetics **46**(2), 100 (2014)

10. Listgarten, J., Lippert, C., Kadie, C.M., Davidson, R.I., Eskin, E., Heckerman, D.: Improved linear mixed models for genome-wide association studies. Nature methods **9**(6), 525 (2012)

11. Lippert, C., Quon, G., Kang, E.Y., Kadie, C.M., Listgarten, J., Heckerman, D.: The benefits of selecting phenotype-specific variants for applications of mixed models in genomics. Scientific reports **3**, 1815 (2013)

12. Tucker, G., Price, A.L., Berger, B.: Improving the power of gwas and avoiding confounding from population stratification with pc-select. Genetics **197**(3), 1045–1049 (2014)

13. Canela-Xandri, O., Rawlik, K., Tenesa, A.: An atlas of genetic associations in uk biobank. Nature genetics **50**(11), 1593–1599 (2018)

14. Kadie, C., Heckerman, D.: Ludicrous speed linear mixed models for genome-wide association studies. bioRxiv doi: 10.1101/154682 (2019)

15. Tam, V., Patel, N., Turcotte, M., Bossé, Y., Paré, G., Meyre, D.: Benefits and limitations of genome-wide association studies. Nature Reviews Genetics **20**(8), 467–484 (2019)

16. Khera, A.V., Chaffin, M., Aragam, K.G., Haas, M.E., Roselli, C., Choi, S.H., Natarajan, P., Lander, E.S., Lubitz, S.A., Ellinor, P.T., *et al.*: Genome-wide polygenic scores for common diseases identify individuals with risk equivalent to monogenic mutations. Nature genetics **50**(9), 1219–1224 (2018)

17. Torkamani, A., Wineinger, N.E., Topol, E.J.: The personal and clinical utility of polygenic risk scores. Nature Reviews Genetics **19**(9), 581 (2018)

18. Yanes, T., Young, M.-A., Meiser, B., James, P.A.: Clinical applications of polygenic breast cancer risk: a critical review and perspectives of an emerging field. Breast Cancer Research **22**(1), 1–10 (2020)

19. Michailidou, K., Lindström, S., Dennis, J., Beesley, J., Hui, S., Kar, S., Lemaçon, A., Soucy, P., Glubb, D., Rostamianfar, A., *et al.*: Association analysis identifies 65 new breast cancer risk loci. Nature **551**(7678), 92 (2017)

20. Bahcall, O.: Common variation and heritability estimates for breast, ovarian and prostate cancers. Nature Genetics **35**, 23–25 (2013)

21. Lambert, S.A., Abraham, G., Inouye, M.: Towards clinical utility of polygenic risk scores. Human molecular genetics **28**(R2), 133–142 (2019)

22. Duncan, L., Shen, H., Gelaye, B., Meijsen, J., Ressler, K., Feldman, M., Peterson, R., Domingue, B.: Analysis of polygenic risk score usage and performance in diverse human populations. Nature communications **10**(1), 1–9 (2019)

23. Jiang, L., Zheng, Z., Qi, T., Kemper, K.E., Wray, N.R., Visscher, P.M., Yang, J.: A resource-efficient tool for mixed model association analysis of large-scale data. Technical report, Nature Publishing Group (2019)

24. Dudbridge, F.: Power and predictive accuracy of polygenic risk scores. PLoS genetics **9**(3) (2013)

25. Neuhaus, J.M.: Estimation efficiency with omitted covariates in generalized linear models. Journal of the American Statistical Association **93**(443), 1124–1129 (1998)

26. Yengo, L., Sidorenko, J., Kemper, K.E., Zheng, Z., Wood, A.R., Weedon, M.N., Frayling, T.M., Hirschhorn, J., Yang, J., Visscher, P.M., *et al.*: Meta-analysis of genome-wide association studies for height and body mass index in 700000 individuals of european ancestry. Human molecular genetics **27**(20), 3641–3649 (2018)

27. Hou, K., Burch, K.S., Majumdar, A., Shi, H., Mancuso, N., Wu, Y., Sankararaman, S., Pasaniuc, B.: Accurate estimation of snp-heritability from biobank-scale data irrespective of genetic architecture. Nature genetics, 1 (2019)

28. Nolte, I.M., Jansweijer, J.A., Riese, H., Asselbergs, F.W., Van Der Harst, P., Spector, T.D., Pinto, Y.M., Snieder, H., Jamshidi, Y.: A comparison of heritability estimates by classical twin modeling and based on genome-wide genetic relatedness for cardiac conduction traits. Twin Research and Human Genetics **20**(6), 489–498 (2017)

29. Watanabe, K., Stringer, S., Frei, O., Mirkov, M.U., de Leeuw, C., Polderman, T.J., van der Sluis, S., Andreassen, O.A., Neale, B.M., Posthuma, D.: A global overview of pleiotropy and genetic architecture in complex traits. Nature genetics **51**(9), 1339–1348 (2019)

30. Martin, A.R., Daly, M.J., Robinson, E.B., Hyman, S.E., Neale, B.M.: Predicting polygenic risk of psychiatric disorders. Biological psychiatry **86**(2), 97–109 (2019)

31. Mars, N., Koskela, J.T., Ripatti, P., Kiiskinen, T.T., Havulinna, A.S., Lindbohm, J.V., Ahola-Olli, A., Kurki,
M., Karjalainen, J., Palta, P., et al.: Polygenic and clinical risk scores and their impact on age at onset and
prediction of cardiometabolic diseases and common cancers. Nature Medicine, 1–9 (2020)

32. Gibson, G.: On the utilization of polygenic risk scores for therapeutic targeting. PLoS genetics **15**(4) (2019)

33. Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M., Lee, J.J.: Second-generation plink: rising to
the challenge of larger and richer datasets. Gigascience **4**(1), 13742–015 (2015)

34. Yang, J., Lee, S.H., Goddard, M.E., Visscher, P.M.: Gcta: a tool for genome-wide complex trait analysis. The
American Journal of Human Genetics **88**(1), 76–82 (2011)

35. Robin, X., Turck, N., Hainard, A., Tiberti, N., Lisacek, F., Sanchez, J.-C., Müller, M.: proc: an open-source
package for r and s+ to analyze and compare roc curves. BMC bioinformatics **12**(1), 77 (2011)

36. Choi, S.W., O'Reilly, P.F.: Prsice-2: Polygenic risk score software for biobank-scale data. Gigascience **8**(7), 082
(2019)

37. Watanabe, K., Taskesen, E., Van Bochoven, A., Posthuma, D.: Functional mapping and annotation of genetic
associations with fuma. Nature communications **8**(1), 1–11 (2017)

38. de Leeuw, C.A., Mooij, J.M., Heskes, T., Posthuma, D.: Magma: generalized gene-set analysis of gwas data.
PLoS computational biology **11**(4) (2015)

39. Lloyd-Jones, L.R., Zeng, J., Sidorenko, J., Yengo, L., Moser, G., Kemper, K.E., Wang, H., Zheng, Z., Magi, R.,
Esko, T., et al.: Improved polygenic prediction by bayesian multiple regression on summary statistics. Nature
communications **10**(1), 1–11 (2019)

Figures

Additional Files

Additional file 1 — Supplementary tables & figures

PDF file of supplementary figures S1-S6 and supplementary tables 1-3.

Figure 1 Recovery of causal variants in fixed simulations. (A) Scatterplot showing proportion of causal variants with P-values below a threshold of $5x10^{-8}$ in 100 simulations using the PGS-LMM (y-axis) and fastGWA (x-axis) methods. The upper and lower horizontal lines show the median number of causal variants recovered with the PGS-LMM and fastGWA methods, respectively. The black line represents the line through the origin with slope 1. (B) Proportion of causal variants identified by fastGWA, PGS-LMM and BOLT-LMM.

Figure 2 Sensitivity analysis. Difference in sensitivity (between the PGS-LMM and fastGWA methods) as a function of specificity. The red line shows the mean difference over all simulations.

Figure 3 Recovery of causal variants varying parameters. Proportion of causal variants identified as a function of (a) the narrow-sense heritability of the simulated trait and (b) the number of causal variants.

Figure 4 Effects of sample size on PGS-LMM. Proportion of causal variants identified as a function of sample size.

Figure 5 PGS-LMM prediction analysis P+T prediction (A) across range of p value thresholds for each method. Bayesian inference prediction (B) across PIP scores. Summary statistics are calculated on the 80% training data