1

Factors enforcing the species boundary between the human pathogens *Cryptococcus*

neoformans* and *Cryptococcus deneoformans*

Shelby J. Priest[1], Marco A. Coelho[1], Verónica Mixão[2,3], Shelly Clancey[1], Yitong Xu[4], Sheng

Sun[1], Toni Gabaldón[2,3,5], and Joseph Heitman[1]*

[1] Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, NC, USA

[2] Life Sciences Department, Barcelona Supercomputing Center, Barcelona, Spain

[3] Institute for Research in Biomedicine, Barcelona Institute of Science and Technology, Barcelona, Spain

[4] Program in Cell and Molecular Biology, Duke University Medical Center, Durham, NC, USA

[5] Catalan Institution for Research and Advanced Studies (ICREA), Barcelona, Spain

Short title: Factors enforcing the *Cryptococcus* species boundary

* Corresponding author

Email: heitm001@duke.edu (JH)

Abstract

Hybridization has resulted in the origin and variation in extant species, and hybrids continue to arise despite pre- and post-zygotic barriers that limit their formation and evolutionary success. One important system that maintains species boundaries in prokaryotes and eukaryotes is the mismatch repair pathway, which blocks homeologous recombination between divergent DNA sequences. Previous studies illuminated the role of the mismatch repair component Msh2 in blocking genetic recombination between divergent DNA during meiosis in yeast and during conjugational and transductional crosses in bacteria. Loss of Msh2 resulted in increased interspecific genetic recombination in bacterial and yeast models, and increased viability of progeny derived from yeast hybrid crosses. Hybrid isolates of two pathogenic fungal *Cryptococcus* species, *Cryptococcus neoformans* and *Cryptococcus deneoformans*, are isolated regularly from both clinical and environmental sources. In the present study, we sought to determine if loss of Msh2 would relax the species boundary between *C. neoformans* and *C. deneoformans*. We found that crosses between these two species in which both parents lack Msh2 produced hybrid progeny with increased viability and high levels of aneuploidy. Whole-genome sequencing revealed few instances of recombination among hybrid progeny and did not identify increased levels of recombination in progeny derived from parents lacking Msh2. Several hybrid progeny from bilateral *msh2Δ* crosses produced structures associated with sexual reproduction when incubated alone on nutrient-rich medium in light, a novel transgressive phenotype in *Cryptococcus*. These findings represent a unique, unexpected case where rendering the mismatch repair system defective did not result in increased meiotic recombination across a species boundary. This suggests that alternative pathways limit meiotic recombination between

homeologous DNA and enforce species boundaries in the basidiomycete *Cryptococcus* species

and possibly in others as well.

Author summary

Several mechanisms enforce species boundaries by either preventing the formation of

hybrids, known as pre-zygotic barriers, or preventing the viability and fecundity of hybrids,

known as post-zygotic barriers. Despite these barriers, interspecies hybrids form at an

appreciable frequency, such as hybrid isolates of the human fungal pathogenic species,

Cryptococcus neoformans and *Cryptococcus deneoformans*, which are regularly isolated from

both clinical and environmental sources. *C. neoformans* x *C. deneoformans* hybrids are typically

highly aneuploid, sterile, and display phenotypes intermediate to those of either parent, although

self-fertile isolates and transgressive phenotypes have been observed. One important mechanism

known to enforce species boundaries or lead to incipient speciation is the DNA mismatch repair

system, which blocks recombination between divergent DNA sequences during meiosis. The aim

of this study was to determine if genetically deleting the DNA mismatch repair component Msh2

would relax the species boundary between *C. neoformans* and *C. deneoformans*. Progeny derived

from *C. neoformans* x *C. deneoformans* crosses in which both parental strains lacked Msh2 had

higher viability, several progeny displayed a novel transgressive phenotype, and unlike previous

studies in *Saccharomyces*, these *Cryptococcus* hybrid progeny had higher levels of aneuploidy

and no observable increase in meiotic recombination at the whole-genome level.

Introduction

The mixing of species through sexual reproduction can result in the production of hybrid

offspring. While hybridization can have beneficial consequences in some cases (e.g. hybrid vigor

and the emergence of novel hybrid species), sexual reproduction between diverging lineages or

different species is typically deleterious and results in hybrid progeny with reduced fitness or

sterility [1-3]. Thus, mechanisms preventing such events, such as pre- and post-zygotic

reproductive barriers, tend to be favored by natural selection. Pre-zygotic species barriers block

the formation of a hybrid zygote and include spatial, temporal, behavioral, mechanical, and

gametic isolation [4]. In the event that a hybrid zygote forms, several post-zygotic barriers exist

that inhibit the viability or fecundity of the hybrid, including 1) gross chromosomal

rearrangements that prevent effective meiotic recombination, 2) Bateson-Dobzhansky-Muller

incompatibilities in which interactions between nuclear elements or between mitochondrial and

nuclear factors are detrimental or lethal, and finally, 3) the mismatch repair (MMR) pathway,

which has an important role in blocking meiotic recombination between diverged DNA

sequences.

The MMR pathway was first identified in prokaryotes as a mechanism to repair

replication errors or damage-induced mismatches in DNA [5]. The MMR pathway is highly

conserved, playing similar roles in unicellular eukaryotes, such as *Saccharomyces* species, and in

multicellular eukaryotes, including humans [6]. Prokaryotic and eukaryotic cells lacking

functional MMR components typically have increased mutation rates and therefore display a

hypermutator phenotype [7]. The MMR pathway plays an additional role in maintaining species

boundaries by inhibiting homeologous chromosome pairing and subsequent recombination

during meiosis. The inability of chromosomes to properly and stably pair, and subsequently

undergo recombination, can lead to chromosome nondisjunction during meiosis, resulting in high

frequencies of aneuploidy and ultimately a decrease in the viability of progeny. Rayssiguier et al.

demonstrated that mutation of the MutL, MutS, or MutH MMR components relaxed the species

boundary between *Escherichia coli* and *Salmonella typhimurium*, two bacterial species whose

genomes are 20% divergent, such that recombination during conjugational and transductional

crosses increased up to 1,000-fold [8]. The involvement of MMR in recombination is conserved

and has also been shown to enforce the species boundary between *Saccharomyces cerevisiae* and

the closely related species *Saccharomyces paradoxus*, whose genomes are ~11% divergent in

coding regions and ~20% divergent in non-coding regions. Mutants lacking the eukaryotic MutS

homolog, Msh2, or the MutL homolog, Pms1, produce hybrid *S. cerevisiae* x *S. paradoxus*

progeny with higher rates of viability as well as approximately ten-fold increased frequencies of

meiotic recombination [9,10].

Despite the numerous pre- and post-zygotic species barriers and the robust fitness defects

associated with hybrids, hybridization occurs at an appreciable frequency. For example, the ursid

hybrids produced by matings between grizzly and polar bears are frequently observed in nature

and genomic analyses suggest ancient hybridization events between brown and polar bears, as

the mitochondrial genomes of brown bears are derived from polar bears [11]. Ligers, the hybrid

progeny of a male lion and female tiger, and mules, the hybrid progeny of a male donkey and

female horse, are examples of hybrids that normally only occur through human intervention.

Ligers are bred for their size, as they are larger than either parent (an instance of hybrid vigor),

while mules are bred for their endurance, docile demeanor, and intelligence [12]. In an exciting

recent study, researchers have even witnessed the origin and monitored the evolution of a novel

finch species that arose through a hybridization event in the Galapagos Islands [13].

Hybridization has impacted recent human evolution too, with several modern human populations

having introgressed genomic regions from Neanderthals or Denisovans [14].

Hybridization is also important in many microbial pathogens and model organisms. For

instance, the diploid model organism *S. cerevisiae* is thought to have arisen following a whole-

genome duplication that was a direct consequence of interspecies hybridization [15-17]. An

instance of relatively recent hybridization is also thought to have led to the emergence of the

novel widespread fungal plant pathogen species *Zymoseptoria pseudotritici*, originating from

fusion between two diverged haploids followed by mitosis and meiosis to generate a

recombinant haploid F_1 hybrid [18]. In recent years, the hybrid nature of several emerging

human opportunistic pathogens has been uncovered [19,20], suggesting that hybridization might

be a mechanism underlying the emergence of novel pathogens [21].

Several *Cryptococcus* species are microbial human fungal pathogens and are responsible

for over 200,000 infections in both immunocompromised and immunocompetent individuals

annually [22]. Cryptococcal infections are associated with high mortality rates and occur

globally. The pathogenic *Cryptococcus* species complex was recently divided into seven distinct

species based on phylogenetic analysis of multiple genomic loci [23]. The pathogenic

Cryptococcus species complex is comprised of two phylogenetically-based subgroups, the

Cryptococcus neoformans species complex and the *Cryptococcus gattii* species complex, which

consist of two and six species, respectively. The present study focuses on the two members of the

C. neoformans species complex: *C. neoformans* and *C. deneoformans* [23,24].

Previously, *C. neoformans* and *C. deneoformans* were considered to be a single species

with two varieties and two serotypes: *C. neoformans* var. *grubii* (serotype A) and *C. neoformans*

var. *neoformans* (serotype D) [23]. These two species are the most commonly isolated

Cryptococcus species from clinical and environmental settings and both species serve as model

pathogenic eukaryotic organisms [25]. Delineating *C. neoformans* and *C. deneoformans* as two

distinct species is in relative agreement with each of the three most common species delimitation

approaches: the phylogenetic, morphological, and biological species concepts [26-29]. In line

with the phylogenetic definition, there is clear genetic evidence separating these two groups and

molecular phylogenetics along with whole-genome sequencing suggests they diverged ~18

million years ago [30-33]. There are also several phenotypes that differentiate *C. neoformans* and

C. deneoformans as distinct species, such as differences in thermotolerance, capsular

agglutination reactions, morphology during murine infection, and human disease manifestations

and outcomes [34-38].

Despite the genetic and phenotypic differences between *C. neoformans* and *C.*

deneoformans, these two groups produce hybrids in the laboratory and in nature. *C. neoformans*

x *C. deneoformans* hybrids, also known as AD hybrids, make up ~7.5% of environmental

isolates and up to 30% of clinical isolates in Europe and North America [39-42]. Spores

produced by genetic crosses between *C. neoformans* and *C. deneoformans* isolates are known to

have poor germination frequencies (~5%), and are typically highly aneuploid or heterozygous

diploids [43]. This poor viability is likely due to a combination of gross chromosomal

rearrangements between the two parental genomes along with ~15% sequence divergence

between the parental species [31,33], leading to a compromised meiosis that produces genetically

imbalanced meiotic progeny. *C. neoformans* x *C. deneoformans* hybrids also have unstable

karyotypes, can be self-fertile, and display phenotypes intermediate between either parent,

although instances of transgressive phenotypes and hybrid vigor have been observed [43-46].

Several studies of *C. neoformans* x *C. deneoformans* hybrid genomes have utilized restriction

fragment length polymorphism analysis or PCR with sequence-specific primers to assess their

genomes. Through these methods it has been demonstrated that although *C. neoformans* x *C.*

deneoformans hybrids are heterozygous at most loci, some chromosomes seem to be

recombinant, indicative of potential mitotic or meiotic recombination [45,47-49].

In the present study, we generated *C. neoformans* and *C. deneoformans* strains lacking

Msh2 to determine if loss of MMR relaxed the boundary between these two species. As

expected, *C. neoformans* and *C. deneoformans msh2Δ* mutants displayed hypermutator

phenotypes. Hybrid progeny derived from genetic crosses in which both parental strains lacked

Msh2 displayed increased germination frequencies compared to wild-type crosses and also

exhibited phenotypes and genotypes in accordance with previous findings [43]. Several instances

of genetic recombination were observed in hybrid progeny derived from both wild-type *C.*

neoformans x *C. deneoformans* crosses and *msh2Δ* mutant crosses, although interestingly,

increased frequencies of meiotic recombination were not observed in hybrid progeny derived

from crosses involving *msh2Δ* mutants. Additionally, lower rates of loss of heterozygosity

(LOH) and higher rates of aneuploidy were observed in hybrid progeny from *msh2Δ* mutant

crosses. These results suggest other pathways and mechanisms besides MMR may enforce

species boundaries in *Cryptococcus*.

Results

***C. deneoformans msh2Δ* mutants are hypermutators**

To assess the role of the MMR pathway in maintaining species boundaries in

Cryptococcus, we first determined if Msh2 plays a similar role in DNA MMR in *C.*

deneoformans as in other fungi. Deletion mutants lacking *MSH2* were generated via biolistic

transformation and homologous recombination in the *C. deneoformans* JEC20**a** genetic

background. A schematic depicting the deletion is shown in Figure S1A. After transformation,

mutants were selected on medium containing nourseothricin, and PCR was employed to confirm

that the deletion allele had replaced the wild-type *MSH2* allele at its endogenous locus (Figure

S1B).

Following isolation and confirmation of the desired *msh2Δ* mutants, we determined if the

mutants displayed hypermutator phenotypes similar to those observed in *msh2* mutants of other

fungi [50,51]. To assess mutation rate, fluctuation assays were performed on either YPD medium

supplemented with a combination of rapamycin and FK506 at 37°C (where calcineurin, the

target of FKBP12-FK506, is essential) or YNB medium supplemented with 5-fluoroorotic acid

(5-FOA) at 30°C. In this analysis, the parental strain of the genetic deletion mutants, JEC20**a**,

served as the negative control, and a *C. neoformans msh2Δ* mutant in the KN99α genetic

background from the *Cryptococcus* deletion mutant collection [52] served as the positive control.

Although fluctuation assays do not provide genome-wide mutation rates, the assays allow us to

compare the mutation rates at two unique coding loci of the JEC20**a** *msh2Δ* mutants to those of

the JEC20**a** parental strain and the KN99α *msh2Δ* positive control. On the rapamycin and

FK506 antifungal drug combination, the *msh2Δ-1, msh2Δ-2, msh2Δ-3,* and *msh2Δ-4* mutants

exhibited hypermutator phenotypes with significantly higher mutation rates (1.02×10^{-6}, $1.30 \times$

10^{-6}, 1.16×10^{-6}, and 1.16×10^{-6} mutations per cell per generation, respectively) than the parental

JEC20**a** strain (8.59×10^{-8} mutations per cell per generation) (Figure 1A).

Insertion/deletion mutations (INDELs) in homopolymeric nucleotide runs are a hallmark

mutation pattern of *msh2Δ* mutants [50,53]. To determine if the four independent JEC20**a** *msh2Δ*

mutants also displayed similar mutation patterns, *FRR1*, the gene encoding FKBP12 (the

common target of FK506 and rapamycin), was PCR amplified and mutations were identified

through Sanger sequencing. This analysis revealed single base pair INDELs in homopolymeric

nucleotide runs within *FRR1* in ten of ten independent colonies analyzed for of the *msh2Δ-1*,

msh2Δ-2, and *msh2Δ-4* strains and nine of ten independent colonies analyzed for the *msh2Δ-3*

strain (Figure 1B). This is significantly different from the frequency of 1-bp INDEL mutations

observed in the wild-type JEC20**a** parental strain, in which only one of nine colonies had a 1-bp

INDEL mutation in a homopolymer run (p<0.001, Fisher's exact test). As anticipated, the

KN99α *msh2Δ* mutant produced resistant colonies with 1-bp INDEL mutations in homopolymer

runs in *FRR1* in all (10/10) colonies analyzed (Figure 1B).

　　　Mutation rates of three of the four independent *msh2Δ* mutants (*msh2Δ-2*, *msh2Δ-3*, and

msh2Δ-4) were also significantly higher than the mutation rate of the parental strain, JEC20**a**, on

5-FOA (Figure 1C). Interestingly, the *msh2Δ-1* mutant failed to produce any 5-FOA resistant

isolates during the fluctuation assay, which is unusual as even JEC20**a** yielded 5-FOA resistant

colonies in this assay despite its relatively low mutation rate. The inability of the *msh2Δ-1*

mutant to produce 5-FOA-resistant colonies, which is typically mediated by loss-of-function

mutations in *URA5* or *URA3*, was explained by an INDEL in a homopolymeric nucleotide run in

FUR1, which encodes a uracil phosphoribosyl transferase involved in the pyrimidine salvage

pathway (deletion of a thymine at position 570229 on Chr5 (AE017345.1), GeneID: CNE02100

(AE017345.1:569,042-570,826)). Mutations in components of the pyrimidine *de novo*

biosynthesis pathway, such as *URA3*, have been demonstrated to be synthetically lethal with

FUR1 mutations in yeast [54,55]. Independent overnight cultures of each of the independent

JEC20**a** *msh2Δ* mutants, a KN99α *fur1Δ* mutant, the JEC20**a** parental strain (negative control),

and a KN99α *msh2Δ* mutant (positive control) were swabbed on YNB plates supplemented with

5-FOA to illustrate that both JEC20**a** *msh2Δ-1* and KN99α *fur1Δ* mutants are incapable of

producing 5-FOA resistant colonies (Figure 1D). Furthermore*, fur1Δ* mutants are also known to

be resistant to the antifungal drug 5-fluorouridine (5FU) and the clinically relevant antifungal

drug 5-fluorocytosine (5FC) [56]. Papillation assays with the same strains used on 5-FOA were

also conducted on YNB plates supplemented with 5FC or 5FU. As anticipated, the JEC20**a**

msh2Δ-1 and the KN99α *fur1Δ* mutant displayed congruent growth phenotypes and were

resistant to both 5FC and 5FU, unlike the parental JEC20**a**, the KN99α *msh2Δ* strain, or any of

the other JEC20**a** *msh2Δ* mutants (Figure 1E and Figure 1F).

Progeny from *msh2Δ* hybrid genetic crosses display increased viability

Interspecific crosses involving MMR-deficient *Saccharomyces* strains produce progeny

with increased viability [9,10]. To determine if a similar increase in viability would be observed

in *Cryptococcus*, genetic crosses were conducted between *C. neoformans* and *C. deneoformans*

wild-type strains as well as corresponding *msh2Δ* mutants. A cross between H99α (a laboratory

standard *C. neoformans* reference strain) and JEC20**a** served as a control for the *msh2Δ* hybrid

crosses. H99α x JEC20**a** interspecific crosses produced robust mating structures, including

hyphae, basidia, and basidiospores (Figure 2A). Interestingly, bilateral crosses in which both

parent strains lacked *MSH2* produced similarly abundant hyphae as those produced by the H99α

x JEC20**a** cross, but exhibited more bald basidia and therefore fewer basidiospore chains (Figure

2A, 2B). In these bilateral crosses, 66%, 64%, 63%, and 57% of the basidia produced by each of

the four independent JEC20**a** *msh2Δ* mutants, respectively, lacked basidiospores compared to the

34% of bald basidia produced by wild-type H99α x JEC20**a** crosses (Figure 2A, 2B).

Basidiospores from wild-type, unilateral *msh2Δ* x wild type, and bilateral *msh2Δ* x

msh2Δ C. neoformans x *C. deneoformans* genetic crosses involving each of the four independent

JEC20**a** *msh2Δ* mutants and the KN99α *msh2Δ* strain were randomly dissected via

micromanipulation onto nutrient-rich YPD medium. The spores were allowed to germinate for

up to three weeks on YPD medium at room temperature, and total germination frequencies were

calculated (Figure 2B and Table S1). Dissected basidiospores from wild-type H99α x JEC20**a**

crosses germinated at an average frequency of 5.8%, which is similar to the previously published

frequency of 5% [43]. Basidiospores from unilateral crosses in which only one parent lacked

MSH2 germinated at highly variable frequencies, ranging from an average of 2% to 11% (Figure

2B and Table S1). Progeny from bilateral crosses involving *msh2Δ* mutants in both parents

displayed increased germination frequencies compared to the progeny from the respective

unilateral crosses. Progeny germination frequencies were as high as 41% for some bilateral

msh2Δ x *msh2Δ* sexual crosses (Table S1), nearing the expected upper germination frequency

limit for *C. neoformans* x *C. deneoformans* crosses based on the presence of one large reciprocal

chromosomal translocation between *C. neoformans* H99α and *C. deneoformans* JEC21α, a strain

congenic to JEC20**a** with the exception of the *MAT* locus [33,57].

Hybrid progeny are capable of hyphal growth and sporulation on nutrient-rich medium

and display phenotypes and genotypes typically associated with *C. neoformans* x *C.*

***deneoformans* hybrids**

Due to the potentially unstable nature of hybrid progeny from *C. neoformans* x *C.*

deneoformans matings, progeny from crosses were allowed to germinate for up to three weeks

following dissection on YPD plates, compared to the standard 5-day incubation time [44,58].

During this prolonged incubation period, YPD agar plates were kept sealed with parafilm on the

benchtop. Surprisingly, after 14 days of incubation, aerial filaments began to emerge from the

periphery of several germinated progeny dissected from bilateral *msh2Δ C. neoformans* x *msh2Δ*

C. deneoformans crosses. Through microscopic analysis we found that these progeny were

producing structures resembling those observed during sexual reproduction under mating-

inducing conditions. Continued incubation and additional microscopic analysis revealed that

after approximately three weeks of incubation, these progeny produced basidia and sparse

basidiospores under conditions previously not known to support sexual reproduction of

Cryptococcus (rich medium, incubated in the light in sealed plates) (Figure 3).

The hybrid progeny from both wild-type *C. neoformans* x *C. deneoformans* crosses, as

well as those from unilateral and bilateral *msh2Δ* crosses, exhibited typical *Cryptococcus* hybrid

genotypes and phenotypes. With sequence-specific primers for the gene *STE20*, which exists as

two mating-type specific alleles and encodes a kinase involved in the pheromone response

signaling cascade, the mating types of the hybrid progeny were determined. Nearly all progeny

inherited and maintained both the *STE20*α allele from the *C. neoformans* parent and the *STE20***a**

allele from the *C. deneoformans* parent, which was expected due to the diploid hybrid

characteristic of *C. neoformans* x *C. deneoformans* hybrids (Figure S2) [43,46]. Based on flow

cytometry, the ploidy of the majority of the hybrid progeny was diploid or aneuploid (Figure S3).

Three out of the 27 progeny for which whole genome sequencing was obtained (progeny YX1,

YX3, and YX6) were, however, estimated to be close to haploid, and all three were from a wild-

type *C. neoformans* x *C. deneoformans* cross (Figure S3).

Although the majority of naturally isolated *C. neoformans* x *C. deneoformans* hybrid

strains are not self-fertile, *C. neoformans* x *C. deneoformans* hybrids produced under laboratory

conditions, similar to those presented in this study, are often self-fertile [43,44]. This fertility is

characterized by the ability to produce hyphae when incubated alone on mating-inducing media,

such as MS agar plates, at room temperature in the dark. All but one of the hybrid progeny

assessed produced hyphae on MS medium and many progeny were also capable of producing

basidia and basidiospores (Figure S4). The one progeny that was not self-fertile (YX3) had lost

the *C. deneoformans MAT*a locus (Figure S2 and S4).

Hybrid progeny from *msh2Δ* genetic crosses are highly aneuploid or diploid and do not

exhibit increased meiotic recombination frequencies

Following isolation and characterization of hybrid progeny from the wild-type and

unilateral and bilateral *msh2Δ* mutant crosses, high molecular weight DNA was isolated from 27

hybrid progeny derived from wild-type (H99α x JEC20**a**), unilateral (KN99α *msh2Δ* x JEC20**a**

and H99α x JEC20**a** *msh2Δ-1*), and bilateral (KN99α *msh2Δ* x JEC20**a** *msh2Δ-1*) crosses for

whole-genome sequencing. The chromosomal composition of the hybrid progeny, along with the

identification of potential sites of meiotic recombination, LOH events, and heterozygosity across

the genome, was assessed by employing the analytic pipeline described in Figure S5 (see

methods for details). To determine ploidy, read depth was assessed in conjunction with flow

cytometry data (Figure 4, S2, and S5).

Hybrid progeny from crosses between wild-type *C. neoformans* and *C. deneoformans*

parental strains, as well as those from unilateral and bilateral *msh2Δ* hybrid crosses, displayed

high rates of aneuploidy (Figure 4, Figure S6). Whole-genome sequencing revealed five

instances of potential meiotic recombination in the hybrid progeny, in which a single copy of a

chromosome was inherited and was chimeric, meaning that various loci along the same

chromosome were homologous to either the *C. neoformans* or the *C. deneoformans* parent, but

not both. For example, in progeny YX3, a portion of *C. neoformans* Chr7 (homologous to *C.*

deneoformans chromosome 6) was inherited solely from the H99α parent while the remaining

portion of the chromosome was inherited from the JEC20**a** parent (Figure 4A). Other examples

of these instances of recombination are seen in progeny YX1 on Chr2 and Chr10, in progeny

YX6 on *C. neoformans* Chr5/*C. deneoformans* Chr4, and also again in progeny YX3 on *C.*

neoformans Chr14/*C. deneoformans* Chr8. All five of these instances of potential meiotic

recombination identified were in progeny derived from the wild-type cross, rather than from

crosses involving *msh2Δ* mutants. Interestingly, hybrid progeny from bilateral *msh2Δ* mutant

crosses had ploidies closer to 2N, were nearly completely heterozygous across a majority of the

genome, and displayed fewer LOH events (Figure 4, S3, S6, S7). Quantifying this

heterozygosity, hybrid progeny derived from bilateral *msh2Δ* x *msh2Δ* mutant crosses had

significantly more heterozygosity ($p < 0.05$, Kruskal-Wallis test, Dunn's test) across the entirety

of their genomes with an average of 96% heterozygosity relative to hybrid progeny derived from

wild-type H99α x JEC20**a** crosses, which had an average of 67% heterozygosity across their

genomes (Figure S7 and Table S2). Another unexpected finding was a phenomenon associated

only with hybrid progeny derived from unilateral or bilateral *msh2Δ* crosses in which

chromosome breaks occurring at centromeres, transposable elements, and at other regions in the

genome appeared to have been repaired by *de novo* addition of telomeric repeats at breakpoints

(Figure 4C, Figure S6, and Figure S8).

Due to the high levels of heterozygosity across the genomes of many of the *C.*

neoformans x *C. deneoformans* hybrid progeny, it is possible that some recombination events

might not be detected by merely assessing the depth of reads aligning to each parental reference

strain. For instance, if a hybrid progeny inherited both homeologous chromosomes involved in a

meiotic reciprocal recombination event, read depth coverage would be equal across both parental

chromosomes. To ensure that additional recombination events like these were not being missed,

Illumina paired-end reads were aligned to a combined reference that included both parental

genomes to identify additional possible recombination sites. This analysis was based on the

assumption that if a recombination event occurred, the two reads of a pair would align to

different chromosomes. Different filtering thresholds were applied based on mapping quality and

the number of read-pairs supporting the event. The results of these analyses, using more or less

stringent filtering thresholds to detect recombination events, show similar frequencies of

recombination events for each of the different types of crosses: wild-type *C. neoformans* x *C.*

deneoformans crosses, unilateral crosses in which one parent lacks Msh2, and bilateral crosses in

which both of the parental strains lack Msh2 (Figure 5 and Table S3). With the strictest filtering

thresholds (at least 15 read-pairs supporting the event and a mapping quality of 60), a median of

1 recombination event is detected across the whole genomes of progeny derived wild-type,

unilateral KN99α *msh2Δ* x JEC20**a**, and bilateral crosses, while a median of 1.5 events are

detected in progeny from unilateral H99α x JEC20**a** *msh2Δ-1* crosses. Furthermore, with the

least strict thresholds (at least 5 read-pairs supporting and mapping quality greater than or equal

to 0), a median of 10 recombination events are detected in progeny derived from a wild-type

cross, a median of 8.5 and 6.5 events are detected from both unilateral crosses (H99α x JEC20**a**

msh2Δ-1 and KN99α *msh2Δ* x JEC20**a**, respectively), and a median of 5.5 events are detected in

progeny from bilateral crosses (Figure 5, Table S3).

Discussion

In this study, we investigated the role of MMR in maintaining the species boundary

between two closely related and prevalent human fungal pathogens, *C. neoformans* and *C.*

deneoformans. In addition to its role in DNA repair, the MMR pathway plays a highly conserved

role in blocking recombination between homeologous DNA sequences during meiosis, serving as

a post-zygotic barrier. Findings from previous studies in prokaryotic and eukaryotic models

indicated that rendering the MMR pathway nonfunctional by genetically deleting pathway

components, particularly Msh2, allows increased homoleogous recombination during meiosis [8-

10]. Therefore, we hypothesized that lack of Msh2 in *Cryptococcus* would allow increased

pairing and recombination between homeologous chromosomes during meiosis, leading to

decreased rates of chromosome nondisjunction, and ultimately the production of more viable

progeny with fewer instances of aneuploidy.

We genetically engineered *C. deneoformans* strains lacking *MSH2* and confirmed

through fluctuation analysis and characterization of mutational spectra that Msh2 plays a similar

role in DNA MMR in *C. deneoformans* as has been previously observed across eukaryotes and in

other pathogenic *Cryptococcus* species [50,51]. Although all of the independent *C.*

deneoformans msh2Δ mutants displayed increased rates of mutation that conferred resistance to

FK506 and rapamycin, the *msh2Δ-1* mutant failed to produce 5-FOA resistant colonies, which

was attributable to a frameshift mutation identified in the *FUR1* that is synthetically lethal with

mutations that can confer 5-FOA resistance. This conclusion was further supported by the

inability of a *C. neoformans fur1Δ* mutant to produce 5-FOA resistant colonies as well as the

inherent 5FC and 5FU resistance displayed by both the *msh2Δ-1* and *fur1Δ* mutants [56]. This

finding illustrates that, in some instances, *Cryptococcus* is unable to acquire concurrent

resistance to both 5-FOA and the clinically relevant drug 5-flucytosine (5FC) as well as the

antimetabolite 5-fluorouracil (5FU). However, there are additional mutations that confer 5FC

and 5FU resistance, such as those in the *UXS1* gene, which encodes a UDP-glucuronate

decarboxylase, which are not presumed to be synthetically lethal with mutations conferring 5-

FOA resistance [56].

Similar to results from previous studies in prokaryotic and eukaryotic microorganisms,

germination frequencies of *C. neoformans* x *C. deneoformans* hybrid progeny derived from

bilateral *msh2Δ* crosses increased substantially relative to wild-type hybrid germination

frequencies [8-10]. Although loss of *MSH2* could lead to increased frequencies of homeologous

recombination in *Cryptococcus* and subsequently increased viability of progeny, it is also

possible that loss of *MSH2* contributes to spore inviability due to the hypermutator phenotype of

msh2Δ mutants, as has been observed in previous studies in *S. cerevisiae* [59]. Despite the low

germination frequencies of some progeny isolated from unilateral *msh2Δ* crosses, progeny

derived from crosses that involved the same independent JEC20**a** *msh2Δ* mutant (i.e. H99α x

JEC20**a** *msh2Δ-1* and KN99α *msh2Δ* x JEC20**a** *msh2Δ-1*) all had increased viability in bilateral

crosses compared to the corresponding unilateral crosses.

Hybrid progeny derived from genetic crosses between *C. neoformans* and *C.*

deneoformans isolates are known to display unique phenotypes. For example, hybrids, especially

those generated under laboratory conditions, can exhibit self-fertility, producing hyphae and in

some cases basidia and spores under mating-inducing conditions [43,46]. The majority of the

hybrid progeny derived from the crosses described here displayed phenotypes in accord with

previously published findings for *C. neoformans* x *C. deneoformans* hybrids [43,44]. However, a

novel transgressive phenotype was exhibited by multiple progeny isolated from bilateral *msh2Δ*

mutant crosses: the ability to produce sexual reproduction structures on nutrient-rich YPD

medium in sealed plates in the light. This was surprising, because environmental cues known to

suppress mating in *Cryptococcus* include nutrient-rich environments, such as YPD medium, as

well as light and high levels of humidity and carbon dioxide [37,60-63]. The production of

sexual reproduction structures by these hybrid progeny under nutrient-rich, light, sealed

conditions represents sexual reproduction of *Cryptococcus* in a novel environment. The isolation

of self-fertile progeny provides an important example of how hybridization can enable

Cryptococcus to access a dimorphic state under conditions that are not conducive for the parental

isolates to sexually reproduce. The ability to produce spores, the infectious propagules of

Cryptococcus human pathogens, in a previously prohibitive environment also provides the

opportunity to produce more infectious propagules, which could potentially result in a higher rate

of infection. This is in line with previous studies which found that up to 30% of clinical isolates

in Europe were *C. neoformans* x *C. deneoformans* hybrids [40-42]. Future studies using

quantitative trait loci mapping as well as other molecular genetic techniques could provide

insight into the genetic determinants contributing to this novel self-fertility phenotype.

Unlike their haploid parents, *C. neoformans* x *C. deneoformans* hybrid progeny typically

have relatively unstable aneuploid or diploid karyotypes [44-46]. All hybrid progeny derived

from *C. neoformans* x *C. deneoformans* crosses in this study were aneuploid based on both

FACS analysis and whole-genome sequencing data. We observed instances of low read-depth

associated with several aneuploid chromosomes, which are thought to represent loss of

chromosomes in fractions of populations of cells that were sequenced at the whole-genome level,

reflecting the karyotypic instability of these progeny. The majority (21/27) of the hybrid progeny

were diploid (11/27 progeny) or nearly diploid (2N+1 or 2N-1) (10/27 progeny), and all progeny

derived from unilateral and bilateral *msh2Δ* crosses were close to diploid. The higher ploidy

levels in progeny derived from crosses involving *msh2Δ* mutants were unexpected because

MMR mutants in yeast produce progeny with lower levels of aneuploidy [9]. It is possible that

the diploids or near diploid *C. neoformans* x *C. deneoformans* progeny were the meiotic products

of a tetraploid, generated by the fusion of two diploids that formed within the mating patch.

However, interspecific progeny derived from divergent *Saccharomyces* tetraploids display much

higher germination frequencies (>90%), which is very different from the frequencies associated

with these *C. neoformans* x *C. deneoformans* hybrid progeny, making it unlikely that they are

derived from tetraploids [64]. It is also possible that the high frequency of diploid progeny could

be indicative of failed meiosis. However, *Cryptococcus* mutants lacking known meiotic genes,

such as those encoding the meiosis-specific endonuclease Spo11 or the key meiotic regulator

Dmc1, are unable to produce basidiospores at an appreciable frequency and their basidia lack the

typical four chains of basidiospores [65,66]. In contrast to the abnormal sexual reproduction

structures produced by these meiosis-deficient mutants, the basidia produced by hybrid crosses

in this study generated four lengthy spore chains, albeit at a lower frequency than intra-species

crosses in some cases. The higher ploidy in hybrid progeny from bilateral *msh2Δ* x *msh2Δ*

crosses was also associated with significantly more heterozygosity across their genomes than

their counterparts from wild-type *C. neoformans* x *C. deneoformans* crosses. It is possible that

this higher level of heterozygosity contributes to the increased viability by masking deleterious

alleles or overcoming Bateson-Dobzhansky-Muller genetic incompatibilities, as has been

observed in previous studies of *C. neoformans* x *C. deneoformans* hybrids [49,67].

An interesting phenomenon associated with progeny derived from unilateral and bilateral

msh2Δ crosses was the *de novo* addition of telomeric repeat sequences to various locations in the

genome including centromeric repeat sequences, non-centromeric transposable elements, and the

rDNA repeat locus. Although the addition of telomeric repeats at non-telomeric sites can

promote the stabilization of a broken chromosome, it also leads to the loss of a large portion of a

chromosome, which would normally threaten cell viability. However, the presence of both

homeologous chromosomes from each parental species in these hybrid progeny alleviates this

problem. *De novo* telomeric repeat addition has also been observed in *Cryptococcus* following

CRISPR-mediated double-stranded breaks at centromeres [68]. Overall, our results suggest that

in *C. neoformans* x *C. deneoformans* crosses involving *msh2Δ* mutants, chromosomes may be

more prone to double-stranded breaks, or that the normally occurring double-stranded breaks are

unable to be properly repaired.

Many instances of loss of heterozygosity and potential recombination were observed in

the hybrid progeny assessed in this study. Three hybrid progeny isolated from wild-type *C.*

neoformans x *C. deneoformans* crosses were close to haploid with five or fewer aneuploid

chromosomes. Conversely, hybrid progeny derived from unilateral or bilateral *msh2Δ* crosses

were more highly aneuploid or diploid. Although instances of recombination were detected in

progeny derived from each of the different types of crosses (wild-type, unilateral, and bilateral

msh2Δ), it is difficult to determine if these recombination events occurred during meiosis or

during subsequent mitotic growth during the culturing required for whole-genome sequencing.

Despite this inherent difficulty, we expected to see significantly higher rates of potential

instances of meiotic recombination based on previous studies in both prokaryotes and eukaryotes

[8-10]. Surprisingly, based on read-depth analysis as well as detecting recombination events by

aligning whole-genome sequencing from the progeny to a combined reference genome with both

parental species, no increase in recombination was observed in crosses involving either a single

parent lacking Msh2 or both parents lacking Msh2. One potential explanation for this

observation could be that the level of sequence divergence between *C. deneoformans* and *C.*

neoformans is large enough that meiotic recombination will be inefficient, even in the absence of

MMR. However, previous studies on meiotic recombination in *S. cerevisiae* indicate that

although recombination is less efficient at high levels of sequence divergence, loss of MMR

leads to approximately a 24-fold increase in meiotic recombination between 85% divergent

sequences, the same level of divergence as between *C. neoformans* and *C. deneoformans*

[31,33,69]. Furthermore, studies on mitotic recombination in *S. cerevisiae* also similarly indicate

that even at 26% sequence divergence, loss of MMR leads to a 55-fold increase in mitotic

recombination [70,71].

In summary, this study illustrates several key findings on the roles of MMR in

Cryptococcus. In *C. deneoformans*, *msh2Δ* mutants behave as hypermutators on various selective

media and acquire INDELs in homopolymeric nucleotide runs. Hybrid *C. neoformans* x *C.*

deneoformans progeny dissected from genetic crosses involving *msh2Δ* mutants generally

displayed increased germination frequencies compared to those from wild-type crosses. Hybrid

progeny derived from *msh2Δ* crosses displayed phenotypes and karyotypes characteristic of *C.*

neoformans x *C. deneoformans* hybrid strains, such as diploidy/aneuploidy and self-fertility,

although some progeny derived from bilateral *msh2Δ* crosses displayed a novel, transgressive

phenotype in which they were capable of producing hyphae, basidia, and basidiospores on a

glucose- and nutrient-rich medium in the light. The increased viability of hybrid progeny derived

from bilateral *msh2Δ* crosses suggests that loss of Msh2 in *Cryptococcus* may allow

homologous chromosomes to pair more efficiently during meiosis. However, it is interesting to

note that loss of Msh2 did not seem to increase the frequency of meiotic recombination between

C. neoformans and *C. deneoformans* homeologous chromosomes. These results suggest that

alternative pathways or additional MMR components may play different or more important roles

in maintaining species boundaries in *Cryptococcus* than in other previously studied organisms,

an intriguing avenue for future studies. Unlike bacterial and budding yeast studies, loss of Msh2

did not relax the species boundary between *C. neoformans* and *C. deneoformans,* illustrating

robust post-zygotic mechanisms that ultimately maintain integrity between these two closely

related species.

Materials and methods

Strains and growth

The *C. neoformans* and *C. deneoformans* strains described in this study are listed in Table

S4. Strains were stored at -80°C in liquid yeast extract peptone dextrose (YPD) supplemented

with 15% glycerol. Strains were inoculated on YPD agar plates, initially grown at 30°C for 3

days, and then maintained at 4°C. Due to the hypermutator phenotypes associated with *msh2Δ*

strains and the genomic instability associated with *C. neoformans* x *C. deneoformans* hybrids,

strains used in the experiments of this study were not maintained for more than two weeks at 4°C

on YPD agar plates and fresh cells from frozen glycerol stocks were inoculated to new YPD agar

plates at the end of each two-week period.

Generation of *msh2Δ* deletion mutants

The open-reading frame of the *MSH2* gene (gene ID: CNA07480) in the *C. deneoformans*

JEC20**a** genetic background was replaced with the gene encoding the dominant drug-resistance

marker for nourseothricin resistance, *NAT*, by homologous recombination via biolistic

transformation as previously described [72]. Following transformation and selection on YPD +

100 µg/mL nourseothricin agar medium, genomic DNA was isolated from candidate mutants

with the MasterPure DNA purification kit (Epicentre) and PCR followed by gel electrophoresis

confirmed correct integration of the *NAT* dominant resistance marker. The locations of the

primers used to generate the deletion allele and confirm deletion mutants are depicted in Figure

S1A and their sequences are given in Table S5.

Fluctuation analysis to quantify mutation rates

Fluctuation analysis was utilized to quantify the mutation rates, or the number of

mutations per cell per generation, of the JEC20**a** *msh2Δ* mutants. The wild-type strain JEC20**a**

served as a negative control and the *msh2Δ* mutant from the 2015 Madhani deletion collection in

the KN99α genetic background served as a positive control [52]. For each strain, including

controls, ten 5 mL YPD liquid cultures were each inoculated with a single colony from a YPD

agar stock plate. Cultures were incubated overnight at 30°C. After incubation, cultures were

pelleted at 3,000 x *g*, washed twice with 5 mL dH$_2$O, and resuspended in 4 mL dH$_2$O. 100 µL of

undiluted, washed cells were plated directly to YPD + 100 ng/mL rapamycin + 1 µg/mL FK506

or yeast nitrogen base (YNB) + 1 mg/mL 5-fluoroorotic acid (5-FOA) solid agar medium.

Washed cells were diluted 1:100,000 and 100 µL of the dilution was plated to YPD solid agar

medium. Inoculated YPD and YNB+5-FOA plates were incubated at 30°C for 4 or 14 days,

respectively. Inoculated YPD+rapamycin+FK506 plates were incubated at 37°C for 14 days.

Following incubation, colonies on each of the media were counted and mutation frequencies

were calculated with the FluCalc program, which utilizes the Ma-Sandri-Sarkar maximum-

likelihood estimation (MSS-MLE) equation for calculations [73].

Mutation spectra analysis

Single resistant colonies from fluctuation analyses were streak purified to YPD +

rapamycin + FK506 medium and grown for 3 days at 37°C. Genomic DNA was isolated using

the MasterPure Yeast DNA Purification Kit (Epicenter Biotechnologies, Madison, WI), and the

FRR1 gene was PCR-amplified with Phusion High-Fidelity DNA Polymerase (NEB, Ipswich

MA, USA). PCR products were subjected to gel electrophoresis, extracted using a QIAgen gel

extraction kit, and mutations were identified through classical Sanger Sequencing at Genewiz.

Fisher's exact probability test was used to calculate statistically significant differences between

the frequencies of 1-bp INDEL mutations compared to other types of mutations in

YPD+rapamycin+FK506-resistant colonies from strains lacking *MSH2* compared to the wild-

type JEC20**a** strain using the VassarStats online software (http://vassarstats.net).

Papillation assays

Papillation assays were conducted on YNB solid agar medium supplemented with 1

mg/mL 5-FOA, 100 µg/mL 5-fluorocytosine (5FC), or 100 µg/mL 5-fluorouridine (5FU). For

this assay, ten independent YPD liquid cultures were inoculated with ten single colonies from

YPD agar plates and incubated overnight at standard laboratory conditions. Following overnight

culture, cells were pelleted at 3,000 x *g* and resuspended in 2 mL dH$_2$O. Sterile cotton swabs

were then used to inoculate quadrants of YNB + 5-FOA agar plates with each independent

overnight culture. The inoculated agar plates were incubated at 30°C for 6 days to allow

sufficient growth to visualize resistant colonies.

Genetic crosses and progeny dissection

All genetic crosses were conducted on Murashige and Skoog (MS) agar medium

following Basic Protocol 1 for mating assays as previously described [58]. The frequency of bald

basidia was calculated by imaging random areas surrounding two mating patches per cross,

counting the number of bald basidia and basidia producing basidiospores to determine the

frequency of bald basidia, and the frequencies from the two mating patches were averaged. For

each genetic cross, two independent mating patches were assessed; for each mating patch, over

130 total basidia were assessed across at least 11 images. Images used to quantify bald basidia

were taken on a Zeiss Axio Scope.A1 with camera. Random basidiospore dissection was

performed as described in Basic Protocol 2 [58]. Following dissection, the micromanipulated

basidiospores were germinated for up to 3 weeks on YPD agar plates sealed with parafilm and

incubated on the laboratory benchtop. Images of self-fertile hybrid progeny derived from

bilateral *msh2Δ* crosses producing sexual structures on YPD agar plates following dissection

were taken with an Accu-Scope EXC-500 microscope with an attached Nikon DXM1200F

microscope camera.

Phenotyping and genotyping of hybrid progeny

Primers designed to specifically amplify only the *C. neoformans STE20***a**, *C. neoformans*

*STE20*α, *C. deneoformans STE20***a**, or *C. deneoformans STE20*α alleles aided in identifying the

mating types of the hybrid progeny (primers listed in Table S5). To assess self-filamentation,

hybrid progeny were spotted onto MS agar plates and incubated for 14 days at room temperature

(approximately 24°C) in the dark as described in Basic Protocol 1 [58]. Filamentation was

assessed via microscopy after 14 days of incubation with an Accu-Scope EXC-500 microscope

with an attached Nikon DXM1200F microscope camera.

Flow Cytometry

Cells were patched onto YPD agar medium and incubated at 30°C overnight; strains that

exhibited slow growth were incubated at 30°C for three days. Cells were harvested by scraping a

2 mm sized colony with a toothpick and resuspending in 1 mL PBS buffer. Cells were washed

once with 1 mL PBS and then fixed in 1 mL 70% ethanol overnight at 4°C. After fixation, cells

were washed once with 1 mL NS buffer (10 mM Tris-HCl pH=7.6, 150 mM sucrose, 1 mM

EDTA pH=8.0, 1 mM $MgCl_2$, 0.1 mM $ZnCl_2$, 0.4 mM phenylmethylsulfonyl fluoride, 1 mM β-

mercaptoethanol) and resuspended in 180 μl NS buffer supplemented with 5 μl propidium iodide

(0.5 mg/mL) and 20 μl RNase A (10 mg/mL). Cells were incubated covered, overnight at 4°C

with shaking. Prior to analysis, 50 μl of cells were diluted in 2 mL of 50 mM Tris-HCl (pH=8.0)

in a 5 mL falcon tube. Flow cytometry was performed on 10,000 cells and analyzed on the FL1

channel on a Becton-Dickinson FACScan.

Whole-genome sequencing

Single colonies of strains for whole-genome sequencing were inoculated into 50 mL of

liquid YPD and grown overnight at 30°C in standard laboratory conditions. Overnight cultures

were then pelleted at 3,000 x *g* in a tabletop centrifuge and subsequently lyophilized. High

molecular weight DNA was extracted from lyophilized cells following the CTAB protocol as

previously described [74]. Genomic DNA libraries were prepared for sequencing with a Kapa

Hyper plus library kit for 300 bp inserts and sequenced at the Duke Sequencing and Genomic

Technologies Shared Research core facility. Libraries were sequenced using paired-end, 2 x 150

bp reads on an Illumina HiSeq 4000 platform.

Whole-genome and chromosome composition analyses of the hybrid progeny

C. deneoformans strain JEC21α is the congenic mating partner of JEC20**a**, which was

obtained in an earlier study by selecting *MAT*α progeny after ten rounds of backcrossing to

JEC20**a** [75]. The genomes of *C. deneoformans* JEC21α and JEC20**a** strains are therefore nearly

identical, with the exception of 5,322 SNPs mainly distributed over three genomic regions

(including the mating-type locus) [76]. Because a highly contiguous *de novo* genome assembly

of JEC20**a** is not currently available, the genome assembly of strain JEC21α

(GCA_000091045.1) was used for all comparisons. The nuclear genomes of two parental strains,

C. neoformans H99α and *C. deneoformans* JEC21α were compared by performing whole-

genome alignments with Satsuma (https://github.com/bioinfologics/satsuma2) [77], using default

parameters. The output of Satsuma was input to the visualization tools "BlockDisplaySatsuma"

and "ChromosomePaint", included in the same package to generate a postscript file. For

representation purposes, chromosome color codes were modified in Adobe Illustrator, and

centromeres and other genomic features (rDNA and *MAT* loci) were superimposed at scale based

on their respective genomic coordinates. Characterization of the chromosome composition of the

hybrid progeny followed the procedure summarized in Figure S5A. The combined nuclear

reference genome used in this study was built with the genome assemblies of the two parental

strains after reordering and reorienting the JEC21α contigs to maximize collinearity with the

H99α assembly (Figure S5B), using the Mauve Contig Mover tool [78]. A dot plot analysis

comparing the H99α assembly with the JEC21α rearranged assembly (Figure S5B) was

performed with D-Genies application [79], which uses minimap2 [80] for aligning the two

genomes. Raw Illumina paired-end reads of the selected progeny and the combined reference

genomes were input into the sppIDer pipeline [81], which is a wrapper that sequentially maps the

Illumina short reads to the combined reference, performs quality filtering (MQ > 3), and

generates depth of coverage plots (Figure 4 and S5). For each progeny, the number of

chromosomes and the ploidy were estimated from the sppIDer plots in conjunction with the

flow-cytometry data. Chromosomal aberrations, e.g. due to recombination, or chromosome

breakage followed by *de novo* telomere addition, were inferred from the sppIDer plots, and

further validated by visual inspection of the mapped reads in IGV [82].

Heterozygosity of the hybrid progeny

To inspect the heterozygosity levels of the hybrid progeny, the same set of Illumina

paired-end reads were mapped to the *C. neoformans* H99α reference genome, using the BWA-

MEM short-read aligner (v0.7.17-r1188) with default settings [83]. SNP discovery, variant

evaluation, and further refinements were carried out with the Genome Analysis Toolkit (GATK)

best-practices pipeline [84,85] (v4.0.1.2), including the use of Picard tools to convert SAM to

sorted BAM files, fix read groups (module: 'AddOrReplaceReadGroups';

SORT_ORDER=coordinate), and mark duplicates. Variant sites were identified with

HaplotypeCaller from GATK and only high-confidence variants that passed filtration were

retained (the "VariantFiltration" module used the following criteria: DP < 20 || QD < 2.0 || FS >

60.0 || MQ < 40.0 || SOR > 4.0). The genome-wide level of heterozygosity was defined as the

ratio of the number of heterozygous SNPs divided by the total number of SNPs (i.e.

heterozygous and non-reference homozygous SNPs), and was calculated from the resulting VCF

files on a per-individual basis after extracting the corresponding sites using the module

VariantsToTable from GATK. Due to the high variance in heterogeneity across the genomes of

progeny from the different genetic crosses and unequal numbers of progeny analyzed from each

different genetic cross, the non-parametric Kruskal-Wallis test followed by Dunn's test were

performed using JMP v15 (SAS Institute).

Detection of variants in the genome of the JEC20a *msh2Δ-1* mutant

To predict variants in the JEC20**a** *msh2Δ-1* mutant, we employed the variant calling

procedure described above, using the JEC21 genome (GCA_000091045.1) as reference. The

effect of a variant (SNPs and INDELs) was predicted and annotated using SnpEff [86]. Only

variants of moderate and high impact were considered (i.e. excluding synonymous and non-

coding variants), and variants in common between the mutant and the JEC20**a** parent were

flagged background mutations and excluded. Given the inability of the *msh2Δ-1* mutant to

produce 5-FOA resistant colonies, we specifically focused on mutations in genes involved in the

de novo and salvage pathways of pyrimidine biosynthesis.

Detection of recombination events based on read-pair alignment

To detect possible recombination events in *C. neoformans* x *C. deneoformans* hybrids,

the respective reference genomes were retrieved from NCBI (accession: ASM301198v1 for H99

and ASM9104v1 for JEC21 as no genome assembly for JEC20 was available). As the H99 strain

was not sequenced in this project, the respective Illumina paired-end reads were retrieved from

SRA (SRR7042283). All Illumina paired-end libraries were filtered with the default parameters

of Trimmomatic v0.36 [87]. Filtered reads of each strain (including the parental strains) were

aligned on the combined reference (containing both H99 and JEC21 genome assemblies) with

HaploTypo pipeline [88], using BWA-MEM v0.7015 [83], samtools v1.9 (Li 2011) and GATK

v4.0.2.1 [85].

To detect recombination events, we assumed that in recombinant sites it would be

possible to detect read-pairs with the two reads aligning to different chromosomes. Therefore, for

each strain (including the parental strains), we obtained the coordinates of all the pairs with each

read aligning to a chromosome of a different parental species. Only reads with at least 75 bp and

a maximum of 3 mismatches were considered. As there were some differences between the

parental JEC20 and the reference JEC21, variant calling was performed with HaploTypo pipeline

[88], using GATK v4.0.2.1 [85]. Bedtools v2.26 [90] intersect was used to determine how many

SNPs overlapped each read, and in cases in which more than 3 SNPs would overlap, the filter of

mismatches was adjusted to that value. The final results were filtered based on mapping quality

(MQ) and number of pairs supporting a given recombination event (Bedtools v2.26 [90] merge

with a distance of 100bp). Four different analyses were performed using more or less stringent

filters: i) MQ = 60 and at least 15 reads supporting the event (stricter); ii) MQ = 60 and at least

five reads supporting the event; iii) MQ >= 0 and at least 15 reads supporting the event; iv) MQ

>= 0 and at least five reads supporting the event (less strict). The script that automatizes this

pipeline is available on GitHub (https://github.com/Gabaldonlab/detect_recombination/). As it

was possible to detect recombination in the parental strains (false positives), Bedtools v2.26 [90]

subtract was used to remove all the regions detected in hybrid progeny that were also detected in

the parental strains.

Data availability

715   Raw sequence data has been deposited into the National Center for Biotechnology

716 Information Sequence Read Archive under BioProject accession no. PRJNA626676. The

717 accession numbers for each sample are provided in Table S6.

Acknowledgements

We thank Sue Jinks-Robertson, Tom Petes, and Paul Magwene for critical reading and comments on the manuscript. We thank Jay Jawahar for his assistance in the initial phases of constructing the JEC20**a** *msh2Δ* mutants. We also thank the Madhani Laboratory and NIH grant R01 AI100272 for the KN99α *msh2Δ* and *fur1Δ* deletion strains.

References

1. Arnold ML, Bulger MR, Burke JM, Hempel AL, Williams JH. Natural hybridization: how low can you go and still be important? *Ecology* 1999;80(2): 371-381. doi: 10.1890/0012-9658(1999)080[0371:NHHLCY]2.0.CO;2

2. Goulet BE, Roda F, Hopkins R. Hybridization in plants: old ideas, new techniques. *Plant Physiol*. 2017;173(1):65-78. PMID: 27895205.

3. Vallejo-Marin M, Hiscock SJ. Hybridization and hybrid speciation under global change. *New Phytol*. 2016;211(4):1170-87. PMID: 27214560.

4. Fu C, Coelho MA, David-Palma M, Priest SJ, Heitman J. Genetic and genomic evolution of sexual reproduction: echoes from LECA to the fungal kingdom. *Curr Opin Genet Dev*. 2019;58-59:70-75. PMID: 31473482.

5. Lu AL, Clark S, Modrich P. Methyl-directed repair of DNA base-pair mismatches in vivo. *Proc Natl Acad Sci U S A*. 1983;80(15):4639-43. PMID: 6308634.

6. Grilley M, Holmes J, Yashar B, Modrich P. Mechanisms of DNA-mismatch correction. *Mutat Res*. 1990;236(2-3):253-67. PMID: 2144613.

7. Harfe BD, Jinks-Robertson S. Mismatch repair proteins and mitotic genome stability. *Mutat Res*. 2000;451(1-2):151-67. PMID: 10915870.

8. Rayssiguier C, Thaler DS, Radman M. The barrier to recombination between *Escherichia coli* and *Salmonella typhimurium* is disrupted in mismatch-repair mutants. *Nature*. 1989;342(6248):396-401. PMID: 2555716.

9. Hunter N, Chambers SR, Louis EJ, Borts RH. The mismatch repair system contributes to meiotic sterility in an interspecific yeast hybrid. *EMBO J*. 1996;15(7):1726-33. PMID: 8612597.

10. Chambers SR, Hunter N, Louis EJ, Borts RH. The mismatch repair system reduces meiotic

homeologous recombination and stimulates recombination-dependent chromosome loss. *Mol*

Cell Biol. 1996;16(11):6110-20. PMID: 8887641.

11. Pongracz JD, Paetkau D, Branigan M, Richardson E. Recent hybridization between a polar

bear and grizzly bears in the Canadian Arctic. *Arctic*. 2017;70(2):151-60. doi:

10.14430/arctic4643

12. Darwin C. *What Mr. Darwin saw in his voyage round the world in the ship "Beagle."* New

York: Harper & Bros; 1879.

13. Lamichhaney S, Han F, Webster MT, Andersson L, Grant BR, Grant PR. Rapid hybrid

speciation in Darwin's finches. *Science*. 2018;359(6372):224-8. PMID: 29170277.

14. Kelso J, Prufer K. Ancient humans and the origin of modern humans. *Curr Opin Genet Dev*.

2014;29:133-8. PMID: 25286439.

15. Dietrich FS, Voegeli S, Brachat S, Lerch A, Gates K, Steiner S, et al. The *Ashbya gossypii*

genome as a tool for mapping the ancient *Saccharomyces cerevisiae* genome. *Science*.

2004;304(5668):304-7. PMID: 15001715.

16. Wolfe KH. Origin of the yeast whole-genome duplication. *PLOS Biol*. 2015;13(8):e1002221.

PMID: 26252643.

17. Marcet-Houben M, Gabaldón T. Beyond the whole-genome duplication: phylogenetic

evidence for an ancient interspecies hybridization in the baker's yeast lineage. *PLOS Biol*.

2015;13(8):e1002220. PMID: 26252497.

18. Stukenbrock EH, Christiansen FB, Hansen TT, Dutheil JY, Schierup MH. Fusion of two

divergent fungal individuals led to the recent emergence of a unique widespread pathogen

species. *Proc Natl Acad Sci U S A*. 2012;109(27):10954-9. PMID: 22711811.

19. Pryszcz LP, Nemeth T, Saus E, Ksiezopolska E, Hegedusova E, Nosek J, et al. The genomic

aftermath of hybridization in the opportunistic pathogen *Candida metapsilosis*. *PLOS Genet*.

2015;11(10):e1005626. PMID:26517373.

20. Mixão V, Hansen AP, Saus E, Boekhout T, Lass-Florl C, Gabaldón T. Whole-genome

sequencing of the opportunistic yeast pathogen *Candida inconspicua* uncovers its hybrid

origin. *Front Genet*. 2019;10:383. PMID: 31105748.

21. Mixão V, Gabaldón T. Hybridization and emergence of virulence in opportunistic human

yeast pathogens. *Yeast*. 2018;35(1):5-20. PMID: 28681409.

22. Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, et al. Global

burden of disease of HIV-associated cryptococcal meningitis: an updated analysis. *The*

Lancet Infect Dis. 2017;17(8):873-81. PMID: 28483415.

23. Hagen F, Khayhan K, Theelen B, Kolecka A, Polacheck I, Sionov E, et al. Recognition of

seven species in the *Cryptococcus gattii*/*Cryptococcus neoformans* species complex. *Fungal*

Genet Biol. 2015;78:16-48. PMID: 25721988.

24. Farrer RA, Chang M, Davis MJ, van Dorp L, Yang D, Shea T, et al. A new lineage of

Cryptococcus gattii (VGV) discovered in the Central Zambezian Miombo Woodlands. *mBio*.

2019;10(6):e02306-19. PMID: 31719178.

25. Chayakulkeeree M, Perfect JR. Cryptococcosis. *Infect Dis Clin North Am*. 2006;20(3):507-

44. PMID: 16984867.

26. Cracraft J. Species concepts and speciation analysis. In: Johnston RF, editors. *Current*

Ornithology. New York, NY: Springer; 1983. p. 159-187.

27. Cracraft J. Species concepts and the ontology of evolution. *Biol Philos*. 1987;2:329-346.

doi:10.1007/BF00128837

28. Cronquist A. Once again, what is a species? In: Knutson LV, editor. *Biosystems in*

Agriculture. Montclair, NJ: Allanheld Osmun; 1978. p. 3-20.

29. Mayr E, Ashlock PD. *Principles of Systematic Zoology*. New York: McGraw-Hill; 1991.

30. Xu J, Vilgalys R, Mitchell TG. Multiple gene genealogies reveal recent dispersion and

hybridization in the human pathogenic fungus *Cryptococcus neoformans*. *Mol Ecol*.

2000;9(10):1471-81. PMID: 11050543.

31. Kavanaugh LA, Fraser JA, Dietrich FS. Recent evolution of the human pathogen

Cryptococcus neoformans by intervarietal transfer of a 14-gene fragment. *Mol Biol Evol*.

2004;23(10):1879-90. PMID: 16870684.

32. Loftus BJ, Fung E, Roncaglia P, Rowley D, Amedeo P, Bruno D, et al., The genome of the

basidiomycetous yeast and human pathogen *Cryptococcus neoformans*. *Science*.

2005;307(5713):1321-4. PMID: 15653466.

33. Janbon G, Ormerod KL, Paulet D, Byrnes EJ 3rd, Yadav V, Chatterjee G, et al. Analysis of

the genome and transcriptome of *Cryptococcus neoformans* var. *grubii* reveals complex RNA

expression and microevolution leading to virulence attenuation. *PLOS Genet*.

2014;10(4):e1004261. PMID: 24743168.

34. Martinez LR, Garcia-Rivera J, Casadevall A. *Cryptococcus neoformans* var. *neoformans*

(serotype D) strains are more susceptible to heat than *C. neoformans* var. *grubii* (serotype A)

strains. *J Clin Microbiol*. 2001;39(9):3365-7. PMID: 11526180.

35. Feldmesser M, Kress Y, Casadevall A. Dynamic changes in the morphology of *Cryptococcus*

neoformans during murine pulmonary infection. *Microbiology*. 2001;147(Pt 8):2355-65.

PMID: 11496012.

36. Desnos-Ollivier M, Patel S, Raoux-Barbot D, Heitman J, Dromer F, The French

Cryptococcosis Study Group. Cryptococcosis serotypes impact outcome and provide

evidence of *Cryptococcus neoformans* speciation. *mBio*. 2015;6(3):e00311-15. PMID:

26060271.

37. Casadevall A, Perfect JR. *Cryptococcus neoformans*. Washington, DC: ASM Press; 1998.

38. Dromer F, Mathoulin-Pelissier S, Launay O, Lortholary O, The French Cryptococcosis Study

Group. Determinants of disease presentation and outcome during cryptococcosis: the Crypto

A/D Study. *PLOS Med*. 2007;4(2):e21. PMID: 17284154.

39. Litvintseva AP, Kestenbaum L, Vilgalys R, Mitchell TG. Comparative analysis of

environmental and clinical populations of *Cryptococcus neoformans*. *J Clin Microbiol*.

2005;43(2):556-64. PMID: 15695645.

40. Cogliati M, Esposto MC, Clarke DL, Wickes BL, Viviani MA. Origin of *Cryptococcus*

neoformans var. *neoformans* diploid strains. *J Clin Microbiol*. 2001;39(11): 3889-94. PMID:

11682503.

41. Frases S, Ferrer C, Sanchez M, Colom-Valiente MF. Molecular epidemiology of isolates of

the *Cryptococcus neoformans* species complex from Spain. *Rev Iberoam Micol*.

2009;26(2):112-7. PMID: 19631160.

42. Maduro AP, Mansinho K, Teles F, Silva I, Meyer W, Martins ML, et al. Insights on the

genotype distribution among *Cryptococcus neoformans* and *C. gattii* Portuguese clinical

isolates. *Curr Microbiol*. 2014;68(2):199-203. PMID: 24077953.

43. Lengeler KB, Cox GM, Heitman J. Serotype AD strains of *Cryptococcus neoformans* are

diploid or aneuploid and are heterozygous at the mating-type locus. *Infect Immun*.

2001;69(1):115-22. PMID: 11119496.

44. Lin X, Litvintseva AP, Nielsen K, Patel S, Floyd A, Mitchell TG, et al. αADα hybrids of

Cryptococcus neoformans: evidence of same-sex mating in nature and hybrid fitness. *PLOS*

Genet. 2007;3(10):1975-90. PMID: 17953489.

45. Kwon-Chung KJ, Varma A. Do major species concepts support one, two or more species

within *Cryptococcus neoformans*. *FEMS Yeast Res*. 2006;6(4):574-87. PMID: 16696653.

46. Li W, Averette AF, Desnos-Ollivier M, Ni M, Dromer F, Heitman J. Genetic diversity and

genomic plasticity of *Cryptococcus neoformans* AD hybrid strains. *G3 (Bethesda)*.

2012;2(1):83-97. PMID: 22384385.

47. Sun S, Xu J. Genetic analyses of a hybrid cross between serotypes A and D strains of the

human pathogenic fungus *Cryptococcus neoformans*. *Genetics*. 2007;177(3):1475-1486.

PMID: 17947421.

48. Vogan A, Khankhet J, Xu J. Evidence for mitotic recombination within the basidia of a

hybrid cross of *Cryptococcus neoformans*. *PLOS One*. 2013;8(5):e62790. PMID: 23690954.

49. Samarasinghe H, Vogan A, Pum N, Xu J. Patterns of allele distribution in a hybrid

population of the *Cryptococcus neoformans* species complex. *Mycoses*. 2019;63(3):275-83.

PMID: 31774582.

50. Billmyre RB, Clancey SA, Heitman J. Natural mismatch repair mutations mediate

phenotypic diversity and drug resistance in *Cryptococcus deuterogattii*. *eLife* 2017;6:e28802.

PMID: 28948913.

51. Boyce KJ, Wang Y, Verma S, Shakya VPS, Xue C, Idnurm A. Mismatch repair of DNA

replication errors contributes to microevolution in the pathogenic fungus *Cryptococcus*

neoformans. *mBio*. 2017;8(3):e00595-17. PMID: 28559486.

52. Liu OW, Chun CD, Chow ED, Chen C, Madhani HD, Noble S. Systematic genetic analysis

of virulence in the human fungal pathogen *Cryptococcus neoformans*. *Cell*. 2008;135(1):174-

88. PMID: 18854164.

53. Tran HT, Keen JD, Kricker M, Resnick MA, Gordenin DA. Hypermutability of

homonucleotide runs in mismatch repair and DNA polymerase proofreading yeast mutants.

Mol Cell Biol. 1997;17(5):2859-65. PMID: 9111358.

54. Koren A, Ben-Aroya S, Steinlaur R, Kupiec M. Pitfalls of the synthetic lethality screen in

Saccharomyces cerevisiae: an improved design. *Curr Genet*. 2003;43(1):62-9. PMID:

12684846.

55. Paluszynski JP, Klassen R, Meinhardt F. Genetic prerequisites for additive or synergistic

actions of 5-fluorocytosine and fluconazole in baker's yeast. *Microbiology*. 2008;154(Pt

10):3154-64. PMID: 18832321.

56. Billmyre RB, Clancey SA, Xi LX, Doering TL, Heitman J. 5-fluorocytosine resistance is

associated with hypermutation and alterations in capsule biosynthesis in *Cryptococcus*. *Nat*

Commun. 2020;11(1):127. PMID: 31913284.

57. Sun S, Xu J. Chromosomal rearrangements between serotype A and D strains in

Cryptococcus neoformans. *PLOS One*. 2009.4(5):e5524. PMID: 19436753.

58. Sun S, Priest SJ, Heitman J. *Cryptococcus neoformans* mating and genetic crosses. *Curr*

Protoc Microbiol. 2019;53(1):e75. PMID: 30661293.

59. Alani E, Reenan RAG, Kolodner RD. Interaction between mismatch repair and genetic

recombination in *Saccharomyces cerevisiae*. *Genetics*. 1994;137(1):19-39. PMID: 8056309.

60. Alspaugh JA, Perfect JR, Heitman J. *Cryptococcus neoformans* mating and virulence are

regulated by the G-protein alpha subunit GPA1 and cAMP. *Genes Dev*. 1997;11(23):3206-

17. PMID: 9389652.

61. Idnurm A, Heitman J. Light controls growth and development via a conserved pathway in the

fungal kingdom. *PLOS Biol*. 2005;3(4):e95. PMID: 15760278.

62. Heitman J, Kozel TR, Kwon-Chung KJ, Perfect JR, Casadevall A, editors. *Cryptococcus:*

From Human Pathogen to Model Yeast. Washington, DC: ASM Press; 2011.

63. Sun S, Coelho MA, David-Palma MA, Priest SJ, Heitman J. The evolution of sexual

reproduction and the mating-type locus: links to pathogenesis of *Cryptococcus* human

pathogenic fungi. *Annu Rev Genet*. 2019;53:417-44. PMID: 31537103.

64. Greig D, Borts RH, Louis EJ, Travisano M. Epistasis and hybrid sterility in *Saccharomyces*.

Proc Biol Sci. 2002;269(1496):1167-71. PMID: 12061961.

65. Feretzaki M, Heitman J. Genetic circuits that govern bisexual and unisexual reproduction in

Cryptococcus neoformans. *PLOS Genet*. 2013;9(8):e1003688. PMID: 23966871.

66. Lin X, Hull CM, Heitman J. Sexual reproduction between partners of the same mating type

in *Cryptococcus neoformans*. *Nature*. 2005;434(7036):1017-21. PMID:15846346.

67. Vogan AA, Xu J. Evidence for genetic incompatibilities associated with post-zygotic

reproductive isolation in the human fungal pathogen *Cryptococcus neoformans*. *Genome*.

2014;57(6):335-44. PMID: 25187972.

68. Yadav V, Sun S, Coelho MA, Heitman J. Centromere scission drives chromosome shuffling

and reproductive isolation. *PNAS*. 2020;117(14):7917-28. PMID:32193338.

69. Chen W, Jinks-Robertson S. The role of the mismatch repair machinery in regulating mitotic

and meiotic recombination between diverged sequences in yeast. *Genetics*.

1999;151(4):1299-313. PMID:10101158.

70. Datta A, Adjiri A, New L, Crouse GF, Jinks-Roberston S. Mitotic crossovers between

diverged sequences are regulated by mismatch repair proteins in *Saccharomyces cerevisiae*.

Mol Cell Biol. 1996;16(3):1085-93. PMID: 8622653.

71. Datta A, Hendrix M, Lipsitch M, Jinks-Robertson S. Dual roles for DNA sequence identity

and the mismatch repair system in the regulation of mitotic crossing-over in yeast. *Proc Natl*

Acad Sci U S A. 1997;94(18):9757-62. PMID: 9275197.

72. Toffaletti DL, Rude TH, Johnston SA, Durack DT, Perfect JR. Gene transfer in *Cryptococcus*

neoformans by use of biolistic delivery of DNA. *J Bacteriol*. 1993;175(5):1405-11. PMID:

8444802.

73. Radchenko EA, McGinty RJ, Aksenova AY, Neil AJ, Mirkin SM. Quantitative analysis of

the rates for repeat-mediated genome instability in a yeast experimental system. *Methods Mol*

Biol. 2018;1672:421-38. PMID: 29043640.

74. Pitkin JW, Panaccione DG, Walton JD. A putative cyclic peptide efflux pump encoded by

the *TOXA* gene of the plant-pathogenic fungus *Cochliobolus carbonum*. *Microbiology*.

1996;142(Pt 6):1557-65. PMID: 8704997.

75. Kwon-Chung KJ, Edman JC, Wickes BL. Genetic association of mating types and virulence

in *Cryptococcus neoformans*. *Infect Immun*. 1992;60(2):602-5. PMID: 1730495.

76. Hua W, Vogan A, Xu J. Genotypic and phenotypic analyses of two "isogenic" strains of the

human fungal pathogen *Cryptococcus neoformans* var. *neoformans*. *Mycopathologia*.

2019;184(2):195-212. PMID: 30891668.

77. Grabherr MG, Russell P, Meyer M, Mauceli E, Alfoldi J, Di Palma F, et al. Genome-wide

synteny through highly sensitive sequence alignment: Satsuma. *Bioinformatics*.

2010;26(9):1145-51. PMID: 20208069.

78. Darling ACE, Mau B, Blattner FR, Perna NT. Mauve: multiple alignment of conserved

genomic sequence with rearrangements. *Genome Res*. 2004;14(7):1394-403. PMID:

15231754.

79. Cabanettes F, Klopp C. D-GENIES: dot plot large genomes in an interactive, efficient and

simple way. *PeerJ*. 2018;6:e4958. PMID: 29888139.

80. Li H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*.

2018;34(18):3094-3100. PMID: 29750242.

81. Langdon QK, Peris D, Kyle B, Hittinger CT. spplDer: a species identification tool to

investigate hybrid genomes with high-throughput sequencing. *Mol Biol Evol*.

2018;35(11):2835-49. PMID: 30184140.

82. Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G, et al.

Integrative genomics viewer. *Nat Biotechnol*. 2011;29(1):24-6. PMID: 21221095.

83. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv:1303.3997v2 [q-bio.GN] [Preprint]. 2013 [cited 2020 May 11]. Available from:

arxiv.org/abs/1303.3997

84. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A framework

for variation discovery and genotyping using next-generation DNA sequencing data. *Nat*

Genet. 2011;43(5):491-8. PMID: 21478889

85. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The

Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA

sequencing data. *Genome Res*. 2010;20(9):1297-303. PMID: 20644199.

86. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. A program for

annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. *Fly*

(Austin). 2012;6(2):80-92. PMID: 22728672.

87. Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence

data. *Bioinformatics*. 2014;30(15):2114-20. PMID: 24695404.

88. Pegueroles C, Mixão V, Carreté L, Molina M, Gabaldón T. HaploTypo: A variant-calling

pipeline for phased genomes. *Bioinformatics*. 2019;36(8):2569-71. PMID: 31834373.

89. Li H. A statistical framework for SNP calling, mutation discovery, association mapping and

population genetical parameter estimation from sequencing data. *Bioinformatics*.

2011;27(21):2987-93. PMID: 21903627.

90. Quinlan AR, Hall IM. BEDTools: A flexible suite of utilities for comparing genomic

features. *Bioinformatics*. 2010;26(6):841-2. PMID: 20110278.

91. Idnurm A. Tetrad analysis of the basidiomycete fungus *Cryptococcus neoformans*. *Genetics*.

2010;185(1):153-63. PMID: 20157004.

92. Fraser JA, Huang JC, Pukkila-Worley R, Alspaugh JA, Mitchell TG, Heitman J.

Chromosomal translocation and segmental duplication in *Cryptococcus neoformans*.

Eukaryot Cell. 2005;4(2):401-406. PMID: 15701802.

Figure Captions

Figure 1. Hypermutator phenotypes of JEC20a *msh2Δ* mutants. (A) Fluctuation analysis on

YPD+rapamycin+FK506 medium was performed to quantify the mutation rates (number of

mutations per cell per generation) of four independent JEC20**a** *msh2Δ* mutants: *msh2Δ-1*,

msh2Δ-2, *msh2Δ-3*, and *msh2Δ-4*. The JEC20**a** parental strain in which the *msh2Δ* mutants were

constructed served as the negative control and an *msh2Δ* mutant in the KN99α genetic

background served as the positive control. Points indicate mean mutation rates and error bars

indicate 95% confidence intervals for the mean; 10 independent replicates of each strain were

included in mutation rate calculation. **(B)** Spectra of mutations identified through sequencing of

the *FRR1* gene in rapamycin+FK506-resistant colonies from fluctuation analysis conducted in

panel A. For JEC20**a** n = 9, for all other strains, n = 10. **(C)** Fluctuation analysis on YNB+5-

FOA medium was performed in a similar manner to the analysis in panel A. **(D-F)**

Representative images of plates used in papillation assays with independent JEC20**a** *msh2Δ*

mutants and a KN99α *fur1Δ* mutant on **(D)** YNB+5-FOA medium, **(E)** YNB+5FC medium, and

(F) YNB+5FU medium with the JEC20**a** parental strain and a KN99α *msh2Δ* mutant as controls.

Strains were incubated on YNB+5-FOA medium for 6 days at 30°C before imaging. Strains were

incubated on YNB+5FC and YNB+5FU media for 3 days at 30°C before imaging.

Figure 2. Mating structure formation in *C. neoformans* x *C. deneoformans* genetic crosses

and germination frequencies of hybrid progeny. (A) *C. neoformans* x C. *deneoformans*

genetic crosses produced robust hyphal filamentation after 6 days of incubation in dark

conditions on MS medium. Scale bars in colony border images (left image of each set) represent

200 μm. Scale bars in basidia and basidiospore morphology images (right image of each set)

represent 50 μm **(B)** Average frequencies of basidia lacking basidiospores, or bald basidia, in *C.*

neoformans x *C. deneoformans* genetic crosses. Error bars represent standard deviation. **(C)**

Mean germination frequencies of *C. neoformans* x *C. deneoformans* hybrid progeny in wild-

type, unilateral, and bilateral *msh2Δ* genetic crosses. Following random basidiospore dissection

and micromanipulation onto YPD agar medium, progeny were allowed to germinate for up to

three weeks before germination frequencies were assessed. X-axis labels are the same for (B)

and (C).

Figure 3. Unique phenotypes of *C. neoformans* x *C. deneoformans* hybrid progeny. Sexual

reproduction structures, including hyphae, basidia, and basidiospores, produced by *C.*

neoformans x *C. deneoformans* hybrid progeny derived from bilateral *msh2Δ* genetic crosses

following dissection and micromanipulation onto YPD agar medium. Sexual structures were

formed on YPD agar plates sealed with parafilm and incubated in light conditions at room

temperature. From left to right, scale bars on microscopy images represent 100 μm, 50 μm, and

10 μm.

1004

Figure 4. Nuclear genome composition of *C. neoformans* x *C. deneoformans* hybrid progeny

reveals substantial aneuploidy and recombination. Representative sequencing read-depth

coverage and inheritance patterns of progeny derived from **(A)** a wild-type H99α x JEC20**a**

genetic cross, **(B)** a unilateral H99α x JEC20**a** *msh2Δ* genetic cross, and **(C)** a bilateral KN99α

msh2Δ x JEC20**a** *msh2Δ* genetic cross. For each progeny, sequencing coverage plots (normalized

to the genome-wide average coverage) are colored according to each parental species

contribution as shown in the key on the top right, and a schematic representation of the inferred

karyotype is depicted on the right. Homeologous chromosomes are color coded based on the H99

reference and asterisks in JEC21 indicate chromosomes in reverse-complement orientation (See

Fig S4 for details). Red arrowheads mark recombination breakpoints between homeologous

chromosomes and/or loss of heterozygosity (also highlighted by red boxes in the karyotype

panels). Circular black labels: (a) indicate changes in ploidy in a subset of the population of cells

that were sequenced; (b) and (c) mark, respectively, chromosome breaks at a transposable

element near the end of the *C. deneoformans* Chr6 and at the centromere of Chr9, which were

both repaired by *de novo* telomere addition (green arrowheads). Note the chromosome order of

each parent; JEC21α contigs have been reordered to maximize collinearity with the H99α

contigs.

1022

Figure 5. Distribution of the number of recombination events detected in the hybrid

progeny based on read-pairs aligning to different chromosomes. Four different filtering

thresholds were applied to detect potential instances of recombination across the genomes of 27

total *C. neoformans* x *C. deneoformans* hybrid progeny: (A) Mapping quality (MQ) = 60 and at

least 15 read-pairs (Cov) supporting each event; (B) MQ = 60 and at least 5 read-pairs supporting

each event; (C) MQ >= 0 and at least 15 read-pairs supporting each event; and (D) MQ >= 0 and

at least 5 read-pairs supporting each event. Each point represents the number of recombination

events detected across the whole genome of a single hybrid progeny. In the box and whisker

plots, red lines represent the median, shaded boxes represent the interquartile ranges (IQRs),

upper and lower whiskers show the largest or smallest observations, respectively, that lie 1.5 *

IQR beyond the upper and lower quartiles, respectively. Outliers are included.

1034

Supporting information captions

Table S1. Germination frequencies of hybrid progeny from individual *C. neoformans* x *C. deneoformans* genetic crosses.

Table S2. (A)Heterozygosity levels of the hybrid progeny, and (B) Kruskal-Wallis and Dunn's statistical tests.

Table S3. Recombination events detected in *C. neoformans* x *C. deneoformans* hybrid progeny.

Table S4. Strains used in this study.

Table S5. Oligonucleotides used in this study.

Table S6. NCBI data submissions related to this study.

Figure S1. Genetic deletion and confirmation of the *MSH2* gene in JEC20a. (A) Genetic deletion mutants lacking *MSH2* were engineered in JEC20**a** by replacing the *MSH2* open reading frame (ORF) with the dominant drug resistance marker *NEO* which confers resistance to nourseothricin via biolistic transformation. 5' and 3' UTRs are depicted as yellow boxes. Arrows depict locations of primers used to generate and verify *MSH2* deletion. The sets of primers denoted by gray arrows located upstream and downstream of the *MSH2* ORF were used to amplify flanking sequences homologous to the JEC20**a** *MSH2* endogenous locus to mediate

homologous recombination (JOHE45551, 45552 and JOHE4555, 45556, respectively). Yellow

arrows depict primers (JOHE45553 and JOHE45554) that amplified the *NAT* cassette and share

homology with JOHE45552 and JOHE45556, respectively. The red arrows (JOHE45559,45560)

and blue arrows (JOHE45822,45823) indicate primers that confirmed integration of the deletion

allele and loss of the *MSH2* ORF, respectively. Gray lines indicate syntenic regions shared

between the deletion allele and endogenous locus. **(B)** Gel electrophoresis of PCR products was

used to confirm integration of a single copy of the *NAT* gene at the correct locus and that the

wild-type *MSH2* gene was absent in the JEC20**a** mutant strains. (Spanning: JOHE45559,45560;

WT in-gene: JOHE45822,45823; WT 5' junction: JOHE45559,45823; Δ5' junction:

JOHE45559;45554; WT 3' junction: JOHE45822,45560; Δ3' junction: JOHE45553,45560).

1068

1069 **Figure S2. PCR analysis of mating-type allele inheritance in *C. neoformans* x *C.**

1070 ***deneoformans* hybrid progeny.** Sequence-specific primers for the mating-type (*MAT*) locus

1071 gene *STE20*, which can differentiate between *C. neoformans MAT***a**, *C. neoformans MAT*α, *C.*

1072 *deneoformans MAT***a**, and *C. deneoformans MAT*α, were used to characterize which *MAT* alleles

1073 each of the hybrid progeny inherited.

1074

1075 **Figure S3. FACS analysis of *C. neoformans* x *C. deneoformans* hybrid progeny.** The *C.*

1076 *neoformans* strain H99α was used as a 1N haploid control and the *C. deneoformans* XL143 strain

1077 was used as a diploid 2N control.

1078

1079 **Figure S4. Self-filamentation in *C. neoformans* x *C. deneoformans* hybrid progeny**.

1080 Filamentation of hybrid progeny selected for whole-genome sequencing on MS medium after

1081 incubation for 14 days. AI187 is a self-filamentous, stable diploid *C. neoformans* strain and

1082 served as a positive control. Scale bars represent 100 μm.

1083

Figure S5. Whole-genome comparisons of *C. neoformans* H99α and *C. deneoformans*

JEC21α strains. (A) Workflow to assess the genomic contribution of each parental species in

the hybrid progeny. The sppIDer pipeline uses short-read sequencing data and a combined

genome built from reference genomes of the two parental *Cryptococcus* species. **(B)** Dot-plot

comparing the H99α assembly with the JEC21α reordered and reoriented assembly. Red and blue

lines represent sequence similarities in the same and reverse orientations, respectively. **(C)**

Linear plots showing overall synteny between the H99α and JEC21α genomes. The

chromosomal positions of centromeres and the *MAT* locus are indicated by black and yellow

bars, respectively. Chromosomes of JEC21α are color coded based on their synteny with the

H99α chromosomes. Three major gross chromosomal changes previously documented

distinguishing the two strains correspond to color changes within the same chromosome: TR

indicates a reciprocal chromosomal translocation; INT indicates an introgression of a 14-gene

region from *C. neoformans* to *C. deneoformans* that was mediated by transposable elements

common to both lineages [31]; and SD indicates a segmental duplication following a

nonreciprocal translocation involving the subtelomeric regions of JEC21 chromosomes 8 and 12

that presumably occurred during the construction of the congenic strain pair JEC21α/JEC20**a**

[92]. Chromosomal inversions are not indicated expect for a large inversion on Chr3 of JEC21α

(dashed arrow).

1102

Figure S6. Nuclear genome composition of additional *C. neoformans* x *C. deneoformans*

hybrid progeny. (A) H99α x JEC20**a**. **(B)** H99α x JEC20**a** *msh2Δ-1* **(C)** KN99α *msh2Δ* x

JEC20**a (D)** H99α *msh2Δ* x JEC20**a** *msh2Δ-1*. For each progeny, read-depth plots (normalized to

the genome-wide average coverage) are colored according to each parental species contribution

as shown in the key on the top right, and a schematic representation of the inferred karyotype is

depicted on the right. Homeologous chromosomes are color coded based on the H99 reference

(see Fig S4 for details) and asterisks in JEC21α indicate chromosomes in reverse-complement

orientation. Red arrowheads mark recombination breakpoints between homeologous

chromosomes and/or loss of heterozygosity (also highlighted by red boxes in the karyotype

panels). Where detected, the breakpoints of additional recombination events within the same

chromosome are indicated by light blue arrowheads. Circular black labels: (a) marks changes in

ploidy in a subset of the population of cells that were sequenced; (b) marks chromosome

breakage events repaired by *de novo* telomere addition (see Fig S6 for details); (c) indicates

recombination events next to the *MAT* locus; (d) marks a break at rDNA locus; (e) marks a

complex chromosomal aberration that cannot be explained by simple rearrangements and

required further investigation.

1119

Figure S7. Quantification of heterozygosity across genomes of *C. neoformans* x *C.*

***deneoformans* hybrid progeny.** Plot showing the percentage of heterozygosity for each

individual progeny (represented by different dots) grouped by type of cross (see Table S2 for

details). The horizontal red line depicts the mean heterozygosity values. Genomes of progeny

derived from bilateral *msh2Δ* × *msh2Δ* mutant crosses were significantly more heterozygous

only when compared to hybrid progeny derived from wild-type H99α x JEC20**a** crosses

(Statistical analysis: Kruskal-Wallis test, followed by Dunn's test, $p < 0.05$).

1127

Figure S8. Chromosome breaks in the hybrid progeny can be repaired by *de novo* telomere

addition. Breakpoints were detected in different chromosomal locations, including *CEN14* and

CEN9 of JEC21α (A and B), a T1 transposable element located at the end of Chr6 of JEC21α

(C), or in other genomic locations (D – F). Each panel shows the result of read mapping for one

or more progeny that underwent chromosome breakage and healing via *de novo* telomere

addition (strains names in boldface type) and a control strain in which no breaks were detected

on the same region (strain names in normal font type). Breakage and *de novo* telomere addition

was inferred, respectively, by abrupt changes in read coverage (depicted as bars on the top and

colored as shown in the key) and by the presence of reads with telomeric repeats at the

breakpoints. When two copies of the same chromosome are present, only a subset of reads are

expected to contain telomeric repeats (as shown e.g. in panel F).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5