Playing the piano with a robotic third thumb: Assessing constraints of human augmentation

Ali Shafti[1,2,3]†, Shlomi Haar[1,3]†, Renato Mio[1], Pierre Guilleminot[1], and A. Aldo Faisal[1,2,3,4]*

1. Brain and Behaviour Lab: Dept. of Bioengineering, Imperial College London, SW7 2AZ, London, UK
2. Dept. of Computing, Imperial College London, SW7 2AZ, London, UK
3. Behaviour Analytics Lab, Data Science Institute, SW7 2AZ, London, UK
4. UKRI CDT in AI for Healthcare, Imperial College London, SW7 2AZ, London, UK
5. MRC London Institute of Medical Sciences, W12 0NN, London, UK
† These authors contributed equally to this work as co first authors

* Correspondence: aldo.faisal@imperial.ac.uk

Abstract

Contemporary robotics gives us mechatronic capabilities for augmenting human bodies with extra limbs. However, how our brains and bodies pose limits on such augmentation is an open question. We developed Supernumerary Robotic 3rd Thumbs (SR3T) with two degrees-of-freedom controlled by the user's body to endow them with an extra contralateral thumb on the hand. We demonstrate that a pianist can learn to play the piano with 11 fingers within an hour. We then evaluate 6 naïve and 6 experienced piano players in their prior motor coordination and their capability in piano playing with the robotic augmentation. Intriguingly, individuals' augmented performance did not depend on prior piano experience but could be predicted by our new custom motor coordination assessment, the Human Augmentation Motor Coordination Assessment (HAMCA) performed pre-augmentation. Our work demonstrates how supernumerary robotics can augment humans in skilled tasks and that individual differences in their augmentation capability are predictable by their individual brains' motor coordination abilities.

Introduction

From ancient myths, such as the many-armed goddess Shiva to modern comic book characters, augmentation with supernumerary (i.e. extra) limbs has captured our common imagination. In real life, Human Augmentation is emerging as the result of the confluence of robotics and neurotechnology. We are mechatronically able to augment the human body; from the first myoelectric prosthetic hand developed in the 1940s [1] to the mechanical design, control and feedback interfaces of modern bionic prosthetic hands [e.g. 2–5]. Robots have been used to augment the bodies of disabled humans, restoring some of their original capabilities [e.g. 6–9]. Similar setups can augment healthy users beyond their capabilities, e.g. augmenting workers in industrial settings through intelligent collaboration [e.g. 10–12], or equipping them with additional arms to perform several tasks concurrently [e.g. 13,14]. The latter fits within a particular area of human augmentation robotics which is referred to as supernumerary robotics. These are robotic systems, typically worn by the user, to extend their body and its physical capabilities. However, a major question is, to what extent do the human brain and body have the capability of adapting and learning to use such technologies efficiently [15]. The supernumerary and augmentative nature of this area of research presents an interesting challenge on how to map human motor commands to robot control.

Supernumerary robotic limbs [e.g. 16] are envisioned to assist human factory workers, and adapted for different types of applications [e.g. 14]. The introduction of supernumerary robotic fingers developed as a grasp support device [17] led to further exploration on optimal materials and mechanical designs for supernumerary robotics [e.g. 9,18,19]. Supernumerary robotic fingers have been particularly envisioned,

and successful in grasp restoration for stroke patients [e.g. 9,20,21]. However, regardless of the mechanism,
material, and use case, given the presence of a human within these devices' control loops, the control
interface is of essential importance.

It was recently shown that for polydactyly subjects who possess six fingers on their hands, the
control interface involves a cortical representation of the supernumerary finger [22]. Unlike polydactyly,
supernumerary robotic limbs and fingers must utilize indirect control interfaces to achieve the same
goal – to enable more complex movements and better task performance. Abdi et. al investigate the
feasibility of controlling a supernumerary robotic hand with the foot [23]. Others have focused on
electromyography (EMG) as the control interface – both in supernumerary robotic fingers [e.g. 9], and
supernumerary robotic limbs [e.g. 24]. Other interfaces used for supernumerary robotics include inertial
measurement units [e.g. 21], voice [e.g. 25], pushbuttons [e.g. 18], and graphical user interfaces [e.g. 26]. Researchers
have also explored indirect control interfaces, e.g. using the concept of grasp synergies [27] to assume
that the supernumerary robotic finger's posture will be highly correlated with that of natural fingers
during manipulation, allowing supernumerary robotic finger control through natural movement of
existing fingers [28]. Importantly, all these user interfaces focus on the interface and not the user.

While extensive research has been conducted on the mechanical design, interface, and control of
supernumerary robotics, there is a gap in understanding the role of human motor control in the
success and adoption of these robotic human augmentation systems. In the rapid development of
human augmentation little attention is devoted to how humans interact with the technology and learn
to control it [15]. There are clear needs for neuroscience and robotics research to come together in
analysing such scenarios [e.g. 29]. Learning to control a supernumerary robot limb or finger is a complex
process which involves learning to utilize one movement (set muscles activations) to perform a new
movement. The field of motor neuroscience has extensively studied the control mechanisms and
learning processes of perturbed movements, where we utilize arm movement in one direction to
move a cursor on the screen in a different direction, accounting for a rotation perturbation [30–33] or a
mirror reversal perturbation [e.g. 34–36]. In these settings, one can predict subjects learning from the task-
relevant variability in their unperturbed movements [e.g. 37,38]. Nevertheless, these studies were done on
simplistic lab-based tasks and only recently the field is starting to address the complexities of real-
world movement and to ask to what extent those lab-based findings generalize to real world motor
control and learning [39,40]. While the relationship between task-relevant variability and learning
performance seems to generalize to real-world tasks, defining task relevance is less trivial [e.g. 39] and
the learning mechanism can differ between users [e.g. 41]. In the case of augmentation technology, the
relevant features can be either those related to performing the task itself without the augmented
device, or features related to the control interface of the augmented device.

In human performance research, such as sports science and rehabilitation, there are significant
efforts to predict future performance. In sports science, there is an attempt to predict athletes' future
success for talent identification purposes. Motor coordination and motor learning are often used as
predictors [e.g. 42–45]. Similar approaches are used in rehabilitation research to predict skill learning
capacity following traumatic brain injury, stroke, or neurodegenerative disease [e.g. 46,47]. Here we are
looking into the predictability of future performance with augmentation technology. We specifically

ask which aspect of motor coordination is a better predictor of performance with the device – i.e.
performance in task related tests versus performance in control interface related tests.

Fig. 1. Piano playing task setup. (A) Top view rendering of the SR3T, showing the horizontal motion DOF and relevant motor. **(B)** Side view rendering of the SR3T showing the vertical motion DOF and relevant motor. **(C)** Top view rendering of the SR3T control interface for the 1st degree of freedom (DOF); the participant controls the motion of the SR3T using their right foot, captured through an inertial measurement unit (IMU) worn on the foot. **(D)** Side view rendering of the SR3T control interface for the 2nd DOF. **(E)** Work surface of a human thumb end-point projected on a sphere for comparison with **(F)** the work surface of the SR3T end-point projected on a sphere – augmenting work surface range for the human (see methods). **(G, H)** Top and side view of the unconstrained pilot experiment: an experienced piano player freely improvising on the piano while wearing and making use of the SR3T, effectively playing 11-fingered piano within 1 hour of use. **(I)** Systematic experiments: playing the piano sequence using 5 fingers of the right hand plus the left-hand index finger (LHIF) and **(J)** Playing the sequence using the SR3T. **(K)** A participant plays the sequence of notes as displayed on the monitors in front of them, using the SR3T.

To address this, we have created an experimental setup to study how different parameters within
the remit of human motor control contribute to successful control, coordination and usage of a human
augmentative robotic system, using a set of motor coordination tests that we developed: the Human
Augmentation Motor Coordination Assessment (HAMCA). We have created a 2 degrees of freedom

(DOF) robotic finger, worn on the side of the hand, to augment human finger count to 11, effectively
giving them a 3rd thumb. We call this the supernumerary robotic 3rd thumb (SR3T – engineering and
design previously described in [48]), and we study its usage in a skilled human task: playing the piano.
The piano is a setting which involves the use of all fingers of the hand, and hence a good environment
to consider for testing the augmentation of fingers. Furthermore, piano playing is structured both in
spatial and temporal dimensions, allowing for quantification of the performance in both aspects.

Fig. 2. Human Augmentation Motor Coordination Assessment (HAMCA) – a set of simple behavioural tasks to predict the ability of human augmentation (see Methods for details). (A) Balance board force measurement platform for the Foot-balance task. **(B)** Foot motion trajectory (curve and arrows) during the Foot-tracking task required to move the foot in a figure of 8 in a plane perpendicular to the resting foot's major axis. (Inset) visual feedback to the participant on the computer screen in front of them, showing the desired trajectory (green curve), the red dot indicating the desired location of the foot tip for pacing the foot movement and the current location of the foot tip (blue dot). **(C)** See-saw like foot motion in the sagittal plane of the foot during the Foot up-down task. **(D,E)** Measurement of motor coordination complexity in the fingers of the right hand by assembly of a toy car and of a toy train in the Toy Assembly Task. **(F)** Piano-position task: Piano key sequences to be played with individual fingers to capture hand and finger positional acuity. Performance is assessed by timing and key board press down errors movement between 3 keys (spaced 1 octave apart) labelled "Left", "Middle" and "Right".

Results

We developed a mechanically powerful supernumerary robotic 3rd thumb (SR3T) and means for interfacing it with human users, initially through a combination of foot and thumb motions, directly controlling the two degrees of freedom of the SR3T. We then tested an experienced piano player in an unconstrained pilot experiment, allowing them to freely play the piano and improvise while wearing the SR3T. We observed that within 1 hour of playing the piano, the participant incorporated the SR3T in their piano playing, effectively playing the piano with 11 fingers (see Fig. 1G-H, and supplementary video). Based on this outcome, and feedback from the participant, we upgraded the control interface to be solely based on foot motions (see Fig. 1, and Methods), for more robust control, and to limit the interface to one limb. We then set out to understand the constraints affecting success with the SR3T, by devising protocols and behavioural markers for motor coordination evaluation: the Human Augmentation Motor Coordination Assessment (HAMCA). We also developed a piano sequence playing task as well as measures for assessing the quality of playing. Finally, we systematically evaluated the SR3T on human subjects, and predicted how well subjects would be able to perform in playing the piano sequence with an augmented additional finger, based on the basic motor coordination assay from the HAMCA. Twelve right-handed participants (6 experienced pianists and 6 naïve players), attended 2 experimental sessions held on separate days in the lab. In the first session they performed the HAMCA set of 8 tasks to assess their hand and foot motor coordination. This set was developed to investigate the possibility of a priori prediction of how well each individual human user can learn to use an augmented device. From the HAMCA we extracted 8 scores as measures of hand and foot motor coordination (see Methods). In the second session the subjects learned to play a sequence on the piano and then repeated it with our human augmentation device, the supernumerary robotic 3rd thumb (SR3T), operated through foot motions as the interface (Fig. 1).

The hand and foot motor-coordination scores from the HAMCA, recorded during the first experimental session, showed moderate differences between the pianists and the naïve players (Fig.

Fig. 3. Performance of all subjects in *the HAMCA*. Showing results for 6 naïve (grey dots) and 6 experienced piano players (black dots). **(A)** Accuracy in HAMCA tasks (foot Balance - FB, foot tracking – FT, foot up down spatial – FUD$_S$, foot up down temporal – FUD$_T$, hand dimensionality – HD, piano positioning – PP, piano timing – PT, piano loudness - PL). **(B)** Pearson's correlations between the accuracies of all HAMCA tasks across subjects. The 6 naïve and 6 experienced players were lumped together as individual performances were not different between the groups (see main text).

3A). There were no significant group differences in any of the piano-based tasks (Piano Position, Piano
Timing, and Piano Loudness). The only measure where the pianists performed significantly different
to the naïve players was Hand Dimensionality (p = 0.049), which is based on the assembly of toys (see
Methods). On the other hand, in the Foot Up-Down Spatial measure the naïve players showed higher
scores than the pianists (p = 0.012), though we believe this difference is possibly the result of a sample
bias due to the small N. In both groups the inter-subject variabilities were relatively evenly distributed
except for one pianist who was an outlier showing poor performance on the Foot Balance, Foot
Tracking, and the Piano Loudness tasks.

The correlation matrix between the subjects' motor-coordination scores (Fig. 3B) suggests
relatively weak dependencies, i.e. a subject who showed high coordination in one task did not
necessarily show high coordination in any other task. There were no significant dependencies within
the foot measures and the only dependency within the hand measures was between the Piano
Position and Piano Timing scores (r = 0.64, p = 0.026). There were a few intriguing correlations
between foot and hand measures. First, the foot and hand timing scores were highly correlated (Foot
Up-Down Temporal and Piano Timing: r = 0.68, p = 0.015). Second, the Foot Up-Down Temporal also
correlated with the Piano Position (r = 0.66, p = 0.020). This was expected considering the correlation
between the piano position and timing scores. These three tasks are metronome based, thus
measuring rhythmic coordination. Lastly, we found a significant correlation between Foot Balance and
Piano Loudness scores (r = 0.61, p = 0.035), but this was driven by the pianist who was an outlier in
both tasks.

In the second experimental session, all subjects performed 10 trials of the Piano Playing task, using
their left-hand index finger (LHIF) to play notes further to the right of their right-hand. This was
followed by an additional 10 trials of Piano Playing with the SR3T, where subjects use both degrees of
freedom within the SR3T for the horizontal reach for the notes to the right and the vertical motion to
play the note. In both tasks (playing with the LHIF and with the SR3T) subjects showed improvement
over the first 5 trials after which they plateaued (Fig. 4, right). Thus, for all future analysis we averaged
over trials 5 to 10 to have a single piano playing score for each of these tasks. Here as well, there were
no significant differences between the pianists and the naïve players in any of the trials played with

Fig. 4. Piano playing performance. (A) The note sequence played for the piano playing task. Notes
exclusively played with the right hand, and those with the SR3T or LHIF are marked, **(B)** Visualisation of
how each individual note is scored linearly based on delay from the beat. Incorrect notes and skipped
notes are assigned a score of 0, full sequence score is the average of all individual scores, **(C)** Accuracy
over trials with the SR3T (orange) and without, using the LHIF (blue).

their LHIF (t-test p > 0.12) and with the SR3T (t-test p > 0.32). Testing over all plateau trials (5-10),
pianists were significantly better in playing with their LHIF (t-test p = 0.017) but there were no group
differences in playing with the SR3T (t-test p = 0.9). Therefore, we merged the two groups and all
further analysis was done on all 12 subjects together.

The Piano Playing with SR3T score is our metric for performance with the human augmentation
device, and the fundamental question is to what extent can it be explained by motor-coordination
measures. The correlations between the subjects' scores in the Piano Playing tasks and the motor-
coordination measures suggest different dependencies for playing with and without the SR3T (See
supplementary, Fig. S1 & Fig. S2). The scores in the Piano Playing task with the LHIF, which required
no foot interface, were significantly correlated with Foot Tracking and Foot Up-Down Temporal scores
(r = 0.66, p = 0.019 and r = 0.64, p = 0.026, respectively). The scores in Piano Playing with SR3T were
significantly correlated only with the Piano
Loudness scores (r = 0.59, p = 0.044).

The pianist who was an outlier in few
motor-coordination measures was also an
outlier in the Piano Playing with the SR3T
score (but not in Piano Playing without) and
is driving the correlation with the Piano
Loudness. Thus, we further investigated the
correlations between Piano Playing with
SR3T and the motor-coordination scores
with Spearman rank correlation scores (Fig.
S1). Foot Up-Down Temporal was the only
measure which showed significant
Spearman correlation with the Piano Playing
with SR3T score (r(Spearman) = 0.67, p =
0.02). The Piano Playing scores without and
with SR3T were highly correlated even with
the outlier (r = 0.63, p = 0.028), and even
better correlated with Spearman rank
correlation (r(Spearman) = 0.71, p = 0.012).

Since none of the motor-coordination
scores explained the Piano Playing with
SR3T score well, we asked whether a
combination of motor-coordination scores
can explain it. We specifically asked which
combination could better explain it – that of
the hand measures, which includes piano-
based tasks as well as the only score where
the pianists were significantly better than
the naïve players (Hand Dimensionality); or
that of the foot measures, considering it
being the control interface of the SR3T.
Generalized linear models were fitted to the
hand (Fig. 5A) and foot (Fig. 5B) measures
trying to explain the Piano Playing with SR3T

Fig. 5. Model predictions for SR3T piano playing.
N=11 subjects after outlier removal. Naïve subjects marked as grey and experienced subjects as black dots. **(A)** model from the 4 HAMCA hand dexterity measures. **(B)** model from the 4 HAMCA foot dexterity measures. **(C)** The corrected Akaike information criterion (AICc) for the HAMCA foot and HAMCA hand models, as well as the models with additional component of the LHIF piano playing score, the expertise group (experienced vs naïve), or both.

score. To account for the impact of the outlier subject, we removed them and fitted the Generalized
linear models to the remaining 11 subjects. While both models could explain most of the variance in
the SR3T Piano Playing score, the Generalized linear models based on foot measures showed a much
better fit than the one based on the hand measures (r = 0.79 and r = 0.55, respectively) and was the
only significant fit (p = 0.0035 and p = 0.081, respectively). Moreover, while in the foot model all scores
had positive contributions to the model with similar magnitudes, the hand model was dominated by
Piano Position while Piano Timing had negative contribution and Hand Dimensionality and Piano
Loudness had none. We then tried Generalized linear models where we added the LHIF Piano Playing
score (without SR3T) to the hand and foot models (See supplementary, Fig. S3 A&B). While it
performed better for both models (r = 0.87 and r = 0.92 respectively, p < 0.005), it still showed more
contribution from the foot measures than the hand, and thus better performance in the Foot+LHIF
model.

Next, we tested for the contribution of piano expertise to the prediction by adding it to the model
(Fig. S3 C&D). Adding expertise improved both models, though to a lesser extent than adding the LHIF
Piano Playing score (r = 0.63 and r = 0.89 respectively, p < 0.05). The expertise had no contribution to
the models with the LHIF Piano Playing score (Fig. S3 E&F). To compare between these models of
different complexity we used the corrected Akaike information criterion (AICc) which is modified for
small sample sizes. AICc estimate the amount of information that is lost while fitting a model and thus
can measure the quality of different models relative to each other. The information criterion clearly
show that the foot models are better than the hand models (Fig. 5C). While information criteria might
be biased to select models with more parameters and overfit, the AICc was developed to address such
potential overfitting. With this correction, the AICc suggests that the best model is the Foot model
without the LHIF Piano Playing score and the piano expertise.

Lastly, we fitted all models to all 12 subjects, including the outlier subject who drives the
correlations (See supplementary, Fig. S4). While both hand and foot model could now significantly
explain most of the variance in the SR3T Piano Playing score (p < 0.01) the model based on foot
measures showed a better fit than the one based on the hand measures (r = 0.92 and r = 0.71,
respectively).

Discussion

In this study we addressed a gap in our understanding of human augmentation technology which
is how human-in-the-loop interaction with an augmentative device is learned and performed by the
human brain. We had previously described the different neurocognitive barriers to successful
embodiment and use of robotic augmentation devices [15]. Here, following the operational definition
set out in the same work [15] for the embodiment of robotic augmentation as the ability to use extra
limbs in natural tasks, we focused on how different parameters within the remit of human motor
control contribute to successful control of a supernumerary robotic finger in an augmented piano
playing task. We created a supernumerary robotic 3rd thumb (SR3T), controlled through substitution,
initially with a combination of the natural thumb and the foot wearing different sensing modalities.
We first demonstrated, in an unconstrained pilot experiment, the feasibility of human augmentation
with the SR3T, with one experienced piano player using the device to freely play the piano using 11
fingers, within 1 hour of wearing it. We then updated our SR3T interface, so that both degrees of
freedom are controlled through a single IMU worn on the foot, and developed the HAMCA set to
assess the participants' motor coordination within the hand (task-space) and the foot (interface-
space) prior to piano playing. This is followed by our subjects playing a piano sequence on the piano
using their natural fingers, and then using the SR3T along with their natural fingers. Our findings
suggest that it is not your expertise in the task you perform with the supernumerary robotic finger

(i.e. piano playing expertise), nor your task-space coordination (i.e. motor coordination of the hand
and fingers), but your interface-space coordination (i.e. foot coordination) that can predict your level
of task performance with the augmented device.

While half of our subjects were experienced pianists and the other half naïve, there were not many
significant differences between the two groups within our experiments. The only motor-coordination
score in the HAMCA in which the experienced pianists performed significantly different to the naïve
players was hand dimensionality (HD, see Fig. 3A). The notion that as a skill evolves into an expertise
one learns to use more degrees of freedom in the movement, is known since the pioneering work of
Nikolai Bernstein. Bernstein found that professional blacksmiths use high variability in their joint
angles across repetitive trials to achieve low variability in their hammer end-point trajectory [49].
Pianists need to get their hands to posture with independent control of digits which are not common
in daily life. Thus, they should be able to control more degrees of freedom in their hand movement.
We would have expected to see significant differences on the piano-based tasks as well as timing-
based tasks with the foot, given the pianists' experience. This is, however, not the case within our
performance results (see Fig. 3.A), which might be due to the design of the tasks within HAMCA not
capturing that difference.

Looking at correlations between the different motor coordination scores (Fig. 3.B), we see a high
and significant correlation between Piano Position (PP), Piano Timing (PT) and Foot Up-Down
Temporal (FUD_T) tasks. These are the only three tasks relying on a timing-based measure, with a
metronome-controlled beat. Therefore, they are measuring rhythmic coordination, and presumably
relay on a common timing mechanism [50], and a common coordination-dependent timing network [51].
The strong correlations between these measures is suggesting that rhythmic coordination is a personal
trait similarly performed in both the fingers (for PP and PT) and the foot (FUD_T).

In the piano playing task, all subjects (naïve players and experienced pianists alike) initially showed
improvement in accuracy from trial to trial (i.e. learning). This was a short learning process which
plateaued quickly after 5 trials. For all subjects the plateau with the SR3T was significantly lower than
with their LHIF, which is to be expected, particularly in early stage of SR3T use. This fast learning within
a session and low plateau (which leaves much room for improvement in future sessions) are hallmarks
of early motor skill learning. This is in line with many evidence of multiple time scales in skill learning
where fast improvement in performance occurs in the initial training and plateau within a session, and
slow improvement develops across sessions [e.g. 52–55]. Accordingly, learning to play the piano,
augmented with the SR3T, seems to be a novel motor skill learning task. Further support can be found
in the group differences while playing the piano with and without the SR3T. When subjects played
with their own LHIF, across all plateau trials pianists performed better than naïve players, as expected
based on their piano experience. Though, surprisingly, there were no significant group differences on
a trial by trial basis, not during learning nor during plateau. When subjects played with the SR3T there
were also no significant group differences on a trial by trial basis, but also across all plateau trials
pianists did not perform better than naïve players. This suggests that playing augmented with the SR3T
is not a trivial extension of the regular piano sequence playing task with your own finger, but a novel
skill that the subjects need to learn.

The correlations of the SR3T piano playing score with all motor coordination measures (See
supplementary, Fig. S1) suggest no one-to-one mapping. While most measures showed some positive
correlation trend, hand dimensionality – the best metric for piano playing expertise – showed no
correlation, and even a slightly negative trend. This is in line with the lack of difference between
experienced pianists and naïve players in performance with the SR3T. The two measures that showed
the strongest correlations to the SR3T score were Piano Loudness and Foot Up-Down Temporal. Yet,

Piano Loudness correlation was driven by the outlier and Foot Up-Down Temporal showed significant rank correlation but no Pearson or robust correlation. Overall, no motor coordination measure can predict the SR3T piano playing score by itself. The only measure that showed high correlation with the SR3T score was the LHIF piano playing score. Thus, while piano playing experience showed no significant contribution to the performance in the SR3T task, performance in the same task without the SR3T is a good predictor of the performance with the SR3T. Given the good performance of the LHIF score as a predictor of the SR3T score, it is interesting to consider which of the HAMCA measures correlate with it. Looking at correlations between the coordination measures and the LHIF piano playing score, we see Foot Tracking showing a significant and robust correlation with the LHIF score, rather than any of the hand related scores.

Next, we asked if a combination of motor coordination measures from HAMCA can predict performance with the SR3T, and if so, which set of measures would be a better predictor – that of the hand coordination measures, directly linked to playing the piano; or that of the foot coordination measures, which are linked to the control mechanism of the SR3T. Our results suggest that the set of foot coordination measures is a good predictor of performance with the SR3T (Fig. 5). The regression coefficients of the four measures are within the same range, suggesting a balanced contribution of these different measures. The model based on hand coordination measures does not perform as well in prediction. Furthermore, the contributions of the measures are unbalanced relative to how foot measures contributed to the foot-based model. Hand dimensionality which was the best metric to distinguish pianists from naïve subjects shows a minimal contribution to the model. Piano position and timing measures are showing reverse contributions, even though they are highly and significantly correlated (see Fig. 3).

Our results suggest that the human motor coordination skill in using the control interface of the robotic augmentation device (in our case, the foot) is the best predictor of how well the augmented human performs with the robotic system; this is confirmed through AICc in comparison with models arising from different combinations of features. Interestingly, skills otherwise relating to the actual task do not serve as good predictors either, i.e. in the case of piano playing, the hand-related motor coordination measures from HAMCA which the piano task heavily relies on are not good predictors of how well the human will perform, even though one of them (hand dimensionality) serves as the best predictor of piano playing expertise. Previous work on interfaces for supernumerary robotics have shown that the foot can generally serve as a good interface for robotic limbs working collaboratively with the user's hands [e.g. 23,56,57]. Abdi et al. [23] study the control of a third robotic hand via the foot in virtual reality, for robotic surgery applications, showing similar learning trends to what we observe here. We see similar effects where roboticists have used legs and feet as a multi-DOF control interface for successfully teleoperating two [58] or four [59] robotic arms, albeit in less skilled tasks than what we show here. Saraiji et. al. show that subjects significantly increased their self-reported sense of embodiment of the tele-operated robots over the course of the experiments, i.e. 40 minutes [58]. Results obtained with adaptive foot interfaces for robot control [e.g. 60], where data-driven approaches are used to create subject-specific motion mapping, are in line with our findings. Huang et al. [60] report that inter-subject variability decreases once a subject-specific motion mapping is enabled. This confirms the dependency of robot control performance on metrics inherent to each subject, which we present here to be the interface space motor coordination skills.

Our work shows the possibility of humans being able to quickly acquire a skilled behaviour, such as playing piano sequences, with a human augmentative robotic system. Both naïve piano players (i.e. without prior experience) and piano playing experts demonstrated the same ability to integrate the supernumerary robotic limb: We saw no difference in the performance with the SR3T, suggesting that

integrating robotic augmentation is primarily driven by a priori motor coordination skills and not affected significantly by expert motor domain knowledge. We are looking here at a setup where supernumerary robotic thumb is controlled by the foot, therefore leading to a transfer of skills across limbs. While this has not been systematically evaluated before, we can look at the expertise of hand use and its transference to the foot in the domain of handwriting, where the shape of handwriting is recognizably transferred from the hand to the foot, and other limbs [61–63]. Similarly, we were expecting people skilled at piano playing with the hand, would show similar levels of skill when controlling the piano with the foot. However, surprisingly, this is not the case in our results. This observation might be due to crudeness in our setup, leading to piano skills not being carried across. Nevertheless, our unconstrained experiment, and our systematic one both show participants capable of controlling the robotic thumb to play the piano and to achieve high scores. It could also be the case that the selected piano sequence was not complex enough for the transfer of skill to emerge, as it is a one handed, simple melody. We designed our piano piece to ensure comparability with the experimental setup, so that pre-augmentation the participants would play the main notes with their right hand and then play the additional notes with the left hand index finger, therefore limiting us to one-handed pieces. The structure of the music piece itself also needed to be simple enough so that piano-naïve participant could acquire it within a reasonable amount of time and not immediately fail.

It is important to consider the meaning of these results in the context of prosthetics, and human augmentation in general. Prosthetics replace a limb that was lost whereas with the SR3T, and with supernumerary robotics in general, the human is operating a new, additional limb – in our case a thumb. Our augmentation is done through substitution, i.e. we use an existing limb to operate an additional one. We show here that this makes the system reliant on human motor skills, specifically in controlling the interface-space. We also demonstrate the possibility of applying substitution across different levels of the biomechanical hierarchy. The foot, which in terms of the biomechanical hierarchy is equivalent of the entire hand, is used here as the interface-space controlling a thumb, which is further down the biomechanical hierarchy. These results sit at one end of the spectrum of solutions for controlling an augmentative device, which goes from substitution all the way to direct augmentation via higher level control, either brain-machine-interfacing or cognitive interfaces such as eye gaze decoding. We previously showed that the end-point of visual attention (where one looks) can control the spatial end-point of a robotic actuator with centimetre-level precision [8,64,65]. This direct control modality is more effective from a user perspective than voice or neuromuscular signals as a natural control interface [66]. We showed that such direct augmentation can be used to control a supernumerary robotic arm to draw or paint, freeing up the two natural arms to do other activities such as eating and drinking at the same time [13]. But such direct augmentation has to date not achieved augmenting fine motor skills such as playing the piano, as playing this instrument requires not just the execution of a note: it is not a simple button-press exercise, but requires fine grade expression of temporal and spatial motor coordination across robotic and natural fingers. We show that we can predict the degree to which subjects can integrate supernumerary limbs into their natural body movements, as a function of their basic motor skills. Thus, our work shows that we can achieve effective augmentation but also predict the capability of individuals to embody supernumerary robotic limbs in real-world tasks, which has impact for robotic augmentation from healthcare to agriculture and industrial assembly e.g. in the aerospace industry.

Materials and Methods

Experimental design. For our unconstrained pilot experiment, a right-handed piano player was selected to wear the SR3T and freely improvise. For our systematic follow-up experiments we developed a set of measurement protocols and behavioural biomarkers, the Human

Augmentation Motor Coordination Assessment (HAMCA), and ran this set of tests to assess hand and foot coordination (due to the foot being the control interface for our robotic system, described below under *Setup*) and piano-related skills. As opposed to existing motor assessments such as the Purdue pegboard [67], the motor domain of the NIH Toolbox [68], the Jebsen-Taylor hand function test [69] or the Action Research Arm Test [ARAT, 70] among others, which tend to be focused on dexterity, and are mainly used to quantify the extent or progress of motor disabilities, here we are interested in assessing specific human motor coordination aspects which relate to the interface-space (foot) and task-space (hand use over the piano) of our piano playing task. The HACMA set includes both spatial and temporal evaluations. From these coordination tasks we extracted 8 hand and foot motor-coordination scores. Finally, the participants were given specific melodies to play on the piano with and without the SR3T. The melody was designed to require 6 fingers, forcing the participant to either use their left-hand index finger (LHIF), or the SR3T if they are wearing it. Table 1 summarises the experimental setup procedure and how they map to results.

Table 1. Experimental procedure

First Session – 2 hours	Second session – 1 hour	
Foot Balance (15 trials) – 20 mins	Piano Playing with LHIF – 25 mins	
Foot Up-Down (15 trials) – 15 mins	5 Practice Trials	10 trials recorded at 80bpm (last 5 count for score)
Foot Tracking (6 trials) – 10 mins		
Piano Timing (25 trials) – 15 mins	SR3T setup and calibration – 10 mins	
Piano Positioning (15 trials) – 10 mins	Piano Playing with SR3T – 25 mins	
Piano Loudness (25 trials) – 15 mins	5 Practice Trials	10 trials recorded at 80bpm (last 5 count for score)
Hand Dimensionality (2x toy tasks) – 35 mins		

Subjects. Twelve right-handed participants took part in our systematic experiments (mean age 23.3+/-2.8 years). Six of the participants had played the piano for several years (pianists group) and the other six did not have any piano playing experience (naïve group). All of the participants from the pianists group had at least 5 years of piano training (range 5-21 years, mean 10.6+/-5.4 years). Two participants of the naïve group had over 5 years' experience of

guitar playing. All participants gave informed consent prior to participating in the study and
all experimental procedures were approved by Imperial College Research Ethics Committee
and performed in accordance with the declaration of Helsinki.

Setup. We created an experimental setup to investigate how individual motor skills
contribute to the performance of a human user of a supernumerary robotic thumb; i.e. a
robotic augmented human. To this end, we have created a 2 degrees of freedom (DoF) robotic
finger that users can wear on the side of their hand, effectively augmenting them with a third
thumb. The design, creation and initial testing of the supernumerary robotic third thumb
(SR3T) was reported in [48], and is the same setup used for our unconstrained pilot experiment
with a single piano player participant. The design specifications for the SR3T were derived
from the design requirements of a fully spherical operating thumb [71] and the natural
eigenmotions of human thumbs in daily life activities [72]. The SR3T is attached to the user's
right hand and is controlled through the user's right foot. In our original implementation, used
for the pilot experiment, the vertical motion of the foot was measured using an
accelerometer, together with horizontal motion data obtained with a flex sensor worn on the
natural thumb on the augmented hand, to control the vertical and horizontal DoFs of the
SR3T, respectively [48]. For our main experiments, we updated the interface, using a 9DoF
inertial measurement unit (IMU - Bosch BNO055, breakout board by Adafruit) for increased
stability, and to limit the interface to a single limb, i.e. the foot. The unit can provide absolute
orientation measurements (with respect to the earth's magnetic field) thanks to an onboard
sensor fusion algorithm. Absolute orientation can then be extracted as Euler vectors, at
100Hz. In this setup, the SR3T's two DoFs correspond to horizontal and vertical movements
of the robotic fingertip. These are mapped to horizontal and vertical movements of the foot,
i.e. yaw and pitch, respectively. Once the subject is wearing the SR3T on their hand, and the
IMU on their foot, they are asked to sit with their foot on the ground facing the piano. The
SR3T is moved horizontally for the fingertip to face the forward position as well. The values
read by the IMU for the orientation of the foot, and by the motor encoders for the position
of the SR3T are recorded. The subject is then asked to rotate their foot clockwise, with the
heel as the centre of rotation, to their maximum comfortable reach (typically 45 degrees from
the forward-facing pose). The SR3T fingertip is also moved accordingly, to the maximum
horizontal position on the right, and values recorded as before. These are used to map the
horizontal motion of the foot to that of the SR3T, with a similar process for vertical motions.
The setup can be seen in Fig. 1.

In order to investigate the workspace augmentation achieved by the SR3T optical markers
were placed at the tip and base of the SR3T, with the SR3T then activated by the subject to
move in its full range of motion. Similarly, optical markers were placed at the tip and base of
the subject's left-hand thumb with them performing the maximum range of thumb
movement while the motion was optically tracked. We used three OptiTrack Prime 13W
cameras with the Motive software for motion capture (NaturalPoint, Inc. DBA OptiTrack,
Oregon, USA). The results can be seen in Fig. 1.E and F; the thumbs' end-point surface is
mapped onto a sphere, assuming the base of the thumbs are situated at the centre. Based on
these measurements, the SR3T has an end-point work surface that is 4 times that of the

human thumb. Furthermore, we used the same camera system to measure delay between
motor intention and action, by placing markers on the user's foot as well as the SR3T; the
mean delay is measured as 85msec.

For the piano playing tasks and piano related hand coordination tasks we used a digital
piano (Roland RP501R-CB, Roland Corp., Osaka, Japan). The piano was connected to a PC with
a MATLAB script establishing communication through its MIDI interface. Each keystroke on
the piano was received by the MATLAB script as a MIDI message which comprised data
regarding the note played, time of keystroke (with a 1ms resolution) and the keystroke
velocity, which leads to proportional loudness of the note played.

HAMCA foot coordination tasks

Foot balance task

A Wii Balance Board (Nintendo Co. Ltd., Kyoto, Japan) together with the BrainBloX software [73] was
used. The board (Figure 2A) is made of four pressure plates and the software interface displays the
real-time centre of pressure computed by the Wii Balance Board across all four plates, and relative to
the board's coordinate system.

Weight plates (70 N) were placed on the left side of the board, moving the centre of pressure away
from the system origin. Subjects then had to move the centre of pressure back towards the origin by
applying pressure on the right side of the board with their right foot. The plates were placed in three
positions (Figure 2A), with five trials per position, resulting in a total of 15 trials, performed in random
order. Before the beginning of each trial, participants were asked to place the centre of pressure as
close as possible to the origin. Once they stated their readiness and after a 3-seconds countdown, a
15-seconds recording was started. Samples were recorded at 85 Hz. The resulting motor-coordination
score is computed according to equation (1):

$$Accuracy = 1 - \frac{error}{maxError} \quad (1)$$

Where error corresponds to the mean Euclidean distance of the centre of pressure from the origin
of the coordinate system across all recorded samples. The maximum error corresponds to the error
computed if the subject was not acting on the platform.

Foot up-down task

The same setup as in the Foot Balance Task was used, without the weights. A steady beat was
played with a metronome, which the subjects had to match when moving their feet from a toe-lifted
(dorsiflexion) to a heel-lifted (plantarflexion) pose and vice versa (see Figure 2C). The pressure exerted
by the foot had to match an upper and lower target value marked on the screen. Ideally, the output
should resemble a square signal with a period equivalent to that of the beat on the metronome.
Subjects performed 15 trials in random order, five at each selected tempi: 40bpm, 60bpm and 80bpm.

Two types of motor-coordination scores are computed from this task: spatial and temporal, both
using equation (1). For the spatial measure, the error is calculated as the absolute distance between
the target pressure position and the measured position of the centre of pressure. The maximum error
corresponds to the total distance between the upper and lower pressure targets. The temporal
measure's error is based on how precise in time the change between target positions occurs. This is

specifically measured at the time of zero-crossing, respective to the beats of the metronome.
Maximum error is the time corresponding to one full period. Both the spatial and temporal absolute
errors had skewed distributions; therefore, the median of the error was utilised instead of the mean.

Foot tracking task

The subjects controlled the 2D position of a dot on a screen through rotations of their ankle,
captured with an inertial measurement unit (IMU) attached to their shoe (see Figure 2B) -- the same
setup used as the control interface of the SR3T. The subjects were directed to use ankle rotations only,
the result of which they could see as a blue dot on a screen. They had to make the blue dot follow the
position of a red dot moving along a figure-of-eight path, as shown in Figure 2B. The path, the red and
blue dots were shown to subjects on a monitor screen in front of them. Each trial is composed of 6
laps around the figure-of-eight path, lasting 35 seconds total. The subjects sat at a height to have their
foot freely moving in space (see Figure 2B). The motor-coordination score for this task follows
equation (1), with the error defined as the Euclidean distance between the blue and red dots. The
maximum error is taken as the maximum recorded error across all time points in all trials of all
subjects. Once again, due to the skewness in the absolute error distribution, the median of the error
was used in the accuracy calculation.

HAMCA hand coordination tasks

Hand dimensionality

Subjects performed two toy assembly tasks while wearing a Cyberglove II (CyberGlove Systems LLC,
San Jose, CA) to capture the motion of their hand and fingers, with 22 degrees of freedom. The tasks
involved assembling a LEGO DUPLO toy train (LEGO 10874), and assembling a toy car (Take Apart, F1
Racing Car Kit) using a toy drill and screws (see Figure 2, D and E). To ensure the appropriate fit of
Cyberglove II we made sure all participants had a minimum hand length of 18 cm, measured from the
wrist to the tip of the middle finger.

Principal Component Analysis (PCA) was performed on the collected data. We relate a greater
number of principal components needed to explain the variance of the motion, to greater hand
coordination [74]. The resulting motor-coordination score is defined as the number of principal
components required to explain 99% of the recorded motion's variance, normalised by the number of
degrees of freedom recorded: 22.

Piano timing

The subjects used their right-hand index finger to press the same piano key at varying tempi
(40bpm, 60bpm, 80bpm, 100bpm and 120bpm) played by a metronome. In total, subjects performed
25 trials in random order (5 at each tempo) composed of 10 keystrokes.

The relevant motor-coordination score follows the same concept as that of equation (1); for further
clarity we present it in more detail, in equation (2). The normalised error is the absolute time deviation
from the correct tempo divided by its period; that is, the time between keystrokes (inter-onset
intervals or IOI) minus the period of each tempo in seconds, as shown in equation (2).

$$TimingAccuracy = 1 - \frac{|IOI - (60/tempo_{bpm})|}{60/tempo_{bpm}} \quad (2)$$

Where tempo is the beats per minute value, hence making 60/tempo the beat period in seconds. Nine samples were generated in each trial (given that nine IOI are generated by ten keystrokes); hence, there were 45 samples generated at each tempo, which had a skewed distribution. The median of these values was taken as the score at each tempo and then the five tempi's scores were averaged to obtain a single value for their motor-coordination score in the task.

Piano positioning

The right-hand index finger was used to move back and forth between two keys and press them at a rate given by the metronome (fixed tempo of 60bpm). Three piano keys were selected, one positioned in the middle of the piano and the other two spaced 7 whole notes to the left and right of it. Three combinations of two keys were to be followed: left and middle, middle and right, left and right (see Figure 2F). In total, subjects performed 15 trials in random order (5 at each key combination) composed of 12 keystrokes each. The relevant motor-coordination score is defined the same way as in the piano timing task. Timings are measured between two consecutive and correct key presses - timings relating to incorrect keypresses were discarded. The latter is done automatically as the note values will be different to what is expected. In order to make up for cases where participants might have pressed the incorrect key, or missed a beat, we consider a window of size of the tempo period (1 second) centred on the correct beat time. If notes are played outside of this window, we assume that the first keystroke of the interval is a wrong one. As the incorrect notes are already removed, a time before the window would mean that the same key was pressed twice consecutively and a time after it would mean that a keystroke was missed. Most of the subjects had no misses or 1 miss out of 165 samples.

Piano loudness

On the digital piano, the loudness of a note depends on the velocity with which the relevant key is pressed. A fast press will produce a louder sound and vice versa. Subjects were instructed to press a single key at a target level of loudness, with both the target level and the level at which they pressed shown to them visually. Before starting the experiment, participants were instructed on how to set their minimum (0%) and maximum (100%) keystroke loudness values. The piano's recorded loudness values range from 0 to 127. A very slow key press corresponds to values around 2-8, whereas fast presses fall within values of 120-127. Participants were allowed to familiarise themselves with the visual interface by the experiment runner doing one block of trials on themselves, with the participant watching the interface. Then, they are given up to 5 unrecorded trials to familiarise themselves with how the key presses relate to numerical values, and for the experiment runner to ensure that they cover the full range of values in their key presses. They are then asked each to define their own range, by pressing the key at 0% and 100%. These values are recorded and used to define their range for the experiment. The loudness values for levels 25%, 50% and 75% are obtained by linear interpolation between the 0% and 100% values defined for each participant.

In total, subjects performed 25 trials, 5 at each loudness level (randomised): 0%, 25%, 50%, 75% and 100%, composed of 10 keystrokes each. The motor-coordination score is calculated following equation (1), with the error defined as the deviation from the target values (in percentage loudness) and maximum error as the maximum committed among all the trials of all of the participants for each loudness level (these are as follows, Level 0: 34.0833, Level 25: 42.6250 , Level 50: 34.6667, Level 75: 28.4583 and Level 100: 24.4167). After analysing the results, the average motor-coordination score was calculated using only the results at the 25%, 50% and 75% loudness levels given that these targets

562   required more skilled velocity control than the 0% and 100% levels. Thus, their use would enhance
563   individual differences between participants.

Piano playing

To assess the participants' performance on actual piano playing, a sequence with 38 notes played
at a constant tempo (isochronous) of 80bpm was devised. Subjects were able to learn and follow the
sequence while playing, aided by the software Synthesia (Synthesia LLC). Synthesia showed the notes
of the sequence as coloured blocks scrolling on-screen. The participants had to press the keys
corresponding to the positions on the keyboard with which the Synthesia blocks were aligned in time
to the music in order to score points (see Figure 1). The sequence of notes was designed to be played
mainly with the right hand, plus one finger for notes that were too far to the right side of the right
hand. These notes could then be reached either using the index finger of the left hand, or, if wearing
the SR3T, by activating the robotic finger. On Synthesia, the notes to be played by the right-hand
fingers were coloured in green, and the notes to be reached with the extra finger were coloured blue.
Similarly, the relevant keys on the keyboard were marked with the same colours (see Figure 2F).

Subjects played the sequence first without and then with the SR3T for 15 trials in each block. The
first five trials were considered as practice trials (not recorded), the next ten were recorded but only
the last 5 are used for computing the mean piano playing scores per subject due to the fact that
subjects were still learning the sequence, especially the ones with no piano playing experience. For
the first block of trials, without the SR3T, subjects played using their right hand for green coloured
notes while blue coloured ones were played with the left-hand index finger. To achieve this, subjects
had to cross their left hand over the right one. For the second block of trials, the left index finger was
replaced by the SR3T. We score each individual keypress's timing as follows:

$$Score = 1 - \frac{\Delta T}{IOI/2} \quad (3)$$

where ΔT is the absolute time difference between the keypress and metronome beat, and IOI is
the corresponding beat time period. Therefore, the participants receive a full score for each correct
keypress at the exact correct time, with the score linearly decreasing for time deviations, up to half
the beat period on each side, at which point the score is 0. Incorrect notes within this window are
obviously marked as 0 too. We then average the note scores for the entire sequence.

In some trials, recordings were stopped prematurely due to technical errors. In all such cases, the
score is calculated with respect to the recorded section only. However, if less than 50% of the notes
are recorded, then the trial is discarded. This happened only in two trials from the same subject which
were removed. No other trials for any subjects had any missed recordings. There were also cases
where participants missed one initial beat, leading to them being off-beat for the entire sequence. To
adjust for this, we calculate the scores for the sequence as originally timed, plus if it were started one
beat early, or one beat late. We then take the highest score of the three cases to represent the piano
playing score for that trial. This only occurred twice.

Statistical Analysis

599   We first tested for statistically significant differences between the pianists and the naïve players in
600   each of the HAMCA motor-coordination scores, using student t test. We then calculated the
601   correlation matrix between the motor-coordination scores of all subjects, looking for dependencies
602   between the HAMCA tasks and scores.

For the analysis of the piano playing performance with the LHIF and with the SR3T we addressed trials 5 to 10, after the initial fast learning plateau. Using t test, we looked for significant differences between the pianists and the naïve players in the scores of the trials played with their LHIF and with the SR3T. We then averaged over trials 5 to 10 to get a single piano playing score for each subject in each task to be used in all following analysis. We also merged the two groups (pianists and the naïve players) so that all test of interactions between the HAMCA motor-coordination scores and the piano playing scores were across all subjects. We then calculated the correlations between the subjects' scores in the piano playing tasks and the HAMCA motor-coordination scores. To account for an outlier subject, we further investigated the correlations with Spearman rank correlation scores.

We then fitted generalized linear models to explain the piano playing scores using two different sets of HAMCA motor-coordination scores: the HAMCA hand measures and the HAMCA foot measures. We first removed the outlier subject before fitting the models (in the main text), and later repeated the analysis with the outlier (in the supplementary figures). We then tested for the contribution of piano expertise and the LHIF playing score to the prediction by adding them to the model (each of them separately and both together). To compare between these models of different complexity we used the corrected Akaike information criterion (AICc) which is modified for small sample sizes.

References

1. Reiter, R. Eine neue elektrokunsthand. *Grenzgeb. Med.* **1**, 133–135 (1948).

2. Zollo, L., Roccella, S., Guglielmelli, E., Carrozza, M. C. & Dario, P. Biomechatronic design and control of an anthropomorphic artificial hand for prosthetic and robotic applications. *IEEE/ASME Trans. Mechatronics* **12**, 418–429 (2007).

3. Farina, D. *et al.* The extraction of neural information from the surface EMG for the control of upper-limb prostheses: emerging avenues and challenges. *IEEE Trans. Neural Syst. Rehabil. Eng.* **22**, 797–809 (2014).

4. Bensmaia, S. J. Biological and bionic hands: natural neural coding and artificial perception. *Philos. Trans. R. Soc. B Biol. Sci.* **370**, 20140209 (2015).

5. Godfrey, S. B. *et al.* The SoftHand Pro: Functional evaluation of a novel, flexible, and robust myoelectric prosthesis. *PLoS One* **13**, (2018).

6. Aszmann, O. C. *et al.* Elective amputation and bionic substitution restore functional hand use after critical soft tissue injuries. *Sci. Rep.* **6**, (2016).

7. Benabid, A. L. *et al.* An exoskeleton controlled by an epidural wireless brain–machine interface in a tetraplegic patient: a proof-of-concept demonstration. *Lancet Neurol.* **18**, 1112–1122 (2019).

8. Shafti, A., Orlov, P. & Faisal, A. A. Gaze-based, context-aware robotic system for assisted reaching and grasping. in *Proceedings - IEEE International Conference on Robotics and Automation* vols 2019-May 863–869 (Institute of Electrical and Electronics Engineers Inc., 2019).

9. Hussain, I., Salvietti, G., Spagnoletti, G. & Prattichizzo, D. The Soft-SixthFinger: A Wearable EMG Controlled Robotic Extra-Finger for Grasp Compensation in Chronic Stroke Patients. *IEEE Robot. Autom. Lett.* **1**, 1000–1006 (2016).

10. Haddadin, S. & Croft, E. Physical human-robot interaction. in *Springer Handbook of Robotics*

645   1835–1874 (Springer International Publishing, 2016). doi:10.1007/978-3-319-32552-1_69.

646 11. Maurtua, I. *et al.* FourByThree: Imagine humans and robots working hand in hand. in *IEEE*
647   *International Conference on Emerging Technologies and Factory Automation, ETFA* vols 2016-
648   November (Institute of Electrical and Electronics Engineers Inc., 2016).

649 12. Shafti, A. *et al.* Real-time robot-assisted ergonomics. in *2019 International Conference on*
650   *Robotics and Automation (ICRA)* 1975–1981 (IEEE, 2019).

651 13. Dziemian, S., Abbott, W. W. & Faisal, A. A. Gaze-based teleprosthetic enables intuitive
652   continuous control of complex robot arm use: Writing & drawing. *Proc. IEEE RAS EMBS Int.*
653   *Conf. Biomed. Robot. Biomechatronics* **2016**-**July**, 1277–1282 (2016).

654 14. Parietti, F. & Asada, H. Supernumerary Robotic Limbs for Human Body Support. *IEEE Trans.*
655   *Robot.* **32**, 301–311 (2016).

656 15. Makin, T. R., De Vignemont, F. & Faisal, A. A. Neurocognitive barriers to the embodiment of
657   technology. *Nat. Biomed. Eng.* **1**, 1–3 (2017).

658 16. Llorens-Bonilla, B., Parietti, F. & Asada, H. H. Demonstration-based control of supernumerary
659   robotic limbs. in *IEEE International Conference on Intelligent Robots and Systems* 3936–3942
660   (2012). doi:10.1109/IROS.2012.6386055.

661 17. Wu, F. & Asada, H. Supernumerary robotic fingers: An alternative Upper-limb prosthesis. in
662   *ASME 2014 Dynamic Systems and Control Conference, DSCC 2014* vol. 2 (American Society of
663   Mechanical Engineers, 2014).

664 18. Hussain, I. *et al.* Design and prototype of supernumerary robotic finger (SRF) inspired by fin
665   ray® effect for patients suffering from sensorimotor hand impairment. in *RoboSoft 2019 - 2019*
666   *IEEE International Conference on Soft Robotics* 398–403 (Institute of Electrical and Electronics
667   Engineers Inc., 2019). doi:10.1109/ROBOSOFT.2019.8722748.

668 19. Tiziani, L. *et al.* Empirical characterization of modular variable stiffness inflatable structures for
669   supernumerary grasp-assist devices. *Int. J. Rob. Res.* **36**, 1391–1413 (2017).

670 20. Hussain, I., Spagnoletti, G., Salvietti, G. & Prattichizzo, D. Toward wearable supernumerary
671   robotic fingers to compensate missing grasping abilities in hemiparetic upper limb. *Int. J. Rob.*
672   *Res.* **36**, 1414–1436 (2017).

673 21. Wu, F. & Asada, H. 'Hold-and-manipulate' with a single hand being assisted by wearable extra
674   fingers. in *Proceedings - IEEE International Conference on Robotics and Automation* vols 2015-
675   June 6205–6212 (Institute of Electrical and Electronics Engineers Inc., 2015).

676 22. Mehring, C. *et al.* Augmented manipulation ability in humans with six-fingered hands. *Nat.*
677   *Commun.* **10**, 2401 (2019).

678 23. Abdi, E., Burdet, E., Bouri, M. & Bleuler, H. Control of a supernumerary robotic hand by foot:
679   An experimental study in virtual reality. *PLoS One* **10**, (2015).

680 24. Parietti, F. & Asada, H. Independent, voluntary control of extra robotic limbs. in *Proceedings -*
681   *IEEE International Conference on Robotics and Automation* 5954–5961 (Institute of Electrical
682   and Electronics Engineers Inc., 2017). doi:10.1109/ICRA.2017.7989702.

683 25. Liang, X., Cheong, H., Chui, C. K. & Yeow, C. H. A fabric-based wearable soft robotic limb. *J.*
684   *Mech. Robot.* **11**, (2019).

685 26. Al-Sada, M., Höglund, T., Khamis, M., Urbani, J. & Nakajima, T. Orochi: investigating
686   requirements and expectations for multipurpose daily used supernumerary robotic limbs. in

ACM International Conference Proceeding Series (Association for Computing Machinery, 2019).
doi:10.1145/3311823.3311850.

27. Santello, M. *et al.* Hand synergies: Integration of robotics and neuroscience for understanding
the control of biological and artificial hands. *Physics of Life Reviews* vol. 17 1–23 (2016).

28. Wu, F. & Asada, H. Implicit and Intuitive Grasp Posture Control for Wearable Robotic Fingers:
A Data-Driven Method Using Partial Least Squares. *IEEE Trans. Robot.* **32**, 176–186 (2016).

29. Valero-Cuevas, F. J. & Santello, M. On neuromechanical approaches for the study of biological
and robotic grasp and manipulation. *J. Neuroeng. Rehabil.* **14**, 101 (2017).

30. Bromberg, Z., Donchin, O. & Haar, S. Eye movements during visuomotor adaptation represent
only part of the explicit learning. *eNeuro* **6**, (2019).

31. Haar, S., Donchin, O. & Dinstein, I. Dissociating visual and motor directional selectivity using
visuomotor adaptation. *J. Neurosci.* **35**, 6813–6821 (2015).

32. Krakauer, J. W., Pine, Z. M., Ghilardi, M. F. & Ghez, C. Learning of visuomotor transformations
for vectorial planning of reaching trajectories. *J. Neurosci.* **20**, 8916–8924 (2000).

33. Taylor, J. A., Krakauer, J. W. & Ivry, R. B. Explicit and implicit contributions to learning in a
sensorimotor adaptation task. *J. Neurosci.* **34**, 3023–3032 (2014).

34. Telgen, S., Parvin, D. & Diedrichsen, J. Mirror reversal and visual rotation are learned and
consolidated via separate mechanisms: Recalibrating or learning de novo? *J. Neurosci.* **34**,
13768–13779 (2014).

35. Wilterson, S. A. & Taylor, J. A. Implicit visuomotor adaptation remains limited after several days
of training. *bioRxiv* 711598 (2019) doi:10.1101/711598.

36. Yang, C. S., Cowan, N. J. & Haith, A. M. De novo learning versus adaptation of continuous
control in a manual tracking task. *bioRxiv* 2020.01.15.906545 (2020)
doi:10.1101/2020.01.15.906545.

37. Van Der Vliet, R. *et al.* Individual differences in motor noise and adaptation rate are optimally
related. *eNeuro* **5**, (2018).

38. Wu, H. G., Miyamoto, Y. R., Castro, L. N. G., Ölveczky, B. P. & Smith, M. A. Temporal structure
of motor variability is dynamically regulated and predicts motor learning ability. *Nat. Neurosci.*
17, 312–321 (2014).

39. Haar, S., van Assel, C. M. & Faisal, A. A. Motor learning in real-world pool billiards. *Sci. Rep.* **10**,
20046 (2020).

40. Maselli, A. *et al.* Where Are You Throwing the Ball? I Better Watch Your Body, Not Just Your
Arm! *Front. Hum. Neurosci.* **11**, 505 (2017).

41. Haar, S. & Faisal, A. A. Brain Activity Reveals Multiple Motor-Learning Mechanisms in a Real-
World Task. *Front. Hum. Neurosci.* **14**, (2020).

42. Di Cagno, A. *et al.* Motor learning as young gymnast's talent indicator. *J. Sport. Sci. Med.* **13**,
767–773 (2014).

43. Johnston, K., Wattie, N., Schorer, J. & Baker, J. Talent Identification in Sport: A Systematic
Review. *Sports Medicine* vol. 48 97–109 (2018).

44. Lopes, V. P., Rodrigues, L. P., Maia, J. A. R. & Malina, R. M. Motor coordination as predictor of
physical activity in childhood. *Scand. J. Med. Sci. Sport.* **21**, 663–669 (2011).

45.   Vandorpe, B. *et al.* The value of a non-sport-specific motor test battery in predicting
729   performance in young female gymnasts. *J. Sports Sci.* **30**, 497–505 (2012).

730   46.   Olivier, G. N. *et al.* Predicting Motor Sequence Learning in People With Parkinson Disease. *J.*
731   *Neurol. Phys. Ther.* **43**, 33–41 (2019).

732   47.   Wadden, K. P. *et al.* Predicting Motor Sequence Learning in Individuals With Chronic Stroke.
733   *Neurorehabil. Neural Repair* **31**, 95–104 (2017).

734   48.   Cunningham, J., Hapsari, A., Guilleminot, P., Shafti, A. & Faisal, A. A. The Supernumerary
735   Robotic   3rd   Thumb   for   Skilled   Music   Tasks.   in   *BioRob*   (2018).
736   doi:10.1109/BIOROB.2018.8487609.

737   49.   Bernstein, N. A. The co-ordination and regulation of movements. (1967).

738   50.   Ivry, R. B. & Hazeltine, R. E. Perception and Production of Temporal Intervals Across a Range of
739   Durations: Evidence for a Common Timing Mechanism. *J. Exp. Psychol. Hum. Percept. Perform.*
740   **21**, 3–18 (1995).

741   51.   Jantzen, K. J., Steinberg, F. L. & Kelso, J. A. S. Functional MRI reveals the existence of modality
742   and coordination-dependent timing networks. *Neuroimage* **25**, 1031–1042 (2005).

743   52.   Karni, A. *et al.* The acquisition of skilled motor performance: Fast and slow experience-driven
744   changes in primary motor cortex. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 861–868 (1998).

745   53.   Dayan, E. & Cohen, L. G. Neuroplasticity subserving motor skill learning. *Neuron* **72**, 443–454
746   (2011).

747   54.   Costa, R. M., Cohen, D. & Nicolelis, M. A. L. Differential corticostriatal plasticity during fast and
748   slow motor skill learning in mice. *Curr. Biol.* **14**, 1124–1134 (2004).

749   55.   Papale, A. E. & Hooks, B. M. Circuit changes in motor cortex during motor skill learning.
750   *Neuroscience* **368**, 283–297 (2018).

751   56.   Carrozza, M. C. *et al.* A wearable biomechatronic interface for controlling robots with voluntary
752   foot movements. *IEEE/ASME Trans. Mechatronics* **12**, 1–11 (2007).

753   57.   Dougherty, Z. & Winck, R. C. Evaluating the performance of foot control of a supernumerary
754   robotic limb. in *ASME 2019 Dynamic Systems and Control Conference, DSCC 2019* vol. 3
755   (American Society of Mechanical Engineers (ASME), 2019).

756   58.   Yamen Saraiji, M. H. D., Sasaki, T., Kunze, K., Minamizawa, K. & Inami, M. MetaArms: Body
757   remapping using feet-controlled artificial arms. in *UIST 2018 - Proceedings of the 31st Annual*
758   *ACM Symposium on User Interface Software and Technology* 65–74 (Association for Computing
759   Machinery, Inc, 2018). doi:10.1145/3242587.3242665.

760   59.   Sanchez, J. H. *et al.* Four-Arm Manipulation via Feet Interfaces. (2019).

761   60.   Huang, Y. *et al.* A subject-specific four-degree-of-freedom foot interface to control a surgical
762   robot. *IEEE/ASME Trans. Mechatronics* 1–1 (2020) doi:10.1109/tmech.2020.2964295.

763   61.   Raibert, M. H. Motor control and learning by the state space model. in (Massachusetts Institute
764   of Technology, 1977).

765   62.   Wright, C. E. Evaluating the special role of time in the control of handwriting *. *Acta Psychol.*
766   *(Amst).* **82**, 5–52 (1993).

767   63.   Castiello, U. & Stelmach, G. E. Generalized representation of handwriting: Evidence of effector
768   independence. *Acta Psychol. (Amst).* **82**, 53–68 (1993).

64. Maimon-Mor, R. O. *et al.* Towards free 3D end-point control for robotic-assisted human
reaching using binocular eye tracking. in *Rehabilitation Robotics (ICORR), 2017 International*
Conference on 1049–1054 (IEEE, 2017).

65. Tostado, P. M., Abbott, W. W. & Faisal, A. A. 3D gaze cursor: Continuous calibration and end-
point grasp control of robotic actuators. *Proc. - IEEE Int. Conf. Robot. Autom.* **2016**-**June**, 3295–
3300 (2016).

66. Noronha, B., Dziemian, S., Zito, G. A., Konnaris, C. & Faisal, A. A. "Wink to grasp"—comparing
eye, voice & EMG gesture control of grasp with soft-robotic gloves. in *Rehabilitation Robotics*
(ICORR), 2017 International Conference on 1043–1048 (IEEE, 2017).

67. Tiffin, J. & Asher, E. J. The Purdue Pegboard: norms and studies of reliability and validity. *J.*
Appl. Psychol. **32**, 234–247 (1948).

68. Reuben, D. B. *et al.* Motor assessment using the NIH Toolbox. *Neurology* **80**, S65–S75 (2013).

69. Jebsen, R. H., Taylor, N., Trieschmann, R. B., Trotter, M. J. & Howard, L. A. An objective and
standardized test of hand function. *Arch. Phys. Med. Rehabil.* **50**, 311–319 (1969).

70. Lyle, R. C. A performance test for assessment of upper limb function in physical rehabilitation
treatment and research. *Int. J. Rehabil. Res.* **4**, (1981).

71. Konnaris, C., Gavriel, C., Thomik, A. A. C. & Faisal, A. A. Ethohand: A dexterous robotic hand
with ball-joint thumb enables complex in-hand object manipulation. in *Biomedical Robotics*
and Biomechatronics (BioRob), 2016 6th IEEE International Conference on 1154–1159 (IEEE,
2016).

72. Konnaris, C., Thomik, A. A. C. & Faisal, A. A. Sparse eigenmotions derived from daily life
kinematics implemented on a dextrous robotic hand. in *2016 6th IEEE International Conference*
on Biomedical Robotics and Biomechatronics (BioRob) 1358–1363 (IEEE, 2016).

73. Cooper, J., Siegfried, K. & Ahmed, A. BrainBLoX: Brain and Biomechanics Lab in a Box Software.
(2014).

74. Belic, J. J. & Faisal, A. A. Decoding of human hand actions to handle missing limbs in
neuroprosthetics. *Front. Comput. Neurosci.* **9**, 1–11 (2015).

Acknowledgments:
The authors thank James Cunningham and Anita Hapsari for their contribution to the SR3T platform
(see Cunningham et al., 2018).
799

800 **Funding:** This research was supported by eNHANCE (http://www.enhance-motion.eu) under the
801 European Union's Horizon 2020 research and innovation programme grant agreement No. 644000.
802

803 **Author contributions:** AS, PG and AAF did the initial conception of this work; AS, PG, RM and AAF
804 developed the experimental protocol and hardware/software requirements for it; RM ran the
805 experiments; AS, SH, RM, AAF analysed and interpreted the data; AS, SH wrote the manuscript; AS, SH
806 and AAF revised the manuscript.
807

808 **Additional information:** AS, SH, RM and PG declare no competing financial interests. AAF has
809 consulted for Airbus.
810

811 **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present
812 in the paper and/or the Supplementary Materials. Our full dataset will be made available on FigShare
813 upon publication. We can also make these available to reviewers upon request.

Supplementary Materials

Fig. S1. Correlations between accuracies. The first eight panels shows correlations between accuracies in piano playing with the SR3T and in the motor coordination tasks. The ninth panel shows correlations between accuracies in piano playing with and without the SR3T. Naïve subjects marked as grey and experienced subjects as black dots.

Fig. S2. Correlations between accuracies. *C*orrelations between accuracies in piano playing with the LHIF and in the motor coordination tasks.

Fig. S3. Model predictions for SR3T piano playing. N=11 subjects after outlier removal. Naïve subjects marked as grey and experienced subjects as black dots. **(A-B)** model from the 4 hand **(A)** or foot **(B)** dexterity measures and the LHIF piano playing score. **(C-D)** model from the 4 hand **(C)** or foot **(D)** dexterity measures and the expertise group (experienced vs naïve). **(E-F)** model from the 4 hand **(E)** or foot **(F)** dexterity measures and the LHIF piano playing score and the expertise group.

Fig. S4. Model predictions for SR3T piano playing.
N=12 subjects. Naïve subjects marked as grey and experienced subjects as black dots. **(A)** model from the 4 HAMCA hand dexterity measures. **(B)** model from the 4 HAMCA foot dexterity measures.