SARS-CoV-2 mutations and where to find them: An *in silico*

perspective of structural changes and antigenicity of the Spike protein

Ricardo Lemes Gonçalves [1,4], Túlio César Rodrigues Leite[1,4], Bruna de Paula Dias[1,4],

Camila Carla da Silva Caetano[1,5], Ana Clara Gomes de Souza[1], Ubiratan da Silva

Batista[1], Camila Cavadas Barbosa[1,5], Arturo Reyes-Sandoval[3], Luiz Felipe Leomil

Coelho[2], Breno de Mello Silva[1,4,5].

(1) Laboratório de Biologia e Tecnologia de Micro-organismos, Departamento de
Ciências Biológicas, Universidade Federal de Ouro Preto, Brazil.
(2) Laboratório de Vacinas, Departamento de Microbiologia e Imunologia,
Universidade Federal de Alfenas, Brazil.
(3) The Jenner Institute, Nuffield Department of Medicine, University of Oxford,
Oxford OX1 2JD, UK
(4) Programa de pós-graduação em Biotecnologia, Universidade Federal de Ouro
Preto, Brazil
(5) Programa de pós-graduação em Ciências Biológicas, Universidade Federal de
Ouro Preto, Brazil

Corresponding author: Professor Breno de Mello Silva, Universidade Federal de Ouro

Preto, Departamento de Ciências Biológicas, Universidade Federal de Ouro Preto, Ouro

Preto, Minas Gerais, 35.400-000, Brazil. Tel: +553135591259. E-mail:

breno@ufop.edu.br

SARS-CoV-2 mutations and where to find them: An *in silico*

perspective of structural changes and antigenicity of the Spike protein

The recent emergence of a novel coronavirus (SARS-CoV-2) is causing a severe
global health threat characterized by severe acute respiratory syndrome (Covid-
19). At the moment, there is no specific treatment for this disease, and vaccines
are still under development. The structural protein Spike is essential for virus
infection and has been used as the main target for vaccine and serological
diagnosis test development. We analysed 2363 sequences of the Spike protein
from SARS-CoV-2 isolates and identified variability in 44 amino acid residues
and their worldwide distribution in all continents. We used the three-dimensional
structure of the homo-trimer model to predict conformational epitopes of B-cell,
and sequence of Spike protein Wuhan-Hu-1 to predict linear epitopes of T-
Cytotoxic and T-Helper cells. We identified 45 epitopes with amino acid
variations. Finally, we showed the distribution of mutations within the epitopes.
Our findings can help researches to identify more efficient strategies for the
development of vaccines, therapies, and serological diagnostic tests based on the
Spike protein of Sars-Cov-2.

Keywords: SARS-CoV-2, Spike protein, Mutations, T-cell epitopes, B-cell
epitopes.

1 Introduction

The SARS-CoV-2, a novel coronavirus, is highly transmissible, leading to high

infection rates and human mortality around the world, turning the disease caused by this

virus (COVID-2019) a huge public health concern (Li et al., 2020). Up to now, SARS-

CoV-2 has been spreading in several continents and causing more than 4,629,000

confirmed cases, with mortality rates of 6,74% (World Health Organization, 2020 –

access at 18/05/2020).

Coronaviruses have the largest genome among RNA viruses (26 to 32 kilobases in

length) with 14 ORFs encoding 27 proteins. At 5' end of the genome, there are the 1ab

and 1a ORFs, which encode 16 mature non-structural proteins (nsp1 to nsp16). These

proteins play crucial functions during viral RNA replication and transcription[1]. At 3'

end of the genome, there are genes encoding four structural proteins: Spike (S),

Envelope (E), Membrane (M), and Nucleoprotein (N) as well as genes for 8 accessory

proteins named 3a, 3b, p6, 7a, 8b, 9b, and orf14[2].

Although recent findings suggest that SARS-CoV-2 has stronger transmissibility when

compared to SARS-CoV, the molecular mechanisms responsible for this difference

remain unclear[3]. However, among the structural proteins, the Spike glycoprotein that is

present as a homo-trimer on the coronaviruses surface, has been pointed as the most

important factor responsible for this stronger transmissibility since this protein is able to

bind cell receptors. The S protein has two subunits named S1 and S2. The S1 subunit is

responsible for binding to the host cell receptor. It has a signal peptide, an N-terminal

domain (NTD) and a receptor-binding domain (RBD). The S2 subunit is responsible for

fusion of the viral and cellular membranes and consists of a conserved fusion peptide

(FP), two hepta-repeats (HR1 and HR2), a transmembrane (TM) and a cytoplasmic (CP)

domains[4]. Structural analysis of the receptor-binding domain (RBD) of SARS-CoV-2

Spike protein and the human receptor angiotensin-converting enzyme 2 (ACE2) revealed that the RBD can induce the spillover to other animals as well as human-to-human transmissions[5,6]. This protein has been shown as a key factor for coronaviruses entry into cells, as well as a target for neutralizing antibodies, due to its role in binding to cellular receptors and fusion of viral and cellular membranes [4,7,8].

The RBD of SARS-CoV S1 domain undergoes conformational changes that hide or expose the determinants of receptor binding[6,9,10]. Then, upon RBD binding to cellular receptors, the Spike protein is cleaved by proteases and the signal peptide is released. This cleavage triggers conformational changes in the S1 and S2 subunits, leading to the exposure of the fusion loop and its interaction into target cell membrane. This fact turns this domain a target to virus neutralization by monoclonal antibodies. Thus, conformational changes of the Spike protein are a necessary step to viral membrane fusion, and it allows the entry of viral nucleocapsids into the host cell to initiate replication[7,9,11,12]. Additionally, the S glycoprotein of SARS-CoV-2 has a furin-like cleavage site at the S1/S2subunits, which highlights the essential differences in Spike protein between SARS-CoV and SARS-CoV-2. Thus, this mechanism is proving to be a potential target for vaccines, therapeutic approaches, and diagnosis for coronaviruses[13,14].

Since the Spike protein has a crucial role in the initial steps of SARS-CoV-2 replication, research studies have been focusing on its structure, function, and antigenicity to gain a better understanding ofthe Spike protein[14,15]. A bioinformatics analysis has shown that the S2 subunit of SARS-CoV-2 is highly conserved and shares 99% identity with those of the two bat SARS-like CoVs (SL-CoV ZXC21 and ZC45) and human SARS-CoV[8]. In addition, the tri-dimensional model of Spike protein structure has recently been

published, showing that the SARS-CoV-2 Spike protein has more affinity for binding

ACE2 than the SARS-CoV Spike protein[16].

However, some studies have shown that cellular receptors used to viral attachment and

entry can vary among host species of different coronaviruses [9,14,17,18]. A recent study

reported that SARS-CoV-2 is most closely related to the bat SARS-CoV RaTG13 that

forms a distinct lineage of SARS-CoVs, and their Spike glycoproteins share 98% amino

acid sequence identity[19]. Despite the amino acid level appears to be similar, there are

important differences in Spike protein between these viruses that might explain some

viral differences regarding the pathogenicity [14]. Therefore, more studies are needed to

understand how those differences can change SARS-CoV-2 functionality.

Data about the genomic variability of the SARS-CoV-2, especially on the region

encoding the Spike protein, could provide important support for the accuracy of

structural predictions. In this present study, we analysed 2363 sequences of the Spike

protein in order to study the frequency of mutations on the protein domains and motifs

and to identify potential epitopes on this protein. Additionally, we also determined the

epitope variability of the Spike protein circulating in all continents. This work can be

used to support long-term studies to identify Spike protein mutations emergence and

understand how it can affect vaccine trials and serological diagnosis.

1   Materials and methods

2   *Sequence retrieval and Structural analysis*

Only complete genomes of SARS-CoV-2 were collected in the GISAID

(https://www.gisaid.org/). Complete sequences of Spike protein of SARS-CoV-2 were

obtained from NCBI (https://www.ncbi.nlm.nih.gov/) and ViPR

(https://www.viprbrc.org). All sequences are aligned using MEGA-X[20] software with

MUSCLE algorithm[21].

The homo-trimer model was obtained through the Robetta server[22]

(http://robetta.bakerlab.org/) referenced by the partial crystal of the "closed-state" of

SARS-CoV-2 Spike protein6VXX[23], representing about 77% of the actual structure

(https://www.rcsb.org/structure/6VXX). Subsequently, the structure of model was

subjected to geometry optimization steps by the Discovery Studios software[24], where

was considered eleven outlier residues from the favorable/acceptable regions in the

Ramachandran plot: 59PHE[A,B,C], 62VAL[A,B,C], 365TYR[A,B,C], 544ASN[B] and 744GLY[B].

So that all residues with a radius distance of 4A from each of the outliers was also

considered in the optimization steps.

17   *T and B cell epitope prediction*

All obtained sequences were used to predict T and B cell epitopes. The Wuhan-Hu-1

Spike protein of reference sequence NC_045512.2[25]

(https://www.ncbi.nlm.nih.gov/protein/1796318598) was used as reference for the

epitope prediction. The NetCTL1.2[26] (http://www.cbs.dtu.dk/services/NetCTL/) was

used to predict MHC-I binding epitopes. The MHC class I was considered for the

prediction of epitopes for cytotoxic T cells through artificial neural networks, using the

standard set of Weight on C terminal cleavage score (0.15), Weight on TAP

(Transporter Associated with Antigen Processing) efficiency matrix (0.05) and

Threshold for epitope identification (0.75). NetMHCpan 4.0[27]

(http://www.cbs.dtu.dk/services/NetMHCpan/),was also used to predict MHC class I

epitopes for cytotoxic T cells. Peptides with mers of 8-11 was pointed through artificial

neural networks, with the threshold of <2% better in the binding score rank. The

NetMHCII 2.3[27] (http://www.cbs.dtu.dk/services/NetMHCII/), was used to prediction

epitopes for T Helper cells with 15 mers. The *loci*HLA-DR was used, with standard

threshold (<2% better in the binding rank affinity) to identify the peptides that best

bound to MHCII.

Linear epitopes for B cells of different sizes were predicted using BepiPred-2.0

(http://www.cbs.dtu.dk/services/BepiPred/)[28]. The standard threshold of 0.5 was used to

ensure the better Specificity/Sensitivity ratio of the epitope. Finally, the conformational

epitopes for B cells were predicted using the validated model of the three-dimensional

structure of the spike protein homo-trimer through two web servers (DiscoTope and

hroughElliPro). DiscoTope (http://tools.iedb.org/discotope/)[29] was used with the

threshold for specify epitope identification (-3.7). The prediction of conformational

epitopes using ThroughElliPro[30](http://tools.iedb.org/ellipro/) was made with the

maximum score threshold of 0.5 and the maximum distance for ligation of 6 angstroms.

Only those epitopes located in the "extracellular" region of the protein (13-1213) were

considered. The identified epitopes were visualized on Spike protein tri-dimensional

model using the software Visual Molecular Dynamics[31] (figures 1 to 3) or Discovery

Studios (supplementary figure 2) [32]. The image processing for figures 1 and 3 was done

using the software Visual Molecular Dynamics[31]. The flowchart of the three main stages

of the study is on figure 1.

Results

Structural modeling

The monomeric Spike protein model was represented according to their respective

domains, disulfide bridges, and glycosylation points indicated by Expasy's P0DTC2

annotations reports (https://viralzone.expasy.org/resources/Coronav/P0DTC2.txt)

(figure 2 and video 1 in supplemental material). The FP region, reported from Expasy's,

has a little difference than that is reported for Sars-Cov. The three-dimensional Spike

homo-trimer model showed a clash score of 1.13 and a molprobity

(http://molprobity.biochem.duke.edu/) score of 1.05 corresponding to the 99th and

100th percentiles, respectively, when compared to high-resolution 3D structures. The

model revealed 99.9% of its residues in acceptable/favorable regions on the

Ramachandram Plot (http://molprobity.biochem.duke.edu/), where only 59PHE residues

from chains A, B and C continue as an outlier (supplementary figure 1). After ensuring

the model quality, the homo-trimer 3D structure was used to the structural/antigenic

SARS-CoV-2 Spike protein variability mapping.

Variations of protein Spike

After Spike protein sequences retrieval and comparison, 856 amino acid variations were

identified in 32.8% of sequences. Europe and South America showed the highest

variation rates among worldwide sequences, showing variation on 47.4% and 44.1% of

sequences, respectively. (data not showed). Sequence comparison analysis identified 42

amino acid residues with variation that occurred at least twice on SARS-Cov-2

sequences. Twenty three variations was mapped in the S1 and 19 in S2 domain (table

1). Some of these variations (28) are represented by residues with non-homologous

physical-chemical characteristics in their respective continents of occurrence (figure 3).

Epitopes prediction of protein Spike

In the present analysis, it was predicted 282 epitopes: 95 for T-cytotoxic cells, 135 for

T-Helper cells and 52 for B cells (30 linear and 22 conformational epitopes). All

epitopes are shown in supplementary table 1. Only those variations whose showed

amino acid changes to residues with non-homologous physical-chemical characteristics

was considered and represented in table 2. As a result, there are 11 predicted epitopes

for T-Cytotoxic cells, 16 for T-Helper cells, 18 for B cells (10 linear and 8 structural

epitopes). Forty-five epitopes with mapped variations were represented in the 3D model

of the SARS-CoV-2 protein Spike structure (figure 4). The S1 domain gathers the

majority counts (33) while the NTD region showed 20 epitopes, of which 7 were

predicted for B cells (table 2).

Discussion

Amino acids variations

Our findings suggest that the majority of residue variations arose punctually on their

respective continents, with the exception of position 614, which occurs in all continents.

The 614-residue variation has the most expressive frequency and has been reported by

recent published/preprinted studies [33–35]. The H49Y variation has been identified in

Asia, Europe, and North America. In contrast, L5F, V367F, R765L, and S940F

variations, with the exception the last, were observed in at least two continents with

<1% frequency. (table 1).

Overall, the largest number of variants was found in S1 protein subunit, with 10 variant

positions found in Asia, 9 in Europe, 9 in North America, and 2 in Oceania (table 1).

South America and Africa did not show any other variation in the S1 domain besides

that identified in residue 614, probably due to the few numbers of analyzed genomes

from these regions. The sequences from Europe showed a higher number of variant

residues in the S2 domain (12) in comparison to the average from the other continents

(1.8). However, the S1 domain is the region more propense to appear new variations

due its most prominent variability between coronaviruses species[36].

Due to the short period since its dissemination, it is difficult to find a specific variation

pattern in Spike protein of SARS-Cov-2 on the continents. But even so, the variant

position D1259H in S2 domain was seen in North America sequences but not in Asia

and Europe (table 1). Future works with a higher number of high-quality sequences are

needed to better understand the possible heterogeneous distribution of South/Central

Americas, Africa and Oceania amino acid variations on these regions.

Mapping of variations

S1 DOMAIN

Our analysis showed that the NTD domain has many variations (figure 3), where

changes on teen residues were identified. However, only two residues variations showed

frequency above 1% (S50L and S247R). The S50L residue variation was found in

2.37% of sequences from Asia and is internally located in the homo-trimer structure. A

recent study demonstrated that the L50 residue found in SARS-Cov / SARS-like CoV's

was replaced by S50 in SARS-CoV-2 [2]. However, some SARS-Cov-2 sequences remain

with the L50 residue. Another particular residue variation (S247) showed 5.26%

frequency on Oceania. This residue is located at the peripheral end of the NTD portion

and was reported in a recent preprint study that five variations of Spike protein may be

related to variation of replication of SARS-CoV-2 in Vero-E6 cells[37].

In the RBD region, four residues variations were identified (table 1).A recent in silico

preprint study shows the variation V367F as a probable factor to the high increase of

binding affinity with the ACE2 receptor[40]. The V483A variation, found only in North

American sequences, are co-located in the binding region of the main residues involved

in the interaction of the SARS-Cov Spike protein with the ACE2 receptor [38,39].The

R408I variation was identified only in the Asian continent and it was reported in a

recent in silico preprint study as a probable factor of decreased binding affinity with the

ACE2 receptor[40] (figure 3-front & table 1). Additionally, the arginine at position 408 in

RBD core region has already been cited as an important point of interaction with N-

glycan, for both SARS and SARS-CoV-2[41,42]. Thus, future investigations should be

done in order to evaluate if these changes on Spike protein residues could be correlated

to functional changes.

A second G476S variation observed in the North American genomic sequences was

found in peripheral and accessible region of the structure (figure1-front) and is also co-

located with the region of the main interaction residues between the SARS-CoV-2

protein Spike and the ACE2 receptor [43]. This was also observed inSARS-CoV[39]. As

mentioned before, the highest frequency mutation rate was identified at position 614 in

almost all continents (except Central America). This variation could be used to group

the continents into either D or G predominantly variations. The group D includes Asia,

North America, and Oceania while group G includes Europe, South America, and

Africa (table 1). However, although residue D614 in SARS and SARS-CoV-2 reference

sequences indicates a positive selection of G614 mutation in those continents, there is

no evidence of how this mutation frequency could impact virus functionality.

Coronavirus SARS-CoV-2 sequencing analysis from diverse continents demonstrates

that the genes encoding the Spike proteins undergo diverse and frequent mutations[44].

First, it was identified a mutation (241C>T), which was developed gradually, and

reported with three other co-mutations (3037C>T, 14408C>T, and 23403A>G)[33]. These

mutations culminated on amino acid variations in Spike protein, nsp3, and RNA

primase, which are responsible for RNA replication[33]. Moreover, all associated-mutations observed were prevalent in Europe isolates, giving insights for SARS-CoV-2 severity in this region. Therefore, the SNP mutation (23403A>G/D614G) in Spike protein D614G, was pointed out by Yin, C.[33]. This residue is near to this furin recognition site (S1/S2 region) of SARS and could affect enzyme activity. However, the distance between alpha carbons of residues 614 and 685 turns the cleavage site for SARS-CoV-2 (S1/S2) greater than 35Å (supplementary figure 1), which makes it difficult to infer that this mutation might have a direct action on cleavage process on S1/S2 domains. Furthermore, a recent study showed possible changes in pathogenicity derived from mutations[37]. However, the D614G variation was not observed in the sequences analyzed by this study.

S2 DOMAIN

Five amino acid variations (T791I, D839Y, V1176F, C1254F, and P1263L) were found in the S2 domain with frequency >1%. These variations do not happen more than one continent. Located on the HR1 region periphery (figure 3 - side in), S940F is present in Europe (0.69% of sequences) and Oceania (3.51% of sequences). T791I that have 1,09% of frequency is occurs on FP region and it is present only in Asian sequences. The V1176 variation has been found in South America with considerable frequency (6.98% of sequences). Furthermore, C1254F and P1263L variations in the IC region at the cytoplasmic end structure (figure 3 - side in) appear in Oceania (3.51% of sequences) and Europe (1.97% of sequences), respectively. These latter verified variations in IC region are residues with non-homologous physical-chemical characteristics and they might carry changes on intra-cellular portion interaction with the cellular components as well as lipid bilayer. Thus, further studies must be performed to verify the influence of these changes on SARS-CoV-2 replication.

Epitope Variations Mapping

Our antigenic predictions show that the ELIPRO server identify the entire NTD domain

as an antigen (table 2 and figure 4). In this domain, seven variations of amino acid

residues were identified with non-homologous physical-chemical characteristics. The

remaining 19 epitopes in the NTD had one or two variations (table 2). In contrast, the

RBD domain showed only five epitopes (table 2). Recently the cryo-electron

microscopy structure of the SARS-CoV-2 protein Spike trimer was determined[8,16],

showing that RBD can undergo movements between its "up" or "down" conformations.

This suggest that the target epitope of the neutralizing antibody (CR3022) is accessible

when the RBD is in the "up" conformation. Additionally, the ACE2 host can interact

with the RBD when protein Spike is in the "up" conformation[45]. Consequently, the

RBD domain has been identified as the most promising target for vaccine prototypes

development[14]. However, up to now, little information about the natural variability of

SARS-CoV-2 residues involved in ACE2 binding has been elucidated in the literature.

Finally, the S2 domain presented 14 epitopes (three are shared with the S1 domain), two

of these S2's epitopes where found on the fusion peptide (FP) region, and another two

where located on the transmembrane (TM) region.

T cell epitopes

Among all epitopes identified, only $S^{610-620}$ and $S^{612-620}$ epitopes have amino acid

variation at high frequencies and worldwide prevalence (residue 614 in table1). These

two epitopes have been identified by two prediction methods used in this study for T-

cytotoxic cells. However, future studies are needed to find out if there is the importance

of this variation in the response to the SARS-CoV-2 infection.

Three more predicted epitopes for T-cytotoxic cells with variations (>1% frequency)

were identified in our analyzes, two at NTD (S^{50-58} in Asia and $S^{240-249}$ in Oceania) and a

third epitope $S^{786-794}$ in S2 domain on Asia (Highlighted in Tab 2). The prediction for T-

Helper cells epitopes identified seven epitopes with variations that haven >1%frequency

(Highlighted in Tab 2). The first T-Helper epitope S^{50-64} is found in Asia. The last six

epitopes are co-located among $S^{233-253}$ region in Oceania sequences (figure 4). This

analysis identified the overlapping of antigenic regions for T cells in the NTD domain,

but not in the RBD domain (figure 4).

Conversely, most of epitope predictions with variations >1% frequency was observed

on a single continent. This suggests the existence of a random Spike protein

diversification on these regions. However, the limited available literature of serological

responses diversity to SARS-CoV-2 does not allow to better clarify this context at this

time.

B cell epitopes

The total of six epitopes predicted for B cells was identified with variations at

frequencies> 1%. Three linear and three conformational (highlighted in table 2). Of

these, four epitopes are present in the RBD domain (only in North America), one linear

($S^{455-478}$), and three conformational ($S^{434-511}$, $S^{434-580}$, and $S^{434-584}$). The RBD polypeptide

chain is classically recognized for its potential to generate neutralizing antibodies to

SARS [39,46]and recently observed also being the target of a neutralizing response for

SARS-CoV-2 [38,47]. A recent in silico study highlighted the region 524-598, which is

partially present in the RBD, as one of the dominant epitopes for SARS and SARS-

CoV-2 sharing an 80% identity[15]. Finally, the last two linear epitopes identified by our

analysis are in domain S2 ($S^{786-800}$, $S^{828-843}$) with variations in Asia and Europe,

respectively. Furthermore, the fusion peptide structure has been shown as a target for

neutralizing immune response to SARS and SARS-CoV-2 [16]. Variations in the S^{786-}

^{800}epitope of FP region described in this study may also cause changes in its

antigenicity, but, as already mentioned, new studies should be done to verify this hypothesis.

The potential variations in physicochemical characteristics brought by a amino acid residue exchange can generate an eventual condition for the viral particle escaping from the immune system via a non-neutralizing cross-reactive response from previous infections. As example, previous mutational assays with the polypeptide chain of the SARS-Cov Spike protein have shown that changes in the physical-chemical characteristics of various residues have resulted in impaired functions[5,7,48].

Thus, there is a clear need to monitor the antigenic variation by the newly emerged coronavirus, as variations in the RBD domain may change over time, due to selective pressure on the virus through future therapies given to host populations. Although we did not identify variations in potential epitopes capable of inducing humoral response against SARS-CoV-2 with high frequencies and/or wide distribution on the continents, these slight variations covered by four months of SARS-CoV-2 spread just strengths the necessity to understand the potential of antigenic variation that the Spike protein might present. However, given the small variation in the epitopes pointed out in this work, we can suggest that vaccine approaches using the spike protein structure are unlikely to be impaired by the variability presented by theSARS-CoV-2. Still, it would be interesting to formulate therapies that focus on the S2 domain since it presented the lower number of variants. Moreover, among the S1 domain, RBD is the less affected by the variants, and therefore, more promising than NTD.

The great global engagement in tackling this pandemic has generated hundreds of vaccine prototypes and potential drugs in the short/medium term. However, prototype generation needs to rely on and be developed based on antigenic and structural variability studies, especially protein S, which is essential for virus/host interactions..

Ideally, drug and vaccine developments should take into account the virus entire

diversity of its population.. Thus, efforts should focus on a continuous study of the

genomic variations and their implications for the change in antigens, to guide the

production of next-generation vaccines and drugs effective against all strains of SARS-

CoV-2.

Acknowledgements

This work was supported by the Conselho Nacional de Desenvolvimento Científico e

Tecnológico under Grant 432611/2016-9. R.L.G., T.C.R.L., B.P.D., C.C.S.C. and

C.C.B. received fellowships from Coordenação de Aperfeiçoamento de Pessoal de

Nível Superior, Brasil (CAPES). A.C.G.S. received fellowships from Conselho

Nacional de Desenvolvimento Científico e Tecnológico (CNPq - PIBIC).

Declaration of interest statement

The authors declare that there is no conflict of interest.

Author contributions

Ricardo Lemes Gonçalves: Conception of the methodology, data collection, processing,

analysis and writing. Túlio César Rodrigues Leite: Data collection, processing and

writing. Camila Carla da Silva Caetano: Data analysis and writing. Ana Clara Gomes de

Souza, Bruna de Paula Dias, Camila Cavadas Barbosa and Ubiratan da Silva Batista:

Data collection and processing. Arturo Reyes-Sandoval: Writing. Luiz Felipe Leomil

Coelho: Data analysis and writing. Breno de Mello Silva: Conception of the

methodology, data analysis and writing.

References

Lokugamage KG, Narayanan K, Nakagawa K *et al.* Middle East Respiratory Syndrome Coronavirus nsp1 Inhibits Host Gene Expression by Selectively Targeting mRNAs Transcribed in the Nucleus while Sparing mRNAs of Cytoplasmic Origin. *J Virol* 2015; **89**: 10970–10981.

Wu A, Peng Y, Huang B *et al.* Genome Composition and Divergence of the Novel Coronavirus (2019-nCoV) Originating in China. *Cell Host Microbe* 2020; **27**: 325–328.

Guan W, Ni Z, Hu Y *et al.* Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020; **382**: 1708–1720.

Beniac DR, Andonov A, Grudeski E, Booth TF. Architecture of the SARS coronavirus prefusion spike. *Nat Struct Mol Biol* 2006; **13**: 751–752.

Jeffers SA, Tusell SM, Gillim-Ross L *et al.* CD209L (L-SIGN) is a receptor for severe acute respiratory syndrome coronavirus. *Proc Natl Acad Sci U S A* 2004; **101**: 15748–15753.

Gui M, Song W, Zhou H *et al.* Cryo-electron microscopy structures of the SARS-CoV spike glycoprotein reveal a prerequisite conformational state for receptor binding. *Cell Res* 2017; **27**: 119–129.

Belouzard S, Chu VC, Whittaker GR. Activation of the SARS coronavirus spike protein via sequential proteolytic cleavage at two distinct sites. *Proc Natl Acad Sci U S A* 2009; **106**: 5871–5876.

Chan JF-W, Kok K-H, Zhu Z *et al.* Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes Infect* 2020; **9**: 221–236.

Kirchdoerfer RN, Cottrell CA, Wang N *et al.* Pre-fusion structure of a human coronavirus spike protein. *Nature* 2016; **531**: 118–121.

Pallesen J, Wang N, Corbett KS *et al.* Immunogenicity and structures of a rationally designed prefusion MERS-CoV spike antigen. *Proc Natl Acad Sci U S A* 2017; **114**: E7348–E7357.

Hofmann H, Pöhlmann S. Cellular entry of the SARS coronavirus. Trends Microbiol. 2004; **12**: 466–472.

Bertram S, Glowacka I, Muller MA *et al.* Cleavage and Activation of the Severe Acute Respiratory Syndrome Coronavirus Spike Protein by Human Airway Trypsin-Like Protease. *J Virol* 2011; **85**: 13363–13372.

Coutard B, Valle C, de Lamballerie X, Canard B, Seidah NG, Decroly E. The spike glycoprotein of the new coronavirus 2019-nCoV contains a furin-like cleavage site absent in CoV of the same clade. *Antiviral Res* 2020; **176**: 104742.

Walls AC, Park Y-J, Tortorici MA, Wall A, Mcguire AT, Veesler D. Structure, function and antigenicity of

the SARS-CoV-2 spike glycoprotein. doi:10.1101/2020.02.19.956581.

Tai W, He L, Zhang X *et al.* Characterization of the receptor-binding domain (RBD) of 2019 novel coronavirus: implication for development of RBD protein as a viral attachment inhibitor and vaccine. *Cell Mol Immunol* 2020; : 1–8.

Wrapp D, Wang N, Corbett KS *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science (80-)* 2020; **367**: 1260–1263.

Hulswit RJG, Lang Y, Bakkers MJG *et al.* Human coronaviruses OC43 and HKU1 bind to 9-O-acetylated sialic acids via a conserved receptor-binding site in spike protein domain A. *Proc Natl Acad Sci U S A* 2019; **116**: 2681–2690.

Earp LJ, Delos SE, Park HE, White JM. The many mechanisms of viral membrane fusion proteins. Curr. Top. Microbiol. Immunol. 2004; **285**: 25–66.

Xu X, Chen P, Wang J *et al.* Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission. Sci. China Life Sci. 2020; **63**: 457–460.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol* 2018; **35**: 1547–1549.

Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. doi:10.1093/nar/gkh340.

Kim DE, Chivian D, Baker D. Protein structure prediction and analysis using the Robetta server. *Nucleic Acids Res* 2004; **32**: 526–531.

Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veesler D. Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* 2020; **181**: 281-292.e6.

Spassov VZ, Yan L. Accelrys Software Inc., Discovery Studio Modeling Environment, Release 4.0. *Proteins Struct Funct Bioinforma* 2013; **81**: 704–714.

Wu F, Zhao S, Yu B *et al.* A new coronavirus associated with human respiratory disease in China. *Nature* 2020; **579**: 265–269.

Larsen M V., Lundegaard C, Lamberth K, Buus S, Lund O, Nielsen M. Large-scale validation of methods for cytotoxic T-lymphocyte epitope prediction. *BMC Bioinformatics* 2007; **8**: 1–12.

Jensen KK, Andreatta M, Marcatili P *et al.* Improved methods for predicting peptide binding affinity to MHC class II molecules. *Immunology* 2018; **154**: 394–406.

Jespersen MC, Peters B, Nielsen M, Marcatili P. BepiPred-2.0: Improving sequence-based B-cell epitope prediction using conformational epitopes. *Nucleic Acids Res* 2017; **45**: W24–W29.

Kringelum JV, Lundegaard C, Lund O, Nielsen M. Reliable B Cell Epitope Predictions: Impacts of Method Development and Improved Benchmarking. *PLoS Comput Biol* 2012; **8**. doi:10.1371/journal.pcbi.1002829.

Ponomarenko J, Bui HH, Li W *et al.* ElliPro: A new structure-based tool for the prediction of antibody

epitopes. *BMC Bioinformatics* 2008; **9**. doi:10.1186/1471-2105-9-514.

31   Humphrey W, Dalke A, Schulten K. VMD: visual molecular dynamics. *J Mol Graph* 1996; **14**: 33–8, 27–8.

32   BIOVIA DISCOVERY STUDIO ® VISUALIZER DATASHEET BIOVIA DISCOVERY STUDIO 4.0
VISUALIZER: FREE VERSUS COMMERCIAL COMPARISON.
http://accelrys.com/products/datasheets/discovery-studio-visualizer.pdf (accessed 2 Aug2017).

33   Yin C. Genotyping coronavirus SARS-CoV-2: methods and implications. .

34   Mohammad Lokman S, Rasheduzzaman M, Salauddin A *et al.* Exploring the genomic and proteomic
variations of SARS-CoV-2 spike glycoprotein: a computational biology approach.
doi:10.1101/2020.04.07.030924.

35   Phan T. Genetic diversity and evolution of SARS-CoV-2. *Infect Genet Evol* 2020; **81**.
doi:10.1016/j.meegid.2020.104260.

36   Madu IG, Roth SL, Belouzard S, Whittaker GR. Characterization of a Highly Conserved Domain within the
Severe Acute Respiratory Syndrome Coronavirus Spike Protein S2 Domain with Characteristics of a Viral
Fusion Peptide. *J Virol* 2009; **83**: 7411–7421.

37   Yao H, Lu X, Chen Q *et al.* Patient-derived mutations impact pathogenicity of SARS-CoV-2.
doi:10.1101/2020.04.14.20060160.

38   Wan Y, Shang J, Graham R, Baric RS, Li F. Receptor Recognition by the Novel Coronavirus from Wuhan:
an Analysis Based on Decade-Long Structural Studies of SARS Coronavirus. *J Virol* 2020; **94**.
doi:10.1128/jvi.00127-20.

39   Li W, Shi Z, Yu M *et al.* Bats are natural reservoirs of SARS-like coronaviruses. *Science (80-)* 2005; **310**:
676–679.

40   Ou J, Zhou Z, Zhang J *et al.* RBD mutations from circulating SARS-CoV-2 strains enhance the structure
stability and infectivity of the spike protein. *bioRxiv* 2020; : 2020.03.15.991844.

41   Tian X, Li C, Huang A *et al.* Potent binding of 2019 novel coronavirus spike protein by a SARS
coronavirus-specific human monoclonal antibody. Emerg. Microbes Infect. 2020; **9**: 382–385.

42   Shang J, Ye G, Shi K *et al.* Structural basis of receptor recognition by SARS-CoV-2. *Nature* 2020; **581**:
221–224.

43   Ortega JT, Serrano ML, Pujol FH, Rangel HR. Unrevealing sequence and structural features of novel
coronavirus using in silico approaches: The main protease as molecular target. *EXCLI J* 2020; **19**: 400–409.

44   Li Q, Guan X, Wu P *et al.* Early transmission dynamics in Wuhan, China, of novel coronavirus-infected
pneumonia. N. Engl. J. Med. 2020; **382**: 1199–1207.

45   Yuan M, Wu NC, Zhu X *et al.* A highly conserved cryptic epitope in the receptor-binding domains of
SARS-CoV-2 and SARS-CoV. *Science (80-)* 2020; **7269**: eabb7269.

46   He Y, Zhou Y, Liu S *et al.* Receptor-binding domain of SARS-CoV spike protein induces highly potent

neutralizing antibodies: Implication for developing subunit vaccine. *Biochem Biophys Res Commun* 2004; **324**: 773–781.

Qiu T, Mao T, Wang Y *et al.* Identification of potential cross-protective epitope between a new type of coronavirus (2019-nCoV) and severe acute respiratory syndrome virus. J. Genet. Genomics. 2020; **47**: 115–117.

Follis KE, York J, Nunberg JH. Furin cleavage of the SARS coronavirus spike glycoprotein enhances cell-cell fusion but does not affect virion entry. *Virology* 2006; **350**: 358–369.

Table 1 - SARS-CoV-2 Spike protein amino acid variations: All variations are

identified by their position in the polypeptide chain, as well as the physical-chemical

characteristic of the consensus (C.) / Variant (M.) Residue. The residues were pooled on

continents from which the samples were isolated. Protein domains are indicated on the

left and the subdomains and motifs on the right. Mutations that occur at frequencies<

1% are indicated by light gray, > 1% mutations by gray and 614 variations by dark gray.

Domain	Position	C.	M.	Asia (548)	N. America (680)	Oceania (57)	Europe (1014)	S. America (43)	Africa (21)	Subdomain
						%				
S1	5	L	F		0,44		0,59			
	8	L	V	2,01						
	28	Y	N	0,36						NTD
	49	H	Y	0,91	0,44		0,20			
	50	S	L	2,37						
	71	S	F		0,29					
	120	V	I				0,30			
	138	D	Y	0,36						
	153	M	T	0,36						
	157	F	L			0,44				
	181	G	V			0,44				
	221	S	W	0,55						
	239	Q	K				0,59			
	247	S	R			5,26				
	254	S	F				0,69			
	367	V	F	0,36			0,49			RBD
	408	R	I	0,55						
	476	G	S		1,03					
	483	V	A		2,35					
	614	D	G	3,28	20,00	14,04				
	614	G	D				32,45	32,56	9,52	
	653	A	V				0,20			
	655	H	Y		0,29					
	675	Q	H				0,49			
S2	765	R	L	0,36			0,20			FP
	791	T	I	1,09						
	797	F	C				0,30			
	831	A	V				0,99			
	839	D	Y				1,18			
	852	A	V				0,20			
	930	A	V	0,55						HR1
	939	S	F				0,30			
	940	S	F			3,51	0,20			
	941	T	A				0,20			
	943	S	P				0,59			
	1040	V	F	0,36						
	1143	P	L				0,20			
	1176	V	F					6,98		
	1216	I	T	0,36						TM
	1229	M	I				0,30			
	1254	C	F			3,51				
	1259	D	H		0,29					IC
	1263	P	L				1,97			

Hydrophobic	L	V	A	I	M	F
Neutral polar	S	T	N	Q	W	
Positively charged	R	K	P			
Negatively charged	D	E				
Aromatic positively charged	H					
Neutral aromatic	Y					
Neutral non-polar	G					

Table 2. SARS-CoV-2 protein Spike identified epitopes with amino acid variants: The

servers in green represent the T-cytotoxic cells and yellow for T-Helper cells

predictions. Antigenic prediction servers of B cells are shown in red and orange colors,

for conformational and linear epitopes, respectively. All epitopes are identified by their

positions in the polypeptide chain, as well as by their amino acid sequences. A

highlighted letter in red represents the variation present in each epitope. The highlight in

light gray represents the presence of mutations with frequencies on the coding region >

1% and the epitopes with the mutations in residue 614 are represented by dark gray.

Server	Position	Peptide
NetMHCpan 4.0	240-249	TLLALHR**S**YL
	610-620	VLYQ**D**VNCTEV
	612-620	YQ**D**VNCTEV
	786-794	KQIYK**T**PPI
	1209-1218	YIKWPWY**I**WL
	1215-1224	Y**I**WLGFIAGL
NetCTL 1.2	50-58	**S**TQDLFLPF
	136-144	CN**D**PFLGVY
	152-160	W**M**ESEFRVY
	612-620	YQ**D**VNCTEV
	652-660	GAE**H**VNNSY
NetMHCII 2.3	50-64	**S**TQDLFLPFFSNVTW
	167-181	TFEYVSQPFLMDLE**G**
	232-246	GINITRF**Q**TLLALHR
	233-247	INITRF**Q**TLLALHR**S**
	234-248	NITRF**Q**TLLALHR**S**Y
	235-249	ITRF**Q**TLLALHR**S**YL
	236-250	TRF**Q**TLLALHR**S**YLT
	237-251	RF**Q**TLLALHR**S**YLTP
	238-252	F**Q**TLLALHR**S**YLTPG
	239-253	**Q**TLLALHR**S**YLTPGD
	758-772	SFCTQLN**R**ALTGIAV
	759-773	FCTQLN**R**ALTGIAVE
	760-774	CTQLN**R**ALTGIAVEQ
	761-775	TQLN**R**ALTGIAVEQD
	762-776	QLN**R**ALTGIAVEQDK
	763-777	LN**R**ALTGIAVEQDKN
BepiPred-2.0	14-33	QCVNLTTRTQLPPA**Y**TNSFT
	59-81	FSNVTWFHAIHV**S**GTNGTKRFDN
	141-163	LGVYYHKNNKSW**M**ESEFRVYSSA
	178-191	DLE**G**KQGNFKNLRE
	248-260	YLTPGD**S**SSGWTA
	404-424	GDEV**R**QIAPGQTGKIADYNYK
	455-478	LFRKSNLKPFERDISTEIYQA**G**ST
	673-691	SY**Q**TQTNSPRRARSVASQS
	786-800	KQIYK**T**PPIKD**F**GGF
	828-843	LADAGFIKQYG**D**CLGD
ElliPro	14-38	QCVNLTTRTQLPPA**Y**TNSFTRGVYY
	56-270	LPF...**S**...**D**...**M**...**G**...**Q**...**S**...**S**...GYL
	391-424	CFT...**R**QIAPGQTGKIADYNYK
	434-511	IAW...YQA**G**STP...RVV
	434-580	IAW...YQA**G**STP...DPQ
	434-584	IAW...YQA**G**STP...LEI
	675-690	**Q**TQTNSPRRARSVASQ
	675-691	**Q**TQTNSPRRARSVASQS
Discotope -2.0	1138-1160	YDPLQ**P**ELDSFKEELDKYFKNHT

Figure 1. The general outline of the methodology. Flowchart of the three main stages of

the study.

Figure 2. Structural mapping of SARS-CoV-2 Spike protein: On the left side, the homo-

trimer is showed as a newcartoon. The B and C chains are represented in gray and the A

chain in blue. On the right side, the monomer is also showed as a newcartoon with the

representation of domains, subdomains, motifs, N-Glycan's, and disulfide bonds. The

blue and gray shadows above structure represent the portions of S1 and S2 domains,

respectively.

Figure 3. SARS-CoV-2 Spike protein variant amino acid residues: The homo-trimer

structure is represented in newcartoon. The positions with the mutations among residues

of non-homologous physico-chemical characteristics are represented on the surface in

the A chain (light blue) only. In the "SIDE IN" perspective, the B and C chains are

represented in transparent newcartoon. The colors of each residue are represented by the

continent where the variations were identified.

Figure 4. Mapping of epitopes with variants on SARS-CoV-2 Spike protein: The homo-

trimer structure was represented in newcartoon. Epitopes are represented by their

surfaces in the A chain, in colors that represent the type of prediction as highlighted at

the top of the figure. The histogram below the structures represents the Spike protein

polypeptide chain with its domains, subdomains, motifs and epitopes proportionally

represented in relation to the primary sequence of the proteins.

STEP 1

2139 complete genomes collected on the SARS-CoV2 GISAID webserver. In addition, 116 sequences from NCBI of the Spike protein and 108 from ViPR were collected (deposited between 11/12/19 to 03/27/20)

The sequences were divided by continents, aligned and their variants identified in the MEGA-X software.

STEP 2

The three-dimensional model of the SARS-CoV2 Spike protein homo-trimer obtained though the Robetta server using the partial reference crystal 6VXX. Geometry optimization steps made in Discovery Studios software

STEP 3

The sequence of the SARS-CoV2 Spike protein NC_045512.2 used as a reference in the prediction of linear epitopes. The three-dimensional model of the homo-trimer of the Spike protein from SARS-CoV2 was used as a reference for conformational epitopes.

Prediction of linear epitopes for cytotoxic T cells (NetCTL 1.2 and NetMHCpan 4.0 webservers). Prediction of linear epitopes for B cells and T-Helper (NetMHCII 2.3 and BEPIPRED 2.0 webservers)

Prediction of conformational epitopes for B cells (EliPRO and DiscoTop webservers)

1 **Figure 1.**

1 **Figure 2.**

2 (See video 1 in supplemental material)

2 **Figure 3.**

1   **Figure 4.**