Mechanistic insights into ventricular arrhythmogenesis of hydroxychloroquine and azithromycin for the treatment of COVID-19

Authors:

Gongxin Wang, *MSc*[1], Chieh-Ju Lu, *D.Phil*[1], Andrew W. Trafford, BVSc, PhD, MRCVS[2,] Xiaohui Tian, *BSc*[3], Hannali M Flores, *MSc*[4], Piotr Maj[5], *PhD*, Kevin Zhang[6], *BSc*, Yanhong Niu, MD[7], Luxi Wang, *BSc*[1], Yimei Du, *PhD,*[8] Xinying Ji, *PhD, MD*[3], Yanfang Xu, *PhD, MD*[9], Dan Li, *PhD*[10], Neil Herring, *MA, DPhil, FRCP*[10], David Paterson, *D.Phil, DSc, FRSB*[10], Christopher L.-H. Huang, *PhD, MD, DSc, FRSB, FESC*[11,13], Henggui Zhang, *PhD, FRSB, FRSA*[4,12,13,], Ming Lei, *DPhil, MD, FRSB*[10], Guoliang Hao. *PhD, MEng.*[1,10]

Affiliations:

[1]Henan SCOPE Research Institute of Electrophysiology Co. Ltd., China

[2]Unit of Cardiac Physiology, Institute of Cardiovascular Sciences, Manchester Academic Health Sciences Centre, The University of Manchester, UK.

[3]Department of Pharmacy, Huaihe Hospital and College of Medicine, Henan University, China.

[4]Biological Physics Group, Department of Physics and Astronomy, The University of Manchester, UK

[5]Department of Pharmacology, University of Oxford, Oxford, UK

[6]School of Medicine, Imperial College of London, UK

[7] Fuwai Central China Cardiovascular Hospital, Beijing, China

[8]Department of Cardiology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430022, China

[9]Department of Pharmacology, Hebei Medical University, China

[10]Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, UK

[11]Physiological Laboratory and Department of Biochemistry, University of Cambridge, UK

[12]Peng Cheng Laboratory, Shenzhen, China

[13]Key Laboratory of Medical Electrophysiology of the Ministry of Education and Institute of Cardiovascular Research, Southwest Medical University, Luzhou, China

Correspondence:

Dr. Gaoliang Hao, Henan SCOPE Research Institute of Electrophysiology Co. Ltd., Kaifeng 475000, China and Department of Physiology, Anatomy and Genetics, University of Oxford, Tel: 008613552324220, Fax: 008613552324220, E-mail: guoliang.hao@epscopelab.com

Or Prof. Ming Lei, Department of Pharmacology, University of Oxford, Mansfield Road, Oxford OX1 3QT, United Kingdom. Tel: 00441865271850; Fax: 00441865271850 E-mail: ming.lei@pharm.ox.ac.uk

CLHH, HZ, ML and GLH are joint senior authors

Article length: 4711 words, excluding abstract and acknowledgement

CLINICAL PERSPECTIVE

What Is New?

- We explore the cardiac electrophysiological effects of hydroxychloroquine (HCQ) and azithromycin (AZM) alone and in combination, to investigate their safety as a treatment for Covid-19.

- Using the *Comprehensive in-vitro Pro-arrhythmia Assay* (CiPA) guidelines, we combine patch clamp studies of individual ion channels, whole heart voltage, Ca^{2+} and electrocardiographic measurements and human *in-silico* modelling to directly assess the pro-arrhythmic potential of these drugs.

  What Are the Clinical Implications?

- The results provide mechanistic insights for HCQ and its synergistic effects with AZM therapy in terms of QT interval and predisposition to arrhythmia.

- The translatable cardiac safety assessment associated with clinical HCQ/AZM therapy that emerge may have significant implications for its use in Covid-19 management.

ABSTRACT

Background:

Recent reports on the use of hydroxychloroquine (HCQ) alone, or combined with azithromycin (AZM) in the management of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) have raised cardiac safety concerns. Currently, there is limited mechanistic data evaluating cardiac safety with HCQ and AZM therapy.

Methods:

Using comprehensive In Vitro ProArrhythmia Assay (CiPA) Schema IC_{50} paradigms, we examined the cardiac electrophysiological effects of HCQ and HCQ/AZM. Molecular modelling explored HCQ and AZM binding properties to hERG. Langendorff-perfused guinea-pig hearts were electrically and optically mapped by multi-electrode array and voltage (RH237) and Ca^{2+} (Rhod-2 AM) dyes. Human action potential and ion current reconstructions were performed in silico.

Results:

HCQ blocked I_{Kr} and I_{K1} with IC_{50} concentrations (10 ± 0.6 and 34 ± 5.0 μM) within the therapeutic range observed clinically. HCQ also blocked I_{Na} and I_{CaL} but at higher IC_{50}, whilst I_{to} and I_{Ks} were unaffected. Contrastingly, AZM produced minor inhibition of I_{Na}, I_{CaL}, I_{Ks}, and $I_{Kr,}$ with no effect on I_{K1} and I_{to}. HCQ + AZM combined inhibited I_{Kr} and I_{K1} with IC_{50}s of 7.7 ± 0.8 μM and 30.4 ± 3.0 μM, but spared I_{Na}, I_{CaL} and $I_{to,}$. Molecular modelling confirmed potential HCQ binding to hERG. Cardiac mapping and ECG studies in isolated hearts demonstrated that HCQ slowed heart rate and ventricular conduction with associated prolongation of PR, QRS and QT intervals. Optical mapping demonstrated, and prolonged, more heterogeneous, action potential durations and intracellular Ca^{2+} transients. These effects were accentuated with combined HCQ+AZM treatment, which elicited electrical alternans, re-entrant circuits and wave breaks. Reconstruction in a human in-silico model demonstrated that this is attributable to the integrated action of HCQ and AZM reducing I_{Kr}, I_{Ks} and I_{K1}.

Conclusions:

These data provide an electrophysiological basis for recent FDA guidelines cautioning against combined HCQ/AZM administration for the treatment of Covid-19 on the grounds of potential

cardiac safety. We would strongly recommend monitoring of electrocardiographic QT interval with the use of this combination of medications.

INTRODUCTION

The global severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic has now infected >4.13 million individuals causing 285660 deaths as of 12[th] of May 2020 (www.ecdc.eurpoe.eu). Although there are no approved drugs to prevent or treat SARS-CoV-2 infection, recent use of the anti-malarial 4-aminoquinolines chloroquine (CQ) or hydroxychloroquine (HCQ) alone, or combined with the antibiotic azithromycin (AZM) in Covid-19 therapy has attracted global attention and extensive early clinical use and trials[1,2]. An initial nonrandomized open label clinical trial of 20 cases, had associated HCQ treatment with viral load reduction or disappearance in COVID-19 patients, effects reinforced by AZM[3]. This subsequently prompted the US Food and Drugs Administration (FDA) authorization of emergency use of CQ or HCQ, alone or in combination with AZM in Covid-19 therapy[4, 5]. Currently, large number of trials of these combinations, with or without the addition of further drugs, are registered worldwide[1,2].

However, several recent studies have flagged cardiac safety concerns for the CQ/HCQ and AZM regime in Covid-19 therapy. Particular concern has been raised regarding potential synergistic effects of HCQ and AZM on QT duration and cardiovascular mortality[6-10]. Higher mortalities and incidences of prolonged corrected QT (QTc) intervals were reported in patients at a Brazilian hospital receiving the high dose (12 g total dose over 10 days, ~1.2 g/day, giving plasma levels of ~10 µM) than the lower dose of CQ, typified by its use in other conditions such as rheumatoid arthritis and systemic lupus erythematosus (SLE). This phase 2b randomized Covid-19 trial was terminated after enrolment of only 81 out of the target of 440 patients owing to occurrences of adverse events[6]. A retrospective data analysis from United States Veterans Health Administration medical centers suggested increased overall mortality even in patients treated with HCQ alone[7]. The largest available analysis from an international consortium on HCQ safety covering over 950,000 patients from six countries declared HCQ safe for short-term use at doses currently used for conditions such as rheumatoid arthritis. However, it urged caution in its use when combined with AZM in view of potential risks of cardiovascular mortality related to QTc prolongation[8]. Two recent reports of 90 and 201 patients hospitalized with COVID-19 treated with HCQ, with or without AZM, associated these drugs with QTc prolongation, especially when used in combination[9] [11]. Although these studies are small, only one case of torsades de pointes was documented and the therapy was seldom discontinued. They suggested that more QT interval monitoring studies should be undertaken before final recommendations can be made[11]. Based on these findings, the FDA

issued a cautions warning against use of HCQ or CQ for COVID-19 outside of the hospital setting or a clinical trial, due to risks of heart rhythm problems [4]. This does not affect FDA-approved uses in malaria, systemic lupus erythematosus (SLE), and rheumatoid arthritis where they have been used many years. Both HCQ and AZM have been in clinical use alone or in combination in other disease conditions for many years with cardiac safety then not considered a major concern.

Currently, there is very limited data evaluating cardiac safety particularly concerning use of HCQ in combination with AZM, although multiple randomized trials are currently in progress evaluating the efficacy of this combination for Covid-19. We therefore explored the cardiac safety of HCQ alone or in combination with AZM using the FDA recommended Comprehensive in Vitro Proarrhythmia Assay (CiPA) protocols to understand arrhythmia mechanisms[12]. Furthermore, we expanded the CiPA protocol by introducing assessments of effects on membrane and Ca^{2+} clocks, their interaction and cardiac conduction recently suggested as crucial indicators for ventricular arrhythmogenesis[13, 14].

METHODS

All procedures were approved by The Institutional Animals Ethics Committees at Henan University, Kaifeng, China under and the national guidelines under which the institution operates, and also conform with the NIH Guide for the Care and Use of Laboratory Animals. Guinea pigs (male; 300-350 g) were provided by The Experimental Animal Center of the Medical College of Xi'an Jiaotong University and maintained in a pathogen−free facility at Henan University with *ad libitum* access to food and water. Animals were humanely sacrificed by intraperitoneal injection with 3% pentobarbital sodium (50 mg/kg).

Single-cell electrophysiological recordings

Core human cardiac channels were heterologously expressed in recombinant HEK293 or CHO cell lines and studied using conventional patch clamp methods. Stable cell lines expressing Nav1.5 and Cav1.2/β2/α2/δ1, hERG, Kir2.1, Kv4.3, KCNQ1/E1 were used. All currents were recorded using a MultiClamp 700A amplifier, 1550A digitizer and pCLAMP 10.6 software (Molecular devices, USA). Patch electrodes with resistances of 2-5 MΩ were pulled with a P-97 micropipette puller (Sutter Instruments, USA). The conventional whole-cell recordings were obtained by manual suction for hERG, Kir2.1, Kv4.3 and Nav1.5 studies. Perforated

whole-cell recordings of KCNQ1/E1 and Cav1.2 were obtained following perforation with 200 µg/ml Amphotericin included in the pipette solution. After whole-cell mode were obtained, cells were used in voltage clamp mode for the current recordings. The external solution contained (mM)：NaCl 137；MgCl$_2$ 1；KCl 4；glucose 10；HEPES 10；CaCl$_2$ 1.8, pH adjusted to 7.4 with NaOH. The K$^+$-containing pipette solution contained (mM)：KCl 40；KF 100；MgCl$_2$ 2；EGTA 5；HEPES 10，pH adjusted to 7.2 with KOH. The sodium and calcium current pipette solution contained (mM): CsCl 65；CsF 75；MgCl$_2$ 2.5；EGTA 5；HEPES 10, pH adjusted to 7.3 with CsOH.

Molecular modelling and docking studies

The experimental structure of the open hERG channel (PDB ID: 5va1; 1) was selected as a receptor for molecular docking studies. The structure was prepared for docking using Protein Preparation Wizard using default settings including modelling of missing loops using Prime. The loops spanning residues 433-448, 578-582, and 588-602 in each subunit were successfully rebuilt. The centroid of residues Y652 and F656 from every subunit was defined as the centre of the docking binding site. The HCQ structure was obtained from PubChem and prepared for docking using LigPrep. Three separate rigid-receptor docking methods were used including: (i) Glide SP docking, (ii) FlexX docking followed by HYDE re-scoring, and (iii) AutoDock Vina docking with each set to output 20 poses per LigPrep HCQ state. Poses from Glide, FlexX, and AutoDock Vina were also re-scored using RF-Score-VS.

Langendorff- perfused isolated hearts

Guinea pigs (male; 300-350 g) were humanely killed by intraperitoneal injection with pentobarbital sodium (50 mg/kg). Hearts rapidly excised after thoracotomy were mounted onto a Langendorff perfusion system and perfused with a modified Tyrode's solution (119 NaCl, 25 NaHCO$_3$, 4 KCl, 1.2 KH$_2$PO$_4$, 1 MgCl$_2$, 1.8 CaCl$_2$.2H$_2$O, and 10 D-glucose (mM) equilibrated with 5% CO$_2$ and 95% O$_2$) with the flow rate of 8 ml/min at 37°C. Hearts were perfused and monitored for stability for 20 mins before experimental procedures commenced.

Electrical mapping of *ex vivo* heart preparations

A multi-electrical array (MEA) mapping system (EMS64-USB-1003, MappingLab Ltd., UK) was employed to record extracellular potentials (ECP) from the epicardium. Two 64 channel MEAs (MappingLab Ltd., UK) were employed for left atrial and left ventricular simultaneous recordings. ECP signals were recorded at a 10 kHz sampling using synchronized 64 channel EMS64-USB-1003 recording systems and recorded using EMapRecord 5.0 software (MapingLab Ltd., UK) and stored in a computer for subsequent off-line analysis. Electrocardiograms (ECG) were continuously recorded by two ECG electrodes (MappingLab Ltd., UK) positioned on the right atrium and left ventricle respectively.

The epicardial isochronal activation map during sinus rhythm or imposed pacing was plotted as time differences between the earliest activation site and the individual activation site on each channel. Activation times were calculated as the point of maximal negative slope of activation waveforms. The conduction velocity (CV) was calculated from the difference in timing and the known distance between the recording points using EMapScope 5.0 software (MappingLab Ltd.).

Optical mapping of *ex vivo* heart preparations

After the Langendorff-perfused hearts reached steady state, contraction artefacts were mimimised using blebbistatin (10 μM). Dye loading was aided by pre-perfusion with pluronic F127 (20 % w/v in DMSO). Rh237 (1μg/ml) and Rhod2-AM (1μg/ml) were perfused to enable simultaneous membrane potential and Ca^{2+} measurements at 37 °C .

Two 530 nm LEDs (LEDC-2001, MappingLab Ltd) were used to illuminate the heart after their emissions were bandpass filtered (wavelengths 530±20 nm) to minimize stray excitation light reaching the dyes. The fluorescence light was passed through a 550 nm long-pass filter and then a dichroic mirror with a cut off of 638 nm. Fluorescence light with wavelengths above 638 nm was passed through a 700 nm long-pass filter and then imaged by the camera for recording voltage signals. Fluorescence light below 638 nm was passed through a bandpass filter (585±40 nm) then imaged by the camera for recording calcium signals (OMS-PCIE-2002, MappingLab Ltd). The raw spatial resolution was 128-by-128 pixels, the total mapping area was 16×16 mm and the temporal resolution was 900 frames/second.

The cameras of the optical mapping system, LED lights and stimulator and ECG recording were simultaneously driven by an 8 channel TTL analog-digital converter and OMapRecord 4.0 software (MappingLab Ltd.). For the analysis of optical mapping signals and generation of isochronal maps, data were semi-automatically processed using OMapScope 5.0 software

(MappingLab Ltd). In this experimental configuration, stable recordings for voltage (V_m) and intracellular Ca^{2+} (CaT) signals could be obtained for >2 hours in preliminary experiments assessing system stability.

Arrhythmia induction protocol

Stimuli were generated by an isolated constant voltage/current stimulator (VCS3001, MappingLab Ltd., UK). They were delivered with a platinum electrode onto the epicardial apex at an amplitude 1.5 × the diastolic voltage threshold and a 2 ms pulse width. The optical mapping was performed in conjunction with an "alternans mapping" protocol. The heart was consecutively paced at basic cycle lengths (BCL) from 210 with 10 ms decrement until 170 ms. The heart was paced 50 times with 2 ms duration pulses and the optical mapping recording was performed at the last 2 second in each episode. This protocol was performed both before and after drugs were delivered.

In silico model reconstruction

The O'Hara-Rudy (ORd) human ventricular model[15] was modified to incorporate the actions on multi-channel block of HCQ, including the dose-dependent block of I_{Na}, I_{CaL}, I_{to}, I_{Kr}, I_{Ks} I_{K1} and the Na^+-Ca^{2+} exchange current, I_{NCX}. This model was selected as it was developed from and validated against experimental data from human ventricular cells and found suitable for studying actions of drugs. The simulations evoked action potentials by a supra-threshold stimulus in control runs and following application of HCQ, and combined application of HCQ and AZM. Action potential time courses characterized by AP durations at 90% repolarisation (APD_{90}) and maximal upstroke velocity (MUP) and the underlying ion channel currents targeted by the drugs were recorded for analysis.

Statistical analysis

All patch clamp recorded data were analyzed using Clampfit 10.6 (Molecular Devices, USA), OriginPro 8.0 (Origin Lab, USA) and Adobe Illustrator 10 (Adobe, USA). The concentration-response curve was fitted by the logistic equation $y = A_2 + (A_1 - A_2)/[1 + (x/x_0)^p]$, where x is the drug concentration, and p is the Hill coefficient. All data are expressed as the means ± SEM. One-way ANOVA followed by multiple-comparison test was used to evaluate multiple test

treatments. A value of $p < 0.05$ was considered to be statistically significant. IC_{50} denotes the concentration determined for half-maximal inhibitory effects. In the figures, the designations for p values are: $*p < 0.05$, $**P < 0.01$ and $***P < 0.001$ respectively.

RESULTS

The four groups of experiments conducted assessed the mechanistic basis for the clinical reports bearing upon cardiac safety of HCQ, whether applied alone or in combination with AZM.

Patch-clamp studies of single channel characteristics

The first series of experiments assessed the effects of HCQ (Fig. 1) and AZM alone (Fig. 2) and in combination (Fig. 3) on a range of cardiac ion channels. The experiments on HCQ and AZM themselves adopted concentrations fully encompassing plasma levels associated with their clinical use. The same HCQ concentration range was then combined with 10 μM AZM. These examined inward, cardiac channel currents from hNav1.5 (I_{Na}, *A*), Cav1.2 ($I_{Ca,L}$, *B*), mediating or maintaining membrane action potential depolarization. They also studied outward Kv4.3 (I_{to}, *C*), hERG (I_{Kr}, *C*), KCNQ1/E1 (I_{Ks}, *E*) and Kir2.1 (I_{K1}, *F*) normally mediating action potential recovery. This encompassed the range of ion channels recommended by the CiPA protocol.

Fig.1 (left panels, i) exemplifies observed ionic currents in the presence of 0, 1, 10 and 100 μM HCQ. The insets summarize the related voltage clamping protocols, fully described in extended Methods in Supplementary data. Five concentrations ranging from 10^{-8} to 10^{-4} M were tested for each current examined in 3 to 6 cells at each concentration. Mean concentration-response plots (right panels, ii) used the maximum magnitudes of each current. Where applicable, the consequent alterations in such current magnitudes following pharmacological challenge were used to derive IC_{50} values.

Fig.1 (right panels, ii) plots mean (±SEM) fractional reductions in current magnitude against HCQ concentration for I_{Na} (A), I_{CaL} (B), I_{to} (C), I_{Ks} (D), I_{Kr} (E) and I_{K1} (F) . μM-HCQ inhibited both I_{Kr} tail and I_{K1} pulse currents with IC_{50}s of 10±0.6 μM (D) and 34±5 μM (F), respectively, with more marked effects on I_{Kr}, where the IC_{50} fell close to expected therapeutic levels in current Covid-19 treatment regimes[6], than I_{K1}. In contrast, μM-HCQ has less effect on I_{Na} and I_{CaL} with IC_{50}s of 113 ± 78 μM and 209 ± 94 μM respectively, Nevertheless, HCQ caused graded I_{Na} reductions through the entire HCQ concentration range typified in a 22±1.8% reduction at 1 μM. No significant inhibition of I_{to} and I_{Ks} by μM-HCQ was observed .

Fig. 2 presents corresponding results obtained with AZM applied by itself similarly encompassing clinical therapeutic levels. A previously reported dose regime administering 500, 250 and 250 mg/day on each of three successive days gave a day 2 serum concentration of 0.22 μM[16]). However, the entire, 0-100 μM explored AZM concentration range yielded only minimal inhibitory effects. Even 10 μM and 100 μM AZM reduced I_{Na} (A), I_{CaL}, (B) I_{Ks} (E) and I_{Kr} (D) by <17% and <35% respectively and produced negligible effects on I_{K1} (F) and no effect on I_{to} (C).

Finally, Fig. 3 summarizes results of applying the same range of five HCQ concentrations displayed in Fig. 1 now combined with 10 μM AZM. Adding AZM did not significantly alter the effects of HCQ on I_{Na} (A), I_{CaL} (B), I_{to} (C), and I_{Ks} (E), but reduced the IC_{50} for I_{Kr} (IC_{50} = 7.7 $\pm$ 0.8 μM) and I_{K1} (30.4 $\pm$ 3.0 μM).

Computational molecular docking determining the interaction between HCQ, AZM and hERG

The above findings particularly concerning pharmacological effects on hERG prompted computational molecular docking explorations for possible interactions between HCQ and AZM and a recent hERG homotetramer structure (PDB ID: 5va1; 1) (Fig.4). We used scoring docking visual inspection irrespective of the docking protocol. Fig.4 illustrates the HCQ-hERG interaction, derived using three separate rigid-receptor docking methods. HCQ interacts predominantly with Y652 from one or more subunits. Interactions with other residues including F656 but also S624, A653, and S660 occurred less frequently. Variability in observed docking positions could be attributed to the highly symmetric structure of the hERG channel pore constituting the binding site enabling many equivalent positions. In contrast, no interactions were demonstrated between AZM and hERG (data not shown).

Electrical mapping and electrocardiographic studies in *ex vivo* heart preparations

We then explored the effects of HCQ alone and combined with AZM in the whole heart on electrophysiological activation and recovery, their electrocardiographic correlates and the consequent Ca^{2+} signalling processes reflecting excitation-contraction coupling.

Fig. 5A summarises sites of apical stimulation, and multi-electrode array (MEA) and ECG recording in the electrical mapping experiments. Representative left atrial (LA) and left ventricular (LV) isochronal conduction maps (B) obtained in the presence of 0 - 10 µM HCQ revealed a clear reduction of conduction velocity and increased conduction heterogeneities at 10 µM HCQ. Synergistic inhibitory effects on conduction velocity were observed when hearts were treated with the 10 µM HCQ in combination with 10 µM AZM. All these effects were reversed on drug washout (C(i)). 10 µM HCQ significantly decreased conduction velocities of excitation with further decreases in velocity with the 10 µM HCQ and 10 µM AZM combination (C(ii)).

Quantification of the corresponding electrocardiogram (ECG) traces (D(i)) demonstrated that both 10 µM HCQ, and 10 µM HCQ in combination with 10 µM AZM, decreased heart rate (HR)(D(i)), and increased QT interval (D(iii)), PR interval (D(iv)) and QRS duration (D(v)). In particular, AZM when combined with HCQ augmented its effects on HR, and QT and PR interval. Together these results suggested that 10 µM HCQ, and 10 µM HCQ and 10 µM AZM combined, exerted synergistic inhibitory effects on heart rate and conduction, and prolonged cardiac repolarisation.

Optical mapping studies in *ex vivo* heart preparations

We next investigated the effects of HCQ alone and in combination with AZM using optical fluorescence mapping in sinus-paced Langendorff-perfused hearts using the recording and ECG monitoring configurations illustrated in Fig. 6A(i). Voltage mapping measurements of action potential initiation and conduction (Fig. 6A(ii)) demonstrated bradycardic effects (A(iii)), visible conduction heterogeneities and slowed conduction (A(ii)) at 10 µM HCQ, effects were augmented by combination with 10 µM AZM (Fig. 6A (iv)). Similarly, optical potentiometric mapping revealed that 10 µM HCQ prolonged action potential duration at 90% repolarisation (APD_{90}) (B(i and ii)) with 10 µM HCQ, effects further prolonged by its combination with AZM (Fig.5B(iii)).

Ca^{2+} dye, Rhod-2 AM (CaD), mapping of ventricular calcium transient (CaT) characteristics demonstrated increased magnitudes of and dispersions in CaD durations at 90% recovery (CaD_{90}) (Figure 6C(i)). The latter were attributed to changes late in the CaT records (C(ii)). The quantified CaD_{90} magnitudes (B(iii)) and dispersions (B(iv)) both increased with 10 but

not 1.0 µM HCQ. The combination of 10 µM HCQ with either 1 or 10 µM AZM exerted significantly greater effects than 10 µM HCQ alone.

Optical mapping assessments of arrhythmic tendency in *ex vivo* heart preparations

Finally, voltage dye optical mapping studies during ventricular pacing were used to assess susceptibility to arrhythmia induction before and after drug challenge in isolated perfused hearts (Fig.7A.B). Whilst 10 µM HCQ did not exert detectable arrhythmic effects at the cycle lengths (CL) used, a combination of HCQ and AZM produced AP alternans at 180 ms CL and episodes of non-sustained VT at 170 ms CL in 5 out of 5 hearts. The corresponding potentiometric maps obtained during the colour coded timepoints 1-6 are represented in C. At a CL of 170 ms, the conduction maps with HCQ+AZM demonstrated significant disruptions of the normally coherent propagation of excitation waves with evidence of excitation re-entry and wave break. Thus, the electrophysiological changes outlined above associated particularly with HCQ and AZM in combination were associated with a ventricular arrhythmic tendency.

Computational simulations of HCQ and AZM on human ventricular action potentials

To translate the experimental data describing the multi-ion channel blocking effects of HCQ and/or AZM to human ventricular excitation, the ORd model was implemented to reconstruct the impacts of HCQ alone and combined with AZM, on human ventricular action potentials including APD_{90} and maximal upstroke velocities.

Fig. 8A displays predicted action potentials under control conditions and in the presence of HCQ, and HCQ and AZM combined. It was shown that with the 10 µM HCQ alone, the APD was prolonged by 100 ms as compared to the control condition. With the combined action of HCQ and AZM, a synergistic action of the two drugs on APD prolongation was seen, which was manifested in a greater increase of APD90 of ~300 ms (Figure 8B). Such an APD prolongation can be attributable to the integrated action of reduced repolarisation potassium channel currents of I_{Kr}, I_{Ks} and I_{K1}, and reduced depolarising currents of I_{Na}, I_{NaL}, I_{CaL} and I_{NaCa}. A small reduction in action potential maximum upstroke velocities was also observed by both HCQ and HCQ+AZM due to the effects on I_{Na}.

DISCUSSION

Here we demonstrate that HCQ is able to block a variety of cardiac ion channels, in particular hERG (I_{Kr}) and Kir2.1 (I_{K1},) and to a lesser degree hNav1.5 (I_{Na}). The effects of HCQ on hERG and Kir2.1 are potentiated by AZM. At the level of the whole heart this translates to prolongation of ventricular APD and slowing of conduction associated with QT, PR and QRS prolongation on the ECG. This together with an increase in the dispersion and heterogeneity of APD and intracellular calcium transients increases the probability of arrhythmia induction. Moreover, hERG binding by HCQ is predicted by computational molecular docking and human *in-silico* modelling of the ion channel effects of HCQ and AZM recapitulates the observations on APD. This mechanistic electrophysiological data complements ongoing multiple randomized trials evaluating the clinical efficacy and cardiac safety profile of these drugs as a treatment for Covid-19. These findings thus directly align with recent FDA guidelines concerning their use.

Using the approach recommended by the CiPA guidelines [12] we examined the effects of HCQ and AZM on human cardiac ion channels responsible for the main inward (hNav1.5 (I_{Na}) Cav1.2 ($I_{Ca,L}$)) and outward (Kv4.3 (I_{to}), hERG (I_{Kr}), KCNQ1/E1 (I_{Ks}) and Kir2.1 (I_{K1})) currents in the human ventricular myocardium. HCQ and AZM were studied both alone and in combination at a wide concentration range encompassing and extending beyond reported therapeutic levels. The latter typically fell around 1.15-4.5 μM[17] and 0.5-0.7 μM [18] for HCQ in SLE and rheumatoid arthritis patients on a daily 400 mg dose. In contrast, its dose used in Covid-19, 1.2 g/day, would give substantially higher therapeutic concentrations of ~10 μM[6]. Blood AZM levels following successive daily 500 mg and two 250 mg doses were reported at 0.22 μM[16]. HCQ both alone and combined with AZM synergistically inhibited I_{Kr} and I_{K1} with IC$_{50}$s at the high end of reported therapeutic levels, with some minor inhibition of I_{Na} in this range.

Others have also explored the electrophysiological effects of HCQ or AZM on some cardiac ion channels at three times the clinically relevant concentration and above [16,19], however our study is the first to systematically evaluate HCQ and AZM in combination. The only study to have evaluated the electrophysiological effects of HCQ specifically looked at sinoatrial cells and demonstrated a significant reduction in the hyperpolarization-activated current (I_f),

although reductions in I_{CaL} and I_{Kr} were also noted at 3 μM[19]. The action of HCQ on I_f explains the bradycardia in response to HCQ that we also observe here and further illustrates its breadth of action on cardiac electrophysiology. Moreover, administering a bradycardic medication in response to sepsis or SARS where hypotension is present could potentially exacerbate septic shock. Administering a bradycardic medication in response to sepsis or SARS where hypotension is present could potentially exacerbate septic shock.

At the level of the whole heart, our optical mapping studies demonstrate prolongation of ventricular APD as well CaD in response to HCQ, which was further exacerbated in combination with AZM. This would be expected from inhibition of I_{Kr} and I_{K1} and translates to QT prolongation on the ECG. It is interesting to note that even prior to the Covid-19 pandemic, series of case reports have led to both HCQ and AZM being listed as definite causes of prolonged QT interval and risk of torsade de pointes for patients with long QT syndrome on _crediblemeds.org_[20].

Prolonged APD predisposes to early afterdepolarisations (EADs) during the plateau phase of the ventricular action potential, classically thought to be due to reactivation of the I_{CaL}[21]. EADs are also more common during conditions of intracellular Ca^{2+} overload, such as increased sympathetic drive, because I_{CaL} can be enhanced by high intracellular Ca^{2+} levels and by L-type Ca^{2+} channel phosphorylation by CaMKII[22]. EADs may potentially trigger an ectopic beat and initiate an arrhythmic event given a suitable substrate.

Using both multi-electrode arrays and optical voltage mapping, we also demonstrate a reduction in ventricular conduction in response to HCQ and HCQ+AZM as would be expected from blockade of Na channels. Compromised atrioventricular and intraventricular conduction is also evident electrocardiographically, as reflected by a prolonged PR interval and QRS duration. Slow ventricular conduction as well as prolongation of the PR interval and widening of QRS duration is also observed with blockade of Na channels by local anaesthetic agents [23]. Such drugs can be pro-arrhythmic as demonstrated by the CAST trials[24]. Cardiac conduction is also influenced by gap junction expression and function as well as Na channel availability [25] and it is also possible that HCQ and/or AZM may have direct or indirect effects on connexins. Slowed conduction combined with increased dispersion of calcium handling and repolarisation offer a suitable substrate to potentially sustain re-entry. We tested this directly using

potentiometric maps and ECG recording during pacing protocols. Combined exposure to HCQ and AZM, but not HCQ alone provoked alternans, ventricular tachycardic episodes, and re-entrant excitation following progressively accelerated cardiac pacing.

Consistent with experimental findings, computational molecular docking simulations utilizing a recent hERG homotetramer structure (PDB ID: 5va1; 1) demonstrate potential HCQ binding. This is predominantly with Y652 from one or more hERG subunits. Interactions with other residues including F656 but also S624, A653, and S660 occurred less frequently. There was no predicted AZM-hERG binding in the model although this does not discount it occurring *in-vivo*.

We also used a validated human ventricular myocyte mathematical model to integrate our ion channel blockade findings in terms of the overall effects on the action potential and intracellular calcium handling. This reproduced the electrophysiological features we observed in whole heart studies that were either explicable in terms of the single channel properties, or calcium handling. Cardiac electrophysiological and arrhythmic properties depend upon surface ion channel function whose combined interactions generate a 'membrane clock' underlying the measurable electrical activity (Fig. 8C)[14]. The modelling reproduced the action potential prolongation and consequent prolonged QT intervals with HCQ and the synergistic effects of combining HCQ with AZM, attributing these to their combined effects on I_{Kr}, I_{Ks} and I_{K1}. The excitation caused by the 'membrane clock' feeds forward into excitation-contraction coupling processes together forming a 'Ca^{2+} clock', which exerts feedback effects at the membrane level. These interactions can have additional modifying effects on action potential conduction, recovery, and post-recovery stability properties, exemplified by Ca^{2+} feedback effects on Na^+ channel and gap junction function (Fig. 8C), as well as longer-term effects on channel expression[13, 26]. The model also predicted an increase in the magnitude and duration of the calcium transient particularly with HCQ+AZM (Fig 8C) and despite only a small change in maximum upstroke velocity due to Na channel blockade, it predicted that slowing of conduction likely also involves calcium dependent gap junction uncoupling.

These results together demonstrate an integrated basis for the electrophysiological and pharmacological effects of HCQ, AZM and their combination, and support recent FDA guidelines cautioning against combined HCQ/AZM administration for the treatment of Covid-19 on the grounds of potential cardiac safety. These effects occurred with the ~10 µM HCQ

concentrations associated with Covid-19 therapy but not with the lower ~1.0 µM HCQ levels more associated with HCQ therapy for rheumatoid arthritis and SLE. Furthermore, they were accentuated by further inclusion of AZM which now produced arrhythmic outcomes. We would therefore strongly recommend monitoring of electrocardiographic QT interval with the use of this combination of medications, particularly in patients with sepsis or SARS where blood gas, electrolyte and haemodynamic disturbances may exacerbate the electrophysiological effects.

Limitations of the present study

However, full clarification of the underlying mode of this action of HCQ, AZM and the HCQ combination will involve future more detailed molecular docking studies on a full hERG structure, completing the currently limited structural information, once this becomes available. Similarly, the detailed clinical applicability of the present experimental results may entail parallel human investigations exploring the corresponding, potentially cardiotoxic, human blood or plasma concentrations and their relationship to therapeutic levels and to corresponding dosing allowing for human pharmacokinetic properties of both HCQ and AZM. In the therapeutic dosing range, HCQ shows large volumes of distribution, likely reflecting its tissue sequestration, rather than its renal clearance from blood (~96 ml/min), accounting for its long plasma half-life around 40 days[27]. AZM similarly accumulates in some tissues reaching concentrations significantly higher than its serum concentrations, though this may contribute to its effective microbial action[28].

Acknowledgments

The authors are grateful to Prof. Lin Wu, Peking University Hospital, for scientific advice with this article.

Sources of Funding

This work is supported by the Kaifeng Excellent Key Laboratory Grant (Grant no. 20190601A) British Heart Foundation (BHF) (PG/14/79/31102, and PG/15/12/31280 and BHF Centres for Research Excellence (CRE) at Cambridge, CLHH, PG/14/80/31106, PG/16/67/32340, ML, PG. FS/17/52/33113,AT) and Oxford (ML) the Chinese Natural Science Foundation (31171085: ML) and EPSRC (United Kingdom) (EP/J00958X/1 and EP/I029826/1) (HZ).

Disclosures

None.

Legends to figures

Figure 1. Effects of HCQ on ion currents attributable to hNav1.5 (I_{Na}, A), Cav1.2 ($I_{Ca,L}$, B), Kv4.3 (I_{to}, C), hERG (I_{Kr}, D), KCNQ1/E1 (I_{Ks}, E) and Kir2.1 (I_{K1}, F). (A-F) Left panels (i): Typical currents in the presence of 0, 1, 10 and 100 µM HCQ. Insets summarize the voltage clamping protocols. Right panels (ii): Mean concentration-response plots constructed for the maximum magnitude of each current (n cells).

Figure 2. Effects of AZM on ion currents attributable to hNav1.5 (I_{Na}, A), Cav1.2 ($I_{Ca,L}$, B), Kv4.3 (I_{to}, C), hERG (I_{Kr}, *D*), KCNQ1/E1 (I_{Ks}, E) and Kir2.1 (I_{K1}, F). (A-F) Left panels (i): Typical currents in the presence of 0, 1, 10 and 100 µM HCQ. Insets summarize the voltage clamping protocols. Right panels (ii): Mean concentration-response plots constructed for the maximum magnitude of each current ((n cells).

Figure 3. Effects of HCQ in combination with 10 µM AZM on ionic currents attributable to hNav1.5 (I_{Na}, A), Cav1.2 ($I_{Ca,L}$, B), Kv4.3 (I_{to}, C), hERG (I_{Kr}, D), KCNQ1/E1 (I_{Ks}, E) and Kir2.1 (I_{K1}, F). (A-F) Left panels (i): Typical currents in the presence of 0, 1, 10 and 100 µM HCQ with 10 µM AZM. Insets summarize the voltage clamping protocols. Right panels (ii): Mean concentration-response plots constructed for the maximum magnitude of each current (n cells).

Figure 4. Computational docking studies. (A) HCQ (ball-and-stick model) docked in hERG structure. Y652 and F656 residues in stick representation colored accordingly to its parent subunit A, B, C or D (red, green, yellow, and blue, respectively). (B) 2-dimensional (2D) diagram depicting interaction types between presented HCQ pose and hERG. Diagram was prepared using Non-bond Interactions Monitor in Discovery Studio Visualizer (Dassault Systèmes, Vélizy-Villacoublay, France).

Figure 5. Multi-electrode array (MEA) mapping and electrocardiographic (ECG) studies in isolated perfused guinea-pig hearts following HCQ and AZM challenge. (A) Schematic summary of stimulation and MEA and ECG recording configuration. (B) Typical left atrial and left ventricular isochronal conduction maps obtained in the presence of 0, 0.01, 0.1, 1.0 and 10 µM HCQ. (C) (i) Successive LV isochronal maps before and following addition of 10 µM HCQ and 10 µM HCQ and 10 µM AZM in combination followed by drug washout. (ii) Quantified conduction velocities through these conditions. (D) ECG studies: (i) ECG records obtained before and following application of 10 µM HCQ, and 10 µM HCQ and 10 µM AZM combined, followed by drug washout. (ii)-(v): corresponding measured (ii) heart rates (HR), (iii) QT and (iv) PR intervals and (v) QRS durations (n hearts) * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Figure 6. Voltage RH237 and Rhod-2 optical mapping in isolated perfused hearts. Comparisons of results before and following challenge by 1 µM and 10 µM HCQ, and 10 µM HCQ combined with either 1 or 10 µM AZM. (A, B) RH237 mapping: (A)(i) Optical mapping and ECG monitoring configuration (ii) Maps of action potential initiation and conduction; (iii) heart rates and (iv) conduction velocities. (B) Action potential time-courses and recoveries: (i) Maps of AP durations at 90% repolarization (APD_{90}). (ii) Representative APs recorded from defined regions of interest. (iii) APD_{90}s averaged over the field of view. (C) Rhod-2 AM (CaD) mapping of ventricular Ca^{2+} transients (CaT): (i) maps of CaD durations at 90% recovery (CaD_{90}). (ii) Comparison of typical CaT transients averaged over defined regions of interest; (iii) CaD_{90} magnitudes; (iv) CaD_{90} dispersions (n hearts). NS not significant, * $p<0.05$, ** $p<0.01$, *** $p<0.001$

Figure 7. Voltage RH237 optical mapping of re-entry and ventricular arrhythmic tendency in isolated perfused hearts. (A) Recording and programmed stimulation configuration. (B) Optical action potential traces obtained under programmed pacing at 190 ms, 180 ms and 170 ms cycle lengths (CLs), before (control) and following addition of 10 µM HCM, and 10 µM HCM with added 10 µM AZM. (C) Corresponding voltage maps obtained at the numbered colour-coded timepoints (n hearts).

Figure 8. In silico modelling of action potentials before and following HCQ and combined HCQ and AZM challenge. (A) Simulated action potentials of human ventricular cell under control conditions and in the presence of HCQ (10 µM) and HCQ (10 µM) + AZM (1 µM) combined. HCQ and AZM when combined produced a synergistic action in prolonging action potential duration (APD$_{90}$), as well as reducing the repolarization rate following the end of phase 0 depolarisation. (B) Effects of HCQ and HCQ + AZM combined on APD$_{90}$ and the maximal upstroke velocity (MUV) of the AP. (C) Illustration of electrophysiological effects and possible mechanisms underlying the pro-arrhythmic effects of applications of HCQ and HCQ and AZM combined at the ion channel, cellular and tissue levels.

References

1.   NIH. NIH clinical trial of hydroxychloroquine, a potential therapy for COVID-19, begins. 2020.

2.   Pharmafield. Novartis sponsors hydroxychloroquine clinical trial in hospitalised COVID-19 patients. 2020.

3.   Gautret P, Lagier J-C, Parola P, Hoang VT, Meddeb L, Mailhe M, Doudier B, Courjon J, Giordanengo V, Vieira VE, Dupont HT, Honoré S, Colson P, Chabrière E, La Scola B, Rolain J-M, Brouqui P and Raoult D. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J of Antimicro Agent*. 2020:105949.

4.   Hinton D. Food and Drug Administration. FDA Emergency use authorization (EUA) of chloroquine and hydroxychloroquine. 2020.

5.   Lenzer J. Covid-19: US gives emergency approval to hydroxychloroquine despite lack of evidence. *BMJ*. 2020;369:m1335.

6.   Borba MGS, Val FFA, Sampaio VS, Alexandre MAA, Melo GC, Brito M, Mourão MPG, Brito-Sousa JD, Baía-da-Silva D, Guerra MVF, Hajjar LA, Pinto RC, Balieiro AAS, Pacheco AGF, Santos JDO, Jr, Naveca FG, Xavier MS, Siqueira AM, Schwarzbold A, Croda J, Nogueira ML, Romero GAS, Bassat Q, Fontes CJ, Albuquerque BC, Daniel-Ribeiro C-T, Monteiro WM, Lacerda MVG and Team ftC-. Effect of High vs Low Doses of Chloroquine Diphosphate as Adjunctive Therapy for Patients Hospitalized With Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infection: A Randomized Clinical Trial. *JAMA Network Open*. 2020;3:e208857-e208857.

7.   Magagnoli J, Narendran S, Pereira F, Cummings T, Hardin JW, Sutton SS and Ambati J. Outcomes of hydroxychloroquine usage in United States veterans hospitalized with Covid-19. *medRxiv*. 2020:2020.04.16.20065920.

8.   Lane JCE, Weaver J, Kostka K, Duarte-Salles T, Abrahao MTF, Alghoul H, Alser O, Alshammari TM, Biedermann P, Burn E, Casajust P, Conover M, Culhane AC, Davydov A, DuVall SL, Dymshyts D, Fernández Bertolín S, Fišter K, Hardin J, Hester L, Hripcsak G, Kent S, Khosla S, Kolovos S, Lambert CG, ver der Lei J, Londhe AA, Lynch KE, Makadia R, Margulis AV, Matheny ME, Mehta P, Morales DR, Morgan-Stewart H, Mosseveld M, Newby D, Nyberg F, Ostropolets A, Park RW, Prats-Uribe A, Rao GA, Reich C, Reps J, Rijnbeek P, Sathappan SMK, Schuemie M, Seager S, Sena A, Shoaibi A, Spotnitz M, Suchard MA, Swerdel J, Torre CO, Vizcaya D, Wen H, de Wilde M, You SC, Zhang L, Zhuk O, Ryan P and Prieto-Alhambra D. Safety of hydroxychloroquine, alone and in combination with azithromycin, in light of rapid wide-spread use for COVID-19: a multinational, network cohort and self-controlled case series study. *medRxiv*. 2020:2020.04.08.20054551.

9.   Mercuro NJ, Yen CF, Shim DJ, Maher TR, McCoy CM, Zimetbaum PJ and Gold HS. Risk of QT Interval Prolongation Associated With Use of Hydroxychloroquine With or Without Concomitant Azithromycin Among Hospitalized Patients Testing Positive for Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol*. 2020. 10.1001/jamacardio.2020.1834

10.   Chorin E, Dai M, Shulman E, Wadhwani L, Bar-Cohen R, Barbhaiya C, Aizer A, Holmes D, Bernstein S, Spinelli M, Park DS, Chinitz LA and Jankelson L. The QT interval in patients with COVID-19 treated with hydroxychloroquine and azithromycin. *Nat Med*. 2020. 10.1038/s41591-020-0888-2

11.   Saleh M, Gabriels J, Chang D, Kim BS, Mansoor A, Mahmood E, Makker P, Ismail H, Goldner B, Willner J, Beldner S, Mitra R, John R, Chinitz J, Skipitaris N, Mountantonakis S and Epstein LM. The Effect of Chloroquine, Hydroxychloroquine and Azithromycin on the Corrected QT Interval in Patients with SARS-CoV-2 Infection. *Circu: Arrhyth Electrophysiol*. doi.org/10.1161/CIRCEP.120.008662

12.   Strauss DG, Gintant G, Li Z, Wu W, Blinova K, Vicente J, Turner JR and Sager PT. Comprehensive In Vitro Proarrhythmia Assay (CiPA) Update from a Cardiac Safety Research Consortium / Health and Environmental Sciences Institute / FDA Meeting. *Therapeutic Innov & Reg Sci*. 2018;53:519-525.

13.   Lei M, Wu L, Terrar DA and Huang CL. Modernized Classification of Cardiac Antiarrhythmic Drugs. *Circulation*. 2018;138:1879-1896.

14.   Lei M and Huang CL-H. Cardiac arrhythmogenesis: a tale of two clocks? *Cardiovasc Res*. 2019. doi.org/10.1093/cvr/cvz283

15.   O'Hara T, Virág L, Varró A and Rudy Y. Simulation of the undiseased human cardiac ventricular action potential: model formulation and experimental validation. *PLoS Comput Biol*. 2011;7:e1002061.

16.   Lai PC, Ho W, Jain N and Walters JD. Azithromycin concentrations in blood and gingival crevicular fluid after systemic administration. *J Periodontol*. 2011;82:1582-6.

17.   Durcan L, Clarke WA, Magder LS and Petri M. Hydroxychloroquine Blood Levels in Systemic Lupus Erythematosus: Clarifying Dosing Controversies and Improving Adherence. *J Rheumatol*. 2015;42:2092-7.

18.   Miller DR, Fiechtner JJ, Carpenter JR, Brown RR, Stroshane RM and Stecher VJ. Plasma hydroxychloroquine concentrations and efficacy in rheumatoid arthritis. *Arthritis Rheum*. 1987;30:567-71.

19.   Capel RA, Herring N, Kalla M, Yavari A, Mirams GR, Douglas G, Bub G, Channon K, Paterson DJ, Terrar DA and Burton R-AB. Hydroxychloroquine reduces heart rate by modulating the hyperpolarization-activated current If: Novel electrophysiological insights and therapeutic potential. *Heart Rhythm*. 2015;12:2186-2194.

20.   Roden DM, Harrington RA, Poppas A and Russo AM. Considerations for Drug Interactions on QTc in Exploratory COVID-19 (Coronavirus Disease 2019) Treatment. *Circulation*. 2020.

21.   January CT and Riddle JM. Early afterdepolarizations: mechanism of induction and block. A role for L-type Ca2+ current. *Circ Res*. 1989;64:977-90.

22.   Herring N, Kalla M and Paterson DJ. The autonomic nervous system and cardiac arrhythmias: current concepts and emerging therapies. *Nat Rev Cardiol*. 2019;16:707-726.

23.   Reiz S and Nath S. Cardiotoxicity of local anaesthetic agents. *Br J Anaesth*. 1986;58:736-46.

24.   Echt DS, Liebson PR, Mitchell LB, Peters RW, Obias-Manno D, Barker AH, Arensberg D, Baker A, Friedman L, Greene HL and et al. Mortality and morbidity in patients receiving encainide, flecainide, or placebo. The Cardiac Arrhythmia Suppression Trial. *N Engl J Med*. 1991;324:781-8.

25.   King JH, Huang CL and Fraser JA. Determinants of myocardial conduction velocity: implications for arrhythmogenesis. *Front Physiol*. 2013;4:154.

26.   Huang CL. Murine Electrophysiological Models of Cardiac Arrhythmogenesis. *Physiol Rev*. 2017;97:283-409.

27.   Tett SE, Cutler DJ, Day RO and Brown KF. A dose-ranging study of the pharmacokinetics of hydroxy-chloroquine following intravenous administration to healthy volunteers. *Br J Clin Pharmacol*. 1988;26:303-13.

28.   Foulds G, Shepard RM and Johnson RB. The pharmacokinetics of azithromycin in human serum and tissues. *J Antimicrob Chemother*. 1990;25 Suppl A:73-82.