Dynamically evolving novel overlapping gene as a factor in the SARS-CoV-2 pandemic

Chase W. Nelson[1,2,a,*], Zachary Ardern[3,a,*], Tony L. Goldberg[4,5], Chen Meng[6], Chen-Hao Kuo[1], Christina Ludwig[6], Sergios-Orestis Kolokotronis[2,7,8], Xinzhu Wei[9,*]

[1]Biodiversity Research Center, Academia Sinica, Taipei, Taiwan
[2]Institute for Comparative Genomics, American Museum of Natural History, New York, NY, USA
[3]Chair for Microbial Ecology, Technical University of Munich, Freising, Germany
[4]Department of Pathobiological Sciences, University of Wisconsin-Madison, Madison, WI, USA
[5]Global Health Institute, University of Wisconsin-Madison, Madison, WI, USA
[6]Bavarian Center for Biomolecular Mass Spectrometry (BayBioMS), Technical University of Munich, Freising, Germany
[7]Department of Epidemiology and Biostatistics, School of Public Health, SUNY Downstate Health Sciences University, Brooklyn, NY, USA
[8]Institute for Genomic Health, SUNY Downstate Health Sciences University, Brooklyn, NY, USA
[9]Departments of Integrative Biology and Statistics, University of California, Berkeley, CA, USA

[a]These authors contributed equally.
*Corresponding authors: cnelson@gate.sinica.edu.tw, zachary.ardern@tum.de, aprilwei@berkeley.edu

1 Abstract

Understanding the emergence of novel viruses requires an accurate and comprehensive

annotation of their genomes. Overlapping genes (OLGs) are common in viruses and have

been associated with pandemics, but are still widely overlooked. We identify and

characterize *ORF3d*, a novel OLG in SARS-CoV-2 that is also present in Guangxi pangolin-

CoVs but not other closely related pangolin-CoVs or bat-CoVs. We then document evidence

of *ORF3d* translation, characterize its protein sequence, and conduct an evolutionary

analysis at three levels: between taxa (21 members of *Severe acute respiratory syndrome-*

related coronavirus), between human hosts (3978 SARS-CoV-2 consensus sequences), and

within human hosts (401 deeply sequenced SARS-CoV-2 samples). *ORF3d* has been

independently identified and shown to elicit a strong antibody response in COVID-19

patients. However, it has been misclassified as the unrelated gene *ORF3b*, leading to

confusion. Our results liken *ORF3d* to other accessory genes in emerging viruses and

highlight the importance of OLGs.

Introduction

The COVID-19 pandemic raises urgent questions about the properties that allow animal

viruses to cross species boundaries and spread within humans. Addressing these questions

requires an accurate and comprehensive understanding of viral genomes. One frequently

overlooked source of novelty is the evolution of new overlapping genes (OLGs), wherein a

single stretch of nucleotides encodes two distinct proteins in different reading frames. Such

"genes within genes" compress genomic information and allow genetic innovation via

overprinting (Keese and Gibbs 1992), particularly as frameshifted sequences preserve

certain physicochemical properties of proteins (Bartonek et al. 2020). However, OLGs also

entail the cost that a single mutation may alter two proteins, constraining evolution of the

pre-existing open reading frame (ORF) and complicating sequence analysis. Unfortunately,

genome annotation methods typically miss OLGs, instead favoring one ORF per genomic

region (Warren et al. 2010).

OLGs are known entities but remain inconsistently reported in viruses of the species

Severe acute respiratory syndrome-related coronavirus (subgenus *Sarbecovirus*; genus

Betacoronavirus; Gorbalenya et al. 2020). For example, annotations of *ORF3b*, *ORF9b*, and

ORF9c are absent or conflicting in SARS-CoV-2 reference genome Wuhan-Hu-1 (NCBI:

NC_045512.2) and genomic studies (e.g., Chan et al. 2020; F. Wu et al. 2020), and OLGs

are often not displayed in genome browsers (e.g., Flynn et al. 2020). Further, *ORF3b* within

ORF3a, an extensively characterized OLG present in other species members including

SARS-CoV (SARS-CoV-1) (McBride and Fielding 2012), has sometimes been annotated in

SARS-CoV-2 even though it contains four early STOP codons in this virus. Such

inconsistencies complicate research, because OLGs may play key roles in the emergence of

new viruses. For example, in human immunodeficiency virus-1 (HIV-1), the novel OLG *asp*

within *env* is actively expressed in human cells (Affram et al. 2019) and is associated with

the pandemic M group lineage (Cassan et al. 2016).

To identify novel OLGs within the SARS-CoV-2 genome, we first generated a list of

candidate overlapping ORFs in the Wuhan-Hu-1 reference genome (NCBI: NC_045512.2).

Specifically, we used the codon permutation method of Schlub et al. (2018) to detect

unexpectedly long ORFs while controlling for codon usage. One unannotated gene

candidate, here named *ORF3d*, scored highly (p=0.0104), exceeding the significance of two

known OLGs annotated in Uniprot (*ORF9b* and *ORF9c*/*ORF14* within *N*;

https://viralzone.expasy.org/8996) (Figure 1, Figure 1—figure supplement 1, and

Supplementary Table 1).

ORF3d comprises 58 codons (including STOP) near the beginning of *ORF3a* (Table

1), making it longer than the known genes *ORF7b* (44 codons) and *ORF10* (39 codons)

(Supplementary Tables 2 and 3). *ORF3d* was discovered independently by Chan et. al

(2020) as '*ORF3b*' and Pavesi (2020) as, simply, 'hypothetical protein'. Due to this naming

ambiguity and its location within *ORF3a*, *ORF3d* has subsequently been conflated with the

previously documented *ORF3b* in multiple studies (Fung et al. 2020; Ge et al. 2020; Gordon

et al. 2020; Hachim et al. 2020; Helmy et al. 2020; Yi et al. 2020). Critically, *ORF3d* is

unrelated (i.e., not homologous) to *ORF3b*, as the two genes occupy different genomic

positions within *ORF3a*: *ORF3d* ends 39 codons upstream of the genome region

homologous to *ORF3b*, and the *ORF3b* start site corresponds to an ORF encoding only 23

codons in SARS-CoV-2 (A. Wu et al. 2020) (Table 1, Figure 1, Figure 1—figure supplement

1, and Supplementary Table 4). Furthermore, the two genes occupy different reading

frames: codon position 1 of *ORF3a* overlaps codon position 2 of *ORF3d* (frame ss12) but

codon position 3 of *ORF3b* (frame ss13). *ORF3d* is also distinct from other OLGs

hypothesized within *ORF3a* (Table 1). Thus, *ORF3d* putatively encodes a novel protein not

present in other *Severe acute respiratory syndrome-related coronavirus* genomes, while the

absence of full-length *ORF3b* in SARS-CoV-2 distinguishes it from SARS-CoV (Figure 1).

Because *ORF3b* plays a central role in SARS-CoV immune interactions, its absence or

truncation in SARS-CoV-2 may be immunologically important (Konno et al. 2020; Yuen et al.

2020).

ORF3d molecular biology and expression

To assess the expression of *ORF3d*, we re-analyzed the SARS-CoV-2 ribosome profiling

(Ribo-seq) data of Finkel et al. (2020). This approach allows the study of gene expression

and reading frame at single nucleotide (nt) resolution by sequencing mRNA fragments

bound by actively translating ribosomes (ribosome footprints) (Ingolia et al. 2009). Focusing

on samples with reads reliably associated with ribosomes stalled at translation start sites

(i.e., lactimidomycin and harringtonine treatments at 5 hours post-infection) and trimming

reads to their 5′ ends (first nt), we observe a consistent peak in read depth ~12 nt upstream

of the start site of *ORF3d* (Figure 2A), the expected ribosomal P-site offset (Calviello and

Ohler 2017). Similar peaks are also observed for previously annotated genes (Figure 2A,

Figure 2—figure supplement 1).

To investigate the relationship between ribosome profiling read depth and expressed

protein levels, we next re-analyzed five publicly available SARS-CoV-2 mass spectrometry

(MS) datasets (Bezstarosti et al. 2020; Bojkova et al. 2020; Davidson et al. 2020; PRIDE

Project PXD018581; Zecha et al. 2020; Methods). We were unable to detect ORF3c,

ORF3d, ORF3b, ORF9c, or ORF10 when employing a 1% false-discovery threshold. This

result may reflect the limitations of MS for detecting proteins that are very short, weakly

expressed under the specific conditions tested, or lack detectable peptides. Even the

envelope protein E is not detected in some SARS-CoV-2 studies (Bojkova et al. 2020;

Davidson et al. 2020), and the only evidence for ORF3b expression in SARS-CoV comes

from Vero E6 cells (McBride and Fielding 2012). However, we do find a strong correlation

between protein expression as estimated from MS and the ribosome profiling read depth

observed at upstream peaks, with r_S=0.89 (p=0.0004, Spearman's rank) (Figure 2, Figure

2—figure supplement 2). This suggests that the presence and depth of an upstream peak is

a reliable indicator of expression. Specifically, results from both proteomic and ribosome

profiling approaches confirm that N, M, and S are the most highly expressed proteins, with N constituting ~80% of the total viral protein content. We also observe that the number of reads determined to be in the correct reading frame (codon position 1) along the gene is moderately correlated with the MS expression values (r_S=0.60, p=0.056), while this is not true for out-of-frame (codon position 2 and 3) reads (r_S=0.39, p=0.237) (Figure 2B, Figure 2—figure supplement 2).

Ribosome profiling reads have a strong tendency to map with their first nucleotide occupying a particular codon position (i.e., in-frame) (Figure 2—figure supplement 3). We therefore examined codon position mapping across *ORF3a* to explore whether the proportions of reads in alternative frames are higher in the *ORF3c*/*ORF3d* region. Analyses were limited to reads of length 30 nt, the expected size of ribosome-bound fragments, which exhibit positioning indicative of the correct reading frame (codon position 1) of annotated genes and the highest total coverage (Figure 2—figure supplement 3, Figure 2—figure supplement 4). Sliding windows reveal a peak in the fraction of reads mapping to the reading frame of *ORF3d* at its hypothesized locus, with this frame's maximum occurring at the center of *ORF3d*, co-located with a dip in the fraction of reads mapping to the frame of *ORF3a* (Figure 2C). These observations are reproducible across treatments (Figure 2—figure supplement 5), robust to sliding window size (Figure 2—figure supplement 6), and similar to the reading frame perturbations observed for the OLG *ORF9b* within *N* (Figure 2—figure supplement 7). At the same time, the *ORF3d* peak stands in contrast to the remainder of *ORF3a*, where the reading frame of *ORF3a* predominates, and smaller peaks in the frame of *ORF3d* are not accompanied by a large dip in the frame of *ORF3a* (Figure 2—figure supplement 8). These observations suggest that *ORF3d* is actively translated. Similar conclusions can be drawn for *ORF3c*, *ORF3d-2*, and *ORF9b* (Figure 2C, Figure 2—figure supplement 7, Figure 2—figure supplement 9).

Additional experiments have also provided evidence of ORF3d translation. Gordon et al. (2020) used overexpression experiments to demonstrate that ORF3d (referred to as

'ORF3b'; Table 1) can be stably expressed and that it interacts with the mitochondrial protein

STOML2. Most compellingly, ORF3d, ORF8, and N elicit the strongest antibody responses

observed in COVID-19 patient sera, with ORF3d sufficient to accurately diagnose the

majority of COVID-19 cases (Hachim et al. 2020). Because *ORF3d* is restricted to SARS-

CoV-2 and pangolin-CoV (see below), this finding is unlikely to be due to cross-reactivity

with another coronavirus, and provides strong independent evidence of *ORF3d* expression

during infection.

Protein sequence properties

To further investigate the antigenic properties of ORF3d, we predicted linear T-cell epitopes

for each SARS-CoV-2 protein. We employed NetMHCpan (Jurtz et al. 2017) for MHC class I

(cytotoxic CD8+ T-cells), an approach shown to accurately predict SARS-CoV-2 epitopes

shared with SARS-CoV (Grifoni et al. 2020), and NetMHCIIpan (Reynisson et al. 2020) for

MHC class II (helper CD4+ T-cells). Specifically, we tested all 9 amino acid (MHC I) or 15

amino acid (MHC II) substrings of each viral protein for predicted weak or strong binding by

MHC. Epitope density was estimated as the mean number of predicted epitopes per residue

for each protein. We also tested two sets of negative controls: (1) randomized peptides

generated from each protein, representing the result expected given amino acid content

alone; and (2) short unannotated ORFs present in the SARS-CoV-2 genome, representing

the result expected for real ORFs that have been evolving without functional constraint.

For CD8+ T cells, the lowest predicted epitope density occurs in ORF3d (1.5 per

residue), which is unexpected given its own amino acid content (p=0.150) and compared to

short unannotated ORFs (p=0.078; permutation tests). The next lowest densities occur in N

and ORF8 (Figure 3A). Intriguingly, as previously mentioned, these three peptides (ORF3d,

N, and ORF8) also elicit the strongest antibody (B-cell epitope) responses measured in

COVID-19 patient sera (Hachim et al. 2020), suggesting a possible balance between CD8+

T and B cell epitopes. For CD4+ T cells, again, ORF3d has one of the lowest predicted

epitope densities (5.6 per residue; p≥0.291), with lower values seen in only three other

genes. However, focusing instead on the shorter ORF3d-2 isoform, this protein contains

zero predicted epitopes, which is a highly significant depletion given its own amino acid

content (p=0.001) and compared to short unannotated ORFs (p=0.001). These observations

suggest either a predisposition towards immune escape, allowing gene survival, or the

action of selective pressures on *ORF3d* or *ORF3d-2* to remove epitopes. In stark contrast to

ORF3d, ORF3c has the highest observed density (8.4 per residue), apparently as a function

of its highly constrained amino acid content (p=0.996) (see below). The enrichment of

predicted epitopes in unannotated proteins (e.g., ORF3c and randomized peptides) but not

N, for which numerous epitopes are annotated at the Immune Epitope Database (Grifoni et

al. 2020), demonstrates that ascertainment or methodological biases cannot account for the

depletion of predicted epitopes in ORF3d.

Our structural prediction for the ORF3d protein suggests *α*-helices connected with

coils and an overall fold model that matches known protein structures (e.g., Protein Data

Bank IDs 2WB7 and 6A93) with borderline confidence (Figure 3—figure supplement 1),

similar to the predictions of Chan et al. (2020). Remarkably, biochemical properties that

influence the structure of this novel protein appear to be inherited from the pre-existing

ORF3a protein sequence encoded by the overlapping alternate reading frame. It has

recently been shown that frame-shifted nucleotide sequences tend to encode proteins with

similar hydrophobicity profiles as a consequence of the standard genetic code (Bartonek et

al., 2020). To explore whether this is the case for *ORF3d*, we used the VOLPES server

(Bartonek and Zagrovic 2019) to calculate the hydrophobicity profiles (Atchley et al. 2005) of

the peptides encoded by all three frames of *ORF3a*. The maximum correlation observed

occurs between the frames of ORF3a (ss11) and ORF3d (ss12) in the region encoding

ORF3d (ORF3a residues 64-102), with r_S=0.87 (p=5.79×10^{-13}, Spearman's rank), much

stronger than the correlation between these frames observed for the non-OLG residues of

ORF3a (r_S=0.27, p=8.70×10^{-4}) (Figure 3C; Figure 3—figure supplement 2). This

conservation of a structure-related property with the ORF3a protein provides further

evidence for *ORF3d* functionality, and may predispose this region towards *de novo* gene

birth. Again, ORF3c shows the opposite pattern, as the minimum correlation observed

between hydrophobicity profiles occurs between the frames of *ORF3a* (ss11) and *ORF3c*

(ss13) in the region encoding ORF3c (ORF3a residues 22-43), with r_S=-0.40 (p=6.16×10^{-2}).

Such disparate hydrophobicity profiles may be due to the strong conservation of *ORF3c* (see

below).

ORF3d taxonomic distribution and origins

To assess the origin of *ORF3d* and its conservation within and among host taxa, we aligned

21 *Severe acute respiratory syndrome-related coronavirus* genomes reported by Lam et al.

(2020) (Supplementary Table 5), limiting our analysis to those with an annotated *ORF1ab*

and no frameshift variants relative to SARS-CoV-2 in the core genes *ORF1ab*, *S*, *ORF3a*, *E*,

M, *ORF7a*, *ORF7b*, and *N* (Supplementary Table 6; SARS-related-CoV_ALN.fasta,

supplementary data). All other genes are also intact (i.e., there is no mid-sequence STOP

codon) in all genomes (Supplementary Tables 7 and 8), with the exception of *ORF3d*,

ORF3b, and *ORF8* (Figure 4). Specifically, *ORF3d* is intact in only 2 sequences: SARS-CoV-

2 Wuhan-Hu-1 and pangolin-CoVs from Guangxi (GX/P5L). Full-length *ORF3b* is intact in

only 3 sequences: SARS-CoV TW11, SARS-CoV Tor2, and bat-CoV Rs7327, with the

remainder falling into two distinct groups sharing an early or late STOP codon, respectively

(Supplementary Table 9). Finally, *ORF8* is intact in all but 5 sequences, where it contains

premature STOP codons or large-scale deletions (Figure 1B).

The presence of intact *ORF3d* homologs among viruses infecting different host

species (human and pangolin) raises the possibility of functional conservation. However, the

taxonomic distribution of this ORF is incongruent with whole-genome phylogenies in that

ORF3d is intact in the pangolin-CoV more distantly related to SARS-CoV-2 (GX/P5L) but not

the more closely related one (GD/1) (Figure 4), a finding confirmed by the alignment of Boni

et al. (2020). New sequence data reveal similarly puzzling trends: *ORF3d* contains STOP

codons in the closely related bat-CoV RmYN02 (GISAID: EPI_ISL_412977; data not shown),

but it is intact in three more distantly related bat-CoVs discovered in Rwanda and Uganda,

where it is further extended by 20 codons (total 78 codons) but shows no evidence of

conservation with SARS-CoV-2 (Wells et al. 2020 and pers. comm; data not shown).

Further, phylogenies of the 21 *Severe acute respiratory syndrome-related coronavirus*

genomes built on *ORF3a* are incongruent with whole-genome phylogenies, likely due to the

presence of recombination breakpoints in *ORF3a* near *ORF3d* (Boni et al. 2020; Rehman et

al. 2020). Recombination, convergence, and recurrent loss may therefore each have played

a role in the origin and taxonomic distribution of *ORF3d*.

Between-taxa divergence

To examine natural selection on *ORF3d*, we measured viral diversity at three hierarchical

evolutionary levels: between-taxa, between-host, and within-host. Specifically, between-taxa

refers to divergence (d) among the 21 aforementioned viruses infecting bat, human, or

pangolin (Figure 1B); between-host refers to diversity (π) between consensus-level SARS-

CoV-2 genomes infecting different human individuals; and within-host refers to π in deeply

sequenced 'intrahost' SARS-CoV-2 samples from single human individuals. At each level,

we inferred selection by estimating mean pairwise nonsynonymous (d_N or π_N; amino acid

changing) and synonymous (d_S or π_S; not amino acid changing) distances among all

sequences. Importantly, we combined standard (non-OLG) methods (Nei and Gojobori 1986;

Nelson et al. 2015) with OLGenie, a new d_N/d_S method tailored for OLGs (hereafter OLG

d_N/d_S), which we previously used to verify purifying selection on a novel OLG in HIV-1

(Nelson et al. 2020).

22   The only gene to show significant evidence of purifying selection at all three

evolutionary levels is *N* (nucleocapsid-encoding gene) (Figure 5), which undergoes

disproportionately low rates of nonsynonymous change ($d_N/d_S < 1$ and $\pi_N/\pi_S < 1$) specifically in

its non-OLG regions, evidencing strict functional constraint. *N* is also the most highly

expressed gene (Figure 2—figure supplement 2), confirming that selection has more

opportunity to act when a protein is manufactured in abundance (Figure 2B; Supplementary

Table 10) (Methods). Importantly, this signal can be missed if non-OLG methods are applied

to *N* without accounting for its internal OLGs, *ORF9b* and *ORF9c* (e.g., at the between-host

level, p=0.0268 when excluding OLG regions, but p=0.411 when including them;

Supplementary Table 11).

With respect to *ORF3d*, comparison of Wuhan-Hu-1 to pangolin-CoV GX/P5L (NCBI:

MT040335.1) yields OLG d_N/d_S=0.14 (p=0.264) (Figure 5), whereas inclusion of a third allele

found in pangolin-CoV GX/P4L (NCBI: MT040333.1) yields 0.43 (p=0.488) (Supplementary

Table 12). Because this is suggestive of constraint, we performed sliding windows of OLG

d_N/d_S across the length of *ORF3a* to check whether the evidence for purifying selection is

specific to the expected genome comparisons and genome positions. Indeed, pairwise

comparisons of each sequence to SARS-CoV-2 reveal OLG d_N/d_S<1 that is specific to the

reading frame, genome positions, and species in which *ORF3d* is intact (pangolin-CoV

GX/P5L) (Figure 6, left). This signal is independent of whether STOP codons are present, so

its consilience with the only intact ORF in this region and species is highly suggestive of

purifying selection. We note that this conclusion does not contradict studies which fail to find

evidence of *ORF3d* conservation when comparing taxa where *ORF3d* is absent (e.g.,

Jungreis et al. 2020), because *ORF3d* is a novel gene and would by definition lack such

conservation. This contrastive signal is also similar to what is observed for the known OLGs

ORF3b (in comparisons to SARS-CoV; Figure 6, right), and *ORF9b* and *ORF9c* (in both

viruses; Figure 6—figure supplement 1).

Between-host evolution and pandemic spread

Purifying selection can be specific to just one taxon, as in the case of novel genes. Thus, to

measure selection within SARS-CoV-2 only, we obtained *n*=3,978 high-quality human

SARS-CoV-2 consensus sequences from GISAID (accessed April 10, 2020; Supplementary

Table 13; GISAID Acknowledgments Table, supplementary data). Between-host diversity

was sufficient to detect marginally significant purifying selection across all genes

(π_N/π_S=0.50, p=0.0613, *Z*-test; Figure 5—figure supplement 2) but not most individual genes

(Figure 5). Therefore, we instead investigated single mutations over time, limiting to 27 high-frequency variants (minor allele frequency ≥2%; Supplementary Table 14).

One high-frequency mutation occurred in *ORF3d*: G25563U, here denoted *ORF3d*-LOF (*ORF3d*-loss-of-function), which causes a STOP codon in *ORF3d* (ORF3d-E14*) but a nonsynonymous change in *ORF3a* (ORF3a-Q57H). This mutation is not observed in any other species member included in our analysis (Figure 4; SARS-related-CoV_ALN.fasta, supplementary data). During the first months of the COVID-19 pandemic, *ORF3d*-LOF increased in frequency (Supplementary Table 15) in multiple locations (Figure 5—figure supplement 2D; Supplementary Table 16), making this mutation a candidate for natural selection on ORF3a, ORF3d, or both (Kosakovsky-Pond 2020). However, temporal allele frequency trajectories (Figure 5—figure supplement 2D) and similar signals from phylogenetic branch tests may also be caused by founder effects or genetic drift, and are susceptible to ascertainment bias (e.g., preferential sequencing of imported infections and uneven geographic sampling) and stochastic error (e.g., small sample sizes).

To partially account for these confounding factors, we instead constructed the mutational path leading from the SARS-CoV-2 haplotype collected in December 2019 to the haplotype carrying *ORF3d*-LOF. This path involves five mutations (C241U, C3037U, C14408U, A23403G, G25563U), constituting five observed haplotypes (EP–3 → EP–2 → EP → EP+1 → EP+1+LOF), shown in Table 2. Here, EP is suggested to have driven the European Pandemic (detected in German patient #4; see footnote 3 of Table 2; Korber et al. 2020; Rothe et al. 2020); EP–3 is the Wuhan founder haplotype; EP–1 is never observed in our dataset; and LOF refers to *ORF3d*-LOF. We then documented the frequencies and earliest collection date of each haplotype (Table 2) to determine whether *ORF3d*-LOF occurred early on the EP background.

Surprisingly, despite its expected predominance in Europe due to a founder effect, the EP haplotype is extremely rare. By contrast, haplotypes with one additional mutation (C14408U; RdRp-P323L) on the EP background are common in Europe, and *ORF3d*-LOF occurred very early on this background to create EP+1+LOF from EP+1. Neither of these

two haplotypes was initially observed in China (Table 2), suggesting that they might have

arisen in Europe. Thus, we further partitioned the samples into two groups, corresponding to

countries with early founders (January samples) or only late founders (no January samples)

(Figure 7). In the early founder group, EP–3 (Wuhan) is the first haplotype detected in most

countries, consistent with most early COVID-19 cases being related to travel from Wuhan.

Because this implies that genotypes EP–3 and EP had longer to spread in the early founder

group, it is surprising that their spread is dwarfed by the increase of EP+1 and EP+1+LOF

starting in late February. This turnover is most evident in the late founder group, where

multiple haplotypes are detected in a narrow time window, and the number of cumulative

samples is always dominated by EP+1 and EP+1+LOF. Thus, while founder effects and drift

are plausible explanations, it is also worth investigating whether the early spread of *ORF3d*-

LOF may have been caused by its linkage with another driver, either C14408U (+1 variant)

or a subsequent variant(s) occurring on the EP+1+LOF background (Discussion).

Within-host diversity and mutational bias

Examination of within-host variation across multiple samples allows detection of recurrent

mutations, which might indicate mutation bias or within-host selection. To investigate these

possibilities, we obtained *n*=401 high-depth 'intrahost' human SARS-CoV-2 samples from

the Sequence Read Archive (Supplementary Table 17) and called SNPs relative to the

Wuhan-Hu-1 reference genome. Within human hosts, 42% of SNPs passed our false-

discovery rate criterion (Methods), with a median passing minor allele frequency of 2% (21 of

1344 reads). Using these variants to estimate within-host diversity, *ORF3d* does not show

significant evidence of selection, with OLG π_N/π_S= 1.5 (p=0.584) (Figure 5). We also

examined six high-depth samples of pangolin-CoVs from Guangxi, but no conclusions could

be drawn due to low sequence quality (Methods; Supplementary Table 18).

To identify recurrent mutations, we next limited to samples for which the major allele

was also ancestral at each site (Wuhan-Hu-1 genotype). At such sites, precluding

coinfection by multiple genotypes, minor alleles occurring in more than one sample are

expected to be derived and recurrent (i.e., identical by state but not descent). For *ORF3d*,

we observe *ORF3d*-LOF as a minor allele in two (0.9%) of 220 samples (SRR11410536 and

SRR11479046 at frequencies of 1.9% and 4.6%, respectively). This proportion of samples

with a recurrent mutation is high but not unusual, as 1.8% of possible genomic changes

have an equal or higher proportion of recurrence. Additionally, no mutations in *ORF3d* recur

in >2.5% of samples (Figure 8). Thus, we find no evidence that within-host selection or

mutation pressure are involved in the spread of *ORF3d*-LOF. However, a small number of

other loci exhibit high rates of recurrent mutation, with five mutations independently

observed in >10% of samples (Methods; Figure 8). Surprisingly, another STOP mutation

(A404U; NSP1-L47*) is observed at low frequency in the majority of samples (Figure 8,

Figure 8—figure supplement 1), unexplainable by mutational bias and warranting

investigation.

Discussion

Our analyses provide strong evidence that SARS-CoV-2 contains a novel overlapping gene

(OLG), *ORF3d*, that has not been consistently identified or fully analyzed before this study.

The annotation of a newly emerged virus is difficult, particularly in genome regions subject to

frequent gene gain and loss. Moreover, due to the inherent difficulty of studying OLGs, they

tend to be less carefully documented than non-OLGs; for example, *ORF9b* and *ORF9c* are

still not annotated in the most-used reference genome, Wuhan-Hu-1 (last accessed

September 26, 2020). Therefore, *de novo* and homology-based annotation are both

essential, followed by careful expression analyses using multi-omic data and evolutionary

analyses within and between species. In particular, we emphasize the importance of using

whole-gene or genome alignments when inferring homology for both OLGs and non-OLGs,

taking into account genome positions and all reading frames. Unfortunately, in the case of

SARS-CoV-2, the lack of such inspections has led to mis-annotation and a domino effect.

For example, homology between *ORF3b* (SARS-CoV) and *ORF3d* (SARS-CoV-2) has been

implied, leading to unwarranted inferences of shared functionality and subsequent claims of

homology between *ORF3b* and other putative OLGs within *ORF3a* (Table 1). Given the rapid

growth of the SARS-CoV-2 literature, it is likely this mistake will be further propagated. We

therefore provide a detailed annotation of Wuhan-Hu-1 protein-coding genes and codons in

Supplementary Tables 2 and 3, respectively, as a resource for future studies.

Our study highlights the highly dynamic process of frequent losses and gains of

accessory genes in the species *Severe acute respiratory syndrome-related coronavirus*, with

the greatest functional constraint typically observed for the most highly expressed genes

(Figure 5—figure supplement 1). With respect to gene loss, while many accessory genes

may be dispensable for viruses in cell culture, they often play an important role in natural

hosts (Forni et al. 2017). Thus, their loss may represent a key step in adaptation to new

hosts after crossing a species barrier (Gorbalenya et al. 2006). For example, the absence of

full-length *ORF3b* in SARS-CoV-2 has received attention from only a few authors (e.g.,

Lokugamage et al. 2020), even though it plays a central role in SARS-CoV infection and

early immune interactions as an interferon antagonist (Kopecky-Bromberg et al. 2007), with

effects modulated by ORF length (Zhou et al. 2012). Thus, the absence or truncation of

ORF3b in SARS-CoV-2 may be immunologically important (Yuen et al. 2020), e.g., in the

suppression of type I interferon induction (Konno et al. 2020). With respect to gene gain, the

apparent presence of the *ORF3d* coincident with the inferred entry of SARS-CoV-2 into

humans from a hitherto undetermined reservoir host suggests that this gene is functionally

relevant for the emergent properties of SARS-CoV-2, analogous to *asp* for HIV-1-M (Cassan

et al. 2016). The mechanisms and detailed histories of gene birth, neo-functionalization, and

survival of novel genes remain unclear. However, our findings that ORF3d remarkably

inherited its hydrophobicity profile from the overlapping region of ORF3a, that ORF3d is

depleted for predicted CD8+ and CD4+ T cell epitopes, and that *ORF3d* exhibits potential

purifying selection between SARS-CoV-2 and pangolin-CoV GX/P5L, provide strong leads

for further research. Indeed, both the structural and immunogenic properties of new viral

proteins deserve further attention and are important parts of their evolutionary story.

Failure to account for OLGs can also lead to erroneous inference of (or failure to

detect) natural selection in known genes, and *vice versa*. For example, a synonymous

variant in one reading frame is very likely to be nonsynonymous in a second overlapping

frame. As a result, purifying selection in the second frame usually lowers d_S (raise d_N/d_S) in

the first frame, increasing the likelihood of mis-inferring positive selection (Holmes et al.

2006; Sabath et al. 2008; Nelson et al. 2020). Such errors could, in turn, lead to

mischaracterization of the genetic contributions of OLG loci to important viral properties such

as incidence and persistence. One potential consequence is misguided countermeasure

efforts, e.g., through failure to detect functionally conserved or immunologically important

regions.

Although our study focuses on *ORF3d*, other OLGs have been proposed in SARS-

CoV-2 and warrant investigation. *ORF3d-2*, a shorter isoform of *ORF3d* (Table 1), shows

evidence of higher overall expression than *ORF3d* (Figure 2A). However, analysis of full-

length *ORF3d* is confounded by its 20 codon triple overlap with both *ORF3a* and *ORF3c*,

making expression and selection results difficult to compare between the two isoforms.

Although not evolutionarily novel, the recently discovered *ORF3c* (Table 1) shows deep

conservation among viruses of this species (Cagliana et al. 2020; Firth 2020; Jungreis et al.

2020), the lowest π_N/π_S ratio observed in our between-host analysis (Figure 5), strong

evidence for translation in SARS-CoV-2 (Figure 2), and the highest predicted CD8+ T cell

epitope density (Figure 3). Within *S*, *S-iORF2* (genome positions 21768-21863) also shows

evidence of translation from ribosome profiling (Finkel et al. 2020), and between-host

comparisons suggest purifying selection (π_N/π_S=0.22, p=0.0278) (Table 1). We therefore

suggest that the SARS-CoV-2 genome could contain additional undocumented or new

OLGs.

Our comprehensive evolutionary analysis of the SARS-CoV-2 genome demonstrates

that many genes are under relaxed purifying selection, consistent with the exponential

growth of the virus (Gazave et al. 2013). At the between-host level, nucleotide diversity

increases somewhat over the initial period of the COVID-19 pandemic, tracking the number

of locations sampled, while the π_N/π_S ratio remains relatively constant at 0.46 (±0.030 SEM)

(Figure 5—figure supplement 2B). Other genes differ in the strength and direction of

selection at the three evolutionary levels, e.g. *ORF9c* (Figure 5), suggesting a shift in

function or importance over time or between different host species or individuals. *ORF3d*

and *ORF8* are both among the youngest genes in SARS-CoV-2, being taxonomically

restricted to a subset of betacoronaviruses (Cui et al. 2019), and both exhibit high rates of

change and turnover (Figure 1; Figure 5; SARS-related-CoV_ALN.fasta, supplementary

data). High between-host π_N/π_S was also observed in *ORF8* of SARS-CoV, perhaps due to a

relaxation of purifying selection upon entry into civet cats or humans (Forni et al. 2017).

However, *ORF3d* and *ORF8* both exhibit strong antibody (B-cell epitope) responses (Hachim

et al. 2020) and predicted T-cell epitope depletion (Figure 3) in SARS-CoV-2. This highlights

the important connection between evolutionary and immunologic processes (Daugherty and

Malik 2012), as antigenic peptides may impose a fitness cost for the virus by allowing

immune detection. Thus, the loss or truncation of these genes may share an immunological

basis and deserves further attention.

The quick expansion of *ORF3d*-LOF (EP+1+LOF) and its haplotype (EP+1) during

this pandemic is surprising, given that a founder effect would have favored other variants

that arrived earlier. However, newer data show that *ORF3d*-LOF has remained at relatively

low frequencies compared to EP+1 variants (Kosakovsky-Pond 2020), and plausible

explanations exist which do not require invoking natural selection. First, the slower 'spread'

of EP–3 and EP may be an artifact of a sampling policy bias in case isolation: testing and

quarantine were preferentially applied to travellers who had recently visited Wuhan, which

may have led to selective detection, isolation, and tracing of these haplotypes rather than

mutations occurring within Europe (e.g., C14408U and G25563U). However, the EP+1 and

EP+1+LOF haplotypes also increase faster in the late founder group, where it is unclear

which haplotype was more travel-related. Second, it is possible that the EP haplotype was

effectively controlled, while EP+1 was introduced independently (Worobey et al. 2020).

Third, although G25563U itself simultaneously causes ORF3a-Q57H and *ORF3d*-LOF, its

spread seems to be linked with RdRp-P323L, and both mutations are linked with Spike-

D614G, a variant with predicted functional relevance (Korber et al. 2020; Yurkovetskiy et al.

2020). Thus, it is possible that G25563U is neutral while a linked variant(s) is under

selection. These observations highlight the necessity of empirically evaluating the effects of

these mutations and their interactions. Lastly, because only five major polymorphisms are

considered in this analysis (Supplementary Table 14), it is possible that the spread of

ORF3d-LOF or other haplotypes was further assisted or hindered by subsequent mutations.

Our study has several limitations. First, we were not able to confirm the translation of

ORF3d using mass spectrometry (MS). This may be due to several reasons: (1) ORF3d is

short, and tryptic digestion of ORF3d generates only two peptides potentially detectable by

MS; (2) ORF3d may be expressed at low levels; (3) the tryptic peptides derived from ORF3d

may not be amenable to detection by MS even under the best possible conditions, as

suggested by its relatively low MS intensity even in an overexpression experiment ('ORF3b'

in Gordon et al. 2020); and (4) hitherto unknown post-translational modifications of ORF3d

could also prohibit detection. Other possibilities for validation of ORF3d include MS with

other virus samples; affinity purification MS; fluorescent tagging and cell imaging; Western

blotting; and the sequencing of additional genomes in this viral species, which would

potentiate more powerful tests of purifying selection and a better understanding of the history

and origin of *ORF3d*. With respect to between-host diversity, we focused on consensus-level

sequence data; however, this approach can miss important variation (Holmes 2009),

stressing the importance of deeply sequenced within-host samples using technology

appropriate for calling within-host variants (Grubaugh et al. 2019). As we use Wuhan-Hu-1

for reference-based read mapping and remove duplicate reads as possible PCR artifacts,

reference bias (Degner et al. 2009) or bias against natural duplicates at high read depth loci

(Zhou et al. 2014) could potentially affect our within-host results. Additionally, we detected

natural selection using counting methods that examine all pairwise comparisons between or

within specific groups of sequences, which may have less power than methods that trace

changes over a phylogeny. However, this approach is robust to errors in phylogenetic and

ancestral sequence reconstruction, and to artifacts due to linkage or recombination (Hughes

et al. 2006; Nelson and Hughes 2015). Additionally, although our method for measuring

selection in OLGs does not explicitly account for mutation bias, benchmarking with other

viruses suggests inference of purifying selection is conservative (Nelson et al. 2020). Finally,

given multiple recombination breakpoints in *ORF3a* (Boni et al. 2020) and the relative

paucity of sequence data for viruses closely related to SARS-CoV-2, our analysis could not

differentiate between convergence, recombination, or recurrent loss in the origin of *ORF3d*.

In conclusion, OLGs are an important part of viral biology and deserve more

attention. We document several lines of evidence for the expression and functionality of a

novel OLG in SARS-CoV-2, *ORF3d*, and compare it to other hypothesized OLG candidates

in *ORF3a*. Finally, as a resource for future studies, we provide a detailed annotation of the

SARS-CoV-2 genome and highlight mutations of potential relevance to the within- and

between-host evolution of SARS-CoV-2.

Methods

Data and software

Supplementary scripts and documentation are freely available on GitHub at

https://github.com/chasewnelson/SARS-CoV-2-ORF3d. Supplementary data not included in main

figure source data are freely available on Zenodo at https://zenodo.org/record/4052729.

Genomic features and coordinates

All genome coordinates are given with respect to the Wuhan-Hu-1 reference sequence (NCBI:

NC_045512.2; GISAID: EPI_ISL_402125) unless otherwise noted. SARS-CoV (SARS-CoV-1)

genome coordinates are given with respect to the Tor2 reference sequence (NC_004718.3). SARS-

CoV-2 Uniprot peptides were obtained from https://viralzone.expasy.org/8996, where *ORF9c* is

referred to as *ORF14*. Nucleotide sequences were translated using R::Biostrings (Lawrence et al.

2013), Biopython (Cock et al. 2009), or SNPGenie (Nelson et al. 2015). Alignments were viewed and

edited in AliView v1.20 (Larsson 2014). To identify OLGs using the codon permutation method of

Schlub et al. (2018), all 12 ORFs annotated in the Wuhan-Hu-1 reference genome were used as a

reference (NCBI=NC_045512.2; *ORF1a*, *ORF1b*, *S*, *ORF3a*, *E*, *M*, *ORF6*, *ORF7a*, *ORF7b*, *ORF8*, *N*,

and *ORF10*) (Figure 1; Supplementary Tables 1-3). Only forward-strand (same-strand; sense-sense)

OLGs were considered in this analysis.

SARS-CoV-2 genome data, alignments, and between-host analyses

SARS-CoV-2 genome sequences were obtained from GISAID on April 10, 2020 (GISAID

Acknowledgments Table, supplementary data). Whole genomes were aligned using MAFFT v7.455

(Katoh and Standley 2013), and subsequently discarded if they contained internal gaps (-) located

>900 nt from either terminus, a length sufficient to exclude sequences with insertions or deletions

(indels) in protein-coding regions. A total of 3,978 sequences passed these filtering criteria, listed in

Supplementary Table 13. Coding regions were identified using exact or partial homology to SARS-

CoV-2 or SARS-CoV annotations. To quantify the diversity and evenness of sample locations, we

quantified their entropy as $-\sum p*\ln(p)$, where p is the number of distinct (unique) locations or countries

reported for a given window (Ewens and Grant 2001) (Supplementary Table 19).

 Severe acute respiratory syndrome-related coronavirus genome data and alignments

Severe acute respiratory syndrome-related coronavirus genome IDs were obtained from Lam et al.

(2020) and downloaded from GenBank or GISAID. Genotype Wuhan-Hu-1 was used to represent

SARS-CoV-2. Except for pangolin-specific analyses, isolates GX/P5L (NCBI: MT040335.1; GISAID:

EPI_ISL_410540) and GD/1 (GISAID: EPI_ISL_410721) were used to represent pangolin-COVs;

GD/1 was chosen as a representative because the other Guangdong (GD) sequence lacks the *S*

gene and contains 27.76% Ns, while GX/P5L was chosen because it is one of two high-coverage

Guangxi (GX) sequences derived from lung tissue that also contains no Ns. Other sequences were

excluded if they lacked an annotated *ORF1ab* with a ribosomal slippage, or contained a frameshift

indel in any gene (Supplementary Table 6), leaving 21 sequences for analysis (Supplementary Table

5).

To produce whole-genome alignments, we first aligned all genome sequences using MAFFT.

Then, coding regions were identified using exact or partial sequence identity to SARS-CoV-2 or

SARS-CoV annotations, translated, and individually aligned at the amino acid level using ProbCons

v1.12 (Do et al. 2005). In the case of OLGs, the longest (reference) protein was used (e.g., N rather

than ORF9b or ORF9c). Amino acid alignments were then imposed on the coding sequence of each

gene using PAL2NAL v14 (Suyama et al. 2006) to maintain intact codons. Finally, whole genomes

were manually shifted to match the individual codon alignments in AliView. Codon breaks were

preferentially resolved to align S/Q/T at 3337-3339, and L/T/I at 3343-3345. This preserved all

nucleotides of each genome while concurrently producing codon-aware alignments. The alignment is

available in the supplementary data as SARS-related-CoV_ALN.fasta, where pangolin-CoV GD/1 has

been removed because it is only available from GISAID (i.e., permission is required for data access).

Phylogenetic relationships among isolates were explored using maximum likelihood

phylogenetic inference, as implemented in IQ-tree (Nguyen et al. 2015), using the generalized time-

reversible (GTR; Tavaré 1986) substitution model combined with the FreeRate model (Soubrier et al.

2012) to account for among-site rate heterogeneity. Trees were rooted *a priori* following Lam et al.

(2020).

Proteomics analysis

We used MaxQuant (Tyanova et al. 2016) to re-analyze five publicly available SARS-CoV-2 mass

spectrometry (MS) datasets: Bezstarosti et al. 2020 (PRIDE accession PXD018760); Bojkova et al.

2020 (PXD017710); Davidson et al. 2020 (PXD018241); PRIDE Project PXD018581; and Zecha et al.

2020 (PXD019645). However, peptide spectrum matches for ORF3d (possible tryptic peptides

CTSCCFSER and FQNHNPQK) did not pass our 1% false-discovery threshold in any of these

datasets, and ORF3d-2 (Table 1) does not encode any peptides detectable by this method. ORF3c,

ORF9c, and ORF10 could also not be reliably detected. For peptides that were successfully detected

in the datasets of Bezstarosti et al. (2020) and Davidson et al. (2020), protein concentrations were

estimated using intensity-based absolute quantification (iBAQ) (Schwanhäusser et al. 2011). iBAQ

values were computed using the Max-Quant software (Cox and Mann 2008) as the sum of all peptide

intensities per protein divided by the number of theoretical peptides per protein. Thus, iBAQ values

are proportional estimates of the molar protein quantity of a protein in a given sample, allowing

relative quantitative comparisons of protein expression levels.

Ribosome profiling analysis

The 16 ribosome profiling (Ribo-seq) datasets of Finkel et al. (2020) using SARS-CoV-2 infected Vero

E6 cells were downloaded from the Sequence Read Archive (accession numbers SRR11713354 to

SRR11713369). These samples comprise four treatments for ribosome stalling: (1) lactimidomycin

(LTM) or (2) harringtonine (Harr), which are biased towards stalling at translation initiation sites; (3)

cyclohexamide (CHX), which stalls along a gene during active translation; and (4) mRNA, which

serves as a control (i.e., total RNA content rather than ribosome footprints). Two time points are

represented: (1) 5 hours and (2) 24 hours post infection (hpi), labelled "05hr" and "4hr", respectively,

in the SRA data. The FASTQ format reads were mapped to the Wuhan-Hu-1 reference genome using

Bowtie2 local alignment (Langmead et al. 2019), with a seed length of 20 and up to one mismatch

allowed, after substituting the isolate's mutations, as listed in Finkel et al. (2020). Mapped reads were

then classified by read length and trimmed to the first (5′-end) nucleotide for downstream analyses.

NetMHCpan T-cell epitope analysis

For predicted MHC class I binding, viral protein sequences were analyzed in 9 aa substrings using

NetMHCpan4.0 (Jurtz et al. 2017). Twelve (12) representative HLA class I alleles (Sidney et al. 2008)

were tested: HLA-A*01:01 (A1), HLA-A*02:01 (A2), HLA-A*03:01 (A3), HLA-A*24:02 (A24), HLA-

A*26:01 (A26), HLA-B*07:02 (B7), HLA-B*08:01 (B8), HLA-B*27:05 (B27), HLA-B*39:01 (B39), HLA-

B*40:01 (B44), HLA-B*58:01 (B58), and HLA-B*15:01 (B62). NetMHCpan4.0 returns percentile ranks

that characterize a peptide's likelihood of antigen presentation compared to a set of random natural

peptides. We employed the suggested threshold of 2% to determine potential presented peptides, and

0.5% to identify strong MHC class I binder. Both strong and weak binders were considered predicted

epitopes. For predicted MHC class II binding, viral protein sequences were analyzed in 15 aa

substrings using NetMHCIIpan4.0 (Reynission et al. 2020). Twenty six (26) representative HLA class

II alleles (Greenbaum et al. 2011; Paul et al. 2016) were tested: HLA-DPA1*0201-DPB1*0401, HLA-

DPA1*0103-DPB1*0201, HLA-DPA1*0201-DPB1*0101, HLA-DPA1*0201-DPB1*0501, HLA-

DPA1*0301-DPB1*0402, HLA-DQA1*0101-DQB1*0501, HLA-DQA1*0102-DQB1*0602, HLA-

DQA1*0301-DQB1*0302, HLA-DQA1*0401-DQB1*0402, HLA-DQA1*0501-DQB1*0201, HLA-

DQA1*0501-DQB1*0301, DRB1*0101, DRB1*0301, DRB1*0401, DRB1*0405, DRB1*0701,

DRB1*0802, DRB1*0901, DRB1*1101, DRB1*1201, DRB1*1302, DRB1*1501, DRB3*0101,

DRB3*0202, DRB4*0101, DRB5*0101. NetMHCIIpan4.0 returns percentile ranks that characterize a

percentile rank of eluted ligand prediction score. We employed the suggested threshold of 10% to

determine potential presented peptides, and 2% to identify strong MHC class II binder. Both strong

and weak binders were considered predicted epitopes.

Hydrophobicity profile analysis

The sequence of *ORF3a* was translated in all three forward-strand reading frames and uploaded to

the VOLPES server (http://volpes.univie.ac.at/; Bartonek and Zagrovic 2019) to determine

hydrophobicity profiles using the unitless scale "FAC1" (Factor 1; Atchley et al. 2005) with a window

size of 25 codons. The correlations between the profiles for the reading frames were calculated for

peptides encoded by the following subregions of *ORF3a*, from 5′ to 3′: (1) *ORF3a*, codons not

overlapping any known or hypothesized overlapping genes; (2) *ORF3a*/*ORF3c*, encoding residues of

both *ORF3a* and *ORF3c*; (3) *ORF3a*/*ORF3c*/*ORF3d*, encoding residues of *ORF3a*, *ORF3c*, and

ORF3d; (4) *ORF3a*/*ORF3d*, encoding residues of both *ORF3a* and *ORF3d*; and (5) *ORF3a*/*ORF3b*,

encoding residues of both *ORF3a* and *ORF3b*. Spearman's rank correlation was computed using

stats::cor.test() with method='spearman' in R, treating mean 25-residue window values as the

observational unit.

1. *Statistics and tests of natural selection*

2. All p-values reported in this study are two-tailed. Statistical and data analyses and visualization were

3. carried out in R v3.5.2 (R Core Team 2018) (libraries: boot, feather, ggrepel, patchwork,

4. RColorBrewer, scales, tidyverse), Python (BioPython, pandas) (McKinney 2010), Microsoft Excel,

5. Google Sheets, and PowerPoint. Colors were explored using Coolors (https://coolors.co). Copyright-

6. free images of a bat, human, and pangolin were obtained from Pixabay (https://pixabay.com). All

7. d_N/d_S or π_N/π_S ratios were estimated for non-OLG regions using SNPGenie scripts snpgenie.pl or

8. snpgenie_within_group.pl (Nelson et al. 2015; https://github.com/chasewnelson/SNPGenie), and for

9. OLG regions using OLGenie script OLGenie.pl (Nelson et al. 2020;

10. https://github.com/chasewnelson/OLGenie). All OLG d_N/d_S (π_N/π_S) estimates refer to d_{NN}/d_{SN} (π_{NN}/π_{SN})

11. for the reference frame and d_{NN}/d_{NS} (π_{NN}/π_{NS}) for the alternate frame (ss12 or ss13), as described in

12. Nelson et al. (2020), because the number of SS (synonymous/synonymous) sites was insufficient to

13. estimate d_{SS} (π_{SS}). The null hypothesis that $d_N-d_S=0$ ($\pi_N-\pi_S=0$) was evaluated using both Z and

14. achieved significance level (ASL) tests (Nei and Kumar 2000) with 10,000 and 1,000 bootstrap

15. replicates for genes and sliding windows, respectively, using individual codons (alignment columns)

16. as the resampling unit (Nei and Kumar 2000). The main text reports only Z-test results, because this

17. test was used to benchmark OLGenie (Nelson et al. 2020) and appears to be more conservative. For

18. ASL, p-values of 0 were reported as the lowest non-zero value possible given the number of bootstrap

19. replicates. Benjamini-Hochberg (Benjamini and Hochberg 1995) or Benjamini-Yekutieli (Benjamini

20. and Yekutieli 2001) false-discovery rate corrections (Q-values) were used for genes (independent

21. regions) and sliding windows (contiguous overlapping regions), respectively.

22. *Between-species analyses*

23. Because uncorrected d values >0.1 were observed in between-taxa comparisons, a Jukes-Cantor

24. correction (Jukes and Cantor 1969) was applied to d_N and d_S estimates. For each ORF, sequences

25. were only used to estimate d_N/d_S if a complete, intact ORF (no STOPs) was present. Most notably, for

26. *ORF3b*, only the region corresponding to the first ORF in SARS-CoV-2 was analyzed, and these sites

27. were also considered an OLG region of *ORF3a*. The remaining 3'-proximal portion of *ORF3a* (codons

28. 165-276) was considered a non-OLG region, and only genomes lacking the full-length *ORF3b* (i.e.,

29. those with mid-sequence STOP codons) were included in these estimates. Similarly, for *ORF3d*, only

30. SARS-CoV-2 and pangolin-CoV GX/P5L were analyzed for Figure 5. For this analysis, *ORF3a* codon

71 in SARS-CoV-2 (CTA) differed by two nucleotides from the same codon in pangolin-CoV GX/P5L

(TTT), resulting in a multi-hit approximation being employed by the OLGenie method (Nelson et al.

2020). Thus, for greater accuracy, we instead estimated changes in this codon using the method of

Wei and Zhang (2015) across two mutational pathways, which yields 0.5

nonsynonymous/nonsynonymous, 0.5 nonsynonymous/synonymous, and 1

synonymous/nonsynonymous changes. The region occupied by the triple overlap of

ORF3a/*ORF3c*/*ORF3d* (*ORF3a* codons 44-64; Table 1) was excluded from analysis for all three

genes. The following codons (or their homologous positions) were also excluded from all analyses: all

codons occupying a non-OLG/OLG boundary; codons 4460-4466 of *ORF1ab*, which constitute either

nsp11 or *nsp12* depending on a ribosomal frameshift; codons 1-13 of *E*, which overlap *ORF3b* in

some genomes; codons 62-64 of *ORF6*, which follow a premature STOP in some genomes; codons

122-124 of *ORF7a* and 1-3 of *ORF7b*, which overlap each other; and codons 72-74 of *ORF9c*, which

follow a premature STOP in some genomes.

Cumulative haplotype frequencies

We defined haplotypes along the mutational path to EP+1+LOF using all five high-derived allele

frequency (DAF) mutations from Wuhan-Hu-1 to *ORF3d*-LOF. Subsequent mutations after *ORF3d*-

LOF were ignored in the haplotype analysis. Samples with missing data at any of the five loci were

removed from the haplotype analysis. We calculated the cumulative frequency of each haplotype in

Germany (where the EP haplotype is a documented founder) and five other countries with the most

abundant samples by the time of data accession. Cumulative frequencies were calculated as the total

number of occurrences of each haplotype collected each day, divided by the total number of samples

from the same country. Countries were subsequently divided into early founders and late founders to

investigate founder effects, where early founder countries tend to have several samples from January,

and late founder countries tend to have samples collected only after mid-February.

Within-host diversity

For within-host analyses, we obtained *n*=401 high-depth (at least 50-fold mean coverage) human

SARS-CoV-2 samples from the Sequence Read Archive (listed in Supplementary Table 17). Only

Illumina samples were used, as some Nanopore samples exhibit apparent systematic bias in calling

putative intrahost SNPs, and this technology has also been shown to be unsuitable for intra-host

analysis (Grubaugh et al. 2019). Reads were trimmed with BBDUK (Bushnell B. 2017. BBTools.

https://jgi.doe.gov/data-and-tools/bbtools/) and mapped against the Wuhan-Hu-1 reference sequence

using Bowtie2 (Langmead and Salzberg 2012) with local alignment, seed length 20, and up to 1

mismatch. SNPs were called from mapped reads using the LoFreq (Wilm et al. 2012) variant caller

with sequencing quality and MAPQ both at least 30. Only single-end or the first of paired-end reads

were used. Variants were dynamically filtered based on each site's coverage using a binomial cutoff

to ensure a false-discovery rate of ≤1 within-host variant across our entire study (401 samples),

assuming a mean sequencing error rate of 0.2% (Schirmer et al. 2016).

To estimate π, numbers of nonsynonymous and synonymous differences and sites were first

calculated individually for all genes in each of the 401 samples using SNPGenie (Nelson et al. 2015;

snpgenie.pl, https://github.com/chasewnelson/SNPGenie). OLG regions were then analyzed

separately using OLGenie (Nelson et al. 2020; OLGenie.pl,

https://github.com/chasewnelson/OLGenie). Because OLGenie requires a multiple sequence

alignment as input, a pseudo-alignment of 1,000 sequences was constructed for each OLG region in

each sample by randomly substituting single nucleotide variants according to their within-host

frequencies into the Wuhan-Hu-1 reference genome. For non-OLG and OLG regions alike, average

within-host numbers of differences and sites were calculated for each codon by taking the mean

across all samples. For example, if a particular codon contained nonsynonymous differences in two of

401 samples, with the two samples exhibiting mean numbers of 0.01 and 0.002 pairwise differences

per site, this codon was considered to exhibit a mean of (0.01+0.002)/401=0.0000299 pairwise

differences per site across all samples. These codon means were then treated as independent units

of observation during bootstrapping.

Pangolin samples refer to Sequence Read Archive records SRR11093266, SRR11093267,

SRR11093268, SRR11093269, SRR11093270, and SRR11093271. Only 179 single nucleotide

variants could be called prior to our FDR filtering, and samples SRR11093271 and SRR11093270

were discarded entirely due to low mapping quality. We also note that after our quality filtering, four

samples contain consensus alleles that do not match their reference sequence (available from

GISAID): P1E, P4L, P5E, and P5L (Supplementary Table 18).

Within-host recurrent mutations analyses

We assume that each host was infected by a single genotype. Under this assumption, the minor allele

of each segregating site within-host is either due to genotyping or sequencing artifacts, or new

mutations. Because there are very few loci with high-frequency derived alleles between hosts, and

because the Wuhan-Hu-1 genome is used as the reference in read mapping, we here only consider

within-host mutations that differ from this reference background. There are four possible bases at

each locus, A, C, G, and U, and three possible mutational changes away from the Wuhan-Hu-1

reference genome. For each locus, we calculate the number of samples matching the reference allele

as $N=N_1+N_2$, where N_1 is the number of samples in which the Wuhan reference allele is the only

observed allele, and N_2 is the number of samples in which the Wuhan reference allele is major allele.

Given N_2, we further determined the number of samples carrying each of the three possible non-

reference alleles as N_A, N_C, N_G, and N_U. For example, if the reference allele was U, we calculated p_A,

p_C, and p_G, where $p_{All}=p_A+p_C+p_G$. If A was an observed non-reference allele, we calculated the

frequency of A as $p_A=N_A/N$. Thus, a larger frequency indicates the derived allele is observed in a high

proportion of samples (Figure 8). The within-host derived allele frequency (DAF) was estimated as the

total number of reads matching the observed minor allele divided by the total number of reads

mapped to the locus. If all reads matched the Wuhan-Hu-1 reference allele, then DAF=0. Five

mutations occur in more than 10% of samples, four of which are nonsynonymous, with their DAF

plotted in Figure 8—figure supplement 1. For this analysis, we did not apply the per-site FDR cutoff,

thus a DAF=0 is equivalent to the absence of reads mapped to the mutation, after reads are filtered

by sequence quality, mapping quality, and LoFreq's default significance threshold (p=0.01).

Author Contributions

X.W. conceived the study. C.W.N., Z.A., and X.W. designed the study. T.G., S.-O.K., and X.W., advised on the study. C.W.N. and Z.A. obtained and processed data. C.W.N., Z.A., C-H.K., M.C., C.L., S.-O.K., and X.W., analyzed data. C.W.N., Z.A., S.-O.K., and X.W. conceived, discussed, and illustrated figures. All authors discussed and interpreted results. C.W.N., Z.A., T.G., S.-O.K., and X.W. wrote the first draft. All authors read, discussed, and revised the manuscript.

Acknowledgements

This work was supported by a Postdoctoral Research Fellowship from Academia Sinica (to C.W.N. under P.I. Wen-Hsiung Li); funding from the Bavarian State Government and National Philanthropic Trust (to Z.A. under P.I. Siegfried Scherer); NSF IOS grants #1755370 and #1758800 (to S.-O.K.); and the University of Wisconsin-Madison John D. MacArthur Professorship Chair (to T.L.G). Copyright-free images were obtained from Pixabay. The authors thank the GISAID platform and the originating and submitting laboratories who kindly uploaded SARS-CoV-2 sequences to the GISAID EpiCov™ Database for public access (see GISAID Acknowledgments Table in supplement). The authors thank Maciej F. Boni, Reed A. Cartwright, John Flynn, Kyle Friend, Dan Graur, Robert S. Harbert, Cheryl Hayashi, David G. Karlin, Niloufar Kavian, Kin-Hang (Raven) Kok, Wen-Hsiung Li, Meiyeh Lu, David A. Matthews, Lisa Mirabello, Apurva Narechania, Felix Li Jin, and attendees of the UC Berkeley popgen journal club for useful information and discussion; Andrew E. Firth, Alexander Gorbalenya, Irwin Jungreis, Manolis Kellis, Raven Kok, Angelo Pavesi, Kei Sato, Manuela Sironi, and Noam Stern-Ginossar for an invaluable discussion regarding standardizing nomenclature; Helen Piontkivska, Patricia Wittkopp, Antonis Rokas, and one anonymous reviewer for critical suggestions; Ming-Hsueh Lin for immense feedback on figures; and special thanks to Priya Moorjani, Jacob Tennessen, Montgomery Slatkin, Yun S. Song, Jianzhi George Zhang, Xueying Li, Hongxiang Zheng,

Qinqin Yu, Meredith Yeager, and Michael Dean for commenting on earlier drafts of this

manuscript.

References

Affram Y, Zapata JC, Gholizadeh Z, Tolbert WD, Zhou W, Iglesias-Ussel MD, Pazgier M, Ray K, Latinovic OS, Romerio F. 2019. The HIV-1 Antisense Protein ASP Is a Transmembrane Protein of the Cell Surface and an Integral Protein of the Viral Envelope. *J Virol* 93:e00574-19.

Atchley WR, Zhao J, Fernandes AD, Druke T. 2005. Solving the protein sequence metric problem. *Proceedings of the National Academy of Sciences* 102:6395–6400.

Bartonek L, Braun D, Zagrovic B. 2020. Frameshifting preserves key physicochemical properties of proteins. *Proc Natl Acad Sci USA* 117:5907–5912.

Bartonek L, Zagrovic B. 2019. VOLPES: an interactive web-based tool for visualizing and comparing physicochemical properties of biological sequences. *Nucleic Acids Research* 47:W632–W635.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society B* 57:289–300.

Benjamini Y, Yekutieli D. 2001. The control of the false discovery rate in multiple testing under dependency. *The Annals of Statistics* 29:1165–1188.

Bezstarosti K, Lamers MM, Haagmans BL, Demmers JAA. 2020. Targeted Proteomics for the Detection of SARS-CoV-2 Proteins. Biochemistry Available from: http://biorxiv.org/lookup/doi/10.1101/2020.04.23.057810

Bojkova D, Klann K, Koch B, Widera M, Krause D, Ciesek S, Cinatl J, Münch C. 2020. SARS-CoV-2 infected host cell proteomics reveal potential therapy targets. In Review Available from: https://www.researchsquare.com/article/rs-17218/v1

Boni MF, Lemey P, Jiang X, Lam TT-Y, Perry B, Castoe T, Rambaut A, Robertson DL. 2020. Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage responsible for the COVID-19 pandemic. Nature Microbiology, doi: 10.1038/s41564-020-0771-4

Boussau B, Gouy M. 2006. Efficient likelihood computations with nonreversible models of evolution. *Syst Biol* 55:756–768.

Cagliani R, Forni D, Clerici M, Sironi M. 2020. Coding potential and sequence conservation of SARS-CoV-2 and related animal viruses. *Infection, Genetics and Evolution* 83:104353.

Calviello L, Ohler U. 2017. Beyond Read-Counts: Ribo-seq Data Analysis to Understand the Functions of the Transcriptome. *Trends in Genetics* 33:728–744.

Cassan E, Arigon-Chifolleau A-M, Mesnard J-M, Gross A, Gascuel O. 2016. Concomitant emergence of the antisense protein gene of HIV-1 and of the pandemic. *Proc Natl Acad Sci USA* 113:11537–11542.

Chan JF-W, Kok K-H, Zhu Z, Chu H, To KK-W, Yuan S, Yuen K-Y. 2020. Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerging Microbes & Infections* 9:221–236.

Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, Friedberg I, Hamelryck T, Kauff F, Wilczynski B, et al. 2009. Biopython: freely available Python tools for computational molecular biology and bioinformatics. *Bioinformatics* 25:1422–1423.

Cui J, Li F, Shi Z-L. 2019. Origin and evolution of pathogenic coronaviruses. *Nature Reviews Microbiology* 17:181–192.

Daugherty MD, Malik HS. 2012. Rules of Engagement: Molecular Insights from Host-Virus Arms Races. *Annual Review of Genetics* 46:677–700.

Davidson AD, Williamson MK, Lewis S, Shoemark D, Carroll MW, Heesom KJ, Zambon M,

Ellis J, Lewis PA, Hiscox JA, et al. 2020. Characterisation of the transcriptome and proteome of SARS-CoV-2 reveals a cell passage induced in-frame deletion of the furin-like cleavage site from the spike glycoprotein. *Genome Med* 12:68.

Degner JF, Marioni JC, Pai AA, Pickrell JK, Nkadori E, Gilad Y, Pritchard JK. 2009. Effect of read-mapping biases on detecting allele-specific expression from RNA-sequencing data. *Bioinformatics* 25:3207–3212.

Do CB, Mahabhashyam MSP, Brudno M, Batzoglou S. 2005. ProbCons: Probabilistic consistency-based multiple sequence alignment. *Genome Research* 15:330–340.

Ewens WJ, Grant GR. 2001. Statistical Methods in Bioinformatics. New York: Springer-Verlag

Fernandes JD, Faust TB, Strauli NB, Smith C, Crosby DC, Nakamura RL, Hernandez RD, Frankel AD. 2016. Functional Segregation of Overlapping Genes in HIV. *Cell* 167:1762–1773.

Finkel Y, Mizrahi O, Nachshon A, Weingarten-Gabbay S, Morgenstern D, Yahalom-Ronen Y, Tamir H, Achdout H, Stein D, Israeli O, et al. 2020. The coding capacity of SARS-CoV-2. *Nature*, in press, https://doi.org/10.1038/s41586-020-2739-1

Firth AE. 2020. A putative new SARS-CoV protein, 3c, encoded in an ORF overlapping ORF3a. *Journal of General Virology*, https://doi.org/10.1099/jgv.0.001469

Flynn JA, Purushotham D, Choudhary MNK, Zhou X, Fan C, Matt G, Li D, Wang T. 2020. Exploring the coronavirus pandemic with the WashU Virus Genome Browser. *Nat Genet* [Internet]. Available from: http://www.nature.com/articles/s41588-020-0697-z

Forni D, Cagliani R, Clerici M, Sironi M. 2017. Molecular Evolution of Human Coronavirus Genomes. *Trends in Microbiology* 25:35–48.

Forster P, Forster L, Renfrew C, Forster M. 2020. Phylogenetic network analysis of SARS-CoV-2 genomes. *Proc Natl Acad Sci USA*:202004999.

Fung S-Y, Yuen K-S, Ye Z-W, Chan C-P, Jin D-Y. 2020. A tug-of-war between severe acute respiratory syndrome coronavirus 2 and host antiviral defence: lessons from other pathogenic viruses. *Emerging Microbes & Infections* 9:558–570.

Gazave E, Chang D, Clark AG, Keinan A. 2013. Population Growth Inflates the Per-Individual Number of Deleterious Mutations and Reduces Their Mean Effect. *Genetics* 195:969–978.

Ge H, Wang X, Yuan X, Xiao G, Wang C, Deng T, Yuan Q, Xiao X. 2020. The epidemiology and clinical information about COVID-19. *European Journal of Clinical Microbiology & Infectious Diseases* 39:1011–1019.

Gorbalenya AE, Enjuanes L, Ziebuhr J, Snijder EJ. 2006. Nidovirales: Evolving the largest RNA virus genome. *Virus Research* 117:17–37.

Gorbalenya AE, Baker SC, Baric RS, de Groot RJ, Drosten C, Gulyaeva AA, Haagmans BL, Lauber C, Leontovich AM, Neuman BW, et al. 2020. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nature Microbiology* 5:536–544.

Gordon DE, Jang GM, Bouhaddou M, Xu J, Obernier K, White KM, O'Meara MJ, Rezelj VV, Guo JZ, Swaney DL, et al. 2020. A SARS-CoV-2 protein interaction map reveals targets for drug repurposing. *Nature* [Internet]. Available from: http://www.nature.com/articles/s41586-020-2286-9

Greenbaum J, Sidney J, Chung J, Brander C, Peters B, Sette A. 2011. Functional classification of class II human leukocyte antigen (HLA) molecules reveals seven different supertypes and a surprising degree of repertoire sharing across supertypes. *Immunogenetics* 63:325–335.

Grifoni A, Sidney J, Zhang Y, Scheuermann RH, Peters B, Sette A. 2020. A Sequence Homology and Bioinformatic Approach Can Predict Candidate Targets for Immune Responses to SARS-CoV-2. *Cell Host & Microbe* 27:671-680.e2.

Grubaugh ND, Gangavarapu K, Quick J, Matteson NL, De Jesus JG, Main BJ, Tan AL, Paul LM, Brackney DE, Grewal S, et al. 2019. An amplicon-based sequencing framework for accurately measuring intrahost virus diversity using PrimalSeq and iVar. *Genome Biol* 20:8.

Hachim A, Kavian N, Cohen CA, Chin AWH, Chu DKW, Mok CKP, Tsang OTY, Yeung YC, Perera RAPM, Poon LLM, et al. 2020. ORF8 and ORF3b antibodies are accurate serological markers of early and late SARS-CoV-2 infection. *Nat Immunol* 21:1293–1301.

Helmy YA, Fawzy M, Elaswad A, Sobieh A, Kenney SP, Shehata AA. 2020. The COVID-19 Pandemic: A Comprehensive Review of Taxonomy, Genetics, Epidemiology, Diagnosis, Treatment, and Control. *JCM* 9:1225.

Holmes EC. 2009. The Evolution and Emergence of RNA Viruses. New York: Oxford University Press.

Holmes EC, Lipman DJ, Zamarin D, Yewdell JW. 2006. Comment on "Large-Scale Sequence Analysis of Avian Influenza Isolates." *Science* 313:1573b–1573b.

Hughes AL, Friedman R, Glenn NL. 2006. The Future of Data Analysis in Evolutionary Genomics. *Current Genomics* 7:227–234.

Ingolia NT, Ghaemmaghami S, Newman JRS, Weissman JS. 2009. Genome-Wide Analysis in Vivo of Translation with Nucleotide Resolution Using Ribosome Profiling. *Science* 324:218–223.

Jukes TH, Cantor CR. 1969. Evolution of Protein Molecules. In: Munro HN, editor. Mammalian Protein Metabolism. New York: Academic Press. p. 21–132. Available from: https://linkinghub.elsevier.com/retrieve/pii/B9781483232119500097

Jungreis I, Sealfon R, Kellis M. 2020. Sarbecovirus comparative genomics elucidates gene content of SARS-CoV-2 and functional impact of COVID-19 pandemic mutations. Genomics Available from: http://biorxiv.org/lookup/doi/10.1101/2020.06.02.130955

Jurtz V, Paul S, Andreatta M, Marcatili P, Peters B, Nielsen M. 2017. NetMHCpan-4.0: Improved Peptide–MHC Class I Interaction Predictions Integrating Eluted Ligand and Peptide Binding Affinity Data. *J.I.* 199:3360–3368.

Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* 30:772–780.

Keese PK, Gibbs A. 1992. Origins of genes: "big bang" or continuous creation? *Proceedings of the National Academy of Sciences* 89:9489–9493.

Konno Y, Kimura I, Uriu K, Fukushi M, Irie T, Koyanagi Y, Sauter D, Gifford RJ, Nakagawa S, Sato K. 2020. SARS-CoV-2 ORF3b Is a Potent Interferon Antagonist Whose Activity Is Increased by a Naturally Occurring Elongation Variant. *Cell Reports* 32:108185.

Kopecky-Bromberg SA, Martínez-Sobrido L, Frieman M, Baric RA, Palese P. 2007. Severe Acute Respiratory Syndrome Coronavirus Open Reading Frame (ORF) 3b, ORF 6, and Nucleocapsid Proteins Function as Interferon Antagonists. *JVI* 81:548–557.

Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, Hengartner N, Giorgi EE, Bhattacharya T, Foley B, et al. 2020. Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. *Cell* 182:812-827.e19.

Kosakovsky-Pond SL. 2020. Natural selection analysis of SARS-CoV-2/COVID-19. *usegalaxy* [Internet]. Available from: https://covid19.galaxyproject.org/evolution/

Lam TT-Y, Shum MH-H, Zhu H-C, Tong Y-G, Ni X-B, Liao Y-S, Wei W, Cheung WY-M, Li W-J, Li L-F, et al. 2020. Identifying SARS-CoV-2 related coronaviruses in Malayan pangolins. *Nature* [Internet] in press. Available from: http://www.nature.com/articles/s41586-020-2169-0

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359.

Langmead B, Wilks C, Antonescu V, Charles R. 2019. Scaling read aligners to hundreds of threads on general-purpose processors.Hancock J, editor. *Bioinformatics* 35:421–432.

Larsson A. 2014. AliView: a fast and lightweight alignment viewer and editor for large datasets. *Bioinformatics* 30:3276–3278.

Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, Morgan MT, Carey VJ. 2013. Software for Computing and Annotating Genomic Ranges.Prlic A, editor. *PLoS Comput Biol* 9:e1003118.

Lokugamage KG, Hage A, Schindewolf C, Rajsbaum R, Menachery VD. 2020. SARS-CoV-2 is sensitive to type I interferon pretreatment. Microbiology Available from: http://biorxiv.org/lookup/doi/10.1101/2020.03.07.982264

McBride R, Fielding B. 2012. The Role of Severe Acute Respiratory Syndrome (SARS)-Coronavirus Accessory Proteins in Virus Pathogenesis. *Viruses* 4:2902–2923.

McKinney W. 2010. Data structures for statistical computing in python. In: Proceedings of the 9th Python in Science Conference. Vol. 445. p. 51–56.

Michel CJ, Mayer C, Poch O, Thompson JD. 2020. Characterization of accessory genes in coronavirus genomes. *Virol J* 17:131.

Nei M, Gojobori T. 1986. Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Molecular Biology and Evolution* 3:418–426.

Nei M, Kumar S. 2000. Molecular Evolution and Phylogenetics. New York, NY: Oxford University Press

Nelson CW, Ardern Z, Wei X. 2020. OLGenie: Estimating Natural Selection to Predict Functional Overlapping Genes. *Molecular Biology and Evolution* in press:msaa087.

Nelson CW, Hughes AL. 2015. Within-host nucleotide diversity of virus populations: Insights from next-generation sequencing. *Infection, Genetics and Evolution* 30:1–7.

Nelson CW, Moncla LH, Hughes AL. 2015. SNPGenie: estimating evolutionary parameters to detect natural selection using pooled next-generation sequencing data. *Bioinformatics* 31:3709–3711.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and Evolution* 32:268–274.

Paul S, Sidney J, Sette A, Peters B. 2016. TepiTool: A Pipeline for Computational Prediction of T Cell Epitope Candidates. *Current Protocols in Immunology* 114:18.19.1-18.19.24.

R Core Team. 2018. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing Available from: https://www.R-project.org/

Rehman S ur, Shafique L, Ihsan A, Liu Q. 2020. Evolutionary Trajectory for the Emergence of Novel Coronavirus SARS-CoV-2. *Pathogens* 9:240.

Reynisson B, Barra C, Kaabinejadian S, Hildebrand WH, Peters B, Nielsen M. 2020. Improved Prediction of MHC II Antigen Presentation through Integration and Motif Deconvolution of Mass Spectrometry MHC Eluted Ligand Data. *Journal of Proteome Research* 19:2304–2315.

Rothe C, Schunk M, Sothmann P, Bretzel G, Froeschl G, Wallrauch C, Zimmer T, Thiel V, Janke C, Guggemos W, et al. 2020. Transmission of 2019-nCoV Infection from an Asymptomatic Contact in Germany. *N Engl J Med* 382:970–971.

Sabath N, Landan G, Graur D. 2008. A Method for the Simultaneous Estimation of Selection Intensities in Overlapping Genes. *PLoS ONE* 3:e3996.

Schirmer M, D'Amore R, Ijaz UZ, Hall N, Quince C. 2016. Illumina error profiles: resolving fine-scale variation in metagenomic sequencing data. *BMC Bioinformatics* 17:125.

Schlub TE, Buchmann JP, Holmes EC. 2018. A simple method to detect candidate overlapping genes in viruses using single genome sequences. *Molecular Biology and Evolution* 35:2572–2581.

Schwanhäusser B, Busse D, Li N, Dittmar G, Schuchhardt J, Wolf J, Chen W, Selbach M. 2011. Global quantification of mammalian gene expression control. *Nature* 473:337–342.

Sidney J, Peters B, Frahm N, Brander C, Sette A. 2008. HLA class I supertypes: a revised and updated classification. *BMC Immunology* 9:1.

Soubrier J, Steel M, Lee MSY, Der Sarkissian C, Guindon S, Ho SYW, Cooper A. 2012. The

Influence of Rate Heterogeneity among Sites on the Time Dependence of Molecular Rates. *Molecular Biology and Evolution* 29:3345–3358.

Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein sequence alignments into the corresponding codon alignments. *Nucleic Acids Research* 34:W609–W612.

Tavaré S. 1986. Some probabilistic and statistical problems in the analysis of DNA sequences. *Lectures on Mathematics in the Life Sciences* 17:57–86.

Tyanova S, Temu T, Cox J. 2016. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nat Protoc* 11:2301–2319.

Warren AS, Archuleta J, Feng W, Setubal JC. 2010. Missing genes in the annotation of prokaryotic genomes. *BMC Bioinformatics* 11:131.

Wei X, Zhang J. 2015. A Simple Method for Estimating the Strength of Natural Selection on Overlapping Genes. *Genome Biology and Evolution* 7:381–390.

Wells HL, Letko M, Lasso G, Ssebide B, Nziza J, Byarugaba DK, Navarrete-Macias I, Liang E, Cranfield M, Han BA, et al. 2020. The evolutionary history of ACE2 usage within the coronavirus subgenus *Sarbecovirus*. Evolutionary Biology Available from: http://biorxiv.org/lookup/doi/10.1101/2020.07.07.190546

Wilm A, Aw PPK, Bertrand D, Yeo GHT, Ong SH, Wong CH, Khor CC, Petric R, Hibberd ML, Nagarajan N. 2012. LoFreq: a sequence-quality aware, ultra-sensitive variant caller for uncovering cell-population heterogeneity from high-throughput sequencing datasets. *Nucleic Acids Research* 40:11189–11201.

Worobey M, Pekar J, Larsen BB, Nelson MI, Hill V, Joy JB, Rambaut A, Suchard MA, Wertheim JO, Lemey P. 2020. The emergence of SARS-CoV-2 in Europe and North America. *Science*:eabc8169.

Wu A, Peng Y, Huang B, Ding X, Wang X, Niu P, Meng J, Zhu Z, Zhang Z, Wang J, et al. 2020. Genome Composition and Divergence of the Novel Coronavirus (2019-nCoV) Originating in China. *Cell Host & Microbe* 27:325–328.

Wu F, Zhao S, Yu B, Chen Y-M, Wang W, Song Z-G, Hu Y, Tao Z-W, Tian J-H, Pei Y-Y, et al. 2020. A new coronavirus associated with human respiratory disease in China. *Nature* 579:265–269.

Yang Z. 1995. A Space-Time Process Model for the Evolution of DNA Sequences. *Genetics* 139:993–1005.

Yi Y, Lagniton PNP, Ye S, Li E, Xu R-H. 2020. COVID-19: what has been learned and to be learned about the novel coronavirus disease. *Int. J. Biol. Sci.* 16:1753–1766.

Yuen K-S, Ye Z-W, Fung S-Y, Chan C-P, Jin D-Y. 2020. SARS-CoV-2 and COVID-19: The most important research questions. *Cell Biosci* 10:40.

Yurkovetskiy L, Wang X, Pascal KE, Tomkins-Tinch C, Nyalile T, Wang Y, Baum A, Diehl WE, Dauphin A, Carbon\e C, et al. 2020. Structural and Functional Analysis of the D614G SARS-CoV-2 Spike Protein Variant. *Cell*, in press. Available from: https://www.cell.com/cell/fulltext/S0092-8674(20)31229-0

Zecha J, Lee C-Y, Bayer FP, Meng C, Grass V, Zerweck J, Schnatbaum K, Michler T, Pichlmair A, Ludwig C, et al. 2020. Data, Reagents, Assays and Merits of Proteomics for SARS-CoV-2 Research and Testing. *Mol Cell Proteomics* 19:1503–1522.

Zhou W, Chen T, Zhao H, Eterovic AK, Meric-Bernstam F, Mills GB, Chen K. 2014. Bias from removing read duplication in ultra-deep sequencing experiments. *Bioinformatics* 30:1073–1080.

Zhou P, Li H, Wang H, Wang L-F, Shi Z. 2012. Bat severe acute respiratory syndrome-like coronavirus ORF3b homologues display different interferon antagonist activities. *Journal of General Virology* 93:275–281.

1   Tables

1   **Table 1.** Nomenclature and reading frames for overlapping gene candidates in SARS-CoV-2 *ORF3a*.

Gene[a]	Reading frame[b]	Genome positions, Wuhan-Hu-1 (CDS positions, (*ORF3a*)[c]	Length	Description	References
ORF3a	ss11 (reference)	25393-26220 (1-828)	276 codons (828 nt)	Ion channel formation and virus release in SARS-CoV infection; host cell apoptosis; triggers inflammation; antagonizes interferon	Lu et al. (2010); Cui et al. (2019)
ORF3c	ss13	25457-25582 (65-190)	42 codons (126 nt)	Features suggestive of a viroporin (Cagliani et al. 2020); lowest π_N/π_S ratio estimated for any gene in our between-host selection analysis (Figure 5); overlaps codons 22-64 of *ORF3a*	First discovered by Cagliani et al. (2020) as *ORF3h*; *ORF3c* in Firth (2020); *ORF3c* in Jungreis et al. (2020); *ORF3a.iORF1* in Finkel et al. (2020); *ORF3b* in Pavesi (2020)
ORF3d	ss12	25524-25697 (132-305)	58 codons (174 nt)	Aligned to and named *ORF3b* by Chan et al. (2020) but is not homologous to *ORF3b*; interferon antagonism has not been demonstrated; binds STOML2 mitochondrial protein (Gordon et al. 2020); contains a predicted signal peptide in the region encoding *ORF3d-2*; contains an *X* motif in pangolin-CoV but not SARS-CoV-2 (Michel et al. 2020); may contribute to differences between SARS-CoV and SARS-CoV-2 in immune response as a unique antigenic target (Hachim et al. 2020; Niloufar Kavian, pers. comm.); overlaps codons 44-102 of *ORF3a*	Present study; first discovered by Chan et al. (2020) but misclassified as *ORF3b*; *ORF3b* in Gordon et al. (2020), Hachim et al. (2020), and citing studies; 'hypothetical protein' in Pavesi (2020); 'a completely different ORF' in Michel et al. (2020)
ORF3d-2	ss12	25596-25697 (204-305)	34 codons (102 nt)	A shorter version of *ORF3d* that excludes its first 24 codons, where the majority of premature STOP codons in SARS-CoV-2 are located; contains a predicted signal peptide (Finkel et al. 2020); likely expressed at higher levels than full-length *ORF3d* (Figure 2A) overlaps codons 68-102 of *ORF3a*	Discovered by Finkel et al. (2020) as *ORF3a-iORF2*
ORF3a-2	ss11 (reference)	25765-26220 (373-828)	152 codons (456 nt)	A shorter version of *ORF3a* that excludes its first 124 codons; evidence of expression separate from that of *ORF3a* (Davidson et al. 2020); has also been conflated with *ORF3b*; equivalent to codons 124-276 of *ORF3a*	Discovered by Davidson et al. (2020; pers. comm.) but referred to as *ORF3b*
ORF3b region[d]	ss13	25814-26281; four ORFs at 25814-82, 25910-84, 26072-170, and 26183-281 (422-889; ORFs at 422-	156 codons (468 nt); the four ORFs are 23, 25, 33, and 33 codons (69, 75, 99, and 99 nt)	Full-length in some related viruses, but truncated by multiple in-frame STOP codons in SARS-CoV-2; longer forms function as interferon antagonist in SARS-related viruses; may contribute to differences between SARS-CoV and SARS-CoV-2 in immune response; although aligned to	Konno et al. (2020) claim functionality for the first (23-codon) ORF in SARS-CoV-2; the first ORF is also mentioned by Wu et al. (2020)

90, 518-92, 680-778, and 791-889)	*ORF3d* by Chan et al., the two are not homologous; overlaps codons 141-276 of *ORF3a* (first ORF overlaps codons 141-164)

[a]Genes are listed by position of start site in the genome from 5′ (top) to 3′ (bottom).

[a]Nomenclature as described in Wei and Zhang (2015) and Nelson et al. (2020): ss=sense-sense (same strand);

ss12=codon position 1 of the reference frame overlaps codon position 2 of the overlapping frame on the same

strand; ss13=codon position 1 of the reference frame overlaps codon position 3 of the overlapping frame on the

same strand. Frame is indicated from the perspective of *ORF3a* as the reference gene, i.e., *ORF3d* starts at

codon position 3 of *ORF3a*, while *ORF3c* and *ORF3b* start at codon position 2 of *ORF3a*.

[c]Positions and counts include STOP codons. Positions or sequences were indicated by the original publications

and verified by personal communication if ambiguous.

[d]The SARS-CoV-2 region homologous to *ORF3b* of SARS-CoV contains 4 premature STOP codons and four

distinct ORFs (AUG-to-STOP); see Figure 4, Figure 6, and Supplementary Table 4.

Table 2. The mutational path to European pandemic founder haplotypes[a]

Variant	EP–3	EP–2	EP[b]	EP+1[b]	EP+1+LOF
C241U (5′-UTR)	0	0	1	1	1
C3037U (nsp3-F106F)	0	0	1	1	1
C14408U (RdRp-P323L)	0	0	0	1	1
A23403G (Spike-D614G)[c]	0	1	1	1	1
G25563U (ORF3a-Q57H/ORF3d-E14*)	0	0	0	0	1
Earliest collection[d]	24-Dec	7-Feb	28-Jan	20-Feb	21-Feb
Earliest location[d]	Wuhan	Wuhan	Munich (Shanghai)[b]	Lombardy	Hauts de France
Occurrence in China	233	1	1 (2)[b]	0	0
Occurrence in Europe	458	0	21	1153	310
Occurrence in Italy	1	0	0	27	0
Occurrence in Germany	15	0	1	11	21
Occurrence in Belgium	27	1	20	187	27
Occurrence in UK	210	0	0	338	38
Occurrence in Iceland	56	0	0	212	54
Occurrence in France	14		0	72	102
Occurrence in US	467	0	0	88[e]	326[e]
Occurrence in GISAID[f]	1610	2	22	1455	752

[a]Haplotypes are here defined by five high-frequency variants (rows 1-5), and other variants with lower frequency on these backgrounds are ignored. EP-1 is not observed in the dataset.

[b]This EP haplotype is first detected in German patient #4 and is a documented founder for coronavirus spread in Germany (Rothe et al. 2020). Neither the EP nor EP+1 haplotypes were detectable between January 28 and February 20, although they immediately became a major haplotype once EP+1 was detectable. Failure to detect these two haplotypes during these three weeks could potentially be explained by ascertainment bias, e.g., lack of testing for travel-independent cases.

[c]This Shanghai sample (GISAID: EPI_ISL_416327) comprises 1.32% poly-Ns and failed our quality control criteria, but is added here since it is potentially relevant to the origin of EP haplotype. Including this sample, the EP haplotype is observed in Shanghai twice.

[d]The earliest collection location and time are highly subject to collection and submission bias and do not necessarily reflect where the mutation/haplotype first occurred.

[e]There is likely a testing bias in the United States, as EP+1+LOF haplotype was often detected in Washington, and EP+1 haplotype was not.

[f]These numbers are based on 3853 samples from December 24 to April 1 at the time of GISAID accession that passed both our quality control criteria for alignment and for this particular analysis (i.e., no ambiguous genotype calls among the five SNPs in this table), unless otherwise stated.

1 Figure 1

A SARS-CoV-2 (Wuhan-Hu-1)

B *Severe acute respiratory syndrome-related coronavirus* species members

2
Figure 1. Gene repertoire and evolutionary relationships of *Severe acute respiratory syndrome-related*
coronavirus species members. Only genes downstream of *ORF1ab* are shown, beginning with *S* (Spike-
encoding). **(A)** Four types of genes and their relative positions in the SARS-CoV-2 Wuhan-Hu-1 genome
(NCBI: NC_045512.2). Genes are colored by type: novel overlapping genes (OLGs) (gold; *ORF3d* only);
conserved OLGs (burgundy); accessory (green); and structural (blue). Note that *ORF3b* has been
truncated relative to SARS-CoV genomes, whereas *ORF8* remains intact (i.e., has not been split into
ORF8a and *ORF8b*). **(B)** Genes with intact ORFs in each of the the 21 *Severe acute respiratory*
syndrome-related coronavirus genomes. Gene positions are shown relative to each genome, i.e.,
homologous genes are not precisely aligned. Only the full-length versions of *ORF3a* and *ORF3d* are
shown (for shorter isoforms, see Table 1). Note that the first 20 codons of *ORF3d* overlap the last
codons of *ORF3c* (supplementary file SARS-CoV-2.gtf), such that the beginning of *ORF3d* involves a
triple overlap (*ORF3a*/*ORF3c*/*ORF3d*). *ORF3b* is full-length in only 3 sequences (SARS-CoV TW11,
SARS-CoV Tor2, and bat-CoV Rs7327), while the remaining sequences have either an early or a late
premature STOP codon (Supplementary Table 9). *ORF8* is not novel in SARS-CoV-2 (contra Chan et

al. 2020), but is intact in all but 5 sequences (split into *ORF8a* and *ORF8b* in SARS-CoVs TW11 and Tor2; deleted in bat-CoVs BtKY72, BM48-31, and JTMC15). *ORF9b* and *ORF9c* are found throughout this virus species, yet rarely annotated in genomes at NCBI.

Figure 1—source data 1. SARS-related-CoV_ALN.fasta. Whole-genome multiple sequence alignment of 21 genomes of the species *Severe acute respiratory syndrome-related coronavirus*. Note that the pangolin-CoV GD/1 sequence has been masked as N (license constraint), but can be acquired with GISAID registration.

Figure 1—source data 2. SARS-related-CoV_ALN.gtf.txt. Gene Transfer Format (GTF) file giving gene positions within SARS-related-CoV_ALN.fasta. Genes were extracted and gaps removed from each sequence before displaying.

Figure 1—figure supplement 1. Codon permutation analysis to identify candidate overlapping genes
in all three forward-sense reading frames of the SARS-CoV-2 genome. Genome positions of known
and hypothesized genes are indicated with horizontal red lines. Reading frames 1, 2, and 3 begin at
position 1, 2, or 3 of the Wuhan-Hu-1 reference genome, respectively, with genome coordinates shown
at the bottom (Supplementary Tables 2 and 3). Yellow indicates low p-values (natural logarithm scale),
while gray indicates absence of a stop-to-stop ORF longer than 30 codons or presence outside of a
known canonical gene (not tested). Note that low p-values are expected for the non-overlapping genes.
Only the first of four SARS-CoV-2 ORFs homologous to *ORF3b* is shown (Figure 4; Supplementary
Table 4).
12
13

Figure 2. SARS-CoV-2 gene expression in ribosome profiling and mass spectrometry datasets.
Ribosome profiling (Ribo-seq) data were obtained from Finkel et al. (2020); mass spectrometry data
were obtained from Davidson et al. (2020) and Bezstarosti et al. (2020). Reads were trimmed to their
first (5′) nucleotide to minimize statistical dependence while preserving reading frame. Results are
shown after pooling samples treated with harringtonine (SRR11713360, SRR11713361) or
lactimidomycin (SRR11713358, SRR11713359) at 5 hours post-infection. (**A**) Ribosome profiling
coverage (read depth) near translation initiation sites, measured as mean reads per million mapped
reads. Only reads of length 29-31 nucleotides were used, chosen for their enrichment at the start sites
of highly expressed annotated genes (Figure 2—figure supplement 4). Light blue backgrounds denote

fully (internal) overlapping genes. Annotated genes show an accumulation of 5′ ends of protected
reads upstream of the gene's start site (grey dotted vertical line), peaking near the ribosome P-site
offset of -12 nt (red=maximum depth). Distributions are largely consistent across individual samples
(Figure 2—figure supplement 1). The ranges of the y axes vary according to expression level, with the
most highly expressed gene (*N*; Figure 2—figure supplement 2) having the largest range. (**B**)
Correlation between protein expression as estimated by mass spectrometry and ribosomal profiling.
'iBAQ prop' refers to the relative (proportion of maximum) protein intensity-based absolute
quantification (iBAQ) value (Schwanhäusser et al. 2011). Genes are denoted by shape and ordinally
colored by iBAQ prop from high (red) to low (blue). 'Upstream peak' refers to the maximum read depth
observed at the approximate P-site offset (red bars in Figure 2A), while mean read depths were
measured across each gene using 30 nt reads separately for in-frame (codon position 1) and out-of-
frame (codon positions 2 and 3) sites (non-overlapping gene regions only, except for *ORF9b*). (**C**)
Reading frame of ribosome profiling reads in the *ORF3c*/*ORF3d* region of *ORF3a*. Solid lines show
the fraction of reads in each frame, summed across samples in sliding windows of 30 nt (step size=1
nt; read length=30 nt). Color denotes frame: green=reference frame (*ORF3a*); burgundy=ss13, the
sense frame encoding *ORF3c*, whose codon position 3 overlaps codon position 1 of *ORF3a*; and
gold=ss12, the sense frame encoding *ORF3d*, whose codon position 2 overlaps codon position 1 of
ORF3a. Values are shown for the central nucleotide of each window, with shaded regions
corresponding to 95% binomial confidence intervals. Alternative frame translation is suggested where
a given frame (solid line) exceeds its average across the remainder of the gene (horizontal dashed
line; non-OLG regions of *ORF3a*). Vertical gray dotted lines indicate gene start and end sites. Gray
bars show read depth for each window, with a maximum of 2889 reads at genome position 25442.
23
24 **Figure 2—source data 1.** riboseq_upstream_peaks.txt. Source data for Figure 2A.
25
26 **Figure 2—source data 2.** expression_data_by_gene_frame.txt. Source data for Figure 2B.
27
28 **Figure 2—source data 3.** riboseq_ORF3d_sliding_window.txt. Source data for Figure 2C.

Figure 2—figure supplement 1

3 **Figure 2—figure supplement 1.** Ribosome profiling coverage near translation initiation sites for
4 individual samples. Details as in Figure 2A, except samples were not pooled and genes are arranged
5 vertically by start site in the genome from 5′ to 3′ (top to bottom). Treatments are indicated as
6 Harr=harringtonine and LTM=lactimidomycin at 5 hours post-infection (hpi).
7

Figure 2—figure supplement 2

Figure 2—figure supplement 2. Correlation between protein expression as estimated by mass
spectrometry and ribosomal profiling for individual treatments and codon positions. Details as in
Figure 2B, except estimates are shown for all sites in a gene (All frames) and out-of-frame reads are
shown separately for codon positions (Codon pos) 2 and 3. *ORF9b* is a fully (internal) overlapping
gene occurring within *N*, with codon position 3 of *ORF9b* corresponding to codon position 1 of *N*. As a
result, codon position 3 of *ORF9b* is predominantly occupied by ribosomes translating *N*, the most
highly expressed gene, leading to overestimation of *ORF9b* expression (Codon pos 3; right).
Ribosome profiling read depth at codon position 1 exhibits a marginal Spearman's rank correlation
with mass spectrometry for cycloheximide (CHX), harringtonine (Harr), and lactimidomycin (LTM)
treatments; however, as expected, all correlations for RNA-seq (mRNA), as well as reads mapping to
other codon positions, were not significant.
14
15

2
3
Figure 2—figure supplement 3. Reading frames occupied by the 5′ ends of ribosome profiling reads
as a function of read length. Processed, aligned reads from Finkel et al. (2020) were trimmed to their
5′ ends (first nt). Fractions of reads mapping to each codon position (x axis) are shown as a function
of read length (y axis; 26-32 nt) after pooling samples according to treatment (CHX=cycloheximide,
Harr=harringtonine, LTM=lactimidomycin, mRNA=RNA-seq) and time (hours post-infection; hpi).
Codon position is defined relative to gene start sites of annotated (canonical) non-overlapping genes,
excluding any overlapping gene regions. Gray bars show relative total counts for reads of different
lengths, with different scales for 5 hpi (lower coverage; max=284,608 reads at CHX/30nt) and 24 hpi
(higher coverage; max=3,149,251 reads at Harr/29nt). At 5 hpi, reads of length 29 and 30 nt tend to
have 5′ ends mapping to the correct protein-coding reading frame (codon position 1) of the gene in
which they occur, and also tend to be most abundant, as expected given the size of ribosome-
protected fragments (~30 nt). Samples from 24 hpi have flatter (left-shifted) read length distributions
and no clear frame signal. The horizontal dashed grey line corresponds to the expectation for RNA
fragments not protected by ribosomes, i.e., 33% of reads mapping to each codon position (random).
18

2
Figure 2—figure supplement 4. Ribosome profiling read accumulation at gene start sites as a
function of read length. Points indicate the percentage of all reads in the region shown whose first
nucleotides (5′ ends) map to each genome position, colored according to read length. In each case,
the longer read lengths have 5′ ends that predominate at approximately -12 nt from the gene start site
(horizontal dashed lines, right), corresponding to the ribosome P-site offset (center of yellow
highlight). A peak in total read depth (grey bars) is also observed at this position, as seen in Figure
2A. Highly expressed canonical (*N*, *M*, *S*, and *ORF7b*) and overlapping (*ORF9b*) genes are shown on
the left, while *ORF3a* and its hypothesized overlapping genes are shown on the right. Note that
ORF9b begins shortly after *N*, such that the upstream peak of *N* occurs at position -22 nt relative to
the start site of *ORF9b*.
13

 Figure 2—figure supplement 5

Figure 2—figure supplement 5. Reading frame of ribosome profiling reads in the *ORF3c*/*ORF3d*
region of *ORF3a* for individual treatments. Details as in Figure 2C, except results are shown
separately after pooling the two biological replicates for each treatment (CHX=cycloheximide,
Harr=harringtonine, LTM=lactimidomycin, mRNA=RNA-seq) (Finkel et al. 2020). Genes and their
reading frames are denoted by color according to Figure 1, with Frames 1, 2, and 3 corresponding to
the frames beginning at codon positions 1, 2, and 3 of *ORF3a*, respectively. Clear evidence for
translation in the appropriate reading frame is observed for *ORF3c* (burgundy) and *ORF3d*/*ORF3d-2*
(gold). The status of the first half of *ORF3d* is obscured by the signal of *ORF3c*, but the increase in
Frame 1 (*ORF3a*) reads and drop in proportion of Frame 2 (*ORF3c*) reads at the start of *ORF3d* is
suggestive of an otherwise-unexpected biological change at this locus.

Figure 2—figure supplement 6. Reading frame of ribosome profiling reads in the *ORF3c*/*ORF3d*
region of *ORF3a* for individual treatments using a smaller sliding window. Details as in Figure 2C and
Figure 2—figure supplement 5, except results are shown for a sliding window size of 19 nt (step
size=1 nt).

2
Figure 2—figure supplement 7. Reading frame of ribosome profiling reads in the *N*/*ORF9b*/*ORF9c*
region. Details as in Figure 2C and Figure 2—figure supplement 5, except Frames 1, 2, and 3
correspond to the frames beginning at positions 1, 2, and 3 of the Wuhan-Hu-1 reference genome,
respectively. Expression of *ORF9b* is evidenced by a peak in the fraction of reads mapping to Frame
3 (burgundy) at the appropriate region, as well as a peak in read depth (gray bars) immediately
upstream (Figure 2A). Translational status of *ORF9c* is ambiguous, with minimal enrichment of reads
in the appropriate frame (burgundy) along the gene.

3 **Figure 2—figure supplement 8.** Reading frame of ribosome profiling reads in the full *ORF3a* gene.
4 Details as in Figure 2C and Figure 2—figure supplement 5. No clear evidence is observed for the
5 expression of any of the *ORF3b* ORFs present in SARS-CoV-2, but the translational status of these
6 ORFs requires further investigation.
7
8
9

2
Figure 2—figure supplement 9. Reading frame of ribosome profiling reads in the *ORF3c*/*ORF3d*
region of *ORF3a* for reads of length 29 nt. Details as in Figure 2C and Figure 2—figure supplement 5,
except results are shown for reads of length 29 nt, which are also indicative of frame but have slightly
lower coverage (Figure 2—figure supplement 3) and exhibit frame-related periodicity. Evidence for
ORF3c (Frame 2; burgundy) is more pronounced using this read length, while the peak for *ORF3d*
(Frame 3; gold) occurs slightly later, overlapping *ORF3d-2*.

9
10
11

1   Figure 3

2
Figure 3. SARS-CoV-2 protein sequence properties. **(A)** Predicted densities of MHC class I-bound
CD8+ T cell 9 aa epitopes (top) and MHC class II-bound CD4+ T cell 15 aa epitopes (bottom). Results
for proteins encoded downstream of *ORF1ab* are shown. Mean numbers of predicted epitopes per
residue (blue bars) are calculated as the number of epitopes overlapping each amino acid position
divided by protein length. Error bars show 95% confidence intervals. Two sets of negative controls were
also tested: (1) *n*=1,000 randomized peptides generated from each protein by randomly sampling its
amino acids with replacement (dark gray bars), representing the result expected given amino acid
content alone; and (2) short unannotated ORFs, the peptides encoded by *n*=103 putatively
nonfunctional ORFs present in the SARS-CoV-2 genome, representing the result expected for real
ORFs that have been evolving without functional constraint. For the short unannotated ORFs, the
horizontal gray dotted line shows the mean number of epitopes per residue, and the gray shaded region
shows the 95% confidence interval (*i.e.*, 2.5% to 97.5% quantiles). ORF3d and N have the lowest MHC
class I epitope density; ORF3d, ORF3b, ORF7b, and ORF10 have the lowest MHC class II epitope
density. **(B)** Hydrophobicity profiles of amino acid sequences in the three forward-strand reading frames
of the *ORF3a*/*ORF3c*/*ORF3d* gene region, as calculated by the VOLPES server, using the unitless
"Factor 1" consensus hydrophobicity scale. Frame is reported using *ORF3a* as the reference, e.g., ss12
refers to the frame encoding *ORF3d*, for which codon position 2 overlaps codon position 1 of *ORF3a*
(Figure 2C).

Figure 3—source data 1. epitope_summary_MHCI_LONG.txt. Source data for MHC class I in Figure 3A.

Figure 3—source data 2. epitope_summary_MHCII_LONG.txt. Source data for MHC class II in Figure 3A.

Figure 3—source data 3. hydrophobicity_profiles_ORF3a_corr.txt. Source data for hydrophobicity profiles in Figure 3B.

Figure 3—figure supplement 1. Structural prediction for the ORF3d protein. Independent computational modeling predictions of *α*-helices at the secondary (left inset, carried out in NetSurf v2) and tertiary structure levels (right inset, carried out in I-TASSER). Folding concordance with the closest protein structure match is shown (rightmost inset, aligned with TM-align). For explanation of shown metrics, see https://zhanglab.ccmb.med.umich.edu/I-TASSER. Chan et al. (2020) also predict a fold with α-helices (Raven Kok, pers. comm.).

3
Figure 3—figure supplement 2. Correlations between hydrophobicity profiles of amino acid sequences encoded by the three forward-strand reading frames of the *ORF3a* region. Vertical dotted lines denote gene start sites. **(A)** Hydrophobicity profiles of the amino acid sequences encoded by all three forward-strand reading frames of *ORF3a*. Details as in Figure 3B, except the entire region encoding *ORF3a* is shown. **(B)** Correlations (Spearman's rank) between hydrophobicity profiles for the three forward-strand reading frames. Color denotes significance.

3
4
5   **Figure 3—figure supplement 3**. Correlation between hydrophobicity profiles in the amino acid
6   sequences encoded by all three forward-strand frames of *ORF3a* by gene subregion. Details as in
7   Figure 3—figure supplement 2. *ORF3a* (left) refers to those codons of *ORF3a* that encode residues not
8   overlapping any hypothesized overlapping genes. Codons counts refer to *ORF3a*, i.e, 21 codons of
9   *ORF3a* (20 codons of *ORF3d*) are involved in a triple overlap (at least one nucleotide) with *ORF3c*.
10
11
12

Figure 4. Amino acid variation in proteins encoded by genes overlapping *ORF3a* in viruses of the species *Severe acute respiratory syndrome-related coronavirus*. **(A)** Amino acid sequences of ORF3c, ORF3d, and ORF3b, as encoded by SARS-CoV-2 (Wuhan-Hu-1; NCBI: NC_045512.2). Note that, while ORF3b encodes a protein of 154 amino acids (aa) in SARS-CoV (NCBI: NC_004718.3), a premature STOP codon in SARS-CoV-2 has resulted in an ORF encoding only 22 aa. **(B)** Amino acid alignments of ORF3c, ORF3d, and ORF3b (3b) show their sequence conservation. Black lines indicate STOP codons in *ORF3d* and *ORF3b*, showing their restricted taxonomic ranges. Intact *ORF3d* is restricted to SARS-CoV-2 and pangolin-CoV GX/P5L; however, note that ORF3d-2 (denoted 3d-2), a shorter isoform of ORF3d, could have a slightly wider taxonomic range if TTG or GTG are permitted as translation initiation codons. Full-length ORF3b (ORF3b region) is found throughout members of this virus species, but truncated early in most genomes outside of SARS-CoV (Supplementary Tables 2, 3, 4, and 8), with the shortest version (denoted 3b) found in SARS-CoV-2 and closely related viruses.

Figure 4—source data 1. aa_alignments_3c_3d_3b.xlsx. Amino acid alignments shown in Figure 4. The full translation of the *ORF3b* region is provided for ORF3b. Note that the pangolin-CoV GD/1 sequence has been masked as X (license constraint), but can be acquired with GISAID registration.

    Figure 5

2
Figure 5. Natural selection analysis of viral nucleotide differences at three hierarchical evolutionary
levels. Nucleotide differences in each virus gene were analyzed at three host levels: between-taxa
divergence (*d*) among *Severe acute respiratory syndrome-related coronavirus* genomes infecting bat,
human, and pangolin species; between-host diversity (π) for SARS-CoV-2 infecting human individuals
(consensus-level); and within-host diversity (π) for SARS-CoV-2 infecting human individuals (deep
sequencing). Each gene/level is shaded according to the ratio of mean nonsynonymous to synonymous
differences per site to indicate purifying selection ($d_N/d_S<1$ or $\pi_N/\pi_S<1$; blue) or positive selection
($d_N/d_S>1$ or $\pi_N/\pi_S>1$; red), except the value for *ORF3c* was artificially lowered to allow the display of
other ratios. A Jukes-Cantor correction was applied to d_N/d_S values. Values range from a minimum of
$\pi_N/\pi_S=0.04$ (*ORF3c*, between-host; p=0.0410, Q=0.308) to a maximum of 21.0 (*ORF9b*, between-host;
p=0.126, Q=0.453), where significance (Q) was evaluated using a Benjamini-Hochberg false-discovery
rate correction after *Z*-tests of the hypothesis that $d_N-d_S=0$ or $\pi_N-\pi_S=0$ (10,000 bootstrap replicates,
codon unit). The mean of all pairwise comparisons is shown for sequenced genomes only, i.e., no
ancestral sequences were reconstructed or inferred. For each gene, sequences were only included in
the between-species analysis if a complete, intact ORF (no STOPs) was present. Genes containing an
overlapping gene (OLG) in a different frame were analyzed separately for non-OLG and OLG regions
using SNPGenie and OLGenie, respectively. For *ORF3b*, only the region corresponding to the first ORF
in SARS-CoV-2 (Table 1) was analyzed. The short overlap between *ORF1a* and *ORF1b* (*nsp11* and
nsp12) was excluded from the analysis. Error bars represent the standard error of mean pairwise
differences. See Methods for further details.

23
24   **Figure 5—source data 1.** selection_three_levels.txt. Source data for Figure 5.

25
26

Figure 5—figure supplement 1. Correlation between natural selection and gene expression. A weak negative Spearman correlation between d_N/d_S and gene expression is observed across all three evolutionary levels (between-taxa, between-host, and within-host). The x axes of different panels differ by method for inferring gene expression, which are those shown in Figure 2B: (1) 'Mass spec' refers to $\log_{10}$(iBAQ prop), where iBAQ prop is the proportion of the maximum intensity-based absolute quantification (iBAQ) observed; (2) 'Upstream peak' is the peak in ribosome profiling read depth upstream of the gene start site, measured for reads of length 29-31nt (Figure 2A); and (3) 'In-frame mean' is the mean read depth at Codon pos 1, measured for reads of length 30nt (Figure 2B). A given gene (color) has only one selection value (y-axis) per evolutionary level, and only one gene expression value (x-axis) per method for determining expression. Selection was calculated as d_N/d_S (π_N/π_S) except for the overlapping gene *ORF9b*, where the OLG-appropriate measure d_{NN}/d_{NS} (π_{NN}/π_{NS}) was used (see Methods).

2
Figure 5—figure supplement 2. SARS-CoV-2 between-host nucleotide diversity and allele
frequencies as a function of time. Analyses refer to human SARS-CoV-2 consensus sequences
obtained from GISAID. Results show sliding windows of 14 days (step size=7 days) representing 13
time points since the first GISAID sample was collected (EPI_ISL_402123 on 12/24/2019). Regions
with overlapping genes were excluded from estimates of π. Shaded regions show standard error of
mean π_N, π_S, or π_N/π_S (10,000 bootstrap replicates, codon unit). **(A)** Nonsynonymous (π_N) and
synonymous (π_S) nucleotide diversity. **(B)** π_N/π_S, where the horizontal dashed gray line denotes the
π_N/π_S ratio expected under neutrality (1.0). **(C)** Diversity of sampling locations measured as entropy,
defined as $-\sum p*\ln(p)$, where p is the number of distinct (unique) locations or countries reported for a
given window (Ewens and Grant 2001). **(D)** Allele frequency trajectories of the *ORFd*-LOF mutation,
colored by continent and limited to sufficient sample sizes (Supplementary Table 19).
14

2
Figure 6. Between-taxa sliding window analysis of natural selection on overlapping frames of *ORF3a*.
Pairwise sliding window analysis (windows size=50 codons; step size=1 codon) of selection across
members of the species *Severe acute respiratory syndrome-related coronavirus*. OLG-appropriate
d_N/d_S values were computed using OLGenie (Nelson et al. 2020), a method that is conservative (non-
conservative) for detecting purifying (positive) selection, and a Jukes-Cantor correction for multiple hits
was employed. On the left-hand side, each genome is compared to SARS-CoV-2 in the ss12 reading
frame of *ORF3a*, which contains *ORF3d* (Table 1 and Figure 2C). This frame shows evidence for
purifying selection specific to the *ORF3d* region that is limited to the comparison with pangolin-CoV
GX/P5L. On the right-hand side, this analysis is repeated for the ss13 reading frame of *ORF3a*, which
contains *ORF3c* and *ORF3b* (Table 1 and Figure 2C), this time with respect to SARS-CoV, where full-
length *ORF3b* is functional. This frame shows constraint across much of this gene and virus species,
and that *ORF3c* in particular is deeply conserved.
15
16   **Figure 6—source data 1.** SARS-CoV-2-ref_ORF3a_ss12_windows.txt. OLGenie sliding window d_N/d_S
17   results for comparisons to SARS-CoV-2 in the frame of *ORF3d*, shown on the left side of Figure 6.
18
19   **Figure 6—source data 2.** SARS-CoV-ref_ORF3a_ss13_windows.txt. OLGenie sliding window d_N/d_S
20   results for comparisons to SARS-CoV in the frame of *ORF3b*, shown on the right side of Figure 6.
21

3   **Figure 6—figure supplement 1.** Between-taxa sliding window of genes overlapping *N*. Pairwise
4   OLGenie analysis of the N gene's ss13 reading frame across members of the species *Severe acute*
5   *respiratory syndrome-related coronavirus*. Details as in Figure 6, except only the ss13 frame is shown.
6
7

Figure 7. Pandemic spread of the EP+1 haplotype and the hitchhiking of *ORF3d*-LOF. The mutational
path leading to EP+1+LOF is shown in the upper panel. Cumulative frequencies of haplotypes in
samples from Germany and five other countries with the most abundant sequence data are shown in
the lower panel. Countries are grouped into early founder (left) and late founder (right) based on the
presence or absence of SARS-CoV-2 samples from January, respectively. In the early founder group,
EP–3 (gray) is observed earlier than other haplotypes in France and the US, and EP (red) is observed
early in Germany, giving them the advantage of a founder effect. However, neither EP nor EP–3
dominate later spread; instead, EP+1 (yellow) and EP+1+LOF (blue) increase much faster despite their
later appearance in these countries. In the late founder group, multiple haplotypes appear at
approximately the same time, but EP-3 and EP spread more slowly. The green dashed line shows the
combined frequencies of EP+1 and EP+1+LOF (yellow and blue, respectively). Note that EP–1 is never
observed in our dataset.

1
2
3

Figure 8. High-frequency within-host mutations. For each site, mutations that occur in more than 2.5% of samples are shown, limiting to samples where the major allele matches Wuhan-Hu-1 (differs by site). The y-axis shows the proportion of such samples having the indicated minor (derived) allele. Each locus has up to three possible single-nucleotide derived alleles compared to the reference background. Open circles (black outlines) show the proportion of samples having any of the three possible derived alleles ("All"), while solid circles (color fill) show the proportion of samples having a specific derived allele (equivalent if only one variant is observed). For most sites, only one derived mutation type (e.g., C→U) was observed across all samples. Precluding co-infection by multiple genotypes and sequencing errors, derived mutations occurring in more than one sample (y-axis) must be identical by state but not descent (i.e., recurrent). Genome positions are plotted on the x-axis, with distinct genes shown in different colors and overlapping genes shown as black blocks within reference genes. Nonsynonymous and nonsense mutations ("Nonsyn") are indicated with a red dot. Source data available in supplementary material.

Figure 8—figure supplement 1. Recurrent nonsynonymous mutations observed in multiple human hosts. Histogram of the derived allele frequency (DAF) of the four most common recurrent within-host protein-coding mutations across 401 samples (bin width=0.01). Mutation A404U introduces a premature stop codon in *nsp1*, whereas the remainder are nonsynonymous.