BRAF^V600E induces reversible mitotic arrest in human melanocytes via microRNA-mediated

suppression of AURKB.

Andrew S. McNeal[1], Rachel L. Belote[2,], Hanlin Zeng[2,], Marcus Urquijo[2], Kendra Barker[2], Rodrigo

Torres[1,3], Meghan Curtin[2], A. Hunter Shain[1,3], Robert H. I. Andtbacka[2,4], Sheri L. Holmen[2,4,5], David H.

Lum[2], Timothy H. McCalmont[6], Matthew W. VanBrocklin[2,4,5], Douglas Grossman[2,5,7], Maria L. Wei[1,3,8],

Ursula E. Lang[1,3,6], and Robert L. Judson-Torres[2,5,7,†]

[1] Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, San

Francisco, CA, 94158, USA.

[2] Huntsman Cancer Institute, University of Utah, Salt Lake City, UT, 84112, USA.

[3] Department of Dermatology, University of California San Francisco, San Francisco, CA, 94115, USA.

[4] Department of Surgery, University of Utah, Salt Lake City, UT, 84132, USA.

[5] Department of Oncological Sciences, University of Utah, Salt Lake City, UT, 84132, USA.

[6] Department of Pathology, University of California San Francisco, San Francisco, CA, 94143, USA.

[7] Department of Dermatology, University of Utah, Salt Lake City, UT, 84102, USA.

[8] San Francisco Veterans Affairs Medical Center, San Francisco, San Francisco, CA, 94121, USA.

† To whom correspondence should be addressed. Tel: +1 801-213-8436; Email:

judsontorreslab@gmail.com

20
Abstract

Benign melanocytic nevi frequently emerge when an acquired *BRAF^V600E* mutation triggers unchecked

proliferation and subsequent arrest in melanocytes. Recent observations have challenged the role of

oncogene-induced senescence in melanocytic nevus formation, necessitating investigations into

alternative mechanisms for the establishment and maintenance of proliferation arrest in nevi. We

compared the transcriptomes of melanocytes from healthy human skin, nevi, and melanomas arisen from

nevi and identified a set of microRNAs as highly expressed nevus-enriched transcripts. Two of these

microRNAs – MIR211-5p and MIR328-3p – induced mitotic failure, genome duplication and proliferation

arrest in human melanocytes through convergent targeting of AURKB. We demonstrate that *BRAF^V600E*

induces a similar proliferation arrest in primary human melanocytes that is both reversible and conditional.

Specifically, *BRAF^V600E* expression stimulates either arrest or proliferation depending on the differentiation

state of the melanocyte. We report genome duplication in human melanocytic nevi, reciprocal expression

of AURKB and microRNAs in nevi and melanomas, and rescue of arrested human nevus cells with

AURKB expression. Together, our data describe an alternative molecular mechanism for melanocytic

nevus formation that is congruent with both experimental and clinical observations.

Introduction

Cutaneous melanoma is a potentially deadly skin cancer arising from the pigment-producing melanocytes

of the human epidermis. An activating mutation in the BRAF proto-oncogene (BRAF[V600E]) drives over half

of all cutaneous melanomas[1–3]. Yet, when a melanocyte acquires a BRAF[V600E] mutation, the cell does not

immediately transform to cancer. Instead, it usually undergoes clonal proliferation followed by stable

arrest resulting in a benign skin lesion known as a melanocytic nevus or "mole"[3–6]. Despite the continued

expression of BRAF[V600E], the majority of melanocytic nevi remain innocuous for the lifespan of the

individual, suggesting that nevus cells have robust intrinsic defenses against hyperproliferation. By

inducing hyperproliferation and subsequent arrest, the BRAF[V600E] mutation elicits divergent, biphasic

phenotypes within a single cell – a poorly understood phenomenon. Characterization of the mechanisms

and environmental factors that distinguish the two phenotypes could illuminate candidate strategies for

chemoprevention of melanocytic nevus formation or interception of melanoma initiation.

The BRAF[V600E] mutation is clonally present in ~80% of human melanocytic nevi[3] and melanocyte-specific

expression of the oncogene is sufficient to induce nevus formation in mice[6–8], suggesting that BRAF[V600E]

drives proliferation arrest in melanocytes. The prevailing theory to explain this seemingly paradoxical role

for BRAF[V600E] is oncogene-induced senescence (OIS)[9]. Cellular senescence is defined as the permanent

cell-cycle arrest of previously replication-competent cells[10,11]. Senescence is associated with a variety of

molecular hallmarks including elevated expression of p16[INK4A] and other cyclin-dependent-kinase

inhibitors, TP53, H2AX, lysosomal beta-galactosidase and senescence-associated secretory proteins

(SASP) as well as a DNA content profile indicative of arrest in the G0/G1 phases of the cell cycle[11–16].

The concept of OIS as a barrier to tumorigenesis derived from observations of a durable proliferation

arrest induced by overexpression of oncogenic RAS or RAF in immortalized human fibroblasts[12,17].

More recent studies have questioned whether the term "senescence" aptly describes proliferation arrest

of melanocytic nevi. Cell cycle re-entry that accompanies nevus recurrence[18], eruption[19], or

transformation to primary melanoma suggests that the nevus arrest phenotype is reversible rather than

permanent. Similarly, the expression of senescence markers do not readily distinguish human

melanocytic nevi from primary or transformed melanocytes in humans (16[INK4A], TP53, H2AX and beta-

galactosidase) or mice (any known hallmark of senescence)[8,20,21]. Setting aside the permanence of OIS,

melanocytic nevus arrest was largely thought to be driven by induction of p16[INK4A] expression from the

CDKN2A gene. Indeed, early studies demonstrated that BRAF[V600E] induces p16[INK4A] expression and that

melanocytic nevi frequently express high levels of p16[INK4A] [9]. Despite this, p16[INK4A] expression is neither

ubiquitous across melanocytic nevi, nor uniform within the cells of a single nevus[22,23], and neither knock

down of the 16[INK4A] transcript nor ablation of the CDKN2A locus prevents the onset or rescues BRAF[V600E]-

induced proliferation arrest[9,23–25]. BRAF[V600E] also stimulates expression of p15[INK4B] - the translational

product of the CDKN2B gene[25] which is situated adjacent to the CDKN2A locus - suggesting BRAF[V600E]

might orchestrate arrest via multiple cyclin dependent kinase inhibitors that induce arrest in the G1 phase

of the cell cycle, prior to genome synthesis. While the loss of the *CDKN2A/B* locus is a formative event in

melanoma progression, genetic evolution studies of melanoma progression suggest that selection against

the *CDKN2A/B* locus occurs after BRAF[V600E] melanocytes have already escaped nevus-associated

arrest[23,26]. Collectively, these data complicate the straightforward model wherein BRAF[V600E] induces the

expression of cell cycle regulators leading to proliferation arrest, which is later circumvented by the

genetic loss of those regulators.

Given these observations, we reasoned that the arrest response of human melanocytes to BRAF[V600E]

might be distinct from previous reports of BRAF[V600E]-induced OIS in other cell types. Specifically, we

hypothesized that nevus-associated arrest is conditional and coordinated by reversible changes in gene

expression. Here, we identify a transcriptional program that is consistently elevated in melanocytic nevi

and includes microRNAs (miRNAs) previously shown to have diagnostic value[27,28]. In parallel, we observe

that, unlike RAF-induced OIS in fibroblasts, BRAF[V600E]–induced arrest in human melanocytes is

reversible and conditional. Merging these two observations, we discover that in melanocytes, regulation

of miRNAs coupled to differentiation status permits toggling between BRAF[V600E]-driven hyperproliferative

to arrested phenotypes.

Results

Melanocytic nevus- enriched miRNAs induce mitotic failure and proliferation arrest in primary

human melanocytes

Since melanocytic nevi do not consistently express known senescence markers, we sought to

characterize the transcripts that are specifically elevated within nevus melanocytes (Fig. 1A). We

analyzed previously published datasets comparing RNA expression between benign and malignant

melanocytic lesions[26,28]. Each clinical specimen contained matched RNA samples derived from

melanocytic nevi and melanomas that arose from those nevi. We reasoned that down-regulated

transcripts within these newly-formed melanomas could represent the most immediate barriers to

melanocyte growth. Of the differentially expressed genes (adjusted p value <0.05), six of the top ten with

elevated expression in nevi were non-coding miRNAs (Fig. 1B and Table S1). Among these were

miRNAs previously shown as enriched in nevi compared to melanomas, including MIR211-5p, MIR125A-

5p, MIR125B-5p and MIR328-3p[28,29]. To determine whether elevated expression of these miRNAs

distinguished nevus melanocytes from normal epidermal melanocytes, we profiled melanocytes from

fresh healthy skin and nevi (Table S2). Three of the miRNAs (MIR211-5p, MIR328-3p, MIR125B-5p) were

also more highly expressed in melanocytic nevi as compared to healthy melanocytes or melanomas - an

expression pattern consistent with enrichment in arrested melanocytes - whereas one miRNA (MIR125A-

5p) presented a pattern consistent with lineage specific expression that is lost upon

112
113

Figure 1: Melanocytic nevus enriched miRNAs induce arrest in primary human melanocytes

(A) Schematic of hypothesized plastic nature of BRAF[V600E]- and nevus- associated proliferation arrest. Expression of the predicted nevus enriched transcriptional program is highlighted in blue. **(B)** Scatter plot depicting gene expression combining publicly available sequencing datasets totaling 14 melanoma-arising-from-nevus specimens, dissected into benign and malignant portions (28 total samples). Plotted are the adjusted p values (DESeq2) comparing matched nevus and melanoma samples against average normalized expression in nevus portion. Only genes exhibiting both high expression and significant differential expression in nevus portions are plotted (see Methods). The full list of genes is included as Table S1. **(C)** Left two columns, the relative expression of indicated miRNAs in freshly isolated human melanocytes from BRAF[V600E] melanocytic nevi (n = 6) compared to BRAF[WT] healthy skin melanocytes (n = 8). Right two columns, the relative expression of indicated miRNAs in melanocytes dissected from FFPE nevus (n = 7) or melanoma (n = 7) specimens. Data represent mean and standard deviation of normalized RNA sequencing read counts relative to nevus-derived samples. **(D)** Mean and standard deviation for population fold change over seven days of melanocytes nucleofected with indicated miRNA mimics compared to non-targeting control (Control RNA) (n = 3). **(E)** Mean and standard deviation for fraction of EdU positive cells 4 days after nucleofection with indicated concentrations of miRNA mimics (n= 3). P values indicate comparisons to concentration matched control. **(F)** Representative QPI images of melanocytes nucleofected with indicated miRNA mimics. Pixel color indicates low (black), mid (blue) or high (red) optical density. Scale bar = 200µm. **(G)** Mean and standard deviation for relative QPI-derived cell number over time (n = 3). **(H)** Mean and standard deviation for QPI-derived cell death counts as percentage of total cells imaged on day 1 (n = 3). Cells treated with 0.1 µM staurosporine were included as a control (n = 1) for induction of cell death. **(I)** Mean and standard deviation for percent Annexin V/PI positive cells nucleofected with indicated miRNA mimics (n = 3). Cells treated with 5µg/ml mitomycin C were included as a control (n = 3) for induction of apoptosis. **(J)** Mean and standard deviation for relative cell number over time (n = 3). **(K)** Mean and standard deviation for relative QPI-derived dry mass per cell (n = 3). **(L)** Representative histograms of DNA content profiling via propidium iodide incorporation measured by flow cytometry 5 days post- nucleofection of miRNA mimics. **(M)** Mean and standard deviation for percent of cells in indicated phases of cell cycle based upon profiles as in (K) (n = 3-6, individual datapoints shown). P values for D-L calculated by unpaired t tests comparing experimental to Control RNA. N.S. = not significant (P ≥ 0.05)

transformation (Fig. 1C). We conclude that MIR211-5p, MIR328-3p, MIR125B-5p are transcripts specifically enriched in arrested melanocytic nevi.

To investigate whether elevated expression of MIR211-5p, MIR328-3p, or MIR125B-5p induce melanocytic arrest, we introduced synthesized mimetics of the mature miRNAs into primary human melanocytes and tracked proliferation over seven days in culture. The introduction of either MIR211-5p or MIR328-3p resulted in significantly fewer melanocytes compared to a non-targeting control (Fig. 1D). Increased concentrations of these mimics resulted in a dose-dependent decrease in EdU uptake, consistent with a cell cycle defect (Fig. 1E). To better characterize the effect of MIR211-5p or MIR328-3p expression on cell cycle and cell viability, we performed time-lapsed quantitative phase imaging (QPI). We and others have previously established QPI as a method for adherent cell cytometry that can simultaneously measure melanocyte proliferation, cell death, and cell cycle stage with single cell resolution[30–32]. Images were taken every hour for 5 days post-nucleofection (Fig. 1F). Cells expressing MIR211-5p or MIR328-3p exhibited significantly reduced growth curves (Fig. 1G). We observed no evidence of increased cell death (Fig. 1H), and confirmed this result with Annexin V staining (Fig. 1I). Experiments performed using three additional melanocyte preparations counted over seven days post

nucleofection yield similar results (Fig. 1J). Thus, MIR211-5p and MIR328-3p induce proliferation arrest in

human melanocytes.

Further characterization of the QPI images revealed a distinct increase in intra-cellular dry mass in

proliferation-arrested conditions (Fig. 1K). The dry mass of a cell is a measurement of total biomass

(DNA, RNA, proteins, lipids, etc) and is quantitatively measured by QPI. Relative dry mass fluctuates with

cell cycle stage in a predictable manner[30]. The 20-40% increase we observed here is characteristic of

cells that have doubled their DNA content but failed to complete mitosis. We confirmed this finding by

measuring DNA content of the arrested populations (Fig. 1L). We observed significant increases of cells

with both 4N DNA content and greater than 4N DNA content, indicative of cytokinetic failure following

DNA replication (Fig. 1M). Consistent with this interpretation, time-lapsed imaging revealed incidences of

tripolar mitosis and anaphase bridging in miRNA-expression melanocytes – both processes associated

with cytokinetic failure[33] - which were not observed in control conditions or standard culture (Video 1 and

Figure 1 – figure supplement 1). We conclude that MIR211-5p and MIR328-3p induce mitotic failure and

proliferation arrest when introduced into primary human melanocytes.

Video 1: Cytokinesis failure associated with MIR211-5p expression.

Four day time-lapse quantitative phase imaging of cytokinesis failure in primary melanocytes transduced with MIR211-5p.

Figure 1-figure supplement 1

Still frames from Video 1 documenting representative cytokinesis failure in primary melanocytes transduced with MIR211-5p. Yellow arrow heads follow a single cell and its progeny through a tripolar mitosis. Red arrow head follows an anaphase bridge connecting two daughter cells that then remerge.

Inhibition of AURKB by MIR211-3p and MIR328-5p restricts proliferation in nevi

To investigate the mechanism by which MIR211-5p and MIR328-3p induce arrest in primary human melanocytes, we applied our previously established pipeline for identification of relevant miRNA targets (Fig. 2A)[34]. First, we nucleofected MIR211-5p, MIR328-3p, or control RNA mimics into primary human melanocytes freshly isolated from four independent donors and conducted RNA sequencing (Table S3). We next selected all genes that were significantly downregulated by either miRNA as compared to the control RNA, and further refined the set to the computationally predicted targets of MIR211-5p and/or MIR328-3p. Finally, we nucleofected siRNAs targeting the 115 resulting genes and assayed for reduced EdU incorporation that would phenocopy MIR211-5p or MIR328-3p expression (Table S4). As expected, computationally predicted targets of MIR211-5p and MIR328-3p were enriched in the set of downregulated genes following MIR211-5p or MIR328-3p nucleofection, respectively (Fig. 2B). AURKB, GPR3, and MAPKAPK2 showed the greatest magnitude of repression due to miRNA expression with significant EdU reduction upon siRNA-mediated knockdown (Fig. 2C). Of these, AURKB (predicted MIR211-5p target) and GPR3 (predicted MIR328-3p target), were significantly inhibited upon nucleofection of either miRNA (Fig. 2D). We reasoned these genes might represent convergent nodes driving proliferation arrest. Further exploration of the AURKB transcript revealed a potential MIR328-3p binding site just before the 3' untranslated region (Fig. 2E). In contrast, we were unable to identify a candidate MIR211-5p binding site in the GPR3 transcript, suggesting indirect inhibition of this gene by MIR211-5p.

Figure 2: Inhibition of AURKB by MIR211-3p and MIR328-5p restricts proliferation in nevi

(A) Schematic of experimental and computational pipeline for identifying targets of MIR211-3p and/or MIR328-3p responsible for proliferation arrest. **(B)** GSEA analysis comparing predicted MIR211-5p and MIR328-3p target mRNAs to changes in gene expression after nucleofection with indicated mimics compared to nucleofection control. NES = normalized enrichment score. P = nominal P value. Table S3 provides full differentially expressed gene lists. **(C)** Plot of siRNA screen results for phenocopying proliferation arrest versus log2 fold change in expression after miRNA mimic nucleofection. Strictly

standardized mean difference (SSMD) reports the magnitude and significance of inhibition of Edu incorporation (n = 3, see
Table S4 for list of genes and screen results). Dotted lines indicate cut-off values for further evaluation. Orange points
indicate genes meeting cut-off criteria. **(D)** Mean and standard deviation for expression of indicated genes after
nucleofection of indicated mimics (n = 4). P values calculated by unpaired t tests compared to No RNA. **(E)** Schematics
showing predicted binding sites of MIR211-5p and MIR328-3p in AURKB and GPR3 transcripts. **(F)** Representative images
of immunohistochemical staining for AURKB (brown chromagen) and GPR3 (red chromagen) expression in FFPE samples
of nevi and melanoma. Scale bars = 50μm. **(G)** Mean and standard deviation for immunohistochemical staining scores as
in (F) for cohorts of 11 nevi and 11 melanomas. P values calculated by unpaired t tests. **(H)** Mean and standard deviation
for read counts for cohorts of 11 nevi and 12 melanomas. P values calculated by unpaired t tests. **(I)** Design of pLVX vectors
expressing either AURKB or GPR3. **(J)** Western Blot of AURKB expression in human melanocytes with or without lentiviral
AURKB overexpression. HSP90 is the loading control. Source data provided as supplemental files (Fig_2_J_Source). **(K)**
Representative photomicrographs (20X) of immunofluorescence for GPR3 (green) or (DAPI) (purple) in human melanocytes
with GPR3 lentiviral overexpression (GPR3 OE) or without (Control) over-expression. **(L)** Mean and standard deviation for
cell number fold change over 7 days after nucleofection with indicated miRNA mimics in melanocytes transduced with
lentiviral constructs expressing zsGreen, AURKB, or GPR3 (n = 6 melanocyte preps, type 2 t test, N.S. = not significant (P
≥ 0.05)). P values calculated by unpaired t tests.

To determine whether decreased AURKB and/or GPR3 expression was associated with melanocytic nevi,

we acquired 23 clinical specimens and measured both the transcript and protein expression of these

genes (Fig. 2F-H). The expression of both AURKB and GPR3 protein and mRNA were enriched in

melanomas as compared to nevi (Fig. 2G-H). Together, these data demonstrate an inverse expression

between two miRNAs (MIR211-5p and MIR328-3p) and two targeted mRNAs (AURKB and GPR3) that

toggles concurrently with the transition of arrested melanocytic nevi to melanomas. To further investigate

whether either mRNA was sufficient to rescue miRNA-mediated arrest, primary melanocytes were

transduced with constructs expressing either zsGreen (control), AURKB or GPR3 (Fig. 1I-K) then

subsequently nucleofected with either MIR211-5p or MIR328-3p. Expression of either mRNA partially

rescued MIR211-5p-induced arrest, while AURKB, but not GPR3, partially rescued MIR328-3p-induced

arrest (Fig. 2L). Thus, while other MIR211-5p and MIR328-3p targets, including GPR3, may play a role in

the induction of melanocyte proliferation arrest, these data suggest that AURKB inhibition is a critical

convergent node of both miRNAs in facilitating proliferation arrest in nevi. This interpretation is further

supported by the well-established role of AURKB as an essential orchestrator of successful mitosis[35].

BRAF^V600E induces a reversible and conditional proliferation arrest in human melanocytes

We hypothesized that AURKB inhibition by MIR211-5p and MIR328-3p may be the underlying

mechanism by which BRAF[V600E] induces proliferation arrest in melanocytes. BRAF[V600E] expression has

been demonstrated to result in an accumulation of growth-arrested fibroblasts with 2N DNA content,

presumably in a permanent G0/G1 senescent state[9,12,17]. In contrast, upon overexpression of MIR211-5p

or MIR328-3p, we observed an accumulation of arrested melanocytes with ≥4N DNA content, and a

significant decrease in cells with 2N DNA content (Fig. 1M). To determine the DNA content profile of

primary human melanocytes under BRAF[V600E]-induced arrest, we utilized an established lentiviral

construct for dose-responsive doxycycline inducible BRAF[V600E] expression (Fig. 3A)[25]. We monitored

melanocyte growth under a titration of doxycycline and identified the lowest concentration (15.6 ng/mL)

that induced sustained arrest after 40 hours exposure (Fig. 3B). Similar to our observations for miRNA-

induced arrest, relative per cell dry mass increased by 50% concurrent with BRAF^{V600E} induced- arrest

(Fig. 3C). Cell cycle profiling confirmed a significant and dose-dependent accumulation of cells with ≥4N

Figure 3: BRAF^{V600E} induces a reversible and conditional arrest in human melanocytes

(A) Representative (of n = 3) Western Blot analysis of BRAF^{V600E}, phospho-ERK1/2, and HSP90 (loading control) in
diBRAF^{V600E} melanocytes in response to increasing concentration of doxycycline. Source data provided in supplemental
files (Fig_3A_Source). **(B)** Mean and standard deviation for relative QPI-derived cell number at indicated hours of

diBRAF[V600E] melanocytes in response to increasing concentration of doxycycline (n = 6). P values from unpaired t tests

comparing to 0 ng/mL condition. **(C)** Mean and standard deviation for relative QPI-derived dry mass per cell of diBRAF[V600E]

melanocytes in response to increasing concentration of doxycycline at indicated hours (n = 6). P values from unpaired t

tests comparing to 0 ng/mL condition. **(D)** Representative histograms of DNA content profiling via propidium iodide

incorporation measured by flow cytometry 5 days after treatment with indicated molecules. AZD2811 included as a control

for induced cytokinesis failure. **(E)** Mean and standard deviation for percent of cells in indicated phases of cell cycle after

treatment with doxycycline based upon profiles as in (D) (n = 4). AZD2811 (n = 4) is a control for induced cytokinesis failure.

PD0332991 (n = 2) is a control for G0/G1 arrest. P values from unpaired t tests comparing to Control (no treatment)

condition. **(F)** Mean and standard deviation for relative QPI-derived cell number of diBRAF[V600E] melanocytes at indicated

hours after treatment doxycycline. Dotted line indicates change in media to either refresh doxycycline or add media without

doxycycline (n = 3). P values from unpaired t tests comparing final timepoints. **(G)** Mean and standard deviation for cell

number as percent of control of diBRAF[V600E] melanocytes cultured with or without TPA after 5 days of treatment with

indicated concentrations of doxycycline. P values from unpaired t tests comparing to 0 ng/mL conditions. Baseline

population doubling times were 3.2 days (+TPA) and 4.2 days (TPA-free). **(H)** Mean and standard deviation for relative

QPI-derived cell number of diBRAF[V600E] melanocytes at indicated hours after treatment doxycycline. Dotted line indicates

change in media to either refresh TPA-containing media or add media without TPA (n = 3). P values from unpaired t tests

comparing final timepoints. **(I)** Schematic of experimental design testing the reversibility of growth arrest after 30 days

exposure to high doxycycline (60 ng/mL). Days of live imaging represented in (J) and (K) are indicated by grey bars. **(J)**

Mean and standard deviation for relative QPI-derived cell number of diBRAF[V600E] melanocytes at 14 days after addition of

doxycycline (n = 3). P values from unpaired t tests comparing final timepoints. **(K)** Mean and standard deviation for relative

QPI-derived cell number of diBRAF[V600E] melanocytes at 30 days after addition of doxycycline (n = 3). Media was changed

just prior to imaging to either refresh doxycycline (blue), +TPA media without doxycycline (black) or add TPA-free media

with doxycycline (red). P values from unpaired t tests comparing final timepoints.

DNA content and a decrease in cells with 2N DNA content (Fig. 3D-E). The cell cycle profiles were

notably dissimilar to that of melanocytes treated with a pharmacologic CDK4/6 inhibitor (PD0332991,

Palbociclib)[36] known to induce G0/G1 arrest. Treatment with a pharmacologic AURKB inhibitor (AZD2811,

Barasertib)[37] that induces G2/M arrest and mitotic failure also resulted in an accumulation of melanocytes

with ≥4N DNA content and a decrease in cells with 2N DNA content.

In addition to the accumulation of 2N cells, previous reports of RAF expression in fibroblasts show that

transient expression can induce a durable arrest phenotype[17]. To test the durability of BRAF[V600E]-induced

arrest in human melanocytes, we induced doxycycline-dependent BRAF[V600E] expression for 48 hours and

monitored cell number for an additional two days to confirm arrest. Media was then replaced to remove

doxycycline. Within twelve hours of doxycycline removal, the previously arrested melanocytes began to

proliferate (Fig. 3F), demonstrating that BRAF[V600E]-induced proliferation arrest in melanocytes is

reversible and dependent on continued oncogene expression.

Nevi recurrence after excision[18] and eruption upon external stimuli[19], together with our observation that

BRAF[V600E] induced arrest is reversible in melanocytes, suggested that BRAF[V600E]-induced arrest may

depend on secondary signals from the microenvironment. Tetradecanoylphorbol acetate (TPA) is a

protein kinase C (PKC) activator commonly used as a mitogen in primary melanocyte media to substitute

for endothelin receptor type B activation[38]. In contrast, TPA also inhibits the cell cycle of melanoma

cells[39,40]. BRAF^{V600E} expression permits growth of the Melan-A cell line in the absence of TPA[41]. We

therefore wanted to determine the relative contributions of TPA and BRAF^{V600E} to the BRAF^{V600E}-induced

primary melanocyte arrest *in vitro*. We induced BRAF^{V600E} expression in primary human melanocytes in

two medias – one containing TPA (+TPA) or one containing the endothelin receptor type B ligand,

endothelin-1 (TPA-free) and observed divergent dose-responsive phenotypes (Fig. 3G). In the presence

of TPA, BRAF^{V600E} expression inhibited cell proliferation. By contrast, in TPA-free media, BRAF^{V600E}

expression induced proliferation. We next subjected melanocytes to the arrest-inducing conditions of

doxycycline treatment in +TPA media for four days, then replaced with doxycycline-containing, TPA-free

media. We observed a rapid shift to the proliferative phenotype even in the continued presence of

doxycycline (Fig. 3H). To determine whether the observed reversibility of BRAF^{V600E}-induced growth

arrest was due to insufficient duration or degree of expression of the oncogene, we exposed melanocytes

to a high concentration of doxycycline (60ng/mL) for 30 days prior to switching medias (Fig. 3I).

Melanocytes in +TPA media remained arrested for the duration of doxycycline treatment then grew

readily upon removal of either doxycycline or TPA (Fig. 3J-K). From these results, we conclude that

BRAF^{V600E} expression induces divergent and interconvertible phenotypes – either hyperproliferation or a

reversible arrest – in human melanocytes conditional on extrinsic stimuli.

BRAF^{V600E}-induced arrest is dependent on melanocyte growth conditions and differentiation state

To investigate the relative contributions of BRAF^{V600E} and TPA to the expression of MIR211-5p and

MIR328-3p, we conducted small RNA sequencing of melanocytes grown in +TPA or TPA-free media and

with or without BRAF^{V600E} expression. The expression of MIR328-3p was relatively stable, demonstrating

less then 2-fold changes across conditions (Fig. 4A-B, Table S5). In contrast, MIR211-5p was one of the

most differentially expressed genes in +TPA conditions compared to TPA-free conditions (adj. P = 6.8E-

23) (Fig. 4A-B, Table S5). Induction of BRAF^{V600E} in +TPA conditions did not further alter expression of

either miRNA. Direct comparison of the arrested melanocytes in +TPA +Dox conditions to the proliferative

melanocytes in -TPA +Dox conditions revealed MIR211-5p as the most significantly downregulated

miRNA in the proliferative condition (Fig. 4B). Thus, in comparison to TPA, BRAF^{V600E} induction had

minimal effect on the expression of the miRNAs. The most notable effect was a decrease of MIR328-3p

expression coinciding with the induction of BRAF^{V600E} in TPA free conditions (Fig. 4A). While MIR328-3p

downregulation likely contributes to increased proliferation in this condition, the more pronounced effect of

TPA on MIR211-5p expression suggests that the mechanisms governing whether BRAF^{V600E} induces

arrest or proliferation are a function of the cell's transcriptional state rather than directly downstream of

BRAF^{V600E} itself.

To better characterize the two transcriptional states influencing the BRAF^{V600E} phenotype, we conducted

mRNA sequencing of the same specimens. Principal component (PC) analysis revealed that 97.4% of the

variance between samples was explained by two principal components – PC1 (81.7% variance)

separated transcriptomes based upon exposure to TPA and PC2 (15.7% variance) separated

transcriptomes based upon BRAF[V600E] expression (Fig. 4C, Table S5). We conducted differential

expression analyses to compare all samples exposed to +TPA versus TPA-free media, regardless of

BRAF[V600E] expression (Fig. 4D, top) and to compare all samples expressing BRAF[V600E], regardless of

TPA exposure (Fig. 4D, bottom). As expected from the PC analysis and similar to the miRNA profiles,

most gene expression variance was due to media conditions with a smaller, but consistent, set of genes

activated by BRAF[V600E].

One initially surprising result of this analysis was the static expression of both MIR211-5p and MIR328-3p

in +TPA conditions, regardless of BRAF[V600E] expression. However, as inhibitors of translation, miRNA

function depends on the expression level of its targets[42,43], effectively functioning as buffers of gene

expression that prevent aberrantly high levels of a transcript[44]. BRAF[V600E] has been previously

demonstrated as an upstream activator of AURKB transcription in melanoma cells[45]. We therefore

reasoned that MIR211-5p expression in +TPA conditions might dampen AURKB activation by BRAF[V600E].

Consistent with this hypothesis, the addition of doxycycline induced AURKB expression in TPA-free /

MIR211-5p low conditions, but this effect was not observed in +TPA / MIR211-5p high conditions (Fig.

4E).

We next conducted gene set enrichment analyses (GSEA) comparing TPA-regulated genes and

BRAF[V600E]- regulated genes to Molecular Signature Database Hallmark gene sets[46]., the KEGG

PATHWAY database[47], and signatures important in BRAF[V600E] signaling[48], human melanocyte

differentiation[49] or melanoma cell state[50–52]. Among the pathways uniquely enriched with BRAF[V600E]

expression were the expected increases in signatures downstream of BRAF[V600E], MAPK (KRAS)

signaling, and glycolysis[53] (Fig. 4F and Table S6). Gene sets uniquely enriched in +TPA media included

signatures associated with cell cycle regulation, such as the G2-M checkpoint, oxidative phosphorylation,

the MITF program, and differentiated melanocytes. In contrast, TPA-free media induced gene sets

associated with cell adhesion and melanocyte progenitor and stem cells. Corroborating this observation,

melanocytes grown in TPA-free media expressed higher levels of AXL (Fig. 4G-H), lower levels of

MLANA, a melanocyte differentiation marker and presented greater activity of aldehyde dehydrogenase,

enzymatic activity associated with stemness (Fig. 4I). These results indicate that exposure to TPA

induces a more differentiated melanocyte phenotype, consistent with previous reports[54–56].

To look for synergist effects of both BRAF[V600E] expression and media conditions, we performed pair-wise

differential expression analyses of each condition (Table S6 and summarized in Figure 4-figure

supplement 1). Generally, most gene sets identified in Figure 4F were regulated by only one of the two

stimuli. For example, genes downstream of BRAF[V600E] signaling were only influenced by doxycycline

exposure. Interesting exceptions to this trend were the inflammatory response, which required both
stimuli; cell cycle regulation gene sets, which were activated by TPA alone but further elevated when
supplemented with doxycycline; and epithelial to mesenchymal transition genes, which were activated by
either stimulus. A subset of melanocyte dedifferentiation gene sets, specifically neural crest genes, the
melanoma invasion program and the AXL program, but not markers of melanocyte stem cells, were
activated by either TPA-free media or BRAF^{V600E} expression, albeit the activation was more pronounced
in the former. This observation suggests that BRAF^{V600E} expression alone does induce some programs
associated with melanocyte dedifferentiation, but with the retention of MIR211-5p expression, the
induction is insufficient to overcome growth arrest.
386
Collectively, our transcriptomic analyses support a model whereby growth in +TPA conditions induces a
more differentiated state in melanocytes coinciding with an increase in MIR211-5p expression (Fig. 4J).
This increased MIR211-5p combined with relatively stable MIR328-3p expression, dampens entry into
mitosis and/or successful completion of mitosis at least in part through AURKB inhibition. The attenuation
of cell division results in an accumulation of cells in G2/M – an effect that is more pronounced in
conditions stimulating accelerated growth, such as with BRAF^{V600E} expression. To test this model, we
transduced doxycycline-inducible BRAF^{V600E} melanocytes in +TPA conditions with lentiviral constructs
constitutively expressing either AURKB (Fig. 2I-J) or miRNA sponges (Fig. 4K) to either MIR211-5p or
MIR328-3p. AURKB expression had no baseline effect on melanocyte growth, but rescued BRAF^{V600E}-
induced arrest (Fig. 4L), suggesting the inability of the oncogene to induce AURKB in this context serves
as a bottleneck in cell division. Also consistent with the model, sponges against either MIR211-5p or
MIR328-3p increased baseline division rate and rescued BRAF^{V600E}-induced arrest, demonstrating that
both miRNAs contribute to inhibition of melanocyte division, with affects that are more pronounced in the
presence of growth signals.
401

Figure 4: BRAF^{V600E}-induced arrest is dependent on melanocyte growth conditions and differentiation state

(**A**) Normalized read counts for MIR328-3p (top) and MIR211-5p (bottom) in TPA-containing (+TPA) or TPA-free media, with or without 15.6 ng/mL doxycycline (Dox). See Table S5 for individual values and exact P values. (**B**) Volcano plots depicting log2 fold change against adjusted P value (-Log10) from differential expression (DE) analysis of small RNA sequencing of indicated comparisons. MIR211-5p labelled and shown in red. MIR328-3p shown in blue. (**C**) First and second

principal components (PC) separating diBRAF^V600E melanocytes cultured in indicated conditions. **(D)** Volcano plots depicting

log2 fold change against adjusted P value (-Log10) from differential expression (DE) analysis of mRNA sequencing of all

+TPA specimens versus all TPA-free specimens (top) or all Dox specimens versus all no-Dox specimens (bottom). **(E)**

Normalized read counts for AURKB normalized to no Dox conditions (n = 3). P values from unpaired t tests. Sequencing

data represented in A-F are from n=3 per condition analyzed with DESeq2. **(F)** Gene set enrichment analysis (GSEA)

comparing DE gene sets from (D) to Molecular Signature Database Hallmark gene sets, KEGG Pathway gene sets and

published signatures of melanocyte signaling and differentiation from Belote, 2021; Tsoi, 2018; Tirosh, 2016; Ryu, 2011;

and Hoek, 2010. All enrichments with adjusted P value < 0.025 are depicted. See Table S6 for all associations. Normalized

enrichment score (NES). Enrichment differences between pathways across the four conditions in (C) are depicted in Figure

4-figure supplement 1. **(G)** Representative photomicrographs of immunofluorescence for MLANA or AXL in human

melanocytes cultured in +TPA or TPA-free conditions. White scale bar = 30μm. **(H)** Quantification of fluorescence intensity

(arbitrary pixel intensity units, a.u.) from experiments represented in (G). Box (median, 25th and 75th percentiles) and

whiskers (10th and 90th percentiles) of 84, 88, 113, and 96 cells respectively. P values from unpaired t tests. **(I)**

Representative plots (top) and quantification (bottom, n = 4) of relative aldehyde dehydrogenase activity of melanocytes

grown in +TPA or TPA-free conditions. **(J)** Schematic of model supported by transcriptomic analyses. TPA induces a more

differentiated state in human melanocytes. Due to increased MIR211-5p and base-line MIR328-3p expression, AURKB

expression is capped, resulting in G2/M accumulation upon BRAF^V600E expression. **(K)** Design of pLVX vectors expressing

microRNA sponges fused to zsGreen. **(L)** Mean and standard deviation for cell number as percent of time point 0 hr after 7

days of culture in +TPA media with or without 15.6 ng/mL Dox (n = 3 per condition). Melanocytes were transduced with

either pLVX-control, -AURKB, -anti-MIR211-5p, or -anti-MIR328-3p. P values from unpaired t tests comparing to matched

no Dox condition (horizontal), Control no Dox condition (vertical in green bar) or Control Dox condition (vertical in red bar).

AURKB inhibition drives melanocytic nevus-associated proliferation arrest

To further investigate whether this mechanism of arrest occurs in clinical nevus specimens, we examined

an independent cohort of 15 nevi for evidence of mitotic failure. Histopathologic analysis revealed that all

nevi within the cohort exhibited binucleation (Fig. 5A). The percent observed binucleated melanocytes

ranged from 3-12% (mean = 6.3%). It is important to note that since binucleation can only be observed in

the fraction of cells sectioned perpendicular to the plane of the adjacent nuclei, 3-12% is likely an under-

estimate. We also observed giant multinucleated cells in 10 of the 15 nevi, indicative of multiple rounds of

DNA synthesis and cytokinetic failure and consistent with previous reports[57].

We sought to determine whether AURKB inhibition was necessary and sufficient for the multi-nucleated

arrest we observed in melanocytic nevi. We obtained three fresh skin biopsies containing nevi,

microdissected the nevus portion, and isolated melanocytes (Fig. 5B). Nevus-derived melanocytes

appeared healthy in culture. Consistent with the histopathologic cohort, co-staining with melanocyte

marker, MLANA, and Hoechst revealed frequent 4N and 8N melanocytes (Fig. 5C). Transduction of

nevus melanocytes with an AURKB and mCherry expressing lentiviral vector at low multiplicity of infection

(MOI) showed that the mCherry positive cells divided significantly more than the their mCherry negative

counterparts (Fig. 5D-E) in cells from all three nevus donors. This result suggests that expression of

AURKB is also rate limiting for the proliferation of human nevus melanocytes, as we observed in primary

melanocytes expressing BRAF[V600E].

To test the sufficiency of AURKB inhibition to limit BRAF[V600E] hyperproliferation in melanoma, we

assessed normalized growth rate inhibition of three cultures - primary human melanocytes, an

established BRAF[V600E] human melanoma cell line, and a short-term BRAF[V600E] human melanoma culture

- when exposed to a pharmacologic AURKB inhibitor (Barasertib). Normalized growth rate measurements

(such as GR_{50}) are independent of baseline cell division rate and are demonstrably more reproducible

than the commonly calculated half maximal inhibitor concentration (IC_{50}) when comparing different cell

cultures[58]. An additional advantage of GR_{50} is the ability to distinguish growth rate inhibition (GR_{50}

approaches 0) versus cell death (GR_{50} is negative). Since GR_{50} calculations require proliferation curves,

Figure 5: AURKB inhibition is critical for BRAF^{V600E}-induced arrest of human nevi

(A) Representative image of 15 H&E-stained melanocytic nevus. Border colors indicate two consecutive magnifications. Arrowheads indicate bi- (red) or multi- (blue) nucleation. **(B)** Example images of skin specimen containing a melanocytic nevus (circled) and phase contrast microscopy of melanocytes isolated from nevus portion. Scale bar = 400μm. **(C)** Images of MLANA (purple) and Hoechst (green) co-staining of melanocytic nevus derived melanocytes in culture. Zoomed images show representative 2N, 4N, and 8N cells. **(D)** Representative of images of QPI coupled with fluorescence microscopy to identify adjacent mCherry -expressing and mCherry-negative melanocytes. Individual cells were identified at time zero, tracked and monitored for division for 96 hrs. **(E)** Mean and individual matched datapoints for the percent of n = 3 mCherry-negative or mCherry-positive cells identified at time zero that divided over 96 hrs. P value from paired t test. **(F)** Mean and standard deviation for 48 hr growth rates of indicated cells treated with Barasertib. P values from unpaired t tests comparing the 30nM samples of melanoma lines to primary melanocytes (n = 3). **(G)** Mean and standard deviation for relative QPI-derived dry mass per 501MEL cell at initial time point (black) compared to 48 hr treatment with Barasertib (grey) (n = 3). P values from unpaired t tests.

we conducted live QPI over 48 hours with a range of Barasertib concentrations (0-300nM). Primary

melanocytes were largely resistant to the compound with reduced growth apparent only at

supraphysiological levels (Fig. 5F). In contrast, both melanoma cultures responded to ten-fold lower

dosage of the compound with GR$_{50}$ approaching zero, indicating primarily arrest, not death. In these

conditions, the per cell dry mass also increased significantly, consistent with a G2/M arrest (Fig. 5G).

Taken together, these data demonstrate that AURKB inhibition limits the rate of BRAF[V600E]-induced

proliferation in human melanocytic nevi and melanomas.

Discussion

Cellular senescence is conventionally defined as a permanent cell-cycle arrest associated with a variety

of molecular markers and an irreversible arrest in the G0/G1 phase of the cell cycle[10–12]. The vast majority

of melanocytic nevi are stably arrested and harbor BRAF[V600E], an oncogenic mutation that induces

cellular senescence when expressed in a variety of mammalian cells[9,12,17]. While these observations

seemingly implicate OIS as the mechanism driving nevus- associated proliferation arrest, a substantial

body of evidence has challenged this paradigm in recent years[8,9,18–25]. Here, we provide further evidence

to challenge the model that acquisition of BRAF[V600E] induces premature cellular senescence in

melanocytes and uncover an alternative mechanism driving nevus formation and stability. We propose

that acquisition of the BRAF[V600E] mutation does not necessitate growth arrest, but instead permits

melanocytes to toggle between hyperproliferation and mitotic arrest. Our transcriptomic analyses

demonstrate that external stimuli can modulate the differentiation state of BRAF[V600E] melanocytes,

orchestrate the expression level of MIR211-5p, and inform which phenotype is triggered by the oncogene

– before or after the oncogene is introduced.

The senescence model is largely rooted in seminal work demonstrating that sustained ectopic expression

of BRAF[V600E] induces proliferative arrest in primary human melanocytes[9] - an experiment reproduced by

several independent labs[24,25,59,60], including our own[23]. In each of these studies, melanocytes were

cultured in TPA containing media. Thus, while observations presented here – that BRAF[V600E] -induced

proliferative arrest is reversible and conditional on the presence of TPA – provide new insights that

challenge the senescence model, they do not contradict the data presented in these previous reports.

One inherent limitation of using any *in vitro* approach to investigate the mechanisms underlaying clinical

phenomena is an inherent incongruity in the magnitude of duration. Melanocytic nevi frequently persist for

several decades, whereas *in vitro* experiments are generally conducted on the order of days to weeks. In

previous *in vitro* studies, the periods of time melanocytes were monitored subsequent to BRAF[V600E]

introduction ranged from one to three weeks. In studies utilizing an inducible RAF in human fibroblasts,

24 hours of expression was sufficient to drive a durable proliferative arrest, even upon removal of the

oncogene[17]. In this study, we induced growth arrest with high levels of BRAF[V600E] for thirty days and

observed the arrest was still reversible. It remains possible that an even longer exposure to BRAF[V600E]

would elicit a permanent proliferative arrest consistent with senescence. It is noteworthy, however, that

the identified mechanisms driving proliferative arrest *in vitro* were also supported by analyses of clinical

nevi specimens.

While supported by analyses of both primary human melanocytes and clinical specimens, one limitation

to this report is the lack of *in vivo* confirmation using animal models. One well-established genetically

engineered mouse model of melanoma expresses BRAF^{V600E} from the endogenous locus in tyrosinase

expressing cells with spatiotemporal control. The predominant phenotype of BRAF^{V600E} activation in this

model is melanocytic hyperplasia, which increases in prominence with BRAF^{V600E} dose[6]. While growth-

arrested melanocytic "nevus-like" lesions also form in this model, they are not associated with markers of

senescence[8]. More recently, an elegant study utilizing a zebrafish model of melanoma demonstrated that

driving BRAF^{V600E} expression from promoters associated with melanocyte progenitors, but not

differentiated melanocytes, resulted in melanoma initiation[61]. These observations are all consistent with

our data and support the conclusion that BRAF^{V600E}-induced proliferation in melanocytes is dependent on

differentiation state.

The presented model offers an explanation for several clinical phenomena. First, melanocytic nevi form

when BRAF^{V600E} drives hyperproliferation followed by proliferation arrest. Second, nevus eruption

involves expansion of previously stable nevi over a short period of time. Third, incomplete excision of

melanocytic nevi can result in subsequent regrowth of the benign lesion[18]. Each event requires

modulation of the BRAF^{V600E} melanocyte between a proliferative and arrested phase. Our data

demonstrate that BRAF^{V600E} melanocytes can indeed oscillate between proliferative and arrested states

and are consistent with each of these phenomena. Our interpretation assumes that once acquired,

BRAF^{V600E} expression is constant and that the environmental context of the melanocytes is variable,

serving as a "gatekeeper" to BRAF^{V600E}-induced proliferation. However, if BRAF^{V600E} expression also

fluctuates, then an equally plausible interpretation of our data is that BRAF^{V600E} serves as a "gatekeeper"

to environmentally induced proliferation. Interestingly, nevus eruption has been observed in metastatic

melanoma patients treated with single agent selective BRAF inhibitors[62–66], consistent with our

observation that proliferation arrest in melanocytes requires sustained expression of the oncogene. A

more thorough understanding of the responsible environmental stimuli in physiologic conditions,

discussed in more detail below, would help to clarify which of the mechanisms – variable BRAF^{V600E},

variable environment, or both – underlie nevus formation, eruption and recurrence.

It is important to note that while our data support a model of reversible proliferation arrest in nevi, we do

not suggest this reversal is commonplace. Unquestionably, the vast majority of melanocytic nevi present

with sustained arrest. However, we do propose that the duration of arrest is dynamic and modulated by

exposure to external signals. Since our data suggest that loss of MIR211-5p and MIR328-3p expression

and restoration of AURKB expression is concurrent with progression to melanoma, identification of the

source of cell-extrinsic signals might suggest new strategies for chemoprevention or therapy. We have

identified TPA, a known activator of PKC signaling, as one extrinsic factor that induces expression of

MIR211-5p and a more dedifferentiated state. Previous studies have identified TPA as either a mitogen,

inhibitor of the G1/S transition, or inhibitor of the G2/M transition, dependent on the context[39,40,67–69]. We

demonstrate that *in vitro*, TPA activation of MIR211-5p is required for BRAF[V600E]-induced mitotic failure.

Yet, the inclusion of TPA in primary melanocyte culture is artificial and the environmental stimulus that

regulates MIR211-5p expression in skin remains unknown. One hypothesis is that PKC activation from

keratinocyte interactions serves as the extrinsic stimulus. MIR211-5p is reported to be transcriptionally

regulated by both MITF[70] and UV exposure[71]. However, TPA-free media containing an endothelin

receptor type B ligand, endothelin-1, did not elicit MIR211-5p expression or BRAF[V600E] arrest, suggestive

that the downstream signaling pathways regulated by the two molecules are not identical. PKC activation

can also result in increased MAPK signaling[72,73] raising the possibility that internal MAPK fluctuations may

play a role in toggling between BRAF[V600E]-induced proliferation versus arrest. The term "mitogenic

window" describes the specific range of oncogene doses that elicit a hyperproliferative phenotype. The

mitogenic window for MAPK signaling activation can be remarkably fine. Here, we conducted a two-fold

dilution series of doxycycline but observed no evidence of a mitogenic window. However, it remains

possible that we simply missed the "sweet spot" in TPA conditions, and that a finer dilution series would

reveal a mitogenic window even in the presence of TPA. Regardless, our data demonstrate that in TPA

conditions the mitogenic window for BRAF[V600E] expression is, at least, exceedingly narrow, whereas in

non-TPA conditions the window is wide open. An important future direction will be to identify the nature

and physiological source of this signal for melanocytic nevi in human skin.

Diminished MIR211-5p/MIR328-3p expression concurrent with increased AURKB expression occurs in a

substantial portion of melanomas. AURKB expression was previously reported as significantly elevated in

melanoma metastases[45] and here we report increased expression in primary melanomas. Similarly,

MIR211-5p expression is consistently observed as lost in melanoma as compared to melanocytic nevi

across independent studies[27–29]. Indeed, MIR211-5p expression levels classify nevi from melanomas with

a high degree of accuracy[27,28]. We have reported on the diagnostic potential of MIR328-3p expression as

well, although multi-study validation was impeded by the absence of hybridization probes in earlier

microarray-based profiling datasets[28]. *In vitro*, we have now demonstrated that inhibition of either miRNA

is sufficient to block BRAF[V600E]-induced arrest in melanocytes. These data suggest that mitotic failure

serves as a barrier to melanoma progression and is consistent with the genome duplication and copy

number alterations associated with invasive melanoma[4,74]. In melanoma, the specific cause of this

genomic instability is unknown, but reentry of cells into cell cycle after mitotic failure has been shown to

cause genome duplication and accumulation of copy number alterations in other cancers[75]. Further

experiments will be necessary to determine whether dysregulation of the spindle checkpoint in nevi

contributes directly to the copy number variations present in melanomas.

In summary, we have identified that BRAF[V600E] induces a reversible arrest in human melanocytes

orchestrated by MIR211-5p/MIR328-3p regulation of AURKB and conditional on the melanocyte

differentiation state. We present a mechanistic model that allows for melanocytic nevus formation,

eruption and recurrence. Moreover, our observations provide an explanation for the genetic duplication

and instability inherent to invasive melanomas and open new directions for novel chemopreventative and

therapeutic strategies.

Competing Interests

R.L.J and M.L.W are inventors on international patent application no. PCT/US2019/023834 concerning

the use of microRNAs investigated in this article as molecular diagnostics for melanoma.

Acknowledgments

This work was supported by the National Institutes of Health (DP5OD019787 and R01CA229896 to

R.L.J., 5F31CA236377 to A.S.M., P30CA042014 to University of Utah), the University of California, San

Francisco Program for Breakthrough Biomedical Research (PBBR) Sandler Fellowship to R.L.J. and the 5

for the Fight fellowship to R.L.J. We acknowledge the use of the HCI Shared Resources for Research

Informatics (RI), Cancer Biostatistics (CB), High-Throughput Genomics and Bioinformatics Analysis

(GBA), and the Biorepository and Molecular Pathology (BMP) Research Histology Section supported by

P30CA042014 awarded to HCI from the National Cancer Institute.

Author Contributions

A.S.M: Investigation, Formal analysis, Writing – original draft, Visualization, Funding acquisition; R.L.B.:

Methodology, Investigation, Resources, Writing – review & editing; H.Z.: Methodology, Investigation,

Validation, Writing – review & editing; M.U.: Investigation; K.B.: Investigation; R.T.: Formal analysis; M.C.

– Formal analysis; A.H.S.: Data curation, Writing – review & editing; R.A., Writing – review & editing:

Resources; S.H.: Conceptualization; D.H.L.: Conceptualization, Writing – review & editing; T.H.M.:

Resources; M.W.V.: Conceptualization; D.G.: Resources, Writing – review & editing; M.W.: Resources,

Writing – review & editing; U.E.L.: Investigation, Formal Analysis, Resources, Writing – review & editing;

R.L.J.: Conceptualization, Formal analysis, Investigation, Writing – original draft; Visualization,

Supervision, Funding acquisition.

Materials and Methods

Transcriptomic profiling

For FFPE samples, histopathologic review, microdissection, targeted exon sequencing, phylogenetic

analysis, and RNA and small RNA sequencing of each tumor area were previously described

(phs001550.v2.p1)[23,26,28]. For this study, small RNA reads were aligned to human reference (hg37) with

Bowtie[76] and small RNA reference groups (miRBase21) were counted. For mRNA sequencing, reads

were aligned to human reference (hg37) with Bowtie2 and counted with HTSeq. Differential expression

analysis was performed using DESeq2[77] with p-values adjusted by Benjamini-Hochberg method (p-adj).

Previously published RNA sequencing datasets from *Shain et al. 2018* and *Torres, et al.* 2020 were re-

analyzed here with DESeq2 the combined for visualization. For RNA profiling from primary melanocytes,

total RNA was extracted using TRIzol Reagent (Thermo Fisher, 15596-026) and further purified with the

RNeasy Power Clean Pro Cleanup Kit (Qiagen, 13997-50) to remove melanin. For small RNAseq of

nevus melanocytes sequencing libraries were constructed with the TailorMix Small RNA Library

Preparation Kit (SeqMatic, CA) and sequencing was performed on the Illumina HiSeq2500 platform at

single-end 50bp. For small RNAseq of cultured BRAF^{V600E} transduced melanocytes, sequencing libraries

were constructed with the Qiagen QIAseq miRNA Library Prep kit and sequencing was performed on the

NovaSeq SP platform at paired-end 50bp. For mRNAseq of microRNA mimic nucleofected melanocytes,

150bp paired end sequencing was conducted by GeneWiz. For mRNAseq of cultured BRAF^{V600E}

transduced melanocytes, sequencing libraries were constructed with the Illumina TruSeq Stranded mRNA

Library Prep kit and sequencing was performed on the NovaSeq S4 platform at paired-end 150bp.

Cell derivation and culture

Benign human melanocytic nevi were excised with informed consent from patient donors at the UCSF

Dermatology clinic (San Francisco, CA) or HCI Dermatology clinic according to Institutional Review

Board- approved protocols. Nevus tissue was minced and enzymatically dissociated in 1mg/mL

collagenase (Sigma 11213857001) and 3.3mg/mL dispase (Sigma, 4942078001) in DMEM (Thermo

Fisher, 10569044) for 1 hour at 37ºC. Melanocytes were further isolated by 5- day exposure to 10µg/mL

G418 (InvivoGen, ant-gn-1). BRAF status was confirmed via sanger sequencing (Quintarabio) using the

primer set: BRAF forward: 5'- GCA CGA CAG ACT GCA CAG GG -3'; BRAF reverse: 5'- AGC GGG CCA

GCA GCT CAA TAG -3'. BRAF wildtype (normal) human melanocytes were isolated from de-identified

and IRB consented neonatal foreskins or adult skin. Foreskin tissue was incubated overnight at 4ºC in

dispase and epithelia were mechanically separated from the dermis the following morning. Epithelial

tissue was minced and incubated in 0.25% trypsin (Gibco 25200056) for 4min at 37ºC. Trypsin was

quenched and tissue centrifuged at 500xg for 5min at room temperature. The cell/tissue pellet was

resuspended in Melanocyte medium (ThermoFisher, M254500) with HMGS (ThermoFisher, S0025) and

plated in low volume.

The 501Mel human melanoma line (Gift from Dr. Boris Bastian, CVCL_4633) were cultured in RPMI with

10% Fetal Bovine Serum (FBS, VWR, S107G), 1% Penicillin-Streptomycin (Gibco, 15140122), 1% L-

Glutamine (Gibco, 25030149), and 1% Non-Essential Amino Acids (Gibco, 11140050). HCIMel019 was

derived from patient-derived xenograft (PDX) tumors propagated in mice. An HCIMel019 (P2)

subcutaneous tumor was resected from mouse, minced in digestion buffer (100µM HEPES (Gibco,

15630-080), 5% FBS (DENVILLE, FB5001-H), 20µg/ml gentamicin (Gibco, 15710-064), 1x insulin (Gibco,

51500-056), and 1mg/ml collagenase IV (Gibco, 17104-019) in DMEM (Gibco, 11965-092)), and digested

overnight at 37˚C. Cells were filtered through a 100µm filter (Falcon, 352360) and red blood cells

removed with RBC lysis buffer (0.5M EDTA, 0.5M KHCO3 (Sigma, 237205), and 5M NH4CL (Sigma,

A9434)). Remaining cells were washed with PBS (Gibco, 10010023), and cultured at 37ºC and 5% CO2

in Mel2 media consisting of 80%MCDB153 (Sigma, M7403), 20% L15 (Gibco, 11415-064), 2% FBS

(DENVILLE, FB5001-H), 1.68mM CaCL (Sigma, C4901), 1x insulin (Gibco, 51500-056) 5ng/ml EGF

(Sigma Aldrich, E9644), 15µg/ml Bovine Pituitary Extract (Gibco 13028-014), and 1x Pen/strep (Gibco,

15070-063).

Small RNA Nucleofection

Human melanocytes were trypsinized (Gibco, 25300062), quenched, and centrifuged at 300xg. Cells

were resuspended in R Buffer (ThermoFisher, MPK1025) at 10,000 cells/µL. 10µL of cell slurry was

mixed with miRIDIAN microRNA mimics (Dharmacon: hsa-MIR211-5p C-300566-03-0005, hsa-MIR328-

3p C-300695-03-0005, MIRControl 1 CN-001000-01-05 or MIRControl 2 CN-002000-01-05); or On-

TargetPlus siRNA (Dharmacon, custom library); at 4µM final concentration and nucleofected using the

NEON Transfection System and protocol (Thermo Fisher, MPK5000).

Generation of Lentiviral vectors

pTRIPZ-diBRAFV600E was a gift from Todd Ridky[25]. pLVX-AURKB (Addgene #153316) and pLVX-

GPR3 (Addgene #153317) were generated by subcloning the respective human cDNA (from Addgene

#100142 and #66350) into the MluI and BamHI sites] of the pLVX-Che-hi3 vector (a gift of Sanford

Simon)[78]. pLVX-anti-MIR211-5p (Addgene #153318), pLVX-anti-MIR328-3p (Addgene #153319), and

pLVX-Che-zsGreen (Addgene #153320) were generated by inserting zsGreen with or without a 3'UTR

into the MluI and XbaI sites of the pLVX-Che-hi3 vector. The 3'UTRs contained 7 tandem microRNA

binding sites to report and inhibit microRNA function, as previously described[34]. ZsGreen was subcloned

from pHIV-zsGreen (gift from Bryan Welm & Zena Werb (Addgene #18121))[79].

Lentiviral transduction

2.75×10^6 HEK293T cells were plated on 10cm tissue culture dishes and grown for ~24 hours in DMEM

with 10% FBS. For each 10cm plate, 5µg lentiviral vector, 3.3µg of pMLV-GagPol and 1.7µg of pVSV-G

packaging plasmids were added to 500µL of jetPRIME buffer and 20µL of jetPRIME transfection reagent

(Polyplus, 712-60), and transfected according to manufacturer's instructions. 48 hours post-transfection,

viral supernatant was collected and filtered using 0.45µm syringe filters (Argos 4395-91). Human

melanocytes seeded at $1.0 \times 10^5 – 2.0 \times 10^5$ cells/well density in 6-well plates were incubated in viral

supernatant with 10µg/mL polybrene (Sigma, TR-1003) and centrifuged at 300xg for 60 min at room

temperature. The viral supernatant was removed and replaced with growth media. Transduced cells were

either selected with puromycin (1μg/mL for 5 days, Sigma P8833) or sorted for mCherry expression using

a BD FACSAria II. Cells expressing pTRIPZ-diBRAFV600E were treated with doxycycline (Sigma, D9891)

at indicated concentrations.

Live Quantitative Phase Imaging

Live quantitative imaging was performed using either the HoloMonitor M4 imaging cytometers (Phase

Holographic Imaging, Lund, Sweden) or the Livecyte platform (Phasefocus, Sheffield, UK). Analyses of

cell proliferation, dry cell mass, and death were acquired with the M4 platform and analyzed using

HStudio (v2.6.3) as previously described[31]. For each experiment, human melanocytes were seeded into

6-well plates (Sarstedt, 83.3920) at 100,000 cells/well and either live-imaged or serially imaged as

indicated. Analysis of growth rate were acquired with the M4 platform and analyzed using App Suite

(v3.2.0.60) as previously described[58]. For each experiment, 100,000 melanocytes, 60,000 501Mel cells or

150,000 HCIMel019 cells were plated per well and media containing indicated concentrations of

barasertib (AURKB inhibitor; AZD1152-HQPA | AZD2811, Selleckchem, A1147) was added. Cells were

imaged for 48-72 hours. Analyses of fluorescent reporters coupled with quantitative phase imaging was

conducted using the Livecyte platform and analyzed using Analyze (v3.1) (Phasefocus, Sheffield, UK).

EdU assays

Normal human melanocytes were nucleofected with microRNA mimics, LNA microRNA inhibitors, or

infected with lentiviral vectors (as above) and seeded in 48-well plates at a density of 50,000 cells/well. 4

days post-seeding EdU was added to culture media at a 10μM final concentration. 24 hours after EdU

addition, media was removed and cells were stained for EdU incorporation and nuclei using the Click-iT

EdU Imaging Kit (Thermo Fisher, C10337) and protocol. Images of EdU and nuclei staining were

acquired using the Evos FL microscope and quantified with FIJI.

Western Blotting

Protein was collected using RIPA Buffer (Thermo Fisher, 89901) supplemented with HALT Protease

Inhibitor (Thermo Fisher, 87786). Immunoblotting was carried out as previously described[23]. Membranes

were incubated overnight at 4 °C with primary antibodies at the following dilutions: anti-HSP90 (CST,

4874) 1:1000, anti-BRAFV600E (Spring Bioscience Corp, E19292) 1:1000, anti-phosphoERK1/2 (CST,

4970) 1:1000, anti-AURKB (Abcam, ab2254) 1:1000. Membranes were washed 4X with TBS and 0.5%

Tween20, incubated with HRP-conjugated secondary antibody (1:2000) for 30 minutes at room

temperature, and visualized with Lumina Forte Western HRP substrate (Millipore, WBLUF0500).

Flow analyses

For assessing DNA content, trypsinized melanocytes (as above) were resuspended in 400 PBS. 1mL -20

ºC 200 proof ethanol was added dropwise while gently vortexing to achieve 70% final concentration for

fixation and incubated overnight at -30 ºC. Cells were centrifuged at 800xg for 5 min, washed with fresh

PBS, and incubated in 500µL FxCycle PI/RNase Staining Solution (Invitrogen, F10797) for 20 min at 37

ºC. Data were collected on the Fortessa (BD) at low speed and analyzed with FlowJo v10.7.1. Annexin

V/PI staining was performed by trypsinizing melanocytes as above. Cells were washed with PBS and

resuspended with Annexin V binding buffer (BioLegend, 422201) at a concentration of 1X10^6 cell/mL.

5µL APC Annexin V antibody (BioLegend, 640919) and 10µL PI were added. Cells were gently vortexed

and incubated in the dark at room temperature for 15 min. Cells were resuspended in 400µL of Annexin

V Binding Buffer. Data were collected on the FACs Verse (BD) and analyzed with FlowJo v10.7.1.

Aldehyde dehydrogenase activity was measured first by trypsinizing melanocytes as above and then

using the ALDEFLUOR kit (Stemcell Technologies, 01700) as described in the manufacture's protocol.

Human nevus and melanoma tissue immunohistochemistry

The archived clinical specimens used in the study were procured with IRB approval from the UCSF

Dermatopathology Pathology Service archives. For assessment of AURKB and GPR3 expression, 11

melanoma and 11 melanocytic nevi with previous diagnosis were de-identified and re-verified

histologically (UEL). For assessment of DNA content, 15 additional melanocytic nevi with previous

diagnosis were re-verified histologically (UEL). Tissue was fixed in 10% neutral-buffered formalin,

processed, embedded in paraffin, and stained with hematoxylin and eosin. 4µm FFPE sections were

stained with AURKB (1:400 dilution, Abcam ab2254) or GPR3 (1:125 dilution, Abnova H00002827-M01)

antibodies. UEL reviewed immunohistochemical stains with semiquantitative grading for GPR3 (0=none;

1=patchy positive; 2=strong positive) and AURKB (0=none; 1=rare positive; 2=scattered positive; 3=

frequent positive; 4=many positive). For in vitro immunofluorescence, cells were fixed and stained as

previously described[80] with AXL (1:250 dilution, Cell Signaling #8661) or MLANA (1:100, Abcam ab731).

Statistical analyses

For differential gene expression, p values were calculated with the DESeq2 (v1.30.1) default Wald test

adjusted by the Benjamini-Hochberg method using a 5% false discovery rate (FDR)[81]. Pathway analyses

were analyzed using the fast gene set enrichment package[82] in R with a 10% FDR. Gene set enrichment

analyses were conducted using GSEA (v4.0.3, Broad Institute) Preranked tool with 1000 permutations.

For *in vitro* experiments, pilot studies were initially conducted in triplicate. The required minimal sample

size to assess the difference between independent means with independent standard deviations with an

alpha error probability of 0.05 and a power of 0.95 was calculated using G*Power (v. 3.1.9.4). P-values

were calculated using either paired or unpaired two-tailed T Tests or Wilcoxon tests via Prism 8

(Graphpad) as indicated in figure legends.

Experimental set-up

For cohorts of clinical specimens, sample number (n) refers to the number of specimens, each from a
different patient. For *in vitro* experiments, n refers to independent experiments conducted on different
days. In cases where fresh primary melanocytes were used, the cells in each experiment are derived from
an independent donor. In cases where cell lines are used, the cells in each experiment represent a
different passage and/or day of that line. For different conditions within experiments, replicate wells were
plated and arbitrarily chosen for each condition (e.g. not treated versus treated; different concentrations of
a compounds, etc.). All data were included.
787
Ethics statement
The archived clinical specimens used in the study were procured with IRB approval from the UCSF
Dermatopathology Pathology Service archives (11-07951). Fresh human tissue samples used in this
study were obtained with informed consent and consent to publish from each patient. Samples were first
coded before being provided to the researchers. The study protocol conformed to the ethical guidelines of
the Declaration of Helsinki (1975). Samples were processed following standard operating procedures
approved by IRB at University of California, San Francisco (16-20299) and at the Huntsman Cancer
Institute (00124195).
796
797 *Data availability*
798 The data which support these findings are available within the article and its supplementary files, as well
799 as from the corresponding author upon reasonable request. The sequencing data produced in this
800 manuscript is available from GEO under the accession number GSE150849.
801
802 **References**
803

1.   Garnett MJ, Marais R. Guilty as charged: B-RAF is a human oncogene. Cancer Cell.
2004;6(4):313–9.
2.   Davies H, Bignell GR, Cox C, Stephens P, Edkins S, Clegg S, Teague J, Woffendin H, Garnett
MJ, Bottomley W, Davis N, Dicks E, Ewing R, Floyd Y, Gray K, Hall S, Hawes R, Hughes J,
Kosmidou V, Menzies A, Mould C, Parker A, Stevens C, Watt S, Hooper S, Jayatilake H,
Gusterson BA, Cooper C, Shipley J, Hargrave D, Pritchard-Jones K, Maitland N, Chenevix-Trench
G, Riggins GJ, Bigner DD, Palmieri G, Cossu A, Flanagan A, Nicholson A, Ho JWC, Leung SY,
Yuen ST, Weber BL, Seigler HF, Darrow TL, Paterson H, Wooster R, Stratton MR, Futreal PA.
Mutations of the BRAF gene in human cancer. Nature. Nature; 2002 Jun 27 [cited 2021 Mar
1];417(6892):949–954.
3.   Pollock PM, Harper UL, Hansen KS, Yudt LM, Stark M, Robbins CM, Moses TY, Hostetter G,
Wagner U, Kakareka J, Salem G, Pohida T, Heenan P, Duray P, Kallioniemi O, Hayward NK,
Trent JM, Meltzer PS. High frequency of BRAF mutations in nevi. Nat Genet. Nat Genet; 2003 Jan

1 [cited 2021 Mar 1];33(1):19–20.

4.   Shain AH, Yeh I, Kovalyshyn I, Sriharan A, Talevich E, Gagnon A, Dummer R, North J, Pincus L,
Ruben B, Rickaby W, D'Arrigo C, Robson A, Bastian BC. The Genetic Evolution of Melanoma
from Precursor Lesions. N Engl J Med. Massachussetts Medical Society; 2015 Nov 12 [cited 2020
Jun 27];373(20):1926–1936.

5.   Bastian BC. The molecular pathology of melanoma: An integrated taxonomy of melanocytic
neoplasia. Annu Rev Pathol Mech Dis. Annual Reviews Inc.; 2014 [cited 2020 Aug 21];9:239–271.

6.   Dankort D, Curley DP, Cartlidge RA, Nelson B, Karnezis AN, Damsky Jr WE, You MJ, DePinho
RA, McMahon M, Bosenberg M. BrafV600E cooperates with Pten loss to induce metastatic
melanoma. Nat Genet. 2009 May 12 [cited 2019 Jul 17];41(5):544–552.

7.   Damsky W, Micevic G, Meeth K, Muthusamy V, Curley DP, Santhanakrishnan M, Erdelyi I, Platt
JT, Huang L, Theodosakis N, Raza Zaidi M, Tighe S, Davies MA, Dankort D, McMahon M, Merlino
G, Bardeesy N, Bosenberg M. mTORC1 activation blocks brafV600E-induced growth arrest but is
insufficient for melanoma formation. Cancer Cell. Cell Press; 2015;27(1):41–56.

8.   Ruiz-Vega R, Chen CF, Razzak E, Vasudeva P, Krasieva TB, Shiu J, Caldwell MG, Yan H,
Lowengrub J, Ganesan AK, Lander AD. Dynamics of nevus development implicate cell
cooperation in the growth arrest of transformed melanocytes. Elife. eLife Sciences Publications
Ltd; 2020 Oct 1 [cited 2021 Mar 1];9:1–24.

9.   Michaloglou C, Vredeveld LCW, Soengas MS, Denoyelle C, Kuilman T, van der Horst CMAM,
Majoor DM, Shay JW, Mooi WJ, Peeper DS. BRAFE600-associated senescence-like cell cycle
arrest of human naevi. Nature. 2005 Aug 4 [cited 2019 Jul 17];436(7051):720–724.

10.   He S, Sharpless NE. Senescence in Health and Disease [Internet]. Cell. Cell Press; 2017 [cited
2021 Mar 2]. p. 1000–1011.

11.   Roy AL, Sierra F, Howcroft K, Singer DS, Sharpless N, Hodes RJ, Wilder EL, Anderson JM. A
Blueprint for Characterizing Senescence. Cell. Cell Press; 2020 Nov 25 [cited 2021 Mar
2];183(5):1143–1146.

12.   Serrano M, Lin AW, McCurrach ME, Beach D, Lowe SW. Oncogenic ras provokes premature cell
senescence associated with accumulation of p53 and p16(INK4a). Cell. Cell Press; 1997 Mar 7
[cited 2021 Mar 2];88(5):593–602.

13.   Campisi J, D'Adda Di Fagagna F. Cellular senescence: When bad things happen to good cells
[Internet]. Nature Reviews Molecular Cell Biology. Nat Rev Mol Cell Biol; 2007 [cited 2021 Mar 2].
p. 729–740.

14.   Collado M, Serrano M. The power and the promise of oncogene-induced senescence markers
[Internet]. Nature Reviews Cancer. Nat Rev Cancer; 2006 [cited 2021 Mar 2]. p. 472–476.

15.   Afshari CA, Vojta PJ, Annab LA, Futreal PA, Willard TB, Barrett JC. Investigation of the role of
G1/S cell cycle mediators in cellular senescence. Exp Cell Res. Exp Cell Res; 1993 [cited 2021
Mar 2];209(2):231–237.

16.   Chatsirisupachai K, Palmer D, Ferreira S, de Magalhães JP. A human tissue-specific
transcriptomic analysis reveals a complex relationship between aging, cancer, and cellular
senescence. Aging Cell. Blackwell Publishing Ltd; 2019 Dec 1 [cited 2021 Mar 2];18(6).

17.   Zhu J, Woods D, McMahon M, Bishop JM. Senescence of human fibroblasts induced by
oncogenic Raf. Genes Dev. Cold Spring Harbor Laboratory Press; 1998 Oct 1 [cited 2021 Mar
1];12(19):2997–3007.

18.   King R, Hayzen BA, Page RN, Googe PB, Zeagler D, Mihm MC. Recurrent nevus phenomenon: A
clinicopathologic study of 357 cases and histologic comparison with melanoma with regression.
Mod Pathol. Mod Pathol; 2009 May [cited 2021 Mar 2];22(5):611–617.

19.   Burian EA, Jemec GBE. Eruptive Melanocytic Nevi: A Review. Am J Clin Dermatol. 2019 May 22
[cited 2019 Jun 21];20(5):669–682.

20.   Cotter MA, Florell SR, Leachman SA, Grossman D. Absence of senescence-associated β-
galactosidase activity in human melanocytic nevi in vivo [3] [Internet]. Journal of Investigative
Dermatology. J Invest Dermatol; 2007 [cited 2021 Mar 2]. p. 2469–2471.

21.   Tran SL, Haferkamp S, Scurr LL, Gowrishankar K, Becker TM, Desilva C, Thompson JF, Scolyer
RA, Kefford RF, Rizos H. Absence of distinguishing senescence traits in human melanocytic nevi.
J Invest Dermatol. Nature Publishing Group; 2012 [cited 2021 Mar 2];132(9):2226–2234.

22.   Oaxaca G, Billings SD, Ko JS. p16 Range of expression in dermal predominant benign epithelioid
and spindled nevi and melanoma. J Cutan Pathol. Blackwell Publishing Ltd; 2020 Sep 6 [cited
2021 Mar 2];47(9):815–823.

23.   Zeng H, Jorapur A, Shain AH, Lang UE, Torres R, Zhang Y, McNeal AS, Botton T, Lin J, Donne
M, Bastian IN, Yu R, North JP, Pincus L, Ruben BS, Joseph NM, Yeh I, Bastian BC, Judson RL.
Bi-allelic Loss of CDKN2A Initiates Melanoma Invasion via BRN2 Activation. Cancer Cell. Cell
Press; 2018 Jul 9 [cited 2021 Mar 2];34(1):56-68.e9.

24.   Haferkamp S, Scurr LL, Becker TM, Frausto M, Kefford RF, Rizos H. Oncogene-induced
senescence does not require the p16 INK4a or p14ARF melanoma tumor suppressors. J Invest
Dermatol. J Invest Dermatol; 2009 Aug [cited 2021 Mar 2];129(8):1983–1991.

25.   McNeal AS, Liu K, Nakhate V, Natale CA, Duperret EK, Capell BC, Dentchev T, Berger SL, Herlyn
M, Seykora JT, Ridky TW. CDKN2B loss promotes progression from benign melanocytic nevus to
melanoma. Cancer Discov. American Association for Cancer Research Inc.; 2015 Oct 1 [cited
2021 Mar 2];5(10):1072–1085.

26.   Shain AH, Joseph NM, Yu R, Benhamida J, Liu S, Prow T, Ruben B, North J, Pincus L, Yeh I,
Judson R, Bastian BC. Genomic and Transcriptomic Analysis Reveals Incremental Disruption of
Key Signaling Pathways during Melanoma Evolution. Cancer Cell. 2018 Jul 9 [cited 2019 Jul
17];34(1):45-55.e4.

27.   Babapoor S, Horwich M, Wu R, Levinson S, Gandhi M, Makkar H, Kristjansson A, Chang M,
Dadras SS. microRNA in situ hybridization for miR-211 detection as an ancillary test in melanoma

diagnosis. Mod Pathol. Nature Publishing Group; 2016 May 1 [cited 2021 Apr 27];29(5):461–475.

28.   Torres R, Lang UE, Hejna M, Shelton SJ, Joseph NM, Shain AH, Yeh I, Wei ML, Oldham MC,

Bastian BC, Judson-Torres RL. MicroRNA Ratios Distinguish Melanomas from Nevi. J Invest

Dermatol. Elsevier B.V.; 2020 Jan 1 [cited 2021 Mar 9];140(1):164-173.e7.

29.   Latchana N, Ganju A, Howard JH, Carson WE. MicroRNA dysregulation in melanoma. Surg

Oncol. Elsevier Ltd; 2016 Sep 1 [cited 2021 Mar 9];25(3):184–189.

30.   Miniotis MF, Mukwaya A, Wingren AG. Digital holographic microscopy for non-invasive monitoring

of cell cycle arrest in L929 cells. PLoS One. Public Library of Science; 2014 Sep 1 [cited 2021 Mar

9];9(9).

31.   Hejna M, Jorapur A, Song JS, Judson RL. High accuracy label-free classification of single-cell

kinetic states from holographic cytometry of human melanoma cells. Sci Rep. Nature Publishing

Group; 2017 Dec 1 [cited 2021 Mar 9];7(1).

32.   Barker KL, Boucher KM, Judson-Torres RL. Label-Free Classification of Apoptosis, Ferroptosis

and Necroptosis Using Digital Holographic Cytometry. Appl Sci. MDPI AG; 2020 Jun 27 [cited

2021 Mar 9];10(13):4439.

33.   Normand G, King RW. Understanding cytokinesis failure. Adv Exp Med Biol. NIH Public Access;

2010 [cited 2021 Mar 9];675:27–55.

34.   Judson RL, Greve TS, Parchem RJ, Blelloch R. MicroRNA-based discovery of barriers to

dedifferentiation of fibroblasts to pluripotent stem cells. Nat Struct Mol Biol. Nat Struct Mol Biol;

2013 Oct [cited 2020 Aug 30];20(10):1227–1237.

35.   D'Avino PP, Capalbo L. New auroras on the roles of the chromosomal passenger complex in

cytokinesis: Implications for cancer therapies. Front Oncol. Frontiers Media S.A.; 2015 [cited 2021

Mar 9];5(OCT).

36.   Clark AS, Karasic TB, DeMichele A, Vaughn DJ, O'Hara M, Perini R, Zhang P, Lal P, Feldman M,

Gallagher M, O'Dwyer PJ. Palbociclib (PD0332991) - A selective and potent cyclin-dependent

kinase inhibitor: A review of pharmacodynamics and clinical development [Internet]. JAMA

Oncology. American Medical Association; 2016 [cited 2021 Mar 9]. p. 253–260.

37.   Helfrich BA, Kim J, Gao D, Chan DC, Zhang Z, Tan AC, Bunn PA. Barasertib (AZD1152), a small

molecule Aurora B inhibitor, inhibits the growth of SCLC cell lines in vitro and in vivo. Mol Cancer

Ther. American Association for Cancer Research Inc.; 2016 Oct 1;15(10):2314–2322.

38.   Hsu MY, Li L, Herlyn M. Cultivation of normal human epidermal melanocytes in the absence of

phorbol esters. Methods Mol Med. Methods Mol Med; 2005 [cited 2021 May 6];107:13–28.

39.   Coppock DL, Buffolino P, Kopman C, Nathanson L. Inhibition of the melanoma cell cycle and

regulation at the G1/S transition by 12-O-tetradecanoylphorbol-13-acetate (TPA) by modulation of

CDK2 activity. Exp Cell Res. 1995 Nov [cited 2021 May 6];221(1):92–102.

40.   Arita Y, Buffolino P, Coppock DL. Regulation of the cell cycle at the G2/M boundary in metastatic

melanoma cells by 12-O-tetradecanoyl phorbol-13-acetate (TPA) by blocking p34(cdc2) kinase

activity. Exp Cell Res. Academic Press Inc.; 1998 Aug 1 [cited 2021 May 6];242(2):381–390.

41.   Wellbrock C, Ogilvie L, Hedley D, Karasarides M, Martin J, Niculescu-Duvaz D, Springer CJ,
Marais R. V599EB-RAF is an Oncogene in Melanocytes. Cancer Res. Cancer Res; 2004 Apr 1
[cited 2021 Mar 9];64(7):2338–2342.

42.   Mukherji S, Ebert MS, Zheng GXY, Tsang JS, Sharp PA, Van Oudenaarden A. MicroRNAs can
generate thresholds in target gene expression. Nat Genet. Nat Genet; 2011 Sep [cited 2021 May
6];43(9):854–859.

43.   Ovando-Vázquez C, Lepe-Soltero D, Abreu-Goodger C. Improving microRNA target prediction
with gene expression profiles. BMC Genomics. BioMed Central Ltd.; 2016 May 17 [cited 2021 May
6];17(1).

44.   Ebert MS, Sharp PA. Roles for MicroRNAs in conferring robustness to biological processes
[Internet]. Cell. Elsevier B.V.; 2012 [cited 2021 May 6]. p. 515–524.

45.   Sharma A, Madhunapantula S V., Gowda R, Berg A, Neves RI, Robertson GP. Identification of
Aurora Kinase B and Wee1-Like Protein Kinase as Downstream Targets of V600EB-RAF in
Melanoma. Am J Pathol. Am J Pathol; 2013 Apr [cited 2021 Apr 27];182(4):1151–1162.

46.   Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The Molecular
Signatures Database Hallmark Gene Set Collection. Cell Syst. Cell Press; 2015 Dec 23 [cited
2021 May 6];1(6):417–425.

47.   Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. KEGG as a reference resource for
gene and protein annotation. Nucleic Acids Res. Oxford University Press; 2016;44(D1):D457–
D462.

48.   Ryu B, Moriarty WF, Stine MJ, DeLuca A, Kim DS, Meeker AK, Grills LD, Switzer RA, Eller MS,
Alani RM. Global Analysis of BRAFV600E Target Genes in Human Melanocytes Identifies Matrix
Metalloproteinase-1 as a Critical Mediator of Melanoma Growth. J Invest Dermatol. NIH Public
Access; 2011 [cited 2021 Sep 19];131(7):1579.

49.   Belote RL, Le D, Maynard A, Lang UE, Sinclair A, Lohman BK, Planells-Palop V, Baskin L, Tward
AD, Darmanis S, Judson-Torres RL. Human melanocyte development and melanoma
dedifferentiation at single-cell resolution. Nat Cell Biol 2021 239. Nature Publishing Group; 2021
Sep 2 [cited 2021 Sep 19];23(9):1035–1047.

50.   Hoek KS, Goding CR. Cancer stem cells versus phenotype-switching in melanoma [Internet].
Pigment Cell and Melanoma Research. Pigment Cell Melanoma Res; 2010 [cited 2021 May 6]. p.
746–759.

51.   Tirosh I, Izar B, Prakadan SM, Wadsworth MH, Treacy D, Trombetta JJ, Rotem A, Rodman C,
Lian C, Murphy G, Fallahi-Sichani M, Dutton-Regester K, Lin JR, Cohen O, Shah P, Lu D,
Genshaft AS, Hughes TK, Ziegler CGK, Kazer SW, Gaillard A, Kolb KE, Villani AC, Johannessen
CM, Andreev AY, Van Allen EM, Bertagnolli M, Sorger PK, Sullivan RJ, Flaherty KT, Frederick DT,
Jané-Valbuena J, Yoon CH, Rozenblatt-Rosen O, Shalek AK, Regev A, Garraway LA. Dissecting

the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. Science (80-).

American Association for the Advancement of Science; 2016 Apr 8 [cited 2021 May

6];352(6282):189–196.

52.   Tsoi J, Robert L, Paraiso K, Galvan C, Sheu KM, Lay J, Wong DJL, Atefi M, Shirazi R, Wang X,

Braas D, Grasso CS, Palaskas N, Ribas A, Graeber TG. Multi-stage Differentiation Defines

Melanoma Subtypes with Differential Vulnerability to Drug-Induced Iron-Dependent Oxidative

Stress. Cancer Cell. Cell Press; 2018 May 14 [cited 2021 May 6];33(5):890-904.e5.

53.   Haq R, Fisher DE, Widlund HR. Molecular pathways: BRAF induces bioenergetic adaptation by

attenuating oxidative phosphorylation. Clin Cancer Res. American Association for Cancer

Research Inc.; 2014 May 1 [cited 2021 May 6];20(9):2257–2263.

54.   S P, T W, PA H, SH K. Stimulation of melanogenesis by tetradecanoylphorbol 13-acetate (TPA) in

mouse melanocytes and neural crest cells. Pigment cell Res. Pigment Cell Res; 2003 Feb [cited

2021 Sep 19];16(1):26–34.

55.   Vidács DL, Veréb Z, Bozó R, Flink LB, Polyánka H, Németh IB, Póliska S, Papp BT, Manczinger

M, Gáspár R, Mirdamadi S, Kemény L, Bata-Csörgő Z. Phenotypic plasticity of melanocytes

derived from human adult skin. Pigment Cell Melanoma Res. John Wiley & Sons, Ltd; 2021 Aug

31 [cited 2021 Sep 19];00:1–14.

56.   Kormos B, Belső N, Bebes A, Szabad G, Bacsa S, Széll M, Kemény L, Bata-Csörgő Z. In Vitro

Dedifferentiation of Melanocytes from Adult Epidermis. PLoS One. Public Library of Science; 2011

[cited 2021 Sep 19];6(2):17197.

57.   Rogers T, Marino ML, Raciti P, Jain M, Busam KJ, Marchetti MA, Marghoob AA. Biologically

distinct subsets of nevi [Internet]. Giornale Italiano di Dermatologia e Venereologia. Edizioni

Minerva Medica; 2016 [cited 2021 Apr 27]. p. 365–384.

58.   Hafner M, Niepel M, Chung M, Sorger PK. Growth rate inhibition metrics correct for confounders in

measuring sensitivity to cancer drugs. Nat Methods. Nature Publishing Group; 2016 Jun 1 [cited

2021 Mar 8];13(6):521–527.

59.   C L, A H, M S, S M. Oncogene activation in melanocytes links reactive oxygen to multinucleated

phenotype and senescence. Oncogene. Oncogene; 2008 Nov 27 [cited 2021 Sep

19];27(56):7070–7082.

60.   H L, S L, G Z, LN K, Y Z, JP M, Y H, W Z, J Z, L W, C K, K S, M X, W X, GC K, LM S, J F, PJ Z, M

H, X X, W G. Oncogenic BRAF-Mediated Melanoma Cell Invasion. Cell Rep. Cell Rep; 2016 May

31 [cited 2021 Sep 19];15(9):2012–2024.

61.   A B, SJ C, E M, JM W, T T, MM T, SE T, RM W, S S, Y F, NR C, SC P, N S, MV H, T S-V, TH H,

Y M, T H, SK T, BS T, E K, RP K, L S, RM W. Developmental chromatin programs determine

oncogenic competence in melanoma. Science. Science; 2021 Sep 3 [cited 2021 Sep

19];373(6559).

62.   Cohen PR, Bedikian AY, Kim KB. Appearance of New Vemurafenib-associated Melanocytic Nevi

on Normal-appearing Skin: Case Series and a Review of Changing or New Pigmented Lesions in
Patients with Metastatic Malignant Melanoma After Initiating Treatment with Vemurafenib. J Clin
Aesthet Dermatol. Matrix Medical Communications; 2013 May [cited 2021 Apr 27];6(5):27–37.

63.   Chen FW, Tseng D, Reddy S, Daud AI, Swetter SM. Involution of eruptive melanocytic nevi on
combination BRAF and MEK inhibitor therapy. JAMA Dermatology. American Medical Association;
2014 Nov 1 [cited 2021 Apr 27];150(11):1209–1212.

64.   Chu EY, Wanat KA, Miller CJ, Amaravadi RK, Fecher LA, Brose MS, McGettigan S, Giles LR,
Schuchter LM, Seykora JT, Rosenbach M. Diverse cutaneous side effects associated with BRAF
inhibitor therapy: A clinicopathologic study. J Am Acad Dermatol. J Am Acad Dermatol; 2012 Dec
[cited 2021 Apr 27];67(6):1265–1272.

65.   Dalle S, Poulalhon N, Debarbieux S, Zaharia D, Mihm MC, Lacouture ME, Rosen A, Marghoob
AA, Busam KJ, Depaepe L, Bringuier PP, Richez P, Baurain JF, Bressac-de Paillerets B, Balme B,
Thomas L. Tracking of second primary melanomas in vemurafenib-treated patients. JAMA
Dermatology. JAMA Dermatol; 2013 Apr [cited 2021 Apr 27];149(4):488–490.

66.   Zimmer L, Hillen U, Livingstone E, Lacouture ME, Busam K, Carvajal RD, Egberts F, Hauschild A,
Kashani-Sabet M, Goldinger SM, Dummer R, Long G V., McArthur G, Scherag A, Sucker A,
Schadendorf D. Atypical melanocytic proliferations and new primary melanomas in patients with
advanced melanoma undergoing selective BRAF Inhibition. J Clin Oncol. J Clin Oncol; 2012 Jul 1
[cited 2021 Apr 27];30(19):2375–2383.

67.   Arita Y, O'Driscoll KR, Weinstein IB. Growth of Human Melanocyte Cultures Supported by 12-0-
Tetradecanoylphorbol-13-acetate Is Mediated through Protein Kinase C Activation. Cancer Res.
1992 [cited 2021 May 6];52(16):4514–4521.

68.   Stavroulaki M, Kardassis D, Chatzaki E, Sakellaris G, Lindschau C, Haller H, Tosca A, Krasagakis
K. Exposure of normal human melanocytes to a tumor promoting phorbol ester reverses growth
suppression by transforming growth factor beta. J Cell Physiol. J Cell Physiol; 2008 Feb [cited
2021 May 6];214(2):363–370.

69.   Chao-Hsing KAO, Hsin-Su YU. A study of the effects of phorbol 12-myristate-13-acetate on cell
differentiation of pure human melanocytes in vitro. Arch Dermatol Res. Springer-Verlag; 1991 Apr
[cited 2021 May 6];283(2):119–124.

70.   Mazar J, de Young K, Khaitan D, Meister E, Almodovar A, Goydos J, Ray A, Perera RJ. The
Regulation of miRNA-211 Expression and Its Role in Melanoma Cell Invasiveness. PLoS One.
PLoS One; 2010 [cited 2021 Apr 27];5(11).

71.   Su M, Miao F, Jiang S, Shi Y, Luo L, He X, Wan J, Xu S, Lei TC. Role of the
p53-TRPM1/miR-211-MMP9 axis in UVB-induced human melanocyte migration and its potential in
repigmentation. Int J Mol Med. Spandidos Publications; 2020 [cited 2021 Apr 27];45(4):1017–
1026.

72.   Herrera R, Hubbell S, Decker S, Petruzzelli L. A role for the MEK/MAPK pathway in PMA-induced

cell cycle arrest: Modulation of megakaryocytic differentiation of K562 cells. Exp Cell Res.

Academic Press Inc.; 1998 Feb 1 [cited 2021 Apr 26];238(2):407–414.

73. Ueda Y, Hirai SI, Osada SI, Suzuki A, Mizuno K, Ohno S. Protein kinase C δ activates the MEK-

ERK pathway in a manner independent of Ras and dependent on Raf. J Biol Chem. J Biol Chem;

1996 [cited 2021 May 6];271(38):23512–23519.

74. Vergara IA, Mintoff CP, Sandhu S, McIntosh L, Young RJ, Wong SQ, Colebatch A, Cameron DL,

Kwon JL, Wolfe R, Peng A, Ellul J, Dou X, Fedele C, Boyle S, Arnau GM, Raleigh J, Hatzimihalis

A, Szeto P, Mooi J, Widmer DS, Cheng PF, Amann V, Dummer R, Hayward N, Wilmott J, Scolyer

RA, Cho RJ, Bowtell D, Thorne H, Alsop K, Cordner S, Woodford N, Leditschke J, O'Brien P,

Dawson SJ, McArthur GA, Mann GJ, Levesque MP, Papenfuss AT, Shackleton M. Evolution of

late-stage metastatic melanoma is dominated by aneuploidy and whole genome doubling. Nat

Commun. Nature Research; 2021 Dec 1 [cited 2021 May 12];12(1).

75. Telentschak S, Soliwoda M, Nohroudi K, Addicks K, Klinz FJ. Cytokinesis failure and successful

multipolar mitoses drive aneuploidy in glioblastoma cells. Oncol Rep. Spandidos Publications;

2015 Apr 1 [cited 2021 Mar 9];33(4):2001–2008.

76. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of short

DNA sequences to the human genome. Genome Biol. Genome Biol; 2009 Mar 4 [cited 2021 Apr

27];10(3).

77. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. Genome Biol. BioMed Central Ltd.; 2014 Dec 5 [cited 2021 Apr 27];15(12).

78. Takacs CN, Andreo U, Dao Thi VL, Wu X, Gleason CE, Itano MS, Spitz-Becker GS, Belote RL,

Hedin BR, Scull MA, Rice CM, Simon SM. Differential Regulation of Lipoprotein and Hepatitis C

Virus Secretion by Rab1b. Cell Rep. Elsevier B.V.; 2017 Oct 10 [cited 2021 Apr 27];21(2):431–

441.

79. Welm BE, Dijkgraaf GJP, Bledau AS, Welm AL, Werb Z. Lentiviral Transduction of Mammary

Stem Cells for Analysis of Gene Function during Development and Cancer. Cell Stem Cell. Cell

Stem Cell; 2008 Jan 10 [cited 2021 Apr 27];2(1):90–102.

80. Zeng H, Jorapur A, Shain AH, Lang UE, Torres R, Zhang Y, McNeal AS, Botton T, Lin J, Donne

M, Bastian IN, Yu R, North JP, Pincus L, Ruben BS, Joseph NM, Yeh I, Bastian BC, Judson RL.

Bi-allelic Loss of CDKN2A Initiates Melanoma Invasion via BRN2 Activation. Cancer Cell. 2018 Jul

9 [cited 2019 Jul 17];34(1):56-68.e9.

81. Benjamini Y, Drai D, Elmer G, Kafkafi N, Golani I. Controlling the false discovery rate in behavior

genetics research. Behav Brain Res. Behav Brain Res; 2001 [cited 2021 May 12]. p. 279–284.

82. Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov M, Sergushichev A. Fast gene set

enrichment analysis. bioRxiv. Cold Spring Harbor Laboratory; 2016 Oct 22 [cited 2021 May

12];060012.

1075

1076

Supplemental Table Legends

1078

1079   **Supplemental Table S1 (related to Figure 1b)**
1080   Results of differential expression analyses performed on previously published databases of matched
1081   melanocytic nevi and melanoma-arising-from-nevi.

1082

1083   **Supplemental Table S2 (related to Figure 1c)**
1084   Normalized read counts from small RNA sequencing performed on melanocytes derived from healthy
1085   human skin and melanocytic nevi.

1086

1087   **Supplemental Table S3 (related to Figure 2)**
1088   Results of differential expression analyses performed on primary melanocytes nucleofected with
1089   microRNA or control mimics.

1090

1091   **Supplemental Table S4 (related to Figure 2c)**
1092   Results from siRNA screen for EdU incorporation performed in primary melanocytes. Mean and standard
1093   deviation of triplicate screens shown for each gene. Column D indicates the microRNA predicted to
1094   targeted the gene.

1095

1096   **Table S5 (related to Figure 4)**
1097   Results of differential expression analyses performed on primary melanocytes with or without BRAF^{V600E}
1098   in +TPA or TPA-free conditions.

1099

1100   **Table S6 (related to Figure 4)**
1101   Results of GSEA analyses performed on primary melanocytes with or without BRAF^{V600E} in +TPA or TPA-
1102   free conditions.

1103

1104   **Fig_3A_Source and Fig_2J_Source**
1105   Zip files containing raw and annotated images of western blots.

1106

1107

A

100µm

100µm · 100µm · 10µm

B

C MLANA / Hoechst · 200µm

8N · 20µm · 1
4N · 20µm · 2
2N · 20µm · 3 · 4
4N · 20µm
50µm

D Low MOI transduction

pCMV | AURKB | pPGK | mCherry

↓

96 hour imaging

68 hr · 74 hr · 80 hr · 82 hr · 83 hr · 84 hr · 90 hr

200 µm

Division

E

P = 0.0141

Percent of cells divided over 96 hr

mCherry negative · mCherry positive

F

☐ Melanocytes (GR50: 149.9nM)
☐ 501MEL (GR50: 25.38nM, P = 0.0119)
☐ MTG019 (GR50: 12.72nM, P = 0.0203)

Growth rate

Barasertib (nM): 0 3 30 300 · 0 3 30 300 · 0 3 30 300

G

■ 0 hr ☐ 48 hr

Relative dry mass

N.S. · P < 0.05 · P < 0.01

DMSO · 30 nM · 300 nM