BRAF[V600E] induces reversible mitotic arrest in human melanocytes via microRNA-mediated

suppression of AURKB.

Andrew S. McNeal[1], Rachel L. Belote[2,*], Hanlin Zeng[2,*], Kendra Barker[2], Rodrigo Torres[1,3], Meghan

Curtin[2], A. Hunter Shain[1,3], Robert H. I. Andtbacka[2,4], Sheri L. Holmen[2,4,5], David H. Lum[2], Timothy H.

McCalmont[6], Matthew W. VanBrocklin[2,4,5], Douglas Grossman[2,5,,7], Maria L. Wei[1,3,8], Ursula E. Lang[1,3,6],

and Robert L. Judson-Torres[2,5,7,†]

[1] Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, San

Francisco, CA, 94158, USA.

[2] Huntsman Cancer Institute, University of Utah, Salt Lake City, UT, 84112, USA.

[3] Department of Dermatology, University of California San Francisco, San Francisco, CA, 94115, USA.

[4] Department of Surgery, University of Utah, Salt Lake City, UT, 84132, USA.

[5] Department of Oncological Sciences, University of Utah, Salt Lake City, UT, 84132, USA.

[6] Department of Pathology, University of California San Francisco, San Francisco, CA, 94143, USA.

[7] Department of Dermatology, University of Utah, Salt Lake City, UT, 84102, USA.

[8] San Francisco Veterans Affairs Medical Center, San Francisco, San Francisco, CA, 94121, USA.

*These authors contributed equally

† To whom correspondence should be addressed. Tel: +1 801-213-8436; Email:

judsontorreslab@gmail.com

21
Abstract

Benign melanocytic nevi frequently emerge when an acquired *BRAF[V600E]* mutation triggers unchecked

proliferation and subsequent arrest in melanocytes. Recent observations have challenged the role of

oncogene-induced senescence in melanocytic nevus formation, necessitating investigations into

alternative mechanisms for the establishment and maintenance of proliferation arrest in nevi. We

compared the transcriptomes of melanocytes from healthy human skin, nevi, and melanomas arisen from

nevi and identified a set of microRNAs as highly expressed nevus-enriched transcripts. Two of these

microRNAs – MIR211-5p and MIR328-3p – induced mitotic failure, genome duplication and proliferation

arrest in human melanocytes through convergent targeting of AURKB. We demonstrate that *BRAF[V600E]*

induces a similar proliferation arrest in primary human melanocytes that is both reversible and conditional.

Specifically, *BRAF[V600E]* expression stimulates either arrest or proliferation depending on the differentiation

state of the melanocyte. We report genome duplication in human melanocytic nevi, reciprocal expression

of AURKB and microRNAs in nevi and melanomas, and rescue of arrested human nevus cells with

AURKB expression. Together, our data describe an alternative molecular mechanism for melanocytic

nevus formation that is congruent with both experimental and clinical observations.

Introduction

Cutaneous melanoma is a potentially deadly skin cancer arising from the pigment-producing melanocytes of the human epidermis. An activating mutation in the BRAF proto-oncogene (BRAF[V600E]) drives over half of all cutaneous melanomas[1–3]. Yet, when a melanocyte acquires a BRAF[V600E] mutation, the cell does not immediately transform to cancer. Instead, it usually undergoes clonal proliferation followed by stable arrest resulting in a benign skin lesion known as a melanocytic nevus or "mole"[3–6]. Despite the continued expression of BRAF[V600E], the majority of melanocytic nevi remain innocuous for the lifespan of the individual, suggesting that nevus cells have robust intrinsic defenses against hyperproliferation. By inducing hyperproliferation and subsequent arrest, the BRAF[V600E] mutation elicits divergent, biphasic phenotypes within a single cell – a poorly understood phenomenon. Characterization of the mechanisms and environmental factors that distinguish the two phenotypes could illuminate candidate strategies for chemoprevention of melanocytic nevus formation or interception of melanoma initiation.

The BRAF[V600E] mutation is clonally present in ~80% of human melanocytic nevi[3] and melanocyte-specific expression of the oncogene is sufficient to induce nevus formation in mice[6–8], suggesting that BRAF[V600E] drives proliferation arrest in melanocytes. The prevailing theory to explain this seemingly paradoxical role for BRAF[V600E] is oncogene-induced senescence (OIS)[9]. Cellular senescence is defined as the permanent cell-cycle arrest of previously replication-competent cells[10,11]. Senescence is associated with a variety of molecular hallmarks including elevated expression of p16[INK4A] and other cyclin-dependent-kinase inhibitors, TP53, H2AX, lysosomal beta-galactosidase and senescence-associated secretory proteins (SASP) as well as a DNA content profile indicative of arrest in the G0/G1 phases of the cell cycle[11–16]. The concept of OIS as a barrier to tumorigenesis derived from observations of a durable proliferation arrest induced by overexpression of oncogenic RAS or RAF in immortalized human fibroblasts[12,17].

More recent studies have questioned whether the term "senescence" aptly describes proliferation arrest of melanocytic nevi. Cell cycle re-entry that accompanies nevus recurrence[18], eruption[19], or transformation to primary melanoma suggests that the nevus arrest phenotype is reversible rather than permanent. Similarly, the expression of senescence markers do not readily distinguish human melanocytic nevi from primary or transformed melanocytes in humans (16[INK4A], TP53, H2AX and beta-galactosidase) or mice (any known hallmark of senescence)[8,20,21]. Setting aside the permanence of OIS, melanocytic nevus arrest was largely thought to be driven by induction of p16[INK4A] expression from the *CDKN2A* gene. Indeed, early studies demonstrated that BRAF[V600E] induces p16[INK4A] expression and that melanocytic nevi frequently express high levels of p16[INK4A] [9]. Despite this, p16[INK4A] expression is neither ubiquitous across melanocytic nevi, nor uniform within the cells of a single nevus[22,23], and neither knock down of the 16[INK4A] transcript nor ablation of the *CDKN2A* locus prevents the onset or rescues BRAF[V600E]-induced proliferation arrest[9,23–25]. BRAF[V600E] also stimulates expression of p15[INK4B] - the translational product of the *CDKN2B* gene[25] which is situated adjacent to the *CDKN2A* locus - suggesting BRAF[V600E]

might orchestrate arrest via multiple cyclin dependent kinase inhibitors that induce arrest in the G1 phase
of the cell cycle, prior to genome synthesis. While the loss of the *CDKN2A/B* locus is a formative event in
melanoma progression, genetic evolution studies of melanoma progression suggest that selection against
the *CDKN2A/B* locus occurs after BRAF[V600E] melanocytes have already escaped nevus-associated
arrest[23,26]. Collectively, these data complicate the straightforward model wherein BRAF[V600E] induces the
expression of cell cycle regulators leading to proliferation arrest, which is later circumvented by the
genetic loss of those regulators.
83
Given these observations, we reasoned that the arrest response of human melanocytes to BRAF[V600E]
might be distinct from previous reports of BRAF[V600E]-induced OIS in other cell types. Specifically, we
hypothesized that nevus-associated arrest is conditional and coordinated by reversible changes in gene
expression. Here, we identify a transcriptional program that is consistently elevated in melanocytic nevi
and includes microRNAs (miRNAs) previously shown to have diagnostic value[27,28]. In parallel, we observe
that, unlike RAF-induced OIS in fibroblasts, BRAF[V600E]–induced arrest in human melanocytes is
reversible and conditional. Merging these two observations, we discover that in melanocytes, regulation
of miRNAs coupled to differentiation status permits toggling between BRAF[V600E]-driven hyperproliferative
to arrested phenotypes.
93
94   **Results**
95
Melanocytic nevus- enriched miRNAs induce mitotic failure and proliferation arrest in primary
human melanocytes
Since melanocytic nevi do not consistently express known senescence markers, we sought to
characterize the transcripts that are specifically elevated within nevus melanocytes (Fig. 1A). We
analyzed previously published datasets comparing RNA expression between benign and malignant
melanocytic lesions[26,28]. Each clinical specimen contained matched RNA samples derived from
melanocytic nevi and melanomas that arose from those nevi. We reasoned that down-regulated
transcripts within these newly-formed melanomas could represent the most immediate barriers to
melanocyte growth. Of the differentially expressed genes (adjusted p value <0.05), six of the top ten with
elevated expression in nevi were non-coding miRNAs (Fig. 1B and Table S1). Among these were
miRNAs we and others have previously shown to be enriched in nevi compared to melanomas, including
MIR211-5p, MIR125A-5p, MIR125B-5p and MIR328-3p[28,29]. To determine whether elevated expression of
these miRNAs distinguished nevus melanocytes from normal epidermal melanocytes, we profiled
melanocytes from fresh healthy skin and nevi (Table S2). Three of the miRNAs (MIR211-5p, MIR328-3p,
MIR125B-5p) were also more highly expressed in melanocytic nevi as compared to healthy melanocytes
or melanomas - an expression pattern consistent with enrichment in arrested melanocytes - whereas one
miRNA (MIR125A-5p) presented a pattern consistent with lineage specific expression that is lost upon

Figure 1

Figure 1: Melanocytic nevus enriched miRNAs induce arrest in primary human melanocytes

(A) Schematic of hypothesized plastic nature of BRAF^{V600E}- and nevus- associated proliferation arrest. Expression of the predicted nevus enriched transcriptional program is highlighted in blue. (B) Scatter plot depicting gene expression combining publicly available sequencing datasets totaling 14 melanoma-arising-from-nevus specimens, dissected into benign and malignant portions (28 total samples). Plotted are the adjusted p values (DESeq2) comparing matched nevus and melanoma samples against average normalized expression in nevus portion. Only genes exhibiting both high expression and significant differential expression in nevus portions are plotted (see Methods). The full list of genes is included as Table S1. (C) Left two columns, the relative expression of indicated miRNAs in freshly isolated human melanocytes from BRAF^{V600E} melanocytic nevi (n = 6) compared to BRAF^{WT} healthy skin melanocytes (n = 8). Right two columns, the relative expression of indicated miRNAs in melanocytes dissected from FFPE nevus (n = 7) or melanoma (n = 7) specimens. Data represent mean and standard deviation of normalized RNA sequencing read counts relative to nevus-derived samples. (D) Mean and standard deviation for population fold change over seven days of melanocytes nucleofected with indicated miRNA mimics compared to non-targeting control (Control RNA) (n = 3). (E) Mean and standard deviation for fraction of EdU positive cells 4 days after nucleofection with indicated concentrations of miRNA mimics (n= 3). P values indicate comparisons to concentration matched control. (F) Representative QPI images of melanocytes nucleofected with indicated miRNA mimics. Pixel color indicates low (black), mid (blue) or high (red) optical density. (G) Mean and standard deviation for relative QPI-derived cell number over time (n = 3). (H) Mean and standard deviation for QPI-derived cell death counts as percentage of total cells imaged on day 1 (n = 3). Cells treated with 0.1 µM staurosporine were included as a control (n = 1) for induction of cell death. (I) Mean and standard deviation for percent Annexin V/PI positive cells nucleofected with indicated miRNA mimics (n = 3). Cells treated with 5µg/ml mitomycin C were included as a control (n = 3) for induction of apoptosis. (J) Mean and standard deviation for relative QPI-derived dry mass per cell (n = 3). Colors as in (E). (K) Representative histograms of DNA content profiling via propidium iodide incorporation measured by flow cytometry 5 days post- nucleofection of miRNA mimics. (L) Mean and standard deviation for percent of cells in indicated phases of cell cycle based upon profiles as in (K) (n = 3-6, individual datapoints shown). P values for D-L calculated by unpaired t tests comparing experimental to Control RNA. N.S. = not significant (P ≥ 0.05)

transformation (Fig. 1C). We conclude that MIR211-5p, MIR328-3p, MIR125B-5p are transcripts specifically enriched in arrested melanocytic nevi.

To investigate whether elevated expression of MIR211-5p, MIR328-3p, or MIR125B-5p induce melanocytic arrest, we introduced synthesized mimetics of the mature miRNAs into primary human melanocytes and tracked proliferation over seven days in culture. The introduction of either MIR211-5p or MIR328-3p resulted in significantly fewer melanocytes compared to a non-targeting control (Fig. 1D). Increased concentrations of these mimics resulted in a dose-dependent decrease in EdU uptake, consistent with a cell cycle defect (Fig. 1E). To better characterize the effect of MIR211-5p or MIR328-3p expression on cell cycle and cell viability, we performed time-lapsed quantitative phase imaging (QPI). We and others have previously established QPI as a method for adherent cell cytometry that can simultaneously measure melanocyte proliferation, cell death, and cell cycle stage with single cell resolution[30–32]. Images were taken every hour for 5 days post-nucleofection (Fig. 1F). Cells expressing MIR211-5p or MIR328-3p exhibited significantly reduced growth curves (Fig. 1G). We observed no evidence of increased cell death (Fig. 1H), and confirmed this result with Annexin V staining (Fig. 1I). Thus, MIR211-5p and MIR328-3p induce proliferation arrest in human melanocytes.

Further characterization of the QPI images revealed a distinct increase in intra-cellular dry mass in proliferation-arrested conditions (Fig. 1J). The dry mass of a cell is a measurement of total biomass (DNA, RNA, proteins, lipids, etc) and is quantitatively measured by QPI. Relative dry mass fluctuates with cell cycle stage in a predictable manner[30]. The 20-40% increase we observed here is characteristic of cells that have doubled their DNA content but failed to complete mitosis. We confirmed this finding by measuring DNA content of the arrested populations (Fig. 1K). We observed significant increases of cells with both 4N DNA content and greater than 4N DNA content, indicative of cytokinetic failure following DNA replication. Consistent with this interpretation, time-lapsed imaging revealed incidences of tripolar mitosis and anaphase bridging in miRNA-expression melanocytes – both processes associated with cytokinetic failure[33] - which were not observed in control conditions or standard culture (Supplemental Files 1 and 2). We conclude that MIR211-5p and MIR328-3p, induced mitotic failure and proliferation arrest when introduced into primary human melanocytes.

Inhibition of AURKB by MIR211-3p and MIR328-5p restricts proliferation in nevi

To investigate the mechanism by which MIR211-5p and MIR328-3p induce arrest in primary human melanocytes, we applied our previously established pipeline for identification of relevant miRNA targets (Fig. 2A)[34]. First, we nucleofected MIR211-5p, MIR328-3p, or control RNA mimics into primary human melanocytes freshly isolated from four independent donors and conducted RNA sequencing (Table S3). We next selected all genes that were significantly downregulated by either miRNA as compared to the control RNA, and further refined the set to the computationally predicted targets of MIR211-5p and/or MIR328-3p. Finally, we nucleofected siRNAs targeting the 115 resulting genes and assayed for reduced EdU incorporation that would phenocopy MIR211-5p or MIR328-3p expression (Table S4). As expected, computationally predicted targets of MIR211-5p and MIR328-3p were enriched in the set of downregulated genes following MIR211-5p or MIR328-3p nucleofection, respectively (Fig. 2B). AURKB, GPR3, and MAPKAPK2 showed the greatest magnitude of repression due to miRNA expression with significance of EdU reduction upon siRNA-mediated knockdown (Fig. 2C). Of these, AURKB (predicted MIR211-5p target) and GPR3 (predicted MIR328-3p target), were significantly inhibited upon nucleofection of either miRNA (Fig. 2D). We reasoned these genes might represent convergent nodes driving proliferation arrest. Further exploration of the AURKB transcript revealed a potential MIR328-3p binding site just before the 3' untranslated region (Fig. 2E). In contrast, we were unable to identify a candidate MIR211-5p binding site in the GPR3 transcript, suggesting indirect inhibition of this gene by MIR211-5p.

Figure 2: Inhibition of AURKB by MIR211-3p and MIR328-5p restricts proliferation in nevi

(A) Schematic of experimental and computational pipeline for identifying targets of MIR211-3p and/or MIR328-3p responsible for proliferation arrest. **(B)** GSEA analysis comparing predicted MIR211-5p and MIR328-3p target mRNAs to changes in gene expression after nucleofection with indicated mimics compared to nucleofection control. NES = normalized enrichment score. P = nominal P value. Table S3 provides full differentially expressed gene lists. **(C)** Plot of siRNA screen results for phenocopying proliferation arrest versus log2 fold change in expression after miRNA mimic nucleofection. Strictly standardized mean difference (SSMD) reports the magnitude and significance of inhibition of Edu incorporation (n = 3, see

Table S4 for list of genes and screen results). Dotted lines indicate cut-off values for further evaluation. Orange points
indicate genes meeting cut-off criteria. **(D)** Mean and standard deviation for expression of indicated genes after
nucleofection of indicated mimics (n = 4). P values calculated by unpaired t tests compared to No RNA. **(E)** Schematics
showing predicted binding sites of MIR211-5p and MIR328-3p in AURKB and GPR3 transcripts. **(F)** Representative images
of immunohistochemical staining for AURKB (brown chromagen) and GPR3 (red chromagen) expression in FFPE samples
of nevi and melanoma. **(G)** Mean and standard deviation for immunohistochemical staining scores as in (F) for cohorts of
11 nevi and 11 melanomas. P values calculated by unpaired t tests. **(H)** Mean and standard deviation for read counts for
cohorts of 11 nevi and 12 melanomas. P values calculated by unpaired t tests. **(I)** Mean and standard deviation for cell
number fold change over 7 days after nucleofection with indicated miRNA mimics in melanocytes transduced with lentiviral
constructs expressing zsGreen, AURKB, or GPR3 (n = 6 melanocyte preps, type 2 t test, N.S. = not significant (P ≥ 0.05)).
P values calculated by unpaired t tests.

To determine whether decreased AURKB and/or GPR3 expression was associated with melanocytic nevi,

we acquired 23 clinical specimens and measured both the transcript and protein expression of these

genes (Fig. 2F-G). The expression of both AURKB and GPR3 protein and mRNA were enriched in

melanomas as compared to nevi (Fig. 2G-H). Together, these data demonstrate an inverse expression

between two miRNAs (MIR211-5p and MIR328-3p) and two targeted mRNAs (AURKB and GPR3) that

toggles concurrently with the transition of arrested melanocytic nevi to melanomas. To further investigate

whether either mRNA was sufficient to rescue miRNA-mediated arrest, primary melanocytes were

transduced with constructs expressing either zsGreen (control), AURKB or GPR3 (Supplemental File 3)

then subsequently nucleofected with either MIR211-5p or MIR328-3p. Expression of either mRNA

partially rescued MIR211-5p-induced arrest, while AURKB, but not GPR3, partially rescued MIR328-3p-

induced arrest (Fig. 2I). Thus, while other MIR211-5p and MIR328-3p targets, including GPR3, may play

a role in the induction of melanocyte proliferation arrest, these data suggest that AURKB inhibition is a

critical convergent node of both miRNAs in facilitating proliferation arrest in nevi. This interpretation is

further supported by the well-established role of AURKB as an essential orchestrator of successful

mitosis[35].

BRAF[V600E] induces a reversible and conditional proliferation arrest in human melanocytes

We hypothesized that AURKB inhibition by MIR211-5p and MIR328-3p may be the underlying

mechanism by which BRAF[V600E] induces proliferation arrest in melanocytes. BRAF[V600E] expression has

been demonstrated to result in an accumulation of growth-arrested fibroblasts with 2N DNA content,

presumably in a permanent G0/G1 senescent state[9,12,17]. In contrast, upon overexpression of MIR211-5p

or MIR328-3p, we observed an accumulation of arrested melanocytes with ≥4N DNA content, and a

significant decrease in cells with 2N DNA content (Fig. 1L). To determine the DNA content profile of

primary human melanocytes under BRAF[V600E]-induced arrest, we utilized an established lentiviral

construct for dose-responsive doxycycline inducible BRAF[V600E] expression (Fig. 3A)[25]. We monitored

melanocyte growth under a titration of doxycycline and identified the lowest concentration (15.6 ng/mL)

that induced sustained arrest after 40 hours exposure (Fig. 3B). Similar to our observations for miRNA-

induced arrest, relative per cell dry mass increased by 50% concurrent with BRAF^V600E induced- arrest

(Fig. 3C). Cell cycle profiling confirmed a significant and dose-dependent accumulation of cells with ≥4N

Figure 3

Figure 3: BRAF^V600E induces a reversible and conditional arrest in human melanocytes

(A) Representative (of n = 3) Western Blot analysis of BRAF^V600E, phospho-ERK1/2, and HSP90 (loading control) in

diBRAF^V600E melanocytes in response to increasing concentration of doxycycline. **(B)** Mean and standard deviation for

relative QPI-derived cell number at indicated hours of diBRAF^V600E melanocytes in response to increasing concentration of

doxycycline (n = 6). P values from unpaired t tests comparing to 0 ng/mL condition. **(C)** Mean and standard deviation for

relative QPI-derived dry mass per cell of diBRAF^V600E melanocytes in response to increasing concentration of doxycycline

at indicated hours (n = 6). P values from unpaired t tests comparing to 0 ng/mL condition. **(D)** Representative histograms of

DNA content profiling via propidium iodide incorporation measured by flow cytometry 5 days after treatment with indicated

molecules. AZD2811 included as a control for induced cytokinesis failure. **(E)** Mean and standard deviation for percent of

cells in indicated phases of cell cycle after treatment with doxycycline based upon profiles as in (D) (n = 4). AZD2811 (n =

4) is a control for induced cytokinesis failure. PD0332991 (n = 2) is a control for G0/G1 arrest. P values from unpaired t

tests comparing to Control (no treatment) condition. **(F)** Mean and standard deviation for relative QPI-derived cell number

of diBRAF^{V600E} melanocytes at indicated hours after treatment doxycycline. Dotted line indicates change in media to either

refresh doxycycline or add media without doxycycline (n = 3). P values from unpaired t tests comparing final timepoints. **(G)**

Mean and standard deviation for cell number as percent of control of diBRAF^{V600E} melanocytes cultured with or without TPA

after 5 days of treatment with indicated concentrations of doxycycline. P values from unpaired t tests comparing to 0 ng/uL

condition. **(H)** Mean and standard deviation for relative QPI-derived cell number of diBRAF^{V600E} melanocytes at indicated

hours after treatment doxycycline. Dotted line indicates change in media to either refresh TPA-containing media or add

media without TPA (n = 3). P values from unpaired t tests comparing final timepoints.

DNA content and a decrease in cells with 2N DNA content (Fig. 3D-E). The cell cycle profiles were

notably dissimilar to that of melanocytes treated with a pharmacologic CDK4/6 inhibitor (PD0332991,

Palbociclib)[36] known to induce G0/G1 arrest. Treatment with a pharmacologic AURKB inhibitor (AZD2811,

Barasertib)[37] that induces G2/M arrest and mitotic failure also resulted in an accumulation of melanocytes

with ≥4N DNA content and a decrease in cells with 2N DNA content.

In addition to the accumulation of 2N cells, previous reports of RAF expression in fibroblasts show that

transient expression can induce a durable arrest phenotype[17]. To test the durability of BRAF^{V600E}-induced

arrest in human melanocytes, we induced doxycylince-dependent BRAF^{V600E} expression for 48 hours and

monitored cell number for an additional two days to confirm arrest. Media was then replaced to remove

doxycycline. Within twelve hours of doxycycline removal, the previously arrested melanocytes began to

proliferate (Fig. 3F), demonstrating that BRAF^{V600E}-induced proliferation arrest in melanocytes is

reversible and dependent on continued oncogene expression.

Nevi recurrence after excision[18] and eruption upon external stimuli[19], together with our observation that

BRAF^{V600E} induced reversible arrest in melanocytes, suggested that BRAF^{V600E}-induced arrest may

dependent on secondary signals from the microenvironment. Tetradecanoylphorbol acetate (TPA) is a

protein kinase C (PKC) activator commonly used as a mitogen in primary melanocyte media to substitute

for endothelin receptor type B activation[38]. In contrast, TPA also inhibits the cell cycle of melanoma

cells[39,40]. BRAF^{V600E} expression permits growth of the Melan-A cell line in the absence of TPA[41]. We

therefore wanted to determine the relative contributions of TPA and BRAF^{V600E} to the BRAF^{V600E}-induced

primary melanocyte arrest *in vitro*. We induced BRAF^{V600E} expression in primary human melanocytes in

two medias – one containing TPA (+TPA) or one containing the endothelin receptor type B ligand,

endothelin-1 (TPA-free) and observed divergent dose-responsive phenotypes (Fig. 3G). In the presence

of TPA, BRAF^{V600E} expression inhibited cell proliferation. By contrast, in TPA-free media, BRAF^{V600E}

expression induced proliferation. We next subjected melanocytes to the arrest inducing conditions of

doxycycline treatment in +TPA media for four days, then replaced with doxycycline – containing, TPA-free

media. We observed a rapid shift to the proliferative phenotype even in the continued presence of

doxycycline (Fig. 3H). From these results, we conclude that BRAF^{V600E} expression induces divergent and

interconvertible phenotypes – either hyperproliferation or a reversible arrest – in human melanocytes

conditional on extrinsic stimuli.

BRAF^{V600E}-induced arrest is dependent on melanocyte growth conditions and differentiation state

To investigate the relative contributions of BRAF^{V600E} and TPA to the expression of MIR211-5p and

MIR328-3p, we conducted small RNA sequencing of melanocytes grown in +TPA or TPA-free media and

with or without BRAF^{V600E} expression. The expression of MIR328-3p was relatively stable, demonstrating

less then 2-fold changes across conditions (Fig. 4A-B, Table S5). In contrast, MIR211-5p was one of the

most differentially expressed genes in +TPA conditions compared to TPA-free conditions (adj. P = 6.8E-

23) (Fig. 4A-B, Table S5). Induction of BRAF^{V600E} in +TPA conditions did not further alter expression of

either miRNA. Direct comparison of the arrested melanocytes in +TPA +Dox conditions to the proliferative

melanocytes in -TPA +Dox conditions revealed MIR211-5p as the most significantly downregulated

miRNA in the proliferative condition (Fig. 4B). Thus, in comparison to TPA, BRAF^{V600E} induction had

minimal effect on the expression of the miRNAs. The most notable effect was a decrease of MIR328-3p

expression coinciding with the induction of BRAF^{V600E} in TPA free conditions (Fig. 4A). While MIR328-3p

downregulation likely contributes to increased proliferation in this condition, the more pronounced effect of

TPA on MIR211-5p expression suggests that the mechanisms governing whether BRAF^{V600E} induces

arrest or proliferation are a function of the cell's transcriptional state rather than directly downstream of

BRAF^{V600E} itself.

To better characterize the two transcriptional states influencing the BRAF^{V600E} phenotype, we conducted

mRNA sequencing of the same specimens. Principal component (PC) analysis revealed that 97.4% of the

variance between samples was explained by two principal components – PC1 (81.7% variance)

separated transcriptomes based upon exposure to TPA and PC2 (15.7% variance) separated

transcriptomes based upon BRAF^{V600E} expression (Fig. 4C, Table S5). We conducted differential

expression analyses to compare all samples exposed to +TPA versus TPA-free media, regardless of

BRAF^{V600E} expression (Fig. 4D, top) and to compare all samples expressing BRAF^{V600E}, regardless of

TPA exposure (Fig. 4D, bottom). As expected from the PC analysis and similar to the miRNA profiles,

most gene expression variance was due to media conditions with a smaller, but consistent, set of genes

activated by BRAF^{V600E}. We conducted gene set enrichment analyses (GSEA) comparing the TPA

regulated genes to known signatures important in human melanocyte or melanoma differentiation[42–44].

Signatures associated with differentiation were significantly enriched (NOM P-value < 0.0001) for the

genes expressed more highly in TPA-free conditions (Fig. 4E). In contrast, signatures associated with

dedifferentiation were significantly enriched (NOM P-value < 0.0001) for the genes expressed more highly

in +TPA conditions, together suggesting that exposure to TPA induces a more dedifferentiated phenotype
in melanocyte culture.

329
Figure 4: BRAF^V600E-induced arrest is dependent on melanocyte growth conditions and
differentiation state
(**A**) Normalized read counts for MIR328-3p (top) and MIR211-5p (bottom) in TPA-containing (+TPA) or TPA-free media,
with or without 15.6 ng/mL doxycycline (Dox). See Table S5 for individual values and exact P values. **(B)** Volcano plots
depicting log2 fold change against adjusted P value (-Log10) from differential expression (DE) analysis of small RNA
sequencing of indicated comparisons. MIR211-5p labelled and shown in red. MIR328-3p shown in blue. **(C)** First and second
principal components (PC) separating diBRAF^V600E melanocytes cultured in indicated conditions. **(D)** Volcano plots depicting

log2 fold change against adjusted P value (-Log10) from differential expression (DE) analysis of mRNA sequencing of all

+TPA specimens versus all TPA-free specimens (top) or all Dox specimens versus all no-Dox specimens (bottom). **(E)**

Gene set enrichment analysis (GSEA) comparing DE gene sets from (D) to published signatures of melanocyte

differentiation from Tsoi, 2018 (NCP, neural crest progenitors, MEL, differentiated melanocytes); Tirosh, 2016 (AXL, AXL-

associated program; MITF, MITF-associated program); Hoek, 2010 (INV, invasive program, PROL, proliferative program).

Normalized enrichment score (NES); All NOM P-values < 0.0001. **(F)** GSEA comparing DE genes sets from (D) to Hallmark

gene sets. Only significant enrichments (NES > 1.5 or NES < -1.5 and NOM P-value < 0.01) shown. **(G)** Normalized read

counts for AURKB normalized to no Dox conditions (n = 3). P values from unpaired t tests. Sequencing data represented in

A-G from n=3 per condition analyzed with DESeq2. **(H)** Schematic of model supported by transcriptomic analyses. TPA

induces a more stem-like state in human melanocytes. Due to increased f MIR211-5p and base-line MIR328-3p expression,

BRAF[V600E] does not induce AURKB, resulting in G2/M accumulation. **(I)** Mean and standard deviation for cell number as

percent of time point 0 hr after 7 days of culture in +TPA media with or without 15.6 ng/mL Dox (n = 3 per condition).

Melanocytes were transduced with either pLVX-control, -AURKB, -anti-MIR211-5p, or -anti-MIR328-3p. P values from

unpaired t tests comparing to matched no Dox condition (horizontal), Control no Dox condition (vertical in green bar) or

Control Dox condition (vertical in red bar).

To identify biological processes associated with differentially expressed genes we conducted GSEAs to

compare TPA-regulated genes and BRAF[V600E]- regulated genes to the Molecular Signature Database

Hallmark gene sets[45]. Despite fewer over-all differentially expressed genes, we observed a greater

number of enriched pathways among the BRAF[V600E]- regulated genes (absolute NES > 1.5, NOM P-value

<0.01) (Fig. 4F). Among the enriched pathways were the expected increases in MAPK (KRAS) signaling,

glycolysis[46], and MTOR signaling[47]. Among the few enriched pathways of TPA- regulated genes were

gene sets associated with increased proliferation (MYC and E2F Targets) and, intriguingly, with the G2-M

Checkpoint. Combined with our observation that BRAF[V600E]-induced proliferation arrest is +TPA

dependent and characterized by cells with ≥4N DNA content (Fig. 3E), this result suggests that

stimulating growth in +TPA conditions results in accumulation of cells in the G2/M phase of the cell cycle.

One initially surprising result of this analysis was the static expression of both MIR211-5p and MIR328-3p

in +TPA conditions, regardless of BRAF[V600E] expression. However, as inhibitors of translation, miRNA

function depends on the expression level of its targets[48,49], effectively functioning as buffers of gene

expression that prevent aberrantly high levels of a transcript[50]. BRAF[V600E] has been previously

demonstrated as an upstream activator of AURKB transcription in melanoma cells[51]. We therefore

reasoned that MIR211-5p expression in +TPA conditions might dampen AURKB activation by BRAF[V600E].

Consistent with this hypothesis, the addition of doxycycline induced AURKB expression in TPA-free /

MIR211-5p low conditions, but this effect was not observed in +TPA / MIR211-5p high conditions (Fig.

4G).

Collectively, our transcriptomic analyses support a model whereby growth in +TPA conditions induces a

more dedifferentiated state in melanocytes coinciding with an increase in MIR211-5p expression (Fig.

4H). This increased MIR211-5p combined with relatively stable MIR328-3p expression, dampens entry

into mitosis and/or successful completion of mitosis at least in part through AURKB inhibition. The
attenuation of cell division results in an accumulation of cells in G2/M – an effect that is more pronounced
in conditions stimulating accelerated growth, such as with BRAF^{V600E} expression. To test this model, we
transduced doxycycline-inducible BRAF^{V600E} melanocytes in +TPA conditions with lentiviral constructs
constitutively expressing either AURKB (Supplemental File 2A) or miRNA sponges (Supplemental File
2D) to either MIR211-5p or MIR328-3p. AURKB expression had no baseline effect on melanocyte growth,
but rescued BRAF^{V600E}-induced arrest (Fig. 4I), suggesting the inability of the oncogene to induce AURKB
in this context serves as a bottleneck in cell division. Also consistent with the model, sponges against
either MIR211-5p or MIR328-3p increased baseline division rate and rescued BRAF^{V600E}-induced arrest,
demonstrating that both miRNAs contribute to inhibition of melanocyte division, with affects that are more
pronounced in the presence of growth signals.
388
AURKB inhibition drives melanocytic nevus-associated proliferation arrest
To further investigate whether this mechanism of arrest occurs in clinical nevus specimens, we examined
an independent cohort of 15 nevi for evidence of mitotic failure. Histopathologic analysis revealed that all
nevi within the cohort exhibited binucleation (Fig. 5A). The percent observed binucleated melanocytes
ranged from 3-12% (mean = 6.3%). It is important to note that since binucleation can only be observed in
the fraction of cells sectioned perpendicular to the plane of the adjacent nuclei, 3-12% is likely an under-
estimate. We also observed giant multinucleated cells in 10 of the 15 nevi, indicative of multiple rounds of
DNA synthesis and cytokinetic failure and consistent with previous reports[52].
397
We sought to determine whether AURKB inhibition was necessary and sufficient for the multi-nucleated
arrest we observed in melanocytic nevi. We obtained three fresh skin biopsies containing nevi,
microdissected the nevus portion, and isolated melanocytes (Fig. 5B). Nevus-derived melanocytes
appeared healthy in culture. Consistent with the histopathologic cohort, co-staining with melanocyte
marker, MLANA, and Hoechst revealed frequent 4N and 8N melanocytes (Fig. 5C). Transduction of
nevus melanocytes with an AURKB and mCherry expressing lentiviral vector at low multiplicity of infection
(MOI) showed that the mCherry positive cells divided significantly more than the their mCherry negative
counterparts (Fig. 5d-e) in cells from all three nevus donors. This result suggests that expression of
AURKB is also rate limiting for the proliferation of human nevus melanocytes, as we observed in primary
melanocytes expressing BRAF^{V600E}.
408
To test the sufficiency of AURKB inhibition to limit BRAF^{V600E} hyperproliferation in melanoma, we
assessed normalized growth rate inhibition of three cultures - primary human melanocytes, an
established BRAF^{V600E} human melanoma cell line, and a short-term BRAF^{V600E} human melanoma culture
- when exposed to a pharmacologic AURKB inhibitor (Barasertib). Normalized growth rate measurements
(such as GR$_{50}$) are independent of baseline cell division rate and are demonstrably more reproducible

than the classically calculated half maximal inhibitor concentration (IC_{50}) when comparing different cell

cultures[53]. An additional advantage of GR_{50} is the ability to distinguish growth rate inhibition (GR_{50}

approaches 0) versus cell death (GR_{50} is negative). Since GR_{50} calculations require proliferation curves,

Figure 5

Figure 5: AURKB inhibition is critical for BRAF^{V600E}-induced arrest of human nevi

(A) Representative image of 15 H&E-stained melanocytic nevus. Border colors indicate two consecutive magnifications.

Arrowheads indicate bi- (red) or multi- (blue) nucleation. **(B)** Example images of skin specimen containing a melanocytic

nevus (circled) and phase contrast microscopy of melanocytes isolated from nevus portion. **(C)** Images of MLANA (purple)

and Hoeschst (green) co-staining of melanocytic nevus derived melanocytes in culture. Zoomed images show

representative 2N, 4N, and 8N cells. **(D)** Representative of images of QPI coupled with fluorescence microscopy to identify

adjacent mCherry -expressing and mCherry-negative melanocytes. Individual cells were identified at time zero, tracked and

monitored for division for 96 hrs. **(E)** Mean and individual matched datapoints for the percent of n = 3 mCherry-negative or

mCherry-positive cells identified at time zero that divided over 96 hrs. P value from paired t test. **(F)** Mean and standard

deviation for 48 hr growth rates of indicated cells treated with Barasertib. P values from unpaired t tests comparing the 30nM

samples of melanoma lines to primary melanocytes (n = 3). **(G)** Mean and standard deviation for relative QPI-derived dry

mass per 501MEL cell at initial time point (black) compared to 48 hr treatment with Barasertib (grey) (n = 3). P values from

unpaired t tests.

we conducted live QPI over 48 hours with a range of Barasertib concentrations (0-300nM). Primary

melanocytes were largely resistant to the compound with reduced growth apparent only at

supraphysiological levels (Fig. 5f). In contrast, both melanoma cultures responded to ten-fold lower

dosage of the compound with GR_{50} approaching zero, indicating primarily arrest, not death. In these

conditions, the per cell dry mass also increased significantly, consistent with a G2/M arrest (Fig. 5g).

Taken together, these data demonstrate that AURKB inhibition limits the rate of BRAF^{V600E}-induced

proliferation in human melanocytic nevi and melanomas.

Discussion

Cellular senescence is conventionally defined as a permanent cell-cycle arrest associated with a variety

of molecular markers and an irreversible arrest in the G0/G1 phase of the cell cycle[10–12]. The vast majority

of melanocytic nevi are stably arrested and harbor BRAF^{V600E}, an oncogenic mutation that induces

cellular senescence when expressed in a variety of mammalian cells[9,12,17]. While these observations

seemingly implicate OIS as the mechanism driving nevus- associated proliferation arrest, a substantial

body of evidence has challenged this paradigm in recent years[8,9,18–25]. Here, we uncover an alternative

mechanism driving nevus formation and stability whereby acquisition of the BRAF^{V600E} mutation does not

necessitate proliferation arrest, but instead permits melanocytes to toggle between hyperproliferation and

arrest. Our transcriptomic analyses suggest that external stimuli can modulate the differentiation state of

BRAF^{V600E} melanocytes, orchestrate the expression level of MIR211-5p, and inform which phenotype is

triggered by the oncogene – before or after the oncogene is introduced. Our data suggest that

hyperproliferation requires AURKB – a kinase necessary for successful cytokinesis that is directly

downstream of BRAF^{V600E} [54]. Whereas sufficient levels of AURKB permit proliferation, AURKB

suppression by MIR211-5p and MIR328-3p results in mitotic failure.

The presented model offers an explanation for several clinical phenomena. First, melanocytic nevi form

when BRAF^{V600E} drives hyperproliferation followed by proliferation arrest. Second, nevus eruption

involves expansion of previously stable nevi over a short period of time. Interestingly, nevus eruption has

been observed in metastatic melanoma patients treated with single agent selective BRAF inhibitors[55–59],

consistent with our observation that proliferation arrest in melanocytes requires sustained expression of

the oncogene. Third, incomplete excision of melanocytic nevi can result in subsequent regrowth of the

benign lesion[18]. Each event requires modulation of the BRAF^{V600E} melanocyte between a proliferative and

arrested phase. Our data demonstrate that BRAF^{V600E} melanocytes can indeed induce proliferative or

arrested states and is consistent with each of these events.

Diminished MIR211-5p/MIR328-3p expression concurrent with increased AURKB expression occurs in a

substantial portion of melanomas. AURKB expression was previously reported as significantly elevated in

melanoma metastases[51] and here we report increased expression in primary melanomas. Similarly,

MIR211-5p expression is consistently observed as lost in melanoma as compared to melanocytic nevi

across independent studies[27–29]. Indeed, MIR211-5p expression levels classify nevi from melanomas with

a high degree of accuracy[27,28]. We have reported on the diagnostic potential of MIR328-3p expression as

well, although multi-study validation was impeded by the absence of hybridization probes in earlier

microarray-based profiling datasets[28]. *In vitro*, we have now demonstrated that inhibition of either miRNA

is sufficient to block BRAF^{V600E}-induced arrest in melanocytes. These data suggest that mitotic failure

serves as a barrier to melanoma progression and is consistent with the genome duplication and copy

number alterations associated with invasive melanoma[4,60]. In melanoma, the specific cause of this

genomic instability is unknown, but reentry of cells into cell cycle after mitotic failure has been shown to

cause genome duplication and accumulation of copy number alterations in other cancers[61]. Further

experiments will be necessary to determine whether dysregulation the spindle checkpoint in nevi

contributes directly to the copy number variations present in melanomas.

It is important to note that while our data support a model of reversible proliferation arrest in nevi, it does

not suggest this reversal is commonplace. Unquestionably, the vast majority of melanocytic nevi present

with sustained arrest. However, we do propose that the duration of arrest is dynamic and modulated by

exposure to external signals. Since our data suggest that loss of MIR211-5p and MIR328-3p expression

and restoration of AURKB expression is concurrent with progression to melanoma, identification of the

source of cell-extrinsic signals might suggest new strategies for chemoprevention or therapy. We have

identified TPA, a known activator of PKC signaling, as one extrinsic factor that induces expression of

MIR211-5p and a more dedifferentiated state. Previous studies have identified TPA as either a mitogen,

inhibitor of the G1/S transition, or inhibitor of the G2/M transition, dependent on the context[39,40,62–64]. We

demonstrate that *in vitro*, TPA activation of MIR211-5p is required for BRAF^{V600E}-induced mitotic failure.

Yet, the inclusion of TPA in primary melanocyte culture is artificial and the environmental stimulus that

regulates MIR211-5p expression in skin remains unknown. One hypothesis is that PKC activation from

keratinocyte interactions serves as the extrinsic stimulus. However, TPA-free media containing an

endothelin receptor type B ligand, endothelin-1, did not elicit MIR211-5p expression or BRAF^{V600E} arrest,

suggestive that the downstream signaling pathways regulated by the two molecules are not identical.

PKC activation can also result in increased MAPK signaling[65,66] raising the possibility that internal MAPK

fluctuations may play a role in toggling between BRAF^{V600E}-induced proliferation versus arrest. MIR211-

5p is also reported to be transcriptionally regulated by both MITF[67] and UV exposure[68]. Additionally,

autocrine signaling might be at play, such that melanocyte density triggers the arrested state as predicted

by recent mathematic modeling of nevus formation[8]. An important future direction will be to identify the

nature and physiological source of this signal for melanocytic nevi in human skin.

In summary, we have identified that BRAF^{V600E} induces a reversible and conditional arrest in human

melanocytes orchestrated by MIR211-5p/MIR328-3p regulation of AURKB. We present a mechanistic

model that allows for melanocytic nevus formation, eruption and recurrence. Moreover, our observations

provide an explanation for the genetic duplication and instability inherent to invasive melanomas and

open new directions for novel chemopreventative and therapeutic strategies.

Competing Interests

R.L.J and M.L.W are inventors on international patent application no. PCT/US2019/023834 concerning

the use of microRNAs investigated in this article as molecular diagnostics for melanoma.

Acknowledgments

This work was supported by the National Institutes of Health (DP5OD019787 to R.L.J., 5F31CA236377 to

A.S.M., P30CA042014 to University of Utah) and the University of California, San Francisco Program for

Breakthrough Biomedical Research (PBBR) Sandler Fellowship to R.L.J. We acknowledge the use of the

HCI Shared Resources for Research Informatics (RI), Cancer Biostatistics (CB), High-Throughput

Genomics and Bioinformatics Analysis (GBA), and the Biorepository and Molecular Pathology (BMP)

Research Histology Section supported by P30CA042014 awarded to HCI from the National Cancer

Institute.

Author Contributions

A.S.M: Investigation, Formal analysis, Writing – original draft, Visualization, Funding acquisition; R.L.B.:

Methodology, Investigation, Resources, Writing – review & editing; H.Z.: Methodology, Investigation,

Validation, Writing – review & editing; K.B.: Investigation; R.T.: Formal analysis; M.C. – Formal analysis;

A.H.S.: Data curation, Writing – review & editing; R.A., Writing – review & editing: Resources; S.H.:

Conceptualization; D.H.L.: Conceptualization, Writing – review & editing; T.H.M.: Resources; M.W.V.:

Conceptualization; D.G.: Resources, Writing – review & editing; M.W.: Resources, Writing – review &

editing; U.E.L.: Investigation, Formal Analysis, Resources, Writing – review & editing; R.L.J.:

Conceptualization, Formal analysis, Investigation, Writing – original draft; Visualization, Supervision,

Funding acquisition.

Materials and Methods

Transcriptomic profiling

For FFPE samples, histopathologic review, microdissection, targeted exon sequencing, phylogenetic

analysis, and RNA and small RNA sequencing of each tumor area were previously described

(phs001550.v2.p1)[23,26,28]. For this study, small RNA reads were aligned to human reference (hg37) with

Bowtie[69] and small RNA reference groups (miRBase21) were counted. For mRNA sequencing, reads

were aligned to human reference (hg37) with Bowtie2 and counted with HTSeq. Differential expression

analysis was performed using DESeq2[70] with p-values adjusted by Benjamini-Hochberg method (p-adj).

Previously published RNA sequencing datasets from *Shain et al. 2018* and *Torres, et al.* 2020 were re-

analyzed here with DESeq2 the combined for visualization. For RNA profiling from primary melanocytes,

total RNA was extracted using TRIzol Reagent (Thermo Fisher, 15596-026) and further purified with the

RNeasy Power Clean Pro Cleanup Kit (Qiagen, 13997-50) to remove melanin. For small RNAseq of

nevus melanocytes sequencing libraries were constructed with the TailorMix Small RNA Library

Preparation Kit (SeqMatic, CA) and sequencing was performed on the Illumina HiSeq2500 platform at

single-end 50bp. For small RNAseq of cultured BRAF^{V600E} transduced melanocytes, sequencing libraries

were constructed with the Qiagen QIAseq miRNA Library Prep kit and sequencing was performed on the

NovaSeq SP platform at paired-end 50bp. For mRNAseq of microRNA mimic nucleofected melanocytes,

150bp paired end sequencing was conducted by GeneWiz. For mRNAseq of cultured BRAF^{V600E}

transduced melanocytes, sequencing libraries were constructed with the Illumina TruSeq Stranded mRNA

Library Prep kit and sequencing was performed on the NovaSeq S4 platform at paired-end 150bp.

Cell derivation and culture

Benign human melanocytic nevi were excised with informed consent from patient donors at the UCSF

Dermatology clinic (San Francisco, CA) or HCI Dermatology clinic according to Institutional Review

Board- approved protocols. Nevus tissue was minced and enzymatically dissociated in 1mg/mL

collagenase (Sigma 11213857001) and 3.3mg/mL dispase (Sigma, 4942078001) in DMEM (Thermo

Fisher, 10569044) for 1 hour at 37ºC. Melanocytes were further isolated by 5- day exposure to 10µg/mL

G418 (InvivoGen, ant-gn-1). BRAF status was confirmed via sanger sequencing (Quintarabio) using the

primer set: BRAF forward: 5'- GCA CGA CAG ACT GCA CAG GG -3'; BRAF reverse: 5'- AGC GGG CCA

GCA GCT CAA TAG -3'. BRAF wildtype (normal) human melanocytes were isolated from de-identified

and IRB consented neonatal foreskins or adult skin. Foreskin tissue was incubated overnight at 4ºC in

dispase and epithelia were mechanically separated from the dermis the following morning. Epithelial

tissue was minced and incubated in 0.25% trypsin (Gibco 25200056) for 4min at 37ºC. Trypsin was

quenched and tissue centrifuged at 500xg for 5min at room temperature. The cell/tissue pellet was

resuspended in Melanocyte medium (ThermoFisher, M254500) with HMGS (ThermoFisher, S0025) and

plated in low volume.

The 501Mel human melanoma line (Gift from Dr. Boris Bastian, CVCL_4633) were cultured in RPMI with

10% Fetal Bovine Serum (FBS, VWR, S107G), 1% Penicillin-Streptomycin (Gibco, 15140122), 1% L-

Glutamine (Gibco, 25030149), and 1% Non-Essential Amino Acids (Gibco, 11140050). HCIMel019 was

derived from patient-derived xenograft (PDX) tumors propagated in mice. An HCIMel019 (P2)

subcutaneous tumor was resected from mouse, minced in digestion buffer (100µM HEPES (Gibco,

15630-080), 5% FBS (DENVILLE, FB5001-H), 20µg/ml gentamicin (Gibco, 15710-064), 1x insulin (Gibco,

51500-056), and 1mg/ml collagenase IV (Gibco, 17104-019) in DMEM (Gibco, 11965-092)), and digested

overnight at 37˚C. Cells were filtered through a 100µm filter (Falcon, 352360) and red blood cells

removed with RBC lysis buffer (0.5M EDTA, 0.5M KHCO3 (Sigma, 237205), and 5M NH4CL (Sigma,

A9434)). Remaining cells were washed with PBS (Gibco, 10010023), and cultured at 37ºC and 5% CO2

in Mel2 media consisting of 80%MCDB153 (Sigma, M7403), 20% L15 (Gibco, 11415-064), 2% FBS

(DENVILLE, FB5001-H), 1.68mM CaCL (Sigma, C4901), 1x insulin (Gibco, 51500-056) 5ng/ml EGF

(Sigma Aldrich, E9644), 15µg/ml Bovine Pituitary Extract (Gibco 13028-014), and 1x Pen/strep (Gibco,

15070-063).

Small RNA Nucleofection

Human melanocytes were trypsinized (Gibco, 25300062), quenched, and centrifuged at 300xg. Cells

were resuspended in R Buffer (ThermoFisher, MPK1025) at 10,000 cells/µL. 10µL of cell slurry was

mixed with miRIDIAN microRNA mimics (Dharmacon: hsa-MIR211-5p C-300566-03-0005, hsa-MIR328-

3p C-300695-03-0005, MIRControl 1 CN-001000-01-05 or MIRControl 2 CN-002000-01-05); or On-

TargetPlus siRNA (Dharmacon, custom library); at 4µM final concentration and nucleofected using the

NEON Transfection System and protocol (Thermo Fisher, MPK5000).

Generation of Lentiviral vectors

pTRIPZ-diBRAFV600E was a gift from Todd Ridky[25]. pLVX-AURKB (Addgene #153316) and pLVX-

GPR3 (Addgene #153317) were generated by subcloning the respective human cDNA (from Addgene

#100142 and #66350) into the MluI and BamHI sites] of the pLVX-Che-hi3 vector (a gift of Sanford

Simon)[71]. pLVX-anti-MIR211-5p (Addgene #153318), pLVX-anti-MIR328-3p (Addgene #153319), and

pLVX-Che-zsGreen (Addgene #153320) were generated by inserting zsGreen with or without a 3'UTR

into the MluI and XbaI sites of the pLVX-Che-hi3 vector. The 3'UTRs contained 7 tandem microRNA

binding sites to report and inhibit microRNA function, as previously described[34]. ZsGreen was subcloned

from pHIV-zsGreen (gift from Bryan Welm & Zena Werb (Addgene #18121))[72].

Lentiviral transduction

2.75×10^6 HEK293T cells were plated on 10cm tissue culture dishes and grown for ~24 hours in DMEM

with 10% FBS. For each 10cm plate, 5µg lentiviral vector, 3.3µg of pMLV-GagPol and 1.7µg of pVSV-G

packaging plasmids were added to 500µL of jetPRIME buffer and 20µL of jetPRIME transfection reagent

(Polyplus, 712-60), and transfected according to manufacturer's instructions. 48 hours post-transfection,

viral supernatant was collected and filtered using 0.45µm syringe filters (Argos 4395-91). Human

melanocytes seeded at $1.0 \times 10^5 – 2.0 \times 10^5$ cells/well density in 6-well plates were incubated in viral

supernatant with 10µg/mL polybrene (Sigma, TR-1003) and centrifuged at 300xg for 60 min at room

temperature. The viral supernatant was removed and replaced with growth media. Transduced cells were

either selected with puromycin (1µg/mL for 5 days, Sigma P8833) or sorted for mCherry expression using

a BD FACSAria II. Cells expressing pTRIPZ-diBRAFV600E were treated with doxycycline (Sigma, D9891)

at indicated concentrations.

Live Quantitative Phase Imaging

Live quantitative imaging was performed using either the HoloMonitor M4 imaging cytometers (Phase

Holographic Imaging, Lund, Sweden) or the Livecyte platform (Phasefocus, Sheffield, UK). Analyses of

cell proliferation, dry cell mass, and death were acquired with the M4 platform and analyzed using

HStudio (v2.6.3) as previously described[31]. For each experiment, human melanocytes were seeded into

6-well plates (Sarstedt, 83.3920) at 100,000 cells/well and either live-imaged or serially imaged as

indicated. Analysis of growth rate were acquired with the M4 platform and analyzed using App Suite

(v3.2.0.60) as previously described[53]. For each experiment, 100,000 melanocytes, 60,000 501Mel cells or

150,000 HCIMel019 cells were plated per well and media containing indicated concentrations of

barasertib (AURKB inhibitor; AZD1152-HQPA | AZD2811, Selleckchem, A1147) was added. Cells were

imaged for 48-72 hours. Analyses of fluorescent reporters coupled with quantitative phase imaging was

conducted using the Livecyte platform and analyzed using Analyze (v3.1) (Phasefocus, Sheffield, UK).

EdU assays

Normal human melanocytes were nucleofected with microRNA mimics, LNA microRNA inhibitors, or

infected with lentiviral vectors (as above) and seeded in 48-well plates at a density of 50,000 cells/well. 4

days post-seeding EdU was added to culture media at a 10µM final concentration. 24 hours after EdU

addition, media was removed and cells were stained for EdU incorporation and nuclei using the Click-iT

EdU Imaging Kit (Thermo Fisher, C10337) and protocol. Images of EdU and nuclei staining were

acquired using the Evos FL microscope and quantified with FIJI.

Western Blotting

Protein was collected using RIPA Buffer (Thermo Fisher, 89901) supplemented with HALT Protease

Inhibitor (Thermo Fisher, 87786). Immunoblotting was carried out as previously described[23]. Membranes

were incubated overnight at 4 °C with primary antibodies at the following dilutions: anti-HSP90 (CST,

4874) 1:1000, anti-BRAFV600E (Spring Bioscience Corp, E19292) 1:1000, anti-phosphoERK1/2 (CST,

4970) 1:1000, anti-AURKB (Abcam, ab2254) 1:1000. Membranes were washed 4X with TBS and 0.5%

Tween20, incubated with HRP-conjugated secondary antibody (1:2000) for 30 minutes at room

temperature, and visualized with Lumina Forte Western HRP substrate (Millipore, WBLUF0500).

DNA content and apoptosis profiling

For assessing DNA content, trypsinized melanocytes (as above) were resuspended in 400 PBS. 1mL -20

ºC 200 proof ethanol was added dropwise while gently vortexing to achieve 70% final concentration for

fixation and incubated overnight at -30 ºC. Cells were centrifuged at 800xg for 5 min, washed with fresh

PBS, and incubated in 500µL FxCycle PI/RNase Staining Solution (Invitrogen, F10797) for 20 min at 37

ºC. Data were collected on the Fortessa (BD) at low speed and analyzed with FlowJo v10.7.1. Annexin

V/PI staining was performed by trypsinizing melanocytes as above. Cells were washed with PBS and

resuspended with Annexin V binding buffer (BioLegend, 422201) at a concentration of 1X10^6 cell/mL.

5µL APC Annexin V antibody (BioLegend, 640919) and 10µL PI were added. Cells were gently vortexed

and incubated in the dark at room temperature for 15 min. Cells were resuspended in 400µL of Annexin

V Binding Buffer. Data were collected on the FACs Verse (BD) and analyzed with FlowJo v10.7.1.

Human nevus and melanoma tissue immunohistochemistry

The archived clinical specimens used in the study were procured with IRB approval from the UCSF

Dermatopathology Pathology Service archives. For assessment of AURKB and GPR3 expression, 11

melanoma and 11 melanocytic nevi with previous diagnosis were de-identified and re-verified

histologically (UEL). For assessment of DNA content, 15 additional melanocytic nevi with previous

diagnosis were re-verified histologically (UEL). Tissue was fixed in 10% neutral-buffered formalin,

processed, embedded in paraffin, and stained with hematoxylin and eosin. 4µm FFPE sections were

stained with AURKB (1:400 dilution, Abcam ab2254) or GPR3 (1:125 dilution, Abnova H00002827-M01)

antibodies. UEL reviewed immunohistochemical stains with semiquantitative grading for GPR3 (0=none;

1=patchy positive; 2=strong positive) and AURKB (0=none; 1=rare positive; 2=scattered positive; 3=

frequent positive; 4=many positive).

Statistical analyses

For differential gene expression, p values were calculated with the DESeq2 (v1.30.1) default Wald test

adjusted by the Benjamini-Hochberg method using a 5% false discovery rate (FDR)[73]. Pathway analyses

were analyzed using the fast gene set enrichment package[74] in R with a 10% FDR. Gene set enrichment

analyses were conducted using GSEA (v4.0.3, Broad Institute) Preranked tool with 1000 permutations.

For *in vitro* experiments, pilot studies were initially conducted in triplicate. The required minimal sample

size to assess the difference between independent means with independent standard deviations with an

alpha error probability of 0.05 and a power of 0.95 was calculated using G*Power (v. 3.1.9.4). P-values

were calculated using either paired or unpaired two-tailed T Tests or Wilcoxon tests via Prism 8

(Graphpad) as indicated in figure legends.

Experimental set-up

For cohorts of clinical specimens, sample number (n) refers to the number of specimens, each from a

different patient. For *in vitro* experiments, n refers to independent experiments conducted on different

days. In cases where fresh primary melanocytes were used, the cells in each experiment are derived from

an independent donor. In cases where cell lines are used, the cells in each experiment represent a

different passage and/or day of that line. For different conditions within experiments, replicate wells were

plated and arbitrarily chosen for each condition (e.g. not treated versus treated; different concentrations of

a compounds, etc.). All data were included.

Ethics statement

The archived clinical specimens used in the study were procured with IRB approval from the UCSF

Dermatopathology Pathology Service archives (11-07951). Fresh human tissue samples used in this

study were obtained with informed consent and consent to publish from each patient. Samples were first

coded before being provided to the researchers. The study protocol conformed to the ethical guidelines of

the Declaration of Helsinki (1975). Samples were processed following standard operating procedures

approved by IRB at University of California, San Francisco (16-20299) and at the Huntsman Cancer

Institute (00124195).

Data availability

The data which support these findings are available within the article and its supplementary files, as well

as from the corresponding author upon reasonable request. The sequencing data produced in this

manuscript is available from GEO under the accession number GSE150849.

References

1.   Garnett MJ, Marais R. Guilty as charged: B-RAF is a human oncogene. Cancer Cell.

2004;6(4):313–9.

2.   Davies H, Bignell GR, Cox C, Stephens P, Edkins S, Clegg S, Teague J, Woffendin H, Garnett

MJ, Bottomley W, Davis N, Dicks E, Ewing R, Floyd Y, Gray K, Hall S, Hawes R, Hughes J,

Kosmidou V, Menzies A, Mould C, Parker A, Stevens C, Watt S, Hooper S, Jayatilake H,

Gusterson BA, Cooper C, Shipley J, Hargrave D, Pritchard-Jones K, Maitland N, Chenevix-Trench

G, Riggins GJ, Bigner DD, Palmieri G, Cossu A, Flanagan A, Nicholson A, Ho JWC, Leung SY,

Yuen ST, Weber BL, Seigler HF, Darrow TL, Paterson H, Wooster R, Stratton MR, Futreal PA.

Mutations of the BRAF gene in human cancer. Nature. Nature; 2002 Jun 27 [cited 2021 Mar

1];417(6892):949–954.

3.   Pollock PM, Harper UL, Hansen KS, Yudt LM, Stark M, Robbins CM, Moses TY, Hostetter G,

Wagner U, Kakareka J, Salem G, Pohida T, Heenan P, Duray P, Kallioniemi O, Hayward NK,

Trent JM, Meltzer PS. High frequency of BRAF mutations in nevi. Nat Genet. Nat Genet; 2003 Jan

1 [cited 2021 Mar 1];33(1):19–20.

4.   Shain AH, Yeh I, Kovalyshyn I, Sriharan A, Talevich E, Gagnon A, Dummer R, North J, Pincus L,

Ruben B, Rickaby W, D'Arrigo C, Robson A, Bastian BC. The Genetic Evolution of Melanoma

from Precursor Lesions. N Engl J Med. Massachussetts Medical Society; 2015 Nov 12 [cited 2020

Jun 27];373(20):1926–1936.

5.   Bastian BC. The molecular pathology of melanoma: An integrated taxonomy of melanocytic

neoplasia. Annu Rev Pathol Mech Dis. Annual Reviews Inc.; 2014 [cited 2020 Aug 21];9:239–271.

6.   Dankort D, Curley DP, Cartlidge RA, Nelson B, Karnezis AN, Damsky Jr WE, You MJ, DePinho

RA, McMahon M, Bosenberg M. BrafV600E cooperates with Pten loss to induce metastatic

melanoma. Nat Genet. 2009 May 12 [cited 2019 Jul 17];41(5):544–552.

7.   Damsky W, Micevic G, Meeth K, Muthusamy V, Curley DP, Santhanakrishnan M, Erdelyi I, Platt

JT, Huang L, Theodosakis N, Raza Zaidi M, Tighe S, Davies MA, Dankort D, McMahon M, Merlino

G, Bardeesy N, Bosenberg M. mTORC1 activation blocks brafV600E-induced growth arrest but is

insufficient for melanoma formation. Cancer Cell. Cell Press; 2015;27(1):41–56.

8.   Ruiz-Vega R, Chen CF, Razzak E, Vasudeva P, Krasieva TB, Shiu J, Caldwell MG, Yan H,

Lowengrub J, Ganesan AK, Lander AD. Dynamics of nevus development implicate cell

cooperation in the growth arrest of transformed melanocytes. Elife. eLife Sciences Publications

Ltd; 2020 Oct 1 [cited 2021 Mar 1];9:1–24.

9.   Michaloglou C, Vredeveld LCW, Soengas MS, Denoyelle C, Kuilman T, van der Horst CMAM,

Majoor DM, Shay JW, Mooi WJ, Peeper DS. BRAFE600-associated senescence-like cell cycle

arrest of human naevi. Nature. 2005 Aug 4 [cited 2019 Jul 17];436(7051):720–724.

10.   He S, Sharpless NE. Senescence in Health and Disease [Internet]. Cell. Cell Press; 2017 [cited

2021 Mar 2]. p. 1000–1011.

11.   Roy AL, Sierra F, Howcroft K, Singer DS, Sharpless N, Hodes RJ, Wilder EL, Anderson JM. A

Blueprint for Characterizing Senescence. Cell. Cell Press; 2020 Nov 25 [cited 2021 Mar

2];183(5):1143–1146.

12.   Serrano M, Lin AW, McCurrach ME, Beach D, Lowe SW. Oncogenic ras provokes premature cell

senescence associated with accumulation of p53 and p16(INK4a). Cell. Cell Press; 1997 Mar 7

[cited 2021 Mar 2];88(5):593–602.

13.   Campisi J, D'Adda Di Fagagna F. Cellular senescence: When bad things happen to good cells

[Internet]. Nature Reviews Molecular Cell Biology. Nat Rev Mol Cell Biol; 2007 [cited 2021 Mar 2].

p. 729–740.

14.   Collado M, Serrano M. The power and the promise of oncogene-induced senescence markers

[Internet]. Nature Reviews Cancer. Nat Rev Cancer; 2006 [cited 2021 Mar 2]. p. 472–476.

15.   Afshari CA, Vojta PJ, Annab LA, Futreal PA, Willard TB, Barrett JC. Investigation of the role of

G1/S cell cycle mediators in cellular senescence. Exp Cell Res. Exp Cell Res; 1993 [cited 2021

Mar 2];209(2):231–237.

16.   Chatsirisupachai K, Palmer D, Ferreira S, de Magalhães JP. A human tissue-specific

transcriptomic analysis reveals a complex relationship between aging, cancer, and cellular

senescence. Aging Cell. Blackwell Publishing Ltd; 2019 Dec 1 [cited 2021 Mar 2];18(6).

17.   Zhu J, Woods D, McMahon M, Bishop JM. Senescence of human fibroblasts induced by

oncogenic Raf. Genes Dev. Cold Spring Harbor Laboratory Press; 1998 Oct 1 [cited 2021 Mar

1];12(19):2997–3007.

18.   King R, Hayzen BA, Page RN, Googe PB, Zeagler D, Mihm MC. Recurrent nevus phenomenon: A
clinicopathologic study of 357 cases and histologic comparison with melanoma with regression.
Mod Pathol. Mod Pathol; 2009 May [cited 2021 Mar 2];22(5):611–617.

19.   Burian EA, Jemec GBE. Eruptive Melanocytic Nevi: A Review. Am J Clin Dermatol. 2019 May 22
[cited 2019 Jun 21];20(5):669–682.

20.   Cotter MA, Florell SR, Leachman SA, Grossman D. Absence of senescence-associated β-
galactosidase activity in human melanocytic nevi in vivo [3] [Internet]. Journal of Investigative
Dermatology. J Invest Dermatol; 2007 [cited 2021 Mar 2]. p. 2469–2471.

21.   Tran SL, Haferkamp S, Scurr LL, Gowrishankar K, Becker TM, Desilva C, Thompson JF, Scolyer
RA, Kefford RF, Rizos H. Absence of distinguishing senescence traits in human melanocytic nevi.
J Invest Dermatol. Nature Publishing Group; 2012 [cited 2021 Mar 2];132(9):2226–2234.

22.   Oaxaca G, Billings SD, Ko JS. p16 Range of expression in dermal predominant benign epithelioid
and spindled nevi and melanoma. J Cutan Pathol. Blackwell Publishing Ltd; 2020 Sep 6 [cited
2021 Mar 2];47(9):815–823.

23.   Zeng H, Jorapur A, Shain AH, Lang UE, Torres R, Zhang Y, McNeal AS, Botton T, Lin J, Donne
M, Bastian IN, Yu R, North JP, Pincus L, Ruben BS, Joseph NM, Yeh I, Bastian BC, Judson RL.
Bi-allelic Loss of CDKN2A Initiates Melanoma Invasion via BRN2 Activation. Cancer Cell. Cell
Press; 2018 Jul 9 [cited 2021 Mar 2];34(1):56-68.e9.

24.   Haferkamp S, Scurr LL, Becker TM, Frausto M, Kefford RF, Rizos H. Oncogene-induced
senescence does not require the p16 INK4a or p14ARF melanoma tumor suppressors. J Invest
Dermatol. J Invest Dermatol; 2009 Aug [cited 2021 Mar 2];129(8):1983–1991.

25.   McNeal AS, Liu K, Nakhate V, Natale CA, Duperret EK, Capell BC, Dentchev T, Berger SL, Herlyn
M, Seykora JT, Ridky TW. CDKN2B loss promotes progression from benign melanocytic nevus to
melanoma. Cancer Discov. American Association for Cancer Research Inc.; 2015 Oct 1 [cited
2021 Mar 2];5(10):1072–1085.

26.   Shain AH, Joseph NM, Yu R, Benhamida J, Liu S, Prow T, Ruben B, North J, Pincus L, Yeh I,
Judson R, Bastian BC. Genomic and Transcriptomic Analysis Reveals Incremental Disruption of
Key Signaling Pathways during Melanoma Evolution. Cancer Cell. 2018 Jul 9 [cited 2019 Jul
17];34(1):45-55.e4.

27.   Babapoor S, Horwich M, Wu R, Levinson S, Gandhi M, Makkar H, Kristjansson A, Chang M,
Dadras SS. microRNA in situ hybridization for miR-211 detection as an ancillary test in melanoma
diagnosis. Mod Pathol. Nature Publishing Group; 2016 May 1 [cited 2021 Apr 27];29(5):461–475.

28.   Torres R, Lang UE, Hejna M, Shelton SJ, Joseph NM, Shain AH, Yeh I, Wei ML, Oldham MC,
Bastian BC, Judson-Torres RL. MicroRNA Ratios Distinguish Melanomas from Nevi. J Invest
Dermatol. Elsevier B.V.; 2020 Jan 1 [cited 2021 Mar 9];140(1):164-173.e7.

29.   Latchana N, Ganju A, Howard JH, Carson WE. MicroRNA dysregulation in melanoma. Surg

Oncol. Elsevier Ltd; 2016 Sep 1 [cited 2021 Mar 9];25(3):184–189.

30.   Miniotis MF, Mukwaya A, Wingren AG. Digital holographic microscopy for non-invasive monitoring

of cell cycle arrest in L929 cells. PLoS One. Public Library of Science; 2014 Sep 1 [cited 2021 Mar

9];9(9).

31.   Hejna M, Jorapur A, Song JS, Judson RL. High accuracy label-free classification of single-cell

kinetic states from holographic cytometry of human melanoma cells. Sci Rep. Nature Publishing

Group; 2017 Dec 1 [cited 2021 Mar 9];7(1).

32.   Barker KL, Boucher KM, Judson-Torres RL. Label-Free Classification of Apoptosis, Ferroptosis

and Necroptosis Using Digital Holographic Cytometry. Appl Sci. MDPI AG; 2020 Jun 27 [cited

2021 Mar 9];10(13):4439.

33.   Normand G, King RW. Understanding cytokinesis failure. Adv Exp Med Biol. NIH Public Access;

2010 [cited 2021 Mar 9];675:27–55.

34.   Judson RL, Greve TS, Parchem RJ, Blelloch R. MicroRNA-based discovery of barriers to

dedifferentiation of fibroblasts to pluripotent stem cells. Nat Struct Mol Biol. Nat Struct Mol Biol;

2013 Oct [cited 2020 Aug 30];20(10):1227–1237.

35.   D'Avino PP, Capalbo L. New auroras on the roles of the chromosomal passenger complex in

cytokinesis: Implications for cancer therapies. Front Oncol. Frontiers Media S.A.; 2015 [cited 2021

Mar 9];5(OCT).

36.   Clark AS, Karasic TB, DeMichele A, Vaughn DJ, O'Hara M, Perini R, Zhang P, Lal P, Feldman M,

Gallagher M, O'Dwyer PJ. Palbociclib (PD0332991) - A selective and potent cyclin-dependent

kinase inhibitor: A review of pharmacodynamics and clinical development [Internet]. JAMA

Oncology. American Medical Association; 2016 [cited 2021 Mar 9]. p. 253–260.

37.   Helfrich BA, Kim J, Gao D, Chan DC, Zhang Z, Tan AC, Bunn PA. Barasertib (AZD1152), a small

molecule Aurora B inhibitor, inhibits the growth of SCLC cell lines in vitro and in vivo. Mol Cancer

Ther. American Association for Cancer Research Inc.; 2016 Oct 1;15(10):2314–2322.

38.   Hsu MY, Li L, Herlyn M. Cultivation of normal human epidermal melanocytes in the absence of

phorbol esters. Methods Mol Med. Methods Mol Med; 2005 [cited 2021 May 6];107:13–28.

39.   Coppock DL, Buffolino P, Kopman C, Nathanson L. Inhibition of the melanoma cell cycle and

regulation at the G1/S transition by 12-O-tetradecanoylphorbol-13-acetate (TPA) by modulation of

CDK2 activity. Exp Cell Res. 1995 Nov [cited 2021 May 6];221(1):92–102.

40.   Arita Y, Buffolino P, Coppock DL. Regulation of the cell cycle at the G2/M boundary in metastatic

melanoma cells by 12-O-tetradecanoyl phorbol-13-acetate (TPA) by blocking p34(cdc2) kinase

activity. Exp Cell Res. Academic Press Inc.; 1998 Aug 1 [cited 2021 May 6];242(2):381–390.

41.   Wellbrock C, Ogilvie L, Hedley D, Karasarides M, Martin J, Niculescu-Duvaz D, Springer CJ,

Marais R. V599EB-RAF is an Oncogene in Melanocytes. Cancer Res. Cancer Res; 2004 Apr 1

[cited 2021 Mar 9];64(7):2338–2342.

42.   Hoek KS, Goding CR. Cancer stem cells versus phenotype-switching in melanoma [Internet].

Pigment Cell and Melanoma Research. Pigment Cell Melanoma Res; 2010 [cited 2021 May 6]. p.
746–759.

43. Tirosh I, Izar B, Prakadan SM, Wadsworth MH, Treacy D, Trombetta JJ, Rotem A, Rodman C,
Lian C, Murphy G, Fallahi-Sichani M, Dutton-Regester K, Lin JR, Cohen O, Shah P, Lu D,
Genshaft AS, Hughes TK, Ziegler CGK, Kazer SW, Gaillard A, Kolb KE, Villani AC, Johannessen
CM, Andreev AY, Van Allen EM, Bertagnolli M, Sorger PK, Sullivan RJ, Flaherty KT, Frederick DT,
Jané-Valbuena J, Yoon CH, Rozenblatt-Rosen O, Shalek AK, Regev A, Garraway LA. Dissecting
the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. Science (80-).
American Association for the Advancement of Science; 2016 Apr 8 [cited 2021 May
6];352(6282):189–196.

44. Tsoi J, Robert L, Paraiso K, Galvan C, Sheu KM, Lay J, Wong DJL, Atefi M, Shirazi R, Wang X,
Braas D, Grasso CS, Palaskas N, Ribas A, Graeber TG. Multi-stage Differentiation Defines
Melanoma Subtypes with Differential Vulnerability to Drug-Induced Iron-Dependent Oxidative
Stress. Cancer Cell. Cell Press; 2018 May 14 [cited 2021 May 6];33(5):890-904.e5.

45. Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The Molecular
Signatures Database Hallmark Gene Set Collection. Cell Syst. Cell Press; 2015 Dec 23 [cited
2021 May 6];1(6):417–425.

46. Haq R, Fisher DE, Widlund HR. Molecular pathways: BRAF induces bioenergetic adaptation by
attenuating oxidative phosphorylation. Clin Cancer Res. American Association for Cancer
Research Inc.; 2014 May 1 [cited 2021 May 6];20(9):2257–2263.

47. Prabowo AS, Iyer AM, Veersema TJ, Anink JJ, Schouten-Van Meeteren AYN, Spliet WGM, Van
Rijen PC, Ferrier CH, Capper D, Thom M, Aronica E. BRAF V600E mutation is associated with
mTOR signaling activation in glioneuronal tumors. Brain Pathol. Brain Pathol; 2014 Jan [cited
2021 May 6];24(1):52–66.

48. Mukherji S, Ebert MS, Zheng GXY, Tsang JS, Sharp PA, Van Oudenaarden A. MicroRNAs can
generate thresholds in target gene expression. Nat Genet. Nat Genet; 2011 Sep [cited 2021 May
6];43(9):854–859.

49. Ovando-Vázquez C, Lepe-Soltero D, Abreu-Goodger C. Improving microRNA target prediction
with gene expression profiles. BMC Genomics. BioMed Central Ltd.; 2016 May 17 [cited 2021 May
6];17(1).

50. Ebert MS, Sharp PA. Roles for MicroRNAs in conferring robustness to biological processes
[Internet]. Cell. Elsevier B.V.; 2012 [cited 2021 May 6]. p. 515–524.

51. Sharma A, Madhunapantula S V., Gowda R, Berg A, Neves RI, Robertson GP. Identification of
Aurora Kinase B and Wee1-Like Protein Kinase as Downstream Targets of V600EB-RAF in
Melanoma. Am J Pathol. Am J Pathol; 2013 Apr [cited 2021 Apr 27];182(4):1151–1162.

52. Rogers T, Marino ML, Raciti P, Jain M, Busam KJ, Marchetti MA, Marghoob AA. Biologically
distinct subsets of nevi [Internet]. Giornale Italiano di Dermatologia e Venereologia. Edizioni

875   Minerva Medica; 2016 [cited 2021 Apr 27]. p. 365–384.

876 53. Hafner M, Niepel M, Chung M, Sorger PK. Growth rate inhibition metrics correct for confounders in
877   measuring sensitivity to cancer drugs. Nat Methods. Nature Publishing Group; 2016 Jun 1 [cited
878   2021 Mar 8];13(6):521–527.

879 54. Bonet C, Giuliano S, Ohanna M, Bille K, Allegra M, Lacour JP, Bahadoran P, Rocchi S, Ballotti R,
880   Bertolotto C. Aurora B is regulated by the mitogen-activated protein kinase/extracellular signal-
881   regulated kinase (MAPK/ERK) signaling pathway and is a valuable potential target in melanoma
882   cells. J Biol Chem. J Biol Chem; 2012 Aug 24 [cited 2021 Mar 10];287(35):29887–29898.

883 55. Cohen PR, Bedikian AY, Kim KB. Appearance of New Vemurafenib-associated Melanocytic Nevi
884   on Normal-appearing Skin: Case Series and a Review of Changing or New Pigmented Lesions in
885   Patients with Metastatic Malignant Melanoma After Initiating Treatment with Vemurafenib. J Clin
886   Aesthet Dermatol. Matrix Medical Communications; 2013 May [cited 2021 Apr 27];6(5):27–37.

887 56. Chen FW, Tseng D, Reddy S, Daud AI, Swetter SM. Involution of eruptive melanocytic nevi on
888   combination BRAF and MEK inhibitor therapy. JAMA Dermatology. American Medical Association;
889   2014 Nov 1 [cited 2021 Apr 27];150(11):1209–1212.

890 57. Chu EY, Wanat KA, Miller CJ, Amaravadi RK, Fecher LA, Brose MS, McGettigan S, Giles LR,
891   Schuchter LM, Seykora JT, Rosenbach M. Diverse cutaneous side effects associated with BRAF
892   inhibitor therapy: A clinicopathologic study. J Am Acad Dermatol. J Am Acad Dermatol; 2012 Dec
893   [cited 2021 Apr 27];67(6):1265–1272.

894 58. Dalle S, Poulalhon N, Debarbieux S, Zaharia D, Mihm MC, Lacouture ME, Rosen A, Marghoob
895   AA, Busam KJ, Depaepe L, Bringuier PP, Richez P, Baurain JF, Bressac-de Paillerets B, Balme B,
896   Thomas L. Tracking of second primary melanomas in vemurafenib-treated patients. JAMA
897   Dermatology. JAMA Dermatol; 2013 Apr [cited 2021 Apr 27];149(4):488–490.

898 59. Zimmer L, Hillen U, Livingstone E, Lacouture ME, Busam K, Carvajal RD, Egberts F, Hauschild A,
899   Kashani-Sabet M, Goldinger SM, Dummer R, Long G V., McArthur G, Scherag A, Sucker A,
900   Schadendorf D. Atypical melanocytic proliferations and new primary melanomas in patients with
901   advanced melanoma undergoing selective BRAF Inhibition. J Clin Oncol. J Clin Oncol; 2012 Jul 1
902   [cited 2021 Apr 27];30(19):2375–2383.

903 60. Vergara IA, Mintoff CP, Sandhu S, McIntosh L, Young RJ, Wong SQ, Colebatch A, Cameron DL,
904   Kwon JL, Wolfe R, Peng A, Ellul J, Dou X, Fedele C, Boyle S, Arnau GM, Raleigh J, Hatzimihalis
905   A, Szeto P, Mooi J, Widmer DS, Cheng PF, Amann V, Dummer R, Hayward N, Wilmott J, Scolyer
906   RA, Cho RJ, Bowtell D, Thorne H, Alsop K, Cordner S, Woodford N, Leditschke J, O'Brien P,
907   Dawson SJ, McArthur GA, Mann GJ, Levesque MP, Papenfuss AT, Shackleton M. Evolution of
908   late-stage metastatic melanoma is dominated by aneuploidy and whole genome doubling. Nat
909   Commun. Nature Research; 2021 Dec 1 [cited 2021 May 12];12(1).

910 61. Telentschak S, Soliwoda M, Nohroudi K, Addicks K, Klinz FJ. Cytokinesis failure and successful
911   multipolar mitoses drive aneuploidy in glioblastoma cells. Oncol Rep. Spandidos Publications;

2015 Apr 1 [cited 2021 Mar 9];33(4):2001–2008.

62. Arita Y, O'Driscoll KR, Weinstein IB. Growth of Human Melanocyte Cultures Supported by 12-0-

Tetradecanoylphorbol-13-acetate Is Mediated through Protein Kinase C Activation. Cancer Res.

1992 [cited 2021 May 6];52(16):4514–4521.

63. Stavroulaki M, Kardassis D, Chatzaki E, Sakellaris G, Lindschau C, Haller H, Tosca A, Krasagakis

K. Exposure of normal human melanocytes to a tumor promoting phorbol ester reverses growth

suppression by transforming growth factor beta. J Cell Physiol. J Cell Physiol; 2008 Feb [cited

2021 May 6];214(2):363–370.

64. Chao-Hsing KAO, Hsin-Su YU. A study of the effects of phorbol 12-myristate-13-acetate on cell

differentiation of pure human melanocytes in vitro. Arch Dermatol Res. Springer-Verlag; 1991 Apr

[cited 2021 May 6];283(2):119–124.

65. Herrera R, Hubbell S, Decker S, Petruzzelli L. A role for the MEK/MAPK pathway in PMA-induced

cell cycle arrest: Modulation of megakaryocytic differentiation of K562 cells. Exp Cell Res.

Academic Press Inc.; 1998 Feb 1 [cited 2021 Apr 26];238(2):407–414.

66. Ueda Y, Hirai SI, Osada SI, Suzuki A, Mizuno K, Ohno S. Protein kinase C δ activates the MEK-

ERK pathway in a manner independent of Ras and dependent on Raf. J Biol Chem. J Biol Chem;

1996 [cited 2021 May 6];271(38):23512–23519.

67. Mazar J, de Young K, Khaitan D, Meister E, Almodovar A, Goydos J, Ray A, Perera RJ. The

Regulation of miRNA-211 Expression and Its Role in Melanoma Cell Invasiveness. PLoS One.

PLoS One; 2010 [cited 2021 Apr 27];5(11).

68. Su M, Miao F, Jiang S, Shi Y, Luo L, He X, Wan J, Xu S, Lei TC. Role of the

p53-TRPM1/miR-211-MMP9 axis in UVB-induced human melanocyte migration and its potential in

repigmentation. Int J Mol Med. Spandidos Publications; 2020 [cited 2021 Apr 27];45(4):1017–

1026.

69. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of short

DNA sequences to the human genome. Genome Biol. Genome Biol; 2009 Mar 4 [cited 2021 Apr

27];10(3).

70. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. Genome Biol. BioMed Central Ltd.; 2014 Dec 5 [cited 2021 Apr 27];15(12).

71. Takacs CN, Andreo U, Dao Thi VL, Wu X, Gleason CE, Itano MS, Spitz-Becker GS, Belote RL,

Hedin BR, Scull MA, Rice CM, Simon SM. Differential Regulation of Lipoprotein and Hepatitis C

Virus Secretion by Rab1b. Cell Rep. Elsevier B.V.; 2017 Oct 10 [cited 2021 Apr 27];21(2):431–

441.

72. Welm BE, Dijkgraaf GJP, Bledau AS, Welm AL, Werb Z. Lentiviral Transduction of Mammary

Stem Cells for Analysis of Gene Function during Development and Cancer. Cell Stem Cell. Cell

Stem Cell; 2008 Jan 10 [cited 2021 Apr 27];2(1):90–102.

73. Benjamini Y, Drai D, Elmer G, Kafkafi N, Golani I. Controlling the false discovery rate in behavior

genetics research. Behav Brain Res. Behav Brain Res; 2001 [cited 2021 May 12]. p. 279–284.

74.   Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov M, Sergushichev A. Fast gene set

enrichment analysis. bioRxiv. Cold Spring Harbor Laboratory; 2016 Oct 22 [cited 2021 May

12];060012.

Supplemental Files and Legends

Supplemental File 1 (related to Figure 1 and Supplemental File 2)

4 day time-lapse quantitative phase imaging of cytokinesis failure in primary melanocytes transduced with

MIR211-5p. Annotations provided in Figure S1.

Supplemental Figure S1

Supplemental File 2 (related to Figure 1 and Supplemental File 1)

Still frames from Supplemental Video 1 documenting representative cytokinesis failure in primary

melanocytes transduced with MIR211-5p. Yellow arrow heads follow a single cell and its progeny through

a tripolar mitosis. Red arrow head follows an anaphase bridge connecting two daughter cells that then

remerge.

Supplemental Figure S2

Supplemental File 3 (related to Figure 2 and Figure 4)

(A) Design of pLVX vectors expressing either AURKB or GPR3. **(B)** Western Blot of AURKB expression in

human melanocytes with or without lentiviral AURKB overexpression. HSP90 is the loading control. **(C)**

Representative photomicrographs (20X) of immunofluorescence for GPR3 (green) or (Hoechst) (purple) in

human melanocytes with GPR3 lentiviral overexpression (GPR3 OE) or without (Control) over-expression.

(D) Design of pLVX vectors expressing microRNA sponges fused to zsGreen.

Supplemental Table S1 (related to Figure 1b)

Results of differential expression analyses performed on previously published databases of matched

melanocytic nevi and melanoma-arising-from-nevi.

Supplemental Table S2 (related to Figure 1c)

Normalized read counts from small RNA sequencing performed on melanocytes derived from healthy

human skin and melanocytic nevi.

Supplemental Table S3 (related to Figure 2)

Results of differential expression analyses performed on primary melanocytes nucleofected with

microRNA or control mimics.

Supplemental Table S4 (related to Figure 2c)

Results from siRNA screen for EdU incorporation performed in primary melanocytes. Mean and standard

deviation of triplicate screens shown for each gene. Column D indicates the microRNA predicted to

targeted the gene.

Table S5 (related to Figure 4)

996 Results of differential expression analyses performed on primary melanocytes with or without BRAF^{V600E}

997 in +TPA or TPA-free conditions.