Translocation of an arctic seashore plant reveals signs of maladaptation to altered climatic

conditions

Hällfors M.H.[*a,b], Lehvävirta S. [a,c], Aandahl T.R.[d], Lehtimäki I.-M.[a], Nilsson L.O. [d,e],

Ruotsalainen A.L.[f], Schulman L.E.[a], Hyvärinen M.-T.[a]

[a] Botany Unit, Finnish Museum of Natural History, P.O. Box 7, FI-00014 University of Helsinki,

Finland.

[b] Research Centre for Environmental Change, Organismal and Evolutionary Biology Research

Programme, Faculty of Biological and Environmental Sciences, P.O. Box 65, FI-00014

University of Helsinki, Finland.

[c] Department of Landscape Architecture, Planning and Management, Swedish University of

Agricultural Sciences, Alnarp, Sweden

[d] Division of Environment and Natural Resources, Norwegian Institute of Bioeconomy Research

(NIBIO), P.O. Box 115, NO-1431 Ås, Norway.

[e] Halmstad University, PO Box 823, SE-30118 Halmstad, Sweden.

[f] Department of Ecology and Genetics, P.O. Box 3000, FI-90014 University of Oulu, Finland.

[*] corresponding author. Contact information: Maria Hällfors, Research Centre for Environmental

Change, Organismal and Evolutionary Biology Research Programme, Faculty of Biological and

Environmental Sciences, P.O. Box 65, FI-00014 University of Helsinki, Finland, e-mail:

maria.hallfors@helsinki.fi; phone: +358 40 721 34 74

Abstract

Ongoing anthropogenic climate change alters the local climatic conditions to which species may

be adapted. Information on species' climatic requirements and their intraspecific variation is

necessary for predicting the effects of climate change on biodiversity. We used a climatic

gradient to test whether populations of two allopatric varieties of an arctic seashore herb

(*Primula nutans* ssp. *finmarchica*) show adaptation to their local climates and how a future

warmer climate may affect them. Our experimental set-up combined i) a reciprocal translocation

within the distribution range of the species with ii) an experiment testing performance of the

sampled populations in warmer climatic conditions south of their range. We monitored survival,

size, and flowering over four growing seasons as measures of performance and, thus, proxies of

fitness. We found that both varieties performed better in experimental gardens towards the north.

Interestingly, highest up in the north, the southern variety outperformed the northern one.

Supported by weather data, this suggests that the climatic optima of both varieties has moved at

least partly outside their current range. Further warming would make the current environments of

both varieties even less suitable. We conclude that *Primula nutans* ssp. *finmarchica* is already

suffering from adaptation-lag due to climate change, and that further warming may be increase

this maladaptation, especially for the northern variety. The study also highlights that it is not

sufficient to run only reciprocal translocation experiments. Climate change is already shifting the

optimum conditions for many species and adaptation needs also to be tested outside the current

range of the focal taxon in order to include both historic conditions and future conditions.

Keywords

botanic garden, conservation, global change, local adaptation, Siberian primrose, threatened

species, transplant experiment

Introduction

Ongoing climate change threatens biodiversity and is predicted, and even shown, to lead to

declines in populations and to extinctions of species (Ceballos et al., 2015; Dawson et al., 2011;

Díaz et al., 2019; Settele et al., 2014; Urban, 2015). The ability of species to respond, *in situ*, to

shifting environmental conditions can be a result of tolerance of a broad range of conditions, or it

can be mediated through evolutionary adaptation (Chevin et al., 2010; Gao et al., 2018; Gienapp

et al., 2008; Merilä & Hendry, 2014) sometimes facilitated by gene flow from congeneric

populations. Understanding through which mechanisms the impact of climate change is

manifested in the responses of species and populations is crucial for anticipating their abilities to

cope with increasing pressures.

Phenotypic plasticity can allow organisms to respond rapidly to changes in their environment

and, hence, it has often been viewed as the main strategy to respond to environmental changes

(Merilä, 2012). In contrast, evolutionary processes are often slow. Some species are showing

signs of acclimatization or adaptation to new conditions, whereas others are either declining or

dispersing to new areas with a favourable environment (Lenoir & Svenning, 2015; Parmesan &

Yohe, 2003; Pöyry et al., 2009). To predict the effects of climate change on biodiversity and to

plan effective conservation strategies, we need both evaluations of how the current distribution

areas of species are changing, and approximations of the genetic and phenotypic potential of

species to endure climatic changes *in situ*.

Species are often used as the basic taxonomic unit in plant conservation, and predictions on the

effects of environmental change usually relate to the species level (Frankham et al., 2012).

Populations of a particular species may, however, be locally adapted, and therefore have

divergent affinities to abiotic conditions (Banta et al., 2012) and show intraspecific differences in

climatic tolerance (Atkins & Travis, 2010; Hill et al., 2011). However, in their review on local

adaptation in plant species, Leimu and Fischer (2008) found that local adaptation of populations

to environmental conditions is not as widespread as often assumed. Local adaptation may be

favoured mostly in cases where populations occur in differing environments and gene flow

between them is restricted (Kawecki & Ebert, 2004). Understanding and taking into account

intraspecific variation in the adaptation to climatic conditions is necessary to reach balanced

estimates of climate change impact at the level of populations (Hällfors et al., 2016; Souther and

McGraw 2011; Valladares et al., 2014). The speed of climate change may outpace the ability of

populations to respond adaptively, thereby causing adaptation lag (McGraw et al. 2015) and,

hence, maladaptation instead of adaptation or acclimatization. Thus, the need to estimate the

effect that climate change has had on natural populations to date, has become increasingly

evident (Anderson & Wadgymar, 2020; Franks et al., 2007; Kooyers et al., 2019; McGraw et al.

2015; Thomann et al., 2015; Wilczek et al., 2014; Zenni et al., 2014).

Manipulative experiments have the potential to provide information on how strong a role climate

plays in defining favourable conditions of a species or population (Hargreaves et al., 2014;

Kawecki & Ebert, 2004; Kreyling et al., 2014; Pelini et al., 2012). Results from experiments

where the climatic conditions for the focal taxa are altered can be incorporated into species

distribution models to improve predictions (Greiser et al., 2020). By reciprocally transplanting

individuals of two or more populations to conditions representing a foreign environment and a

home environment, we can effectively measure intraspecific variation that allows maintenance of

fitness under varying temperature conditions (Blanquart et al., 2013; Kawecki & Ebert, 2004).

Experiments can also test species' responses to future conditions, for example, through

treatments mimicking increased temperatures, and inform us about the extent to which plasticity

contributes towards population resilience under climate change. Furthermore, such experiments

can reveal locally adaptive traits, which would signify that the survival of the species in the

future would require either rapid microevolution *in situ*, a range shift or, in the absence of these,

assisted migration (*sensu* Hällfors et al., 2014) to new favourable areas.

Here, we describe a study on Siberian primrose (*Primula nutans* ssp. *finmarchica* (Jacq.) Á.

Löve & D.), in which we sampled populations of two of its varieties (the northern var.

finmarchica and the southern var. *jokelae* L. Mäkinen & Y. Mäkinen); while acknowledging that

our sampling did not cover the varieties' full distribution, for clarity we refer to the populations

of the different varieties by 'variety'. We performed 1) a reciprocal transplant experiment and 2)

an experiment of out-of-range transplantation. The former tests the importance of intraspecific

local adaptation versus insensitivity towards climatic conditions (Fig. 1 (b) versus (c)), (Fig. 1 (c)

versus (d)) and the latter the varieties' capability to survive in future climate conditions. We

conducted the experiment across four growing seasons using a network of botanic gardens in

Finland, Norway, and Estonia (Fig. 1). Two of the experimental gardens represent the home

environments of each variety, forming the reciprocal component in the experiment (Oulu and

Svanvik; Fig. 1). Gardens outside of the range of the species (Tartu, Helsinki, Rauma; Fig. 1)

were included to simulate the potential future climates. We tested the effect of the variety and the

environmental conditions (garden or annual temperature) and their interaction on the

performance of plant individuals.

Specifically, we hypothesised that

i)   The varieties are adapted to local climatic conditions and thus perform better in their

123   home environments (Fig. 1 (b)).

124   ii)   Both varieties perform less well in the out-of-range environments, but the southern

125   variety would have higher performance than the northern one in locations with higher

126   temperatures outside the species' natural range (Fig. 1 (b)).

Equally high performance of both varieties in the test environments (Fig. 1 (c)) would reveal

a lack of local adaptation, while no affinity towards conditions within the species range

should result in an equally high performance of the species within *and* outside of its range

(Fig. 1 (d)). However, because climate change has already proceeded (IPCC, 2019), the

process of adaptation may be lagging behind and populations may in fact be maladapted to

current climatic condition (Anderson & Wadgymar, 2020; Kooyers et al., 2019; Wilczek et

al., 2014). This scenario of climate change induced maladaptation (Fig. 1 (e)) needs to be

taken into account in contemporary translocation studies. Our study set-up thus allowed us to

disentangle climatic affinity and infer the occurred and predicted effects of global warming

on these two populations of the Siberian primrose.

Materials and Methods

 <u>Study species and seed sampling</u>

Siberian primrose (*Primula nutans*) is a small-statured perennial herb with a discontinuous,

circumpolar distribution. The Fennoscandian subspecies *P. nutans* ssp. *finmarchica* (Jacq.) Á.

Löve & D. Löve is a red-listed species (VU in Norway and NT in Finland with a disjunct

distribution (Fig. 1). It comprises two morpho-ecological varieties (Mäkinen & Mäkinen, 1964):

var. *finmarchica* (the northern variety) grows on the shores of the Arctic Sea, while var. *jokelae*

(the southern variety) occurs by the Bothnian Bay and the White Sea (Fig. 1 (a)). It is a habitat

specialist that mainly grows in seashore and riverside meadows (Kreivi, Aspi, & Leskinen, 2011;

Mäkinen & Mäkinen, 1964). It propagates sexually by seeds and vegetatively by stolons growing

from the wintering buds and the rosettes (Mäkinen & Mäkinen, 1964). On the basis of matrix

models published by Björnström et al. (2011) individuals especially sterile rosettes can be

relatively persistent and hence live for several years. The species is insect-pollinated and

produces copious seeds that spread by gravitation, via water flows, and possibly also with birds

(Ulvinen 1997).

Due to the geographic distance (c. 400-550 km) between its two varieties var. *jokelae* and var.

finmarchica, and the lack of obvious adaptations for efficient gene flow between the areas, the

varieties are presumably genetically isolated, which is also reflected in their genetic differences

(Kreivi et al., 2011). The main areas where the varieties occur are climatically different (Hällfors

et al., 2016; also see Fig. 2 (a)), and there is a clear difference in the requirement of colder night

conditions for flower induction in the northern variety compared to southern variety (Mäkinen &

Mäkinen, 1964).

We sampled seeds from wild populations of both varieties of *Primula nutans* ssp. *finmarchica* in

August to September 2012 (Fig. 1 (a), Table S1). Seeds of the southern variety were collected in

five sites in Haukipudas and Ii in Finland (seed sampling approval number:

POPELY/346/07.01/2012.), and seeds of the northern variety from six sites in Sør-Varanger in

Norway (no seed sampling permit required as the species is not protected in Norway). Voucher

specimens from each sampling site are deposited at the herbarium of the Finnish Museum of

Natural History (H *sensu* Thiers 2016). From each site we collected seeds from 10 - 53

individuals (as available, following seed collecting guidelines; ENSCONET, 2009; Table S1) to

obtain a representative sample. The number of seeds obtained from the collected capsules of

each sampled individual varied from 0 to *c.* 200. We stored the seed lot from each maternal

individual separately and left them to dry and ripen at room temperature for a minimum of two

weeks after which we placed them in a freezer (-18 to -20°C) for cold stratification to break

dormancy. The seeds were kept in these conditions for about six months, until used to produce

plant material for the translocation experiment (in spring 2013).

Experimental design and plant material

We set up a common garden experiment in 2013 in five botanic gardens located in Estonia and

Finland, and on research station grounds in Norway (Fig. 1 (a); Table S2). We chose to use

botanic gardens as testing grounds instead of natural sites to avoid genetic contamination of

natural populations by introducing alien genotypes, as well as for logistical reasons and legal

restrictions of introducing species outside their natural range.

We chose the seeds for producing experimental plants for the trial through a hierarchical

randomized method (Table S1; Fig S1). For each experimental garden, we selected material from

three seed sampling sites per variety. From each site, we randomly chose five maternal

individuals. We anticipated that not all seeds would germinate and therefore we sowed eight

seeds per maternal individual (120 seeds per variety for each site) although we needed only three

seedlings per maternal individual (45 seedlings per variety for each site).

We used the F1 offspring for our tests for two main reasons. Producing a 'refresher' generation

between collecting the seeds and growing the experimental plants would have been difficult or

impossible in uniform conditions as the two varieties flower in different temperatures conditions

(Mäkinen & Mäkinen 1964). Second, even in the best case, this would have delayed the

experiment by at least one, possibly 2–3 years, due to the life history of the focal species.

Nevertheless, a recent meta-analysis (Yien et al. 2019) concluded that perennial plants show

hardly any transgenerational responses (i.e., effects on the offspring of the ancestor

environmental conditions), so it is not likely that the use of F1 offspring affected our results

significantly.

We sowed each seed in a 6×6 cm pot with commercial sowing soil (Kekkilä) mixed with sand,

vermiculite and perlite (5:2:2:0.5 litres, respectively), and with a thin layer of sand on top. The

seeds were sown in six cohorts, five weeks before intended planting at each experimental garden

(Table S2). They were left to germinate in a greenhouse in Kumpula Botanic Garden in Helsinki,

first in a small greenhouse inside the actual greenhouse to enable suitable temperatures during

early spring. Temperature conditions varied from 18–24□C and lighting was set to 12 h of extra

light (400 W) during daytime.

Due to poor germination and growth, possibly caused by higher than optimal temperature or

insufficient amount of nutrients or light, the seedlings were moved out of the small greenhouse

into the actual one to allow cooler temperatures and more light (10-18□C; Table S2). The

moving followed the same order and time intervals as the sowing so that each cohort spent the

same amount of time in the small greenhouse. Five days after the move, the pots were given a

one-time nutrient addition with a general commercial fertilizer (Kekkilä Taimi-Superex). This is

a common procedure when using nutrient poor germination soil. Without fertilization of the soil

after seed germination the growth usually comes to a halt.

For each experimental garden we selected 45 seedlings of each variety, so that each variety was

represented by three seed sampling sites, each of which contributed five seedlings from three

different maternal individuals (Fig S1). If there were not enough seedlings from a certain

maternal individual, we complemented the design with seedlings from *another maternal*

individual of the same seed sampling population (in 69 cases). If there was no other such

maternal individual, we used seedlings from *another maternal individual from another seed*

sampling population but of the same variety (in 21 cases).

Because of a delayed spring in 2013, establishing the experimental plots had to be postponed by

three weeks to avoid frozen soil and frost that could kill the seedlings after planting. The

approximately two-month-old seedlings were transported and planted during May–June 2013

(see Table S2 for exact dates), starting from the southernmost garden in mid-May (Tartu,

Estonia) and reaching north (Svanvik, Norway; Table S2) in late June. The intention was to

follow the advance of the season so as to plant as early as possible while avoiding frost damage.

In each experimental garden, we planted the seedlings in three $165 \times 145 \times 20$ cm experimental

plots with a minimum distance of 30 m between the plots, following a randomized block design.

A filter cloth was placed at the bottom to prevent the ground soil from affecting the growing

conditions and to help retain moisture. The plots were filled with fine sand and peat mixture (75

and 25 volume-%, respectively) with some added dolomite lime. The intention was to roughly

mimic the seashore meadow soils where the species grows in the wild while using commercially

available soils that could be obtained in large quantities by a commercial soil provider and

transferred to the all experimental gardens. The aim with using the same substrate was to

homogenise (to certain extent) the growth conditions and thus to be able to separate the thermal

effect from other effects.

In each plot, we planted 30 seedlings (i.e., 90 seedlings per garden) in a grid, c. 21 cm from each

other and 30 cm from the wooden frame. In each plot, 15 seedlings were of the southern and 15

of the northern variety, placed in a random order. We placed light metal cages on top of each plot

to prevent large animals from interfering with the experiment. Plots were irrigated with c. 20

litres of water once a week during the growing season to keep the soil sufficiently moist.

Although precipitation is part of climate, the effects of which we attempt to measure, we

considered basic watering of the trials necessary as this is a sea-shore plant naturally exposed to

high water tables. This watering regime was not intended to directly resemble the seashore

meadows in nature, where the water level can vary substantially and occasionally even flood the

plants. Instead, we applied irrigation to avoid large effects of drought, which this seashore

species growing on occasionally inundated meadows, would not endure. Thus, watering was

done only to the amount to compensate for the fact that our test sites were not at the shoreline,

i.e. to mimic the natural sites of the primrose up to a certain baseline so that the plants would not

die because of an unnatural unfavourable moisture regime. Plots were also weeded when deemed

necessary to avoid effects of competition from other species, as our focus was not on the effect

of competition on plant performance.

Since almost 50% of the plants died during the first summer (139 individuals of the northern

variety and 57 of the Southern; likely due to small seedling size and too little irrigation), in the

autumn of 2013, the experimental plots were supplemented with left-over plants that had been

growing outdoors in Kumpula botanic garden in Helsinki during the summer (dates presented in

Table S2). We chose new plants repeating the plant selection process described above for

situation where there were not enough representatives of the preselected maternal individual for

the experimental garden. We recorded the planting time, to enable differentiating between

original and new plants in subsequent analyses. Altogether, 614 individuals were planted in the

five gardens during the early and late summer visits.

To describe the performance of individuals, we measured their survival, size, and flowering

(whether they flowered, how many flowers they produced, and when) from year 2014 to 2016.

The fitness of genotypes can be described as the relative success with which they transmit their

genes to the next generation (Silvertown & Charlesworth, 2001). Because of the limited temporal

extent of our study and because we do not measure the individual's fitness through its ability to

produce viable progeny, we focus on key plant performance measures likely to correlate with

fitness. Survival is a definite measure of fitness as a dead individual cannot transmit genes, but

survival can also be stochastic. Also, the physical size of an individual and reproductive output,

such as abundance of flowering, tend to correlate with plant fitness (Silvertown & Charlesworth,

2001). Each spring (2014, 2015, and 2016) we recorded flowering presence and abundance.

Flowering was inventoried approximately every second day for 14 days after the first flower

appeared in that garden in the specific year. In the autumn of the same years, we recorded

survival and photographed each surviving plant from above. We assessed plant size from

photographs, through digitally cutting out and measuring the area (cm^2) of the visible vegetative

parts. We used the size at the time of planting as a measurement of original size, but 24

individuals lack data on that due to missing photographs or insufficient resolution. These

individuals were therefore not included in the analyses, and our total N is reduced from 614 to

590.

Weather and climatic data

For describing the climatic conditions at the seed sampling sites and experimental gardens, we

obtained data on historic climatic conditions (1970-2000; 10 minutes resolution) and future

projections (CMIP5 for 2050, 10 minutes resolution, HADGEM2-ES model) through the

Worldclim database (Fick & Hijmans, 2017). These climatic data were downloaded in R using

the *getData* function in the *raster* package (Hijmans, 2019). Additionally, we modelled plant

performance as a function of mean annual temperature in the experimental gardens during 2013-

2016 (see below; Fig 2). The weather data were obtained from the Gridded Agro-Meteorological

Data in Europe (Joint Research Centre, 2014), which contains meteorological parameters from

European weather stations interpolated on a 25x25km grid. We used the function *biovar* in the

package *dismo* (Hijmans et al., 2017) to calculate bioclimatic variables of the weather data in R.

We present historic and predicted mean annual temperature for the seed sampling sites and

experimental gardens and as a mean across experimental years for each garden in Fig. 2. Mean

annual temperature is also shown explicitly for each year in each site in Fig. S4 and S5 together

with several additional bioclimatic variables, including annual precipitation sum and mean

temperature during the warmest quarter of the year. The current climatic conditions in Tartu,

Helsinki, and Rauma roughly correspond to the anticipated future climatic conditions in Oulu

and in the distribution area of the southern variety (Fig. 2, Fig. S4). The historic conditions in

Oulu roughly correspond to the future conditions in Svanvik and in the distribution area of the

northern variety (Fig. 2, Fig. S4)).

In addition to modelling plant performance as a function of experimental garden and other

variables, we conducted an alternative model using wetaher data instead of experimental garden.

To represent weather, we chose annual mean temperature at the experimental gardens across

experimental years. Climate parameters tend to be multicollinear, and we do not know which

specific aspects of temperature these plants respond to. Thus, mean annual temperature is used as

a large-scale descriptor of differences between experimental gardens.

<u>Statistical analyses</u>

To test the performance of the two varieties in the different gardens we analysed their combined

performance using aster models (Geyer, Wagenius, & Shaw, 2007; Shaw et al., 2008). These

models allow the joint analysis of several life history components that together quantify

differences in performance, such as survival, flowering and flowering abundance. The complete

model (Fig. S2) included plant survival and flowering in each year (0 or 1; Bernoulli error

distribution) plus flowering abundance (zero-truncated Poisson distribution of error) that were

conditional on survival in the previous year. Whether a plant flowered or not was also

conditional on it having survived in the year of flowering. The abundance of flowers per

individual was further conditional on that the individual flowered that year.

These combined responses were modelled using the *aster* call in the aster package (Geyer et al.,

2007; Geyer, 2017) in two main models: one testing the effect of *Garden,* and one testing the

effect of *Temperature* (mean annual temperature over 2013-2016). Specifically, the responses

were modelled as a function of *original size* at planting (continuous, square-root transformed and

thereafter centred around the mean of each variety; see Fig. S3 for distribution of original size

per variety and experimental garden), *Variety* (categorical: Southern and Northern) and either

Garden (categorical, 5 levels) or *Temperature* (continuous) and the interaction between *Variety*

and either *Garden* or *Temperature*. To rule out the effect of *planting time* (early or late summer)

we also ran a model including this variable, which did not improve model fit (change in deviance

1.96, change in df=1, p=0.16). The estimated mean value parameters represent overall flowering

output, that is, the expected flower count during the whole experiment for plants of average

original size (Geyer, 2019). We estimate these for each variety at each experimental garden and

hereafter refer to these predicted values as the overall performance.

As we assumed random variation between the three *Plots* in each garden and between the five to

six *Seed sampling sites* of each variety, we also fitted random-effect aster models for both main

fixed effects (i.e. for the *Garden* and *Temperature* model) using the *reaster* function in the *aster*

package (Geyer, 2017). Predicting plant performance based on random effects models is not

straightforward nor directly possible within the aster models package (Geyer et al., 2019; Geyer,

Ridley, Latta, Etterson, & Shaw, 2013; although see Geyer, Ridley, Latta, Etterson, & Shaw,

2010). Nevertheless, as the estimates for the fixed effects were very similar for both the aster

models including only fixed effects and the models also including random effects (see Table S4

for the Garden-models and Table S5 for the Temperature-models) we obtained predicted values

of plant performance based on the fixed effects models using the *predict* function in the *aster*

package (Tables S6 and S7). Finally, for assessing statistical significance of fixed effects, we

conducted likelihood ratio tests (anova) of the full random effects model versus models

excluding individual variables (Table 1).

<u>Summarizing results using local adaptation metrics</u>

For the reciprocal part of the trial, we calculated local adaptation metrics from the mean

estimated combined predicted values obtained through the fixed effects Garden-model (see the

mathematical formulas below and Table S6). We did not include the out-of-range experimental

sites in order to specifically enable quantification of local adaptation patterns of the two varieties

within the distribution area of the variety. This gives an indication of the current situation of

local adaptation, while including the out-of-range gardens would confuse the interpretability of

the metrics over time and space. There are three main approaches to make quantitative

comparisons of local adaptation from performance measures (Blanquart et al., 2013). The

sympatric-allopatric contrast (henceforth called *ΔSA*) focuses on the *difference* between the

average performance of the genotypes in their home environment and non-home environment.

Thus, it describes how well the genotypes overall fit their home-environments. The home vs.

away approach (*HA*) in turn describes the *home environment quality* for each tested genotype,

and is measured by the difference between the fitness of a genotype in its home environment and

the away environment. Finally, the local vs. foreign approach (*LF*) describes the *genotype quality*

for each tested home environment, and is measured by the difference between the fitness of a

genotype in its home environment and the mean fitness of all other genotypes when exposed to

the same environment. These three conceptual approaches relate to different aspects of local

adaptation, all of which lie in the interest of our study: the overall habitat quality, the overall

quality of the variety, and how the two varieties fit the different environmental conditions.

Below, we provide a brief description of the metrics that we used, while a more thorough

definition can be found in the Supplementary Material Text S1 and in Blanquart et al. (2013).

We estimated the mean performance for an average sized plant of each variety in each

experimental garden over four years (based on the aster model predictions). We used these

estimates to calculate local adaptation metrics. Specifically, *ΔSA* for the *subspecies as whole* is

the difference between the mean performance (*p*) of both varieties (northern, N, southern S) in

their home (H) environments and that in the test (away, A) environment:

$$\Delta SA = \frac{p_{NH} + p_{SH}}{2} - \frac{p_{NA} + p_{SA}}{2}$$

386 The Home vs. Away metric (*HA*) for each *variety* is the difference between the performance of

387 each variety in its home environment and that in its away environment.

$$HA_S = p_{SH} - p_{SA}$$
$$HA_N = p_{NH} - p_{NA}$$

390   The Local vs. Foreign metric (*LF*) for each *garden* is the difference between the performance of

391   the local variety and that of the foreign variety.

$$LF_S = p_{SH} - p_{NA}$$
$$LF_N = p_{NH} - p_{SA}$$

The performance estimates range between 0.04 and 5.7 and were therefore scaled to range

between 0 and 1. The *HA*, *LF* and *SA* metrics will, therefore, range between -1 and 1, with values

around zero indicating negligible differences in performance. The closer the value is to 1, the

stronger is the evidence for positive response of the local genotype in the environment, or

suitability of the environment for the genotype, and values close to -1 show the opposite.

All data management and analyses were conducted in R (version 3.5.3; R Core Team 2019).

Results

<u>Realized weather conditions during the experimental years</u>

The weather conditions during the experimental years differed between the experimental gardens

in accordance with what we expected when designing the experiment (Fig. 2, Fig. S4). However,

the observed temperatures were warmer than the historic means. In essence, this means that the

experienced temperature conditions in Svanvik resembled the historic average temperatures in

Oulu (deviation of 0.146°C), while the experienced temperatures in Oulu resembled, or were

even warmer than, the historic average temperatures in the out-of-range gardens in Rauma,

Helsinki, and Tartu (Fig. 2). The northern variety, however, did not experience historical

temperature conditions in any of the experimental sites, as even the coldest site, Svanvik,

deviated from its historic mean annual temperature by 1.146°C . The variation in bioclimatic

variables between gardens and years are shown in Fig. S5. Additional variables describing the

weather conditions during the historic period and experimental years are provided in Figures S4

and S5.

Plant survival

Altogether, 110 out of 590 (18.6%) plant individuals survived in the experiment by 2016. Low

survival is common in transplantation experiments (Dalrymple, Banks, Stewart, & Pullin 2012)

but there were large differences in survival between the varieties and the gardens in our

experiment. The survival percentage across gardens was 8.8% (27/307) for the northern variety

and 29.3% (83/283) for the southern variety. The survival percentage varied between the gardens

from 5.8% in Tartu (9/155) to 51% in Svanvik (52/102; Fig. 3, Table S3).

Individuals of the southern variety survived at a higher percentage every year and in each garden

(Fig. 3; Table S3), yet, the difference in survival between the varieties varied according to the

garden. Both varieties showed roughly the same trend, surviving worst in the out-of-range

gardens (Rauma, Helsinki, and Tartu) and best in the northernmost garden (Svanvik) (Fig. 3;

Table S3).

Flowering and plant size

Mean values of our data on flower presence, flowering abundance and size (cm^2) are presented in

Fig. S6, both across all planted individuals and across all individuals that were alive in the focal

year. Among all the planted individuals, flowering was most frequent and abundant, and plants

grew biggest in Svanvik. The Southern variety also grew well and flowered abundantly in

Helsinki. When considering only the surviving individuals, however, both varieties flowered

frequently and abundantly in Tartu and Rauma and the surviving individuals of the southern

variety grew big in all southern gardens. Yet, in many cases the number of surviving individuals

was very low, for example, only one individual of the northern variety was alive in Tartu in 2016

(Table S3), but it flowered.

<u>Overall performance</u>

Both the Garden and Temperature models revealed similar and significant differences between

the varieties in their overall performance (Table 1; Fig. 4). The Garden model indicated

increased performance of both varieties in gardens towards the north but revealed substantial

differences between the varieties (Fig. 4a; Table S6). Likewise, the Temperature model revealed

a significant negative effect of mean annual temperature on the performance on both varieties,

with a stronger negative effect on the northern variety (Fig 5b; Table S7). Indeed, we found

significant interaction effects between variety vs. experimental garden (Fig. 4a; Table 1) and

annual temperature (Fig. 4b; Table 1). In addition, the original size at planting had a highly

significant effect on plant performance in both models (Table 1).

Of the random effects, *Plot* explained a significant proportion of the residual variance (change in

deviance was 40 and 60 when Plot was omitted from the model for the Garden and Temperature

models, respectively) while *Seed sampling site* explained a marginal and non-significant

proportion of the residual variation (Table 1; Tables S4 and S5).

The aster models provide an overall performance estimate based on flower abundance per variety

in each garden, conditional on survival and presence of flowers. For the southern variety, these

estimates (Fig. 4., Table S6 and S7) were highest in Svanvik, followed by Helsinki (however

with overlapping standard deviations of estimates; Fig. 4 and Table 1). The overall performance

of the northern variety corroborated our hypothesis (Fig. 1b) by showing the highest performance

in its home-environment Svanvik, followed by Oulu, and performing worse in all three out-of-

range gardens (Tartu, Helsinki, and Rauma).

<u>Local adaptation metrics</u>

The local adaptation metrics based on the reciprocal part of the experiment (Tables S6 and S8)

suggested maladaptation to the current home environments of both varieties. For the southern

variety, the *HA*-value of -0.71 indicates that it experiences reduced habitat quality in its home

environment, as it thrives better away (in Svanvik) than at home (in Oulu). The *LF* of 0.26,

however, indicates that the southern variety possesses superior quality in Oulu since it still does

better at home than the tested foreign northern variety does in Oulu. For the northern variety, the

HA of 0.22 indicates that it experiences superior habitat quality in its home environment

(Svanvik) compared to the tested away-environments (Oulu). Nevertheless, the *LF* of -0.75

indicates that this variety is of reduced quality in its home environment, as the foreign (southern)

variety performs better there. Mean *ΔSA* for the subspecies as a whole was -0.25 indicating an

overall negative fit of the environments for the subspecies.

Discussion

Our results reveal clear patterns of adaptation to macroclimatic conditions at the subspecies

level. In general, both varieties showed their best performance within the species' current range.

The southern variety showed higher overall performance in all experimental gardens, compared

to the northern variety. The overall survival, size, and flowering frequency and abundance tended

to be higher for the southern variety. This is in accordance with the results from a greenhouse

experiment conducted by Mäkinen & Mäkinen (1964). According to them the northern variety

flowered only in cold night conditions (12 h 21°C + 12 h 5°C versus 24h 21°C) whereas the

southern variety flowered only in continuously warm conditions but showed increased vegetative

growth in the cold night conditions.

We did not find characteristic indications of local adaptation at the variety level when looking at

the performance of the sampled populations per experimental garden. Individuals of both

varieties survived best, and flowered more frequently and abundantly, in the northernmost

garden Svanvik. This pattern is also evident from the local adaptation metrics, where the *LF*

metric indicated that the southern variety was superior in the home environment of the northern.

Further, the overall performance proxy, the sympatric-allopatric contrast, indicates that the

varieties may have become maladapted to the present conditions prevailing in their current

environments, suggesting the conditions have already changed compared to historical conditions

to which the populations were better adapted. In other words, the varieties are suffering from so

called adaptation-lag (McGraw et al. 2015). When considering the weather conditions during the

experimental years compared to the historic climatic conditions (Fig. 2 (b), Fig. S4), a conclusion

of local adaptation to historical temperature conditions indeed becomes the most parsimonious

explanation. This is revealed by, the fact that the historical mean annual temperature of the

Southern variety was most closely realized in Svanvik during the experimental years while the

historical mean annual temperature for the Northern variety was not realized in any of the

experimental gardens. In addition the mean annual temperature in Oulu resembled that of the

three southern gardens' historical means, that is the climate change treatment that we had

envisioned. Thus, the conditions that we aimed to mimic through our experimental design had in

practise shifted locations one step further north, which we did not take into account initially

when verbalising our hypotheses. Our findings are also supported by a species distribution

modelling study testing the difference between varietal level and subspecies level modelling

(Hällfors et al. 2016). According to the conclusions of this study, the Southern population, if

indeed locally adapted to the historical conditions in its current range, should already by the

2030s find suitable conditions in the current distribution of the Northern variety, while the same

conditions will no longer remain suitable for the Northern variety.

Maladaptation is presumably increasing due to climate change, as the rapid change outpaces the

ability of species to adapt or migrate, and conditions reach beyond the extent of temperature

tolerance (Crespi, 2000; Kooyers et al., 2019). Our findingss are in line with studies showing

signs of maladaptation to current climatic conditions in plant species (see Anderson &

Wadgymar, 2020; Gellie et al., 2016; Kooyers et al., 2019; Wang, et al., 2019; Wilczek et al.,

2014). McGraw et al. (2015) found that the climatic optimum of their study species had shifted

by 140 kilometres. Our study reveals a displacement over a much longer extent: the distance

between the seed sampling sites of the southern variety and the Svanvik experimental sites is

over 500 kilometres. As far as we are aware, this represents the strongest example of climate

change induced maladaptation.

Although maladaptation may be increasing because of climate change, it can also be caused by

underlying population genetic structures, such as genetic drift or founder effects (Crespi, 2000;

Leimu & Fischer, 2008) related to the biogeographic history of the populations. The two

varieties of Siberian primrose probably became spatially separated during the end of the last

glacial period (Mäkinen & Mäkinen, 1964), which first allowed colonization of emerging areas

and later fragmented the population through diminishing suitable habitat as land masses

reformed from underneath the receding ice shelves (see Kreivi et al., 2011). Indeed, genetic and

allelic diversity is known to be remarkably low in all populations of Siberian primrose in

Fennoscandia (Kreivi et al., 2011). However, the consistent pattern of increased performance of

both varieties from south to north, suggests that maladaptation caused by recent change in

climate is a likely explanation for the results of this study. In addition, at least for the Southern

variety, this is not only a question of relative maladaptation (for definition of terms, see Brady et

al., 2019). According to threat assessments, it also shows absolute maladaptation in the form of

population decline (Hyvärinen et al. 2019, results for this species available at

https://punainenkirja.laji.fi/en/results/MX.38659?checklist=MR.424).

Potential confounding effects and general limitations of transplantation experiments

While our results suggest a strong impact of climate on the Siberian primrose, they are not fully

conclusive. Large-scale translocation experiments are complex, and it is important to evaluate

potential methodological pitfalls. Transplantation experiments along the north-south axis

necessarily contain confounding effects, such as day length and other non-thermal environmental

conditions (Saikkonen et al., 2012; Björkman et al., 2016), which cannot be removed in a

transplantation study. Despite the rigorous measures to standardise treatments in all gardens (See

Text S1), we had only one experimental garden at each latitude (albeit with three plots in each

experimental garden). Therefore, random variation in care, pressure from weeds, and weather

events may have confounded the climatic signals. For example, some plots may have

experienced short periods of drought despite our regime of additional watering designed to

mimic the natural conditions experienced by this species of damp seashore meadows. The

roughly three-year observation period of our study may also not fully represent the prevailing

climates in the trial gardens, since inter-annual variation in weather may mask mean climatic

differences, nor does such a short experiment necessarily reveal local adaptation (Bennington et

al. 2012). Long-term studies would even out annual variations, and allow the measurement of

characteristics that become available later in a plant's life cycle and population development

(seed production and germination, adaptation across generations, etc.).

In our study, plants that were larger at the time of planting showed higher overall performance.

Previous studies of reintroductions have shown that the best results (the highest numbers of

surviving plants) are gained by using mature plants or seeds (Dalrymple et al., 2012). However,

in an experimental study, we believe that using mature plants can be confounding since they may

be affected by the climatic conditions of the experimental garden where they are propagated for

the experiment, thus possibly biasing the results through stronger priming effects (see below for

a discussion of the potential effect of Helsinki on the Southern variety). Also, early stages in the

life cycle may be the ones most susceptible to climate change as they define the environments to

which post-germination life-stages of the individuals are exposed (Donohue et al., 2010), and

therefore they should not be omitted from a test setting. On the other hand, while running

experiments from seeds would have allowed us to evaluate performance through germination,

this might have risked the experiment in case of failure to germinate. In fact, even in the

relatively stable greenhouse conditions, the germination percentage of these varieties ranged

between 27-69% (depending on the sampling site; Lehtimäki 2016).

The seeds of each variety were sampled within one region each within a c. 10 km radius (Fig. 1,

Table S1). Thus, from the data at hand we cannot infer patterns across the complete ranges of the

varieties. Nevertheless, the genetic variation within each variety is relatively low (Kreivi et al.,

2011) and the varieties occur in distinct climatic environments (Hällfors et al. 2016), which

makes it plausible to assume that individuals across the range of each variety would respond

similarly to environmental conditions. We did, however, not control for, or try to remove,

maternal carry-over effects, for instance through growing a first generation in identical and

controlled conditions and then utilizing the progeny for the trials. However, the random effect of

seed sampling site did not significantly increase the explanatory power of residual variation. On

the other hand, the lack of replication across the distribution of the varieties inevitably lessens

the potential for drawing strong conclusions about the strength of local adaptation and potential

effects of climate change for the subspecies as a whole.

Another useful performance measure would have been seed production of the tested individuals

from the experimental locations and germination of these seeds. However, because of limited

resources and the possibility of hybridization between the two varieties, we restricted our

observations to the most readily observable performance measures of the planted populations. In

natural conditions, however, each seed capsule usually contains tens to hundreds of seed.

Assuming that pollination was successful (the species is pollinated by a variety of insects

including small flies which are present at all sites), flower number would be a relevant proxy for

seed set. Although germinability of seeds is a more conclusive indicator of reproductive capacity

and population fitness, flowering is also an important indicator of the existence of a necessary

life cycle stage towards offspring. In addition, for this species, flowering has been shown to be

an important fitness-reflecting factor, as failure in flowering is connected to increased extinction

risk of the species (Björnström et al., 2011).

Why the southern variety thrived relatively well in Helsinki remains unclear. The raw proportion

of surviving individuals in Helsinki (31%) was not substantially higher than in Oulu and Rauma

(24.5 and 23%, respectively). However, the overall performance estimate, which describes

population-level flowering output, was substantially higher in Helsinki (4.1) than in the variety's

home environment in Oulu (1.69), and in Rauma (1.01). We suspect that this could have been

caused by the above-mentioned difference in general maintenance or weather conditions that we

were unable to capture through long-term means. Because the individuals were propagated

together in common conditions in Helsinki, the difference between these conditions and those of

the experimental garden, where the individuals ended up, could additionally explain the high

performance in Helsinki. If the individuals planted in Helsinki were primed to the Helsinki

conditions, this might explain the higher performance of them compared to the other southern

gardens and to Oulu. However, as this higher performance was not seen for the northern variety

(although the varieties may be differentially affected by priming) we suspect that the main cause

was stochastic effects.

The effect of climate change on the Siberian primrose

In this study, we have elucidated the role that plasticity towards varying temperature conditions

can play within the boundaries of local adaptation for the Siberian primrose and its two varieties.

We conclude that, especially for the northern variety, the existing plasticity towards warmer

conditions is very limited. As the climate continues to change, migration northwards could

enable the varieties of this species to cope with increasing temperatures. However, because of its

poor dispersal ability, habitat specialisation further reducing chances of range expansion, and

limited temperature tolerance, the available options for this species are restricted to the

production of characteristics suitable for the new climatic conditions *in situ* through evolutionary

adaptation. Nevertheless, it remains unclear how big a role evolutionary adaptation will play

under climatic change in relation to temperature tolerance (Arnold et al., 2019; Charmantier &

Gienapp, 2014; Gienapp et al., 2008; Nicotra et al., 2010). It is also not clear whether selection

can act to increase or decrease plasticity to allow larger variation among phenotypes, which

could then respond adaptively to a range of temperature conditions (Arnold et al., 2019).

Our raw data indicate that those individuals that did survive in the southernmost gardens tended

to be big and to flower frequently and abundantly (Fig. S6). Thus, the elimination of individuals

that are maladapted to the conditions in each experimental garden leaves a survivor population

for which the impact of climate is not as decisive. The adaptive performance of the remnant

plants in the out-of-range gardens highlights how individual features may be a key determinant

for continued survival when a population goes through a bottleneck (Carson, 1990). Whether

such a small founding population of a genetically impoverished species could form a viable

population under rapidly changing conditions is, however, uncertain. However, experiments on

the evolutionary potential in this species are needed to draw more solid conclusions on the

potential for the species to overcome the lag in adaptation that it is likely currently experiencing.

Siberian primrose, a habitat specialist and poor disperser with small and fragmented populations

that harbour low genetic variability and a limited thermal tolerance, is an example of a species

likely to suffer from lack of adaptive capacity under rapid climatic change. Proactive

conservation methods, such as assisted migration, may be needed in order to maintain this

species. Overall, our results cause concern about the viability of the Siberian primrose, especially

the northern variety, for which the detrimental effects of climate change may become evident

within a few decades. In addition, these findings remind us of the need to take the population

level into account when modelling the effect of climate change on species (Souther & McGraw,

2011; Hällfors et al., 2016).

<u>Lessons for translocation trials under altered climate conditions</u>

The results of this study also highlight a critical point in testing for local adaptation and plasticity

under the rapidly changing climate. If populations are not able to track ongoing changes, only

experiments both within and outside the current range of the species (or variety) can provide

accurate information regarding local adaptation and plasticity towards historical, current, and

future conditions. Maladaptation to historical conditions, which currently may prevail only

polewards and even outside of the species range, must be considered a possible scenario (e.g.,

Anderson & Wadgymar, 2020; Gellie et al., 2016; Kooyers et al., 2019; McGraw et al. 2015;

Wilczek et al., 2014). However, the out-of range gardens that were included in this study

(Rauma, Helsinki, and Tartu) can inform us about likely trajectories for future maladaptation in

case the species is not able to adapt evolutionally and climate continues to shift temperature

isoclines. Climate change thus gives cause for re-examining traditional reciprocal and out-of-

range common garden experiments (Anderson & Wadgymar, 2020), and to plan experiments

with climatic and weather data at hand. Comparing responses to future, current, and historical

conditions offers a powerful approach for future studies (Anderson & Wadgymar, 2020; Franks

et al., 2007; Merilä & Hendry, 2014; McGraw et al. 2015; Thomann et al., 2015). Together with

continued monitoring of population abundances (Cotto et al., 2017), such studies can provide

early warning signals for potential feed-back effects on the ecosystem productivity (Schedlbauer

et al. 2018; Curasi et al. 2019) and species that may be close to tipping points and to going

extinct.

Acknowledgements

We are grateful to the botanic gardens that provided staff and premises for conducting the trials:

University of Tartu, University of Helsinki, University of Turku teaching garden in Rauma,

University of Oulu, and NIBIO Svanhovd. We thank Pertti Pehkonen, Outi Pakkanen, and

Toomas Kangro for technical assistance in preparing the experiments. Terhi Ryttäri and Henry

Väre gave advice on the species and required soil properties. Ritva Hiltunen and Tuomas

Kauppila assisted in collection of seeds, Elina Vaara helped with permits for seed and plant

transfers, Hanna Finne with data management, and Bess Hardwick with acquiring weather data.

MHH was supported by the University of Helsinki Research Fund, LUOVA – Doctoral

Programme in Wildlife Biology Research and the Jane and Aatos Erkko Foundation through the

Research Centre for Ecological Change, University of Helsinki. The study was supported by the

Academy of Finland grant 126915 and Societas pro Fauna et Flora Fennica. The authors declare

no conflicts of interest.

Author's contributions

MHH, SL, and MTH conceived the idea and designed methodology, LS contributed to the selection of experimental gardens; MHH and LS collected seeds; MHH, SL, IL, and MTH produced plant material and coordinated the experiments; all authors took part in setting up the experiments and collecting the data; MHH analyzed the data and led the writing of the manuscript. All authors contributed critically to the manuscript drafts and gave final approval for publication.

Data accessibility

The data are available through GitHub at https://github.com/MariaHallfors/Primula-nutans-translocation

References

Anderson, J. T., & Wadgymar, S. M. (2020). Climate change disrupts local adaptation and favours upslope migration. *Ecology Letters*, *23*(1), 181–192. doi: 10.1111/ele.13427

Arnold, P. A., Nicotra, A. B., & Kruuk, L. E. B. (2019). Sparse evidence for selection on phenotypic plasticity in response to temperature. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *374*(1768), 20180185. doi: 10.1098/rstb.2018.0185

Atkins, K. E., & Travis, J. M. J. (2010). Local adaptation and the evolution of species' ranges under climate change. *Journal of Theoretical Biology*, *266*(3), 449–457. doi: 10.1016/j.jtbi.2010.07.014

Banta, J. A., Ehrenreich, I. M., Gerard, S., Chou, L., Wilczek, A., Schmitt, J., … Purugganan, M. D. (2012). Climate envelope modelling reveals intraspecific relationships among

flowering phenology, niche breadth and potential range size in Arabidopsis thaliana.

Ecology Letters, *15*(8), 769–777. doi: 10.1111/j.1461-0248.2012.01796.x

Bennington, C. C., N. Fetcher, M. C. Vavrek, G. R. Shaver, K. J. Cummings, and J. B. McGraw.

(2012). Home site advantage in two long-lived arctic plant species: results from two 30-

year reciprocal transplant studies. *Journal of Ecology* 100: 841-851. doi: 10.1111/j.1365-

2745.2012.01984.x

Björkman, A.D., Vellend, M., Frei, E.R., and Henry, G.H.R. (2016). Climate adaptation is not

enough: warming does not facilitate success of southern tundra plant populations in the

high Arctic. Glob. Chang. Biol. 23: 1540–1551. doi:10.1111/gcb.13417.

Björnström, T., Hyvärinen, M., Siikamäki, P., Strengell, H., Väre, H., & Tuomi, J. (2011).

Demographic variability and population viability of an endemic variety of the Siberian

primrose. *Annales Botanici Fennici*, *48*(2), 107–119. Retrieved from JSTOR.

Blanquart, F., Kaltz, O., Nuismer, S. L., & Gandon, S. (2013). A practical guide to measuring

local adaptation. *Ecology Letters*, *16*(9), 1195–1205. doi: 10.1111/ele.12150

Brady, S. P.et al. (2019) Understanding Maladaptation by Uniting Ecological and Evolutionary

Perspectives. *The American Naturalist* 194:4, 495-515

Carson, H. L. (1990). Increased genetic variance after a population bottleneck. *Trends in Ecology*

& Evolution, *5*(7), 228–230. doi: 10.1016/0169-5347(90)90137-3

Ceballos, G., Ehrlich, P. R., Barnosky, A. D., García, A., Pringle, R. M., & Palmer, T. M.

(2015). Accelerated modern human–induced species losses: Entering the sixth mass

extinction. *Science Advances*, *1*(5), e1400253. doi: 10.1126/sciadv.1400253

Charmantier, A., & Gienapp, P. (2014). Climate change and timing of avian breeding and

migration: evolutionary versus plastic changes. *Evolutionary Applications*, *7*(1), 15–28.

doi: 10.1111/eva.12126

Chevin, L.-M., Lande, R., & Mace, G. M. (2010). Adaptation, Plasticity, and Extinction in a

Changing Environment: Towards a Predictive Theory. *PLOS Biology*, *8*(4), e1000357.

doi: 10.1371/journal.pbio.1000357

Cotto, O., Wessely, J., Georges, D., Klonner, G., Schmid, M., Dullinger, S., … Guillaume, F.

(2017). A dynamic eco-evolutionary model predicts slow response of alpine plants to

climate warming. *Nature Communications*, *8*(1), 15399. doi: 10.1038/ncomms15399

Crespi, B. J. (2000). The evolution of maladaptation. *Heredity*, *84*(6), 623–629. doi:

10.1046/j.1365-2540.2000.00746.x

Curasi, S.R., Parker, T. C., Rocha, A.V., Moody, M.L., J. Tang, and N. Fetcher. (2019).

Differential responses of ecotypes to climate in a ubiquitous arctic sedge: implications for

future ecosystem C cycling. *New Phytologist* 223: 180-192

746
Dalrymple, S.E., Banks, E., Stewart, G.B., & Pullin, A.S. (2012) A Meta-Analysis of Threatened

Plant Reintroductions from Across the Globe. In: Machinski and Haskins (Eds.), *Plant*

Reintroduction in a Changing Climate. Promises and Perils (pp. 31- 50). Island Press.

Dawson, I. K., Vinceti, B., Weber, J. C., Neufeldt, H., Russell, J., Lengkeek, A. G., …

Jamnadass, R. (2011). Climate change and tree genetic resource management:

maintaining and enhancing the productivity and value of smallholder tropical

agroforestry landscapes. A review. *Agroforestry Systems*, *81*(1), 67–78. doi:

10.1007/s10457-010-9302-2

Díaz, S., Settele, J., Brondízio, E., Ngo, H. T., Guèze, M., Agard, J., … Vilá, B. (2019).

Summary for policymakers of the global assessment report on biodiversity and ecosystem

services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem

Services. 44.

Donohue, K., Rubio de Casas, R., Burghardt, L., Kovach, K., & Willis, C. G. (2010).

Germination, Postgermination Adaptation, and Species Ecological Ranges. *Annual*

Review of Ecology, Evolution, and Systematics, *41*(1), 293–319. doi: 10.1146/annurev-

ecolsys-102209-144715

ENSCONET 2009: ENSCONET Seed Collecting Manual for Wild Species. Available at:

http://ensconet.maich.gr/PDF/Collecting_protocol_English.pdf

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces

for global land areas. *International Journal of Climatology*, *37*(12), 4302–4315. doi:

10.1002/joc.5086

Frankham, R., Ballou, J. D., Dudash, M. R., Eldridge, M. D. B., Fenster, C. B., Lacy, R. C., …

Ryder, O. A. (2012). Implications of different species concepts for conserving

biodiversity. *Biological Conservation*, *153*, 25–31. doi: 10.1016/j.biocon.2012.04.034

Franks, S. J., Sim, S., & Weis, A. E. (2007). Rapid evolution of flowering time by an annual

plant in response to a climate fluctuation. *Proceedings of the National Academy of*

Sciences, *104*(4), 1278–1282. doi: 10.1073/pnas.0608379104

Frenne De, P., Graae, B. J., Rodríguez☐Sánchez, F., Kolb, A., Chabrerie, O., Decocq, G., …

Verheyen, K. (2013). Latitudinal gradients as natural laboratories to infer species'

responses to temperature. *Journal of Ecology*, *101*(3), 784–795. doi: 10.1111/1365-

2745.12074

Gao, S., Mo, L., Zhang, L., Zhang, J., Wu, J., Wang, J., … Gao, Y. (2018). Phenotypic plasticity

vs. local adaptation in quantitative traits differences of Stipa grandis in semi-arid steppe,

China. *Scientific Reports*, *8*(1), 3148. doi: 10.1038/s41598-018-21557-w

GBIF (2013). *Global Biodiversity Information Facility Data Portal.*

Gellie, N. J. C., Breed, M. F., Thurgate, N., Kennedy, S. A., & Lowe, A. J. (2016). Local

maladaptation in a foundation tree species: Implications for restoration. *Biological*

Conservation, *203*, 226–232. doi: 10.1016/j.biocon.2016.08.036

Geyer, C. J., Wagenius, S., & Shaw, R. G. (2007). Aster models for life history analysis.

Biometrika, *94*(2), 415–426. doi: 10.1093/biomet/asm030

Geyer, Charles J., Ridley, C. E., Latta, R. G., Etterson, J. R., & Shaw, R. G. (2010). *Aster*

Models with Random Effects via Penalized Likelihood [Technical Report]. Retrieved

from University of Minnesota website: http://conservancy.umn.edu/handle/11299/199694

Geyer, Charles J., Ridley, C. E., Latta, R. G., Etterson, J. R., & Shaw, R. G. (2013). LOCAL

ADAPTATION AND GENETIC EFFECTS ON FITNESS: CALCULATIONS FOR

EXPONENTIAL FAMILY MODELS WITH RANDOM EFFECTS. *The Annals of*

Applied Statistics, *7*(3), 1778–1795.

Geyer, C.J., (2017). R package aster2 (Aster Models), version 0.3. https://cran.r-

project.org/package=aster2.

Geyer, C.J., (2019): The Aster Package Tutorial. Available at: https://cran.r-

project.org/web/packages/aster/vignettes/tutor.pdf

798
Gienapp, P., Teplitsky, C., Alho, J. S., Mills, J. A., & Merilä, J. (2008). Climate change and

evolution: disentangling environmental and genetic responses. *Molecular Ecology*, *17*(1),

167–178. doi: 10.1111/j.1365-294X.2007.03413.x

Greiser, C., Hylander, K., Meineri, E., Luoto, M., & Ehrlén, J. (2020). Climate limitation at the

cold edge: contrasting perspectives from species distribution modelling and a transplant

experiment. *Ecography*, *n/a*(n/a). doi: 10.1111/ecog.04490

Hällfors, M. H., Vaara, E. M., Hyvärinen, M., Oksanen, M., Schulman, L. E., Siipi, H., &

Lehvävirta, S. (2014). Coming to Terms with the Concept of Moving Species Threatened

by Climate Change – A Systematic Review of the Terminology and Definitions. *PLOS*

ONE, *9*(7), e102979. doi: 10.1371/journal.pone.0102979

Hällfors, M. H., Liao, J., Dzurisin, J., Grundel, R., Hyvärinen, M., Towle, K., Wu., G.,

Hellmann, J. J. (2016). Addressing potential local adaptation in species distribution

models: implications for conservation under climate change. *Ecological Applications*,

26(4), 1154–1169. doi: 10.1890/15-0926

Hargreaves, A. L., Samis, K. E., & Eckert, C. G. (2014). Are Species' Range Limits Simply

Niche Limits Writ Large? A Review of Transplant Experiments beyond the Range. *The*

American Naturalist, *183*(2), 157–173. doi: 10.1086/674525

Hill, J. K., Griffiths, H. M., & Thomas, C. D. (2011). Climate Change and Evolutionary

Adaptations at Species' Range Margins. *Annual Review of Entomology*, *56*(1), 143–159.

doi: 10.1146/annurev-ento-120709-144746

Hultén, E., & Fries, M. (1986). *Atlas of North European vascular plants: north of the Tropic of*

Cancer, vol. 1-3. Königstein, Koeltz.

Hijmans, R.J. (2019). raster: Geographic Data Analysis and Modeling. R package version

2.8-19. https://CRAN.R-project.org/package=raster

Hijmans, R.J., Phillips, S., Leathwick, J. & Jane Elith, J. (2017). dismo: Species

Distribution Modeling. R package version 1.1-4. https://CRAN.R-project.org/package=dismo

Hyvärinen, E., Juslén, A., Kemppainen, E., Uddström, A. & Liukko, U.-M. (eds.). The 2019 Red

828   List of Finnish Species. (Ministry of the Environment & Finnish Environment Institute,

829   2019).

830   IPCC, 2019: Summary for Policymakers. In: *Climate Change and Land: an IPCC special report*

831   *on climate change, desertification, land degradation, sustainable land management, food*

832   *security, and greenhouse gas fluxes in terrestrial ecosystems* [P.R. Shukla, J. Skea, E.

833   Calvo Buendia, V. Masson-Delmotte, H.- O. Pörtner, D. C. Roberts, P. Zhai, R. Slade, S.

834   Connors, R. van Diemen, M. Ferrat, E. Haughey, S. Luz, S. Neogi, M. Pathak, J. Petzold,

835   J. Portugal Pereira, P. Vyas, E. Huntley, K. Kissick, M. Belkacemi, J. Malley, (eds.)].

836   Joint Research Centre 2014: *Gridded Agro-Meteorological Data in Europe.* URI:

837   http://data.europa.eu/88u/dataset/jrc-marsop4-7-weather_obs_grid_2015

838   Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology Letters*, *7*(12),

839   1225–1241. doi: 10.1111/j.1461-0248.2004.00684.x

840   Kooyers, N. J., Colicchio, J. M., Greenlee, A. B., Patterson, E., Handloser, N. T., & Blackman,

841   B. K. (2019). Lagging Adaptation to Climate Supersedes Local Adaptation to Herbivory

842   in an Annual Monkeyflower. *The American Naturalist*, *194*(4), 541–557. doi:

843   10.1086/702312

844   Kreivi, M., Aspi, J., & Leskinen, E. (2011). Regional and local spatial genetic structure of

845   Siberian primrose populations in Northern Europe. *Conservation Genetics*, *12*(6), 1551–

846   1563. doi: 10.1007/s10592-011-0252-7

847   Kreivi, M., Huttunen, S., & Aspi, J. (2006). Isolation and characterization of polymorphic

848   microsatellite markers from Primula nutans (Primulaceae). *Molecular Ecology Notes*,

849   *6*(2), 334–336. doi: 10.1111/j.1471-8286.2005.01222.x

Kreyling, J., Buhk, C., Backhaus, S., Hallinger, M., Huber, G., Huber, L., … Beierkuhnlein, C.

(2014). Local adaptations to frost in marginal and central populations of the dominant

forest tree Fagus sylvatica L. as affected by temperature and extreme drought in common

garden experiments. Ecology and Evolution, 4(5), 594–605. doi: 10.1002/ece3.971

Lampinen, R., Lahti, T., Heikkinen, M. 2012. *Kasviatlas 2011*. -- Helsingin Yliopisto,

Luonnontieteellinen keskusmuseo, Helsinki. Distribution maps available at:

http://www.luomus.fi/kasviatlas.

Lehtimäki, I. (2016). *Uhanalaisen ruijanesikon suojelu. Etäsuojelun ja avustetun leviämisen*

näkökulma. Master's thesis [In Finnish]. Department of Biosciences, University of

Helsinki.

Leimu, R., & Fischer, M. (2008). A Meta-Analysis of Local Adaptation in Plants. PLoS ONE,

3(12), e4010. doi: 10.1371/journal.pone.0004010

Lenoir, J., & Svenning, J.-C. (2015). Climate-related range shifts – a global multidimensional

synthesis and new research directions. *Ecography*, *38*(1), 15–28. doi:

10.1111/ecog.00967

Mäkinen, L., & Mäkinen, Y. (1964). The distribution, ecology, morphology and taxonomy of

Primula nutans Georgi ssp. finmarchica (Jacq.) Löve & Löve. *Annales Botanici Fennici*,

1(4), 273–291. Retrieved from JSTOR.

McGraw, J. B., J. B. Turner, S. Souther, C. C. Bennington, M. C. Vavrek, G. R.Shaver, and N.

Fetcher. (2015). Northward displacement of optimal climate conditions for ecotypes of

Eriophorum vaginatum L. across a latitudinal gradient in Alaska. Global Change Biology

21: 3827–3835. doi: 10.1111/gcb.12991

Merilä, J. (2012). Evolution in response to climate change: In pursuit of the missing evidence.

BioEssays, *34*(9), 811–818. doi: 10.1002/bies.201200054

Merilä, J., & Hendry, A. P. (2014). Climate change, adaptation, and phenotypic plasticity: the

problem and the evidence. *Evolutionary Applications*, *7*(1), 1–14. doi: 10.1111/eva.12137

Nicotra, A. B., Atkin, O. K., Bonser, S. P., Davidson, A. M., Finnegan, E. J., Mathesius, U., …

van Kleunen, M. (2010). Plant phenotypic plasticity in a changing climate. *Trends in*

Plant Science, *15*(12), 684–692. doi: 10.1016/j.tplants.2010.09.008

Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts

across natural systems. *Nature*, *421*(6918), 37–42. doi: 10.1038/nature01286

Pelini, S. L., Diamond, S. E., MacLean, H., Ellison, A. M., Gotelli, N. J., Sanders, N. J., & Dunn,

R. R. (2012). Common garden experiments reveal uncommon responses across

temperatures, locations, and species of ants. *Ecology and Evolution*, *2*(12), 3009–3015.

doi: 10.1002/ece3.407

Pöyry, J., Luoto, M., Heikkinen, R. K., Kuussaari, M., & Saarinen, K. (2009). Species traits

explain recent range shifts of Finnish butterflies. *Global Change Biology*, *15*(3), 732–

743. doi: 10.1111/j.1365-2486.2008.01789.x

R Core Team 2019: R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.

Saikkonen, K., Taulavuori, K., Hyvönen, T., Gundel, P. E., Hamilton, C. E., Vänninen, I., …

Helander, M. (2012). Climate change-driven species' range shifts filtered by

photoperiodism. *Nature Climate Change*, *2*(4), 239–242. doi: 10.1038/nclimate1430

Schedlbauer, J. L., N. Fetcher, K. Hood, M. L. Moody, and J. Tang. (2018). Effect of growth

temperature on photosynthetic capacity and respiration in three ecotypes of Eriophorum

vaginatum. *Ecology and Evolution* 8: 3711-3725 doi: 10.1002/ece3.3939

Settele, J., Scholes, R., Betts, R., Bunn, S., Leadley, P., Nepstadt, D., & Taboada, M. A. (2014).

Terrestrial and inland water systems. In: Cambridge, United Kingdom and New York,

NY, USA, pp. 271-359.

Shaw, R. G., Geyer, C. J., Wagenius, S., Hangelbroek, H. H., Etterson, J. R., Taylor, A. E. P. D.,

& DeAngelis, E. D. L. (2008). Unifying Life History Analyses for Inference of Fitness

and Population Growth. *The American Naturalist*, *172*(1), E35–E47. doi: 10.1086/588063

Silvertown, J., & Charlesworth, D. (2001). *Introduction to Plant Population Biology*. 4th edition.

Blackwell Science.

Souther, S., and McGraw, J.B. (2011). Evidence of Local Adaptation in the Demographic

Response of American Ginseng to Interannual Temperature Variation. *Conserv. Biol.* 25:

922–931. doi:10.1111/j.1523-1739.2011.01695.x.

Thiers, B. (2016) Index Herbariorum: A Global Directory of Public Herbaria and Associated

Staff. New York Botanical Garden's Virtual Herbarium.

http://sweetgum.nybg.org/science/ih/

Thomann, M., Imbert, E., Engstrand, R. C., & Cheptou, P.-O. (2015). Contemporary evolution of

plant reproductive strategies under global change is revealed by stored seeds. *Journal of*

Evolutionary Biology, *28*(4), 766–778. doi: 10.1111/jeb.12603

Ulvinen, T. (1997). Rujanesikko. In: *Uhanalaiset kasvimme*. Ryttäri T. and Kettunen T. (eds.).

[In Finnish]. Suomen ympäristökeskus, Helsinki.

Urban, M. C. (2015). Accelerating extinction risk from climate change. *Science*, *348*(6234),

571–573. doi: 10.1126/science.aaa4984

Valladares, F., Matesanz, S., Guilhaumon, F., Araújo, M. B., Balaguer, L., Benito-Garzón, M.,

… Zavala, M. A. (2014). The effects of phenotypic plasticity and local adaptation on

forecasts of species range shifts under climate change. *Ecology Letters*, *17*(11), 1351–

1364. doi: 10.1111/ele.12348

Wang, Y., Pedersen, J. L. M., Macdonald, S. E., Nielsen, S. E., & Zhang, J. (2019).

Experimental test of assisted migration for conservation of locally range-restricted plants

in Alberta, Canada. *Global Ecology and Conservation*, *17*, e00572. doi:

10.1016/j.gecco.2019.e00572

Wilczek, A. M., Cooper, M. D., Korves, T. M., & Schmitt, J. (2014). Lagging adaptation to

warming climate in Arabidopsis thaliana. *Proceedings of the National Academy of*

Sciences, *111*(22), 7906–7913. doi: 10.1073/pnas.1406314111

Zenni, R. D., Bailey, J. K., & Simberloff, D. (2014). Rapid evolution and range expansion of an

invasive plant are driven by provenance -environment interactions. *Ecology Letters*,

17(6), 727–735. doi: 10.1111/ele.12278

Figure captions and tables

FIGURE 1. Geographical distribution of seed sampling sites and experimental gardens (a), and

hypotheses of plant performance in experimental gardens (b-e). Panel (a) shows the geographical

distribution of seed sampling sites and experimental gardens with occurrences of *Primula nutans* ssp.

finmarchica marked by dark grey points: Var. *finmarchica* occurs by the Arctic Sea in N-Norway and var.

jokelae by the Bothnian bay in Finland and Sweden and the White Sea in Russia. Red = seed sampling

sites of the southern variety (var. *jokelae*) in Finland; Blue= seed sampling sites of the northern variety

(var. *finmarchica*) in Norway. Source for occurrences of Siberian primrose: Global Biodiversity

Information Facility (GBIF 2013); Kastikka (Finnish plant distribution database; Lampinen, Lahti, &

Heikkinen 2012); Hertta (database of Finnish Environment Institute, unpublished data); occurrences in

Russia based on information from herbarium specimens (from collections in the herbarium of the Finnish

Museum of Natural History [H *sensu* Thiers 2016] and the herbarium of the University of Turku [TUR

sensu Thiers 2016] and the distribution map by Hultén and Fries [1986]). Panels (b-e) show hypothesized

overall performance of the tested varieties in all experimental gardens following opposing underlying

scenarios of (b) local relative adaptation (*sensu* Brady et al. 2019) at the varietal and subspecies level, (c)

relative adaptation of the subspecies to its current environment vs. areas outside it, (d) tolerance (through

plasticity) towards all tested conditions (including those not currently present within the occurrence area

of the subspecies), and (e) relative maladaptation caused by climate change (see text for hypotheses).

Dashed area demarks within-range gardens, i.e. the reciprocal part of the experiment. Red= southern

variety; blue = northern variety.

FIGURE 2. Mean annual temperature of seed sampling sites and experimental gardens. Black points

show historic mean temperatures (1970-2000; 10 minutes resolution; (Fick & Hijmans, 2017) and gray

points show the future projections of mean temperatures (CMIP5 for 2050, 10 minutes resolution,

HADGEM2-ES model; (Fick & Hijmans, 2017) for each experimental gardens and seed sampling sites

(Northern= seed sampling sites of the northern variety; Southern= seed sampling sites of the southern

variety). For experimental gardens, open circles show mean temperature during the experimental years

2013-2016. The experimental gardens and seed sampling sites within the species range are outlined by a

dashed gray box, whereas the blue box indicates the sites within the range of the northern variety and the

red box the sites within the range of the southern variety. Corresponding figures for precipitation sum and

other climatic variables in Fig. S4. The future temperatures in Oulu approach the historic ones in Tartu,

Helsinki and Rauma, and the future temperatures in Svanvik approach the historic ones in Oulu. During

the experiment, mean temperatures were higher than the historic means. The conditions in Svanvik

corresponded with historical means in Oulu. Those in Oulu approached the historic means in the southern

gardens.

FIGURE 3. Mean survival per variety across gardens and years. The total number of individuals planted

in 2013 appear at the top of the plot (both original plants and additional plants due to high die-off in the

first summer). Red points = southern variety; blue points = northern variety. The size of the point

indicates year: big point= 2014; medium point= 2015; small point = 2016. Please refer to Table S3 for the

exact numbers of individuals per year and garden, and percentage decrease over year.

FIGURE 4. Predicted overall performance of the two varieties across (a) experimental gardens, and (b)

mean annual temperature over the experimental period 2013-2016. Error bars represent 95% confidence

intervals. Predictions are made for plants of average original size and based on the fixed effects aster

model (i.e., not including *Plot* and *Seed sampling site* as random effects). Red= southern variety; blue=

northern variety.

TABLE 1. Change in deviance when a single variable was omitted from the full random effects models.

Summary from comparing aster models testing the fixed effects of experimental garden (left-hand side of

table) or annual temperature (right hand side of table), variety and original size on plant performances, as

well as *Plot* and *Seed sampling site* as random effects. Statistical significance of predictor variables was

assessed using likelihood ratio tests, that is, a model where one variable was omitted was compared to the

full model using anova.

(a)

(b) Local adaptation of varieties

(c) Local adaptation of subspecies

(d) No adaptation

(e) Maladaptation

Garden-model				Annual temperature -model		
p-value	df	Change in deviance	Variable	Change in deviance	df	p-value
<0.001	1	62.4	Original size	60.2	1	<0.001
<0.05	4	11	Variety:Garden/Ann.Temp.	8.8	1	<0.01
<0.01	5	20	Variety (+ Variety:Garden/Ann. Temp.)	17.2	2	<0.001
<0.05	8	19.6	Garden/Ann. Temp. (+ Variety:Garden/Ann. Temp.)	14.2	2	<0.001
<0.001	0	46	Plot	60	0	<0.001
0.09	0	1.8	Seed sampling site	2.2	0	0.07