Translocation of an arctic seashore plant reveals signs of maladaptation to altered climatic

conditions

Hällfors M.H.[*a,b], Lehvävirta S. [a,c], Aandahl T.R.[d], Lehtimäki I.-M.[a], Nilsson L.O. [d,e],

Ruotsalainen A.L.[f], Schulman L.E.[a], Hyvärinen M.-T.[a]

[a] Botany Unit, Finnish Museum of Natural History, P.O. Box 7, FI-00014 University of Helsinki,

Finland.

[b] Research Centre for Environmental Change, Organismal and Evolutionary Biology Research

Programme, Faculty of Biological and Environmental Sciences, P.O. Box 65, FI-00014

University of Helsinki, Finland.

[c] Department of Landscape Architecture, Planning and Management, Swedish University of

Agricultural Sciences, Alnarp, Sweden

[d] Division of Environment and Natural Resources, Norwegian Institute of Bioeconomy Research

(NIBIO), P.O. Box 115, NO-1431 Ås, Norway.

[e] Halmstad University, PO Box 823, SE-30118 Halmstad, Sweden.

[f] Department of Ecology and Genetics, P.O. Box 3000, FI-90014 University of Oulu, Finland.

[*] corresponding author. Contact information: Maria Hällfors, Research Centre for Environmental

Change, Organismal and Evolutionary Biology Research Programme, Faculty of Biological and

Environmental Sciences, P.O. Box 65, FI-00014 University of Helsinki, Finland, e-mail:

maria.hallfors@helsinki.fi; phone: +358 40 721 34 74

Abstract

Ongoing anthropogenic climate change alters the local climatic conditions to which species may

be adapted. Information on species' climatic requirements and on variation in intraspecific

adaptation to climate is necessary for predicting the effects of climate change on biodiversity.

We used a climatic gradient to test whether populations of two allopatric varieties of an arctic

seashore herb (*Primula nutans* ssp. *finmarchica*) show adaptation to their local climatic

conditions and how a future warmer climate may affect them. Our experimental set-up combined

i) a reciprocal translocation within the distribution range of the species with ii) an experiment

testing performance of the sampled populations in warmer climatic conditions south of their

range. We monitored survival, size, and flowering over four growing seasons as measures of

performance. We found that plants of both varieties performed better in experimental gardens

towards the north. Interestingly, highest up in the north, representatives of the southern variety

outperformed the northern one. Supported by weather data, this suggests that the range of both

varieties is at least partly outside their climatic optima. Further warming would make the current

environments of both varieties even less suitable. Our results thus show maladaptation of

populations of both varieties to their current home environments and that their overall

performance is lower in warmer conditions. We conclude that *Primula nutans* ssp. *finmarchica* is

already suffering from reduced viability related to changes in climate, and that further warming

may be detrimental to this subspecies, especially the northern variety. These results reveal that it

is not sufficient to run reciprocal experiments: adaptation needs also to be tested outside the

current range of the focal taxon in order to include historic conditions to which they have

possibly adapted over time.

Keywords

botanic garden, conservation, global change, local adaptation, Siberian primrose, threatened

species, transplant experiment

Introduction

Ongoing climate change threatens biodiversity and is predicted, and even shown, to lead to

declines in populations and to extinctions of species (Ceballos et al., 2015; Dawson et al., 2011;

Díaz et al., 2019; Settele et al., 2014; Urban, 2015). How the threat of climate change on

biodiversity is manifested in the response of species and populations is crucial for understanding

their abilities to cope with the change. Plants offer several key functions in ecosystems, such as

carbon assimilation and biomass production, and thereby provide resources for other species.

The ability of populations to respond, *in situ*, to shifting environmental conditions can be a result

of plasticity towards a broad range of conditions, or it can be mediated through evolutionary

adaptation (Chevin et al., 2010; Gao et al., 2018; Gienapp et al., 2008; Merilä & Hendry, 2014).

Phenotypic plasticity can allow organisms to respond rapidly to changes in the environment, and

hence, it has often been viewed as the main strategy to respond to environmental changes

(Merilä, 2012). In contrast, evolutionary processes are often slow, especially for species with

long generation times, such as perennial plants. Some species are showing signs of

acclimatization or adaptation to new conditions, whereas others are either declining or dispersing

to new areas with a favourable environment (Lenoir & Svenning, 2015; Parmesan & Yohe, 2003;

Pöyry et al., 2009). To predict the effects of climate change on biodiversity and to plan effective

conservation strategies, we need both evaluations of how the current distribution areas of species

are changing, and approximations of the genetic and phenotypic potential of species to endure

these changes.

Species are often used as the basic taxonomic unit in plant conservation and predictions on the

effects of environmental change usually relate to the species level (Frankham et al., 2012).

Populations of a particular species may, however, be locally adapted, and therefore have

divergent affinities to abiotic conditions (Banta et al., 2012) and show intraspecific differences in

climatic tolerance (Atkins & Travis, 2010; Hill et al., 2011). However, in their review on local

adaptation in plant species, Leimu and Fisher (2008) found that local adaptation of populations

to environmental conditions is not as widespread as often assumed. Local adaptation may be

favoured mostly in cases where populations occur in differing environments and gene flow

between them is restricted (Kawecki & Ebert, 2004). According to Kawecki and Ebert (2004)

"the resident genotypes would have on average a higher fitness in their local habitat than

genotypes originating from other habitats". Understanding and taking into account intraspecific

variation in the adaptation to climatic conditions is necessary to reach balanced estimates of

climate change impact at the level of populations (Hällfors et al., 2016; Valladares et al., 2014).

The speed of climate change may outpace the ability of populations to respond adaptively,

thereby causing maladaptation instead of adaptation or acclimatization. Thus, the need to

estimate the effect that climate change has had on natural populations to date, has become

increasingly evident (Anderson & Wadgymar, 2020; Franks et al., 2007; Kooyers et al., 2019;

Thomann et al., 2015; Wilczek et al., 2014; Zenni et al., 2014).

Manipulative experiments have the potential to provide information on how strong a role climate

plays in defining favourable conditions of a species or population (Hargreaves et al., 2014;

Kawecki & Ebert, 2004; Kreyling et al., 2014; Pelini et al., 2012). Results from experiments

where the climatic conditions for the focal taxa are altered can be incorporated into species

distribution models to improve predictions (Greiser et al., 2020). By reciprocally transplanting

individuals of two or more populations to conditions representing a foreign environment and a

home environment, we can effectively measure intraspecific variation that allow maintaining

fitness under varying temperature conditions (Blanquart et al., 2013; Kawecki & Ebert, 2004).

Experiments can also test species' responses to future conditions, for example, through

treatments mimicking increased temperatures, and inform us about the extent to which plasticity

contributes towards population resilience under climate change. Furthermore, such experiments

can reveal locally adaptive fixed traits, which would signify that the survival of the species in the

future would require either rapid microevolution *in situ* or assisted migration (*sensu* Hällfors et

al., 2014) to new favourable areas.

Here, we describe a study on Siberian primrose (*Primula nutans* ssp. *finmarchica* (Jacq.) Á.

Löve & D.), in which we sampled populations of two of its varieties (the northern var.

finmarchica and the southern var. *jokelae* L. Mäkinen & Y. Mäkinen); while acknowledging that

our sampling did not cover the varieties' full distribution, for clarity we refer to the populations

of the different varieties by 'variety'. We performed i) a reciprocal transplant experiment and ii)

an experiment of out-of-range transplantation. The former tests the importance of intraspecific

local adaptation versus insensitivity towards temperature (Fig 1 (b) versus (c)), (Fig 1 (c) versus

(d)). The experiment was conducted across four growing seasons using a botanic garden network

in Finland, Norway, and Estonia (Fig. 1). Our five experimental gardens represent the i) home

environments of both varieties, forming the reciprocal component in the experiment (Oulu and

Svanvik; Fig 1), as well as ii) treatments simulating potential future climates in gardens

occurring outside of the range of the species (Tartu, Helsinki, Rauma; Fig 1). We test the effect

of the genotype G (variety) and the environment E (garden or annual temperature) and their

interaction (GxE) on the performance of plant individuals:

performance ~ G + E + G*E

Specifically, we hypothesised that i) the varieties are adapted to local climatic conditions and

thus would show higher performance in their home environments (Fig 1 (b)). Additionally, ii)

both varieties would show lower performance in the out-of-range environments but the southern

variety would have higher performance than the northern one in locations with higher

temperatures outside the species' natural range (Fig 1 (b)). No intraspecific local adaptation

would be revealed by equally high viability of both varieties in both environments (Fig 1 (c))

while no affinity towards conditions within the species range should result in an equally high

performance of the species within and outside of its range (Fig 1 (d)). However, because climate

change has already proceeded (IPCC, 2019), the process of adaptation may be lagging behind

and populations may in fact be maladapted to current climatic condition (Anderson &

Wadgymar, 2020; Kooyers et al., 2019; Wilczek et al., 2014). This scenario of climate change

induced maladaptation (Fig 1 (e)) needs to be taken into account in contemporary translocations

studies. Our study set-up thus allows us to disentangle climatic affinity and on the occurred and

predicted effects of global warming on the Siberian primrose.

Materials and Methods

<u>Study species and seed sampling</u>

Siberian primrose (*Primula nutans*) is a small-statured perennial herb with a discontinuous,

circumpolar distribution. The Fennoscandian subspecies *P. nutans* ssp. *finmarchica* (Jacq.) Á.

Löve & D. Löve is a red-listed species (VU in Norway and Finland) with a disjunct distribution

(Fig 1). It comprises two morpho-ecological varieties (Mäkinen & Mäkinen, 1964): var.

finmarchica (the northern variety) grows on the shores of the Arctic Sea, while var. *jokelae* (the

southern variety) occurs by the Bothnian Bay and the White Sea (Fig. 1 (a)). It is a habitat

specialist that mainly grows in seashore and riverside meadows (Kreivi, Aspi, & Leskinen, 2011;

Mäkinen & Mäkinen, 1964). It propagates sexually by seeds and vegetatively by stolons growing

from the wintering buds and the rosettes (Mäkinen & Mäkinen, 1964). The species is insect-

pollinated and produces copious seeds that spread by gravitation, via water flows, and possibly

also with birds (Ulvinen 1997).

Due to the geographic distance (c. 400-550 km) between its two varieties var. *jokelae* and var.

finmarchica (referred to in this study as the southern and the northern variety, respectively) and

the lack of obvious adaptations for efficient gene flow between the areas, the varieties are

presumably genetically isolated, which is also reflected in their genetic differences (Kreivi et al.,

2011). Moreover, a recent study showed that the main areas where the varieties occur are

climatically different (Hällfors et al., 2016; also see Fig 2 (a)), and an earlier experiment

indicated a clear difference in the requirement of colder night conditions for flower induction in

the northern variety compared to southern variety (Mäkinen & Mäkinen, 1964).

We sampled seeds from wild populations of both varieties of *Primula nutans* ssp. *finmarchica* in

August to September 2012 (Fig. 1 (a), Table S1). Seeds of the southern variety were collected in

five sites in Haukipudas and Ii in Finland, and seeds of the northern variety from six sites in Sør-

Varanger in Norway. Voucher specimens from each sampling site are deposited at the herbarium

of the Finnish Museum of Natural History (H *sensu* Thiers 2016). From each site we collected

seeds from 10 - 53 individuals (as available, following seed collecting guidelines; ENSCONET,

2009; Table S1) to obtain a representative sample. The number of seeds obtained from the

collected capsules of each sampled individual varied from 0 to *c.* 200. We stored the seed lot

from each maternal individual separately and left them to dry and ripen at room temperature for a

minimum of two weeks after which we placed them in a freezer (-18 to -20°C) for cold

stratification to break dormancy. The seeds were kept in these conditions for about six months,

until used to produce plant material for the translocation experiment (in spring 2013).

<u>Experimental design and plant material</u>

We set up a common garden experiment in 2013 in five botanic gardens located in Estonia and

Finland, and on research station grounds in Norway (Fig 1 (a); Table S2). We chose to use

botanic gardens as testing grounds instead of natural sites to avoid genetic contamination of

natural populations by introducing alien genotypes, as well as for logistical reasons and legal

restrictions of introducing species outside their natural range. For a detailed description on the

experimental set-up, see Supplementary Material Text S1.

To describe the performance of individuals, we measured their survival, size, and flowering

(whether they flowered, how many flowers they produced, and when) from year 2014 to 2016.

The fitness of genotypes can be described as the relative success with which they transmit their

genes to the next generation (Silvertown & Charlesworth, 2001). Because of the limited temporal

extent of our study and because we do not measure the individual's fitness through its ability to

produce viable progeny, we focus on key plant performance measures likely to correlate with

fitness. Survival is a definite measure of fitness as a dead individual cannot transmit genes, but

survival can also be stochastic. Also, the physical size of an individual and reproductive output,

such as abundance of flowering, tend to correlate with plant fitness (Silvertown & Charlesworth,

2001). Each spring (2014, 2015, and 2016) we recorded flowering presence and abundance. In

the autumn of the same years, we recorded survival and photographed each surviving plant from

above. We assessed plant size from photographs, through digitally cutting out and measuring the

area (cm^2) of the visible vegetative parts. We used the size at the time of planting as a

measurement of original size, but 24 individuals lack data on that due to missing photographs or

insufficient resolution. These individuals were therefore not included in the analyses, wherefore

our total N is reduced from 614 to 590.

Weather and climatic data

For describing the climatic conditions at the seed sampling sites and experimental gardens, we

obtained data on historic climatic conditions (1970-2000; 10 minutes resolution) and future

projections (CMIP5 for 2050, 10 minutes resolution, HADGEM2-ES model) through the

Worldclim database (Fick & Hijmans, 2017). These climatic data were downloaded in R using

the *getData* function in the *raster* package. Additionally, we describe weather conditions during

the experimental years (Figs 2b and S5-6) and model plant performance as a function of mean

annual temperature in the experimental gardens during 2013-2016 (see below). The weather data

were obtained from the Gridded Agro-Meteorological Data in Europe (Joint Research Centre,

2014), which contains meteorological parameters from European weather stations interpolated

on a 25x25km grid. We used the function *biovar* in the package *dismo* to calculate bioclimatic

variables of the weather data in R.

We present historic and predicted mean annual temperature and mean temperature of the

warmest quarter for the seed sampling sites and experimental gardens and as a mean across

experimental years for each garden in Figure 2. The same variables are also shown explicitly for

each year in each garden in Figure 3. Several additional bioclimatic variables, including annual

precipitation sum, are shown in figures in Fig S5 and S6.

In addition to modeling plant performance as a function of experimental garden and other

variables, we conducted an alternative model using weather data instead of experimental garden.

As the weather parameter we chose annual mean temperature at the experimental gardens across

experimental years. Weather parameters tend to be highly correlated among each other, and we

do not know which specific aspects of temperature these plants respond to. Thus, mean annual

temperature functions as a large scale descriptor of differences between experimental gardens.

The current climatic conditions in Tartu, Helsinki, and Rauma roughly correspond to the

anticipated future climatic conditions in Oulu and thus the occurrence area of the southern

variety (Fig 2). The historic conditions in Oulu roughly correspond to the future conditions in

Svanvik and thus the occurrence area of the northern variety (Fig 2).

The weather conditions during the experimental years differed between the experimental gardens

in accordance with what we expected when designing the experiment (Fig. 2, Fig. S5). However,

the observed temperatures were warmer than the historic means. In essence, this means that the

experienced temperature conditions in Svanvik resembled the historic average temperatures in

Oulu, while the experienced temperatures in Oulu resembled, or were even warmer than, the

historic average temperatures in the out-of-range gardens in Rauma, Helsinki, and Tartu (Fig 2).

The variation in bioclimatic variables between gardens and years are shown in Fig 3 and Fig S6.

Statistical analyses

To test the performance of the two varieties in the different gardens we analysed their combined

performance using aster models (Geyer, Wagenius, & Shaw, 2007; Shaw et al., 2008). They

allow the joint analysis of several components of life history that together quantify differences in

performance, such as survival, flowering and flowering abundance. The complete model (Fig.

S3) included plant survival and flowering in each year (0 or 1; bernoulli error distribution) plus

flowering abundance (zero-truncated poisson distribution of error) that are conditional on

survival in the previous year. Whether a plant flowered or not is also conditional on that it had

survived in the year of flowering. The abundance of flowers per individual is further conditional

on that the individual flowered that year.

These combined responses were modeled using the *aster* call in the aster package (Geyer et al.,

2007; Geyer, 2017) in two main models: one testing the effect of *Garden,* and one testing the

effect of *Temperature* (mean annual temperature over 2013-2016). Specifically, the responses

were modeled as a function of *original size* at planting (continuous, square-root and centered; see

Fig S4 for distribution of original size per variety and experimental garden), *Variety* (categorical:

Southern and Northern) and either *Garden* (categorical, 5 levels) or *Temperature* (continuous)

and the interaction between *Variety* and either *Garden* or *Temperature*. To rule out an effect of

planting time (early or late summer) we also ran a model including this variable, it did not

improve model fit (results not shown). The estimated mean value parameters represent overall

flowering output, that is, the expected flower count during the whole experiment for plant of

average original size (Geyer, 2019). We estimate these for each variety at each experimental

garden and hereafter refer to these predicted values as the overall performance.

As we assumed that random variation between the three *Plots* in each garden and between the

five to six *Seed sampling sites* of each variety, we also fitted random-effect aster models for both

main fixed effects (i.e. for the *Garden* and *Temperature* model) using the *reaster* function in the

aster package (Geyer, 2017). Predicting plant performance based on random effects models is

not straightforward nor directly possible within the aster models package (Geyer et al., 2019;

Charles J. Geyer, Ridley, Latta, Etterson, & Shaw, 2013; although see Charles J. Geyer, Ridley,

Latta, Etterson, & Shaw, 2010). Nevertheless, as the estimates for the fixed effects were very

similar for both the aster models including only fixed effects and the models also including

random effects (see Table S4 for the Garden-models and Table S5 for the Temperature-models)

we obtain predicted values of plant performance based on the fixed effects- models using the

predict function in the *aster* package (Tables S6 and S7). However, for assessing statistical

significance of fixed effects we conducted likelihood ratio tests (anova) of the full random

effects model versus models excluding individual variables (Table 1).

<u>Summarizing results using local adaptation metrics</u>

For the reciprocal part of the trial, we calculated the local adaptation metrics from the mean

estimated combined predicted values obtained through the fixed effects Garden-model (Table

S6). There are three main approaches to make quantitative comparisons of local adaptation from

performance measures (Blanquart et al., 2013). The sympatric-allopatric contrast (henceforth

called *ΔSA*) focuses on the *difference* between the average performance of the genotypes in their

home environment and non-home environment. Thus, it describes how well the genotypes

overall fit their home-environments. The home vs. away approach (henceforth called *HA*) in turn

describes the *home environment quality* for each tested genotype, and is measured by the

difference between the fitness of a genotype in its home environment and all other tested

environments (away). Finally, the local vs. foreign approach (henceforth called *LF*) describes the

genotype quality for each tested home environment, and is measured by the difference between

the fitness of a genotype in its home environment and the mean fitness of all other genotypes

when exposed to the same environment. These three conceptual approaches relate to different

aspects of local adaptation, all of which lie in the interest of our study: the overall habitat quality,

the overall quality of the variety, and how the two varieties fit the different environmental

conditions.

In our study, we estimated plant performance over four years, by predicting mean performance

for an average sized plant of each variety in each experimental garden (based on the GLM

model). We use these estimated measures of performance to calculate local adaptation metrics.

Specifically, *ΔSA* for the *subspecies as whole* is the difference between the mean performance

(*p*) of both varieties (northern, N, southern S) in their home (H) environments and that in the test

(away, A) environment:

$$\Delta SA = \frac{p_{NH} + p_{SH}}{2} - \frac{p_{NA} + p_{SA}}{2}$$

303   The Home vs. Away metric (*HA*) for each *variety* is the difference between the performance of

304   each variety in its home environment and that in its away environment.

$$HA_S = p_{SH} - p_{SA}$$

$$HA_N = p_{NH} - p_{NA}$$

307   The Local vs. Foreign metric (*LF*) for each *garden* is the difference between the performance of

308   the local variety and that of the foreign variety.

$$LF_S = p_{SH} - p_{NA}$$

$$LF_N = p_{NH} - p_{SA}$$

The performance -estimates range between 0.04 and 5.7 and were therefore scaled to range

between 0 and 1. The *HA*, *LF* and *SA* metrics will, therefore, range between -1 and 1, with values

around zero indicating negligible differences in performance. The closer the value is to 1, the

stronger is the evidence for positive response of the local genotype in the environment, or

suitability of the environment for the genotype, and values close to -1 show the opposite.

All data management and analyses were conducted in R (version 3.5.3; R Core Team 2019).

Results

Plant survival

Altogether, 110 out of 590 (18.6%) plant individuals survived in the experiment by 2016. Low

survival is common in transplantation experiments (Dalrymple, Banks, Stewart, & Pullin 2012)

but there were large differences in survival between the varieties and the gardens in our

experiment. The survival percentage across gardens was from 8.8% (27/307) for the northern

variety and 29.3% (83/283) for the southern variety. The survival percentage varied between the

gardens from 5.8% in Tartu (9/155) to 51% in Svanvik (52/102; Fig. 4, Table S3). Individuals of

the southern variety survived at a higher percentage every year and in each garden (Fig 3; Table

S3).

Overall survival was higher for the southern variety (Fig. 4; Table S3) even though the difference

in survival between the varieties varied according to the garden. The northern variety survived

best in Svanvik (representing its home environment), followed by the next garden further south,

Oulu. Its survival was low in the out-of-range gardens (Rauma, Helsinki, and Tartu). The

southern variety showed roughly the same trend, surviving worst in Tartu and best in the

northernmost garden, Svanvik (Fig 4).

Flowering and plant size

Raw data on flower presence, flowering abundance and size (cm^2) are presented in Figure S7

both as averaged across all planted individuals and across all individuals that were alive in the

specific year. Overall, of all the planted individuals, flowering was most frequent and abundant,

and plants were of the biggest sizes in Svanvik. The Southern variety also grew well and

flowered abundantly in Helsinki. When considering only the surviving individuals, however,

both varieties flowered frequently and abundantly in Tartu and Rauma and the surviving

individuals of the southern variety grew big in all southern gardens. However, in many cases the

number of surviving individuals was very low, for example, only one individual of the northern

variety was alive in Tartu in 2016 (Table S3), but it flowered.

<u>Overall performance</u>

Both the Garden and Temperature models revealed similar and significant differences between

the varieties in their overall performance (Table 1; Fig 5). Specifically, the Garden model

indicated increased performance of both varieties in gardens further towards the north but

revealed substantial differences between the varieties (Fig 5a; Table S6). Likewise, the

Temperature model revealed a significant negative effect of mean annual temperature on the

performance on both varieties, however with stronger negative effect on the northern variety (Fig

5b; Table S7). Indeed, we found a significant interaction effects between variety vs. experimental

garden (Fig 5a; Table 1) and annual temperature (Fig 5b; Table 1). In addition, the original size

at planting had a highly significant effect on plant performance in both models (Table 1).

Of the random effects *Plot* explained a significant proportion of the residual variance (change in

deviance when Plot was omitted from the model 40 versus 60 for the Garden and Temperature

models, respectively) while *Seed sampling site* explained a marginal and non-significant

proportion of the residual variation (Table 1; Tables S4 and S5).

The overall performance estimated using aster models provides an estimate of overall

contribution of flowers per variety in each garden while conditional on survival and presence of

flowers. These predicted values (Fig 5., Table S6 and S7), or overall performance, of the

southern variety were highest in Svanvik, followed by Helsinki (however with overlapping

standard deviations of estimates; Fig. 5 and Table 1). The southern variety performed equally

well in Oulu, Rauma, and Tartu. The overall performance of the northern variety followed our

hypothesis (Fig 1a) by showing the highest performance in its home-environment Svanvik,

followed by Oulu, and performed less well in all three out-of-range gardens (Tartu, Helsinki, and

Rauma).

<u>Local adaptation metrics</u>

We calculated local adaptation metrics (Table S8) based on the estimates of overall performance

for the fixed effect Garden model (Table S6) of the two varieties in the reciprocal part of the

experiment. For the southern variety, the *HA*-value of -0.71 indicates that it experiences reduced

habitat quality in its home environment, as it thrives better away (in Svanvik) than at home (in

Oulu). The *LF* of 0.26, however, indicates that the southern variety possesses superior quality in

Oulu since it still does better at home than the tested foreign northern variety does in Oulu. For

the northern variety, the *HA* of 0.22 indicates that it experiences superior habitat quality in its

home environment (Svanvik) compared to in the tested away-environments (Oulu). Nevertheless,

the *LF* of -0.75 indicates that this variety is of reduced quality in its home environment, as the

foreign (southern) variety performs better there. Mean *ΔSA* for the subspecies as a whole was -

0.25 indicating an overall negative fit of the environments for the subspecies.

Discussion

Our results reveal clear patterns of adaptation to macroclimatic conditions at the subspecies level. In general, both varieties show their best performance within the species' range. The southern variety showed higher overall performance in all experimental gardens. The overall survival, size, and flowering frequency and abundance tended to be higher for the southern variety. This is in line with an earlier greenhouse experiment where the northern variety flowered only in cold night conditions (12 h 21°C, 12 h 5°C versus 24h 21°C) whereas the southern variety flowered only in continuously warm conditions but showed increased vegetative growth in the cold night conditions (Mäkinen & Mäkinen, 1964).

We did not find characteristic indications of local adaptation at the variety level when looking at the performance of the sampled populations per experimental garden. Individuals of both varieties survived best, and flowered more frequently and abundantly, in the northernmost garden Svanvik. This is reflected by the local adaptation metrics, where the *LF* metric indicated that the southern variety was superior in the home environment of the northern. Further, the overall performance proxy, the sympatric-allopatric contrast, suggests that the varieties may already be maladapted to their current conditions. In addition, when considering the weather conditions during the experimental years compared to the historic climatic conditions (Fig 2 (b), Fig S5), a conclusion of local adaptation becomes more parsimonious. This is evidenced by, for example, the fact that the mean annual temperature in Svanvik during the experiment resembled that of Oulu's historic mean, and the mean annual temperature in Oulu resembled that of the three southern gardens' historical means. Thus, the conditions that we aimed to mimic through our experimental design had in practise shifted locations one step towards the north, making the

interpretation of the results in relation to our hypotheses perplexing.

Maladaptation is presumably increasing due to climate change, as the rapid change outpaces the

ability of species to adapt or migrate, and conditions reach beyond the extent of temperature

tolerance (Crespi, 2000; Kooyers et al., 2019). Our finding that the varieties did not show the

highest performance in their home area is in line with studies showing signs of maladaptation to

current climatic conditions in plant species (see Anderson & Wadgymar, 2020; Gellie et al.,

2016; Kooyers et al., 2019; Wang, et al., 2019; Wilczek et al., 2014). Although maladaptation

may be increasing because of climate change, it can also be caused by underlying population

genetic structures, such as genetic drift or founder effects (Crespi, 2000; Leimu & Fischer, 2008)

related to the biogeographic history of the populations. The two varieties of Siberian primrose

probably became spatially separated during the end of the last glacial period (Mäkinen &

Mäkinen, 1964), which first allowed colonization of emerging areas and later fragmented the

population through diminishing suitable habitat as land masses reformed from underneath the

receding ice shelves (see Kreivi et al., 2011). Indeed, genetic and allelic diversity is known to be

remarkably low in populations of Siberian primrose (Kreivi et al., 2011). However, the

consistent pattern of increased performance of both varieties from south to north, suggests that

maladaptation caused by recent change in climate is a major driving force for the results of this

study.

Potential confounding effects

While our results suggest a strong impact of climate on the Siberian primrose, they cannot be

interpreted as fully conclusive. Large-scale translocation experiments are complex, and it is

important to evaluate potential methodological pitfalls. Transplantation experiments along the

north-south axis necessarily contain confounding effects, such as day length (Saikkonen et al.,

2012), which cannot be removed in a transplantation study. Despite the rigorous measures to

standardise treatments in all gardens (See Text S1), we had only one experimental garden at each

latitude (albeit with three plots in each experimental garden). Therefore, random variation in

care, pressure from weeds, and weather events may have confounded the climatic signals. For

example, some plots may have experienced short periods of drought despite our regime of

additional watering. The roughly three-year observation period of our study may also not fully

represent the prevailing climates in the trial gardens, since inter-annual variation in weather may

mask mean climatic differences. Long-term studies would even out annual variations, and allow

the measurement of characteristics that become available later in a plant's life cycle and

population development (seed production and germination, adaptation across generations, etc.).

In our study, plants that were larger at the time of planting showed higher overall performance.

Previous studies of reintroductions have shown that the best results (the highest numbers of

surviving plants) are gained by using mature plants or seeds (Dalrymple et al., 2012). However,

in an experimental study, we believe that using mature plants can be confounded since they may

be affected by the climatic conditions of the experimental garden where they are propagated for

the experiment, thus possibly biasing the results through stronger priming effects (see below for

a discussion of the potential effect of Helsinki on the Southern variety). Also, early stages in the

life cycle may be the ones most susceptible to climate change as they define the environments to

which post-germination life-stages of the individuals are exposed (Donohue et al., 2010),

wherefore they should not be omitted from a test setting. On the other hand, while running

experiments from seeds would have allowed us to evaluate performance through germination,

this might have risked the experiment in case of failure to germinate. In fact, even in the

relatively stable greenhouse conditions, the germination percentage of these varieties ranged

between 27-69% (depending on the sampling site; Lehtimäki 2016).

The seeds of each variety were sampled within one region each within a c. 10 km radius (Fig 1,

Table S1). Thus, from the data at hand we cannot infer patterns across the complete ranges of the

varieties. Nevertheless, previous studies have shown that the genetic variation within each

variety is relatively low (Kreivi et al., 2011) and that the varieties occur in distinct climatic

environments (Hällfors et al. 2016). Thus, it is plausible to assume that individuals across the

range of each variety would respond similarly to environmental conditions. We did, however, not

control for, or try to remove, maternal carry-over effects, for instance through growing a first

generation in identical and controlled conditions and then utilizing the progeny of these for the

trials. However, the random effect of seed sampling site did not significantly increase the

explanatory power of residual variation.

Another useful performance measure would have been seed production of the tested individuals

from the experimental locations and germination of these seeds. However, because of limited

resources and the possibility of hybridization between the two varieties, we restricted our

observations to the most readily observable performance measures of the planted populations.

Although germinability of seeds is a more conclusive indicator of reproductive capacity and

population fitness, flowering is also an important indicator of the existence of a necessary life

cycle stage towards offspring. In addition, for this species, flowering has been shown to be an

important fitness-reflecting factor, as failure in flowering is connected to increased extinction

risk of the species (Björnström et al., 2011).

Why the southern variety thrived relatively well in Helsinki remains unclear. The raw proportion

of surviving individuals in Helsinki (31%) was not substantially higher than that in Oulu and

Rauma (24.5 and 23%, respectively). However, the overall performance estimate, which

describes population-level flowering output, was substantially higher in Helsinki (4.1) than in the

variety's home environment in Oulu (1.69) and in Rauma (1.01). We suspect that this could have

been caused by the above-mentioned difference in general maintenance or weather conditions

that we were unable to capture through long-term means. Because the individuals were

propagated together in common conditions in Helsinki, the difference between these conditions

and those of the experimental garden, where the individuals ended up, could additionally explain

the high performance in Helsinki. If the individuals planted in Helsinki were primed to the

Helsinki conditions, this might explain the higher performance of them compared to the other

southern gardens and to Oulu. However, as this higher performance was not seen for the northern

variety (although the varieties may be differentially affected by priming) we suspect that the

main cause was stochastic effects.

<u>The effect of climate change on the Siberian primrose</u>

In this study, we have elucidated the role that plasticity towards varying temperature conditions

can play within the boundaries of local adaptation for the Siberian primrose and its two varieties.

We conclude that, especially for the northern variety, the existing plasticity towards warmer

conditions is negligible. As the climate continues to change, migration northwards could enable

502 the varieties of this species to cope with increasing temperatures. However, because of its poor

503 dispersal ability and limited temperature tolerance, the available options for this species are

504 restricted to the production of characteristics suitable for the new climatic conditions *in situ*,

505 through evolutionary adaptation. Nevertheless, it remains unclear how big a role evolutionary

506 adaptation will play under climatic change in relation to temperature tolerance (Arnold et al.,

507 2019; Charmantier & Gienapp, 2014; Gienapp et al., 2008; Nicotra et al., 2010). It is also not

508 clear whether selection can act to increase or decrease plasticity to allow larger variation among

509 phenotypes, which could then respond adaptively to a range of temperature conditions (Arnold et

510 al., 2019).

512 Our raw data indicate that those individuals that did survive in the southernmost gardens tended

513 to be big and to flower frequently and abundantly (Fig. S7). Thus, the elimination of individuals

514 that are maladapted to the conditions in each experimental garden leaves a survivor population

515 for which the impact of climate is not as decisive. The adaptive performance of the remnant

516 plants in the out-of-range gardens highlights how individual features may be a key determinant

517 for continued survival when a population goes through a bottleneck (Carson, 1990). Whether

518 such a small founding population of a genetically impoverished species could form a viable

519 population under rapidly changing conditions in natural conditions is, however, unlikely.

520 However, experiments on the evolutionary potential in this species are needed to draw more

521 solid conclusions.

523 Siberian primrose, a poor disperser with small and fragmented populations that harbour low

524 genetic variability and a limited thermal tolerance, is an example of a species likely to suffer

from lack of adaptive capacity under rapid climatic change. Proactive conservation methods,

such as assisted migration, may be needed in order to maintain this species. Overall, our results

give cause for concern about the viability of the Siberian primrose, especially for the northern

variety, for which the detrimental effects of climate change may become evident within the

coming decades.

<u>Lessons for translocation trials under altered climate conditions</u>

The results of this study also highlight a critical point in testing for local adaptation and plasticity

under the rapidly changing climate. If populations are not able to track ongoing changes, only

experiments both within and outside the current range of the species (or variety) can provide

accurate information regarding local adaptation and plasticity, as maladaptation must be

considered as a scenario (e.g., Anderson & Wadgymar, 2020; Gellie et al., 2016; Kooyers et al.,

2019; Wilczek et al., 2014). Climate change thus gives cause for re-examining traditional

reciprocal and out-of-range common garden experiments (Anderson & Wadgymar, 2020), and to

plan experiments with climatic and weather data at hand. Comparing responses to current versus

"historical" conditions offers a powerful approach (Anderson & Wadgymar, 2020; Franks et al.,

2007; Merilä & Hendry, 2014; Thomann et al., 2015). Together with continued monitoring of

population abundances (Cotto et al., 2017) such studies can provide early warning signals for

species that may be close to tipping points and to going extinct.

Acknowledgements

We are grateful to the botanic gardens that provided staff and premises for conducting the trials:

University of Tartu, University of Helsinki, University of Turku teaching garden in Rauma,

University of Oulu, and NIBIO Svanhovd. We thank Pertti Pehkonen, Outi Pakkanen, and

Toomas Kangro for technical assistance in preparing the experiments. Terhi Ryttäri and Henry

Väre gave advice on the species and required soil properties. Ritva Hiltunen and Tuomas

Kauppila assisted in collection of seeds, Elina Vaara helped with permits for seed and plant

transfers, Hanna Finne with data management, and Bess Hardwick with acquiring weather data.

MHH was supported by the University of Helsinki Research Fund, LUOVA – Doctoral

Programme in Wildlife Biology Research and the Jane and Aatos Erkko Foundation through the

Research Centre for Ecological Change, University of Helsinki. The study was supported by the

Academy of Finland grant 126915 and Societas pro Fauna et Flora Fennica. The authors declare

no conflicts of interest.

Author's contributions

MHH, SL, and MH conceived the idea and designed methodology, LS contributed to the

selection of experimental gardens; MHH and LS collected seeds; MHH, SL, IL, and MH

produced plant material and coordinated the experiments; all authors took part in setting up the

experiments and collecting the data; MHH analyzed the data and led the writing of the

manuscript. All authors contributed critically to the manuscript drafts and gave final approval for

publication.

Data accessibility

The data are available through GitHub at https://github.com/MariaHallfors/Primula-nutans-

translocation

Bibliography

Anderson, J. T., & Wadgymar, S. M. (2020). Climate change disrupts local adaptation and

favours upslope migration. *Ecology Letters*, *23*(1), 181–192. doi: 10.1111/ele.13427

Arnold, P. A., Nicotra, A. B., & Kruuk, L. E. B. (2019). Sparse evidence for selection on

phenotypic plasticity in response to temperature. *Philosophical Transactions of the Royal

Society B: Biological Sciences*, *374*(1768), 20180185. doi: 10.1098/rstb.2018.0185

Atkins, K. E., & Travis, J. M. J. (2010). Local adaptation and the evolution of species' ranges

under climate change. *Journal of Theoretical Biology*, *266*(3), 449–457. doi:

10.1016/j.jtbi.2010.07.014

Banta, J. A., Ehrenreich, I. M., Gerard, S., Chou, L., Wilczek, A., Schmitt, J., … Purugganan, M.

D. (2012). Climate envelope modelling reveals intraspecific relationships among

flowering phenology, niche breadth and potential range size in Arabidopsis thaliana.

Ecology Letters, *15*(8), 769–777. doi: 10.1111/j.1461-0248.2012.01796.x

Björnström, T., Hyvärinen, M., Siikamäk, P., Strengell, H., Väre, H., & Tuomi, J. (2011).

Demographic variability and population viability of an endemic variety of the Siberian

primrose. *Annales Botanici Fennici*, *48*(2), 107–119. Retrieved from JSTOR.

Blanquart, F., Kaltz, O., Nuismer, S. L., & Gandon, S. (2013). A practical guide to measuring

local adaptation. *Ecology Letters*, *16*(9), 1195–1205. doi: 10.1111/ele.12150

Carson, H. L. (1990). Increased genetic variance after a population bottleneck. *Trends in Ecology

& Evolution*, *5*(7), 228–230. doi: 10.1016/0169-5347(90)90137-3

Ceballos, G., Ehrlich, P. R., Barnosky, A. D., García, A., Pringle, R. M., & Palmer, T. M.

(2015). Accelerated modern human–induced species losses: Entering the sixth mass

extinction. *Science Advances*, *1*(5), e1400253. doi: 10.1126/sciadv.1400253

Charmantier, A., & Gienapp, P. (2014). Climate change and timing of avian breeding and

migration: evolutionary versus plastic changes. *Evolutionary Applications*, *7*(1), 15–28.

doi: 10.1111/eva.12126

Chevin, L.-M., Lande, R., & Mace, G. M. (2010). Adaptation, Plasticity, and Extinction in a

Changing Environment: Towards a Predictive Theory. *PLOS Biology*, *8*(4), e1000357.

doi: 10.1371/journal.pbio.1000357

Cotto, O., Wessely, J., Georges, D., Klonner, G., Schmid, M., Dullinger, S., … Guillaume, F.

(2017). A dynamic eco-evolutionary model predicts slow response of alpine plants to

climate warming. *Nature Communications*, *8*(1), 15399. doi: 10.1038/ncomms15399

Crespi, B. J. (2000). The evolution of maladaptation. *Heredity*, *84*(6), 623–629. doi:

10.1046/j.1365-2540.2000.00746.x

Dalrymple, S.E., Banks, E., Stewart, G.B., & Pullin, A.S. (2012) A Meta-Analysis of Threatened

Plant Reintroductions from Across the Globe. In: Machinski and Haskins (Eds.), *Plant*

Reintroduction in a Changing Climate. Promises and Perils (pp. 31- 50). Island Press.

Dawson, I. K., Vinceti, B., Weber, J. C., Neufeldt, H., Russell, J., Lengkeek, A. G., …

Jamnadass, R. (2011). Climate change and tree genetic resource management:

maintaining and enhancing the productivity and value of smallholder tropical

agroforestry landscapes. A review. *Agroforestry Systems*, *81*(1), 67–78. doi:

10.1007/s10457-010-9302-2

Díaz, S., Settele, J., Brondízio, E., Ngo, H. T., Guèze, M., Agard, J., … Vilá, B. (2019).

Summary for policymakers of the global assessment report on biodiversity and ecosystem

services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem

Services. 44.

Donohue, K., Rubio de Casas, R., Burghardt, L., Kovach, K., & Willis, C. G. (2010).

Germination, Postgermination Adaptation, and Species Ecological Ranges. *Annual*

Review of Ecology, Evolution, and Systematics, *41*(1), 293–319. doi: 10.1146/annurev-

ecolsys-102209-144715

ENSCONET 2009: ENSCONET Seed Collecting Manual for Wild Species. Available at:

http://ensconet.maich.gr/PDF/Collecting_protocol_English.pdf

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces

for global land areas. *International Journal of Climatology*, *37*(12), 4302–4315. doi:

10.1002/joc.5086

Frankham, R., Ballou, J. D., Dudash, M. R., Eldridge, M. D. B., Fenster, C. B., Lacy, R. C., …

Ryder, O. A. (2012). Implications of different species concepts for conserving

biodiversity. *Biological Conservation*, *153*, 25–31. doi: 10.1016/j.biocon.2012.04.034

Franks, S. J., Sim, S., & Weis, A. E. (2007). Rapid evolution of flowering time by an annual

plant in response to a climate fluctuation. *Proceedings of the National Academy of*

Sciences, *104*(4), 1278–1282. doi: 10.1073/pnas.0608379104

Frenne De, P., Graae, B. J., Rodríguez☐Sánchez, F., Kolb, A., Chabrerie, O., Decocq, G., …

Verheyen, K. (2013). Latitudinal gradients as natural laboratories to infer species'

responses to temperature. *Journal of Ecology*, *101*(3), 784–795. doi: 10.1111/1365-

2745.12074

Gao, S., Mo, L., Zhang, L., Zhang, J., Wu, J., Wang, J., … Gao, Y. (2018). Phenotypic plasticity

638   vs. local adaptation in quantitative traits differences of Stipa grandis in semi-arid steppe,

639   China. *Scientific Reports*, *8*(1), 3148. doi: 10.1038/s41598-018-21557-w

640   GBIF (2013). *Global Biodiversity Information Facility Data Portal.*

641   Gellie, N. J. C., Breed, M. F., Thurgate, N., Kennedy, S. A., & Lowe, A. J. (2016). Local

642   maladaptation in a foundation tree species: Implications for restoration. *Biological*

643   *Conservation*, *203*, 226–232. doi: 10.1016/j.biocon.2016.08.036

644   Geyer, C. J., Wagenius, S., & Shaw, R. G. (2007). Aster models for life history analysis.

645   *Biometrika*, *94*(2), 415–426. doi: 10.1093/biomet/asm030

646   Geyer, Charles J., Ridley, C. E., Latta, R. G., Etterson, J. R., & Shaw, R. G. (2010). *Aster*

647   *Models with Random Effects via Penalized Likelihood* [Technical Report]. Retrieved

648   from University of Minnesota website: http://conservancy.umn.edu/handle/11299/199694

649   Geyer, Charles J., Ridley, C. E., Latta, R. G., Etterson, J. R., & Shaw, R. G. (2013). LOCAL

650   ADAPTATION AND GENETIC EFFECTS ON FITNESS: CALCULATIONS FOR

651   EXPONENTIAL FAMILY MODELS WITH RANDOM EFFECTS. *The Annals of*

652   *Applied Statistics*, *7*(3), 1778–1795.

653   Geyer, C.J., (2017). R package aster2 (Aster Models), version 0.3. https://cran.r-

654   project.org/package=aster2.

655   Geyer, C.J., (2019): The Aster Package Tutorial. Available at: https://cran.r-

656   project.org/web/packages/aster/vignettes/tutor.pdf

658   Gienapp, P., Teplitsky, C., Alho, J. S., Mills, J. A., & Merilä, J. (2008). Climate change and

659   evolution: disentangling environmental and genetic responses. *Molecular Ecology*, *17*(1),

660   167–178. doi: 10.1111/j.1365-294X.2007.03413.x

Greiser, C., Hylander, K., Meineri, E., Luoto, M., & Ehrlén, J. (2020). Climate limitation at the

cold edge: contrasting perspectives from species distribution modelling and a transplant

experiment. *Ecography*, *n/a*(n/a). doi: 10.1111/ecog.04490

Hällfors, M. H., Vaara, E. M., Hyvärinen, M., Oksanen, M., Schulman, L. E., Siipi, H., &

Lehvävirta, S. (2014). Coming to Terms with the Concept of Moving Species Threatened

by Climate Change – A Systematic Review of the Terminology and Definitions. *PLOS*

ONE, *9*(7), e102979. doi: 10.1371/journal.pone.0102979

Hällfors, M. H., Liao, J., Dzurisin, J., Grundel, R., Hyvärinen, M., Towle, K., Wu., G.,

Hellmann, J. J. (2016). Addressing potential local adaptation in species distribution

models: implications for conservation under climate change. *Ecological Applications*,

26(4), 1154–1169. doi: 10.1890/15-0926

Hargreaves, A. L., Samis, K. E., & Eckert, C. G. (2014). Are Species' Range Limits Simply

Niche Limits Writ Large? A Review of Transplant Experiments beyond the Range. *The*

American Naturalist, *183*(2), 157–173. doi: 10.1086/674525

Hill, J. K., Griffiths, H. M., & Thomas, C. D. (2011). Climate Change and Evolutionary

Adaptations at Species' Range Margins. *Annual Review of Entomology*, *56*(1), 143–159.

doi: 10.1146/annurev-ento-120709-144746

Hultén, E., & Fries, M. (1986). *Atlas of North European vascular plants: north of the Tropic of*

Cancer, vol. 1-3. Königstein, Koeltz.

IPCC, 2019: Summary for Policymakers. In: *Climate Change and Land: an IPCC special report*

on climate change, desertification, land degradation, sustainable land management, food

security, and greenhouse gas fluxes in terrestrial ecosystems [P.R. Shukla, J. Skea, E.

Calvo Buendia, V. Masson-Delmotte, H.- O. Pörtner, D. C. Roberts, P. Zhai, R. Slade, S.

Connors, R. van Diemen, M. Ferrat, E. Haughey, S. Luz, S. Neogi, M. Pathak, J. Petzold,

J. Portugal Pereira, P. Vyas, E. Huntley, K. Kissick, M. Belkacemi, J. Malley, (eds.)].

Joint Research Centre 2014: *Gridded Agro-Meteorological Data in Europe.* URI:

http://data.europa.eu/88u/dataset/jrc-marsop4-7-weather_obs_grid_2015

Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology Letters*, *7*(12),

1225–1241. doi: 10.1111/j.1461-0248.2004.00684.x

Kooyers, N. J., Colicchio, J. M., Greenlee, A. B., Patterson, E., Handloser, N. T., & Blackman,

B. K. (2019). Lagging Adaptation to Climate Supersedes Local Adaptation to Herbivory

in an Annual Monkeyflower. *The American Naturalist*, *194*(4), 541–557. doi:

10.1086/702312

Kreivi, M., Aspi, J., & Leskinen, E. (2011). Regional and local spatial genetic structure of

Siberian primrose populations in Northern Europe. *Conservation Genetics*, *12*(6), 1551–

1563. doi: 10.1007/s10592-011-0252-7

Kreivi, M., Huttunen, S., & Aspi, J. (2006). Isolation and characterization of polymorphic

microsatellite markers from Primula nutans (Primulaceae). *Molecular Ecology Notes*,

6(2), 334–336. doi: 10.1111/j.1471-8286.2005.01222.x

Kreyling, J., Buhk, C., Backhaus, S., Hallinger, M., Huber, G., Huber, L., … Beierkuhnlein, C.

(2014). Local adaptations to frost in marginal and central populations of the dominant

forest tree Fagus sylvatica L. as affected by temperature and extreme drought in common

garden experiments. Ecology and Evolution, 4(5), 594–605. doi: 10.1002/ece3.971

Lampinen, R., Lahti, T., Heikkinen, M. 2012. *Kasviatlas 2011*. -- Helsingin Yliopisto,

Luonnontieteellinen keskusmuseo, Helsinki. Distribution maps available at:

http://www.luomus.fi/kasviatlas.

Lehtimäki, I. (2016). *Uhanalaisen ruijanesikon suojelu. Etäsuojelun ja avustetun leviämisen*

näkökulma. Master's thesis [In Finnish]. Department of Biosciences, University of

Helsinki.

Leimu, R., & Fischer, M. (2008). A Meta-Analysis of Local Adaptation in Plants. PLoS ONE,

3(12), e4010. doi: 10.1371/journal.pone.0004010

Lenoir, J., & Svenning, J.-C. (2015). Climate-related range shifts – a global multidimensional

synthesis and new research directions. *Ecography*, *38*(1), 15–28. doi:

10.1111/ecog.00967

Mäkinen, L., & Mäkinen, Y. (1964). The distribution, ecology, morphology and taxonomy of

Primula nutans Georgi ssp. finmarchica (Jacq.) Löve & Löve. *Annales Botanici Fennici*,

1(4), 273–291. Retrieved from JSTOR.

Merilä, J. (2012). Evolution in response to climate change: In pursuit of the missing evidence.

BioEssays, *34*(9), 811–818. doi: 10.1002/bies.201200054

Merilä, J., & Hendry, A. P. (2014). Climate change, adaptation, and phenotypic plasticity: the

problem and the evidence. *Evolutionary Applications*, *7*(1), 1–14. doi: 10.1111/eva.12137

Nicotra, A. B., Atkin, O. K., Bonser, S. P., Davidson, A. M., Finnegan, E. J., Mathesius, U., …

van Kleunen, M. (2010). Plant phenotypic plasticity in a changing climate. *Trends in*

Plant Science, *15*(12), 684–692. doi: 10.1016/j.tplants.2010.09.008

Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts

across natural systems. *Nature*, *421*(6918), 37–42. doi: 10.1038/nature01286

Pelini, S. L., Diamond, S. E., MacLean, H., Ellison, A. M., Gotelli, N. J., Sanders, N. J., & Dunn,

R. R. (2012). Common garden experiments reveal uncommon responses across

temperatures, locations, and species of ants. *Ecology and Evolution*, *2*(12), 3009–3015.

doi: 10.1002/ece3.407

Pöyry, J., Luoto, M., Heikkinen, R. K., Kuussaari, M., & Saarinen, K. (2009). Species traits

explain recent range shifts of Finnish butterflies. *Global Change Biology*, *15*(3), 732–

743. doi: 10.1111/j.1365-2486.2008.01789.x

R Core Team 2019: R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.

Saikkonen, K., Taulavuori, K., Hyvönen, T., Gundel, P. E., Hamilton, C. E., Vänninen, I., …

Helander, M. (2012). Climate change-driven species' range shifts filtered by

photoperiodism. *Nature Climate Change*, *2*(4), 239–242. doi: 10.1038/nclimate1430

Settele, J., Scholes, R., Betts, R., Bunn, S., Leadley, P., Nepstadt, D., & Taboada, M. A. (2014).

Terrestrial and inland water systems. In: Cambridge, United Kingdom and New York,

NY, USA, pp. 271-359.

Shaw, R. G., Geyer, C. J., Wagenius, S., Hangelbroek, H. H., Etterson, J. R., Taylor, A. E. P. D.,

& DeAngelis, E. D. L. (2008). Unifying Life☐History Analyses for Inference of Fitness

and Population Growth. *The American Naturalist*, *172*(1), E35–E47. doi: 10.1086/588063

Silvertown, J., & Charlesworth, D. (2001). *Introduction to Plant Population Biology*. 4th edition.

Blackwell Science.

Thiers, B. (2016) Index Herbariorum: A Global Directory of Public Herbaria and Associated

Staff. New York Botanical Garden's Virtual Herbarium.

http://sweetgum.nybg.org/science/ih/

Thomann, M., Imbert, E., Engstrand, R. C., & Cheptou, P.-O. (2015). Contemporary evolution of

plant reproductive strategies under global change is revealed by stored seeds. *Journal of*

Evolutionary Biology, *28*(4), 766–778. doi: 10.1111/jeb.12603

Ulvinen, T. (1997). Rujanesikko. In: *Uhanalaiset kasvimme*. Ryttäri T. and Kettunen T. (eds.).

[In Finnish]. Suomen ympäristökeskus, Helsinki.

Urban, M. C. (2015). Accelerating extinction risk from climate change. *Science*, *348*(6234),

571–573. doi: 10.1126/science.aaa4984

Valladares, F., Matesanz, S., Guilhaumon, F., Araújo, M. B., Balaguer, L., Benito-Garzón, M.,

… Zavala, M. A. (2014). The effects of phenotypic plasticity and local adaptation on

forecasts of species range shifts under climate change. *Ecology Letters*, *17*(11), 1351–

1364. doi: 10.1111/ele.12348

Wang, Y., Pedersen, J. L. M., Macdonald, S. E., Nielsen, S. E., & Zhang, J. (2019).

Experimental test of assisted migration for conservation of locally range-restricted plants

in Alberta, Canada. *Global Ecology and Conservation*, *17*, e00572. doi:

10.1016/j.gecco.2019.e00572

Wilczek, A. M., Cooper, M. D., Korves, T. M., & Schmitt, J. (2014). Lagging adaptation to

warming climate in Arabidopsis thaliana. *Proceedings of the National Academy of*

Sciences, *111*(22), 7906–7913. doi: 10.1073/pnas.1406314111

Zenni, R. D., Bailey, J. K., & Simberloff, D. (2014). Rapid evolution and range expansion of an

invasive plant are driven by provenance -environment interactions. *Ecology Letters*,

17(6), 727–735. doi: 10.1111/ele.12278

Figure legends and tables

FIGURE 1. Geographical distribution of seed sampling sites and experimental gardens (a), and

hypotheses of plant performance in experimental gardens (b-e). Panel (a) shows the geographical

distribution of seed sampling sites and experimental gardens with occurrences of *Primula nutans* ssp.

finmarchica marked by dark grey points: Var. *finmarchica* occurs by the Arctic Sea in N-Norway and var.

jokelae by the Bothnian bay in Finland and Sweden and the White Sea in Russia. Red = seed sampling

sites of the southern variety (var. *jokelae*) in Finland; Blue= seed sampling sites of the northern variety

(var. *finmarchica*) in Norway. Source for occurrences of Siberian primrose: Global Biodiversity

Information Facility (GBIF 2013); Kastikka (Finnish plant distribution database; Lampinen, Lahti, &

Heikkinen 2012); Hertta (database of Finnish Environment Institute, unpublished data); occurrences in

Russia based on information from herbarium specimens (from collections in the herbarium of the Finnish

Museum of Natural History [H *sensu* Thiers 2016] and the herbarium of the University of Turku [TUR

sensu Thiers 2016] and the distribution map by Hultén and Fries [1986]). Panels (b-e) show hypothesized

overall performance of the tested varieties in all experimental gardens following opposing underlying

scenarios of (b) local adaptation at the varietal and subspecies level, (c) local adaptation at the subspecies

level, (d) tolerance (through plasticity) towards all tested conditions (including those not currently present

within the occurrence area of the subspecies), and (e) maladaptation caused by climate change (see text

for hypotheses). Dashed area demarks within-range gardens, i.e. the reciprocal part of the experiment.

Red= southern variety; blue = northern variety.

FIGURE 2. Climatic and weather conditions of seed sampling sites and experimental gardens. (a) Mean

annual temperature and (b) mean temperature of warmest quarter (°C; mean temperature for the three

months during which temperatures were highest in each year). Black points show historic mean

temperatures (1970-2000; 10 minutes resolution; (Fick & Hijmans, 2017) and gray points show the future

projections of mean temperatures (CMIP5 for 2050, 10 minutes resolution, HADGEM2-ES model; (Fick

& Hijmans, 2017) for each experimental gardens and seed sampling sites (Northern= seed sampling sites

of the northern variety; Southern= seed sampling sites of the southern variety). For experimental gardens,

open circles show mean temperature during the experimental years 2013-2016. The experimental gardens

and seed sampling sites within the species range is outlined by a dashed gray box, whereas the blue box

indicates the sites within the range of the northern variety and the red box the sites within the range of the

southern variety. Corresponding figures for precipitation sum and other climatic variables in Fig S5. The

future temperatures in Oulu approach the historic ones in Tartu, Helsinki and Rauma, and the future

temperatures in Svanvik approach the historic ones in Oulu. During the experiment, mean temperatures

were higher than the historic means. The conditions in Svanvik corresponded with historical means in

Oulu. Those in Oulu approached the historic means in the southern gardens.

FIGURE 3. Temperature conditions in the experimental gardens over the course of the experiment

(2013-2016). (a) Mean annual temperature and (b) mean temperature of the warmest quarter (mean

temperature for the three months during which temperatures were highest in each year) per garden and

year.

FIGURE. 4. Mean survival per variety across gardens and years. The total number of individuals planted

in 2013 appear at the top of the plot (both original plants and additional plants due to high die-off in the

first summer). Red points = southern variety; blue points = northern variety. The size of the point

indicates year: big point= 2014; medium point= 2015; small point = 2016. Exact numbers of individuals

per year and garden and percentage decrease over year are presented in Table S3.

FIGURE 5. Predicted overall performance of the two varieties across (a) experimental gardens, and (b)

mean annual temperature over the experimental period 2013-2016. Predictions are made for plants of

average original size and based on the fixed effects aster model (i.e., not including *Plot* and *Seed*

sampling site as random effects). Red= southern variety; blue= northern variety.

TABLE 1. Change in deviance when a single variable was omitted from the full random effects models.

Summary from comparing aster models testing the fixed effects of experimental garden (left-hand side of

table) or annual temperature (right hand side of table), variety and original size on plant performances, as

well as *Plot* and *Seed sampling site* as random effects. Statistical significance of predictor variables was

assessed using likelihood ratio tests, that is, a model where one variable was omitted was compared to the

full model using anova.

Garden-model				Annual temperature -model		
p-value	df	Change in deviance	Variable	Change in deviance	df	p-value
<0.001	1	62.4	Original size	60.2	1	<0.001
<0.05	4	11	Variety:Garden/Ann.Temp.	8.8	1	<0.01
<0.01	5	20	Variety (+ Variety:Garden/Ann. Temp.)	17.2	2	<0.001
<0.05	8	19.6	Garden/Ann. Temp. (+ Variety:Garden/Ann. Temp.)	14.2	2	<0.001
<0.001	0	46	Plot	60	0	<0.001
0.09	0	1.8	Seed sampling site	2.2	0	0.07

(a)

Mean annual temperature

(b)

Mean temperature of warmest quarter