An embryonic stem cell–specific heterochromatin state allows core histone exchange in the absence of DNA accessibility

Carmen Navarro and Simon J Elsässer

Affiliations

Science for Life Laboratory, Department of Medical Biochemistry and Biophysics, Karolinska Institutet, Stockholm, Sweden

Ming Wai Lau Centre for Reparative Medicine, Stockholm node, Karolinska Institutet, Stockholm, Sweden

Corresponding Author

Simon J Elsässer, simon.elsasser@scilifelab.se

ABSTRACT

Nucleosome turnover concomitant with incorporation of the replication-independent histone variant H3.3 is a hallmark of regulatory regions in the animal genome. In our current understanding, nucleosome turnover is universally linked to DNA accessibility and histone acetylation. In mouse embryonic stem cells, H3.3 is also highly enriched at interstitial heterochromatin, most prominently intracisternal-A particle endogenous retroviral elements. Interstitial heterochromatin is established over confined domains by the TRIM28/SETDB1 corepressor complex and has stereotypical features of repressive chromatin, such as H3K9me3 and recruitment of all HP1 isoforms. Here, we demonstrate that fast histone turnover and H3.3

incorporation is compatible with these hallmarks of heterochromatin. Further, we find that histone H3.3 is required to maintain minimal DNA accessibility in this surprisingly dynamic heterochromatin state. Loss of H3.3 in mouse embryonic stem cells elicits a highly specific opening of interstitial heterochromatin with minimal effects on other silent or active regions of the genome.

INTRODUCTION

Histone H3.3 is an evolutionary conserved variant of the canonical histone H3 proteins, termed H3.1 and H3.2 in mammalian cells [1]. First described to associate with regions of active transcription in *Drosophila* [2], H3.3 has been recognized as the sole substrate for replication-independent chromatin assembly in mammalian cells [3]. While H3.1/2 expression and replication-dependent incorporation is limited to S phase, H3.3 is expressed through the cell cycle and incorporated at sites of dynamic nucleosome turnover. A multitude of genome-wide studies have identified such dynamic chromatin at promoters, enhancers, gene bodies and origins of replication [4–13]. In these instances, nucleosome turnover is thought to be a consequence of chromatin remodeling, transcriptional activity, nucleosome-destabilizing DNA sequences, competition with other chromatin binding factors, or a combination thereof. While histone H3.3 itself has been proposed to destabilize the nucleosome, it is unclear if H3.3 has a causal role in promoting nucleosome turnover [14–16]. Histone H3.3 has been shown to be enriched in histone posttranslational modifications (PTMs) characteristic of euchromatin, i.e. H3K4me3, H3K9ac, H3K27ac [17,18]. Accumulation of active PTMs on histone H3.3 is sufficiently explained by their co-occurrence at sites of active enhancers or transcription, and H3.3 appears not to be required for maintaining those active PTMs. However, a notable exception represents

phosphorylation of Ser31, a H3.3-specific residue. H3.3Ser31 phosphorylation facilitates rapid activation of genes [19,20], thus providing a mechanistic link between H3.3 incorporation and gene activation. In summary, histone H3.3 is a well-established component of euchromatin, acting as a replacement histone for a range of dynamic processes. While H3.3 has to be considered predominantly neutral to the underlying dynamic process, it can – as exemplified in the specific instance above – partake mechanistically in establishing open chromatin.

In mouse embryonic stem cells (ESC), a considerable fraction of H3.3-enriched regions do not fall into the active regions outlined above, but colocalizes with histone H3 Lys 9 trimethylation (H3K9me3) modification at interstitial heterochromatin [21]. Re-ChIP experiments suggested that H3.3 and H3K9me3 coincide in the same nucleosome. Interstitial heterochromatin in mouse ESC spreads over relatively small genomic distance (~10kb), is established by a Tripartite motif-containing 28 (TRIM28) co-represser/SETDB1 histone methyltransferase complex and includes a subset of endogenous retroviral elements (ERVs), most prominently Intracisternal A-type particle (IAP), and imprinted genes [22–27]. Interstitial heterochromatin is highly CpG-methylated, enriched in linker histone and bound by HP1 family proteins, the stereotypical H3K9me3 readers [28–30]. HP1 proteins have been shown to bridge nucleosomes and compact chromatin into phase-separated liquid condensates [31,32]. HP1 and linker histone render DNA in an inaccessible state [32], and underlying genes/repeats are refractory to activating factors [33].

Enrichment of H3.3 at H3K9me3 heterochromatin regions, so far uniquely observed in mouse ESC, poses a conundrum: do these regions show fast nucleosome turnover as known for euchromatic H3.3-enriched regions? Does the presence of H3.3 coincide with DNA accessibility, and how would such dynamic properties be reconciled with a silent, condensed

heterochromatin structure? Here, we follow up on these questions analyzing a set of complementary genome-wide datasets from published studies, together revealing surprisingly dynamic properties of interstitial heterochromatin in mESC. We show that the hallmarks of heterochromatin are compatible with rapid exchange of core histones while maintaining DNA in an inaccessible state. Further, a supply of the replacement histone H3.3 is necessary to maintain the inaccessible state.

RESULTS

Several studies have assessed genome-wide H3.3 dynamics in mESC, either measuring incorporation or turnover of ectopically tagged H3.3 [5,8,9,34]. A recent study assessed nucleosome turnover using a method termed 'time-ChIP' in mouse embryonic stem cells as well as neural stem cells (NSC) [9]. In this method, SNAP-tagged H3.3 is expressed from a Tet-controlled transgene and biotin-labeled using the SNAP-tag at the onset of a time course (0h). The biotin-labeled fraction then is observed over a time range of 3-12 hours (Figure 1a, b). Dynamic regions are marked by H3.3 at 0h and gradually decay towards the genome-wide average, implying replication-independent turnover of H3.3 [9].

H3.3-enriched regions in ESC called from the initial time point (0h) overlapped well with those determined for endogenous histone H3.3, including those peaks sharing H3.3 and H3K9me3 (H3.3+H3K9me3) (Figure 1c,d). As an example, H3.3 enrichment tracked well with known ESC enhancers and H3K27ac marks, but to similar levels with a nearby IAP ERV (Figure 1b, Extended Data Figure 1a). NSC showed globally similar turnover kinetics (Figure 1a, Extended Data Figure 1b, c) and maintained many of the H3.3 peaks defined in ESC (Figure 1c,d). However, among those H3.3 peaks specifically lost in NSC were essentially all H3.3+H3K9me3

peaks (Figure 1c,d), as exemplified by an IAP ERV localized in the Hist1 histone gene cluster (Extended Data Figure 1b). Thus, as previously concluded [21], co-enrichment of H3.3 and H3K9me3 is a feature of pluripotent stem cells which appears to be lost early upon differentiation.

Interestingly, H3K9me3 and H3.3 has also been studied genome-wide during reprogramming of mouse fibroblasts to induced-pluripotent cells [35]; while H3K9me3 becomes enriched over IAP ERV in pre-iPSC, H3.3 is absent at this stage (Extended Data Figure 1c). Pre-iPSC are stable late-stage reprogramming intermediates that can be converted to the pluripotent state through MEK inhibition. Thus, these data corroborate a link between pluripotency and the acquisition of H3.3 at interstitial heterochromatin, where H3K9me3 has already been established during reprogramming.

Annotating H3.3 peaks with a 15-state ChromatinHMM model generated with seven histone modifications and RNA Polymerase II (RNAP2) profiles [36,37], we found that H3.3-only peaks overlapped largely with active transcription (states 1-3) and enhancers (states 4-9) (Figure 1d, Extended Data Figure 2a). H3.3+H3K9me3 peaks were assigned to heterochromatin (states 12-14) (Figure 1d, Extended Data Figure 2a), defined by the absence of H3K4me3, H3K4me1, H3K27ac, H3K36me3 and RNAP2 [37], in line with the fact that H3K9me3 appears mutually exclusive with these signatures of active chromatin.

Binning the genome into 5kb windows, we found a bimodal H3.3 distribution, with regions of very high and very low levels of H3K9me3 coinciding with the highest H3.3 enrichment (Figure 1e). Overlaying the 5kb bins with repeat annotation, we confirmed that the strongest H3.3+H3K9me3 co-enrichment was linked to the presence of an IAP element (Figure 1e,

Extended Data Figure 3). Further, ETn and MusD elements also defined a subset of H3.3+H3K9me3 co-enriched regions (Extended Data Figure 3). It is known that a small fraction of ETn/MusD elements are active and, in fact, highly transcribed. This is evident from elongating RNA Polymerase II running into the 3' flanking region of few elements, and the complete absence of H3K9me3 around these elements (Extended Data Figure 4a). Since we cannot unambiguously attribute ChIP-Seq reads within the highly conserved ETn/MusD internal sequence to specific instances, we are not able to discern how much H3.3 incorporation can be attributed to the small fraction of active versus majority of repressed copies. Therefore, we did not further focus on ETn/MusD elements. Amongst the IAP repeat family, none of the annotated instances showed similar evidence of transcriptional activity (Extended Data Figure 4b). Consistent with this, the transcript level of IAP is extremely low, roughly 100fold lower than ETn (Extended Data Figure 4c).

Using the time-ChIP data, we next assessed the turnover of labeled histone H3.3 at H3.3-only and H3.3+H3K9me3 peaks. Decay of H3.3 towards background levels over the 12h chase period was observed at all peaks irrespective of their H3K9me3 status (Figure 2a). Turnover at H3.3+H3K9me3 enriched regions was also observed in another Tet-OFF dataset [5] (Extended Data Figure 5). A different dataset, applying a pulse of tagged histone H3.3 [34], shows that newly synthesized histone H3.3 is incorporated rapidly at both H3.3-only and H3.3+H3K9me3 peaks (Figure 2b, Extended Data Figure 1a, 5). An orthogonal method to measure nucleosomal H3/H4 incorporation and turnover, CATCH-IT, relies on metabolic labeling of newly synthesized proteins with Azido-homoalanine (AHA) [38]. CATCH-IT previously performed in mouse ESC [6] confirms the dynamic exchange of core histones at both H3.3-only and H3.3+H3K9me3 peaks (Figure 2c). Crucial for this method, all non-histone proteins and H2A/H2B dimers are stripped

of chromatin biochemically before performing the ChIP, revealing the incorporation dynamics of newly synthesized H3/H4 units irrespective of the H3 variant.

Thus, three independent datasets provide complementary evidence for nucleosome turnover at interstitial heterochromatin. While rapid H3.3 turnover has been attributed to 'hyperdynamic' promoter and enhancer nucleosomes [8,9], our analysis suggests that similar dynamics also apply to H3.3+H3K9me3 peaks.

We next compared H3.3 dynamics across 15 ChromHMM states [37] and H3.3+H3K9me3 heterochromatic regions and found that highest histone H3.3 turnover is observed at active promoters and enhancers followed by H3.3+H3K9me3 regions (Figure 2d). Analysing individual repeat families using UCSC RepeatMasker annotation, the ~340bp IAPLTR1/1a long terminal repeats showed comparable turnover to active promoters and enhancers, followed by IAPLTR2 and internal IAP regions (Extended Data Figure 6). Examining H3.3 dynamics across full-length IAPLTR1 elements, it was interesting to note that histone turnover was not confined to individual positioned nucleosome but broadly occurred over LTRs and adjacent ~2kb of internal region (Figure 2e). Remarkably, regions of highest turnover were also highly enriched for H3K9me3, concomitant with HP1 binding (Figure 2f-g). Prior analysis suggested that, as stereotypical for heterochromatin, DNA accessibility at ERVs is generally low [29], and analysis of recent ATAC-Seq data in mESC [20] confirmed that IAPLTRs neither show defined nucleosome-free regions nor broader domains of accessible DNA (Figure 2h). In summary, we find that histone turnover does not require or induce an increase in DNA accessibility at IAP ERVs (Figure 2f), Instead, it appears that histone H3.3/H4 is swapped into chromatin directly replacing an existing H3/H4 histone dimer or tetramer, hinting at the possibility of a concerted mechanism.

To achieve an exchange of core histones without rendering DNA at least transiently accessible, a tightly regulated supply of H3.3 would appear necessary. We thus wondered how a lack of H3.3 would affect DNA accessibility at H3.3+H3K9me3 regions. ATAC-Seq profiles of wildtype and H3.3 knockout cells have previously been compared [20]. The authors note that DNA accessibility is largely unchanged across the genome in the absence of H3.3, including sites of rapid histone exchange, such as promoters and enhancers [20]. Loss of H3.3 was thus fully compensated at promoters and enhancers, presumably by assembly or recycling pathways for H3.1/H3.2. In line with the original report, our re-analysis of the ATAC-Seq signals at H3.3-only peaks showed that DNA accessibility was not altered in H3.3KO cells (Figure 3a, b). Genome-wide analysis of 5kb bins and ChromHMM regions provided no evidence of systematic changes in DNA accessibility (Figure 3c, Extended Data Figure 7), suggesting that by large, loss of H3.3 had no effect on the packaging of the genome.

On the background of this, it was striking to find a roughly threefold increase in DNA accessibility in H3.3KO cells specifically at H3.3+H3K9me3 peaks (Figure 3a, b). Genome-wide analysis of 5kb bins further confirmed in an unbiased manner that H3K9me3+H3.3 co-enrichment regions showed the highest fold-increase in DNA accessibility in the absence of H3.3 (Figure 3d, Extended Data Figure 7). This suggested that H3.3, while dispensable for proper packaging of other regions of the genome, is required for maintaining a closed chromatin state at H3.3+H3K9me3 heterochromatin.

As discussed above, most H3.3+H3K9me3 heterochromatin is contributed by IAP ERVs (Figure 3d). Most IAP ERV containing 5kb bins moved from a low DNA accessibility (roughly genome-average) to the top quartile of accessible regions genome-wide in H3.3KO cells (Figure 3c). Although an increase in DNA accessibility could be indicative of a gain of euchromatic

histone modifications, loss of H3.3 did not lead to a gain in histone H3K18, H3K27, H3K64 or H3K122 acetylation at H3.3+H3K9me3 peaks (Figure 3b, Extended Data Figure 8). This is in line with prior observations that loss of H3.3 leads to only a partial loss of H3K9me3 modifications and a moderate increase in transcriptional activity [21,39].

We next wondered if specific functional DNA elements of the IAP ERV would become accessible in the absence of H3.3. In contrast to the typically narrow peaks over promoters and enhancers corresponding to transcription factor binding sites, ATAC-Seq signal increased broadly across LTR and ~2kb of internal region of IAP ERVs, in a profile that closely matched the distribution of H3.3 in wildtype cells (Figure 3e, see Figure 2e for comparison). Thus, H3.3 appears to play a specific role at IAP ERV in maintaining DNA in a tightly packaged and inaccessible state. Analysis of uniquely mappable paired-end reads within and adjacent to full-length IAP ERVs supports a widespread effect on many individual instances of the IAP ERV family (Figure 3f, Extended Data Figure 9). In addition to full-length or truncated IAP ERVs, the mouse genome contains many single LTRs (predominantly of the IAPLTR2 family). Using only uniquely mappable read pairs, we found that the 651 H3.3-enriched orphan LTRs also gained DNA accessibility in the absence of H3.3 (Figure 3g).

In summary, we find that interstitial heterochromatin maintains underlying DNA in an inaccessible state, a key characteristic of all heterochromatin. Despite this, interstitial heterochromatin is permissive to nucleosome turnover and histone H3.3 incorporation, providing a mechanistic explanation for the co-occurence of H3.3 and H3K9me3. According to the collective evidence from the data analyzed here, nucleosome eviction and assembly of new histone H3.3 nucleosomes must be accomplished through a mechanism that does not allow the underlying DNA to become accessible in the process. Loss of histone H3.3 may uncouple

eviction of existing nucleosomes from immediate reassembly of a new nucleosome and consequently lead to the observed broad gain in DNA accessibility.

DISCUSSION

In the current literature, heterochromatin is invariably associated with an inert chromatin packaging [33]. Dynamic histone exchange is inherently viewed as a mechanism that opposes heterochromatin formation [40]. Here we show that rapid histone exchange can occur even in regions that show traditional hallmarks of heterochromatin, such as H3K9me3, HP1 binding and inaccessibility of DNA. A dynamic turnover of histones provides a mechanistic explanation for the enrichment of histone variant H3.3 at IAP ERVs in mouse embryonic stem cells. We link the loss of H3.3 to a loss of heterochromatin integrity, as indicated by a gain in DNA accessibility. Remarkably, histone exchange appears to reinforce heterochromatin formation in this specific setting. Given that co-enrichment of H3.3 and H3K9me3 is to date a feature uniquely know in mouse pluripotent stem cells, it may be speculated that this new flavor of heterochromatin fulfills a specific function within the context of the highly plastic pluripotent chromatin.

Histone H3.3 is known to be deposited at interstitial heterochromatin through the histone chaperone DAXX [21,41–43]. DAXX has been proposed to contribute to ERV silencing through its histone chaperone activity and, independently, recruitment of histone deacetylases to ERVs [21,39]. Loss of H3.3 also reduces global DAXX levels in mouse ESC, but H3.3+H3K9me3 regions did not gain histone H3 acetylation in the absence of H3.3 (Extended Data Figure 8). This suggests that impaired nucleosome assembly in the absence of H3.3 was the cause of the observed opening of chromatin (Figure 3a). The mechanism how existing nucleosomes are evicted to facilitate exchange with new H3.3 is unknown. The chromatin remodeler ATRX can form a complex with DAXX and has previously been linked to ERV silencing [22,43–46], yet it is unclear if ATRX contributes substantially to nucleosome turnover and H3.3 incorporation [21,39,46]. The chromatin remodeler Smarcad1 has recently been linked to IAP ERVs in mouse ESC [47].

Smarcad1 has further been shown to have nucleosomes sliding and eviction activity, thus representing another candidate for mediating H3.3 turnover at ERVs [48,49]. Structural studies of the HP1-bound dinucleosome corroborate the view that heterochromatin can be accessible for nucleosome remodeling [32]. We show that supply of H3.3 is necessary for maintaining chromatin structure at IAP ERVs, implying that existing nucleosome eviction is directly linked to DAXX-dependent deposition of new H3.3. A major question remains why ATP-dependent turnover of old and incorporation of new histone is required as part of a repressive chromatin domain. Transient eviction of nucleosomes could allow KRAB zinc-finger proteins to sample the DNA more effectively and thereby facilitate recruitment of TRIM28 corepressor complex. Alternative models could be proposed where nucleosome eviction represents a mere side effect of a more widespread chromatin remodeling activity in embryonic stem cells, imposed by the need for a highly dynamic chromatin structure to maintain pluripotency [50]. Irrespective of the purpose of remodeling, nucleosome turnover necessitates sustained SETDB1 methyltransferase activity for maintaining high H3K9me3 levels at ERVs. This is because newly synthesized H3.3 is devoid of H3K9 methylation, and SETDB1 enzyme is known to establish H3K9me3 only on nucleosomal substrates. Histone H3K9me3 levels decrease in the absence of H3.3 [21], showing that histone H3.3 incorporation promotes rather than counteracts establishment of H3K9me3 and the formation of interstitial heterochromatin in mESC.

Collectively, our analyses shed first light into a dynamic heterochromatin state that defies the stereotypical distinction of euchromatin and heterochromatin. While the purpose and mechanism remain only poorly understood, our study provides a novel hypothesis and functional insights for future mechanistic studies.

Figure 1

a Time-ChIP
Genome-wide distribution (5kb bins)

b Chr13

c

d

e 5kb bins

Figure 1 - Genome-wide distribution of ectopic H3.3-SNAP in mouse ESC and NSC shows ESC-specific enrichment at interstitial heterochromatin.

a Violin plots showing genome-wide distribution of biotin-labeled H3.3-SNAP [9] in 5kb windows, in mouse embryonic stem cells (ESC) and neuronal stem cells (NCS). Biotin labeling of H3.3-SNAP was performed at 0h and followed over subsequent 3, 6 and 12h. **b** Genome tracks showing example region of time-ChIP experiment in ESC showing overlap of H3.3 with H3K27ac at enhancers and H3K9me3 at IAP ERVs. All tracks were normalized to Reads Per Genome Coverage (RPGC). See also Extended Data Figure 1a,b for controls and additional tracks. **c** Venn diagrams showing overlap of H3.3-SNAP 0h peaks with endogenous H3.3 peaks [21] in ESC (top) and H3.3-SNAP 0h peaks in NSC. **d** Peak density heatmaps and average profiles of H3.3-SNAP in ESC and NSC, at ESC H3.3 peaks [21]. Peaks were separated in two categories "H3.3-only" and "H3.3+H3K9me3" by the coincidence with H3K9me3 peaks [21]. Overlap with 15 State ChromHMM [36] states, full-length IAP ERV elements (from repMasker) and H3.3-SNAP peaks in ESC and NCS is shown. For additional information on ChromHMM annotation see Extended Data Figure 1, 2 **e** Scatter plot showing relationship between H3.3 and H3K9me3 assessed over 5kb bins genome-wide. Color scale represents density of underlying data points (5kb bins). Bottom graph shows overlap IAP ERVs. Overlap with additional repeat families are shown in Extended Data Figure 3.

a

H3.3-only
H3.3 + H3K9me3

ESC H3.3-SNAP Time-ChIP
0h 3h 6h 12h Input

b

ESC H3.3-HA Tet-ON
3h 6h

c

CATCH-IT

d

ESC H3.3-SNAP Time-ChIP

	1 Txn Elongation	2 Weak Txn	3 Txn Transition	4 Poised Enhancer	5 Active Promoter	6 Strong Enhancer	7 Active Promoter	8 Strong Enhancer	9 Strong Enhancer	10 Poised Promoter	11 Repressed	12 Heterochrom	13 Heterochrom	14 Heterochrom	15 Insulator	16 H3K9 Hetero	17 H3K9 H3 Hetero	
0h	0.72	0.38	1.55	1.91	1.86	2.15	2.41	2.23	1.21	1.51	0.17	-0.39	-0.47	0.44	1.00	0.13	1.81	0h
3h	0.50	0.26	1.11	1.44	1.20	1.55	1.78	1.66	0.86	1.08	0.13	-0.25	-0.29	0.31	0.70	0.22	1.64	3h
6h	0.46	0.24	1.04	1.37	1.12	1.44	1.64	1.53	0.81	1.03	0.12	-0.23	-0.25	0.30	0.65	0.23	1.51	6h
12h	0.25	0.11	0.68	1.01	0.79	0.98	1.16	1.05	0.54	0.78	0.09	-0.14	-0.13	0.18	0.43	0.33	1.17	12h

ESC H3.3-HA Tet-ON

3h	2.69	2.22	5.02	6.48	6.09	9.46	9.19	5.53	3.65	6.19	2.00	1.64	1.09	2.36	3.44	3.22	5.85	3h
6h	3.17	2.46	8.44	11.55	9.89	15.96	15.30	12.44	5.93	8.29	1.96	1.30	0.85	2.65	4.38	3.03	10.16	6h RPGC

ESC H3K9me3

	0.29	0.39	0.29	0.21	0.30	0.21	0.17	0.35	0.34	0.36	0.72	0.86	1.04	0.58	0.45	8.40		RPGC

15 State ChromHMM

log2 FC ChIP/Input

e H3.3-SNAP

0h
3h
6h
12h
input

IAPEz-int/LTR1(a)
(1297)

−2kb   2kb

f H3K9me3

H3K9me3

IAPEz-int/LTR1(a)
(1297)

−2kb   2kb

g HP1 binding

HP1α
HP1β
HP1γ

IAPEz-int/LTR1(a)
(1297)

−2kb   2kb

h ATAC-Seq

IAP ERV
random regions

IAPEz-int/LTR1(a)
(1297)

−2kb   2kb

Figure 2 - Histones are rapidly turned over at interstitial heterochromatin in the absence of DNA accessibility.

a Peak density heatmaps and average profiles of H3.3-SNAP time-ChIP data [9] over two classes of H3.3 peaks, H3.3-only and H3.3+H3K9me3 peaks. **b** Peak density heatmaps and average profiles of H3.3-HA pulse data [34] over H3.3 peaks. **c** Peak density heatmaps and average profiles of CATCH-IT data [6]. **d** Mean read density heatmap showing enrichments of H3.3-SNAP time-ChIP (log2 fold-change over input) over 15 ChromHMM states [37] as well as H3K9me3 and H3.3+H3K9me3 enriched regions [21]. For enrichment analysis by repeat families see also Extended Data Figure 6 **e** Average profiles of H3.3-SNAP time-ChIP over 1297 IAP ERVs. **f** Average profiles of H3K9me3 ChIP [51] over 1297 IAP ERVs. **g** Average profiles of BioChIP for HP1 isoforms [52] over 1297 IAP ERVs. **h** Average profiles of DNA accessibility [20] over 1297 IAP ERVs.

Figure 3

a

H3.3-only
H3.3 + H3K9me3

ATAC H3.3 WT   ATAC H3.3 KO   H3.3 SNAP 0h

RPGC

H3.3-only peaks

H3.3 + H3K9me3

b

Chr13   50,560 kb   50,580 kb   50,600 kb

ESC H3.3-SNAP
0h [0 - 22]
3h [0 - 22]
6h [0 - 22]
12h [0 - 22]

H3K27ac
WT [0 - 34]
KO [0 - 34]

ATAC
WT [0 - 31]
KO [0 - 31]

primed active   primed   active

Nutm2   IAPEz-int / IAPLTR2   RLTR13D6

c

5kb bins   all bins   IAP overlap

ATAC-Seq RPGC

H3.3 WT   H3.3 KO

d

5kb bins   H3.3 KO vs WT

IAP overlap
all bins

H3K9me3 RPGC

ATAC-Seq log2 KO/WT

e

H3.3 WT
H3.3 KO

RPGC

−2kb   IAPEz-int/LTR1(a) (1297)   2kb

f

ATAC-Seq

RPGC

IAP ERV

RPGC

full-length IAP ERV (225)   full-length IAP ERV (225)

-2.5kb   +2.5kb   -2.5kb   +2.5kb

g

ATAC-Seq

H33 WT
H33 KO

-2.5kb   +2.5kb

H3.3 SNAP   H33

-2.5kb   +2.5kb

IAP solo-LTRs (651)

Figure 3 - Interstitial heterochromatin becomes accessible in the absence of H3.3.

a Peak density heatmaps and average profiles of H3.3-SNAP time-ChIP [9] and ATAC-Seq [20] over two classes of H3.3 peaks, H3.3-only and H3.3+H3K9me3 peaks. See Extended Data Figures 7 and 8 for additional analysis **b** Genome tracks showing H3.3-SNAP time-ChIP, H3K27ac and ATAC-Seq signal over example region on Chr13. H3.3 peaks flank DNA accessibility peaks over enhancers while an IAP ERV is not associated with DNA accessibility in wildtype H3.3 cells (WT). ATAC-Seq signal greatly increases over IAP ERV in H3.3 knockout (KO) cells. **C** Violin plots showing genome-wide distribution of ATAC-Seq signal in 5kb bins, in H3.3 wildtype (WT) and knockout (KO) cell line. Overlayed in color are 1713 individual bins overlapping with full-length or truncated IAP ERVs (longer than 2kb). **d** Scatter plot of 5kb bins, showing H3K9me3 level versus log2-fold change in ATAC-Seq signal upon H3.3 knockout. Bins overlapping with IAP ERVs are drawn in black. **e** Average profile of ATAC-Seq signal over 1297 IAP ERVs in H3.3 wildtype (WT) and knockout (KO) cell line. **f** Density heatmaps and average profiles of ATAC-Seq signal using only uniquely mappable reads, over 225 full-length IAP ERVs and flanking regions. See Extended Data Figure 9 for multimap read analysis **g** Average profiles of ATAC-Seq signal (top) and H3.3-SNAP ChIP using only uniquely mappable reads, over 651 IAP solo LTRs (mostly IAPLTR2),

Methods

Primary data analysis pipeline. Original FASTQ files were downloaded from GEO database (see Extended Data Table 1 for list of GEO accessions). Paired-end reads were mapped to mm9 reference genome using bowtie2 (v 2.3.5.1) [53], deduplicated using Picard MarkDuplicates with default settings and filtered for blacklisted regions. Bigwig files were generated from resulting deduplicated and filtered bam files using deepTools (v 3.1.0) [54], normalized to 1x Genome Coverage (Reads Per Genome Coverage) using --normalizeUsing RPGC --effectiveGenomeSize 2150570000. Unless stated otherwise, downstream analysis was performed on the normalized bigwig files. Where available, we evaluated all experimental replicates (see Extended Data Figure 5). For simplicity, figures show the first replicate.

Track visualization. We visualize genomic tracks from 1x normalized BigWig files using Integrated Genome Viewer (IGV) [55]. Y axis corresponds to Reads Per Genome Coverage (RPGC).

Peak calling. Peaks were calculated using MACS2 (v 2.1.2) [56] callpeak using parameters: broadPeak, default --broad-cutoff 0.05, corresponding input alignment as control. The three replicates were combined to obtain a reliable set of peaks. A peak is considered if it is called in at least two out of the three replicates. This was performed for both ESC and NSC at initial timepoint (0h). The region considered as reliable peak is then the result of merging the overlapping called peak regions.

Bin-based analyses. For the genome-wide coverage analyses in 5kb bins, mean coverage bin files were generated with deepTools multiBigWigSummary bins tool from the final bigwig files using a bin size of 5kb.

Annotated bins. Genome-wide 5kb bins were intersected with UCSC RepMasker repetitive element groups using bedtools (v2.27.1) intersect [57]. Bins were assigned the repetitive element group of the element that overlapped the most with each bin, in the unusual case where the same bin overlaps more than one repetitive element.

Annotated peaks. A peak list from endogenous H3.3 ChIP-seq [21] was intersected with peaks called from [9] as described above. Any amount of overlap > 0 was considered. Other loci of interest were also intersected with these peaks in the same fashion: IAPEz and ChromHMM15 regions. Since ChromHMM15 is a partition of the genome, a peak is expected to fall in at least one category. For the cases where a peak overlapped with more than one ChromHMM annotation, the largest overlap was reported.

Peak density heatmaps. Values were calculated with deepTools (v 3.1.0) computeMatrix using parameters --referencePoint center and flanking regions of 2,500 base pairs (-a 2500 -b 2500). The resulting matrices were visualized using matplotlib (v 3.1.1) with the same parameter settings as deepTools plotHeatmap. In combined plots, the heatmap on the left is sorted by mean coverage per locus and sorting order is applied to all other heatmaps. Profile plots on top of each heatmap are calculated using the mean values of the columns in the matrices generated by deepTools.

IAPEz flanking H3.3 density heatmaps. Values were calculated with deepTools. In this case the execution mode was reference-point with --regionBodyLength set to 6500 to match the average length of the studied loci. A set of uniquely-mapped BigWig files was generated from the original aligmnent files using deepTools bamCoverage tool with parameters --minMappingQuality 10 --extendReads 150. For paired-end data, coverage was extended to match fragment length.

Mean read density heatmaps. Mean coverage values over each of the ChromHMM15 regions were calculated with bwtool (v1.0) summary function and normalized to their corresponding input value (log2 ratio). Values obtained were visualized using R package pheatmap.

Genome-wide enrichment over 5kb bins for ES and NS cells, timepoints 0h, 3h, 6h and 12h. Bins were generated using deepTools multiBigWigSummary (see Bins). Each bin value was divided by its corresponding input value. Bins with zero values were excluded from this analysis. Additionally, outliers were removed using Tukey's 1.5 IQR rule for the violin plots.

Profile plots. Average plots over genomic regions were generated from normalized BigWig files using SeqPlots [58].

Datasets: A list of all used datasets is provided as Supplementary Data Table 1.

ACKNOWLEDGEMENTS

Bioinformatics analyses were performed on resources provided by the Swedish National Infrastructure for Computing (SNIC) at Uppmax server (projects SNIC 2020/15-9, SNIC 2020/6-3, uppstore2018208, SNIC 2018/3-669, sllstore2017057, SNIC 2017/1-508).

References

Bibliography

1. Elsaesser, S. J., Goldberg, A. D. & Allis, C. D. New functions for an old variant: no substitute for histone H3.3. *Curr. Opin. Genet. Dev.* **20,** 110–117 (2010).

2. Ahmad, K. & Henikoff, S. The histone variant H3.3 marks active chromatin by replication-independent nucleosome assembly. *Mol. Cell* **9,** 1191–1200 (2002).

3. Tagami, H., Ray-Gallet, D., Almouzni, G. & Nakatani, Y. Histone H3.1 and H3.3 complexes mediate nucleosome assembly pathways dependent or independent of DNA synthesis. *Cell* **116,** 51–61 (2004).

4. Ray-Gallet, D. *et al.* Dynamics of histone H3 deposition in vivo reveal a nucleosome gap-filling mechanism for H3.3 to maintain chromatin integrity. *Mol. Cell* **44,** 928–941 (2011).

5. Ha, M., Kraushaar, D. C. & Zhao, K. Genome-wide analysis of H3.3 dissociation reveals high nucleosome turnover at distal regulatory regions of embryonic stem cells. *Epigenetics Chromatin* **7,** 38 (2014).

6. Banaszynski, L. A. *et al.* Hira-dependent histone H3.3 deposition facilitates PRC2 recruitment at developmental loci in ES cells. *Cell* **155,** 107–120 (2013).

7. Goldberg, A. D. *et al.* Distinct factors control histone variant H3.3 localization at specific genomic regions. *Cell* **140,** 678–691 (2010).

8. Schlesinger, S. *et al.* A hyperdynamic H3.3 nucleosome marks promoter regions in pluripotent embryonic stem cells. *Nucleic Acids Res.* **45,** 12181–12194 (2017).

9. Deaton, A. M. *et al.* Enhancer regions show high histone H3.3 turnover that changes during differentiation. *elife* **5,** (2016).

10. Schneiderman, J. I., Orsi, G. A., Hughes, K. T., Loppin, B. & Ahmad, K. Nucleosome-depleted chromatin gaps recruit assembly factors for the H3.3 histone variant. *Proc Natl Acad Sci USA* **109,** 19721–19726 (2012).

11. Elsaesser, S. J. & Allis, C. D. HIRA and Daxx constitute two independent histone H3. 3-containing predisposition complexes. *Cold Spring Harbor Symposia on Quantitative Biology* (2010).

12. Mito, Y., Henikoff, J. G. & Henikoff, S. Genome-scale profiling of histone H3.3 replacement patterns. *Nat. Genet.* **37,** 1090–1097 (2005).

13. Schwartz, B. E. & Ahmad, K. Transcriptional activation triggers deposition and removal of the histone variant H3.3. *Genes Dev.* **19,** 804–814 (2005).

14. Jin, C. & Felsenfeld, G. Nucleosome stability mediated by histone variants H3.3 and H2A.Z. *Genes Dev.* **21,** 1519–1529 (2007).

15. Flaus, A., Rencurel, C., Ferreira, H., Wiechens, N. & Owen-Hughes, T. Sin mutations alter inherent nucleosome mobility. *EMBO J.* **23,** 343–353 (2004).

16. Thakar, A. *et al.* H2A.Z and H3.3 histone variants affect nucleosome structure: biochemical and biophysical studies. *Biochemistry* **48,** 10852–10857 (2009).

17. Loyola, A., Bonaldi, T., Roche, D., Imhof, A. & Almouzni, G. PTMs on H3 variants before chromatin assembly potentiate their final epigenetic state. *Mol. Cell* **24,** 309–316 (2006).

18. Sidoli, S. *et al.* A mass spectrometry-based assay using metabolic labeling to rapidly monitor chromatin accessibility of modified histone proteins. *Sci. Rep.* **9,** 13613 (2019).

19. Armache, A. *et al.* Phosphorylation of the ancestral histone variant H3.3 amplifies stimulation-induced transcription. *BioRxiv* (2019). doi:10.1101/808048

20. Martire, S. *et al.* Phosphorylation of histone H3.3 at serine 31 promotes p300 activity and enhancer acetylation. *Nat. Genet.* **51,** 941–946 (2019).

21. Elsässer, S. J., Noh, K.-M., Diaz, N., Allis, C. D. & Banaszynski, L. A. Histone H3.3 is required for endogenous retroviral element silencing in embryonic stem cells. *Nature* **522,** 240–244 (2015).

22. Sadic, D. *et al.* Atrx promotes heterochromatin formation at retrotransposons. *EMBO Rep.* **16,** 836–850 (2015).

23. Rowe, H. M. *et al.* KAP1 controls endogenous retroviruses in embryonic stem cells. *Nature* **463,** 237–240 (2010).

24. Matsui, T. *et al.* Proviral silencing in embryonic stem cells requires the histone methyltransferase ESET. *Nature* **464,** 927–931 (2010).

25. Rowe, H. M. *et al.* TRIM28 repression of retrotransposon-based enhancers is necessary to preserve transcriptional dynamics in embryonic stem cells. *Genome Res.* **23,** 452–461 (2013).

26. Stoll, G. A. *et al.* Structure of KAP1 tripartite motif identifies molecular interfaces required for retroelement silencing. *Proc Natl Acad Sci USA* (2019). doi:10.1073/pnas.1901318116

27. Fukuda, K., Okuda, A., Yusa, K. & Shinkai, Y. A CRISPR knockout screen identifies SETDB1-target retroelement silencing factors in embryonic stem cells. *Genome Res.* **28,** 846–858 (2018).

28. Maksakova, I. A. *et al.* Distinct roles of KAP1, HP1 and G9a/GLP in silencing of the two-cell-specific retrotransposon MERVL in mouse ES cells. *Epigenetics Chromatin* **6,** 15 (2013).

29. He, J. *et al.* Transposable elements are regulated by context-specific patterns of chromatin marks in mouse embryonic stem cells. *Nat. Commun.* **10,** 34 (2019).

30. Maksakova, I. A. *et al.* H3K9me3-binding proteins are dispensable for SETDB1/H3K9me3-dependent retroviral silencing. *Epigenetics Chromatin* **4,** 12 (2011).

31. Sanulli, S. *et al.* HP1 reshapes nucleosome core to promote phase separation of heterochromatin. *Nature* **575,** 390–394 (2019).

32. Machida, S. *et al.* Structural basis of heterochromatin formation by human HP1. *Mol. Cell* **69,** 385-397.e8 (2018).

33. Nishibuchi, G. & Déjardin, J. The molecular basis of the organization of repetitive DNA-containing constitutive heterochromatin in mammals. *Chromosome Res.* **25,** 77–87 (2017).

34. Yildirim, O. *et al.* A system for genome-wide histone variant dynamics in ES cells reveals dynamic MacroH2A2 replacement at promoters. *PLoS Genet.* **10,** e1004515 (2014).

35. Chronis, C. *et al.* Cooperative binding of transcription factors orchestrates reprogramming. *Cell* **168,** 442-459.e20 (2017).

36. Ernst, J. & Kellis, M. ChromHMM: automating chromatin-state discovery and characterization. *Nat. Methods* **9,** 215–216 (2012).

37. Bogu, G. K. *et al.* Chromatin and RNA maps reveal regulatory long noncoding rnas in mouse. *Mol. Cell. Biol.* **36,** 809–819 (2015).

38. Deal, R. B. & Henikoff, S. Capturing the dynamic epigenome. *Genome Biol.* **11,** 218 (2010).

39. Hoelper, D., Huang, H., Jain, A. Y., Patel, D. J. & Lewis, P. W. Structural and mechanistic insights into ATRX-dependent and -independent functions of the histone chaperone DAXX. *Nat. Commun.* **8,** 1193 (2017).

40. Henikoff, S. & Shilatifard, A. Histone modification: cause or cog? *Trends Genet.* **27,** 389–396 (2011).

41. Drané, P., Ouararhni, K., Depaux, A., Shuaib, M. & Hamiche, A. The death-associated protein DAXX is a novel histone chaperone involved in the replication-independent deposition of H3.3. *Genes Dev.* **24,** 1253–1265 (2010).

42. Elsässer, S. J. *et al.* DAXX envelops a histone H3.3-H4 dimer for H3.3-specific recognition. *Nature* **491,** 560–565 (2012).

43. Lewis, P. W., Elsaesser, S. J., Noh, K.-M., Stadler, S. C. & Allis, C. D. Daxx is an H3.3-specific histone chaperone and cooperates with ATRX in replication-independent chromatin assembly at telomeres. *Proc Natl Acad Sci USA* **107,** 14075–14080 (2010).

44. He, Q. *et al.* The Daxx/Atrx Complex Protects Tandem Repetitive Elements during DNA Hypomethylation by Promoting H3K9 Trimethylation. *Cell Stem Cell* **17,** 273–286 (2015).

45. Xue, Y. *et al.* The ATRX syndrome protein forms a chromatin-remodeling complex with Daxx and localizes in promyelocytic leukemia nuclear bodies. *Proc Natl Acad Sci U S A* **100,** 10635–40 (2003).

46. Voon, H. P. J. *et al.* ATRX plays a key role in maintaining silencing at interstitial heterochromatic loci and imprinted genes. *Cell Rep.* **11,** 405–418 (2015).

47. Sachs, P. *et al.* SMARCAD1 ATPase activity is required to silence endogenous retroviruses in embryonic stem cells. *Nat. Commun.* **10,** 1335 (2019).

48. Byeon, B. *et al.* The ATP-dependent chromatin remodeling enzyme Fun30 represses transcription by sliding promoter-proximal nucleosomes. *J. Biol. Chem.* **288,** 23182–23193 (2013).

49. Awad, S., Ryan, D., Prochasson, P., Owen-Hughes, T. & Hassan, A. H. The Snf2 homolog Fun30 acts as a homodimeric ATP-dependent chromatin-remodeling enzyme. *J. Biol. Chem.* **285,** 9477–9484 (2010).

50. Schlesinger, S. & Meshorer, E. Open chromatin, epigenetic plasticity, and nuclear organization in pluripotency. *Dev. Cell* **48,** 135–150 (2019).

51. Shi, H. *et al.* ZFP57 regulation of transposable elements and gene expression within and beyond imprinted domains. *Epigenetics Chromatin* **12,** 49 (2019).

52. Ostapcuk, V. *et al.* Activity-dependent neuroprotective protein recruits HP1 and CHD4 to control lineage-specifying genes. *Nature* **557,** 739–743 (2018).

53. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9,** 357–359 (2012).

54. Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A. & Manke, T. deepTools: a flexible platform for exploring deep-sequencing data. *Nucleic Acids Res.* **42,** W187-91 (2014).

55. Robinson, J. T. *et al.* Integrative genomics viewer. *Nat. Biotechnol.* **29,** 24–26 (2011).

56. Liu, T. Use model-based Analysis of ChIP-Seq (MACS) to analyze short reads generated by sequencing protein-DNA interactions in embryonic stem cells. *Methods Mol. Biol.* **1150,** 81–95 (2014).

57. Quinlan, A. R. BEDTools: The Swiss-Army Tool for Genome Feature Analysis. *Curr. Protoc. Bioinformatics* **47,** 11.12.1-34 (2014).

58. Stempor, P. & Ahringer, J. SeqPlots - Interactive software for exploratory data analyses, pattern discovery and visualization in genomics. [version 1; peer review: 2 approved, 1 approved with reservations]. *Wellcome Open Res.* **1,** 14 (2016).

59. Di Tommaso, P. *et al.* Nextflow enables reproducible computational workflows. *Nat. Biotechnol.* **35,** 316–319 (2017).

60. Bunch, H. *et al.* TRIM28 regulates RNA polymerase II promoter-proximal pausing and pause release. *Nat. Struct. Mol. Biol.* **21,** 876–883 (2014).

An embryonic stem cell-specific heterochromatin state allows core histone exchange in the absence of DNA accessibility

Carmen Navarro and Simon J Elsässer

Affiliations

Science for Life Laboratory, Department of Medical Biochemistry and Biophysics, Karolinska Institutet, Stockholm, Sweden

Ming Wai Lau Centre for Reparative Medicine, Stockholm node, Karolinska Institutet, Stockholm, Sweden

Corresponding Author

Simon J Elsässer, simon.elsasser@scilifelab.se

SUPPLEMENTARY INFORMATION

9 Extended Data Figures

1 Extended Data Table (separate file)

Extended Data Figure 1

Extended Data Figure 1 - related to Figure 1a, b - Additional examples and controls of H3.3 enrichment at active regions and interstitial heterochromatin.

a Genome tracks example of time-ChIP data (H3.3-SNAP and H3.1-SNAP timepoints 0h throughout 12h) in mouse embryonic stem cells (ESC), together with H3.3-HA pulse data [9,34], H3K27ac and H3K9me3 [20,51]. H3.3-SNAP and H3K27ac enriched regions show overlap with active and primed enhancers. H3.3-SNAP and H3K9me3 coincide with IAP ERV (see also genome-wide analysis in Extended Data Figure 3). H3.1-SNAP and Input controls show no particular enrichment. All tracks were normalized to Reads Per Genome Coverage (RPGC). **b** Genome tracks example showing time-ChIP H3.3 data in both mouse embryonic stem cells (ESC) and neural stem cells (NSC), together with H3K9me3 ChIP [51] and RNA Polymerase II ChIP [60] at Hist1 histone gene cluster. H3.3 enrichment and turnover at IAP elements is observed in ESC but not NSC, whereas H3.3 enrichment and turnover at the housekeeping genes is maintained in NSC. **c** Peak density heatmaps and average profiles (RPGC) of time-ChIP H3.3-SNAP in embryonic stem cells (ESC) and neural stem cells (NSC) and H3.1-SNAP at timepoint 0h in ESC over endogenous H3.3 peaks in ESC coinciding (H3.3+H3K9me3) or not (H3.3-only) with H3K9me3 peaks [21]. Overlap with 15-State ChromHMM annotation [37] states, IAP ERV elements (IAPEz-int and flanking LTRs from mm9 RepeatMasker track), and H3.3-SNAP peaks called in ESC and NCS is shown. Additionally, ESC and NSC input data is also shown. **d** H3.3 and H3K9me3 profiles along a reprogramming time-course from mouse embryonic fibroblasts (MEF) [35]. Early reprogramming intermediate (48h after expression of reprogramming factors Oct4, Nanog, Klf4, Myc) and late reprogramming intermediate stable cell line (pre-iPSC) are compared to MEF and ESC. All tracks are plotted on the same scale as RPGC. H3K9me3 is first enriched at IAP ERV at the pre-iPSC intermediate

and further increases in ESC. H3.3 is exclusively acquired at IAP ERVs in ESC, while nearby housekeeping gene *Rps12* is enriched consistently in all cell types.

Extended Data Figure 2

a

Extended Data Figure 2 - related to Figure 1d - Annotation of H3.3 peaks using 15-State ChromHMM model

Barplot showing relative overlap of "H3.3-only" and "H3.3+H3K9me3" peak sets [21] with 15 State ChromHMM annotation, grouped by coincidence with H3K9me3 peaks. Each peak is assigned the ChromHMM15 state that overlaps with it the most. Amounts are shown as a percentage of the total peak set. H3.3-only peaks overlap the most with strong enhancers and transcription elongation states, whereas H3.3+H3K9me3 peaks overlap with heterochromatin states 12-14, defined by absence of H3K4me3, H3K4me1, H3K27ac, H3K36me3 and RNAP2 marks.

Extended Data Figure 3

Genome-wide analysis in ~545k 5kb bins (Log2 fold-enrichment over input)

Extended Data Figure 3 - related to Figure 1e - Genome-wide correlation of H3.3 and H3K9me3 is shaped by a subset of ERV families

Scatter plot showing relationship between H3.3 and H3K9me3 signal assessed over 5kb bins genome-wide. Genome was partitioned in 5kb bins and coverage enrichment was calculated as log2 fold change of each signal over their corresponding input. Color of each tile represents point density of all the bins with similar H3K9me3 and H3.3 values (on a tile-resolution of 0.1). RepMasker annotation was intersected with 5kb bins for major repeat families. Bins that overlap with the corresponding repeat type are shown in black. In the rare case where more than one element overlapped with the same bin, the repeat type that overlapped the most was adjudicated.

Extended Data Figure 4

Extended Data Figure 4 - related to Figure 1e - ETn but not IAP ERV families contain a small number of active elements in the annotated genome. a Heatmap showing RNA Polymerase II and H3K9me3 signal [51,60] over unique 5kb flanking regions of ETn repeat elements. Among the annotated ETn elements, a small subset show high RNA Polymerase II signal at their 3' flanking region together with absence of H3K9me3, which suggests that these individual elements are transcribed at high levels, whereas the remaining elements are in a repressed or inactive state. **b** Heatmap showing RNA Polymerase II and H3K9me3 signal over 5kb flanking regions of annotated IAP repeat elements. No actively transcribed IAP ERVs are discernable by RNA Polymerase II signal. **c** Density plot of transcript levels for genes and repeat elements. IAP elements show extremely low transcript level per element as compared to genes. ETn elements show significantly higher transcript levels (over 2 orders of magnitude) than IAP elements, likely arising from the small number of active copies.

Extended Data Figure 5

a

H3.3 Tet-ON (Schlesinger et. al. 2017)   H3.3 Tet-OFF (Ha et. al. 2014)   H3.3 Tet-ON (Yildirim et. al. 2014)

b

Extended Data Figure 5 - related to Figure 2a-c - Analysis of additional H3.3 pulsing datasets. a Peak Density Heatmaps and average profiles of H3.3 pulse or chase data from three different studies in mouse ESC [5,8,34] at "H3.3-only" and "H3.3+H3K9me3" peak sets [21]. **b** Mean read density heatmap showing enrichments of H3.3-SNAP time-ChIP (log2 fold-change over input) over 15 ChromHMM states as well as H3K9me3 and H3.3+H3K9me3 enriched regions [21]. Same as Figure 2d but with additional replicates [9].

Extended Data Figure 6

mm9 RepeatMasker annotation (>200bp elements)

Extended Data Figure 6 - related to Figure 2d - Analysis of repetitive element families

a Mean read density heatmaps of time-ChIP H3.3-SNAP data at a major repetitive elements from RepeatMasker annotation (filtered for elements larger than 200bp). Each category enrichment is calculated as log2 fold change of mean coverage over input. IAPLTR1a shows the highest enrichment at time point 0h and turnover. Most IAP LTRs and internal sequences, as well as ETn and MusD elements show both enrichment and H3.3 turnover.

Extended Data Figure 7

a

ATAC-Seq signal

b

ATAC-Seq

mm9 RepeatMasker annotation (>200bp elements)

c

Genome-wide 5kb bins signal

ATAC-Seq H3.3 KO vs H3.3 WT

Extended Data Figure 7 - related to Figure 3a - Genome-wide maintenance of accessibility in the absence of H3.3KO cells independent of active or inactive chromatin state, and selective gain in accessibility at H3K9me3-H3.3 co-enriched regions.

a Mean read density heatmap of ATAC-seq data of H3.3 wildtype and knockout cells [20]. Both WT and KO show very similar genome-wide read density levels accross all 15 ChromHMM States. While H3K9me3-only regions remain inaccessible, a striking ~3-fold increase in ATAC-Seq signal is found at H3.3 + H3K9me3 regions in H3.3KO cells. **b** Mean read density heatmap of ATAC-seq data of H3.3 wildtype and knockout by repeat family. Increase in accessibility is observed for those repeat families that are enriched in H3.3 in wildtype cells. **c** Tile-plot showing relationship between H3.3 and H3K9me3 signal assessed over 5kb bins genome-wide. Each tile color shows log2 fold enrichment of average ATAC-seq signal for H3.3 KO over WT of all the bins with similar H3K9me3 and H3.3 values (on a tile-resolution of 0.1). A majority of zero values show ATAC-seq signal is very similar genome-wide, with H3.3 KO showing highest enrichment over WT on regions both H3.3- and H3K9me3-enriched.

Extended Data Figure 8

Extended Data Figure 8 - related to Figure 3a - Loss of H3.3 does not lead to an increase in histone acetylations at H3.3+H3K9me3 regions.

a Peak Density Heatmaps and average profiles of H3K18ac, H3K27ac, H3K64ac and H3K122ac marks for H3.3 WT and KO samples [20] at H3.3-only and H3.3+H3K9me3 peak sets. As previously described, histone acetylation, present at a subset of H3.3 peaks, is reduced in the absence of H3.3 [20], but basal levels at H3.3+H3K9m3 regions are not changed.

Extended Data Figure 9

Extended Data Figure 9 - related to Figure 3f - Uniquely mappable reads show widespread increase in DNA accessibility over individual IAP ERV elements.

Density heatmap of uniquely mappable paired-end reads within and adjacent to 225 full-length IAP ERVs for H3K9me3, H3.3 KO and WT ATAC-seq, and Time-ChIP H3.3-SNAP at timepoint 0h. Top row shows average profiles. Uniquely mapped reads at flanking regions allow to distinguish among individual instances of repetitive elements, showing that the observed effect is pervasive across such repeat elements. Lower row shows the corresponding unfiltered signal including multi-mappable reads for the same datasets at the same IAP ERVs set. Multimappable reads were assigned randomly to one of the ambigous matches.