1 **Title: Spatiotemporal structure of sensory-evoked and spontaneous activity revealed by**

2 **mesoscale imaging in anesthetized and awake mice**

3
4

Authors: Navvab Afrashteh[1,3], Samsoon Inayat[1,3], Edgar Bermudez Contreras[1,3], Artur Luczak[1],

Bruce L. McNaughton[1,2], Majid H. Mohajerani[1*]

[1]University of Lethbridge, Faculty of Arts and Sciences, Department of Neuroscience, 4401

University Dr. W., Lethbridge, Alberta, Canada, T1K 3M4.

[2]Center for Neurobiology of Learning and Memory, Department of Neurobiology and Behavior,

University of California, Irvine.

[3]These authors contributed equally.

[*] Corresponding author: Majid H. Mohajerani (mohajerani@uleth.ca)

Abstract

The functional role of cortical spontaneous activity is still not well understood. For instance, it remains controversial, how spontaneous activity relates to sensory evoked patterns, with some studies showing striking similarities between both, while other studies reporting differences. To investigate those discrepancies, we systematically compared several characteristics of visual, auditory and tactile evoked activity elicited with different stimulus strengths and spontaneous activity motifs in lightly anesthetized and awake mice using mesoscale wide-field voltage-sensitive dye and glutamate imaging respectively. In particular, we analyzed: amplitude, speed, and propagation trajectory in spontaneous and evoked activity patterns. We found that spontaneous activity had similar amplitude, and speed when compared to evoked activity elicited with low stimulus strengths. However, this similarity gradually decreased when the strength of stimuli eliciting evoked responses increased. Moreover, stronger stimuli elicited larger amplitude evoked activity with higher speed and with more stereotypical propagation trajectory. Similarly, amplitude of spontaneous activity was positively correlated with the speed of propagation. Interestingly, the trajectories of evoked activity covered a smaller cortical area compared to spontaneous activity. Altogether, these results show that the relationship between spontaneous and sensory evoked activity depends on the strength of sensory stimuli. Thus, this study helps to reconcile the previously reported conflicting findings when comparing evoked and spontaneous activity.

Keywords: wide-field optical imaging, cortical states, sensory-evoked activity, spontaneous activity, optical flow analysis, activity propagation speed, direction, trajectories

1 Introduction

Much of the development of our current understanding of sensory processing comes from the study of evoked activity and the stimulus-response relationships at different stages in the nervous system (Seung and Sompolinsky, 1993; Butts and Goldman, 2006; Jones and Smith, 2014). However, in the absence of stimulus the cortex remains active, even in the primary sensory areas. Until not long ago, spontaneous activity used to be regarded as 'noise' (Parga and Abbott, 2007; Faisal et al., 2008; Stringer et al., 2016). However, this idea has changed significantly and now the study of spontaneous activity is considered to be crucial for the understanding of cortical function (McCormick, 1999; Ringach, 2009; Raichle, 2010; Deco et al., 2013).

Despite significant advances in the understanding of cortical processing underlying both spontaneous and evoked activity, their relationship and interactions remain under debate. In some studies, the spatiotemporal patterns of evoked and spontaneous activity are reported to be similar (Hoffman and McNaughton, 2002; Kenet et al., 2003; Han et al., 2008; Luczak et al., 2009; Berkes et al., 2011), while in others they are reported to have remarkable differences (Stringer et al., 2019). For example, at the microcircuit level, Luczak and colleagues proposed that the temporal patterns of tone-evoked spiking activity occur as part of a larger set of patterns produced during spontaneous activity recorded using silicon probes within auditory cortex in anesthetized rats (Luczak et al., 2009; Luczak et al., 2015). In contrast, more recent studies using large-scale high-density optical imaging as well as silicon probes to record the single-unit activity of large neuronal populations, report that the patterns of activity from evoked responses belong to a different space from the one in spontaneous activity (Stringer et al., 2019).

At the mesoscale level, there are reported similarities in the characteristics of evoked and spontaneous activities. For example, Arieli et al. stated that the amplitude of the ongoing

spontaneous activity was similar to the one in evoked responses in a portion of the anesthetized cat visual cortex (Arieli et al., 1995). Moreover, Kenet et al. suggested that spontaneous activity patterns are similar to the orientation maps observed in visually evoked responses in anesthetized cats (Kenet et al., 2003). More recently, it has been shown that these maps emerge only in anaesthetized states but not in awake states (Omer et al., 2019). Similarly, Han et al. showed that the propagation patterns of spontaneous activity resembles recent sensory-evoked patterns when measured in a portion of the visual cortex in anesthetized rats (Han et al., 2008). At the macro-scale level, the fluctuations of the BOLD signal in functional magnetic resonance imaging (fMRI) in humans has been reported to be larger during spontaneous activity (resting state) than during task-driven activity (He, 2011; Ponce-Alvarez et al., 2015; Ito et al., 2019).

Some of the contradicting results on the relationship between evoked and spontaneous activity in previous studies can be explained by methodological differences in these diverse works. For example, these studies use different animal models which can affect the certain characteristics of cortical processing (e.g. theta oscillations in humans are significantly slower than in rodents) (Alloway et al., 1993; Jacobs, 2014). Moreover, the recording techniques used in these studies have important intrinsic differences in spatial and temporal resolutions, which in turn might impact the measurements to compare evoked and spontaneous activity (Menon and Kim, 1999). Furthermore, there are also inconsistencies in the preparations used in these studies such as brain slices or *in vivo* recordings which have distinct activity dynamics and might impact their conclusions (Azzarelli et al., 2017). Also, the brain states in which the recordings were performed might be different. For example, the cortical dynamics of anesthetized or sleep animals have remarkable differences to the ones from awake or desynchronized brain states (Sellers et al., 2015).

Finally, the sensory modalities from which the brain activity was recorded from might not be

directly comparable.

In this study, we used VSD and glutamate mesoscale imaging of most of the right cerebral

hemisphere in lightly anesthetized and awake mice, respectively. Moreover, we used optical flow

analysis, to characterize and compare the spatiotemporal dynamics of sensory-evoked activity to

to motifs of spontaneous activity that originate in the same cortical region using evoked activity-

derived template matching (Mohajerani et al., 2013; Afrashteh et al., 2017). Overall, this technique

offers the possibility of monitoring activity over large portions of the cortical mantle at high

spatiotemporal resolution which, in turn, permits the application of methods to characterize the

cortical activity dynamics in both regimes. In contrast with previous studies, we used different

sensory modalities (tactile, auditory, and visual), several stimulation strengths, and brain states,

which provide a more complete comparison of ongoing and evoked cortical activity.

We found that the response amplitude, propagation speed, and extent of propagation of the evoked

activity in the supra-granular somatosensory cortex has a positive correlation with the stimulus

intensity. The speed, direction stability, and extent of propagation of spontaneous activity was also

positively correlated with the amplitude. Our results demonstrate that, for the above-mentioned

parameters, sensory-evoked responses at low stimulus strengths are similar to motifs of

spontaneous activity and that this similarity diminishes as stimulus strength increases. Finally, we

showed that the repertoire of directions and trajectories of propagation of sensory-evoked activity

is smaller than the one from spontaneous activity.

In summary, this study helps to clarify the discrepancies in the previous research works comparing

evoked and spontaneous activity by providing a more complete and uniform experimental and

analytical framework.

2 Materials and Methods

2.1 Animals and Surgery

Animal housing, surgery, and all experimental protocols were approved by the University of Lethbridge Animal Care Committee in line with the guidelines from the Canadian Council for Animal Care. For anesthetized preparations, adult male C57BL/6J mice (n=5 animals for forelimb and hindlimb stimulation experiments and n=4 animals for visual stimulation experiments) around ~3 months of age and weighing approximately 25g were used. Mice were anesthetized with isoflurane (1.0–1.5%) for induction and during surgery, and a reduced concentration during data collection (0.5–1.0%). A 7×6 mm unilateral craniotomy (bregma +2.5 to -4.5 mm, lateral to the midline 0 to 6 mm) was made, and the dura was removed, as described previously (Kyweriga and Mohajerani, 2016). The body temperature was maintained at 37°C during surgeries and subsequent imaging sessions. For awake preparations, n=5 adult mice (2 males and 3 females) Ai85-CamKtTA-EMX1-Cre (cross of B6; 129S-Igs7tm85.1 (tetO-gltI/GFP*) Hze/J and B6.129S2-Emx1tm1(cre)Krj/J) with 5–8 months of age and weighing approximately 26g were used. These mice express intensity-based glutamate sensing fluorescent reporter (iGluSnFR) in excitatory neurons and glial cells of cortex (Xie et al., 2016; Karimi Abadchi et al., 2020). One week prior to performing wide-field glutamate imaging, a surgery was done to expose the cranial bone and for implanting a head-plate. A large portion of the cranial bone was exposed by removing the skin. The cranial bone was then cleaned and metabond was applied on top. Finally, a glass coverslip was placed on top to keep the surface clear from accumulating debris from the animal's home cage and the environment.

2.2 Data acquisition

2.2.1 Wide-field optical imaging

Wide-field optical imaging of cortical population activity was utilized to capture the mesoscale dynamics of the cortex in both anesthetized and awake preparations. For voltage-sensitive dye imaging in anesthetized mice, the dye RH-1691 (Optical Imaging, New York, NY) (Shoham et al., 1999), was dissolved in HEPES-buffered saline solution (0.5 mg/ml) and applied to the exposed cortex for 30-40 min, as described previously (Mohajerani et al., 2010; Mohajerani et al., 2011; Lim et al., 2012; Lim et al., 2013). The rest of the dye was carefully washed and the brain was covered with 1.5% agarose in HEPES-buffered saline solution and sealed with a glass coverslip. VSD was excited using a red LED (627 nm center, Luxeon Star LEDs Quadica Developments Inc., Alberta, Canada) and excitation filter (630±15 nm, Semrock, New York, NY). The VSD reflected signal was passed through an emission filter (688±15 nm, Semrock, New York, NY). Images were taken through a macroscope composed of front-to-front video lenses (8.6×8.6 mm field of view, 67 μm per pixel). The focal plane of the camera was 0.5–1 mm below the cortical surface. The images were captured by a 12-bit charged coupled device (CCD) camera (1M60 Pantera Dalsa, Waterloo, ON) and an EPIX E8 frame grabber with XCAP 3.8 imaging software (EPIX, Inc., Buffalo Grove, IL) at 150 Hz frame rate. For glutamate imaging, iGluSnFR was excited using a blue LED (470nm center, Luxeon Star LEDs Quadica Developments Inc., Alberta, Canada) filtered through an excitation filter (470±20nm, Semrock, New York, NY). The fluorescent signal emitted from cortical activity was collected after passing through an emission filter (542±27nm). Images were taken with the same macroscope camera as described above but at 100 Hz frame rate for recording spontaneous activity and at 150 Hz frame rate for recording evoked activity.

2.2.2 Recording of spontaneous and evoked activity

For anesthetized preparations, spontaneous activity was recorded for at least 20 mins prior to

recording evoked activity as sensory stimulation could alter the characteristics of spontaneous

activity (Han et al., 2008). To stimulate left forelimb and hindlimb paws, a thin needle (0.14 mm)

was inserted into the ventral surface of each paw. Different levels of current were presented

ascendingly to stimulate the paws for 1ms duration (0.1 to 1mA for forelimb and 0.1 to 1.1mA for

hindlimb). The stimulation intensities were categorized into three classes: low (Lo), medium

(Med), and high (Hi) strength as follows: the low and high stimulation levels were identified as

the minimum levels that evoke a detectable VSD response and a saturated response, respectively.

The medium stimulation level was a moderate level between low and high stimulation levels that

generate a response with clear distinction from low and high responses. A 1-ms pulse of combined

blue and green lights was used as visual stimulation. For fore and hind paw stimulation, 10–20

trials of each stimulation intensity level were given with an interstimulus interval of 10s. For visual

stimulation experiments, VSD response distribution were calculated from 100–200 trials of

stimulation with an interstimulus interval of 10s.

For awake preparations, spontaneous activity was recorded similarly as described above. Evoked

activity was recorded by presenting a tone of 12 kHz frequency for a duration of 50ms. The sound

levels for Lo and Hi stimulation levels were 40 and 80 dBSPL respectively. 10-20 trials of

stimulation were presented with an intertrial interval of 10-12s. Tones were generated using a

Tucker-Davis Technologies Inc. (TDT, Alachua, FL, USA) RX6 processor and delivered to

animal's left ear via a free-field ES1 (TDT Inc.) speaker.

2.3 Data analysis

2.3.1 Data preprocessing

All the preprocessing was done in Matlab® (Mathworks, Natick, MA, USA). To do fair

comparisons between evoked responses and spontaneous activity, all recordings (image frames)

were treated using similar preprocessing procedures. As staining of non-flat cortical surface with

voltage-sensitive dye is likely non-uniform, a regional bias in the captured signal is possible. To

overcome this bias, the signal was normalized to a baseline ($\Delta F/F_0$). The baseline was estimated

for both, no-stimulation trials and spontaneous activity, using the "locdetrend" function from the

Chronux toolbox (Bokil et al., 2010). The $\Delta F/F_0$ time series of the VSD signal was then filtered

with a zero-phase lag Gaussian low-pass filter (6 dB attenuation at 25 Hz frequency) while the

glutamate signal was filtered with 6 Hz low-pass filter to eliminate the effects of hemodynamics

that appears at 8-14 Hz. At the end, a two-dimensional spatial Gaussian filter ($\sigma \approx 67\ \mu m$) was

applied. Each evoked response trial was preprocessed separately and then averaged to get the

average response of the corresponding cortical activation and stimulation level.

2.3.2 Optical flow analysis

To capture the spatiotemporal dynamics of VSD and glutamate recordings, we used the Combined

Local-Global (CLG) method implemented in Optical-Flow Analysis Toolbox (Afrashteh et al.,

2017). Using this method, velocity vector fields were estimated for each time point. For each pixel

in a frame, we calculated a velocity vector that depicts the instantaneous speed and direction of

movement of such pixel.

2.3.3 Identification of spontaneous activity motifs

For anesthetized mice, spontaneous activity patterns were compared to patterns of evoked activity in the primary forelimb somatosensory (FLS1), hindlimb somatosensory (HLS1), and visual cortices (VC) while for awake mice, similar comparisons were done for activity in the auditory cortex (AC). Motifs of spontaneous activity were identified using the template matching method used in (Mohajerani et al., 2013). For each sensory modality the template was defined as the set of three frames after the onset of the evoked activity elicited by Hi stimulus level. Then, we calculated the Pearson correlation coefficients (PCC) between such template and spontaneous activity. PCC peaks above 0.4 were considered time points of maximum similarity between evoked and spontaneous activity. The consecutive events with PCC peaks above 0.4 were also screened to be separated by at least 100msec. Frames around these time points were then selected as spontaneous activity events or motifs. To determine the onset of a spontaneous activity motif, local minima were found within 50 frames around the PCC peak and fitted with a cubic polynomial curve (as shown in supplementary Fig. SM1A). The onset of a spontaneous activity motif was considered one frame prior to the intersection of the fitted curve and the original signal. We validated this methodology by estimating stimulus onsets for all trials of evoked activity and comparing with actual values. Most of the differences between estimated and actual onsets were close to 0 (supplementary Fig. SM1B, SM1C, SM1D). The estimation was less accurate for glutamate imaging (supplementary Fig SM1E). For all subsequent analyses, "estimated" onsets for both evoked and spontaneous activity were used. To determine the trajectory of the activity we needed to determine the end of such activity. For this, we considered the activity peak as end point since after this frame the activity starts to diminish. Note that for finding the spontaneous motifs in the forelimb and hindlimb regions, the threshold value for finding PCC peaks (see above) was

defined individually for each animal as the mean PCC between forelimb and hindlimb-evoked

activity (first three frames) for all stimulus intensities (total 9 values). If the mean PCC value was

less than 0.4, 0.4 threshold value was used.

2.3.4 Measurement of amplitude, time-to-peak, and propagation speed and direction

To determine characteristics of evoked and spontaneous activity such as amplitude, and time-to-

peak, speed and direction in a cortical area (e.g. FLS1, HLS1, VC, or AC), a region-of-interest

(ROI) was first defined and time series of $\Delta F/F_0$ for all pixels within the ROI were averaged to

obtain one $\Delta F/F_0$ time series which was used in subsequent analyses. A ROI was defined as the

area for which $\Delta F/F_0$ values are above the mean+SD of all $\Delta F/F_0$ values in that frame. The center

of the PTA ROI was located anatomically as -2mm anterior-posterior and +1.5mm lateral from the

bregma and with the size of $0.5 \times 1 mm^2$. To calculate the amplitude of sensory-evoked responses,

the mean of the baseline was subtracted from the maximum of $\Delta F/F_0$ values within the first 10

frames (67ms for VSD and 100ms for glutamate experiments) after stimulus onset. Similarly, to

calculate the amplitude of spontaneous activity motifs, the mean of baseline activity was subtracted

from the peak $\Delta F/F_0$ within ±15 frames around the peak PCC used for identifying a motif (see

above). For both evoked and spontaneous activity, the baseline was considered as $\Delta F/F_0$ of 10

frames before the onset of activity. The time-to-peak was calculated as the time of the peak minus

the time of the estimated activity onset.

To calculate the propagation speed and direction of the activity from the optical flow analysis, we

averaged the velocity vectors of all pixels in each ROI for each time frame. Speed and direction of

propagation were then determined from the velocity vector with the largest magnitude after the

estimated activity onset and within ±15 frames around the peak of $\Delta F/F_0$ signal. For determining

how stable the directions of propagation were for a spontaneous activity motif, we defined the

propagation direction stability (Figures 7 and S7) and calculated as follows. First, for each frame,

we calculated the cosine of the angles between the average velocity vector at peak speed and the

average velocity vectors in the frames before and after such frame (± 6 and ± 10 frames around the

peak speed frame for VSD and glutamate imaging respectively). Then, we used the average cosine

value to estimate how similar the directions of propagation of the motif to its direction at peak

speed were. Therefore, the propagation direction stability measured this way has values between -

1 and 1 with values closer to 1 indicating that on average, the direction of propagation of

spontaneous activity motif was similar to its direction at peak speed.

2.3.5 Determination of propagation trajectories/paths of activity

To compare the propagation of sensory-evoked and spontaneous activity motifs, trajectories were

calculated starting from the onsets of activity to temporal locations of their peak amplitude (Fig.

SM1F). To estimate the activity trajectory, we first determined the extent of activity (i.e. onset and

offset) and its centroid for each frame using contours identified by $\Delta F/F_0$ values above mean +

m$\times$SD of all $\Delta F/F_0$ values in that frame. The value of "m" was 0.6 and 0.5 for the first and

subsequent frames respectively. Next, one spatial point was selected in each frame to represent the

"location of activity" in that frame. The trajectory of the activity propagation was then defined as

the sequence of such activity locations across frames. In the first frame, the centroid of the activity

contour was selected as the "location of activity". For all subsequent frames, the location of activity

was defined as the intersection of the contour line and the extrapolated directional vector joining

the centroid of activity in the previous frame to centroid of activity in the current frame (Fig.

SM1F).

2.4 Statistical Analysis

2.4.1 Statistical Tests

All statistical analyses were done in Matlab®. Paired sample t-test, Wilcoxon rank sum test, or

repeated measures ANOVA with Greenhouse-Geisser adjustment for the potential lack of

sphericity and post-hoc Tukey-Kramer correction for multiple comparisons were used. We report

F-values, p-values and the estimates of the effect size (partial η^2). Also, in Figures 7 and S7 we

fitted a linear regression model to each of the scatter plots. The linear model coefficients and their

p-values are reported. The type of test used is indicated in the text as well as in figure captions.

Error bars represent the standard error of mean (SEM). *, **, and *** indicate significance levels

of $p < 0.05$, $p < 0.01$, and $p < 0.001$ respectively.

2.4.2 Bootstrap Method to generate Distributions of Hausdorff Fractal Dimension

The Hausdorff fractal dimension measures the roughness (complexity) of a trajectory path.

Trajectories that occupy larger space exhibit higher fractal dimension (Singh et al., 2016). We

determined the Hausdorff fractal dimensions using the box counting method for each trajectory

heat map which encode the percentage of the number of activity that passed through a certain point

on the cortical surface. The heat maps were first binarized using Otsu's method (Otsu, 1979) before

determining fractal dimensions. For each animal and stimulus type (e.g. forelimb stimulation), we

generated a distribution of Hausdorff fractal dimension using bootstrapping over trials from all

levels of stimulus strength. Once we calculated these distributions over all trials from all stimulus

levels, we calculated the trajectory for each stimulation trial. At each iteration of this bootstrapping

method, we randomly selected the corresponding trajectories of 10 trials and their average

trajectory was used to represent the trajectory in this condition. Then, we calculated the fractal

dimension for the average trajectory. This process was done 500 times and we estimated the probability density function (PDF) of the calculated fractal dimension. Analogously, we used the same procedure to estimate the PDF of spontaneous motifs for each animal and stimulus-type motifs.

3 Results

3.1 Response amplitudes of sensory-evoked cortical activity are larger than the amplitudes of motifs of spontaneous cortical activity in both anesthetized and awake mice

We investigated how changes in stimulus strength affect the amplitude of sensory-evoked activity and compared it with amplitudes of spontaneous activity events or motifs (see Methods). In previous studies (Petersen et al., 2003b; MacLean et al., 2005; Han et al., 2008; Luczak et al., 2009), the amplitude of spontaneous activity was found to be comparable to the amplitude of evoked responses. Works on the topographic organization and processing characteristics of different sensory cortices using fMRI, electrophysiology, and optical imaging (Jancke et al., 1998; Shoham et al., 1999; Spenger et al., 2000; Petersen et al., 2003a; Ferezou et al., 2007; Polley et al., 2007; Guo et al., 2012), suggested that increasing the stimulus intensity induces a larger response amplitude at the neuronal network level. Here, we revisited the study of the the relation between the amplitude of spontaneous motifs and evoked activity elicited by multiple stimulus strengths to clarify this discrepancy in both, anesthetized and awake animals.

We used voltage-sensitive dye (VSD) imaging in anesthetized mice and modeled different levels of somatosensory stimulation, low (Lo), medium (Med) and high (Hi), to the fore and hind paws. Different level of stimuli strength used here may mimic the statistics of the sensory input experienced by mice during natural behaviour (Simoncelli, 2003; Carriot et al., 2017). Anaesthesia

provided a long-lasting, stable baseline brain state that was free of active behaviours and voluntary

mental processes that may contaminate the analysis of purely spontaneous activity from which to

make our observations (Musall et al., 2019; Stringer et al., 2019). We measured evoked cortical

responses in the contralateral cerebral hemisphere for multiple trials of forelimb or hindlimb

stimulation. We also recorded spontaneous activity and identified events which were similar to

patterns of cortical activation elicited by forelimb and hindlimb stimulation. These spontaneously

occurring motifs were determined using a template matching method with the templates extracted

from evoked responses elicited by Hi forelimb and hindlimb stimulus strength (see Methods).

Using this approach, we identified the spontaneous activity motifs originated in the primary fore

or hindlimb somatosensory (FLS1 or HLS1) cortices (top rows in Fig. 1A and S1A) and

propagated following different trajectories on the cortical surface. Sensory-evoked activity

increased in amplitude and spread when stimulus intensities were increased (Fig. 1A and

supplementary Fig. S1A). For low level forelimb and hindlimb stimuli, the response was local to

the forelimb and hindlimb areas of the primary and secondary somatosensory cortices respectively,

while for medium and high stimuli, the spatial extent of the evoked response was larger than the

primary forelimb and hindlimb somatosensory areas. Moreover, for forelimb stimuli, after an

initial expansion of the evoked response in the FLS1 area, activity traveled in the medio-caudal

direction passing through the HLS1 area (Fig 1A). Similarly, the spread of the initial hindlimb-

evoked response increased with stronger stimuli (supplementary Fig. S1A).

For a quantitative analysis, we first determined the average value of the VSD signal over time in

an ROI (FLS1 or HLS1) by finding the mean $\Delta F/F_0$ for all pixels within an ROI at each time point.

In this way, time series of VSD signals for an ROI were determined for all trials of evoked activity

and for all motifs of spontaneous activity. The average over trials (evoked-activity) and motifs

(spontaneous activity) for all animals (n = 5) of these VSD signals is shown in Fig. 1B for FLS1

ROI (supplementary Fig. S1B for hindlimb stimulation and HLS1 ROI). It is quite evident that

motifs of spontaneous activity have the lowest $\Delta F/F_0$ values while these values increase with

increasing stimulus levels for evoked activity. Next, we calculated the amplitudes of these signals

by subtracting the mean value over the baseline from the peak $\Delta F/F_0$ (see Fig. 1B) and observed

that the amplitudes of evoked activity increased with the intensity of the stimulus (Fig. 1B-C, and

S1B-C). Note that the mean amplitudes of spontaneous activity motifs were similar to those of

evoked activity elicited with Lo stimulus levels while they were significantly smaller than

amplitudes of evoked activity elicited with Med and Hi stimulus levels (Fig. 1D for forelimb,

repeated measures ANOVA: F-value = 29.075921, p = 0.0000086849, η^2 = 0.879066; post-hoc:

spontaneous vs Med p = 0.018772, spontaneous vs Hi p = 0.0059376; Fig. S1D for hindlimb,

repeated measures ANOVA: F-value = 27.93219, p = 0.0000107044, η^2 = 0.874735; post-hoc:

spontaneous vs Med p = 0.048948, spontaneous vs Hi p = 0.011894). We also quantified the time-

to-peak (time from stimulus onset to the VSD signal maximum) and found that spontaneous

activity motifs were the slowest to reach their peak levels as compared to responses of evoked

activity (Fig. 1E for forelimb, repeated measures ANOVA: F-value = 20.520079, p =

0.0000512563, η^2 = 0.836868; post-hoc: spontaneous vs Lo p = 0.0051397, spontaneous vs Med

p = 0.0067666, spontaneous vs Hi p = 0.011896; Fig. S1E for hindlimb, repeated measures

ANOVA: F-value = 8.992695, p = 0.0021416665, η^2 = 0.692135; post-hoc: spontaneous vs Hi p

= 0.038979). In a separate cohort of anesthetized mice (n = 4), we also compared the amplitudes

and time-to-peak of visual evoked activity (elicited with only one stimulus level) with those of

visual motifs in spontaneous activity (supplementary Fig. S2). The mean amplitude of evoked

activity was significantly larger than that of spontaneous activity events (Fig. S2D, paired sample

t-test p = 0.012257). The time-to-peak for evoked activity was significantly smaller than that for

visual motifs in spontaneous activity (Fig. S2E, paired sample t-test p = 0.0078674).

To compare the amplitudes and temporal dynamics of sensory-evoked and spontaneous activity in

the awake brain state, we used glutamate (Glu) imaging and auditory stimulation with two levels

of sound volume (Lo and Hi). The preprocessing to find Glu $\Delta F/F_0$ values was done in a similar

fashion as for VSD imaging except that Glu signal was filtered using a bandpass filter (see

methods). Spontaneous activity motifs were also determined in a similar fashion as described

above for VSD imaging. The auditory-evoked response originated in the primary auditory cortical

region (AC) and expanded into other cortical regions (montage Fig. 2A). The Glu signal was the

largest for evoked response elicited by Hi stimulus level. The average Glu signal (Fig. 2B; average

over ROI pixels, trials, and animals) was also the largest for evoked activity elicited by Hi stimulus

level. However, average Glu signals for evoked activity elicited by Lo stimulus level was

comparable to average Glu signal for spontaneous activity motifs. Next, we compared the

amplitudes of the Glu signal in a similar way as for VSD signal. The amplitudes of auditory motifs

in spontaneous activity were significantly smaller than those of evoked activity elicited with Hi

stimulus levels (Fig. 2C-D, repeated measures ANOVA: F-value = 19.906767, p = 0.0007837121,

η^2 = 0.832683; post-hoc: spontaneous vs Hi p = 0.0040670) but were similar to those elicited by

Lo stimulus level (p = 0.93913). The time-to-peak for motifs of spontaneous activity was

significantly larger than the time-to-peak for responses of evoked activity with Hi and Lo auditory

stimulus levels (Fig. 2E, repeated measures ANOVA: F-value = 201.184997, p = 0.0000001444,

η^2 = 0.980505; post-hoc: spontaneous vs Lo p = 0.00024736, spontaneous vs Hi p = 0.00030659).

These results, from both anesthetized and awake preparations, thus suggest that the amplitudes of

spontaneous activity motifs are smaller than those of evoked activity and their temporal dynamics

is slower.

3.2 The propagation speed of sensory-evoked cortical activity is larger than that of motifs of

spontaneous cortical activity in anesthetized and awake mice

We next investigated how stimuli of different strengths affects the speed of propagation of evoked

activity waves on the cortical surface and whether the speeds of sensory-evoked activity are larger

than those of spontaneous activity. Given that, behaviorally, animals must attend to salient stimuli

for survival (Corbetta et al., 2008), we hypothesized that the propagation speeds of sensory-evoked

activity would increase with highly salient stimuli and will be higher than the speeds of

spontaneous activity. To address these questions, we quantified the propagation speed of sensory-

evoked activity for forelimb and hindlimb modalities in anesthetized mice with VSD imaging.

Similarly, we quantified the speed of auditory evoked responses in awake mice with glutamate

imaging. To determine velocity vector fields on VSD and glutamate imaging recordings, we

applied optical flow analysis (Afrashteh et al., 2017) to obtain the instantaneous speed and

direction of propagation (Fig. 3A).

For anesthetized mice and somatosensory modalities (forelimb and hindlimb), the instantaneous

speeds increased with stronger stimuli (Fig. 3A and S3A). Instantaneous speeds for forelimb motifs

in spontaneous activity and for responses elicited by Lo stimulus strength, however, were very

small (notice different scale factors used for plotting velocity vector fields in first column in Fig.

3A and S3A). Furthermore, we calculated the average speed over time by finding the magnitude

of the vector sum of the velocity vectors of all pixels within an ROI (e.g. FLS1 or HLS1) for each

time point calculated over the average response of all trials. The average over animals (n = 5) was

then determined and shown in Fig. 3B for FLS1 ROI (and in supplementary Fig. S3B for hindlimb

stimulation and HLS1 ROI). The average speeds were the lowest for spontaneous activity motifs

and increased with increasing stimulus levels for evoked activity. The peak speeds in the

corresponding ROIs were also determined from the time series of average speeds. The distributions

of these peak speeds are shown in Fig. 3C (and in supplementary Fig. S3C for hindlimb

stimulation). Motifs of spontaneous activity had peak speeds similar to those of responses of

evoked activity elicited with Lo stimulus strength. The speeds increased with increasing stimulus

levels (Fig. 3C and 3D) in the FLS1 ROI. The mean peak speed of evoked activity elicited by Hi

and Med stimulus strength was significantly larger than the mean peak speed of evoked activity

elicited by Lo stimulus strength (Fig. 3D, repeated measures ANOVA: F-value = 9.226881, p =

0.0019301139, η^2 = 0.697586; post-hoc: Med vs Lo p = 0.012693, Hi vs Lo p = 0.036275). Since

forelimb evoked activity elicited by Med and Hi stimulus levels originated in the FLS1 ROI but

traveled in the medio-caudal direction on the cortical surface, we also compared propagation

speeds in the HLS1 and posterior tegmental area (PTA) (posterior parietal cortex) ROIs. The

difference in speeds of sensory-evoked with Med and Hi stimulus strengths and spontaneous

activity was significant in the HLS1 and PTA ROIs (Fig. 3E for HLS1, repeated measures

ANOVA: F-value = 19.027242, p = 0.0000743228, η^2 = 0.826293; post-hoc: Med vs spontaneous

p = 0.0068222, Hi vs spontaneous p = 0.036175; 3F for PTA, repeated measures ANOVA: F-value

= 14.780107, p = 0.0159957714, η^2 = 0.787009; post-hoc: Med vs spontaneous p = 0.041526, Hi

vs spontaneous p = 0.038762). Also, for HLS1 and PTA ROIs the mean speed for Med and Hi

strength of sensory forelimb stimulation is significantly higher than the mean speed of propagation

for Lo stimulus level (Fig. 3E for HLS1, Med vs Lo p = 0.0062288, Hi vs Lo p = 0.018329; 3F for

PTA, Med vs Lo p = 0.017250, Hi vs Lo p = 0.036929). For hindlimb stimulation also similar

trend is observed in which Hi stimulus level evokes significantly larger speed compared to Lo and

Med stimulus strength and spontaneous activity in all three ROIs (Fig. S3D for HLS1, repeated

measures ANOVA: F-value = 28.555109, p = 0.0008278413, η^2 = 0.877131; post-hoc: Hi vs Lo p

= 0.0067064, Hi vs Med p = 0.020179, Hi vs spontaneous p = 0.017031; Fig. S3E for FLS1,

repeated measures ANOVA: F-value = 23.909933, p = 0.0000238001, η^2 = 0.856682; post-hoc:

Hi vs Lo p = 0.016317, Hi vs Med p = 0.0078756, Hi vs spontaneous p = 0.025145; Fig. S3F for

PTA, repeated measures ANOVA: F-value = 17.738107, p = 0.0069307567, η^2 = 0.815991; post-

hoc: Hi vs Lo p = 0.026711, Hi vs Med p = 0.027323, Hi vs spontaneous p = 0.044441). We also

performed optical flow analysis for visual cortical stimulation in a separate cohort of anesthetized

mice (n = 4) and found that peak speeds of evoked activity are larger than those of spontaneous

activity motifs (Fig. S4, paired sample t-test p = 0.032798). These results thus suggest that in

anesthetized mice, the propagation speeds for spontaneous activity are comparable to those in

evoked activity elicited with Lo stimulus levels but are smaller than those of evoked activity

elicited with Hi stimulus levels.

Next, we compared the propagation speeds of spontaneous activity with those of sensory evoked

activity in awake mice. The velocity fields for spontaneous and evoked activity at Lo stimulus

strength had vectors with comparable magnitudes while sensory evoked propagations with Hi

stimulus level showed velocity vectors with larger magnitudes (Fig. 4A, see scale factors for

plotting vector fields and compare with scales in Fig. 3A). The average speeds (over trials and

animals) were determined as described above and shown in Fig. 4B. All the graphs show a baseline

speed close to zero and then a rise in speed after the stimulus onset or spontaneous motif onset

which gets to its peak and then declines towards baseline. The distribution of peak speeds

determined from the average speed curves (Fig. 4C) were similar for evoked responses by Lo

stimulus level and spontaneous motifs with no significant difference between means of peak

speeds in AC ROI (Fig. 4D, repeated measures ANOVA: F-value = 37.011210, p = 0.0000904960, η^2 = 0.902466; post-hoc: Lo vs spontaneous p = 0.60796). However, the mean of peak speed for evoked responses elicited by Hi stimulus strength is significantly larger than that of both Lo stimulus level and spontaneous motifs (Fig. 4D, Hi vs Lo p = 0.0028434, Hi vs spontaneous p = 0.0045889). These results thus suggest that in awake mice, the propagation speed of sensory-evoked activity increases with increasing stimulus levels (similar to anesthetized mice). Also, spontaneous activity propagates at a speed similar to the speed of sensory-evoked activity elicited by Lo stimulus level.

3.3   Spatiotemporal propagation patterns of spontaneous activity is more complex than those of sensory-evoked activity in both anesthetized and awake mice

Interaction between neuronal populations is essential for adequate brain function. This interaction often requires brain activity to flow over different brain regions. As demonstrated by wide-field imaging with VSD or glutamate, there is a sequential activation of neuronal populations which have been termed as traveling waves – see review (Muller et al., 2018). We observed propagation patterns of traveling waves and asked whether stimulus levels would affect propagation patterns of sensory-evoked activity and whether these evoked patterns were similar or different from propagation patterns of spontaneous activity. Given that there is a known cortical network responsive to saliency (Downar et al., 2002) and stronger stimulus results in higher saliency (Schultz, 2016), we hypothesized that with increasing stimulus levels, evoked activity would converge to a common propagation pattern. Furthermore, based on the functional roles of spontaneous activity such as memory formation, consolidation, and retrieval, and information processing for carrying out endogenous body functions, we hypothesized that propagation patterns of spontaneous activity would be more complex than those of sensory-evoked activity i.e.

spontaneous activity would have a larger repertoire of propagation patterns. To test these

hypotheses, we performed two types of analyses. First, we analyzed the distributions of directions

at peak values of average speeds of cortical activity in different ROIs using optical flow analysis.

The directions were determined from the average velocity vectors. Second, we calculated the

trajectory space of cortical activity by building normalized histograms of trajectories. For sensory-

evoked and spontaneous activity, the trajectories of responses in all trials and in all motifs were

used respectively.

In anesthetized mice, we observed a gradual convergence to a stereotypical direction of

propagation with increasing stimulus levels for sensory forelimb evoked responses (Fig. 5). The

distributions of directions of velocity vectors of forelimb evoked activity elicited with Lo, Med,

and Hi stimulus levels became tighter and the angles became closer to the medio-caudal direction.

This result was evident for directions observed in the FLS1 ROI (Fig. 5A) but was more prominent

for HLS1 and PTA ROIs (Fig. 5C and 5E respectively). In order to quantify the variability of

directions we determined the magnitude of the average of normalized velocity vectors i.e. first we

normalized all velocity vectors such that their magnitude was 1, then we averaged all these vectors

and calculated the magnitude of the vector sum. A magnitude of the average vector closer to 0

would represent a large variability while a magnitude closer to 1 would represent presence of more

dominant direction. The magnitude of the average vector for Lo, Med, and Hi stimulus levels

increased in the FLS1 ROI (Fig. 5B) suggesting that with increasing stimulus levels, propagation

directions converge to a dominant direction. This was also observed for the HLS1 and PTA ROIs

(Fig. 5D and 5F). This matched our visual observation of forelimb evoked VSD activity where for

Med and Hi stimulus levels the response activity travels abruptly in the medio-caudal direction.

For forelimb motifs in spontaneous activity, although we hypothesized that the activity

propagation would be more complex in terms of distributions of directions (i.e. it would have a

uniform distribution with all angles represented equally), we observed that the distributions were

skewed with the majority of angles representing the caudolateral or lateral direction in the FLS1,

HLS1, and PTA ROIs (Fig. 5A, 5C, and 5E). This suggests that spontaneous activity events might

have a dominant direction. Furthermore, the magnitude of the average vector for spontaneous

events activity shows no significant difference with the magnitude of the average vector for evoked

activity elicited with Lo stimulus level in all three ROIs (Fig. 5B, 5D, and 5F). On the other hand,

this measurement for spontaneous motifs is significantly smaller than that of Med and Hi stimulus

strength (Fig 5D for HLS1, repeated measures ANOVA: F-value = 14.081650, p = 0.0003093654,

$\eta^2 = 0.778781$; post-hoc: Med vs spontaneous p = 0.0066370, Hi vs spontaneous p = 0.0091750;

5F for PTA, repeated measures ANOVA: F-value = 6.166851, p = 0.0088490512, $\eta^2 = 0.606565$;

post-hoc: Hi vs spontaneous p = 0.0011688). This observation also asserts the high similarity

between spontaneous activity and evoked activity at Lo stimulus level. To investigate the structure

of trajectories of evoked and spontaneous activity motifs on the cortical mantle, we analyzed

normalized histograms of activity trajectories represented as heat maps (Fig. 5G). Bright and dim

colors indicate larger and smaller percentage of activity respectively. For forelimb evoked activity,

the trajectory space became smaller with increasing stimulus levels (Fig. 5G for representative

animal). In contrast, the trajectories for forelimb motifs in spontaneous activity occupied a larger

space. For a quantitative comparison of trajectories, we determined the Hausdorff fractal

dimension which measures the roughness of the trajectory path. The Hausdorff fractal dimensions

of heat maps for spontaneous activity events were significantly larger than those of evoked activity

(Fig. 5H, repeated measures ANOVA: F-value = 11.937434, p = 0.0111830599, $\eta^2 = 0.749019$;

post-hoc: Med vs spontaneous p = 0.015937). To compare the trajectory space of spontaneous

activity events with all levels of evoked responses, the distributions of Hausdorff fractal dimension

for both spontaneous motifs and evoked responses pooled from all stimulus intensity levels were

generated using a bootstrapping technique (see methods). Fig. 5I shows these distributions for

individual animals as well as the mean distributions. The Hausdorff fractal dimension distribution

of spontaneous motifs spans to higher values and has higher mean value compared to the one from

evoked responses (Wilcoxon rank sum test on the mean values for animals, $p = 0.047619$, $n = 5$).

A larger Hausdorff fractal dimension corresponds to a larger space occupancy for trajectories

(Singh et al., 2016) and thus spontaneous activity has a larger spatial presence as compared to

sensory-evoked activity. For hindlimb stimulation, the distributions of directions observed in the

HLS1 ROI for Lo, Med, and Hi, stimulus levels were similar with comparable average vector

magnitudes (Fig. S5A and S5B). In the FLS1 and PTA ROIs however, the distributions of

directions converged with increasing stimulus levels suggesting a stereotypical expansion of

hindlimb evoked activity with Hi stimulus strength (Fig. S5C and S5E). The hindlimb motifs in

spontaneous activity, similar to forelimb, had the majority of propagation angles in the caudolateral

and lateral directions in the HLS1, FLS1, and PTA ROIs. The average vector magnitudes tend to

be smaller than evoked responses at all levels (Fig. S5). The Hausdorff fractal dimensions of

trajectory heat maps for hindlimb motifs in spontaneous activity were significantly larger than

those of hindlimb-evoked activity (Fig. S5G and S5H, repeated measures ANOVA: F-value =

8.356785, $p = 0.0028671496$, $\eta^2 = 0.676291$; post-hoc: spontaneous vs Med $p = 0.00083979$, :

spontaneous vs Hi $p = 0.014265$). For the visual sensory modality, we observed a stereotypical

distribution of directions for evoked activity towards the anterior brain regions while spontaneous

activity events traveled in all directions (Fig. S6A and S6B). Therefore, contrary to what was

observed for the forelimb and hindlimb somatosensory modalities, the Hausdorff fractal dimension

here was larger for the evoked activity heat maps (Fig. S6C and S6D, paired sample t-test p =

0.0082663862). These results combined suggest that spontaneous activity has more complex

spatiotemporal propagation patterns with a wider range of angles and larger trajectory spaces than

evoked activity elicited with Hi stimulus levels. Although, the distributions of directions of

velocity vectors suggest that spontaneous activity motifs might have a dominant direction, its

average vector magnitude is similar to Lo stimulus level and is significantly smaller than for higher

stimulus strengths. Alternatively, evoked activity converges to a common propagation pattern with

increasing stimulus levels. Finally, spontaneous motifs occupy a larger trajectory space compared

to evoked responses.

Next, we investigated the propagation patterns of spontaneous and evoked activity in awake mice.

For this analysis, we selected two regions over the mouse cortical surface based on visual

observation of trajectories of auditory stimulation responses. These regions are adjacent to auditory

cortical areas, namely anteromedial to AC ROI (AC-AM) near barrel cortex and posteromedial to

AC ROI (AC-PM) near visual cortex. In the selected ROIs, results were similar to the ones

observed for anesthetized mice. Sensory-evoked activity converged to a propagation pattern with

increasing stimulus levels as suggested by tendency to increase in magnitude of the average of

normalized velocity vectors for Lo and Hi stimulus levels (Fig. 6B, 6D, and 6F). However, the

magnitudes are less than 0.5 (i.e. closer to 0), suggesting a large variability in the distributions of

directions, which is evident in Fig. 6A. This matches our visual observation of auditory-evoked

activity originating and expanding in the AC ROI thus giving rise to velocity vectors in all

directions. In the AC-AM ROI (Fig. 6C and 6D) however, the activity propagation patterns were

not as clear either because the evoked activity does not travel strongly to this ROI or because when

the animals are awake, the evoked activity co-exists with spontaneous activity and becomes less

salient compared to the anesthetized case. For AC-PM ROI, it is interesting to note that evoked

activity with Hi stimulus level has a tight distribution of directions (Fig. 6E). This suggests that

with Hi stimulus strength, evoked activity traveled from the AC to AC-PM ROI in a stereotypical

fashion across animals. For spontaneous activity motifs in auditory region, we observed dominant

directions towards the midline of the cortical surface (Fig. 6A, 6C, and 6E). The magnitude of

average velocity vector is less than 0.5, which also suggests a large variability in the direction of

spontaneous activity events (Fig. 6B, 6D, and 6F). In AC-PM ROI there is no significant difference

between the velocity vectors average for Lo and Hi stimulus levels and spontaneous motifs, but Hi

stimulus level tends to show more stereotyped propagation direction. As described above, here

also, we calculated heat maps to study the structure of cortical activity (Fig. 6G). As observed for

anesthetized mice, the trajectory space occupied by spontaneous activity events was larger than

that occupied by evoked activity, which is confirmed by a significantly larger Hausdorff fractal

dimension for spontaneous activity (Fig. 6H, repeated measures ANOVA: F-value = 28.842795,

$p = 0.0002200288$, $\eta^2 = 0.878208$; post-hoc: spontaneous vs Lo p = 0.0011569, : spontaneous vs

Hi p = 0.0051601). Finally, comparing distributions of Hausdorff fractal dimension for

spontaneous motifs and awake evoked responses pooled from both Lo and Hi stimulus levels

indicate that spontaneous motifs exhibit larger values (Wilcoxon rank sum test on the mean values

for animals, p = 0.047619, n = 5).

Altogether, our results suggest that the propagation patterns of spontaneous activity are more

complex by occupying larger trajectory spaces compared to those of evoked activity in awake

mice.

3.4   Propagation speed of spontaneous activity is positively correlated with its amplitude while

its pattern of propagation and direction stabilize as amplitude increases

As reported earlier, we found that the amplitude of evoked activity is positively correlated with its

speed and also with the stability of its propagation. Here, we asked if the propagation speed,

direction, and pattern of spontaneous activity was also correlated with its amplitude. We found

that for evoked and spontaneous activity in the forelimb and auditory regions, the propagation

speed was positively correlated with its amplitude (scatter plots in Fig. 7A and D where each blue

dot represents a single spontaneous activity motif and each red dot represent a single evoked trial).

The fitted linear regression model showed a positive slope (Fig. 7A, spontaneous: linear

coefficient=1.509 and p-value=0.000057533; evoked: linear coefficient=5.869 and p-

value=3.5064×10^{-11}, and Fig. 7D, spontaneous: linear coefficient=3.437 and p-

value=1.2669×10^{-38}; evoked: linear coefficient=3.431 and p-value=9.8767×10^{-10}). A similar

relationship was also found between propagation direction stability and amplitude (Fig. 7B,

spontaneous: linear coefficient=0.296 and p-value=0.047929; evoked: linear coefficient=0.626

and p-value=0.000019484, and Fig. 7E, spontaneous: linear coefficient=3.437 and p-

value=1.2669×10^{-38}; evoked: linear coefficient=3.431 and p-value=9.8767×10^{-10}). The stability

of propagation pattern measured as the Hausdorff fractal dimension however was more stable with

increasing amplitude for spontaneous activity in the forelimb region compared to auditory region

(Fig. 7C, spontaneous: linear coefficient=0.601 and p-value=0.0046286; evoked: linear

coefficient=-0.101 and p-value=0.31972, and Fig. 7F, spontaneous: linear coefficient=0.002 and

p-value=0.83653; evoked: linear coefficient=0.062 and p-value=0.13361).

Similarly spontaneous activity in the forelimb region show significant positive correlation with

amplitude and speed (Fig. S7A, spontaneous: linear coefficient=0.539 and p-value=0.021671;

evoked: linear coefficient=5.725 and p-value= 0.000027819) stability of propagation direction in the PTA ROI (Fig. S7B, spontaneous: linear coefficient=0.336 and p-value=0.0066435; evoked: linear coefficient=0.412 and p-value=0.025690) and fractal dimension of patterns in the entire imaging field (Fig. S7C, spontaneous: linear coefficient=0.622 and p-value=0.0000023820; evoked: linear coefficient=0.057 and p-value=0.77803). For spontaneous motifs in the visual region and the whole imaging cortical area all three measurements were positively correlated with the amplitude (Fig. S7D, spontaneous: linear coefficient=6.038 and p-value=2.6386×10^{-38}; evoked: linear coefficient=0.057 and p-value=1.4251×10^{-60}. Fig. S7E, spontaneous: linear coefficient=0.389 and p-value= 2.4717×10^{-8}; evoked: linear coefficient=0.210 and p-value= 0.0018673. Fig. S7F, spontaneous: linear coefficient=0.202 and p-value=0.0014744; evoked: linear coefficient=0.110 and p-value=0.017036).

4   Discussion

In this paper, we used voltage-sensitive dye and glutamate imaging in anesthetized and awake mice respectively to characterize cortical dynamics of spontaneous and evoked activity at the mesoscale level. Our main conclusions were similar in anesthetized and awake mice preparations, i.e., sensory-evoked activity and spontaneous activity are similar to each other for low stimulus strength but as the stimulus strength increases, they become different from each other. Previous studies have compared spontaneous and evoked activity using different parameters such as the temporal structure of single-unit activity, unit activity correlations, or dimensionality. In contrast, in this study we use more direct parameters of cortical dynamics such as amplitude, direction and speed of propagation, and complexity of the propagation trajectories. This more direct

characterization of the cortical dynamics might be less influenced by confounding effects

introduced in the methods to calculate such more abstract measurements.

4.1 The effect of stimulus strength on sensory-evoked activity

When the stimulus strength was increased, the amplitude, spread, and speed of the evoked cortical

response increased. These results are consistent with previous studies conducted to understand the

relationship between cortical responsiveness, and the strength of sensory stimulation (Jancke et

al., 1998; Spenger et al., 2000; Peeters et al., 2001; Petersen et al., 2003a; Muniak et al., 2007).

The positive correlation between stimulus strength and amplitude, spread and speed of evoked

cortical activity could be explained in two ways. First, since a larger number of subcortical neurons

are activated with stronger stimulus, these networks, in turn, activate a larger region of cortex

through one-to-one connections (Jones, 1998; Sato et al., 2012). Alternatively, the larger spatial

spread of evoked cortical response in the cortex might be attributed to enhanced activation of

cortico-cortical networks driven by stronger localized subcortical inputs. Further experiments are

required to investigate the subcortical-cortical circuit loops involved in shaping the extent of

cortical activation (Jones, 1998). In addition to the relationship between the extent of activity

propagation and stimulus intensity, our results also suggest a positive correlation between the

speed of the activity propagation and the stimulus strength. However, we found a negative

correlation between the stimulus strength and the variability in the direction of activity

propagation. Similar observations have been reported in psychophysics and electrophysiological

research, where the relationship between stimulus strength or contrast and speed, accuracy or

variability of response are negatively correlated (Palmer et al., 2005; Deco and Hugues, 2012;

Sadagopan and Ferster, 2012). An alternative explanation for observing a larger variability of

directions for weaker stimuli compared to when giving stronger stimulation might be attributed to

a less accurate detection of propagation direction by the optical flow method used in this study.

4.2 Comparison between sensory-evoked and spontaneous activity patterns

Our results indicate that the amplitude of sensory-evoked cortical response is larger than the

amplitude of spontaneous activity events for larger stimulus intensity (Med and Hi) in the

somatosensory and visual cortices of anesthetized mice and auditory cortex of awake mice. A

related comparison has been previously reported for the visual cortex of anesthetized cats and

monkeys using electrophysiological recordings (Nauhaus et al., 2009). In their work, low

contrast stimulus yielded high amplitude LFP waves that travelled longer distances than high-

contrast evoked waves. In addition, it has been reported that in V1 of anesthetized cats, the

amplitude of VSD signals of evoked and spontaneous activity are comparable (Grinvald et al.,

2003). The discrepancy between these studies and our results might be due to the level of stimulus

strength used, or scale (e.g., field of view) at which cortical activity was measured (micro for

electrophysiology vs meso for wide-field VSD), or the origin of the signal (mostly subthreshold

membrane depolarization for VSD vs a combination of subthreshold and suprathreshold activity

for LFP signal) (Grinvald and Hildesheim, 2004).

We also found that on average, sensory-evoked activity travels faster than the spontaneous activity.

On average, evoked activity in the FLS1 area travels at 3.35 ± 0.63 mm/s while forelimb motifs in

spontaneous activity travels at 0.91 ± 0.08 mm/s. For hindlimb stimulation and HLS1 area, evoked

activity travels at 2.37 ± 0.43 mm/s and hindlimb spontaneous motifs travel at 0.88 ± 0.08 mm/s.

Sensory evoked activity in the visual cortex and its spontaneous motifs travels at 4.79 ± 1.88 mm/s

and 1.92 ± 0.55 mm/s respectively. It is important to mention that in previous VSD recordings in

the visual cortex of anesthetized rats, it is reported that spontaneous activity travels faster than

evoked activity (16 mm/s and 10 mm/s respectively) (Han et al., 2008). In contrast, we found that

with our method, this is not the case for the somatosensory cortex, nor for visual cortex. Moreover,

(Nauhaus et al., 2009; Sato et al., 2012) reported that in anesthetized monkeys and cats,

spontaneous activity travels at ~0.3m/s in visual cortex using electrophysiological studies. One

reason for the discrepancy between the speeds reported in these studies and our results might be

the differences in methodology used to quantify the propagation of activity in VSD recordings and

the scale of the neuronal population and brain size from which the activity was measured.

In terms of propagation, sensory-evoked cortical activity has more stereotyped directions of

propagation and trajectories (with higher stimulus strengths) compared to spontaneous activity

which shows a broader distribution of possible directions of activity propagation. We also show

that the trajectories of spontaneous activity sample a broader space than during evoked activity.

4.3 Possible mechanisms for the differences between sensory-evoked and spontaneous activity

Why would evoked activity have larger amplitudes than spontaneous activity? And why would

spontaneous activity travel slower and longer distances than evoked activity? The explanation for

this might lie in the difference in their origin and the particular circuitry of the laminar structure

of the cortex. While spontaneous activity spreads upward from deep layers and slowly across

columns, evoked activity, on the other hand, initiates in thalamocortical recipient layers and

spreads rapidly across columns (Sakata and Harris, 2009; Chauvette et al., 2010; Beltramo et al.,

2013; Petersen and Crochet, 2013).

Since in VSD, most of the signal comes from superficial cortical layers that contain dendritic trees

of neurons located in both layer 2/3 (L2/3) and layer 5 (L5) (Chemla and Chavane, 2010;

Mohajerani et al., 2011), the explanation for the differences reported in this work might be found

by revisiting the input circuitry of L2/3 of the cortex (Sakata and Harris, 2009). According to the

canonical circuit, sensory evoked activity flows into and through the cortex first from the thalamus

to layer 4 (L4), then to L2/3 and finally to L5/6 (Douglas and Martin, 2004; DeNardo et al., 2015).

Although it has been shown that the connectivity of the somatosensory cortex largely follows this

canonical circuit, there is recent evidence that L2/3, L5, and L6 also receive inputs directly from

the thalamus (Viaene et al., 2011; Constantinople and Bruno, 2013; DeNardo et al., 2015). Thus,

the larger amplitude and faster propagation observed during evoked activity could be due to strong

thalamic inputs apart from the already existent cortico-cortical connections used during

spontaneous activity (Bruno and Sakmann, 2006). Finally, a factor for the shorter propagation of

evoked activity compared to spontaneous activity might be due to the inhibition triggered by the

thalamocortical system engaged after the initiation of evoked activity (Sheroziya and Timofeev,

2014; McCormick et al., 2015).

4.4 Functional relevance of differences between sensory-evoked and spontaneous activity

While evoked activity is arguably only involved in sensory processing, spontaneous activity has

multiple functions including memory consolidation, refinement of cortical circuitry, and synaptic

homeostasis. Therefore, the possible realm of computations involved during the communication

amongst different cortical circuits during spontaneous activity might be broader than the ones

involved only in processing sensory information (Luczak et al., 2007). Moreover, the differences

in speed and propagation can be strongly related to the functional roles of spontaneous and evoked

activity in learning and memory (Mercado, 2014). These observations are based on evidence about

the sparse action potential firing of L2/3 and which, in combination with its dense subthreshold

inputs can be exploited for associative learning (Petersen and Crochet, 2013). This sparse action

potential firing of L2/3 might be paired with specific sensory events, top-down input, and

neuromodulatory input to regulate plasticity. Therefore, the temporal differences between

spontaneous and evoked activity, in the order of milliseconds, might be directly related to synaptic

processes. For example, the faster propagation of evoked activity might involve different synaptic

processes to facilitate the encoding of relevant events. On the other hand, the longer propagation

of spontaneous activity might involve the 'replay' of coherent experiences or memories distributed

over large cortical areas for consolidation (Wilson and McNaughton, 1994; Hoffman et al., 2007;

Ji and Wilson, 2007). In contrast, the similarities found between spontaneous activity and sensory-

evoked activity with low stimulus strength at both micro and meso scales, might be due to the

restrictions imposed by the brain circuitry (Luczak and Maclean, 2012) i.e., the underlying

common brain hardware.

This study clarifies the discrepancies in similarities and differences previously reported for

spontaneous and evoked activity. Using wide-field imaging allowed for better resolution in space

than previous electrophysiological studies (Luczak et al., 2009; Sakata and Harris, 2009) and in

time than previous fMRI studies (Jancke et al., 1998; Spenger et al., 2000; Huang et al., 2015) , or

imaging studies over a small cortical area (Han et al., 2008). Spontaneous activity characteristics

including amplitude, propagation speed, and direction distribution is similar to evoked activity

elicited with lower stimulus strength. Also, the trajectory space of spontaneous activity covers a

larger cortical area compared to evoked activity.

4.5 Differences in spontaneous and evoked activity as a biomarker for neuropsychiatric

disorders

Alterations in the balance of excitation and inhibition ratio in the cerebral cortex have been

suggested as an explanation for various neurological and psychiatric disorders such as

schizophrenia (Goel and Portera-Cailliau, 2019). Similarly, the differences in sensory-evoked and

spontaneous activity might be good candidates as objective biomarkers for clinical diagnosis of

neurological and psychiatric disorders. Resting-state spontaneous activity alone measured with

fMRI (Fox and Raichle, 2007; Kiviniemi, 2008) has been used to differentiate between normal

individuals and patients with various neurological diseases (Fox and Greicius, 2010) such as

Alzheimer disease (Zeng et al., 2019), Parkinson's disease (Zhang et al., 2019), schizophrenia

(Nejad et al., 2012), tinnitus (Cai et al., 2019), unilateral amblyopia (Dai et al., 2019), amyotrophic

lateral sclerosis (Luo et al., 2012), idiopathic trigeminal neuralgia (Yuan et al., 2018), and corneal

ulcer (Shi et al., 2019). In addition, combined task-evoked and spontaneous activity fMRI have

been used for pre-operative mapping (Fox et al., 2016). In the above-mentioned studies, three main

types of analysis have been used; 1) regional coherence in which similarity of activity of

neighboring voxels is assessed using cross-correlation, 2) power spectrum analysis in which

amplitudes of low-frequency fluctuations are measured, and 3) spatial pattern analysis in which

the topology of activity is compared between controls and diseased individuals (Fox and Greicius,

2010). These analyses lack focus on the dynamics of activity i.e., speed and propagation. A greater

understanding of the parallel time-varying properties of resting-state activity is increasingly

believed to be essential to understanding brain function in health and disease). We propose the use

of task-independent graded stimuli for recording local evoked activity and its comparison with

corresponding local resting-state spontaneous activity for identifying diseased individuals. This

may have particular utility in revealing the neural circuit dysfunction underlying altered sensory

processing in neuropsychiatric disease contexts. For example, individuals with schizophrenia

exhibit deficits in auditory sensory gating as revealed by the suppression of event related potential

markers P50 and N100 (Brockhaus-Dumke et al., 2008; Javitt, 2009). The methods described in

our study allow the characterization of the dynamics of activity, and therefore, would provide

superior discrimination measurements. Moreover, this approach could be extended to human

studies using fMRI by adapting our methods to 3D version of the optical flow analysis for estimation of activity speed and propagation (von Tiedemann et al., 2010; Rajna et al., 2019).

Disclosures

The authors have no relevant financial interests in the manuscript and no other potential conflicts of interest to disclose.

Acknowledgments: *This work was supported by a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant #40352, Campus Alberta for Innovation Program Chair, Alberta Alzheimer Research Program to MHM and NSERC CREATE in BIP doctoral fellowship to NA. We thank Dr. Jianjun Sun for assistance with surgeries, Behroo Mirza Agha and Di Shao for animal husbandry, and Allen Chan and Timothy H. Murphy for critical reading of the manuscript.*

Author Contribution: Conceptualization, M.H.M., N.A., S.I., and E.B.C., A.L., B.L.M.; Methodology, M.H.M., N.A., S.I., and E.B.C.; Formal Analysis, N.A. and S.I.; Writing - Original Draft, M.H.M., N.A., S.I., and E.B.C.; Writing - Review & Editing, M.H.M., N.A., S.I., A.L., E.B.C., and B.L.M.; Funding Acquisition, M.H.M., B.L.M., A.L.; Resources, M.H.M.; Supervision, M.H.M.

Competing financial interests: The authors declare no competing financial interests.

References:

Afrashteh N, Inayat S, Mohsenvand M, Mohajerani MH (2017) Optical-flow analysis toolbox for characterization of spatiotemporal dynamics in mesoscale optical imaging of brain activity. Neuroimage 153:58-74.

Alloway KD, Johnson MJ, Wallace MB (1993) Thalamocortical interactions in the somatosensory system: interpretations of latency and cross-correlation analyses. Journal of neurophysiology 70:892-908.

Arieli A, Shoham D, Hildesheim R, Grinvald A (1995) Coherent spatiotemporal patterns of ongoing activity revealed by real-time optical imaging coupled with single-unit recording in the cat visual cortex. Journal of neurophysiology 73:2072-2093.

Azzarelli R, Oleari R, Lettieri A, Andre V, Cariboni A (2017) In Vitro, Ex Vivo and In Vivo Techniques to Study Neuronal Migration in the Developing Cerebral Cortex. Brain Sci 7.

Beltramo R, D'Urso G, Dal Maschio M, Farisello P, Bovetti S, Clovis Y, Lassi G, Tucci V, De Pietri Tonelli D, Fellin T (2013) Layer-specific excitatory circuits differentially control recurrent network dynamics in the neocortex. Nat Neurosci 16:227-234.

Berkes P, Orban G, Lengyel M, Fiser J (2011) Spontaneous cortical activity reveals hallmarks of an optimal internal model of the environment. Science 331:83-87.

Bokil H, Andrews P, Kulkarni JE, Mehta S, Mitra PP (2010) Chronux: a platform for analyzing neural signals. J Neurosci Methods 192:146-151.

Brockhaus-Dumke A, Schultze-Lutter F, Mueller R, Tendolkar I, Bechdolf A, Pukrop R, Klosterkoetter J, Ruhrmann S (2008) Sensory gating in schizophrenia: P50 and N100 gating in antipsychotic-free subjects at risk, first-episode, and chronic patients. Biol Psychiatry 64:376-384.

Bruno RM, Sakmann B (2006) Cortex is driven by weak but synchronously active thalamocortical synapses. Science 312:1622-1627.

Butts DA, Goldman MS (2006) Tuning curves, neuronal variability, and sensory coding. PLoS biology 4:e92.

Cai WW, Li Z, Yang Q, Zhang T (2019) Abnormal Spontaneous Neural Activity of the Central Auditory System Changes the Functional Connectivity in the Tinnitus Brain: A Resting-State Functional MRI Study. Front Neurosci 13.

Carriot J, Jamali M, Chacron MJ, Cullen KE (2017) The statistics of the vestibular input experienced during natural self-motion differ between rodents and primates. J Physiol 595:2751-2766.

Chauvette S, Volgushev M, Timofeev I (2010) Origin of active states in local neocortical networks during slow sleep oscillation. Cereb Cortex 20:2660-2674.

Chemla S, Chavane F (2010) A biophysical cortical column model to study the multi-component origin of the VSDI signal. NeuroImage 53:420-438.

Constantinople CM, Bruno RM (2013) Deep cortical layers are activated directly by thalamus. Science 340:1591-1594.

Corbetta M, Patel G, Shulman GL (2008) The reorienting system of the human brain: from environment to theory of mind. Neuron 58:306-324.

Dai P, Zhang J, Wu J, Chen Z, Zou B, Wu Y, Wei X, Xiao M (2019) Altered Spontaneous Brain Activity of Children with Unilateral Amblyopia: A Resting State fMRI Study. Neural Plast 2019:3681430.

Deco G, Hugues E (2012) Neural network mechanisms underlying stimulus driven variability reduction. PLoS Comput Biol 8:e1002395.

Deco G, Jirsa VK, McIntosh AR (2013) Resting brains never rest: computational insights into potential cognitive architectures. Trends Neurosci 36:268-274.

DeNardo LA, Berns DS, DeLoach K, Luo L (2015) Connectivity of mouse somatosensory and prefrontal cortex examined with trans-synaptic tracing. Nat Neurosci 18:1687-1697.

Douglas RJ, Martin KA (2004) Neuronal circuits of the neocortex. Annu Rev Neurosci 27:419-451.

Downar J, Crawley AP, Mikulis DJ, Davis KD (2002) A cortical network sensitive to stimulus salience in a neutral behavioral context across multiple sensory modalities. Journal of neurophysiology 87:615-620.

Faisal AA, Selen LP, Wolpert DM (2008) Noise in the nervous system. Nat Rev Neurosci 9:292-303.

Ferezou I, Haiss F, Gentet LJ, Aronoff R, Weber B, Petersen CC (2007) Spatiotemporal dynamics of cortical sensorimotor integration in behaving mice. Neuron 56:907-923.

Fox MD, Raichle ME (2007) Spontaneous fluctuations in brain activity observed with functional magnetic resonance imaging. Nat Rev Neurosci 8:700-711.

Fox MD, Greicius M (2010) Clinical Applications of Resting State Functional Connectivity. Front Syst Neurosci 4.

Fox MD, Qian T, Madsen JR, Wang D, Li M, Ge M, Zuo HC, Groppe DM, Mehta AD, Hong B, Liu H (2016) Combining task-evoked and spontaneous activity to improve pre-operative brain mapping with fMRI. NeuroImage 124:714-723.

Goel A, Portera-Cailliau C (2019) Autism in the Balance: Elevated E-I Ratio as a Homeostatic Stabilization of Synaptic Drive. Neuron 101:543-545.

Grinvald A, Hildesheim R (2004) VSDI: a new era in functional imaging of cortical dynamics. Nature reviews Neuroscience 5:874-885.

Grinvald A, Arieli A, Tsodyks M, Kenet T (2003) Neuronal assemblies: single cortical neurons are obedient members of a huge orchestra. Biopolymers 68:422-436.

Guo W, Chambers AR, Darrow KN, Hancock KE, Shinn-Cunningham BG, Polley DB (2012) Robustness of cortical topography across fields, laminae, anesthetic states, and neurophysiological signal types. The Journal of neuroscience : the official journal of the Society for Neuroscience 32:9159-9172.

Han F, Caporale N, Dan Y (2008) Reverberation of recent visual experience in spontaneous cortical waves. Neuron 60:321-327.

He BJ (2011) Scale-free properties of the functional magnetic resonance imaging signal during rest and task. The Journal of neuroscience : the official journal of the Society for Neuroscience 31:13786-13795.

Hoffman KL, McNaughton BL (2002) Coordinated reactivation of distributed memory traces in primate neocortex. Science 297:2070-2073.

Hoffman KL, Battaglia FP, Harris K, MacLean JN, Marshall L, Mehta MR (2007) The upshot of up states in the neocortex: from slow oscillations to memory formation. The Journal of neuroscience : the official journal of the Society for Neuroscience 27:11838-11841.

Huang Z, Zhang J, Longtin A, Dumont G, Duncan NW, Pokorny J, Qin P, Dai R, Ferri F, Weng X, Northoff G (2015) Is There a Nonadditive Interaction Between Spontaneous and Evoked Activity? Phase-Dependence and Its Relation to the Temporal Structure of Scale-Free Brain Activity. Cereb Cortex.

Ito T, Brincat SL, Siegel M, Mill RD, He BJ, Miller EK, Rotstein HG, Cole MW (2019) Task-evoked activity quenches neural correlations and variability in large-scale brain systems. bioRxiv.

Jacobs J (2014) Hippocampal theta oscillations are slower in humans than in rodents: implications for models of spatial navigation and memory. Philos Trans R Soc Lond B Biol Sci 369:20130304.

Jancke L, Shah NJ, Posse S, Grosse-Ryuken M, Muller-Gartner HW (1998) Intensity coding of auditory stimuli: an fMRI study. Neuropsychologia 36:875-883.

Javitt DC (2009) Sensory processing in schizophrenia: neither simple nor intact. Schizophr Bull 35:1059-1064.

Ji D, Wilson MA (2007) Coordinated memory replay in the visual cortex and hippocampus during sleep. Nat Neurosci 10:100-107.

Jones EG (1998) Viewpoint: the core and matrix of thalamic organization. Neuroscience 85:331-345.

Jones LA, Smith AM (2014) Tactile sensory system: encoding from the periphery to the cortex. Wiley interdisciplinary reviews Systems biology and medicine 6:279-287.

Karimi Abadchi J, Nazari-Ahangarkolaee M, Gattas S, Bermudez-Contreras E, Luczak A, McNaughton BL, Mohajerani MH (2020) Spatiotemporal patterns of neocortical activity around hippocampal sharp-wave ripples. Elife 9.

Kenet T, Bibitchkov D, Tsodyks M, Grinvald A, Arieli A (2003) Spontaneously emerging cortical representations of visual attributes. Nature 425:954-956.

Kiviniemi V (2008) Endogenous brain fluctuations and diagnostic imaging. Hum Brain Mapp 29:810-817.

Kyweriga M, Mohajerani MH (2016) Optogenetic Approaches for Mesoscopic Brain Mapping. Methods Mol Biol 1408:251-265.

Lim DH, Ledue J, Mohajerani MH, Vanni MP, Murphy TH (2013) Optogenetic approaches for functional mouse brain mapping. Front Neurosci 7:54.

Lim DH, Mohajerani MH, Ledue J, Boyd J, Chen S, Murphy TH (2012) In vivo Large-Scale Cortical Mapping Using Channelrhodopsin-2 Stimulation in Transgenic Mice Reveals Asymmetric and Reciprocal Relationships between Cortical Areas. Front Neural Circuits 6:11.

Luczak A, Maclean JN (2012) Default activity patterns at the neocortical microcircuit level. Front Integr Neurosci 6:30.

Luczak A, Bartho P, Harris KD (2009) Spontaneous events outline the realm of possible sensory responses in neocortical populations. Neuron 62:413-425.

Luczak A, McNaughton BL, Harris KD (2015) Packet-based communication in the cortex. Nat Rev Neurosci 16:745-755.

Luczak A, Bartho P, Marguet SL, Buzsaki G, Harris KD (2007) Sequential structure of neocortical spontaneous activity in vivo. Proc Natl Acad Sci U S A 104:347-352.

Luo C, Chen Q, Huang R, Chen X, Chen K, Huang X, Tang H, Gong Q, Shang HF (2012) Patterns of spontaneous brain activity in amyotrophic lateral sclerosis: a resting-state FMRI study. PLoS One 7:e45470.

MacLean JN, Watson BO, Aaron GB, Yuste R (2005) Internal dynamics determine the cortical response to thalamic stimulation. Neuron 48:811-823.

McCormick DA (1999) Spontaneous activity: signal or noise? Science 285:541-543.

McCormick DA, McGinley MJ, Salkoff DB (2015) Brain state dependent activity in the cortex and thalamus. Curr Opin Neurobiol 31:133-140.

Menon RS, Kim S (1999) Spatial and temporal limits in cognitive neuroimaging with fMRI. Trends in Cognitive Sciences 3:208-215.

Mercado EI (2014) Relating Cortical Wave Dynamics to Learning and Remembering. neuroscience 2014, Vol 1, Pages 185-209.

Mohajerani MH, Aminoltejari K, Murphy TH (2011) Targeted mini-strokes produce changes in interhemispheric sensory signal processing that are indicative of disinhibition within minutes. Proc Natl Acad Sci U S A 108:E183-191.

Mohajerani MH, McVea DA, Fingas M, Murphy TH (2010) Mirrored bilateral slow-wave cortical activity within local circuits revealed by fast bihemispheric voltage-sensitive dye imaging in anesthetized and awake mice. The Journal of neuroscience : the official journal of the Society for Neuroscience 30:3745-3751.

Mohajerani MH, Chan AW, Mohsenvand M, LeDue J, Liu R, McVea DA, Boyd JD, Wang YT, Reimers M, Murphy TH (2013) Spontaneous cortical activity alternates between motifs defined by regional axonal projections. Nat Neurosci 16:1426-1435.

Muller L, Chavane F, Reynolds J, Sejnowski TJ (2018) Cortical travelling waves: mechanisms and computational principles. Nat Rev Neurosci 19:255-268.

Muniak MA, Ray S, Hsiao SS, Dammann JF, Bensmaia SJ (2007) The neural coding of stimulus intensity: linking the population response of mechanoreceptive afferents with psychophysical behavior. The Journal of neuroscience : the official journal of the Society for Neuroscience 27:11687-11699.

Musall S, Kaufman MT, Juavinett AL, Gluf S, Churchland AK (2019) Single-trial neural dynamics are dominated by richly varied movements. Nat Neurosci 22:1677-1686.

Nauhaus I, Busse L, Carandini M, Ringach DL (2009) Stimulus contrast modulates functional connectivity in visual cortex. Nat Neurosci 12:70-76.

Nejad AB, Ebdrup BH, Glenthøj BY, Siebner HR (2012) Brain Connectivity Studies in Schizophrenia: Unravelling the Effects of Antipsychotics. In: Curr Neuropharmacol, pp 219-230.

Omer DB, Fekete T, Ulchin Y, Hildesheim R, Grinvald A (2019) Dynamic Patterns of Spontaneous Ongoing Activity in the Visual Cortex of Anesthetized and Awake Monkeys are Different. Cereb Cortex 29:1291-1304.

Otsu N (1979) Threshold Selection Method from Gray-Level Histograms. Ieee T Syst Man Cyb 9:62-66.

Palmer J, Huk AC, Shadlen MN (2005) The effect of stimulus strength on the speed and accuracy of a perceptual decision. J Vis 5:376-404.

Parga N, Abbott LF (2007) Network model of spontaneous activity exhibiting synchronous transitions between up and down States. Front Neurosci 1:57-66.

Peeters RR, Tindemans I, De Shutter E, Van der Linden A (2001) Comparing BOLD fMRI signal changes in the awake and anesthetized rat during electrical forepaw stimulation. Magnetic Resonance Imaging 19:821-826.

Petersen CC, Crochet S (2013) Synaptic computation and sensory processing in neocortical layer 2/3. Neuron 78:28-48.

Petersen CC, Grinvald A, Sakmann B (2003a) Spatiotemporal dynamics of sensory responses in layer 2/3 of rat barrel cortex measured in vivo by voltage-sensitive dye imaging combined with whole-cell voltage recordings and neuron reconstructions. The Journal of neuroscience : the official journal of the Society for Neuroscience 23:1298-1309.

Petersen CC, Hahn TT, Mehta M, Grinvald A, Sakmann B (2003b) Interaction of sensory responses with spontaneous depolarization in layer 2/3 barrel cortex. Proc Natl Acad Sci U S A 100:13638-13643.

Polley DB, Read HL, Storace DA, Merzenich MM (2007) Multiparametric auditory receptive field organization across five cortical fields in the albino rat. Journal of neurophysiology 97:3621-3638.

Ponce-Alvarez A, He BJ, Hagmann P, Deco G (2015) Task-Driven Activity Reduces the Cortical Activity Space of the Brain: Experiment and Whole-Brain Modeling. PLoS Comput Biol 11:e1004445.

Raichle ME (2010) Two views of brain function. Trends Cogn Sci 14:180-190.

Rajna Z, Raitamaa L, Tuovinen T, Heikkila J, Kiviniemi V, Seppanen T (2019) 3D Multi-Resolution Optical Flow Analysis of Cardiovascular Pulse Propagation in Human Brain. IEEE Trans Med Imaging 38:2028-2036.

Ringach DL (2009) Spontaneous and driven cortical activity: implications for computation. Curr Opin Neurobiol 19:439-444.

Sadagopan S, Ferster D (2012) Feedforward origins of response variability underlying contrast invariant orientation tuning in cat visual cortex. Neuron 74:911-923.

Sakata S, Harris KD (2009) Laminar structure of spontaneous and sensory-evoked population activity in auditory cortex. Neuron 64:404-418.

Sato TK, Nauhaus I, Carandini M (2012) Traveling waves in visual cortex. Neuron 75:218-229.

Schultz W (2016) Dopamine reward prediction-error signalling: a two-component response. Nat Rev Neurosci 17:183-195.

Sellers KK, Bennett DV, Hutt A, Williams JH, Frohlich F (2015) Awake vs. anesthetized: layer-specific sensory processing in visual cortex and functional connectivity between cortical areas. Journal of neurophysiology 113:3798-3815.

Seung HS, Sompolinsky H (1993) Simple models for reading neuronal population codes. ProcNatlAcadSci 90:10749-10753.

Sheroziya M, Timofeev I (2014) Global intracellular slow-wave dynamics of the thalamocortical system. The Journal of neuroscience : the official journal of the Society for Neuroscience 34:8875-8893.

Shi WQ, Wu W, Ye L, Jiang N, Liu WF, Shu YQ, Su T, Lin Q, Min YL, Li B, Zhu PW, Shao Y (2019) Altered spontaneous brain activity patterns in patients with corneal ulcer using amplitude of low-frequency fluctuation: An fMRI study. Exp Ther Med 18:125-132.

Shoham D, Glaser DE, Arieli A, Kenet T, Wijnbergen C, Toledo Y, Hildesheim R, Grinvald A (1999) Imaging cortical dynamics at high spatial and temporal resolution with novel blue voltage-sensitive dyes. Neuron 24:791-802.

Simoncelli EP (2003) Vision and the statistics of the visual environment. Curr Opin Neurobiol 13:144-149.

Singh S, Kaur H, Sandhir R (2016) Fractal dimensions: A new paradigm to assess spatial memory and learning using Morris water maze. Behavioural brain research 299:141-146.

Spenger C, Josephson A, Klason T, Hoehn M, Schwindt W, Ingvar M, Olson L (2000) Functional MRI at 4.7 tesla of the rat brain during electric stimulation of forepaw, hindpaw, or tail in single- and multislice experiments. Exp Neurol 166:246-253.

Stringer C, Pachitariu M, Steinmetz N, Reddy CB, Carandini M, Harris KD (2019) Spontaneous behaviors drive multidimensional, brainwide activity. Science 364:255.

Stringer C, Pachitariu M, Steinmetz NA, Okun M, Bartho P, Harris KD, Sahani M, Lesica NA (2016) Inhibitory control of correlated intrinsic variability in cortical networks. Elife 5.

Viaene AN, Petrof I, Sherman SM (2011) Synaptic properties of thalamic input to layers 2/3 and 4 of primary somatosensory and auditory cortices. Journal of neurophysiology 105:279-292.

von Tiedemann M, Fridberger A, Ulfendahl M, de Monvel JB (2010) Brightness-compensated 3-D optical flow algorithm for monitoring cochlear motion patterns. J Biomed Opt 15:056012.

Wilson MA, McNaughton BL (1994) Reactivation of hippocampal ensemble memories during sleep. Science 265:676-679.

Xie Y, Chan AW, McGirr A, Xue S, Xiao D, Zeng H, Murphy TH (2016) Resolution of High-Frequency Mesoscale Intracortical Maps Using the Genetically Encoded Glutamate Sensor iGluSnFR. The Journal of neuroscience : the official journal of the Society for Neuroscience 36:1261-1272.

Yuan J, Cao S, Huang Y, Zhang Y, Xie P, Fu B, Zhang T, Song G, Yu T, Zhang M (2018) Altered Spontaneous Brain Activity in Patients With Idiopathic Trigeminal Neuralgia: A Resting-state Functional MRI Study. Clin J Pain 34:600-609.

Zeng Q, Luo X, Li K, Wang S, Zhang R, Hong H, Huang P, Jiaerken Y, Xu X, Xu J, Wang C, Zhou J, Zhang M (2019) Distinct Spontaneous Brain Activity Patterns in Different Biologically-Defined Alzheimer's Disease Cognitive Stage: A Preliminary Study. Front Aging Neurosci 11:350.

Zhang C, Dou B, Wang J, Xu K, Zhang H, Sami MU, Hu C, Rong Y, Xiao Q, Chen N, Li K (2019) Dynamic Alterations of Spontaneous Neural Activity in Parkinson's Disease: A Resting-State fMRI Study. Front Neurol 10.

Fig. 1 Amplitude of forelimb-evoked activity is larger than that of spontaneous activity in anesthetized mice. **(A)** Experimental paradigm (left) and montages of VSD imaging (right). Montage top row shows a representative average of forelimb- motifs in spontaneous (Spon) activity (averaged from 10 motifs). Bottom three rows show average evoked (Ev) cortical activity in response to forelimb stimulation with low (Lo), medium (Med), and high (Hi) stimuli strengths (averaged over 10 trials). The primary forelimb region of interest (FLS1-ROI) is outlined in the third column of the bottom row. Compass lines indicate anterior (A), posterior (P), medial (M) and lateral (L) directions. Scale bar is 1 mm. **(B)** Plots of the average VSD signal in the FLS1-ROI across animals (n=5). The lines represent the mean over animals calculated from the averages over trials for individual animals. Thick lines indicate the mean values while shaded regions indicate SEM. **(C)** Average distributions of VSD signal amplitudes in the FLS1-ROI (normalized to the percentage of occurrences). VSD Signal Amplitude = (peak $\Delta F/F_0$ – (mean of baseline) as depicted in (**B**). Shaded regions denote the SEM. **(D-E)** Mean±SEM values of VSD signal peak amplitudes and time-to-peak respectively. * and ** indicate $p<0.05$ and $p<0.01$ respectively repeated measure ANOVA with post-hoc Tukey-Kramer correction.

Fig. 2 Amplitude of tone-evoked activity is larger than that of spontaneous activity in awake mice. **(A)** Experimental paradigm (left) and montages of extracellular glutamate imaging (right). Montage top row shows a representative average motif of auditory spontaneous (Spon) activity motifs. Bottom two rows show average evoked (Ev) cortical activity in response to auditory stimulation with low (Lo) and high (Hi) stimuli strengths. The primary auditory region of interest (AC-ROI) is outlined in the third column of the bottom row. Compass lines indicate anterior (A), posterior (P), medial (M) and lateral (L) directions. Scale bar is 1 mm. **(B)** Plots of the average glutamate signal in the AC-ROI (n=5 animals). The Plots represent average over animals calculated from averages over trials for individual animals. Thick lines indicate the mean values while shaded regions indicate SEM. **(C)** Average distributions of glutamate signal amplitudes in the AC-ROI (normalized to the percentage of occurrences). Glutamate signal amplitude = (peak $\Delta F/F_0$) – (mean of baseline). Shaded regions represent the SEM. **(D-E)** Bar graphs show the mean±SEM values of glutamate signal amplitude peaks and time-to-peak respectively. *, **, and *** indicate $p<0.05$, $p<0.01$, and $p<0.001$ respectively for a repeated measure ANOVA with post-hoc Tukey-Kramer correction.

Fig. 3 Propagation speed of forelimb-evoked activity is larger than that of spontaneous activity in anesthetized mice. **(A)** Montage of representative examples of motifs of forelimb spontaneous (Spon) activity (top row) with overlaid velocity vector fields determined by optical flow analysis. The bottom three rows show similar montages for average evoked (Ev) activity in response to contralateral forelimb stimulation with low (Lo), medium (Med), and high (Hi) stimuli strengths. FLS1 and HLS1 ROIs are outlined in the third column of the bottom row. Compass lines indicate anterior (A), posterior (P), medial (M) and lateral (L) directions. Scale is 1 mm. Numbers in the first column indicate scale factor for drawing velocity vector fields. **(B)** Plots of the average speed signal in the FLS1-ROI (n=5 animals). Note that the scale in the y axis is logarithmical. Shaded regions indicate SEM. **(C)** Average distributions of peak average speeds in the FLS1-ROI (normalized to the percentage of occurrences). Shaded regions represent the SEM. **(D-E)**, and **(F)** Mean±SEM values of peak from average speeds in the FLS1, HLS1, and PTA ROIs respectively. Note that different y-axis scales are used for each ROI. * and ** indicate p<0.05 and p<0.01 respectively for a repeated measure ANOVA test with post-hoc Tukey-Kramer correction.

Fig. 4 Propagation speed of auditory tone-evoked and spontaneous activity in awake mice. **(A)** Montages of representative examples of spontaneous (top row) and averaged evoked activity in response to auditory stimulation with low (Lo) and high (Hi) stimuli strengths (two bottom rows) with overlaid velocity vector fields determined by the optical flow analysis. Primary AC ROI is outlined in the third column of the bottom row. Compass lines indicate anterior (A), posterior (P), medial (M) and lateral (L) directions. Scale bar is 1 mm. Numbers in the first column indicate scale factor for drawing the velocity vector fields. **(B)** Plots of the average speed signal in the AC-ROI (n=5 animals). The lines represent the mean while shaded regions indicate the SEM across animals. **(C)** Average distributions of peak average speed in the AC-ROI from 5 animals (normalized to the percentage of occurrences). Shaded regions show the SEM. **(D)** Mean±SEM values of peak of average speeds for the AC ROI. ** indicate $p<0.01$, repeated measure ANOVA with post-hoc Tukey-Kramer correction.

Fig. 5 Propagation patterns of forelimb-evoked activity converge as stimulus levels increases and are less complex than those of spontaneous activity in anesthetized mice. **(A)** Average distributions of directions of the peak velocity vectors in the FLS1 ROI for spontaneous (Spon) activity and evoked (Ev) activity elicited with Lo, Med, and Hi stimulus levels normalized to the percentage of occurrences across animals (n=5). **(B)** Magnitude of the average of normalized velocity vectors. (**C-D)** and (**E-F)** similar to (**A-B**) but for HLS1 and PTA ROIs respectively. **(G)** Normalized histogram of activity trajectories represented as a heat map with warm and cold colors indicating larger and smaller occurrences of activity passing through a given point on the cortical surface. **(H)** Mean±SEM values of Hausdorff fractal dimension of heat maps. **(I)** Distributions of the Hausdorff fractal dimension for evoked and spontaneous activity for each animal and average for all animals (last column). * and ** indicate $p < 0.05$ and $p < 0.01$ respectively, repeated measure ANOVA with post-hoc Tukey-Kramer correction.

Fig. 6 Propagation patterns of auditory tone-evoked activity converge with increasing stimulus levels and are less complex than those of spontaneous activity in awake mice. **(A)** Average distributions of the directions of the peak velocity vectors in the AC ROI for spontaneous (Spon) activity and evoked (Ev) activity elicited with Lo and Hi stimulus levels normalized to the percentage of occurrences over animals (n=5). **(B)** Magnitude of the average of normalized velocity vectors. **(C** and **D)** and **(E** and **F)** similar to **(A** and **B)** but for AC-AM and AC-PM ROIs respectively. **(G)** Normalized histogram of activity trajectories represented as a heat map with warm and cold colors indicating larger and smaller occurrences of activity passing through a given point on the cortical surface. **(H)** Bar plot of the mean±SEM of the Hausdorff fractal dimension of the heat maps. **(I)** Distributions of the Hausdorff fractal dimension for evoked and spontaneous activity for each animal and average for all animals (last column). ** indicates $p < 0.01$, repeated measure ANOVA with post-hoc Tukey-Kramer correction.

Fig. 7 Speed, flow direction stability, and fractal dimension of the propagation of spontaneous and evoked activity are positively correlated with its amplitude in awake and anaesthetized mice. **(A)** Scatter plot of VSD signal amplitudes for forelimb spontaneous motifs (blue) and evoked (red) activity versus peak velocity vectors in the FLS1 ROI for anaesthetized mice. Each dot represents one trial. The data is pooled from all animals (n=5). The lines represent linear regression models fitted to the corresponding data. **(B)** Scatter plot and linear regression fit for VSD signal amplitude for forelimb spontaneous (blue) motifs and evoked activity (red) in the HLS1 ROI vs flow direction stability for anesthetized mice. **(C)** Similar to **(A)** but for VSD signal amplitudes in the entire imaging window vs fractal dimensions of spontaneous motifs (blue) and evoked activity (red). **(D-F)** Similar to (**A-C**) but for auditory spontaneous motifs (blue) and evoked activity in AC in awake mice. The data is pooled from all animals (n=5). m is the estimated slope. *, **, and *** indicates p<0.05, p<0.01, and p<0.001 respectively, for the p-value of the linear regression slope. Note that speed axis in **A** and **D** is logarithmic scale.