Predicting wildlife hosts of betacoronaviruses for SARS-CoV-2 sampling prioritization

Daniel J. Becker[1,†], Gregory F. Albery[2,†], Anna R. Sjodin[3], Timothée Poisot[4], Tad A. Dallas[5], Evan
A. Eskew[6,7], Maxwell J. Farrell[8], Sarah Guth[9], Barbara A. Han[10], Nancy B. Simmons[11], Michiel
Stock[12], Emma C. Teeling[13], and Colin J. Carlson[14,15,*]

† These authors share lead author status
* Corresponding author: colin.carlson@georgetown.edu

1. Department of Biology, Indiana University, Bloomington, IN, U.S.A.
2. Department of Biology, Georgetown University, Washington, D.C., U.S.A.
3. Department of Biological Sciences, University of Idaho, Moscow, ID, U.S.A.
4. Université de Montréal, Département de Sciences Biologiques, Montréal, QC, Canada.
5. Department of Biological Sciences, Louisiana State University, Baton Rouge, LA, U.S.A.
6. Department of Ecology, Evolution, and Natural Resources, Rutgers University, New Brunswick,
NJ, U.S.A.
7. Department of Biology, Pacific Lutheran University, Tacoma, WA, U.S.A.
8. Department of Ecology & Evolutionary Biology, University of Toronto, Toronto, ON, Canada.
9. Department of Integrative Biology, University of California Berkeley, Berkeley, CA, U.S.A.
10. Cary Institute of Ecosystem Studies, Millbrook, NY, U.S.A.
11. Department of Mammalogy, Division of Vertebrate Zoology, American Museum of Natural
History, New York, NY, U.S.A.
12. KERMIT, Department of Data Analysis and Mathematical Modelling, Ghent University, Belgium
13. School of Biology and Environmental Science, Science Centre West, University College Dublin,
Belfield, Dublin, Ireland.
14. Center for Global Health Science and Security, Georgetown University Medical Center,
Washington, D.C., U.S.A.
15. Department of Microbiology and Immunology, Georgetown University Medical Center,
Washington, D.C., U.S.A.

Abstract.

Despite massive investment in research on reservoirs of emerging pathogens, it remains difficult to rapidly identify the wildlife origins of novel zoonotic viruses. Viral surveillance is costly but rarely optimized using model-guided prioritization strategies, and predictions from a single model may be highly uncertain. Here, we generate an ensemble of eight network- and trait-based statistical models that predict mammal-virus associations, and we use model predictions to develop a set of priority recommendations for sampling potential bat reservoirs and intermediate hosts for SARS-CoV-2 and related betacoronaviruses. We find over 200 bat species globally could be undetected hosts of betacoronaviruses. Although over a dozen species of Asian horseshoe bats (*Rhinolophus* spp.) are known to harbor SARS-like coronaviruses, we find at least two thirds of betacoronavirus reservoirs in this bat genus might still be undetected. Although identification of other probable mammal reservoirs is likely beyond existing predictive capacity, some of our findings are surprisingly plausible; for example, several civet and pangolin species were highlighted as high-priority species for viral sampling. Our results should not be over-interpreted as novel information about the plausibility or likelihood of SARS-CoV-2's ultimate origin, but rather these predictions could help guide sampling for novel potentially zoonotic viruses; immunological research to characterize key receptors (e.g., ACE2) and identify mechanisms of viral tolerance; and experimental infections to quantify competence of suspected host species.

Main text.

Coronaviruses are a diverse family of positive-sense, single-stranded RNA viruses, found widely in mammals and birds[1]. They have a broad host range, a high mutation rate, and the largest genomes of any RNA viruses, but they have also evolved mechanisms for RNA proofreading and repair, which help to mitigate the deleterious effects of a high recombination rate acting over a large genome[2]. Consequently, coronaviruses fit the profile of viruses with high zoonotic potential. There are seven human coronaviruses (two in the genus *Alphacoronavirus* and five in *Betacoronavirus*), of which three are highly pathogenic in humans: SARS-CoV, SARS-CoV-2, and MERS-CoV. These three are zoonotic and widely agreed to have evolutionary origins in bats[3–6].

Our collective experience with both SARS-CoV and MERS-CoV illustrate the difficulty of tracing specific animal hosts of emerging coronaviruses. During the 2002–2003 SARS epidemic, SARS-CoV was traced to the masked palm civet (*Paguma larvata*)[7], but the ultimate origin remained unknown for several years. Horseshoe bats (family Rhinolophidae: *Rhinolophus*) were implicated as reservoir hosts in 2005, but their SARS-like viruses were not identical to circulating human strains[4]. Stronger evidence from 2017 placed the most likely evolutionary origin of SARS-CoV in *Rhinolophus ferrumequinum* or potentially *R. sinicus*[8]. Presently, there is even less certainty in the origins of MERS-CoV, although spillover to humans occurs relatively often through contact with dromedary camels (*Camelus dromedarius*). A virus with 100% nucleotide identity in a ~200 base pair region of the polymerase gene was detected in *Taphozous* bats (family Emballonuridae) in Saudi Arabia[9]; however, based on spike gene similarity, other sources treat HKU4 virus from *Tylonycteris* bats (family Vespertilionidae) in China as the closest-related bat virus[10,11]. Several bat coronaviruses have shown close relation to MERS-CoV, with a surprisingly broad geographic distribution from Mexico to China[12,13,14,15].

Coronavirus disease 2019 (COVID-19) is caused by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), a novel virus with presumed evolutionary origins in bats. Although the earliest cases were linked to a wildlife market, contact tracing was limited, and there has been no definitive identification of the wildlife contact that resulted in spillover nor a true "index case." Two bat viruses are closely related to SARS-CoV-2: RaTG13 bat CoV from *Rhinolophus affinis* (96% identical overall), and RmYN02 bat CoV from *Rhinolophus malayanus* (97% identical in one gene but only 61% in the receptor-binding domain and with less overall similarity)[6,16]. The divergence time between these bat viruses and human SARS-CoV-2 has been estimated as 40-50 years[17], suggesting that the main host(s) involved in spillover remain unknown. Evidence of viral recombination in pangolins has been proposed but is unresolved[17]. SARS-like betacoronaviruses have been recently isolated from Malayan pangolins (*Manis javanica*) traded in wildlife markets[18,19], and these viruses have a very high amino acid identity to SARS-CoV-2, but only show a ~90% nucleotide identity with SARS-CoV-2 or Bat-CoV RaTG13[20]. None of these host species are universally accepted as the origin of SARS-CoV-2 or a progenitor virus, and a "better fit" wildlife reservoir could likely still be identified. However, substantial gaps in betacoronavirus sampling

92   across wildlife limit actionable inference about plausible reservoirs and intermediate hosts for
93   SARS-CoV-2[21].
94
Identifying likely reservoirs of zoonotic pathogens is challenging[22]. Sampling wildlife for the
presence of active or previous infection (i.e., seropositivity) represents the first stage of a pipeline
for proper inference of host species[23], but sampling is often limited in phylogenetic, temporal, and
spatial scale by logistical constraints[24]. Given such restrictions, modeling efforts can play a
critical role in helping to prioritize pathogen surveillance by narrowing the set of plausible
sampling targets[25]. For example, machine learning approaches have generated candidate lists of
likely, but unsampled, primate reservoirs for Zika virus, bat reservoirs for filoviruses, and avian
reservoirs for *Borrelia burgdorferi*[26–28]. In some contexts, models may be more useful for
identifying which host or pathogen groups are *unlikely* to have zoonotic potential[29]. However,
these approaches are generally applied individually to generate predictions. Implementation of
multiple modeling approaches collaboratively and simultaneously could reduce redundancy and
apparent disagreement at the earliest stages of pathogen tracing and help advance modeling
work by addressing inter-model reliability, predictive accuracy, and the broader utility (or
inefficacy) of such models in zoonosis research.
109
Because SARS-like coronaviruses (subgenus *Sarbecovirus*) are only characterized from a small
number of bat species in publicly available data, current modeling methods are poorly tailored to
exactly infer their potential reservoir hosts. In this study, we instead conduct two predictive efforts
that may help guide the inevitable search for known and future zoonotic coronaviruses in wildlife:
(1) broadly identifying bats and other mammals that may host any *Betacoronavirus* and (2)
specifically identifying species with a high viral sharing probability with the two *Rhinolophus*
species carrying the closest known wildlife relatives of SARS-CoV-2. To do this, we developed a
standardized dataset of mammal-virus associations by integrating a previously published
mammal-virus dataset[30] with a targeted scrape of all GenBank accessions for Coronaviridae and
their associated hosts. Our final dataset spanned 710 host species and 359 virus genera,
including 107 mammal hosts of betacoronaviruses as well as hundreds of other (non-
coronavirus) association records. We integrated our host-virus data with a mammal phylogenetic
supertree[31] and over 60 ecological traits of bat species[27,32,33]. We used these standardized data
to generate eight predictive models of host-virus associations, including four network-based
approaches, three trait-based approaches, and one hybrid approach using network and trait
information. These efforts generated eight ranked lists of suspected bat hosts of
betacoronaviruses and five ranked lists for other mammals. Each ranked list was scaled
proportionally and consolidated in an ensemble of recommendations for betacoronavirus
sampling and broader eco-evolutionary research (Supplemental Figure 1).
129
In our ensemble, we draw on two popular approaches to identify candidate reservoirs and
intermediate hosts of betacoronaviruses. *Network-based methods* estimate a full set of "true"
unobserved host-virus interactions based on a recorded network of associations (here, pairs of
host species and associated viral genera). These methods are increasingly popular as a way to

identify latent processes structuring ecological networks[34−36], but they are often confounded by
sampling bias and can only make predictions for species within the observed network (i.e., those
that have available virus data; in-sample prediction). In contrast, *trait-based methods* use
observed relationships concerning host traits to identify species that fit the morphological,
ecological, and/or phylogenetic profile of known host species of a given pathogen and rank the
suitability of unknown hosts based on these trait profiles[28,37]. These methods may be more likely
to recapitulate patterns in observed host-pathogen association data (e.g., geographic biases in
sampling, phylogenetic similarity in host morphology), but they more easily correct for sampling
bias and can predict host species without known viral associations (out-of-sample prediction).

Predictions of bat betacoronavirus hosts derived from network- and trait-based approaches
displayed strong inter-model agreement within-group, but less with each other (Figure 1A,B). In-
sample, we identified bat species across a range of genera as having the highest predicted
probabilities of hosting betacoronaviruses, distributed in distinct families in both the Old World
(e.g., Hipposideridae, several subfamilies in the Vespertilionidae) and the New World (e.g.,
Artibeus jamaicensis from the Phyllostomidae; Figure 1C). Out-of-sample, our multi-model
ensemble more conservatively limited predictions to primarily Old World families such as
Rhinolophidae and Pteropodidae (Figure 1D). Of the 1,037 bat hosts not currently known to host
betacoronaviruses, our models identified between 1 and 720 potential hosts based on a 10%
omission threshold (90% sensitivity). Applying this same threshold to our ensemble predictions,
we identified 239 bat species that are likely undetected hosts of betacoronaviruses. These
include more half of bat species in the genus *Rhinolophus* not currently known to be
betacoronavirus hosts (35 of 61), compared to 16 known hosts in this genus. Given known roles
of rhinolophids as hosts of SARS-like coronaviruses, our results suggest that SARS-like
coronavirus diversity could be undescribed for around two-thirds of the potential reservoir bat
species.

Our multi-model ensemble predicted undiscovered betacoronavirus bat hosts with striking
geographic patterning (Figure 2). In-sample, the top 50 predicted bat hosts were broadly
distributed and recapitulated observed patterns of bat betacoronavirus hosts in Europe, parts of
sub-Saharan Africa, and Southeast Asia, although our models also predicted greater-than-
expected richness of likely bat reservoirs in the Neotropics and North America. In contrast, the
top out-of-sample predictions clustered in Vietnam, Myanmar, and southern China.

Because only trait-based models were capable of out-of-sample prediction, the differences in
geographic patterns of our predictions likely reflect distinctions between the network- and trait-
based modeling approaches, which we suggest should be considered qualitatively different lines
of evidence. Network approaches proportionally upweight species with high observed viral
diversity, recapitulating sampling biases largely unrelated to coronaviruses (e.g., frequent
screening for rabies lyssaviruses in vampire bats, which have been sampled in a comparatively
limited capacity for coronaviruses[14,38−40]). Highly ranked species may also have been previously
sampled without evidence of betacoronavirus presence; for example, *Rhinolophus luctus* and

Macroglossus sobrinus from China and Thailand, respectively, tested negative for
betacoronaviruses, but detection probability was limited by small sample sizes[41–43]. In contrast,
trait-based approaches are constrained by their reliance on phylogeny and ecological traits, and
the use of geographic covariates made models more likely to recapitulate existing spatial
patterns of betacoronavirus detection (i.e., clustering in southeast Asia). However, their out-of-
sample predictions are, by definition, inclusive of unsampled hosts[44], which potentially offer
greater return on viral discovery investment.

Multi-model ensemble predictions also clustered taxonomically along parallel lines. Applying a
graph partitioning algorithm (phylogenetic factorization) to the bat phylogeny[45], we found that in-
sample predictions were on average lowest for the Yangochiroptera (Figure 3). This makes
intuitive sense, because this clade does not include the groups known to harbor the majority of
betacoronaviruses detected in bats (e.g., *Rhinolophus*, Hipposideridae). Out-of-sample
predictions were lower in the New World superfamily Noctilionoidea, the emballonurids, and the
Lasiurus genus, whereas the *Rhinolophus* genus and the Old World fruit bats (Pteropodidae) both
had higher mean probabilities of betacoronavirus hosting (Supplemental Table 1).

These clade-specific patterns of predicted probabilities across extant bats could be particularly
applicable for guiding future surveillance. On the one hand, betacoronavirus sampling in
southeast Asian bat taxa (especially the genus *Rhinolophus*) may have a high success of viral
detection but may not improve existing bat sampling gaps[46]. On the other hand, discovery of novel
betacoronaviruses in Neotropical bats or Old World fruit bats could significantly revise our
understanding of the bat-virus association network. Such discoveries would be particularly
important for global health security, given the surprising identification of a MERS-like virus in
Mexican bats[14] and the likelihood that post-COVID pandemic preparedness efforts will focus
disproportionately on Asia despite the near-global presence of bat betacoronaviruses.

Although our ensemble model of potential bat betacoronavirus reservoirs generated strong and
actionable predictions, our mammal-wide predictions were largely uninformative. In particular,
minimal inter-model agreement (Supplemental Figure 2) indicated a lack of consistent,
biologically meaningful findings. Major effects of sampling bias were apparent from the top-
ranked species, which were primarily domestic animals or well-studied mesocarnivores (ED
Figure 2B). Phylogenetic factorization mostly failed to find specific patterns in prediction
(Supplemental Table 2): in-sample, mean predictions primarily confirmed betacoronavirus
detection in the remaining Laurasiatheria (e.g., ungulates, carnivores, pangolins, hedgehogs,
shrews), although nested clades of marine mammals (i.e., cetaceans) were less likely to harbor
these viruses. Although cetaceans are predicted to be susceptible to SARS-CoV-2 based on ACE2
similarity with humans[47], this result is expected given betacoronavirus epidemiology and their
predominance in terrestrial mammals. Our mammal predictions thus reflect a combination of
detection bias and poor performance of network methods on limited data that likely signals the
limits of existing predictive capacity. Our dataset contained only 30 non-bat betacoronavirus
hosts, many of which were identified during sampling efforts following the first SARS outbreak[7].

Although the laurasiatherians include more potential intermediate hosts than other mammals,
likely driven by inclusion of bats in this group, the high species diversity of this clade restricts
insights for sampling prioritization, experimental work, or spillover risk management.
221
Given the unresolved origins of SARS-CoV-2 and significant motivation to identify other SARS-like
coronaviruses and their reservoir hosts for pandemic preparedness[21], we further explored our
only model that could generate out-of-sample predictions for all mammals[48]. This model uses
geographic distributions and phylogenetic relatedness to estimate viral sharing probability.
Where one or more (potential) hosts are known, these sharing patterns can be interpreted to
identify probable reservoir hosts[48]. Because *Rhinolophus affinis* and *R. malayanus* host viruses
that are closely related to SARS-CoV-2[6,16], we used their predicted sharing patterns to identify
possible reservoirs of sarbecoviruses. In doing so, we aimed to work around a major data
limitation: fewer than 20 sarbecovirus hosts were recorded in our dataset, a sample size that
would preclude most modeling approaches.
232
For both presumed bat host species of sarbecoviruses, the most probable viral sharing hosts
were again within the Laurasiatheria. Although bats—especially rhinolophids—unsurprisingly
assumed the top predictions given phylogenetic affinity with known hosts (Supplemental Table
3, Supplemental Figure 3), several notable patterns emerged in the rankings of other mammals.
Pangolins (Pholidota) were disproportionately likely to share viruses with *R. affinis* and *R.*
malayanus (Supplemental Figure 4); the Sunda pangolin (*Manis javanica*) and Chinese pangolin
(*M. pentadactyla*) were in the top 20 predictions for both reservoir species (Supplemental Table
4). This result is promising given the much-discussed discovery of SARS-like betacoronaviruses
in *M. javanica*[18]. The Viverridae were also disproportionately well-represented in the top
predictions (Supplemental Figure 5), most notably the masked palm civet (*Paguma larvata*), which
was identified as an intermediate host of SARS-CoV[49,50] (Supplemental Table 4). Our virus sharing
model thus captured historic patterns of betacoronavirus spillover and predicted cross-species
transmission from only the phylogeography of east Asian mammals. The opportunity for inter-
species contact depends on both animal behavior and habitat selection, which could be better
studied in natural and anthropogenic habitats for these clades[51]. However, over evolutionary
timescales, our results suggest recombination and diversification of novel betacoronaviruses
should be expected among such taxa in nature, without necessarily requiring further contact via
the wildlife trade. For example, recent observations in Gabon demonstrate cohabitation between
pangolins, bats, and various other mammals (e.g., rodents) in burrows[52].
252
Moreover, these findings lend credibility to other predictions of SARS-CoV-2 sharing patterns and
host susceptibility. Many of the model's top predictions were mustelids (i.e., ferrets and weasels),
and the most likely viral sharing partner for both *Rhinolophus* species was the hog badger
(*Arctonyx collaris*; Supplemental Table 4). Taken together with reports of SARS-CoV-2 spread in
mink farms[53], these results highlight the relatively unexplored potential for mustelids to serve as
betacoronavirus hosts and as models for infectivity studies[54]. Similarly, identification of several
deer and Old World monkey taxa as high-probability hosts in our clade-based analysis

(Supplemental Figure 3) meshes with the observation of high binding of SARS-CoV-2 to ACE2
receptors in cervid deer and primates[47]. Felids (especially leopards) also ranked relatively high in
our viral sharing predictions (Supplemental Table 4, Supplemental Figure 5), which is of particular
interest given reports of SARS-CoV-2 susceptibility among cats[55]. However, we caution that this
model was the only approach in our ensemble that could generate out-of-sample prediction
across mammals, and therefore its predictions lacked confirmation (and filtering of potential
spurious results) by other models that were designed and implemented independently.
267
Several limitations apply to our work, most notably the difficulty of empirically verifying
predictions. Although some virological studies have incidentally tested specific hypotheses (e.g.,
filovirus models and bat surveys[27,56], henipavirus models and experimental infections[23,57]), model-
based predictions are nearly never subject to systematic verification or post-hoc efforts to identify
and correct spurious results. Greater dialogue between modelers and empiricists is necessary to
systematically confront the growing set of predicted host-virus associations with experimental
validation or field observation.
275
Already, we have identified roughly a half-dozen new bat reservoirs of betacoronaviruses that our
study correctly predicted. *Scotophilus heathii*, *Hipposideros larvatus*, and *Pteropus lylei*, all highly
predicted bat species in our out-of-sample rankings, have been reported positive for
betacoronaviruses in the literature[43,58]; however, resulting sequences were not annotated to
genus level in GenBank. More recently, the release of 630 novel bat coronaviruses sequences
from China included four additional bat hosts (*Hipposideros pomona, Scotophilus kuhlii, Myotis
pequinius,* and *M. horsfieldii*)[59]. All but *M. horsfieldii* were correctly identified by the ensemble, and
three models correctly identified all seven verification hosts (Trait-based 1 and 3 and Network-
based 1). These results support the idea that our models identified relevant targets correctly but
also highlight an evident limitation of the workflow. Whereas an automated approach was the
ideal method to systematically compile over 30,000 samples on the timescales commensurate
with ongoing efforts to trace SARS-CoV-2 in wildlife, we suggest this discrepancy highlights the
need for careful virological work downstream at every stage of the modeling process, including
the development of hybrid manual-automated data pipelines.
290
Additionally, overcoming underlying model biases that are driven by historical sampling regimes
will require coordinated efforts in field study design. Bat sampling for betacoronaviruses has
prioritized viral discovery[39,40,60–62], but limitations in the spatial and temporal scale (and
replication) of field sampling have likely created fundamental gaps in our understanding of
infection dynamics in bat populations[24]. Limited longitudinal sampling of wild bats suggests
betacoronavirus detection is sporadic over time and space[58,63], implying strong seasonality in
virus shedding pulses[64]. Carefully tailored spatial and temporal sampling efforts for priority taxa
identified here, within the *Rhinolophus* genus or other high-prediction bat clades, will be key to
identifying the environmental drivers of betacoronavirus shedding from wild bats and possible
opportunities for contact between bats, intermediate hosts, and humans.
301

Future field studies will undoubtedly be important to understand viral dynamics in bats but are
inherently costly and labor-intensive. These efforts are particularly challenging during a pandemic,
as many scientific operations have been restricted including field studies of bats in some regions
to limit possible viral spillback from humans. However, various alternative efforts could both
advance basic virology and allow testing model predictions. General open access to viral
association records, including GenBank accessions and the upcoming release of the USAID
PREDICT program's data, could answer open questions and allow updates to our sampling
prioritization (including potentially modeling at subgenus level, with greater data availability).
Appropriately preserved museum specimens and historical collections, from diverse research
programs, also offer key opportunities to retrospectively screen samples from bats and other
mammals for betacoronaviruses and to enhance our understanding of complex host-virus
interactions[65]. Large-scale research networks, such as GBatNet (Global Union of Bat Diversity
Networks; https://gbatnet.blogspot.com) and its member networks, could provide diverse
samples and ensure proper partnerships and equitable access and benefit sharing of knowledge
across countries[66,67]. Whole-genome sequencing through initiatives such as the Bat1K Project
(https://bat1k.ucd.ie) would facilitate fundamental and applied insights into the immunological
pathways by which bats can harbor many virulent viruses without displaying disease[68,69].
Additionally, targeted sequencing efforts could also identify endogenized viral elements in bat
genomes, shedding light on the diversity of bat viruses and the evolution of bat immune
systems[70,71].

To expedite such work, we have made our binary predictions of host-virus associations for all
eight models and all 1,000+ bat species publicly available (Supplementary File 1). Such results
are provided both in the spirit of open science and with the hope that future viral detection,
isolation, or experimental studies might confirm some of these predictions or rule out others[57]. In
ongoing collaborative efforts, we aim to consolidate results from field studies that address these
predictions (e.g., serosurveys) and to track Genbank submissions to expand the known list of
betacoronavirus hosts. In several years, we intend to revisit these predictions as a post-hoc test
of model validation, which would represent the first effort to test the performance of such models
and assess their contribution to basic science and to pandemic preparedness.

It is crucial that our predictions be interpreted as a set of hypotheses about potential host-virus
compatibility rather than strong evidence that a particular mammal species is a true reservoir for
betacoronaviruses. In particular, susceptibility is only one aspect of host competence[22,72], which
encompasses the diverse genetic and immunological processes that mediate within-host
responses following exposure[73]. SARS-CoV-2 in particular may have a broad host range[47], given
hypothesized compatibility with the ACE2 receptor in many mammal species, but this only adds
to the extreme caution with which any data should be used to implicate a potential wildlife
reservoir of the virus, given that rapid interpretation of inconclusive molecular evidence has likely
already generated spurious reservoir identifications[74,75]. Future efforts to isolate live virus from
wildlife or to experimentally show viral replication would more robustly test whether predicted
host species actually play a role in betacoronavirus maintenance in wildlife[57].

Without direct lines of virological evidence, we note that our sampling prioritization scheme also
does not implicate any given mammal species in SARS-CoV-2 transmission to humans. Care
should be taken to communicate this, especially given the potential consequences of
miscommunication for wildlife conservation. The bat research community in particular has
expressed concern that negative framing of bats as the source of SARS-CoV-2 will impact public
and governmental attitudes toward bat conservation[76]. In zoonotic virus research on bats, studies
often over-emphasize human disease risks[77] and rarely mention ecosystem services provided by
these animals [78]. Skewed communication can fuel negative responses against bats, including
indiscriminate culling (i.e., reduction of populations by selective slaughter)[79], which has already
occured in response to COVID-19 even outside of Asia (where spillover occurred)[80].

To minimize potential unintended negative impacts for bat conservation, public health and
conservation responses should act in accordance with substantial evidence suggesting that
culling has numerous negative consequences, not only threatening population viability of
threatened bat species in shared roosts[81] but also possibly increasing viral transmission within
the very species that are targeted[82,83]. Instead, bat conservation programs and long-term
ecological studies are necessary to help researchers understand viral ecology and find
sustainable solutions for humans to live safely with wildlife. From another perspective, policy
solutions aimed at limiting human-animal contact could potentially prevent virus establishment
in novel species (e.g., as observed in mink farms[53]), especially in wildlife that may already face
conservation challenges (e.g., North American bats threatened by an emerging disease, white-
nose syndrome[79,84]). At least two bat species that can be infected by the fungal pathogen
(*Eptesicus fuscus* and *Tadarida brasiliensis*) are in our list of the 239 bat species most likely to be
betacoronavirus hosts.

Substantial investments are already being planned to trace the wildlife origins of SARS-CoV-2.
However, the intermediate progenitor virus may never be isolated from samples
contemporaneous with spillover, and it may no longer be circulating in wildlife. MERS-CoV
circulates continuously in camels[85] and SARS-CoV persisted in civets long enough to seed
secondary outbreaks[49,50], but the limited description of Pangolin-CoV symptoms suggests high
mortality, potentially indicating a more transient epizootic such as Ebola die-offs in red river hogs
(*Potamochoerus porcus*)[18]. In lieu of concrete data, our study provides no additional evidence
implicating any particular species—or any particular pathway of spillover (e.g., wildlife trade,
consumption of hunted animals)—as more or less likely. No specific scenario can be confirmed
or rigorously interrogated by ecological models, and we explicitly warn against misinterpretation
or misuse of our findings as evidence for adjacent policy decisions. Although policies that focus
on particular potential reservoir species or target human-wildlife contact could reduce future
spillovers, they will have a negligible bearing on the ongoing pandemic, as SARS-CoV-2 is highly
transmissible within humans (e.g., unlike MERS-CoV or other zoonoses that are sustained in
people by constant reintroduction). SARS-CoV-2 is likely to remain circulating in human
populations at least until a vaccine is developed (if not much longer, given current challenges to

vaccine production, allocation, governance, and uptake), regardless of immediate actions
regarding wildlife. COVID-19 response must be informed by the best consensus evidence
available and prioritize solutions that address immediate reduction of transmission through
public health and policy channels. Meanwhile, we hope our proposed wildlife sampling priorities
will help increase the odds of preventing the future emergence of novel betacoronaviruses.
391

Acknowledgements

We thank Heather Wells for generously sharing thoughtful comments and code. The VERENA consortium is supported by L'Institut de Valorisation de Données (IVADO) through Université de Montreal. DJB was supported by an appointment to the Intelligence Community Postdoctoral Research Fellowship Program at Indiana University, administered by Oak Ridge Institute for Science and Education through an interagency agreement between the U.S. Department of Energy and the Office of the Director of National Intelligence. MS was supported by the Research Foundation - Flanders (FWO17/PDO/067) and the Flemish Government under the "Onderzoeksprogramma Artificiële Intelligentie (AI) Vlaanderen" program.

 Figures

 Figure 1. An ensemble of predictive models facilitates identification of likely betacoronavirus
 bat hosts. The pairwise Spearman's rank correlations between models' ranked species-level
 predictions were generally substantial and positive (A,B). Models are arranged in decreasing
 order of their mean correlation with other models. In-sample predictions, expressed as host
 species' proportional rank (0 is the most likely host from a given model, 1 is the least likely
 host), varied significantly due to the uncertainty of network approaches (C). In contrast, species'
 proportional ranks were tightly correlated across out-of-sample predictive approaches, which
 relied on species traits (D). Each line represents a different bat species' proportional rank
 across models. The ten species with the highest mean proportional ranks across all models are
 highlighted in shades of purple.

Figure 2. Species richness of known and suspected betacoronavirus bat hosts. Known hosts of betacoronaviruses (*top*) are found worldwide, but particularly in southern Asia and southern Europe. The top 50 predicted bat hosts with viral association records (*middle*) are mostly Neotropical, including several species of vampire bats. In contrast, the top 50 *de novo* bat host predictions based on phylogeny and ecological traits (*bottom*) are mostly clustered in Myanmar, Vietnam, and southern China, with none in the Neotropics or North America.

Figure 3. Phylogenetic distribution of predicted bat and mammal hosts of betacoronaviruses.
Bar height indicates mean predicted rank across the model ensemble (higher values = lower
proportional rank score, more likely to be a host) and black indicates known betacoronavirus
hosts. Colored regions indicate clades identified by phylogenetic factorization as significantly
different in their predicted rank compared to the paraphyletic remainder; those clades more
likely to contain a host are shown in red, whereas those less likely to contain a host are shown
in blue. Results are displayed for bats and all mammals separately, stratified by in- and out-of-
sample predictions. Numbers reference clade names, species richness, and mean predicted
ranks as described in Supplemental Tables 1 and 2.

Methods.

The underlying conceptual aim of this study was to produce and synthesize several different
models that predict and rank candidate reservoir species—each with different methods,
assumptions, and framings—and to rapidly synthesize these into a consensus list. We broadly
structured our study around two modeling targets: (1) produce rankings of likely bat hosts of
betacoronaviruses and (2) identify potential non-bat mammal hosts. We developed a novel
dataset that merged existing knowledge about the broader mammal-virus network with targeted
data collection about coronaviruses; implemented eight modeling methods; synthesized these
into an ensemble; and post-hoc identified taxonomic patterns in prediction using phylogenetic
factorization.

Host-Virus Association Data

Entries were downloaded from GenBank on March 27th 2020 using the following search terms:
Coronavirus, Coronaviridae, Orthocoronavirinae Alphacoronavirus, Betacoronavirus,
Gammacoronavirus, and Deltacoronavirus. Data were sorted using a Python script that saved all
available metadata regarding accession number, division, submission date, entry title, organism,
genus, genome length, host classification, country, collection date, PubMed ID, journal containing
associated publication, publication year, genome completeness, and the gene sequenced. The
dataset was cleaned to remove duplicate entries, using GenBank accession number, and entries
that did not correspond to viral sequences, using GenBank division. After cleaning, 31,473 entries
remained, of which 25,628 had metadata regarding host species.

Data from GenBank were merged with the Host-Pathogen Phylogeny Project (HP3) dataset[30]. The
HP3 dataset consists of 2,805 associations between 754 mammal hosts and 586 virus species,
compiled from the International Committee on Taxonomy of Viruses (ICTV) database, and
manually cleaned over a period of five years. Data collection on HP3 began in 2010 and has been
static since 2017, but it still represents the most complete dataset on the mammal virome
published with a high standard of data documentation. Several recent studies have used the HP3
dataset to produce statistical models of viral sharing or zoonotic potential[29,48,86], making it a
comparable reference for a multi-model ensemble study.

Because of naming inconsistencies both within GenBank and between the two datasets (HP3 and
GenBank), we used a two-step pipeline for taxonomic reconciliation. Viral names were matched
to the ICTV 2019 master species list, up to the sub-genus level. Host species names were
matched against GBIF using their species API with an automated Julia script, and processed to a
fully cleaned set of names. This led to an harmonized dataset representing a global list of
mammal-virus associations, from which the bat-coronavirus data can be extracted for
downstream and specific modeling efforts. Because the HP3 dataset used an older version of the
ICTV master list, and because not all host names in the GenBank metadata could be matched by

the GBIF species API (or could be solved unambiguously to the species level), some host-virus
interactions were lost; this reinforces the need to careful data curation of taxonomic metadata if
they are to enable and support predictive pipelines.
483
484
485   **Predictor Data**
486
487   *Phylogeny*
488
We used a supertree of extant mammals to unify modeling approaches incorporating host
phylogeny[31]. Although more recent mammal supertrees exist, we used this particular phylogeny
for consistency with trait datasets and several of the modeling frameworks included in our
ensemble. We manually matched select bat species names between our edge list and this
particular phylogeny. This included reverting any *Dermanura* to their former *Artibeus* designation
(i.e., *D. phaeotis*, *D. cinerea*, *D. tolteca*)[87], switching *Tadarida* species to either *Mops* or *Chaerophon*
species (i.e., *Tadarida condylura* to *Mops condylurus*, *Tadarida plicata* to *Chaerephon plicatus*,
Tadarida pumila to *Chaerephon pumilus*)[88], and renaming *Myotis pilosus* to the more recent *Myotis*
ricketti. *Chaerephon pusillus* was considered its own species but is now synonymous with
Chaerephon pumilus[88]. Minor discrepancies between virus data and our phylogeny were also
corrected (*Hipposideros commersonii* to *Hipposideros commersoni* [although more recently
changed to *Macronycteris commersoni*], *Rhinolophus hildebrandti* to *Rhinolophus hildebrandtii*,
Neoromicia nana to *Neoromicia nanus*). In other cases, some recently revised genera in our edge
list were modified to match former genera in the mammal supertree: *Parastrellus hesperus* to
Pipistrellus hesperus, and *Perimyotis subflavus* to *Pipistrellus subflavus*[89]. Lastly, some names in
our edge list missing from the mammal supertree represent former subspecies being raised to
full species rank, and names were reverted accordingly: *Artibeus planirostris* to *Artibeus*
jamaicensis, *Miniopterus fuliginosus* to *Miniopterus schreibersii*, *Triaenops afer* to *Triaenops*
persicus, and *Carollia sowelli* to *Carollia brevicauda*. Although we recognize that these are each
now recognized as distinct species, in all cases our synonymized names are thought to be either
sister taxa or very closely related.
510
511   *Ecological traits*
512
We used a previously published dataset of 63 ecological traits describing the morphology, life
history, biogeography, and diet of 1,116 bat species. These data are drawn from a combination
of PanTHERIA[32], EltonTraits[33], and the IUCN Red List range maps, and were previously cleaned in
a study producing predictions of bat reservoirs of filoviruses[27]. Four redundant variables (two for
human population density, mean potential evapotranspiration in range, and body mass) were
eliminated prior to analyses, favoring variables with higher completeness.
519
520   *Correction for sampling bias*
521

522 To correct for sampling bias, in the style of several previous studies[30,86], we used the number of
523 peer-reviewed citations available on a given host as a measure of scientific sampling effort. We
524 used the R package *easyPubMed* to scrape the number of citations in PubMed returned when
525 searching each of the 1,116 bat names in the trait data on April 10, 2020.
526
527 **Modeling Approaches**
528
Our team produced an ensemble of eight statistical models (Supplemental Tables 5 and 6), and
applied them to generate a predictive set of eight models for bats and five for other mammals.
Four use a network-theoretic component (k-nearest neighbors, linear filtering, trait-free plug-and-
play, and scaled phylogeny), while three primarily used ecological traits as predictors (boosted
regression trees, Bayesian additive regression trees, and neutral phylogeographic). A final hybrid
model used a combination approach with a network backbone but incorporating the phylogeny
and bat trait data.
536
All eight approaches were used to generate predictions about potential bat hosts of
betacoronaviruses. A subset of five were used to recommend potential non-bat mammal hosts
of betacoronaviruses (*k*-nearest neighbor, linear filtering, scaled phylogeny, trait-free plug-and-
play, and neutral phylogeographic). We did not use trait-based models to predict non-bat hosts,
because assigning pseudoabsences to the vast majority (~3500 or more) of mammal species
would likely lead to largely uninformative predictions, weighed against the 109 known
betacoronavirus hosts (79 bats and 30 other mammals).
544
545 *Network model 1: k-nearest neighbors recommender*
546
We follow the methodology previously developed for the recommendation of species feeding
interactions[90]. This method builds a recommender system internally based on the *k*-nearest
neighbor (kNN) algorithm, under which candidate hosts are recommended for a virus from a pool
constituted by the hosts of the *k* viruses with which it has the greatest overlap. Overlap (host
sharing) is measured using Tanimoto similarity, which is the cardinality of the intersection of two
sets divided by the cardinality of their union. To obtain the pairwise similarity between two viruses,
this divides the number of shared hosts by the cumulative number of hosts. The *k*NN of a virus
are the *k* other viruses with which it has the highest Tanimoto similarity.
555
Hosts are then recommended by counting how many times they appear in these *k* neighbors, a
quantity that ranges from 1 to *k*. We can impose arbitrary cutoffs by limiting the
recommendations to the hosts that occur in at least *k*, *k-1*, etc, viruses. Previous leave-one-out
validation of this model revealed that it is particularly effective for viruses with a reduced number
of hosts, which is likely to be the case for emerging viruses. Furthermore, the performance of this
model was not significantly improved by the addition of functional traits, making it acceptable to
run on the association data only.
563

This model was run two times: first, by measuring the similarity of viruses, and recommending
hosts; second, by measuring the similarity of hosts, and recommending viruses. In all cases, only
results for betacoronaviruses are reported.

The outcome of this model should be subject to caution, as leave-one-out validation revealed that
the success rate (i.e., ability to recover one interaction that has been removed) remained lower
than 50% even when using k=8, and dropped as low as 5% when using k=1 (the nearest-neighbor
algorithm). This strongly suggests that the dataset of reported host-virus associations is
extremely incomplete; therefore, the identification of the nearest neighbors can be biased by
under-reported interactions, and this can result in noise in the prediction. This noise can be
particularly important when the kNN technique operates on viruses, of which the bat dataset has
only 15.

Network-based model 2: Linear filter recommender

Following Stock *et al.*[91], we used a previously developed linear filter to infer potential missing
interactions. This recommender system assumes that networks tend to be self-similar, and use
this information to generate a score for an un-observed interaction that is a linear combination of
the status of the interaction (relative weight of 1/4), relative degree of host and virus, and of the
observed connectance of the network (all with relative weights of 1); as we are concerned with
ranking interactions as opposed to examining the absolute value of the score, the penalization
coefficient associated to the interaction being presumed absent could be omitted with no change
in the ranking, but has been set to a low value instead. The scores returned by the linear filter are
not directly related to the probability of the interaction existing in this context, but higher scores
still indicate interactions that are more likely to exist. Indeed, known hosts of betacoronavirus
typically scored higher.

We used the zero-one-out approach to assess the performance of this model on the entire
datasets. In all cases, non-interactions ranked lower than positive interactions even when entirely
removing the penalization coefficient from the linear filter parameters, which suggests that the
network structure (degree and connectance) is capturing a lot of information as to which species
can interact.

Network-based model 3: Plug and play

For network problems, the "plug and play" model is a statistical approach that formulates Bayes'
theorem for link prediction around the conditional density of traits of known associations
compared to traits of every possible association in a network. The conditional density function is
measured by using non-parametric kernel density estimators (implemented with the R package
np), and the conditional ratio between them is used to estimate link "suitability", a scale-free ratio.
Compared to other machine learning methods that fit training data iteratively, plug and play is
comparatively simple, and directly infers the most likely extensions of observed patterns in data.

The plug and play was originally developed to forecast missing links in host-parasite networks[36],
but has since been used to model species distributions[92] and predict the global spread of human
infectious diseases[93]. We used this model here to estimate suitability of host-virus interactions
by first modeling the entire estimated network of host-virus interaction suitability, and ranking
hosts that are not infected by betacoronaviruses by their estimated suitability for
betacoronaviruses.

The "plug and play" model is trained using either matched pairs of host and pathogen ecological,
morphological, or phylogenetic traits[36], or by using a latent approach[93] which considers the mean
similarity of pathogens in their host ranges and the mean similarity of hosts in their pathogen
communities as 'traits'. We decided to use the latent approach, as host trait data was far more
available than viral trait data. Further, the taxonomic scale considered for host (species) and virus
(genus) differed, making the resolution of potential trait data different enough to potentially
confound trait-based approaches in this modeling framework.

Relative suitability of a host-virus association, as estimated by the "plug and play" model, is
formulated as a density ratio estimation problem. The suitability of a host-virus association is
quantified as the quotient of the distribution of latent trait values when an association was
recorded over the distribution of all the latent trait values. As an attempt to control for sampling
effort of mammal and bat host species, we included PubMed citation counts for host species (as
described above) in the estimation of host-virus suitability. We explored host-pathogen suitability
using the entire mammal-virus associations dataset, to maximize the available information on
the network's structure, and ranked host-pathogen pairs by their relative suitability value. From
the final predictions, we subset out bat-specific predictions. When predicting, we set citation
counts to the mean of training data, as a sampling bias correction.

Network-based model 4: Scaled-phylogeny

We apply the network-based conditional model of Elmasri *et al.*[94] for predicting missing links in
bipartite ecological networks. The full model combines a hierarchical Bayesian latent score
framework which accounts for the number of interactions per taxon, and a dependency among
hosts based on evolutionary distances. To predict links based on evolutionary distance, the
probability of a host-parasite interaction is taken as the sum of evolutionary distances to the
documented hosts of that parasite. This allocates higher probabilities when a few closely related
hosts, or many distantly related hosts interact with a parasite. In this way phylogenetic distances
are combined with individual affinity parameters per taxa to model the conditional probability of
an interaction.

In ecological studies, it is common to use time-scaled phylogenies to quantify evolutionary
distance among species[95]. We may use these fixed evolutionary distances for link prediction, but
parasite taxa are known to be more or less constrained by phylogenetic distances among hosts[96].
Further, phylogenies are hypotheses about evolutionary relationships and have uncertainties in

the topology and relative distances among species[97]. Rather than treating phylogenetic distances
as fixed, Elmasri *et al.*[94] re-scale the phylogeny by applying a macroevolutionary model of trait
evolution. While any evolutionary model that re-scales the covariance matrix may be used, we use
the early-burst model, which allows evolutionary change to accelerate or decelerate through
time[98]. This different emphasis to be placed on deep versus recent host divergences when
predicting links.
654
We apply the model to a network of associations among host species and viral genera, and the
mammal supertree, which allows us to leverage information from across the network to predict
undocumented bat-betacoronavirus associations. We fit sets of models, applying both the full
model, and the phylogeny-only model to both the bat-viral genera associations, and the mammal-
viral genera associations. For each data-model combination we fit the model using ten-fold cross-
validation holding out links for which there is a minimum of two observed interactions. The
posterior interaction matrices resulting from each of the ten models are then averaged to
generate predictions for all links in the network, with betacoronaviruses subset to generate the
ensemble predictions.
664
To assess predictive performance, we attempted to predict the held out interactions, and
calculated AUC scores by thresholding predicted probabilities per fold, and taking an average
across the 10 folds. In addition to AUC, we also assessed the model based on the percent of
documented interactions accurately recovered. For the bat-viral genera data the full model
resulted in an average AUC of 0.82 and recovered an average of 90.1% of held out interactions,
while the phylogeny-only model showed increased AUC (0.86), but a decreased proportion of held-
out interactions recovered (84.5%). Interestingly, the models for bat-virus genera associations
had marginally worse predictive performance compared to the same models run on the larger
network of mammal-virus associations (full model: AUC 0.88, 84.4% positive interactions
recovered; phylogeny-only model:AUC: 0.88, 88.8% positive interactions recovered), indicating
that predicting bat-betacoronavirus associations may benefit from including data on non-bat
hosts. The models also estimated the scaling parameter (eta) of the early-burst model to be
positive (average eta=7.92 for the full model run on the bat subset), indicating accelerating
evolution compared to the input tree (Supplemental Figure 6). This means that recent divergences
are given more weight than deeper ones for determining bat-viral genera associations, which is
consistent with recent work on viral sharing[48,99].
681
682 *Trait-based model 1: Boosted regression trees*
683
Previous work has been highly successful in predicting zoonotic reservoirs using a combination
of taxonomic, ecological, and geographic traits as predictors. This approach has been previously
used to identify wildlife hosts of filoviruses[27,100], flaviviruses[28,101], henipaviruses[23], *Borrelia*
burgdorferi[26], to predict mosquito vectors of flaviviruses[102], and to predict rodent reservoirs and
tick vectors of zoonotic viruses[37,103]. These approaches treat the presence of a specific virus (or
genus of viruses) or a zoonotic pathogen as an outcome variable, with negative values given for

species not known to be hosts (pseudoabsences), and use machine learning to identify the
characteristics that predispose animals to hosting pathogens of concern. By predicting the
probability that a given pseudoabsence is a false negative, the method can infer potential
undetected or undiscovered host species.
694
This approach has almost exclusively been implemented using boosted regression trees (BRT),
a classification and regression tree (CART) machine learning method that became popular a
decade ago for species distribution modeling.[104] Boosted regression trees develop an ensemble
of classification trees which iteratively explain the residuals of previous trees, up to a fixed tree
depth (usually between 3 and 5 splits). The incorporation of boosting allows the model, as it is fit,
to progressively better explain poorly-fit cases within training data.
701
We used boosted regression trees to identify trait profiles that predict bat hosts of
betacoronaviruses, including all trait predictors from the trait database that met baseline
coverage (< 50% missing values) and variation (< 97% homogenous) thresholds. For all model
fitting, we specified a Bernoulli error distribution for our binary response variable and applied 10-
fold cross validation to prevent overfitting (R package *gbm*). We started by fitting a global model
to our full dataset, first specifying learning rate = 0.01 (shrinks the contribution of each tree to the
model) and tree complexity = 4 (controls tree depth) as per default values and subsequently
tuning to minimize cross-validation error.
710
711 We reduced the variable set by calling the gbm.simp() function, which computes and compares
712 the mean change in cross-validation error (deviance) produced by dropping different sets of least-
713 contributing predictors. The final simplified model included 23 variables, plus citation counts,
714 which we added to correct for sampling bias.
715
We applied bootstrapping resampling methods to estimate uncertainty, using our tuned model to
fit 1000 replicate models. For each model, training sets were assembled by randomly selecting
with replacement 79 bat-coronavirus associations from the set of reported bat hosts and 79
pseudoabsences. Trained models were used to generate relative influence coefficients for trait
predictors and coronavirus host probabilities across all bat species. Partial dependence plots
display relative influence coefficients and bootstrapped confidence intervals for the top ten
contributing trait predictors. The medians of host probabilities were ranked and used to identify
the top ten candidate host species. When predicting, we set citation counts to the mean of training
data, as a sampling bias correction.
725
726 *Trait-based model 2: Bayesian additive regression trees*
727
A similar workflow to trait-based model 1 was implemented using Bayesian additive regression
trees (BART), an emerging machine learning tool that has similarities to more popular methods
like random forests and boosted regression trees. BART adds several layers of methodological
innovation, and performs well in bakeoffs with other advanced machine learning methods.

Several features make BART very convenient for modeling projects like these, including several
easy-to-use implementations in R packages, built-in capacity to impute and predict on missing
data, and easy construction of variable importance and partial dependence plots.
735
Like other classification and regression tree methods, BART assigns the probability of a binary
outcome variable by developing a set of classification trees - in this case, a sum-of-trees model -
that split data ("branches") and assign values to terminal nodes ("leaves"). Whereas other similar
methods generate uncertainty by adjusting data (e.g. random forests bootstrap training data and
fit a tree to each bootstrap; boosted regression trees are usually implemented with iterated
training-test splits to generate confidence intervals), BART generates uncertainty using an MCMC
process. An initial sum-of-trees model is fit to the entire dataset, and then rulesets are adjusted
in a limited and stochastic set of ways (e.g., adding a split; switching two internal nodes), with the
sum-of-trees model backfit to each change. After a burn-in period, the cumulative set of sum-of-
trees models is treated as a posterior distribution. This has some advantages over other methods,
like boosted regression trees or random forests. In particular, posterior width directly measures
model uncertainty (rather than approximating it by permuting training data), and a single model
can be run (instead of an ensemble trained on smaller subsets of training data), allowing the
model to use the full training dataset all at once.[105]
750
Unlike many Bayesian machine learning methods, BART is easily implemented out-of-the-box, due
to a limited set of customization needs. Three main priors control the fitting process: one usually-
uniform prior on variable importance, one two-parameter negative power distribution on tree
depth (preventing overfitting), and an inverse chi-squared distribution on residual variance. A set
of well-performing priors from the original BART study[106] are widely used across R
implementations for out-of-the-box settings, but can be further adjusted relative to modeling
needs. In this study, we implemented BART models using a Dirichlet prior for variable importance
(DART), a specification that is designed for situations with high dimensionality data that probably
reflects a small number of true informative predictors. This often produces a much more reduced
model without going through a stepwise variable selection process, which can be slow and very
subject to stochasticity.[105]
762
763 We implemented this approach using the *BART* package in R, using the bat-virus association
764 dataset to generate an outcome variable, and the bat traits dataset as predictors. BART models
765 were implemented with 200 trees and 10,000 posterior draws, using every trait feature that was
766 at least 50% complete and < 97% homogenous (taken from TBM1).
767
We tried four total implementations, based on two decisions: BART uncorrected and corrected
for citation counts (BART-u, BART-c), and DART uncorrected and corrected for citation counts
(DART-u, DART-c). All four models performed well, with little variation in predictive power
measured by the area under the receiver operator curve calculated on training data (BART-u: AUC
= 0.93; BART-c: AUC = 0.93; DART-u: AUC = 0.93; DART-c: 0.90; Supplemental Figure 7). Across
all models, spatial variables had a high importance, including some regionalization (extent of

range) and some variables capturing larger geographic range sizes, as did a diet of invertebrates
(pulling out the phylogenetic signal of insectivorous bats; Supplemental Figure 8).

All models identified a number of "false negative" hosts that would be suitable based on a 10%
false negative classification threshold for known betacoronavirus hosts (implemented with the R
package 'PresenceAbsence'). BART-u identified 217 missing hosts, BART-c identified 279
missing hosts, DART-u identified 222 missing hosts, and DART-c identified 384 missing hosts,
suggesting that this model most penalized overfitting as intended. As a result, we considered this
model the most rigorous and powerful for inference, and used DART-c in the final model
ensemble. We predicted across all 1,040 bats without recorded betacoronavirus associations,
and ranked predicted probability. When predicting, we set citation counts to the mean of training
data, as a sampling bias correction.

Trait-based model 3: Phylogeographic neutral model

We used a previously published pairwise viral sharing model[48] to predict potential
betacoronavirus hosts based on the sharing patterns of known hosts in a published dataset [30].
We used a generalised additive mixed model (GAMM), which was fitted in the first half of 2019
using the *mgcv* package, with pairwise binary viral sharing (0/1 denoting if a species shares at
least one virus) as a response variable. Explanatory variables include pairwise proportional
phylogenetic distance and geographic range overlap (taken from the IUCN species ranges), with
a multi-membership random effect to control for species-level sampling biases. The model was
then used to predict the probability that a given species pair share at least one virus across 4196
placental mammals with available data, producing a predicted viral sharing network that
recapitulates a number of known macroecological patterns, as well as predicting reservoir hosts
with surprising accuracy[48]. Subsetting this predicted sharing matrix, we listed the rank order of
hosts most likely to share with all known betacoronavirus hosts in our datasets.

Rhinolophus-specific implementation of trait-based model 3

We then repeated this process with sharing patterns of *Rhinolophus affinis* and *R. malayanus*
specifically. Given the strong phylogenetic effect, the top 139 predictions were bat species:
predominantly rhinolophids and hipposiderids. The top 20 predictions for both *R. malayanus* and
R. affinis are displayed in Supplemental Tables 3 and 4. Notable predictions included the hog
badger *Arctonyx collaris* (Carnivora: Mustelidae), which was examined for SARS-CoV antibodies
in 2003 and is reported in wildlife markets[7,107]; a selection of civet cats (Carnivora: Viverridae)
including *Viverra* species; the binturong (*Arctitis binturong*); and the masked palm civet (*Paguma
larvata*), the latter of which were implicated in the chain of emergence for SARS-CoV[49,50]; and
pangolins (Pholidota: Manidae) including *Manis javanica* and *Manis pentadactyla,* which have
been hypothesised to be part of the emergence chain for SARS-CoV-2[18,19].

Alongside these high-ranked species-level predictions, we visually examined how predictions
varied across all mammal orders and families using the whole dataset (Supplemental Figure 5).
Pangolins (Pholidota), treeshrews (Scandentia), carnivores (Carnivora), hedgehogs
(Erinaceomorpha), and even-toed ungulates (Artiodactyla) had high mean predicted probabilities.
Investigating family-level sharing probabilities revealed that civets (Viverridae) and mustelids
(Mustelidae) were responsible for the high Carnivora probabilities, and mouse deer (Tragulidae)
and bovids (Bovidae) were mainly responsible for high probabilities in the Artiodactyla
(Supplemental Figure 5).
823
824 *Hybrid model 1: Two-Step Kernel Ridge Regression*
825
To make predictions using both the traits and the phylogeny of the hosts, we used Two-Step
Kernel Ridge Regression (TSKRR), a kernel-based pairwise learning method[108,109]. This method
can predict interactions between two species, i.e., hosts and viruses, both in-sample and out-of-
sample. Conceptually, TSKRR performs nonlinear regression twice: once to generalize to new
hosts and once to generalize to new viruses (Supplemental Figure 9A). As a kernel method, it can
exploit arbitrary similarity measures to describe the species. At the same time, its conceptual
simplicity permits using efficient shortcuts for tuning and specialized cross-validation.
833
For the hosts, we again restricted ourselves to bats. First, we inputted missing trait values using
the *MissForest* package[110]. We then selected the relevant traits, standardized them, and
computed the squared Euclidean distance. We plugged this distance into a standard radial basis
kernel[111], where we used the median heuristic to set the bandwidth. Similarly, for the bat
phylogeny, we computed the distance matrix. We combined this distance with the radial basis
kernel, again setting the bandwidth by the median of the distances. Bats were described using a
kernel matrix derived from traits, phylogeny, or the average of both (Supplemental Figure 9B). We
corrected for sampling bias by including the number of citations per species as a covariate and
predicted using the average number of citations for all hosts. For viruses, we used a simple kernel
with a "1" if two virus genera belong to the same family, and a "0" elsewise. For all the kernel
matrices, we added 0.1 to each element and an additional 0.1 to the diagonal elements. This
modification serves as an intercept and it gives the TSKRR the same capacity as the linear filtering
method described earlier[108].
847
We fitted the models using the R package *xnet*. We assessed model performance in two ways.
First, we used leave-one-out for every element of the incidence matrix and computed the AUC
over all predicted interactions. Secondly, we used leave-one-out cross-validation on bats(i.e.,
leaving out a single species, making predictions for all the viruses of that bat, and repeating this
for every species). In this setting, we computed the AUC for every virus over all bats and averaged
these values (Supplemental Figure 9C). For both evaluation strategies, we removed viruses with
no hosts in the dataset, although we kept these for training the model (Supplementary Table 7).
Compared to the average performance for the viruses, betacoronaviruses were slightly more
difficult to predict. Phylogeny was slightly more informative than traits, although the effect

between them was synergistic. As would be expected, it was easier to predict missing interactions than to predict new hosts.

To assess which variables are most important, we computed feature importance based on random permutations. We randomly reshuffled one or both of the bats' kernel matrices, destroying its information content. Subsequently, we recorded the decrease in AUC for the setting 'interactions' in the model as an indication of relative importance. We also randomly reshuffled each trait separately and monitored the effect on performance. We repeated each random permutation 100 times. Reshuffling the complete kernel matrix with both traits and phylogeny resulted in the largest average drop in AUC of 0.1904 (Supplementary Figure 10). Reshuffling only the phylogeny kernel matrix had a much more profound impact than the trait matrix (0.07769 vs. 0.05301). Contributions of traits were smaller by comparison. Citation count was the largest effect (0.02119), with diet breadth (0.00520), population group size (0.003802), and annual birth pulse (0.0022944) as the remaining largest contributions. Removing citations resulted in a stronger importance for these variables but did not change the relative importance.

Consensus Methods and Recommendations

Combining and ranking predictions

For all eight models predicting bat hosts of betacoronaviruses, and five models predicting mammal hosts of betacoronaviruses, we combined predictions—generated using the same standardized data—into one standardized dataset. All mammal models were trained on data including bats, but predictions were subset to exclude bats to focus on likely intermediate hosts.

Each study's unique output—a non-intercomparable mix of different definitions of suitability or probability of association—were transformed into proportional rank, where lower rank indicates higher evidence for association out of the total number of hosts examined. By rescaling all results to proportional ranks between zero and one, we also allowed comparison of in-sample and out-of-sample predictions across all models. Proportional ranks were averaged across models to generate one standardized list of predictions. This absorbed much of the variation in model performance (Supplemental Figure 1) and produced a set of rankings that performed well.

We elected not to withhold any "test" data to measure model performance, given that each method deployed in the ensemble has been independently and rigorously tested and validated in previous publications. Instead, to maximize the amount of available training data for every model, we used full datasets in each model and measured performance on the full training data.

For bats, the final ensemble of models spanned a large range of performance on the training data, measured by the area under the receiver operator curve (AUC; Network 1: 0.624; Network 2: 0.987; Network 3: 0.514; Network 4: 0.726; Trait 1: 0.850; Trait 2: 0.902; Trait 3: 0.762; Hybrid 1: 0.924), indicating that it was possible to suitably detect differences in model performance on the full

data. The total ensemble of proportional ranks performed medium well (AUC = 0.832). We used
known betacoronavirus associations to threshold each model and the ensemble predictions
based on a 10% omission threshold (90% sensitivity), and we again found a wide range in the
number of predicted undiscovered bat hosts of betacoronaviruses (Network 1: 162 species;
Network 2: 1; Network 3: 111; Network 4: 44; Trait 1: 425; Trait 2: 384; Trait 3: 720; Hybrid 1: 181;
total ensemble: 239 species). Given concerns about mammal model performance and biological
accuracy (see Main Text), we elected not to apply this exercise to mammal hosts at large.

To visualize the spatial distribution of predicted bat hosts, we used the IUCN Red List database
of species geographic distributions. We took the top 50 ranked in-sample predictions and top 50
ranked out-of-sample predictions and combined these range maps to visualize species richness
of top predicted hosts (Figure 3).

Phylogenetic factorization of ensemble models

We used phylogenetic factorization to flexibly identify taxonomic patterns in the consensus
proportional rankings of likely hosts of SARS-CoV-2. Phylogenetic factorization is a graph-
partitioning algorithm that iteratively partitions a phylogeny in a series of generalized linear
models to identify clades at any taxonomic level (e.g., rather than *a priori* comparing strictly
among genera or family) that differ in a trait of interest [45]. Using the mammal supertree, we used
the *phylofactor* package to partition proportional rank as a Gaussian-distributed variable. We
determined the number of significant phylogenetic factors using a Holm's sequentially rejective
5% cutoff for the family-wise error rate. We applied this algorithm across our four final ensemble
prediction datasets: in-sample bat ranks, out-of-sample bat ranks, in-sample mammal ranks, and
out-of-sample mammal ranks.

Using network and trait-based models within-sample, we identified only one bat clade with
substantially different consensus proportional rankings, the Yangochiroptera ($\bar{x}$=0.55 compared
to 0.39 for the remaining bat phylogeny, the Yinpterochiroptera). Out of sample, using only trait-
based models, we instead identified eight bat clades with different propensities to include unlikely
or likely bat hosts of betacoronaviruses. Subclades of the New World superfamily Noctilionoidea
broadly had higher proportional ranks ($\bar{x}$=0.73), indicating lower predicted probability of being
hosts, as did the Emballanuridae ($\bar{x}$=0.75). In contrast, both the Rhinolophidae and the
Pteropodidae broadly had lower mean ranks ($\bar{x}$=0.29 and $\bar{x}$=0.38).

Using network models within-sample across non-volant mammals, we identified four clades with
different proportional ranks. The largest clade was the Laurasiatheria (Artiodactyla,
Perissodactyla, Carnivora, Pholidota, Soricomorpha, and Erinaceomorpha), which had lower
proportional ranks (higher risk; $\bar{x}$=0.55). Nested within this clade, the Cetacea had greater
proportional ranks ($\bar{x}$=0.89), indicating lower risk. A large subclade of the Murinae (Old World rats
and mice) also had lower ranks ($\bar{x}$=0.52). Out of sample, using only the biogeographic viral sharing
model, we instead identified 15 clades with different proportional ranks. The first clade identified
large swaths of the Muridae as having higher risk ($\bar{x}$=0.38) as well as the Laurasiatheria ($\bar{x}$=0.50).

Old World primates had weakly lower risk (x̄=0.65), as did the Scuridae (x̄=0.67). The Cetacea and
Pinnipedia both had greater proportional ranks (x̄=0.89 and x̄=0.71). Old World porcupines
(Hystricidae) and the Erinaceidae (Paraechinus, Hemiechinus, Mesechinus, Erinaceus, Atelerix)
both had greater risk (x̄=0.48 and x̄=0.39), while the Afrosoricida had higher ranks (x̄=0.97).
946
To assess potential discrepancy between taxonomic patterns in model ensemble predictions and
those of simply host betacoronavirus status itself, we ran a secondary phylogenetic factorization
treating host status as a Bernoulli-distributed variable, with the same procedure applied to
determine the number of significant phylogenetic factors. To assess sensitivity of taxonomic
patterns to sampling effort, we ran phylogenetic factorization with and without square-root
transformed PubMed citations per species as a weighting variable (Supplemental Figure 11).
953
Without accounting for study effort, phylogenetic factorization of betacoronavirus host status
identified one significant clade across the bat phylogeny, the Yangochiroptera, as having fewer
positive species (4.71%) than the paraphyletic remainder (12.12%). When accounting for study
effort, however, the single clade identified by phylogenetic factorization changed, with a subclade
of the family Pteropodidae (the Rousettinae) having a greater proportion of positive species
(28.6%). For non-volant mammals, phylogenetic factorization identified only one clade, the family
Camelidae, as having more positive species (75%) than the tree remainder (0.68%).
961
962   *Phylogenetic factorization of Rhinolophidae virus sharing*
963
964   Because phylogenetic patterns in predictions from our viral sharing model could vary across other
965   taxonomic scales beyond order and family, we also used phylogenetic factorization to more
966   flexibly identify host clades with different propensities to share viruses with *R. affinis* and *R.*
967   *malayanus*. We partitioned rank as a Gaussian-distributed variable and again determined the
968   number of significant phylogenetic factors using Holm's sequentially rejective 5% cutoff.
969
Within the Chiroptera, we identified 10 clades with different propensities to share viruses with *R.*
affinis and 5 clades with different propensities to share viruses with *R. malayanus*. For both bats,
the top clade was the family Rhinolophidae, reinforcing phylogenetic components of the
biogeographic model and highlighting the greater likelihood of viral sharing (mean rank x̄=40 for
R. affinis, x̄=42 for *R. malayanus*). For *R. affinis*, several individual bat species had lower risks of
viral sharing (e.g., *Myotis leibii*, x̄=4100; *Pteropus insularis*, x̄=3157; *Nyctimene aello*, x̄=2497;
Chaerephon chapini, x̄=2497). The Megadermatidae, Nycteridae, and Hipposideridae (under which
the PanTHERIA dataset includes the genus *Rhinonicteris*, although this is now considered a
separate family, the Rhinonycteridae[112]) collectively had greater likelihood of viral sharing
(x̄=557), as did the Vespertillionidae (x̄=704).
980
Across the non-volant mammals, we identified 7 clades with different propensities to share
viruses with *R. affinis* and only 1 clade with different propensities to share viruses with *R.*
malayanus. For both bat species, the first and primary clade was the Ferungulata (Artiodactyla,

Perissodactyla, Carnivora, Pholidota, Soricomorpha, and Erinaceomorpha), which had lower ranks
(higher viral sharing; $\bar{x}$=2084). For viral sharing with *R. affinis*, the Sciuridae was more likely to
share viruses ($\bar{x}$=1948), as was the Scandentia ($\bar{x}$=1416) and many members of the Colobinae
($\bar{x}$=1958). However, members of the tribe Muntiacini (genera *Elaphodus* and *Muntiacus*) had
especially high likelihoods of viral sharing and low rank ($\bar{x}$=361).
989
990   **Data and Code Availability**
991
The standardized data on betacoronavirus associations, and all associated predictor data, is
available from the VERENA consortium's Github (github.com/viralemergence/virionette). All
modeling teams contributed an individual repository with their methods, which are available in
the organizational directory (github.com/viralemergence). All code for analysis, and a working
reproduction of each authors' contributions, is available from the study repository
(github.com/viralemergence/Fresnel).

Supplemental Material

999
1000
1001 **Supplemental Table 1. Results of phylogenetic factorization applied to predicted rank**
1002 **probabilities for bats.** The number of retained phylogenetic factors (following a 5% family-wise
1003 error rate applied to GLMs), taxa corresponding to those clades, number of species per clade, and
1004 mean predicted rank probabilities for the clade compared to the paraphyletic remainder are
1005 shown stratified by models applied in- and out-of-sample.
1006

Sample	Factor	Taxa	Tips	Clade	Other
in	1	Yangochiroptera	160	0.550	0.393
out	1	Mystacinidae, Noctilionidae, Mormoopidae, Phyllostomidae	161	0.730	0.483
out	2	Rhinolophidae	73	0.295	0.538
out	3	Pteropodidae	173	0.379	0.548
out	4	Mosia, Emballonura, Coleura, Rhynchonycteris, Cyttarops, Diclidurus, Centronycteris, Cormura, Saccopteryx, Balantiopteryx, Peropteryx	31	0.755	0.512
out	5	Eonycteris, Macroglossus, Syconycteris, Notopteris, Melonycteris, Harpyionycteris, Aproteles, Dobsonia, Styloctenium, Neopteryx, Pteralopex, Acerodon, Pteropus	99	0.458	0.527
out	6	Thyropteridae, Furipteridae, Natalidae	12	0.818	0.517
out	7	Otomops, Promops, Molossus, Eumops	26	0.669	0.517
out	8	Lasiurus	15	0.717	0.518

1007
1008

1014

Sample	Factor	Taxa	Tips	Clade	Other
in	1	Phocoenidae, Delphinidae, Tursiops, Monodontidae, Physeteridae, Balaenopteridae, Eschrichtiidae	12	0.889	0.611
in	2	Artiodactyla, Perissodactyla, Carnivora, Pholidota, Erinaceomorpha, Soricomorpha	173	0.549	0.661
in	3	Lophuromys, Micaelamys, Apodemus, Arvicanthis, Bandicota, Madromys, Dasymys, Hydromys, Lemniscomys, Mastomys, Mus, Pelomys, Niviventer, Otomys, Praomys, Rattus, Vandeleuria	38	0.520	0.627
out	1	Abditomys, Bullimus, Limnomys, Tarsomys, Tryphomys, Acomys, Lophuromys, Uranomys, Aethomys, Micaelamys, Anisomys, Chiruromys, Coccymys, Crossomys, Hyomys, Leptomys, Lorentzimys, Pseudohydromys, Paraleptomys, Macruromys, Mallomys, Microhydromys, Parahydromys, Pogonomelomys, Abeomelomys, Solomys, Xenuromys, Apodemus, Tokudaia, Apomys, Crunomys, Chrotomys, Rhynchomys, Arvicanthis, Bandicota, Batomys, Carpomys, Crateromys, Berylmys, Bunomys, Chiromyscus, Chiropodomys, Hapalomys, Haeromys, Colomys, Nilopegamys, Conilurus, Leporillus, Mesembriomys, Melomys, Protochromys, Mammelomys, Paramelomys, Uromys, Zyzomys, Leggadina, Notomys, Pseudomys, Mastacomys, Madromys, Cremnomys, Millardia, Dacnomys, Dasymys, Dephomys, Hybomys, Hydromys, Xeromys, Desmomys, Diomys, Diplothrix, Echiothrix, Margaretamys, Melasmothrix, Tateomys, Eropeplus, Lenomys, Golunda, Grammomys, Thallomys, Hadromys, Heimyscus, Hylomyscus, Komodomys, Papagomys, Oenomys, Thamnomys, Lemniscomys, Lenothrix, Leopoldamys, Malacomys, Praomys, Myomyscus, Mastomys, Maxomys, Micromys, Muriculus, Mus, Mylomys, Pelomys, Stenocephalemys, Nesokia, Niviventer, Otomys, Parotomys, Palawanomys, Paruromys, Phloeomys, Pithecheir, Pogonomys, Rattus, Rhabdomys, Srilankamys, Nesoromys, Stochomys, Sundamys, Taeromys, Vandeleuria, Vernaya, Zelotomys	510	0.382	0.672
out	2	Artiodactyla, Perissodactyla, Carnivora, Pholidota	505	0.495	0.651
out	3	Anomaluridae, Pedetidae, Dipodidae, Cricetidae, Muridae, Nesomyidae, Calomyscidae, Spalacidae, Platacanthomyidae	779	0.643	0.622
out	4	Talpidae, Erinaceomorpha, Soricidae	357	0.630	0.627
out	5	Cercopithecidae, Hominidae, Hylobatidae	139	0.649	0.626

1015

Sample	Factor	Taxa	Tips	Clade	Other
out	6	Abrawayaomys, Handleyomys, Aepeomys, Thomasomys, Abrothrix, Akodon, Necromys, Deltamys, Thaptomys, Andalgalomys, Auliscomys, Loxodontomys, Phyllotis, Paralomys, Graomys, Andinomys, Bibimys, Kunsia, Scapteromys, Blarinomys, Calomys, Chelemys, Chilomys, Chinchillula, Delomys, Eligmodontia, Euneomys, Galenomys, Geoxus, Holochilus, Lundomys, Pseudoryzomys, Irenomys, Lenoxus, Melanomys, Microryzomys, Neacomys, Nectomys, Neotomys, Nesoryzomys, Notiomys, Oecomys, Oligoryzomys, Oryzomys, Oxymycterus, Brucepattersonius, Phaenomys, Podoxymys, Punomys, Reithrodon, Rhagomys, Rhipidomys, Scolomys, Sigmodontomys, Thalpomys, Wiedomys, Wilfredomys, Juliomys, Zygodontomys, Anotomys, Chibchanomys, Ichthyomys, Neusticomys, Rheomys, Sigmodon, Nyctomys, Otonyctomys, Ototylomys, Tylomys, Baiomys, Scotinomys, Ochrotomys, Habromys, Neotomodon, Podomys, Osgoodomys, Megadontomys, Peromyscus, Onychomys, Isthmomys, Reithrodontomys, Hodomys, Xenomys, Neotoma, Nelsonia	397	0.703	0.616
out	7	Tamiasciurus, Sciurus, Rheithrosciurus, Microsciurus, Syntheosciurus, Pteromys, Petaurista, Belomys, Biswamoyopterus, Trogopterus, Pteromyscus, Aeromys, Eupetaurus, Aeretes, Glaucomys, Eoglaucomys, Hylopetes, Petinomys, Petaurillus, Iomys, Ratufa, Callosciurus, Glyphotes, Lariscus, Menetes, Rhinosciurus, Funambulus, Tamiops, Dremomys, Exilisciurus, Hyosciurus, Prosciurillus, Rubrisciurus, Nannosciurus, Sundasciurus	139	0.672	0.625
out	8	Phocoenidae, Delphinidae, Tursiops, Monodontidae, Physeteridae, Balaenopteridae, Eschrichtiidae	12	0.889	0.626
out	9	Odobenidae, Otariidae, Phocidae	33	0.714	0.626
out	10	Hystricidae	11	0.482	0.627
out	11	Caprolagus, Poelagus, Lepus, Oryctolagus	33	0.642	0.627
out	12	Paraechinus, Hemiechinus, Mesechinus, Erinaceus, Atelerix	15	0.388	0.628
out	13	Afrosoricida	41	0.970	0.623
out	14	Castoridae, Heteromyidae, Geomyidae, Octodontidae, Ctenodactylidae, Ctenomyidae, Abrocomidae, Caviidae, Dinomyidae, Petromuridae, Dasyproctidae, Myocastoridae, Echimyidae, Erethizontidae, Capromyidae, Cuniculidae, Thryonomyidae, Bathyergidae, Chinchillidae	295	0.872	0.603
out	15	Cheirogaleidae, Indriidae, Daubentoniidae, Lemuridae, Lepilemuridae	48	0.921	0.623

Supplemental Table 3. Predicted high-similarity bat hosts sharing with *Rhinolophus affinis* and
R. malayanus. Species on these lists may be particularly likely to be the ultimate evolutionary
origin of SARS-CoV-2, or a closely-related virus prior to recombination in an intermediate host.
Predictions are made based just on the average viral sharing probability inferred for the two hosts
from the phylogeography model (Trait-based 3). (* Note that the two species have high sharing
probabilities with each other, potentially indicating that efforts to trace the origins of SARS-CoV-
2 are already very close to their target.)

1026

Rhinolophus affinis	Rhinolophus malayanus
1. Rhinolophus macrotis (P=0.84)	1. Rhinolophus shameli (P=0.87)
2. Rhinolophus stheno (P=0.83)	2. Rhinolophus coelophyllus (P=0.84)
3. Rhinolophus malayanus (P=0.82)	3. Rhinolophus thomasi (P=0.84)
4. Rhinolophus acuminatus (P=0.81)	4. Rhinolophus affinis (P=0.82)
5. Rhinolophus pearsonii (P=0.78)	5. Rhinolophus marshalli (P=0.82)
6. Rhinolophus shameli (P=0.78)	6. Rhinolophus pearsonii (P=0.82)
7. Rhinolophus thomasi (P=0.78)	7. Rhinolophus yunanensis (P=0.79)
8. Rhinolophus sinicus (P=0.77)	8. Rhinolophus paradoxolophus (P=0.78)
9. Rhinolophus trifoliatus (P=0.76)	9. Rhinolophus macrotis (P=0.76)
10. Rhinolophus marshalli (P=0.72)	10. Rhinolophus acuminatus (P=0.75)
11. Rhinolophus shortridgei (P=0.71)	11. Rhinolophus siamensis (P=0.75)
12. Rhinolophus luctus (P=0.7)	12. Rhinolophus rouxii (P=0.72)
13. Rhinolophus sedulus (P=0.7)	13. Rhinolophus stheno (P=0.71)
14. Rhinolophus rouxii (P=0.69)	14. Rhinolophus luctus (P=0.69)
15. Rhinolophus pusillus (P=0.68)	15. Rhinolophus trifoliatus (P=0.65)
16. Rhinolophus ferrumequinum (P=0.67)	16. Rhinolophus pusillus (P=0.62)
17. Rhinolophus lepidus (P=0.67)	17. Rhinolophus borneensis (P=0.6)
18. Hipposideros pomona (P=0.66)	18. Hipposideros lylei (P=0.59)
19. Rhinolophus celebensis (P=0.66)	19. Rhinolophus shortridgei (P=0.59)
20. Rhinolophus paradoxolophus (P=0.66)	20. Rhinolophus sinicus (P=0.59)

1027

Supplemental Table 4. Predicted high-similarity non-bat hosts sharing with *Rhinolophus affinis*
and *R. malayanus*. Species on these lists may be particularly suitable as stepping stones for
betacoronavirus transmission from bats into humans, including potentially for SARS-CoV-2 and
other SARS-like viruses. Predictions are made based just on the average viral sharing probability
inferred for the two hosts from the phylogeography model (Trait-based 3). Species' binomial
names are included alongside their families.

1034

Rhinolophus affinis		*Rhinolophus malayanus*	
1. Arctonyx collaris (P=0.33)	Mustelidae	1. Arctonyx collaris (P=0.29)	Mustelidae
2. Budorcas taxicolor (P=0.33)	Bovidae	2. Herpestes urva (P=0.28)	Herpestidae
3. Viverra tangalunga (P=0.32)	Viverridae	3. Lutrogale perspicillata (P=0.28)	Mustelidae
4. Manis javanica (P=0.3)	Manidae	4. Melogale personata (P=0.27)	Mustelidae
5. Mustela altaica (P=0.3)	Mustelidae	5. Viverra megaspila (P=0.26)	Viverridae
6. Ursus thibetanus (P=0.3)	Ursidae	6. Arctictis binturong (P=0.25)	Viverridae
7. Cynogale bennettii (P=0.29)	Viverridae	7. Euroscaptor klossi (P=0.25)	Talpidae
8. Elaphodus cephalophus (P=0.29)	Cervidae	8. Lutra sumatrana (P=0.25)	Mustelidae
9. Lutrogale perspicillata (P=0.29)	Mustelidae	9. Sus scrofa (P=0.25)	Suidae
10. Viverricula indica (P=0.29)	Viverridae	10. Capricornis milneedwardsii (P=0.23)	Bovidae
11. Capricornis sumatraensis (P=0.28)	Bovidae	11. Manis javanica (P=0.23)	Manidae
12. Chimarrogale himalayica (P=0.28)	Soricidae	12. Manis pentadactyla (P=0.23)	Manidae
13. Helarctos malayanus (P=0.28)	Ursidae	13. Mustela nudipes (P=0.23)	Mustelidae
14. Herpestes javanicus (P=0.27)	Herpestidae	14. Paguma larvata (P=0.23)	Viverridae
15. Hylomys suillus (P=0.27)	Erinaceidae	15. Panthera pardus (P=0.23)	Felidae
16. Mustela kathiah (P=0.27)	Mustelidae	16. Viverra zibetha (P=0.23)	Viverridae
17. Capricornis milneedwardsii (P=0.26)	Bovidae	17. Bandicota savilei (P=0.22)	Muridae
18. Catopuma temminckii (P=0.26)	Felidae	18. Chrotogale owstoni (P=0.22)	Viverridae
19. Crocidura negligens (P=0.26)	Soricidae	19. Crocidura fuliginosa (P=0.22)	Soricidae
20. Capricornis thar (P=0.25)	Bovidae	20. Crocidura vorax (P=0.22)	Soricidae

1035

1036 **Supplemental Table 5. Taxonomic scale of model training data and predictive implementation.**
1037 Notes: (1) These models generated predictions of sharing with *Rhinolophus affinis* over all non-
1038 human mammals in the HP3 dataset, then subsetted to bats. (2) In these models, bat-
1039 betacoronavirus predictions are based on a subset of binary outcomes for known association
1040 with betacoronaviruses, without any other viruses included.
1041

Model approach	Training data scale	Bat *Betacoronavirus* predictions	Mammal-wide *Betacoronavirus* predictions
Network-based 1 k-Nearest neighbors	Bat-virus	✓	
Network-based 1 k-Nearest neighbors	Mammal-virus		✓
Network-based 2 Linear filter	Bat-virus	✓	
Network-based 2 Linear filter	Mammal-virus		✓
Network-based 3 Plug-and-play	Mammal-virus[1]	✓	✓
Network-based 4 Scaled-phylogeny	Bat-virus	✓	
Network-based 4 Scaled-phylogeny	Mammal-virus		✓
Trait-based 1 Boosted regression trees	Bat-betacoronavirus[2]	✓	
Trait-based 2 Bayesian additive regression trees	Bat-betacoronavirus[2]	✓	
Trait-based 3 Neutral phylogeographic	Mammal-virus[1]	✓	✓
Hybrid 1 Two-step kernel ridge regression	Bat-betacoronavirus	✓	

1042
1043

1044 **Supplemental Table 6. Data scale of prediction, by method.** Some methods use pseudoabsences
1045 to expand the scale of prediction but still only analyze existing data, with no out-of-sample
1046 inference, while others can predict freshly onto new data. (* Training data from the HP3 database
1047 uses pseudoabsences, but no new ones are generated in this study that modify the model or the
1048 bat-virus association dataset)

1049

Model approach	Prediction on hosts without known associations (out-of-sample)	Predictive extent and use of pseudoabsences
Network-based 1 k-Nearest neighbors	No	Only predicts link probabilities among species in the association data
Network-based 2 Linear filter	No	Only predicts link probabilities among species in the association data
Network-based 3 Plug-and-play	No	Uses pseudoabsences to predict over all mammals in association data, using latent approach
Network-based 4 Scaled-phylogeny	No	Only predicts link probabilities among species in the association data
Trait-based 1 Boosted regression trees	Yes	Uses pseudoabsences for all bats in trait data to predict over all species, including those without known associations
Trait-based 2 Bayesian additive regression trees	Yes	Uses pseudoabsences for all bats in trait data to predict over all species, including those without known associations
Trait-based 3 Neutral phylogeographic	Yes	Trains on a broader network, and predicts sharing probabilities among any mammals in phylogeny and IUCN range map data
Hybrid 1 Two-step kernel ridge regression	Yes	Uses pseudoabsences for all bats in trait data to predict over all species, including those without known associations

1050
1051

1052 **Supplemental Table 7. Model performance for hybrid model.** Area under receiver-operator
1053 curve for predicting interactions, or predicting for betacoronaviruses or for hosts, using bat
1054 traits, phylogeny or a combination of both.
1055
1056

	interactions	Betacoronavirus	hosts (average)
traits	0.8784	0.6742	0.7304
phylogeny	0.8848	0.73389	0.7623
both	0.8975	0.7510	0.7909

1057

1058 **Supplemental Figure 1. Bat models perform more strongly together than in isolation.** Curves
1059 show observed betacoronavirus hosts against predicted proportional ranks from eight individual
1060 models, and incorporated into one multi-model ensemble. Black lines show a binomial GLM fit to
1061 the predicted ranks against the recorded presence or absence of known betacoronavirus
1062 associations. Points are jittered to reduce overlap.
1063

1064

Supplemental Figure 2. Poor concordance among predictive models for mammal hosts of
betacoronaviruses. The pairwise Spearman's rank correlations between models' ranked species-
level predictions were generally low (A). In-sample predictions varied significantly and heavily
prioritized domestic animals and well-studied hosts (B). The ten species with the highest mean
proportional ranks across all models are highlighted in shades of purple. Only in-sample
predictions are displayed because only one model (Trait-based 3) was able to predict out of
sample for all mammals.

1072

1073

Supplemental Figure 3. Results of phylogenetic factorization applied to predicted ranks of virus
sharing with *Rhinolophus affinis* and *Rhinolophus malayanus*. Colored regions indicate clades
identified as significantly different in their predicted rank compared to the paraphyletic remainder;
those more likely to share a virus with the *Rhinolophus* are shown in red, whereas those less likely
to share a virus are shown in blue. Bar height indicates predicted rank (higher values = lower rank
score, more likely share virus). Results are displayed for bats and remaining mammals separately.
Mammal-wide clades with high propensities to share viruses with *R. affinis* based solely on their
phylogeography included the treeshrews (Scandentia), Old World monkeys (Colobinae), and both
tufted and barking deer (Muntiacini).
1083
1084

1085

1086   **Supplemental Figure 4.** Predicted species-level sharing probabilities of A) *Rhinolophus affinis* and
1087   B) *Rhinolophus malayanus*, calculated according to the phylogeographic viral sharing model[48].
1088   Each coloured point is a mammal species. Black points and error bars denote means and
1089   standard errors for each order. Mammal orders are arranged according to their mean sharing
1090   probability.

1091

1092

1093   **Supplemental Figure 5.** Predicted species-level sharing probabilities of A) *Rhinolophus affinis* and
1094   B) *Rhinolophus malayanus*, calculated according to the phylogeographic viral sharing model^.
1095   Each coloured point is a mammal species. Black points and error bars denote means and
1096   standard errors for each order. Mammal orders are arranged according to their mean sharing
1097   probability.
1098

1099
1100

Supplemental Figure 6. To account for uncertainty in the phylogenetic distances among hosts,
the scaled-phylogeny model estimates a tree scaling parameter (eta) based on an early-burst
model of evolution. On the left is the unscaled bat phylogeny for the hosts in the bat-virus genera
network, and on the right is the same tree rescaled according to mean estimated scaling
parameter (eta = 7.92). Eta values above 1 indicate accelerating evolution, suggesting less
phylogenetic conservatism in host-virus associations among closely related taxa than would be
predicted by a Brownian-motion model on the unscaled tree.
1108

Unscaled tree   Scaled tree (eta = 7.92)

1109

Supplemental Figure 7. Four formulations of Bayesian additive regression tree (BART) models
produce slightly different results, but largely agree. Two models use baseline BART, while two
models use a Dirichlet prior on variable importance (DART). Two are uncorrected for sampling
bias (u) while two are corrected using citation counts (c). In the final main-text model ensemble,
we present a DART model including correction for citation bias, which penalizes overfitting and
spurious patterns two ways and leads to predictions with a lower total correlation with the data,
but a still-high performance (AUC = 0.90).
1117

1118

1119   **Supplemental Figure 8.** Variable importance plots for the Bayesian additive regression tree
1120   models with uniform variable importance prior (top) versus Dirichlet prior (bottom), without (left)
1121   and with (right) correction for citations.

1122

Supplemental Figure 9. Overview of the Two-Step Kernel Ridge Regression method. A. The
incidence matrix *Y* is modelled by two kernel matrices *K* and *G*, describing the hosts (here: bats)
and viruses, respectively. The equations for fitting the model and making predictions are given
below. B. Computing the host kernel matrix is done based on traits, phylogeny or both. A standard
radial basis kernel is used to transform a distance to a valid kernel. The two kernels are combined
by averaging. C. The models are validated by specialized cross-validation. The setting
interactions leaves one interaction out at a time and repeats this for every interaction. AUC is
computed in micro-fashion. In the setting hosts, we leave out one host at a time and compute the
AUC per virus, averaging the individual results. D. To assess the variable importance, we first
randomly reshuffle the kernel matrices and monitor how the performance of a model fitted on
these data drops. In a more fine-grained approach, we just reshuffle a single trait (indicated by
the red arrow) and compare the resulting performances.
1135

1136

1137 **Supplemental Figure 10. Illustration of the TSKRR model and the variable importance scores.** A.
1138 Heatmap of the prediction scores, together with a hierarchical clustering based on the kernel
1139 distance of *K* and *G*. B. Variable importance computed by random permutations of the kernel
1140 matrices or a single trait.

1141
1142
1143

1144 **Supplemental Figure 11.** Results of phylogenetic factorization applied to binomial
1145 betacoronavirus data across bats (top) and other mammals (bottom), using raw data (left) and
1146 after weighting by citation counts (right). Any significant clades (5% family-wise error rate) are
1147 displayed in colored sharing on the phylogeny. Bars indicate betacoronavirus detection, and
1148 clades are colored by having more (red) or fewer (blue) positive species.
1149

1150

Bibliography

1. Anthony, S. J. *et al.* Global patterns in coronavirus diversity. *Virus Evol* **3**, vex012 (2017).

2. Denison, M. R., Graham, R. L., Donaldson, E. F., Eckerle, L. D. & Baric, R. S. Coronaviruses: an RNA proofreading machine regulates replication fidelity and diversity. *RNA Biol.* **8**, 270–279 (2011).

3. Ren, W. *et al.* Full-length genome sequences of two SARS-like coronaviruses in horseshoe bats and genetic variation analysis. *J. Gen. Virol.* **87**, 3355–3359 (2006).

4. Li, W. *et al.* Bats are natural reservoirs of SARS-like coronaviruses. *Science* **310**, 676–679 (2005).

5. Yang, X.-L. *et al.* Isolation and Characterization of a Novel Bat Coronavirus Closely Related to the Direct Progenitor of Severe Acute Respiratory Syndrome Coronavirus. *J. Virol.* **90**, 3253–3256 (2015).

6. Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270–273 (2020).

7. Guan, Y. *et al.* Isolation and characterization of viruses related to the SARS coronavirus from animals in southern China. *Science* **302**, 276–278 (2003).

8. Hu, B. *et al.* Discovery of a rich gene pool of bat SARS-related coronaviruses provides new insights into the origin of SARS coronavirus. *PLoS Pathog.* **13**, e1006698 (2017).

9. Memish, Z. A. *et al.* Middle East respiratory syndrome coronavirus in bats, Saudi Arabia. *Emerg. Infect. Dis.* **19**, 1819–1823 (2013).

10. Wang, Q. *et al.* Bat origins of MERS-CoV supported by bat coronavirus HKU4 usage of human receptor CD26. *Cell Host Microbe* **16**, 328–337 (2014).

11. Yang, Y. *et al.* Receptor usage and cell entry of bat coronavirus HKU4 provide insight into bat-to-human transmission of MERS coronavirus. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 12516–12521 (2014).

12. Hu, B., Ge, X., Wang, L.-F. & Shi, Z. Bat origin of human coronaviruses. *Virology Journal* vol. 12 (2015).

13. Anthony, S. J. *et al.* Further Evidence for Bats as the Evolutionary Source of Middle East Respiratory Syndrome Coronavirus. *MBio* **8**, (2017).

14. Anthony, S. J. *et al.* Coronaviruses in bats from Mexico. *J. Gen. Virol.* **94**, 1028–1038 (2013).

15. Yang, L. *et al.* MERS-related betacoronavirus in Vespertilio superans bats, China. *Emerg. Infect. Dis.* **20**, 1260–1262 (2014).

16. Zhou, H. *et al.* A novel bat coronavirus reveals natural insertions at the S1/S2 cleavage site of the Spike protein and a possible recombinant origin of HCoV-19. doi:10.1101/2020.03.02.974139.

17. Nielsen, R., Wang, H. & Pipes, L. Synonymous mutations and the molecular evolution of SARS-Cov-2 origins. doi:10.1101/2020.04.20.052019.

18. Lam, T. T.-Y. *et al.* Identifying SARS-CoV-2 related coronaviruses in Malayan pangolins. *Nature* (2020) doi:10.1038/s41586-020-2169-0.

19. Xiao, K. *et al.* Isolation of SARS-CoV-2-related coronavirus from Malayan pangolins. *Nature* (2020) doi:10.1038/s41586-020-2313-x.

20. Zhang, T., Wu, Q. & Zhang, Z. Probable Pangolin Origin of SARS-CoV-2 Associated with the
COVID-19 Outbreak. *Curr. Biol.* **30**, 1578 (2020).
21. Andersen, K. G., Rambaut, A., Ian Lipkin, W., Holmes, E. C. & Garry, R. F. The proximal origin
of SARS-CoV-2. *Nature Medicine* vol. 26 450–452 (2020).
22. Viana, M. *et al.* Assembling evidence for identifying reservoirs of infection. *Trends Ecol.*
Evol. **29**, 270–279 (2014).
23. Plowright, R. K. *et al.* Prioritizing surveillance of Nipah virus in India. *PLoS Negl. Trop. Dis.*
13, e0007393 (2019).
24. Becker, D. J., Crowley, D. E., Washburne, A. D. & Plowright, R. K. Temporal and spatial
limitations in global surveillance for bat filoviruses and henipaviruses. *Biol. Lett.* **15**,
20190423 (2019).
25. Becker, D. J., Washburne, A. D., Faust, C. L., Mordecai, E. A. & Plowright, R. K. The problem
of scale in the prediction and management of pathogen spillover. *Philos. Trans. R. Soc.*
Lond. B Biol. Sci. **374**, 20190224 (2019).
26. Becker, D. J. & Han, B. A. The macroecology and evolution of avian competence for Borrelia
burgdorferi. doi:10.1101/2020.04.15.040352.
27. Han, B. A. *et al.* Undiscovered Bat Hosts of Filoviruses. *PLoS Negl. Trop. Dis.* **10**, e0004815
(2016).
28. Han, B. A. *et al.* Confronting data sparsity to identify potential sources of Zika virus spillover
infection among primates. *Epidemics* **27**, 59–65 (2019).
29. Washburne, A. D. *et al.* Taxonomic patterns in the zoonotic potential of mammalian viruses.
PeerJ **6**, e5979 (2018).
30. Olival, K. J. *et al.* Host and viral traits predict zoonotic spillover from mammals. *Nature* **546**,
646–650 (2017).
31. Fritz, S. A., Bininda-Emonds, O. R. P. & Purvis, A. Geographical variation in predictors of
mammalian extinction risk: big is bad, but only in the tropics. *Ecol. Lett.* **12**, 538–549
(2009).
32. Jones, K. E. *et al.* PanTHERIA: a species-level database of life history, ecology, and
geography of extant and recently extinct mammals. *Ecology* vol. 90 2648–2648 (2009).
33. Wilman, H. *et al.* EltonTraits 1.0: Species-level foraging attributes of the world's birds and
mammals. *Ecology* vol. 95 2027–2027 (2014).
34. Trifonova, N. *et al.* Spatio-temporal Bayesian network models with latent variables for
revealing trophic dynamics and functional networks in fisheries ecology. *Ecological*
Informatics vol. 30 142–158 (2015).
35. Rohr, R. P., Scherer, H., Kehrli, P., Mazza, C. & Bersier, L.-F. Modeling food webs: exploring
unexplained structure using latent traits. *Am. Nat.* **176**, 170–177 (2010).
36. Dallas, T., Park, A. W. & Drake, J. M. Predicting cryptic links in host-parasite networks. *PLoS*
Comput. Biol. **13**, e1005557 (2017).
37. Han, B. A., Schmidt, J. P., Bowden, S. E. & Drake, J. M. Rodent reservoirs of future zoonotic
diseases. *Proceedings of the National Academy of Sciences* vol. 112 7039–7044 (2015).
38. Brandão, P. E. *et al.* A coronavirus detected in the vampire bat Desmodus rotundus. *Braz. J.*
Infect. Dis. **12**, 466–468 (2008).

39. Corman, V. M. *et al.* Highly diversified coronaviruses in neotropical bats. *J. Gen. Virol.* **94**,
1984–1994 (2013).
40. Moreira-Soto, A. *et al.* Neotropical bats from Costa Rica harbour diverse coronaviruses.
Zoonoses Public Health **62**, 501–505 (2015).
41. Wang, L. *et al.* Discovery and genetic analysis of novel coronaviruses in least horseshoe
bats in southwestern China. *Emerg. Microbes Infect.* **6**, e14 (2017).
42. Lin, X.-D. *et al.* Extensive diversity of coronaviruses in bats from China. *Virology* vol. 507 1–
10 (2017).
43. Wacharapluesadee, S. *et al.* Diversity of coronavirus in bats from Eastern Thailand. *Virol. J.*
12, 57 (2015).
44. Guy, C., Ratcliffe, J. M. & Mideo, N. The influence of bat ecology on viral diversity and
reservoir status. *Ecol. Evol.* **2008**, 209 (2020).
45. Washburne, A. D. *et al.* Phylofactorization: a graph partitioning algorithm to identify
phylogenetic scales of ecological data. *Ecol. Monogr.* **89**, e01353 (2019).
46. Crowley, D., Becker, D., Washburne, A. & Plowright, R. Identifying Suspect Bat Reservoirs of
Emerging Infections. *Vaccines* vol. 8 228 (2020).
47. Damas, J., Hughes, G. M., Keough, K. C. & Painter, C. A. Broad Host Range of SARS-CoV-2
Predicted by Comparative and Structural Analysis of ACE2 in Vertebrates. *bioRxiv* (2020).
48. Albery, G. F., Eskew, E. A., Ross, N. & Olival, K. J. Predicting the global mammalian viral
sharing network using phylogeography. *Nat. Commun.* **11**, 2260 (2020).
49. Wang, M. *et al.* SARS-CoV Infection in a Restaurant from Palm Civet. *Emerging Infectious*
Diseases vol. 11 1860–1865 (2005).
50. Song, H.-D. *et al.* Cross-host evolution of severe acute respiratory syndrome coronavirus in
palm civet and human. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 2430–2435 (2005).
51. Dougherty, E. R., Seidel, D. P., Carlson, C. J., Spiegel, O. & Getz, W. M. Going through the
motions: incorporating movement analyses into disease research. *Ecol. Lett.* **21**, 588–604
(2018).
52. Lehmann, D. *et al.* Pangolins and bats living together in underground burrows in Lopé
National Park, Gabon. *African Journal of Ecology* (2020) doi:10.1111/aje.12759.
53. Oreshkova, N. *et al.* SARS-CoV2 infection in farmed mink, Netherlands, April 2020.
doi:10.1101/2020.05.18.101493.
54. Kim, Y.-I. *et al.* Infection and Rapid Transmission of SARS-CoV-2 in Ferrets. *Cell Host*
Microbe **27**, 704–709.e2 (2020).
55. Shi, J. *et al.* Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS–
coronavirus 2. *Science* (2020) doi:10.1126/science.abb7015.
56. Yang, X.-L. *et al.* Genetically Diverse Filoviruses in Rousettus and Eonycteris spp. Bats,
China, 2009 and 2015. *Emerg. Infect. Dis.* **23**, 482–486 (2017).
57. Seifert, S. N. *et al.* Rousettus aegyptiacus Bats Do Not Support Productive Nipah Virus
Replication. *The Journal of Infectious Diseases* vol. 221 S407–S413 (2020).
58. Wacharapluesadee, S. *et al.* Longitudinal study of age-specific pattern of coronavirus
infection in Lyle's flying fox (Pteropus lylei) in Thailand. *Virol. J.* **15**, 38 (2018).
59. Latinne, A. *et al.* Origin and cross-species transmission of bat coronaviruses in China.

doi:10.1101/2020.05.31.116061.

60. Yang, L. *et al.* Novel SARS-like betacoronaviruses in bats, China, 2011. *Emerg. Infect. Dis.* **19**, 989–991 (2013).

61. Geldenhuys, M. *et al.* A metagenomic viral discovery approach identifies potential zoonotic and novel mammalian viruses in Neoromicia bats within South Africa. *PLoS One* **13**, e0194527 (2018).

62. Memish, Z. A. *et al.* Middle East respiratory syndrome coronavirus in bats, Saudi Arabia. *Emerg. Infect. Dis.* **19**, 1819–1823 (2013).

63. Luo, Y. *et al.* Longitudinal Surveillance of Betacoronaviruses in Fruit Bats in Yunnan Province, China During 2009-2016. *Virol. Sin.* **33**, 87–95 (2018).

64. Peel, A. J. *et al.* Synchronous shedding of multiple bat paramyxoviruses coincides with peak periods of Hendra virus spillover. *Emerg. Microbes Infect.* **8**, 1314–1323 (2019).

65. de Souza Cortez J. L. Dunnum A. W. Ferguson F. A. Anwarali Khan D. L. Paul D. M. Reeder N. B. Simmons B. M. Thiers C. W. Thompson N S. Upham M. P. M. Vanhove P. W. Webala M. Weksler R. Yanagihara P. S. Soltis., C. J. A. S. A. B. A. J. B. C. A. C. B. M. B. Integrating biodiversity infrastructure into pathogen discovery and mitigation of epidemic infectious diseases. *Bioscience* (2020) doi:biaa064.

66. Kingston, T. *et al.* Networking networks for global bat conservation. in *Bats in the Anthropocene: Conservation of Bats in a Changing World* 539–569 (Springer, Cham, 2016).

67. Phelps, K. L. *et al.* Bat Research Networks and Viral Surveillance: Gaps and Opportunities in Western Asia. *Viruses* **11**, (2019).

68. Teeling, E. C. *et al.* Bat Biology, Genomes, and the Bat1K Project: To Generate Chromosome-Level Genomes for All Living Bat Species. *Annu Rev Anim Biosci* **6**, 23–46 (2018).

69. Mandl, J. N., Schneider, C., Schneider, D. S. & Baker, M. L. Going to Bat(s) for Studies of Disease Tolerance. *Front. Immunol.* **9**, 2112 (2018).

70. Skirmuntt, E. C., Escalera-Zamudio, M., Teeling, E. C., Smith, A. & Katzourakis, A. The Potential Role of Endogenous Viral Elements in the Evolution of Bats as Reservoirs for Zoonotic Viruses. *Annu Rev Virol* (2020) doi:10.1146/annurev-virology-092818-015613.

71. Jebb, D. *et al.* Six new reference-quality bat genomes illuminate the molecular basis and evolution of bat adaptations. (2019).

72. Gervasi, S. S., Civitello, D. J., Kilvitis, H. J. & Martin, L. B. The context of host competence: a role for plasticity in host–parasite dynamics. *Trends Parasitol.* **31**, 419–425 (2015).

73. Martin, L. B., Burgan, S. C., Adelman, J. S. & Gervasi, S. S. Host Competence: An Organismal Trait to Integrate Immunology and Epidemiology. *Integr. Comp. Biol.* **56**, 1225–1237 (2016).

74. Callaway, E. & Cyranoski, D. Why snakes probably aren't spreading the new China virus. *Nature* (2020) doi:10.1038/d41586-020-00180-8.

75. Gong, Y., Wen, G., Jiang, J. & Feng, X. Complete title: Codon bias analysis may be insufficient for identifying host(s) of a novel virus. *J. Med. Virol.* (2020) doi:10.1002/jmv.25977.

76. Zhao, H. COVID-19 drives new threat to bats in China. *Science* **367**, 1436 (2020).

77. Fenton, M. B. *et al.* Knowledge gaps about rabies transmission from vampire bats to

humans. *Nature Ecology & Evolution* **4**, 517–518 (2020).

78. López-Baucells, A., Rocha, R. & Fernández-Llamazares, Á. When bats go viral: negative
framings in virological research imperil bat conservation. *Mamm. Rev.* **48**, 62–66 (2018).

79. O'Shea, T. J., Cryan, P. M., Hayman, D. T. S., Plowright, R. K. & Streicker, D. G. Multiple
mortality events in bats: a global review. *Mamm. Rev.* **46**, 175–190 (2016).

80. MB Fenton, S Mubareka, SM Tsang, NB Simmons, DJ Becker. COVID-19 and threats to bats.
FACETS in press (2020).

81. Aguiar, L. M. S., Brito, D. & Machado, R. B. Do current vampire bat (Desmodus rotundus)
population control practices pose a threat to Dekeyser's nectar bat's (Lonchophylla
dekeyseri) long-term persistence in the Cerrado? *Acta Chiropt.* **12**, 275–282 (2010).

82. Streicker, D. G. *et al.* Ecological and anthropogenic drivers of rabies exposure in vampire
bats: implications for transmission and control. *Proc. Biol. Sci.* **279**, 3384–3392 (2012).

83. Blackwood, J. C., Streicker, D. G., Altizer, S. & Rohani, P. Resolving the roles of immunity,
pathogenesis, and immigration for rabies persistence in vampire bats. *Proc. Natl. Acad. Sci.*
U. S. A. **110**, 20837–20842 (2013).

84. Frick, W. F. *et al.* An emerging disease causes regional population collapse of a common
North American bat species. *Science* **329**, 679–682 (2010).

85. Sabir, J. S. M. *et al.* Co-circulation of three camel coronavirus species and recombination of
MERS-CoVs in Saudi Arabia. *Science* **351**, 81–84 (2016).

86. Guth, S., Visher, E., Boots, M. & Brook, C. E. Host phylogenetic distance drives trends in virus
virulence and transmissibility across the animal–human interface. *Philos. Trans. R. Soc.*
Lond. B Biol. Sci. **374**, 20190296 (2019).

87. Redondo, R. A. F., Brina, L. P. S., Silva, R. F., Ditchfield, A. D. & Santos, F. R. Molecular
systematics of the genus Artibeus (Chiroptera: Phyllostomidae). *Mol. Phylogenet. Evol.* **49**,
44–58 (2008).

88. Bouchard, S. Chaerephon pumilus. *Mammalian Species* 1–6 (1998).

89. Hoofer, S. R., Van Den Bussche, R. A. & HoráČek, I. Generic Status of the American
Pipistrelles (Vespertilionidae) with Description of a New Genus. *J. Mammal.* **87**, 981–992
(2006).

90. Desjardins-Proulx, P., Laigle, I., Poisot, T. & Gravel, D. Ecological interactions and the Netflix
problem. *PeerJ* **5**, e3644 (2017).

91. Stock, M., Poisot, T., Waegeman, W. & De Baets, B. Linear filtering reveals false negatives in
species interaction data. *Sci. Rep.* **7**, 45908 (2017).

92. Drake, J. M. & Richards, R. L. Estimating environmental suitability. *Ecosphere* vol. 9 e02373
(2018).

93. Dallas, T. A., Carlson, C. J. & Poisot, T. Testing predictability of disease outbreaks with a
simple model of pathogen biogeography. *R Soc Open Sci* **6**, 190883 (2019).

94. Elmasri, M., Farrell, M. J., Jonathan Davies, T. & Stephens, D. A. A hierarchical Bayesian
model for predicting ecological interactions using scaled evolutionary relationships. *The*
Annals of Applied Statistics vol. 14 221–240 (2020).

95. Cadotte, M. W. *et al.* Phylogenetic diversity metrics for ecological communities: integrating
species richness, abundance and evolutionary history. *Ecol. Lett.* **13**, 96–105 (2010).

96. Park, A. W. *et al.* Characterizing the phylogenetic specialism–generalism spectrum of
mammal parasites. *Proceedings of the Royal Society B: Biological Sciences* vol. 285
20172613 (2018).
97. Harvey, P. H. & Pagel, M. D. *The comparative method in evolutionary biology*. (Oxford
University Press, USA, 1998).
98. Harmon, L. J. *et al.* Early bursts of body size and shape evolution are rare in comparative
data. *Evolution* **64**, 2385–2396 (2010).
99. Mollentze, N. & Streicker, D. G. Viral zoonotic risk is homogenous among taxonomic orders
of mammalian and avian reservoir hosts. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 9423–9430
(2020).
100. Schmidt, J. P. *et al.* Ecological indicators of mammal exposure to Ebolavirus. *Philos. Trans.*
R. Soc. Lond. B Biol. Sci. **374**, 20180337 (2019).
101. Pandit, P. S. *et al.* Predicting wildlife reservoirs and global vulnerability to zoonotic
Flaviviruses. *Nat. Commun.* **9**, 5425 (2018).
102. Evans, M. V., Dallas, T. A., Han, B. A., Murdock, C. C. & Drake, J. M. Data-driven identification
of potential Zika virus vectors. *Elife* **6**, (2017).
103. Yang, L. H. & Han, B. A. Data-driven predictions and novel hypotheses about zoonotic tick
vectors from the genus Ixodes. *BMC Ecol.* **18**, 7 (2018).
104. Elith, J., Leathwick, J. R. & Hastie, T. A working guide to boosted regression trees. *J. Anim.*
Ecol. **77**, 802–813 (2008).
105. Carlson, C. J. embarcadero: Species distribution modelling with Bayesian additive
regression trees in R. doi:10.1101/774604.
106. Chipman, H. A., George, E. I. & McCulloch, R. E. BART: Bayesian additive regression trees.
The Annals of Applied Statistics vol. 4 266–298 (2010).
107. Chen, W. *et al.* The illegal exploitation of hog badgers (Arctonyx collaris) in China: genetic
evidence exposes regional population impacts. *Conservation Genetics Resources* vol. 7
697–704 (2015).
108. Stock, M., Pahikkala, T., Airola, A., De Baets, B. & Waegeman, W. A Comparative Study of
Pairwise Learning Methods Based on Kernel Ridge Regression. *Neural Comput.* **30**, 2245–
2283 (2018).
109. Stock, M., Pahikkala, T., Airola, A., Waegeman, W. & De Baets, B. Algebraic shortcuts for
leave-one-out cross-validation in supervised network inference. *Brief. Bioinform.* (2018)
doi:10.1093/bib/bby095.
110. Stekhoven, D. J. & Bühlmann, P. MissForest—non-parametric missing value imputation for
mixed-type data. *Bioinformatics* **28**, 112–118 (2012).
111. Schölkopf, B., Smola, A. J., Managing Director of the Max Planck Institute for Biological
Cybernetics in Tubingen Germany Profe Bernhard Scholkopf & Bach, F. *Learning with*
Kernels: Support Vector Machines, Regularization, Optimization, and Beyond. (MIT Press,
2002).
112. Foley, N. M., Goodman, S. M., Whelan, C. V., Puechmaille, S. J. & Teeling, E. Towards
navigating the Minotaur's labyrinth: cryptic diversity and taxonomic revision within the
speciose genus Hipposideros (Hipposideridae). *Acta Chiropt.* **19**, 1–18 (2017).