Brain orchestration of pregnancy and maternal behavior in mice

David André Barrière[1,2#], Arsène Ella[1,3], Frédéric Szeremeta[4], Hans Adriaensen[1], William Même[4], Elodie Chaillou[1], Martine Migaud[1], Sandra Même[4], Frédéric Lévy[1], Matthieu Keller[1,#]

[1] *Physiologie de la Reproduction et des Comportements, UMR INRAE/CNRS/Université de Tours/IFCE, Nouzilly, France.*

[2] *Université Paris-Saclay, CEA, CNRS, BAOBAB, NeuroSpin, 91191, Gif-Sur-Yvette, France.*

[3] *MRC Cognition and Brain Sciences Unit, University of Cambridge, UK.*

[4] *Complexes Métalliques et IRM, Centre de Biophysique Moléculaire, UPR44301 CNRS, Orléans, France.*

[#]**Corresponding authors:**

matthieu.keller@inrae.fr and david.a.barriere@gmail.com

Abstract

Reproduction induces changes within the brain to prepare for gestation and motherhood. However, the dynamic of these central changes and their relationships with the development of maternal behavior remain poorly understood. Here, we describe a longitudinal morphometric neuroimaging study in female mice between pre-gestation and weaning, using new magnetic resonance imaging (MRI) resources comprising a high-resolution brain template, its associated tissue priors (60-µm isotropic resolution) and a corresponding mouse brain atlas (1320 regions of interest). Using these tools, we observed transient hypertrophies not only within key regions controlling gestation and maternal behavior (medial preoptic area, bed nucleus of the *stria terminalis*), but also in the amygdala, caudate nucleus and hippocampus. Additionally, unlike females exhibiting lower levels of maternal care, highly maternal females developed transient hypertrophies in somatosensory, entorhinal and retrosplenial cortices among other regions. Therefore, coordinated and transient brain modifications associated with maternal performance occurred during gestation and lactation.

Key words: Gestation, lactation, maternal brain, MRI, atlas, voxel-based morphometry.

<u>Abbreviations:</u>

AC-PC: anterior commissure-posterior commissure

AMBMC: Australian Mouse Brain Mapping Consortium

AOB: accessory olfactory bulb

BNST: bed nucleus of the *stria terminalis*

CNS: central nervous system

CSF: cerebrospinal fluid

DARTEL: diffeomorphic anatomical registration using exponentiated lie algebra

df: degree of freedom

FA: flip angle

FLASE: fast large-angle spin-echo

FoV: field of view

MRI: magnetic resonance imaging

MOB: main olfactory bulb

mPOA: medial preoptic area

GM: gray matter

GMC: gray matter concentration

PVN: paraventricular nucleus of the hypothalamus

RARE: rapid acquisition with relaxation enhancement

ROC: receiver operating characteristic

ROI: region of interest

TE: echo time

TR: repetition time

VBM: voxel-based morphometry

WM: white matter

Introduction

Motherhood is among the most transformative experiences in the lives of female mammals. While virgin females tend to avoid neonates, the end of the gestation period and the birth process lead to a behavioral switch characterized by an attraction towards infant cues, the expression of nurturing behavior and ultimately the establishment of infant bonding[1,2]. Decades of scientific research dedicated to the maternal brain have revealed a core neural circuitry that includes the medial preoptic area (mPOA) and the adjoining ventral part of the bed nucleus of the *stria terminalis* (BNSTv), and that is highly critical for the onset of maternal behavior[1,3–5]. Functional modulations of the mPOA/BNSTv consistently disrupt maternal motivation and expression in numerous species[1,2,6]. This core maternal circuitry regulates maternal behavior through its direct projections to the ventral tegmental area, which promotes reward system activation[1,7], as well as through its connections with cortical regions, including the prefrontal cortex[8–10]. This crucial central circuitry is finely regulated by multiple neural networks that integrate both internal and external stimulations. The proper expression of maternal care towards offspring is prepared through the neuroendocrine action of sex steroids and neuropeptides such as oxytocin among others during the gestation period[11]. These internal factors induce rewiring of the maternal brain, including through structural plasticity through increasing neuronal soma size or astrocytic complexity within the mPOA[12], and changes in neurogenesis mainly in the main olfactory bulb (MOB)[13] but also in the mPOA/BNST in rodents[14]. In human, regional morphological changes of gray matter (GM) within the parahippocampal gyrus, precuneus, cingulate, insula and frontal cortex have been observed in primiparous women using magnetic resonance imaging (MRI)[15]. Additionally, olfactory cues coming from the neonate are integrated by the MOB and the accessory olfactory bulb (AOB) through an amygdalo-hypothalamic pathway, which is responsible for attraction/repulsion behavioral outcomes[16]. Hence, the development of the maternal brain is

dependent on both integration of external and internal cues acting through multiple brain

pathways and regions to prepare the brain to gestation and motherhood. Nevertheless, the

relationship between the brain rewiring over the gestation and lactation periods and the

establishment of the maternal behavior is poorly documented.

To assess the dynamics of the maternal brain, a longitudinal MRI morphometric study

over a complete reproductive experience was performed in mouse to investigate changes in

the gray matter concentration (GMC) using voxel-based morphometry (VBM). VBM is a

well-established and well-validated image analysis technique that provides an unbiased and

comprehensive assessment of anatomical differences throughout the brain, and has been

successfully used to study GM changes within the mouse brain[17–21]. However, available

mouse brain MRI resources are often partial or provided in different spatial orientations or

spatial resolutions (**Table 1**). As an example, the <u>Australian Mouse Brain Mapping</u>

<u>Consortium</u> (AMBMC), offers a high resolutive template and detailed atlases of the mouse

brain including the cerebellum[22], hippocampus[23], diencephalon[24] and cortices[25].

Unfortunately, segmentation of the MOB and the AOB and hindbrain is lacking, and this

resource does not provide associated tissues probabilistic maps necessary for VBM. In other

hand, the <u>Allen Mouse Brain Common Coordinate Framework</u>, is the most advanced mouse

brain atlas[26]. This new atlas delimitates discrete structures within the thalamus, hindbrain,

olfactory system and diencephalon and provides a full segmentation of cortical layers

however, MRI template and brain tissues priors are still lacking for VBM investigations.

Thus, we combined these resources and associated probabilistic maps to emulate a

complete resource dedicated to the mouse brain. Using these resources and a longitudinal

VBM approach, we were able to assess the dynamic morphological changes of the brain

during the whole reproductive period and demonstrate how these changes predict the quality

of maternal behavior.

<u>Results</u>

Mouse MRI atlas

VBM strategies require a template image and its associated priors of gray matter

(GM), white matter (WM) and cerebrospinal fluid (CSF) for brain segmentation and

normalization. In addition, a complete atlas of the mouse brain is mandatory for the

identification of regions of interest (ROIs) highlighted by the VBM analysis. Given the

limitations of available tools to thoroughly study GMC changes during the gestation and

lactation periods in mice, we developed first a new set of resources using the AMBMC, an

ultra-high-resolution template built from *ex vivo* brain images finely normalized within the

same space[25] and the <u>Allen Mouse Brain Common Coordinate Framework</u>[26] (**Figure 1**). Our

resources comprise the following four components: 1) a complete mouse brain template with

a spatial resolution suitable for mouse brain analysis (60-µm isotropic resolution); 2) the

corresponding GM, WM and CSF probabilistic maps for brain normalization together with a

VBM analysis built from 138 T_2-weighted images; 3) a complete mouse brain atlas derived

from Paxinos and Franklin's mouse brain atlas[27] and composed of a mosaic of 1320 ROIs

(**Figure 2A**); 4) a brain mesh permitting brain plot generation and data visualization (**Figure**

2B and **Supplemental Video 1**). We visually inspected and carefully checked the results of

the normalization process against the original coregistered atlas. Then, the labeled structures

were reclassified and aggregated according to the brain regions to which they belonged

(auditory, insular, temporal cortices, etc.), with respect to their anatomical topography (cortex,

basal ganglia, etc.), tissue type (GM, WM and CSF) and hemisphere (left or right). Cortical

structures were subdivided into functional (*e.g.,* primary and secondary motor cortices) or

structural (agranular, dysgranular, agranular/dysgranular, granular and posterior agranular

insular cortices) areas (**Figure 2C**), as well as into different cortical layers (**Figure 2D**).

Subcortical structures, as for example, the hypothalamus (**Figure 2E**) and the hippocampus

(**Figure 2F**), were fully segmented according to Paxinos and Franklin's atlas[27].

Morphometric changes occurred during the gestation and lactation periods

Next, we used our new resources to study the variations of GMC in mouse brain from

the beginning of gestation until weaning. Using MRI T_2-weighted anatomical acquisitions, we

estimated the GMC maps which offer for each animal a global estimation of the GM.

Longitudinal comparison of GMC between virgin mice (control group, n= 11) and mice who

became pregnant and raised their young until weaning (parous group, n=12) permits to

highlight local modifications of GM during the whole reproductive cycle. A comparison of

baseline and early gestation GMC maps between the control and parous groups did not reveal

significant differences. However, at the end of the gestation period, significant increases in

GMC were observed within several brain regions in the parous group compared to the control

group (**Table S1** and **Figure 3A**). Time course analysis revealed differences in GMC profiles

between control and parous groups at the end of the gestation period, early in lactation and at

the end of the lactation period. Specifically, GMCs within the mPOA and the BNST were

consistently significantly higher in the parous group at that times. In addition, within the

agranular insular cortex in the late gestation period and the early lactation period, GMC was

significantly higher in parous group compare to control group (**Figure 4A**). During the early

lactation period, we also found specific and significant increases in GMCs of the parous group

within numerous brain regions (**Table S2** and **Figure 3B**). Among these structures, the

hippocampus (CA1 layer), amygdalar area and piriform area showed a transient increase in

GMC at the early lactation time point that returned to baseline values at the end of the

lactation period in parous group compared to control group (**Figure 4B**). In contrast, the

caudate putamen, arcuate nucleus and paraventricular nucleus of the hypothalamus (PVN)

showed significantly higher GMCs in the parous mice than in the control mice during the

lactation period (**Table S3**and **Figures 3C and 4C**). Together, our data demonstrate that the

late gestation period is associated with a pronounced increase in GMC in the mPOA/BNST,

the core neural system of maternal motivation, lasting up to the late lactation period.

Furthermore, the early lactation period is associated with increased GMC in other key

maternal motivation areas in midbrain regions, including the hypothalamus, caudate putamen

and amygdala. These GMC differences between both groups were no longer observed after

weaning.

Morphometric changes during gestation predict the quality of maternal behavior

In the last part of this work, we evaluated whether these morphometric changes might

reflect differences in maternal performance. Based on 15 min of behavioral observation

during the pup retrieval test performed one week after birth, we evaluated the maternal

performance of each mother by measuring the first, second and third pup retrieval times, pup-

licking duration, crouching time, rearing time, digging time and self-grooming time (**Figure

5A**). Interestingly, we observed a large distribution of values for both crouching (284.2 s, SD

$\pm$ 268.7 s) and digging (98.22 s, SD $\pm$ 159.3 s) durations within the parous group. Whereas

crouching duration relates to maternal behavior, digging duration is widely recognized as a

discriminative marker of stress-related behavior in rodents[28]. Hence, we used crouching and

digging durations to cluster animals using the k-means clustering procedure, thereby

clustering parous animals into a high maternal behavior group (those with a high crouching

time and low digging time) and a low maternal behavior group (those with low crouching

time and high digging time).

A comparison of maternal performance parameters between the two clustered

groups revealed significantly lower crouching and pup-licking times and a significantly higher

digging time (**Figure 5B**) in the low maternal behavior group (n=6) than in the high maternal

behavior group (n=6). Moreover, a comparison of GMC maps revealed both cortical and

subcortical differences between the two clustered groups, mainly in the late gestation period

(**Table S4** and **Figure 5C**) but also during the early lactation period (**Table S5** and **Figure**

5D). Indeed, transient increases in GMCs within the entorhinal area, lateral part of the orbital

area, AOB, and medial preoptic area hat were observed at the end of pregnancy in the high

maternal behavior group were absent in the low maternal behavior group (**Figure 6A**). In

addition, the high maternal behavior group showed consistent higher GMCs in the

hippocampus, retrosplenial area and barrel field of the primary somatosensory cortex (**Figure**

6B) from the end of the gestation until the end of the lactation period.

Interestingly, using a receiver operating characteristic (ROC) analysis, we found that

GMCs values within the entorhinal area and AOB at late gestation are reliable predictors for

mouse maternal performance after birth. These GMC values significantly distinguished low

maternal performance from high maternal performance postpartum (entorhinal area:

sensitivity = 100, confidence interval (CI) = 61% to 100%; specificity = 83, CI = 44% to

99%; likelihood ratio = 6; **Figure 7A**; AOB: sensitivity =100, CI = 61% to 100%; specificity

= 83, CI = 44% to 99%; likelihood ratio = 6, **Figure 7B**). The GMC values of both the

entorhinal area and AOB observed at late gestation were also significantly correlated with

maternal behavior (crouching and digging times). These results reveal that the GMC

differences in olfactory (AOB and entorhinal cortex) and mnesic (entorhinal area)-related

brain regions occurring during the late gestation period significantly predicted the quality of

maternal behavior.

Discussion

Using a new comprehensive neuroimaging resource dedicated to mouse brain, this longitudinal study reveals that pregnancy and lactation coincide with pronounced and transient cerebral changes. Transient increases in GMCs were observed in key regions controlling maternal behavior (mPOA, BNST, and PVN), as well as regions involved in emotions (amygdala), in motivation and reward (caudate nucleus, orbitofrontal cortex) and in mnesic functions (hippocampus). Interestingly, increase in GMC was also revealed in the insular cortex thought to link social and emotional skills. Moreover, we showed that females expressing high levels of maternal behavior had developed specific increases in GMCs in structures involved in olfactory (MOB and AOB) and somatosensory (somatosensory cortex) information processing, in memory (hippocampus, entorhinal cortex, retrosplenial cortex) and in reward and reinforcement (striatum). Interestingly, these hypertrophies were already significant at the end of the gestation period thus being predictive of the quality of maternal care (**Supplemental Video 2**).

Implementation of new resources to support the analysis of mouse brain MRI data

The use of preclinical MRI is a target of growing interest for the study of brain structure and function in both healthy and pathological conditions. The use of advanced MRI techniques, coupled with the development of advanced animal models, is a powerful way to push new breakthroughs in the understanding of brain functioning and pathology. Herein, in the first step in our study, from the recent major advances in the development of brain mouse atlas we generated a new set of neuroinformatic tools offering for the first time a complete resource dedicated to MRI studies of the mouse brain, namely, an accurate brain atlas (1320 ROIs), a high-resolution brain template and the associated GM, WM and CSF priors (60-μm isotropic resolution). The GM, WM and CSF probabilistic maps built and used in this study

were calculated from 138 T_2-weighted anatomical images, resulting in robust tissue class

priors not only for VBM analysis but also for functional MRI and diffusion tensor imaging

analysis in mice.

This comprehensive set of MRI compatible template and atlas for the mouse brain,

allows a unified and standardized analysis of multimodal mouse brain MRI data and paves the

way for the development of multicentric preclinical studies. Indeed, animal models deliver

crucial information for the understanding of brain structure and function both in healthy and

pathological conditions. Our template and mouse brain atlas were conceived to bridge the gap

between basic and clinical neurosciences by providing to the preclinical neuroimaging

community specific resources designed to be used in conjunction with the neuroinformatic

tools and methodologies commonly used in human MRI studies. We anticipate that these

resources will help neuroscientists to conduct their analyses of anatomical and functional

datasets in a more standardized way, with the final goal of reaching more reproducible

conclusions (https://www.nitrc.org/projects/tmbta_2019).

Gestation and lactation periods induce strong but transient GMC hypertrophy

The establishment of accurate mouse MRI resources permit to study the variations of

GMC longitudinally, *in vivo* and during the gestation and lactation periods in female mice.

We observed that several brain regions became transiently hypertrophic during pregnancy or

in the lactation period until weaning. A set of structures comprising the core of the maternal

circuit – especially the mPOA and BNST – displayed long-lasting hypertrophy that started at

late gestation, culminated during the first week of lactation, and then disappeared at weaning.

The initial GMC increase observed during the gestation period probably reflects changes

induced by hormonal priming[11]. Indeed, both the mPOA and BNST express a high number of

steroid hormone and neuropeptide receptors[29,30]. These factors are well known to trigger

significant plasticity changes within the core maternal circuitry that are necessary for the

preparation and adaptation of the brain to motherhood[3,11]. From parturition, the GMC

differences observed during the whole lactation period highlight that mPOA/BNST receives a

variety of sensory inputs from the pups, integrates that information with the females'

endocrine status, and then projects to brain sites involved in socially-relevant motivation,

affective state, and cognition[2,31]. Pup stimulation, electrolytic and neurotoxic lesions and local

steroid hormone injections[32–35] have been shown to modify the intrinsic activity of these

nuclei and consequently responsible for motivation and expression of maternal behavior. The

mPOA is engaged throughout the postpartum period but differentially according to the needs

of the developing pups. It has been shown that neurons of the mPOA in late postpartum

inhibits maternal responses allowing the changing expression and waning of maternal

behavior across postpartum[10]. The sustained increase in GMC reported in late lactation could

reflect the involvement of the mPOA to appropriately influence maternal behavior.

Our study highlights another set of brain structure that became hypertrophic only

during the period from parturition to weaning – specifically, the PVN and arcuate nucleus of

the hypothalamus. These changes illustrate the structural plasticity occurring in these regions.

For instance, at parturition and during lactation oxytocin neurons of the PVN undergo

dramatic neuronal, glial and synaptic changes such as an increase in size of the oxytocin

neurons and an amplification of their synaptic input[36]. Oxytocin release at parturition

facilitates the onset of maternal behavior by acting on the mPOA and is also important for

maternal memory[37]. Finally, lesions of the PVN disrupt the onset of maternal behavior[38–40]. In

the arcuate nucleus, dopaminergic cells are responsible for suckling induced prolactin

release[41] and neurons projecting to the arcuate nucleus are involved in the maintenance of

maternal motivation[42].

Additionally, we report changes in GMCs found in olfactory related structures (main

olfactory bulb, piriform cortex), somatosensory areas and auditory areas which reflect the

multisensory control of maternal behavior. This finding is in accordance with a functional

MRI study performed in rats which demonstrates that pup suckling is associated with

increased neuronal activity within the midbrain, striatum and cortical sensory areas

(somatosensory, olfactory and auditory cortices)[9].

GMC variations within the hippocampus and entorhinal cortex, highlight the role of

two essential structures involved in learning and memory processing during the reproductive

period. Our data support evidence that the hippocampus undergoes profound neural changes

during lactation. Indeed, lactating females have elevated spine densities in the hippocampus[43]

and show significant dendritic remodeling in pyramidal neurons[44]. Changes in hippocampal

neurogenesis occurs during lactation and may support the enhancement of spatial memory

necessary to foraging behavior in lactating females[43,45–47].

Finally, our study also revealed a hypertrophy of the agranular insular cortex, which

has never been reported in this context. The agranular insular cortex is a laminar part of the

insular cortex and can be considered as a hub structure linking large-scale brain systems[48].

Indeed, the insula receives direct thalamic and somatosensory afferents carrying sensitive

stimulations. In addition to its sensory afferents, the insula displays structural connectivity

with the limbic system (basolateral, lateral and central amygdalar nuclei) as well as with the

BNST, mediodorsal nucleus of the thalamus, lateral hypothalamus and perirhinal and lateral

entorhinal cortices[48]. The insula also connects brain regions implicated in motivation and

reward, such as the nucleus accumbens and caudate putamen[48]. Hence, our findings and the

current literature suggest that before and after birth, the insular cortex may integrate and

combine information from both external and internal stimulation and act as a relay between

higher cortical and subcortical structures. Together, our results describe the dynamics of

neurophysiological adaptation occurring in the brain from the early gestation period to weaning, thereby ensuring efficient maternal behavior and, by extension, the development of the offspring.

GMC modifications in the olfactory system at the end of the gestation predict the level of maternal behavior post-partum

All these transient modifications of GMC in parous animals indicate that various brain regions undergo considerable plasticity from birth to weaning. Then, we sought to determine whether inter-individual variations in maternal behavior were associated with similar variations of GMC. Based on their behavioral performance in the pup retrieval test, we used a k-means clustering strategy to divide maternal female mice into two groups displaying high *versus* low levels of maternal behavior. This analysis revealed that several transient and several long-lasting increases in GMCs were observed in the high maternal behavior group that were absent in the low maternal behavior group. Brain regions showing significant differences included the olfactory bulbs, somatosensory system, limbic system, especially the orbitofrontal area, and mnesic system, including the retrosplenial cortex, hippocampus and entorhinal area. Some of these structures are directly responsive to pup stimulation, and the observed dynamics may have been induced by mother-offspring interactions. For example, higher GMC values in the somatosensory cortex and olfactory bulbs in the high maternal behavior group potentially reflected increased suckling duration and proximity between the mother and pups, respectively.

Strikingly, differences in GMCs were detected in the entorhinal area, orbitofrontal area, olfactory bulb, hippocampus, retrosplenial area and primary somatosensory area before parturition. These findings suggest that the maturation of these structures, probably through hormone-dependent plasticity mechanisms, is a key determinant of the intensity of maternal behavior expressed during the lactation period. Using a ROC procedure, we found that GMC

values of the entorhinal area and AOB at the end of gestation were significantly predictive of

the maternal behavior postpartum. Interestingly, previous studies in mice reported an increase

in cell proliferation during gestation in the subventricular zone, the neurogenic niche which

provides newly generated neurons within the olfactory bulb (for review see [13]). These adult-

born olfactory neurons are fully responsive to pup odor exposure[49] and are in part involved in

some components of maternal behavior[50,51].

Hence, the correlations observed between GMC values in AOB at the late gestation

period and maternal behavior performances may suggest that impairments of neural plasticity

of this olfactory region would induce maladaptive neuroendocrine processing of the maternal

brain at the end of the gestation period impacting maternal behavior performance. Taken

together, our data provide the first potential imaging-based predictive biomarkers of the

quality of maternal behavior and suggest the key role of the maturation of the olfactory

system at the end of the pregnancy in the development of adaptative maternal behavior in

mice.

Conclusion

Our study provides a new generation of neuroinformatic tools which will help basic

neuroscientists to conduct structural and functional MRI investigations. Using these

resources, we found that the development of the maternal brain is associated with substantial

mesoscopic changes in critical regions. These modifications can be interpreted as cell size

changes, neural or glial cell genesis/apoptosis, spine density or blood flow modifications[52–54].

As cellular and molecular plasticity events are key for the adaptation of the brain to

motherhood, molecular, cellular and behavioral investigations must be performed to obtain a

more precise view of the physiological mechanisms responsible for GMC variations.

Methods

Animals. Twenty-three female RjOrl:SWISS virgin mice (8 weeks old; 20-25 g; Janvier Laboratory, Le Genest-Saint-Isle, France) were maintained on a 12-h light/dark cycle with access to food (standard chow) and water *ad libitum*. Animals were acclimatized 6 per cage to the housing facility for 7 days prior to manipulation. Females were randomly divided into two groups: a parous group (n = 12), in which each female was exposed to a male (RjOrl:SWISS,8 weeks old; 20-25 g; Janvier Laboratory, Le Genest-Saint-Isle, France) for 5 days, became pregnant, and raised their offspring (litter size: 6 to 14 pups) until weaning at 21 days postpartum, and a control group (n = 11), in which virgin females were not exposed to male mice. Each parous female was individually housed after male exposure. Control females were housed together in a separate room from parous females.

The MRI protocol was optimized to keep mice anesthetized for 2 h during each of the six acquisitions. During lactation MRI acquisitions, pups were kept under a heat lamp. One week after birth, maternal behavior was assessed as described in the behavioral section. All experiments were conducted in accordance with the local research ethics committee (APAFIS #6626-201002281145814V1) and are reported in accordance with the ARRIVE guidelines.

MRI acquisition. In vivo 3D MRI of the entire brain was performed three days before male exposure (baseline), at one week of gestation (early gestation), two days before the expected day of birth (late gestation), one week postpartum (early lactation), three weeks postpartum (late lactation) and two weeks after weaning (weaning). One female in the parous group and one female in the control group were scanned under similar conditions on the same day.

Mice were anesthetized using isoflurane (2.5%; induction in O_2/air mixture 1:1) (TEM-SEGA, F-33600 Pessac, France) and then transferred and placed head first *procubitus* within an MRI-compatible cradle that incorporated a stereotaxic system dedicated for mouse

head MRI, connected to a heater with circulating water to maintain body temperature and

supplied with 1-2% isoflurane *via* a fitted mask. Respiration rate was recorded during all the

experiments using an MRI-compatible monitoring system (PC-SAM model #1025; SA

Instruments Inc., Stony Brook, NY, USA) and used to adjust the isoflurane rate to maintain a

rate between 20 and 40 respirations per minute. After a recovery period of one hour, mouse

returned to her pups. MRI studies were conducted at the Centre de Biophysique Moléculaire

d'Orléans and were performed on a 7T/160 mm PharmaScan spectrometer (Bruker Biospin,

Wissembourg, France) equipped with an actively shielded B-GA09 gradient set, with 90-mm

inner diameter and 300-mT/m gradient intensity. A 23-mm inner diameter Bruker birdcage

coil with a cradle dedicated to a mouse head was used. Data acquisitions were performed on

an Advance III console running ParaVision 5.1 software. T_2-weighted images were acquired

using a 3D fast large-angle spin-echo (FLASE) sequence which allows 3D brain mapping

with a high resolution in a suitable time for *in vivo* acquisition[55,56]. Thus, the sequence with

echo time (TE) = 20 ms, 1 repetition, acquisition matrix =160 × 140 × 95, and a field of

view (FoV) of 19.2 × 16.8 × 11.4 mm^3, resulting in a final resolution of 120 µm isotropic

voxels[55,56]. To obtain the FLASE sequence[55], which is a specific sequence that is not included

in the sequence package provided with ParaVision, the usual rapid acquisition with relaxation

enhancement (RARE) spin-echo sequence was modified; in particular, the RARE-factor was

fixed to 1 allowing a flip angle (FA) higher than 90° for the excitation pulse, while

maintaining a 180° refocusing pulse[55]. Thus, T_2-weighted images were obtained with a

repetition time (TR) as short as 300 ms, 10 times lower than that needed for a classical T_2-

weighted spin-echo sequence. The sequence was optimized for acquisition in 1 h 28 min, with

an isotropic resolution of 120 µm, a TR of 300 ms, an TE of 20 ms, an excitation pulse (FA)

of 115°, with 1 repetition and with a matrix of 160 x 140 x 95 corresponding to a FoV of 19.2

x 16.8 x 11.4 mm^3 and contained the whole mouse brain.

Maternal behavioral test. One week after birth, the maternal behavior of each female was

evaluated using the pup retrieval test[57]. Briefly, three pups were removed from the nest and

placed at three different corners within the home cage. The latency to retrieve each pup and

the time spent licking the pups, crouching in the nest over the pups and performing

nonmaternal behaviors such as self-grooming and digging were recorded over 15 min.

Retrieval was defined as the animal picking up a pup and transporting it to the nest.

Crouching was defined as the animal assuming the nursing posture. Nursing and licking were

permitted whether they took place in the nest. All videos were analyzed using BORIS

software version 4.1.4.

K-means clustering. Clustering analysis of the behavioral data was performed using

MATLAB Simulink 10b (The Mathworks, Inc., USA). Normality was verified and no outlier

subjects were detected. To classify animals according to their maternal performance, a k-

means clustering algorithm was used with crouching and digging times as behavioral markers.

Digging was chosen because it is indicative of high maternal stress[28,58]. This algorithm

iteratively grouped the animals by creating k initial centroids, assigning each animal to the

closest centroid, iteratively re-calculating the centroids from the mean of its assigned animals

and re-assigning the animals to each centroid until there were no more changes across

iterations[59]. This clustering divided parous animals into a high maternal behavior group (with

high crouching and low digging time) and low maternal behavior group (with low crouching

and high digging time).

Mouse brain template and atlas building. For the MRI protocol, we developed a brain

template and an atlas from the AMBMC brain template and the Allen Mouse Brain Common

Coordinate Framework, respectively (**Figure 1**). First, we down-sampled the AMBMC

template and its associated atlases and the Allen Mouse Brain Atlas and its associated Nissl

images to a suitable resolution for MRI analysis (60-µm isotropic resolution; **Figure 1**, step

1). Then, all images were manually aligned to the anterior commissure/posterior commissure

(AC/PC) axis, and the center of the images was defined relative to the AC (**Figure 1**, step 2).

The resulting template was then segmented into GM, WM and CSF probability maps using

the unified segmentation approach[60] of Statistical Parametric Mapping 8 (SPM8) and the

mouse brain priors provided by the SPMMouse toolbox (**Figure 1**, step 3). In parallel, our T_2-

weighted anatomical images were realigned, coregistered, bias-corrected and normalized to

our template. Using the SPMMouse toolbox, we also segmented the images as described

above, and from these preprocessed images, we obtained a large set of 138 images for each

tissue class (**Figure 1**, steps 4-7). From these images, we built population-specific GM, WM

and CSF priors. To build these priors, for each tissue class, we applied a diffeomorphic

anatomical registration using an exponentiated lie algebra (DARTEL) approach, which is an

automated, unbiased, and nonlinear template-building algorithm[61] (**Figure 1**, step 8). This

new set of population-specific tissue priors was used for both atlas building and final VBM

preprocessing.

To normalize the Allen Mouse Brain Atlas to our brain template, we used the

associated Nissl-stained images because (1) Nissl staining corresponds to the GM prior in

terms of histology, and (2) this image was already coregistered to the atlas (**Figure 1**, step 4).

Therefore, we applied the segmentation function provided by SPM8 using the GM prior

previously calculated from the Nissl image to generate the "Nissl2template" normalization

matrix. We used this matrix to normalize the atlas to the template, while avoiding

interpolation to maintain the label indices as integers (**Figure 1**, step 7). Then, a visual

inspection of each normalized label was carried out to assess whether the normalization

process modified the position and volume of the structure too much. When necessary, holes

were filled and labels were redrawn according to Paxinos and Franklin's atlas and using the

FreeSurfer package. Finally, the olfactory bulbs and hind brain regions were completed, the

corpus collosum and ventricles were drawn from the WM and CSF priors, and the cerebellum

labels were replaced by the AMBMC cerebellum labels, which are more accurate. Finally, the

atlas image was symmetrized (left-right). Our mouse brain template, priors and atlas were

normalized within the same space and with the same final resolution (60-µm isotropic

resolution), resulting in our final mouse brain atlas composed of a mosaic of 1320 ROIs

covering the entire brain (**Supplemental Video 1**).

VBM data preprocessing. Previously preprocessed normalized T_2-weighted data were

segmented into GM, WM, and CSF within SPM8 using the images of the population-specific

priors (**Figure 1**, step 9). Then, to produce a more accurate registration within each mouse as

well as across all mice, a longitudinal VBM analysis was applied using the strategy described

by Asami *et al*[62]. First, a *subject-specific* template was created by the DARTEL algorithm

using the previous tissue class images (*i.e.*, GM, WM, and CSF maps) obtained from each

mouse for the six time points. The DARTEL procedure releases *individual-specific* flow field

maps, permitting the application of diffeomorphic normalization on images of each tissue

class to spatially normalize each time point on a *subject-specific* template space. Each

normalized tissue class image was modulated by the Jacobian determinant to account for the

expansion and/or contraction of brain regions over time. Then, a *population-specific* template

was created by the DARTEL algorithm using all *subject-specific* templates of the tissue class

images. Here, the DARTEL procedure releases *population-specific* flow field maps,

permitting the application of diffeomorphic normalization of each animal onto the images of

each tissue class. Finally, tissue class images were modulated by the Jacobian determinant,

and the final modulated GM images were spatially smoothed with an isotropic Gaussian

kernel with a 3-mm full-width at half-maximum and convolved with GMC images to create

GMC maps (**Figure 1**, steps 10-12).

VBM statistics and analysis. SPM8 was used to reveal the temporal and regional changes in

the GMC maps. A second-level SPM analysis comprising a flexible factorial model, which is

equivalent to a 2×2 mixed-model ANOVA with group as the between-subject factor and time

point as the within-subject factor, was used to compare the control *versus* the parous groups

and the low *versus* high maternal behavior groups[63]. The factors included in the analysis were

subjects, group (control *versus* parous, or low maternal behavior *versus* high maternal

behavior), and time points (baseline, early gestation, late gestation, early lactation, late

gestation, and weaning). A brain mask was used to constrain the analysis within the brain. For

each cluster, the significance of the peak voxel was set as $p < 0.01$ ($t_{(126)} = 2.356$, control

versus parous; $t_{(60)} = 2.39$, low *versus* high maternal behavior), and the minimum cluster

extent was set at 25 voxels. The results are presented on axial brain slice series generated by

the Xjview SPM plugin. Corresponding surfacing results were produced with BrainNet

viewer 1.6[64], allowing the generation of both brain meshes and brain plots to visualize data

and create supplemental videos.

Postprocessing statistical analysis. Cluster peaks revealed by the flexible factorial analyses

were identified using our atlas and an *in situ* procedure developed with MATLAB Simulink

10b (The Mathworks, USA). For each comparison, clusters were binarized, and the obtained

masks were used to extract GMC values of corresponding regions from the GMC map using

the REX plugin. GMC data and behavioral measurements were then compiled and analyzed

using GraphPad Prism 6.02 software. GMCs were compared between groups and for each

time point using a two-way ANOVA with repeated measures followed by a two-stage setup

method of Benjamini, Kriegger and Yekutieli as recommended by the software. Maternal and

nonmaternal behaviors of the low maternal and high maternal groups were compared using a

multicomparison t-test with a false discovery rate (FDR) approach (Q = 1%). Correlation

analyses were performed using a parametric two-tailed Pearson test. Specificity and

selectivity analyses were performed using the ROC curve method. Statistical significance was

defined as $p < 0.05$ (*) for these analyses.

Acknowledgments

The authors acknowledge the regional council of Centre Val-de-Loire for funding this research through the IMACERVOREPRO grant (convention 201500104011, 2015-2018) awarded to Matthieu Keller.

Author Contributions

D.A.B. contributed to the atlas and template building, data analysis, drafted and revised the manuscript. **A.E.** contributed to the imaging sequence troubleshooting, data analysis and contributed to the critical revisions. **F.S.** contributed to MRI sequence programming, imaging protocol setting, data acquisitions and contributed to the analysis and critical revisions. **H.A.**, **W.M.**, **E.C.**, **M.M.**, **S.M.** contributed to the study conception and design and contributed to the critical revisions of the manuscript. **F.L.** contributed to raise the funding, study conception and design and contributed to the critical revisions of the manuscript. **M.K.**, is the principal investigator of the study, raised the funding, coordinated the project, revised and validated the manuscript.

References

1. Numan, M. & Insel, T. R. *The neurobiology of parental behavior*. (Springer, 2003).

2. Lonstein, J. S., Lévy, F. & Fleming, A. S. Common and divergent psychobiological mechanisms underlying maternal behaviors in non-human and human mammals. *Horm. Behav.* **73**, 156–185 (2015).

3. Bridges, R. S. Neuroendocrine regulation of maternal behavior. *Front. Neuroendocrinol.* **36**, 178–196 (2015).

4. Kohl, J. & Dulac, C. Neural control of parental behaviors. *Curr. Opin. Neurobiol.* **49**, 116–122 (2018).

5. Kohl, J., Autry, A. E. & Dulac, C. The neurobiology of parenting: A neural circuit perspective. *BioEssays* **39**, e201600159 (2017).

6. Lévy, F. & Keller, M. Olfactory mediation of maternal behavior in selected mammalian species. *Behav. Brain Res.* **200**, 336–345 (2009).

7. Numan, M. & Stolzenberg, D. S. Medial preoptic area interactions with dopamine neural systems in the control of the onset and maintenance of maternal behavior in rats. *Front. Neuroendocrinol.* **30**, 46–64 (2009).

8. Afonso, V. M., Sison, M., Lovic, V. & Fleming, A. S. Medial prefrontal cortex lesions in the female rat affect sexual and maternal behavior and their sequential organization. *Behav. Neurosci.* **121**, 515–526 (2007).

9. Febo, M. A bold view of the lactating brain: functional magnetic resonance imaging studies of suckling in awake dams. *J. Neuroendocrinol.* **23**, 1009–1019 (2011).

10. Pereira, M. & Morrell, J. I. Functional Mapping of the Neural Circuitry of Rat Maternal Motivation: Effects of Site-Specific Transient Neural Inactivation: Mapping maternal motivation circuitry. *J. Neuroendocrinol.* **23**, 1020–1035 (2011).

11. Brunton, P. J. & Russell, J. A. The expectant brain: adapting for motherhood. *Nat. Rev. Neurosci.* **9**, 11–25 (2008).

12. Kinsley, C. H. & Lambert, K. G. Reproduction-Induced Neuroplasticity: Natural Behavioural and Neuronal Alterations Associated with the Production and Care of Offspring. *J. Neuroendocrinol.* **20**, 515–525 (2008).

13. Lévy, F., Gheusi, G. & Keller, M. Plasticity of the Parental Brain: A Case for Neurogenesis: Parental behaviour and neurogenesis. *J. Neuroendocrinol.* **23**, 984–993 (2011).

14. Akbari, E. M., Chatterjee, D., Lévy, F. & Fleming, A. S. Experience-dependent cell survival in the maternal rat brain. *Behav. Neurosci.* **121**, 1001–1011 (2007).

15. Hoekzema, E. *et al.* Pregnancy leads to long-lasting changes in human brain structure. *Nat. Neurosci.* **20**, 287 (2017).

16. Kohl, J. *et al.* Functional circuit architecture underlying parental behaviour. *Nature* **556**, 326–331 (2018).

17. Ashburner, J. & Friston, K. J. Unified segmentation. *NeuroImage* **26**, 839–851 (2005).

18. Sawiak, S. J., Wood, N. I., Williams, G. B., Morton, A. J. & Carpenter, T. A. Voxel-based morphometry with templates and validation in a mouse model of Huntington's disease. *Magn. Reson. Imaging* **31**, 1522–1531 (2013).

19. Sawiak, S. J., Picq, J.-L. & Dhenain, M. Voxel-based morphometry analyses of in vivo MRI in the aging mouse lemur primate. *Front. Aging Neurosci.* **6**, (2014).

20. Keifer Jr, O. P. *et al.* Voxel-based morphometry predicts shifts in dendritic spine density and morphology with auditory fear conditioning. *Nat. Commun.* **6**, (2015).

21. Lein, E. S. *et al.* Genome-wide atlas of gene expression in the adult mouse brain. *Nature* **445**, 168–176 (2007).

22. Ullmann, J. F. P. *et al.* Segmentation of the C57BL/6J mouse cerebellum in magnetic resonance images. *NeuroImage* **62**, 1408–1414 (2012).

23. Richards, K. *et al.* Segmentation of the mouse hippocampal formation in magnetic resonance images. *NeuroImage* **58**, 732–740 (2011).

24. Watson, C. *et al.* An ontologically consistent MRI-based atlas of the mouse diencephalon. *NeuroImage* **157**, 275–287 (2017).

25. Ullmann, J. F. P., Watson, C., Janke, A. L., Kurniawan, N. D. & Reutens, D. C. A segmentation protocol and MRI atlas of the C57BL/6J mouse neocortex. *NeuroImage* **78**, 196–203 (2013).

26. Wang, Q. *et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D Reference Atlas. *Cell* **181**, 936-953.e20 (2020).

27. Paxinos and Franklin's the Mouse Brain in Stereotaxic Coordinates, Compact - 5th Edition. https://www.elsevier.com/books/paxinos-and-franklins-the-mouse-brain-in-stereotaxic-coordinates-compact/franklin/978-0-12-816159-3.

28. Deacon, R. M. J. Digging and marble burying in mice: simple methods for *in vivo* identification of biological
impacts. *Nat. Protoc.* **1**, 122 (2006).

29. Tsuneoka, Y. *et al.* Neurotransmitters and neuropeptides in gonadal steroid receptor-expressing cells in
medial preoptic area subregions of the male mouse. *Sci. Rep.* **7**, 1–16 (2017).

30. McHenry, J. A., Rubinow, D. R. & Stuber, G. D. Maternally responsive neurons in the bed nucleus of the
stria terminalis and medial preoptic area: Putative circuits for regulating anxiety and reward. *Front.*
Neuroendocrinol. **38**, 65–72 (2015).

31. Olazábal, D. E. *et al.* Flexibility and adaptation of the neural substrate that supports maternal behavior in
mammals. *Neurosci. Biobehav. Rev.* **37**, 1875–1892 (2013).

32. Bridges, R. S. & Hays, L. E. Steroid-induced alterations in mRNA expression of the long form of the
prolactin receptor in the medial preoptic area of female rats: Effects of exposure to a pregnancy-like regimen of
progesterone and estradiol. *Brain Res. Mol. Brain Res.* **140**, 10–16 (2005).

33. Lonstein, J. S. & De Vries, G. J. Maternal behaviour in lactating rats stimulates c-fos in glutamate
decarboxylase-synthesizing neurons of the medial preoptic area, ventral bed nucleus of the stria terminalis, and
ventrocaudal periaqueductal gray. *Neuroscience* **100**, 557–568 (2000).

34. Miceli, M. O., Fleming, A. S. & Malsbury, C. W. Disruption of maternal behaviour in virgin and
postparturient rats following sagittal plane knife cuts in the preoptic area-hypothalamus. *Behav. Brain Res.* **9**,
337–360 (1983).

35. Lee, A., Clancy, S. & Fleming, A. S. Mother rats bar-press for pups: effects of lesions of the mpoa and limbic
sites on maternal behavior and operant responding for pup-reinforcement. *Behav. Brain Res.* **100**, 15–31 (1999).

36. El Majdoubi, M., Poulain, D. A. & Theodosis, D. T. Activity-dependent morphological synaptic plasticity in
an adult neurosecretory system: magnocellular oxytocin neurons of the hypothalamus. *Biochem. Cell Biol.*
Biochim. Biol. Cell. **78**, 317–327 (2000).

37. D'Cunha, T. M., King, S. J., Fleming, A. S. & Lévy, F. Oxytocin receptors in the nucleus accumbens shell
are involved in the consolidation of maternal memory in postpartum rats. *Horm. Behav.* **59**, 14–21 (2011).

38. Insel, T. R. & Harbaugh, C. R. Lesions of the hypothalamic paraventricular nucleus disrupt the initiation of
maternal behavior. *Physiol. Behav.* **45**, 1033–1041 (1989).

39. Numan, M. & Corodimas, K. P. The effects of paraventricular hypothalamic lesions on maternal behavior in
rats. *Physiol. Behav.* **35**, 417–425 (1985).

40. Olazábal, D. E., Kalinichev, M., Morrell, J. I. & Rosenblatt, J. S. MPOA cytotoxic lesions and maternal
behavior in the rat: effects of midpubertal lesions on maternal behavior and the role of ovarian hormones in
maturation of MPOA control of maternal behavior. *Horm. Behav.* **41**, 126–138 (2002).

41. Freeman, M. E., Kanyicska, B., Lerant, A. & Nagy, G. Prolactin: structure, function, and regulation of
secretion. *Physiol. Rev.* **80**, 1523–1631 (2000).

42. Cservenák, M. *et al.* Thalamic neuropeptide mediating the effects of nursing on lactation and maternal
motivation. *Psychoneuroendocrinology* **38**, 3070–3084 (2013).

43. Kinsley, C. H. *et al.* Motherhood and the hormones of pregnancy modify concentrations of hippocampal
neuronal dendritic spines. *Horm. Behav.* **49**, 131–142 (2006).

44. Pawluski, J. L. & Galea, L. A. M. Hippocampal morphology is differentially affected by reproductive
experience in the mother. *J. Neurobiol.* **66**, 71–81 (2006).

45. Pawluski, J. L., Walker, S. K. & Galea, L. A. M. Reproductive experience differentially affects spatial
reference and working memory performance in the mother. *Horm. Behav.* **49**, 143–149 (2006).

46. Eid, R. S. *et al.* Early and late effects of maternal experience on hippocampal neurogenesis, microglia, and
the circulating cytokine milieu. *Neurobiol. Aging* **78**, 1–17 (2019).

47. Duarte-Guterman, P., Leuner, B. & Galea, L. A. M. The long and short term effects of motherhood on the
brain. *Front. Neuroendocrinol.* **53**, 100740 (2019).

48. Gogolla, N. The insular cortex. *Curr. Biol.* **27**, R580–R586 (2017).

49. Mak, G. K. & Weiss, S. Paternal recognition of adult offspring mediated by newly generated CNS neurons.
Nat. Neurosci. **13**, 753–758 (2010).

50. Larsen, C. M. & Grattan, D. R. Prolactin-induced mitogenesis in the subventricular zone of the maternal
brain during early pregnancy is essential for normal postpartum behavioral responses in the mother.
Endocrinology **151**, 3805–3814 (2010).

51. Sakamoto, M. *et al.* Continuous neurogenesis in the adult forebrain is required for innate olfactory responses.
Proc. Natl. Acad. Sci. U. S. A. **108**, 8479–8484 (2011).

52. Kelly, J. R. *et al.* Breaking down the barriers: the gut microbiome, intestinal permeability and stress-related
psychiatric disorders. *Front. Cell. Neurosci.* **9**, 392 (2015).

53. Arentsen, T. *et al.* The bacterial peptidoglycan-sensing molecule Pglyrp2 modulates brain development and
behavior. *Mol. Psychiatry* **22**, 257–266 (2017).

54. Barrière, D. A. *et al.* Structural and functional alterations in the retrosplenial cortex following neuropathic
pain. *Pain* (2019) doi:10.1097/j.pain.0000000000001610.

55. DiIorio, G., Brown, J. J., Borrello, J. A., Perman, W. H. & Shu, H. H. Large angle spin-echo imaging. *Magn.*
Reson. Imaging **13**, 39–44 (1995).

56. Ma, Y. *et al.* In vivo 3D digital atlas database of the adult C57BL/6J mouse brain by magnetic resonance
microscopy. *Front. Neuroanat.* **2**, (2008).

57. Keller, M., Pawluski, J. L., Brock, O., Douhard, Q. & Bakker, J. The α-fetoprotein knock-out mouse model
suggests that parental behavior is sexually differentiated under the influence of prenatal estradiol. *Horm. Behav.*
57, 434–440 (2010).

58. van den Brom, C. E. *et al.* Altered myocardial substrate metabolism is associated with myocardial
dysfunction in early diabetic cardiomyopathy in rats: studies using positron emission tomography. *Cardiovasc.*
Diabetol. **8**, 39 (2009).

59. Magalhães, R. *et al.* The dynamics of stress: a longitudinal MRI study of rat brain structure and connectome.
Mol. Psychiatry (2017) doi:10.1038/mp.2017.244.

60. Ashburner, J. & Friston, K. J. Voxel-based morphometry--the methods. *NeuroImage* **11**, 805–821 (2000).

61. Ashburner, J. A fast diffeomorphic image registration algorithm. *NeuroImage* **38**, 95–113 (2007).

62. Asami, T. *et al.* Longitudinal loss of gray matter volume in patients with first-episode schizophrenia:
DARTEL automated analysis and ROI validation. *NeuroImage* **59**, 986–996 (2012).

63. Barrière, D. A. *et al.* In vivo magnetic resonance imaging reveals the effect of gonadal hormones on
morphological and functional brain sexual dimorphisms in adult sheep. *Psychoneuroendocrinology* **109**, 104387
(2019).

64. Xia, M., Wang, J. & He, Y. BrainNet Viewer: a network visualization tool for human brain connectomics.
PloS One **8**, e68910 (2013).

Tables

Ex vivo / In vivo	Strain	Sex	Age (days)	Number of animals	Magnetic field intensity	Anatomical contrast	Spatial resolution	MRI Template	Atlas	Regions of interest	Tissue Probability maps	References
Ex vivo	C57BL/6J	Male	63	6	9.4 Tesla	T2 PD DW	90 x 90 x 90 µm³	Yes	Yes	21	No	Ali et al, 2005
In vivo	C57Bl/6J	Male	100	6	11.7 Tesla	T2	60 x 60 x 60 µm³	Yes	Yes	NA	No	MacKenzie-Graham et al, 2004
Ex vivo	129S1/SvImJ	Male	56	9	7 Tesla	T2	60 x 60 x 60 µm³	Yes	Yes	9	No	Kovačević et al, 2005
Ex vivo	C57BL/6J	NA	P0	8	11.7 Tesla	T2	40 x 40 x 40 µm³	Yes	Yes	12	Yes	Lee et al, 2005
In vivo	C57Bl/6J	Male	84-98	12	9.4 Tesla	T2	100 x 100 x 100 µm³	Yes	Yes	20	No	Ma et al, 2008
In vivo	C3H/HeSnJ	NA	77	15	7 Tesla	T1	156 x 156 x 156 µm³	Yes	Yes	6	No	Bock et al, 2006
Ex vivo	129S1/SvImJ C57/Bl6 CD1	Male	126	27	7 Tesla	T2	60 x 60 x 60 µm³	Yes	Yes	42	No	Chen et al, 2008
Ex vivo	C57Bl/6J	NA	63	6	9.4 Tesla	T1 T2	21.5 x 21.5 x 21.5 µm³ 43 x 43 x 43 µm³	Yes	Yes	33	No	Badea et al, 2008
Ex vivo	C57Bl/6J	20 males 20 females	84	40	7 Tesla	T2	32 x 32 x 32 µm³	Yes	Yes	62	No	Dorr et al, 2008
Ex vivo	C57Bl/6J and BXD	Male	63	12	9.4 Tesla	T1 T2	21.5 x 21.5 x 21.5 µm³ 43 x 43 x 43 µm³	Yes	Yes	20	No	Sharief et al, 2008
Ex vivo	C57BL/6J	Male	66-78	14	9.4 Tesla	T1 T2 T2*	21.5 x 21.5 x 21.5 µm³	Yes	Yes	37	No	Johnson et al, 2010
Ex vivo	C57Bl/6J	Male	84	18	16.4 Tesla	T1/T2*	30 x 30 x 30 µm³	Yes	Yes (Partial: hippocampus)	40	No	Richards et al, 2011
Ex vivo	C57Bl/6J	Male	84	18	16.4 Tesla	T1/T2*	30 x 30 x 30 µm³	Yes	Yes (Partial: cerebellum)	38	No	Ullman et al, 2013
Ex vivo	C57Bl/6J	Male	84	18	16.4 Tesla	T1/T2*	30 x 30 x 30 µm³	Yes	Yes (Partial: basal ganglia)	35	No	Ullman et al, 2013
Ex vivo	C57Bl/6J	Male	84	18	16.4 Tesla	T1/T2*	30 x 30 x 30 µm³	Yes	Yes (Partial: neocortex)	74	No	Ullman et al, 2013
In vivo	C57Bl/6J	Male	126	82	4.7 Tesla	T2	70 x 70 x 70 µm³	No	NA	NA	Yes	Sawiak et al, 2013
Ex vivo	C57Bl/6J	Male	84	18	16.4 Tesla	T1/T2*	30 x 30 x 30 µm³	Yes	Yes (Partial: diencephalon)	89	No	Watson et al, 2017
Ex vivo	C57Bl/6J	1,051 males 621 female	77	1675	NA	NA	10 x 10 x 10 µm³	No	Yes	1327	No	Wang et al, 2020

 Table 1. Comparison of mouse brain resources currently available in the literature (NA = not

 available).

Figures

Figure 1. Processing of mouse brain templates and building an atlas from the AMBMC

template and Allen Brain Atlas for data analysis and visualization. To create our resources,

we used both AMBMC mouse brain template and the mouse Allen Brain Atlas and its

associated Nissl images. (**1**) We down-sampled to a suitable resolution for MRI analysis (60-

µm isotropic resolution) and (**2**) realign in the AC/PC axis. The resulting template was then

segmented into GM, WM and CSF probability maps (**3**). These probability maps were used to

segment all the images which have been previously normalized to the template (**5,6,7**). We

obtained a large set of 138 images for each tissue class which have been used to build a

population-specific GM, WM and CSF priors. using an exponentiated lie algebra (DARTEL)

approach (**8**). This new set of population-specific tissue priors was used to segment again

normalized T_2 images (**9**) for the final VBM preprocessing (**10**, **11**, **12**). To normalize the

Allen Brain Atlas, we manually realign (**13**) and normalized the associated Nissl-stain mouse

brain using the GM priors generated previously (**14**, **15**). Both linear and nonlinear

transformations have been applied to the Allen mouse brain atlas. Then, a visual inspection of

each normalized label was carried out and, when necessary, redrawn according to Paxinos and

Franklin's atlas. Finally, the olfactory bulb and hind brain regions were completed, the *corpus*

callosum and ventricles were drawn from the WM and CSF priors, and the cerebellum labels

were replaced by the AMBMC cerebellum labels.

AMBMC = Australian Mouse Brain Mapping Consortium mouse brain template, AC/PC =

anterior commissure/posterior commissure, CSF = cerebrospinal fluid, GM = gray matter,

WM = white matter.

Figure 2. Details of the mouse brain template and atlas. (**A**) Coronal slices of the anatomical

template of the mouse brain and the corresponding gray matter, white matter and

cerebrospinal fluid probabilistic maps and the associated anatomical atlas (60-µm isotropic

resolution). (**B**) Dorsal (left panel) and lateral (right panel) 3D representations of the

anatomical mouse brain atlas. (**C**) Lateral views of the cortical areas after normalization of the

Allen Mouse Brain Atlas to the AMBMC anatomical template. The cortex was segmented

into cortical areas such as the cingulate, visual, auditory, frontal, visceral, insular, olfactory,

retrosplenial, motor, somatosensory, temporal, entorhinal and gustatory systems. Each area

was subdivided into secondary areas (*e.g.,* primary and secondary motor cortices) or structural

areas (*i.e.,* agranular, dysgranular, agranular/dysgranular, granular and posterior agranular

insular cortices). (**D**) The four images depict the different cortical layers (I, II/III, IV/V and

VIa/VIb). (**E** and **F**) 3D rendering and axial sections of subcortical structures (hypothalamus

and hippocampus).

Legend for labeled regions: <u>Hypothalamus:</u> *ARh = arcuate hypothalamic nucleus; LHA =*

lateral hypothalamic area; LPOA = lateral preoptic area; MPOA = medial preoptic area;

MPOn = medial preoptic nucleus; PH = posterior hypothalamic nucleus. AHN = anterior

hypothalamic nucleus.

<u>Hippocampus:</u> *Apr = area prostriata; CA1, CA2, CA3 = cornu ammonis areas 1, 2 and 3;*

DG = dentate gyrus; FC = fasciola cinerea; PARA = parasubiculum; POST =

postsubiculum; PRO = prosubiculum; PRE = presubiculum. For further details, see

<u>https://www.nitrc.org/projects/tmbta_2019</u>.

Figure 3. Longitudinal effects of the reproductive cycle on brain morphometry. Coronal

slices (at left) and brain plots (at right) showing gray matter concentration (GMC) differences

between control and parous animals at the end of gestation (**A**) and during early lactation (**B**)

and late lactation (**C**).

SPM flexible factorial analysis revealed an interaction between control mice and parous mice

in the late gestation period (**A**), early lactation period (**B**) and late lactation period (**C**); voxel-

712   level threshold $p < 0.01$, $t_{(126)} = 2.356$, cluster threshold = 25 voxels. *BNST = bed nucleus of*

713   *the stria terminalis; Hyp = hypothalamus; mPOA = medial preoptic area; dHippo = dorsal*

714   *hippocampus; ObF/PrL = orbitofrontal/prelimbic area; aOb/Ob = accessory olfactory*

715   *nucleus/olfactory bulb; Pir = piriform cortex; Amyg/Ent = amygdala/entorhinal cortex.*

<u>Figure 4.</u> Longitudinal time course analysis of gray matter concentration (GMC) over the
reproductive cycle. Time course comparisons in GMC between the control (black dots and
lines) and parous (red dots and lines) groups showing 3 different time profiles. (**A**) GMC
values within the medial preoptic area, the bed nucleus of *stria terminalis* (BNST) and the
agranular insular cortex reveal a significant increase in GMC during the late gestation (LG)
period maintained until weaning (W). (**B**) Specific and transient increases in GMCs are
observed in the amygdala, the piriform area and the hippocampus during early lactation (EL).
(**C**) The arcuate nucleus, PVN and caudate putamen display an increase in GMC through both
EL and late lactation (LL) periods.

Orange and blue areas represent the gestation and lactation periods, respectively. Data are
expressed as the mean ± standard error of the mean (SEM); two-way ANOVA followed by
Holm-Sidak multiple comparisons test; *$p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$, compared
with control mice.

<u>Figure 5.</u> Distribution of animals according to the quality of their maternal behavior assessed

with the pup retrieval test and brain morphometric. K-means clustering of parous animals to

classify mice into low and high maternal behavior groups based on behavior during the pup

retrieval test (**A**). Comparisons between the low and high maternal behavior groups revealed

significant differences in crouching, pup-licking and digging times (**B**). Brain slices (left

panel) and brain plots (right panel) comparing gray matter concentration (GMC)

modifications and surface maps of GMC differences between females exhibiting low and

those exhibiting high maternal behavior at the end of the gestation period (**C**) and early

lactation period (**D**).

Low and high maternal behavioral data were compared using a Student's t-test with post hoc

corrections for multiple comparisons using an FDR approach (Q = 1%) and are expressed as

the mean ± SEM; ** $p < 0.01$ and *** $p < 0.001$. SPM flexible factorial analysis revealed an

interaction between low and high maternal behavior parous mice in the late gestation period

(**A**) and early lactation period (**B**); voxel-level threshold $p < 0.01$, $t_{(60)} = 2.39$, cluster

threshold = 25 voxels. *RS = retrosplenial cortex; vHippo = ventral hippocampus; S1 =*

primary somatosensory cortex; aOB = accessory olfactory bulb; OB = olfactory bulb; Pir =

piriform cortex; Enth = entorhinal cortex; Str= striatum; OrbF = orbitofrontal cortex.

Figure 6. Longitudinal analysis in gray matter concentration (GMC) during the reproductive

cycle in control females and females exhibiting low or high maternal behavior. Time-course

comparison of GMC between the control (black dots and lines), low maternal behavior (pink

dots and lines) and high maternal behavior (red dots and lines) groups revealed two types of

time profile. GMC analysis in the entorhinal area, orbitofrontal area, the accessory olfactory

bulb and the medial preoptic nucleus revealed an acute and specific increase in GMC values

in the high maternal behavior group at the late gestation period (**A**). In contrast, GMC analysis

in the ventral hippocampus, the retrosplenial area and the primary somatosensory area

revealed an increase in GMC in the high maternal behavior group at the late gestation period,

and this increase was maintained until weaning (**B**).

Orange and blue areas represent the gestation and lactation periods, respectively. Data were

compared using a two-way ANOVA followed by Holm-Sidak multiple comparisons test and

expressed as the mean ± SEM; * $p < 0.05$, ** $p < 0.01$ when high maternal mice were

compared with control mice and [$$] p < 0.01, [$$$] p < 0.001 when high maternal mice were

compared with low maternal mice.

<u>Figure 7.</u> Estimation of the sensitivity and specificity of late-gestation GMC measures in the
entorhinal area (**A**) and accessory olfactory bulb (**B**) to predict postpartum maternal
performance.

Receiver operating characteristic (ROC) curves were estimated using a Wilson/Brown test
with a 95% confidence interval. Correlations were estimated using a Pearson correlation test.
ROC and correlation analyses were considered significant at $p < 0.05$.

771   **Supplementary Data**

# Label	Region of Interest	Hemisphere	T Value	p Value	x (mm)	y (mm)	z (mm)
712	Visceral area, layer 2/3	Left	2.3789	0.0094	-41.5	-3.5	-3.5
1842	Agranular insular area, dorsal part, layer 2/3	Left	3.9264	< 0.0001	-34.5	16.5	-8.5
652	Gustatory areas, layer 2/3	Left	3.4262	0.0004	-30.5	25.5	-4.5
2562	Cortical amygdalar area, posterior part, medial zone	Left	2.6591	0.0044	-23.5	-16.5	-18.5
3002	Olfactory tubercle	Left	4.2298	< 0.0001	-22.5	13.5	-23.5
2502	Piriform area	Left	3.6483	0.0002	-21.5	26.5	-14.5
3122	Medial amygdalar nucleus	Left	2.5052	0.0067	-21.5	-10.5	-11.5
2972	Caudoputamen	Left	2.5003	0.0068	-20.5	7.5	-6.5
3522	Lateral dorsal nucleus of thalamus	Left	2.3933	0.0090	-17.5	-4.5	4.5
292	Primary somatosensory area, lower limb, layer 2/3	Left	2.4134	0.0086	16.5	4.5	23.5
1692	Orbital area, lateral part, layer 2/3	Left	2.4677	0.0074	-15.5	37.5	-0.5
2462	Anterior olfactory nucleus	Left	2.3608	0.0098	-11.5	33.5	-5.5
4132	Lateral hypothalamic area	Left	2.5217	0.0064	-10.5	-10.5	-18.5
2982	Nucleus accumbens	Left	2.3953	0.0090	-9.5	18.5	-15.5
3192	Diagonal band nucleus	Left	2.5835	0.0054	-5.5	14.5	-22.5
1731	Orbital area, medial part, layer 1	Right	2.5912	0.0053	1.5	37.5	0.5
2421	Main olfactory bulb	Right	3.7602	0.0001	2.5	35.5	-17.5
3861	Medial preoptic area	Right	2.6983	0.0040	6.5	11.5	-21.5
3971	Anterior hypothalamic nucleus	Right	2.4173	0.0085	8.5	-0.5	-20.5
4201	Tuberal nucleus	Right	2.4250	0.0083	8.5	-3.5	-20.5
4131	Lateral hypothalamic area	Right	3.3026	0.0006	11.5	-10.5	-16.5
11	Frontal pole, layer 1	Right	2.5614	0.0058	12.5	41.5	5.5
4141	Lateral preoptic area	Right	2.5685	0.0057	12.5	5.5	-18.5
2461	Anterior olfactory nucleus	Right	2.4144	0.0086	14.5	35.5	7.5
2631	Dentate gyrus, molecular layer	Right	2.4871	0.0071	14.5	-16.5	12.5
3121	Medial amygdalar nucleus	Right	2.6129	0.0050	21.5	-10.5	-11.5
3001	Olfactory tubercle	Right	2.3745	0.0095	22.5	10.5	-24.5
1841	Agranular insular area, dorsal part, layer 2/3	Right	3.0066	0.0016	23.5	32.5	-2.5
2501	Piriform area	Right	2.9740	0.0018	26.5	24.5	-12.5
2971	Caudoputamen	Right	2.7880	0.0031	27.5	-7.5	-5.5
621	Supplemental somatosensory area, layer 6a	Right	2.4045	0.0088	29.5	8.5	2.5
591	Supplemental somatosensory area, layer 2/3	Right	2.5642	0.0057	36.5	16.5	-0.5

773   **Table S1.** Local variation in gray matter concentration between control and parous mice at the

774   end of the gestation period. SPM flexible factorial analysis revealed an interaction between

775   the control and parous groups at the late gestation time point.

# Label	Region of Interest	Hemisphere	T Value	p Value	x {mm}	y {mm}	z {mm}
712	Visceral area, layer 2/3	Left	4.0474	< 0.0001	41.5	3.5	3.5
592	Supplemental somatosensory area, layer 2/3	Left	4.2243	< 0.0001	-40.5	6.5	-2.5
1842	Agranular insular area, dorsal part, layer 2/3	Left	3.9710	< 0.0001	-34.5	16.5	-8.5
352	Primary somatosensory area, mouth, layer 2/3	Left	1.9021	0.0293	-31.5	22.5	5.5
2972	Caudoputamen	Left	1.8455	0.0331	-31.5	-8.5	-0.5
342	Primary somatosensory area, mouth, layer 1	Left	1.7124	0.0437	-30.5	23.5	6.5
72	Primary motor area, layer 2/3	Left	1.7423	0.0411	-29.5	27.5	-0.5
232	Primary somatosensory area, barrel field, layer 2/3	Left	2.0289	0.0220	-27.5	6.5	18.5
532	Primary somatosensory area, unassigned, layer 2/3	Left	1.7753	0.0384	-27.5	11.5	16.5
2812	Subiculum	Left	3.0917	0.0012	-25.5	-24.5	10.5
2642	Dentate gyrus, polymorph layer	Left	1.7318	0.0420	-20.5	-23.5	-9.5
4092	Ventral premammillary nucleus	Left	2.3287	0.0107	-8.5	-14.5	-18.5
5522	Medial vestibular nucleus	Left	1.9545	0.0261	-5.5	-47.5	1.5
4442	Superior colliculus, motor related, intermediate gray layer	Left	2.9857	0.0017	2.5	14.5	13.5
2411	Olfactory areas	Right	2.0350	0.0217	0.5	23.5	-11.5
4031	Medial preoptic area	Right	1.6858	0.0461	5.5	-10.5	-19.5
2421	Main olfactory bulb	Right	2.4342	0.0081	2.5	34.5	-17.5
4541	Olivary pretectal nucleus	Right	2.6632	0.0044	6.5	-12.5	11.5
3211	Bed nuclei of the stria terminalis	Right	1.7760	0.0383	10.5	8.5	-6.5
4131	Lateral hypothalamic area	Right	2.0659	0.0202	11.5	-9.5	-17.5
4211	Zona incerta	Right	1.9015	0.0293	11.5	8.5	11.5
2791	Postsubiculum	Right	3.0783	0.0013	15.5	-18.5	12.5
3121	Medial amygdalar nucleus	Right	1.8183	0.0351	16.5	-10.5	-16.5
2561	Cortical amygdalar area, posterior part, medial zone	Right	3.5177	0.0003	22.5	-15.5	-19.5
401	Primary somatosensory area, upper limb, layer 1	Right	1.7139	0.0435	24.5	9.5	19.5
531	Primary somatosensory area, unassigned, layer 2/3	Right	1.7406	0.0412	26.5	9.5	18.5
61	Primary motor area, layer 1	Right	1.7743	0.0385	27.5	27.5	7.5
221	Primary somatosensory area, barrel field, layer 1	Right	2.2903	0.0118	27.5	5.5	19.5
2731	Entorhinal area, medial part, dorsal zone, layer 1	Right	1.9890	0.0241	27.5	-32.5	-1.5
231	Primary somatosensory area, barrel field, layer 2/3	Right	1.8671	0.0316	29.5	7.5	17.5
2571	Piriform-amygdalar area	Right	3.3855	0.0005	29.5	-6.5	-24.5
351	Primary somatosensory area, mouth, layer 2/3	Right	1.9950	0.0238	32.5	14.5	11.5
171	Primary somatosensory area, nose, layer 2/3	Right	1.8466	0.0330	33.5	12.5	11.5
2501	Piriform area	Right	3.4453	0.0004	36.5	15.5	17.5

Table S2. Local variation in gray matter concentration between control and parous mice at the beginning of the lactation period. SPM flexible factorial analysis revealed an interaction between the control and parous groups at the early lactation time point.

# Label	Region of Interest	Hemisphere	T Value	p Value	x {mm}	y {mm}	z {mm}
712	Visceral area, layer 2/3	Left	2.1741	0.01566	41.5	0.5	2.5
2812	Subiculum	Left	2.0845	0.01938	-28.5	-24.5	9.5
2972	Caudoputamen	Left	2.8735	0.00238	-28.5	-7.5	-3.5
2502	Piriform area	Left	2.3017	0.01143	-27.5	16.5	-16.5
2562	Cortical amygdalar area, posterior part, medial zone	Left	2.2071	0.01445	-23.5	-17.5	-17.5
2992	Fundus of striatum	Left	2.0065	0.0232	-20.5	8.5	-17.5
4442	Superior colliculus, motor related, intermediate gray layer	Left	1.9022	0.02927	-16.5	-21.5	30.5
2142	Anterior visual area, layer 2/3	Left	1.7105	0.04382	-13.5	-0.5	25.5
4082	Dorsal premammillary nucleus	Left	2.6229	0.00488	-5.5	-14.5	-15.5
3011	Lateral septal nucleus, caudal (caudodorsal) part	Right	2.2155	0.01416	4.5	8.5	7.5
3861	Medial preoptic area	Right	2.8305	0.0027	6.5	11.5	-21.5
3211	Bed nuclei of the stria terminalis	Right	1.7490	0.04052	7.5	14.5	12.5
4131	Lateral hypothalamic area	Right	1.7901	0.03721	11.5	-6.5	-17.5
2621	Field CA3	Right	2.6408	0.00465	21.5	-18.5	-6.5
5161	Dorsal cochlear nucleus	Right	1.7477	0.04063	23.5	-43.5	-4.5
2561	Cortical amygdalar area, posterior part, medial zone	Right	1.8753	0.03104	24.5	-18.5	-17.5
2731	Entorhinal area, medial part, dorsal zone, layer 1	Right	1.9285	0.02763	27.5	-32.5	-1.5
2971	Caudoputamen	Right	2.8706	0.0024	30.5	8.5	1.5
701	Visceral area, layer 1	Right	1.8073	0.03589	41.5	3.5	-3.5

781 **Table S3.** Local variation in gray matter concentration between control and parous mice at the

782 end of the lactation period. SPM flexible factorial analysis revealed an interaction between the

783 control and parous groups at the late lactation time point.

# Label	Region of Interest	Hemisphere	T Value	p Value	x {mm}	y {mm}	z {mm}
992	Ventral auditory area, layer 6b	Left	2.4578	0.0084	-36.5	-15.5	5.5
2292	Temporal association areas, layer 6a	Left	2.7134	0.0043	-36.5	-24.5	8.5
242	Primary somatosensory area, barrel field, layer 4	Left	2.4559	0.0084	-33.5	0.5	12.5
2752	Entorhinal area, medial part, dorsal zone, layer 3	Left	3.7707	0.0002	-30.5	-29.5	-9.5
2812	Subiculum	Left	2.6370	0.0053	-29.5	-23.5	10.5
2212	Rostrolateral visual area, layer 4	Left	2.7190	0.0043	-23.5	-7.5	22.5
2622	Field CA3	Left	4.2394	0.0000	21.5	18.5	6.5
4362	Substantia nigra, reticular part	Left	2.4952	0.0076	-17.5	-23.5	-5.5
72	Primary motor area, layer 2/3	Left	2.5921	0.0060	-12.5	13.5	20.5
2162	Retrosplenial area, dorsal part, layer 6a	Left	3.1820	0.0012	-11.5	-0.5	23.5
4132	Lateral hypothalamic area	Left	3.0784	0.0016	-11.5	-6.5	-17.5
2462	Anterior olfactory nucleus	Left	4.0594	0.0001	10.5	31.5	12.5
4212	Zona incerta	Left	3.0013	0.0020	-9.5	-10.5	-10.5
3862	Medial preoptic area	Left	2.6620	0.0050	-8.5	4.5	-20.5
4412	Superior colliculus, motor related, deep gray layer	Left	2.8740	0.0028	-8.5	-23.5	16.5
4012	Medial mammillary nucleus, medial part	Left	2.4376	0.0088	-4.5	-16.5	-16.5
4112	Ventromedial hypothalamic nucleus	Left	2.5173	0.0072	-3.5	-6.5	-22.5
4272	Superior colliculus, zonal layer	Left	2.8049	0.0034	2.5	14.5	17.5
4052	Tuberomammillary nucleus, dorsal part	Left	2.4993	0.0076	-1.5	-15.5	-19.5
4441	Superior colliculus, motor related, intermediate gray layer	Right	2.5749	0.0062	0.5	-26.5	19.5
4451	Periaqueductal gray	Right	2.4815	0.0079	3.5	-32.5	13.5
2651	Dentate gyrus, granule cell layer	Right	2.7718	0.0037	7.5	7.5	10.5
4111	Ventromedial hypothalamic nucleus	Right	2.6915	0.0046	7.5	-6.5	-21.5
4431	Superior colliculus, motor related, intermediate white layer	Right	2.4866	0.0078	9.5	-28.5	18.5
2421	Main olfactory bulb	Right	3.9879	0.0001	10.5	38.5	-8.5
1071	Anteromedial visual area, layer 2/3	Right	4.0275	0.0001	12.5	-4.5	25.5
2971	Caudoputamen	Right	2.4763	0.0080	16.5	10.5	-7.5
2411	Olfactory areas	Right	3.9513	0.0001	17.5	30.5	-7.5
2621	Field CA3	Right	3.7785	0.0002	25.5	-17.5	-8.5
2811	Subiculum	Right	2.4734	0.0081	25.5	23.5	11.5
1241	Posterolateral visual area, layer 1	Right	2.5546	0.0066	27.5	-31.5	19.5
2741	Entorhinal area, medial part, dorsal zone, layer 2	Right	4.0416	0.0001	29.5	-31.5	-8.5

Table S4. Local variation in gray matter concentration between low maternal behavior parous mice and high maternal behavior parous mice at the end of the gestation period. SPM flexible factorial analysis revealed an interaction between the low *versus* high maternal behavior groups at the late gestation time point.

# Label	Region of Interest	Hemisphere	T Value	p Value	x (mm)	y (mm)	z (mm)
2732	Entorhinal area, medial part, dorsal zone, layer 1	Left	2.5882	0.0060	28.5	33.5	1.5
2632	Dentate gyrus, molecular layer	Left	2.5173	0.0072	-20.5	-16.5	6.5
412	Primary somatosensory area, upper limb, layer 2/3	Left	3.1161	0.0014	-19.5	14.5	18.5
2602	Field CA1	Left	2.5276	0.0070	-19.5	-11.5	17.5
122	Secondary motor area, layer 2/3	Left	2.8890	0.0027	-15.5	24.5	15.5
2422	Main olfactory bulb	Left	3.5084	0.0004	-8.5	45.5	6.5
3652	Parafascicular nucleus	Left	3.0525	0.0017	-3.5	-12.5	-1.5
2672	Induseum griseum	Left	2.4662	0.0082	-1.5	19.5	5.5
1482	Anterior cingulate area, dorsal part, layer 1	Left	2.4675	0.0082	-0.5	21.5	14.5
2082	Retrosplenial area, ventral part, layer 1	Left	2.3913	0.0099	-0.5	-3.5	23.5
2421	Main olfactory bulb	Right	2.7201	0.0043	2.5	43.5	-14.5
3651	Parafascicular nucleus	Right	2.9370	0.0023	6.5	12.5	0.5
2411	Olfactory areas	Right	3.2967	0.0008	8.5	42.5	4.5
2601	Field CA1	Right	3.6131	0.0003	18.5	-9.5	17.5
381	Primary somatosensory area, mouth, layer 6a	Right	3.1451	0.0013	24.5	14.5	5.5
2821	Prosubiculum	Right	2.9853	0.0020	24.5	-19.5	14.5
2691	Entorhinal area, lateral part, layer 2	Right	3.0830	0.0015	38.5	29.5	5.5

790 **Table S5.** Local variation in gray matter concentration between low maternal behavior parous

791 mice and high maternal behavior parous mice at the beginning of the lactation period. SPM

792 flexible factorial analysis revealed an interaction between the low *versus* high maternal

793 behavior groups at the early lactation time point.