Dopamine-gated memory selection during slow wave sleep

**One Sentence Summary:**

Dopamine before sleep promotes forgetting of weak memory traces associated with increased

spindle amplitude around the peak of a slow oscillations.

**Authors:**

Hanna Isotalus 1, 2 *

-- hanna.isotalus@bristol.ac.uk

Will J Carr 1

-- wc17397@bristol.ac.uk

George G Averill 1

-- george.g.averill@gmail.com

Oliver Radtke 6

-- Oliver.Radtke@hhu.de

James Selwood 1, 3

-- james.selwood@bristol.ac.uk

Rachel Williams 1

-- Rachel.Williams@nbt.nhs.uk

Elizabeth Ford 1

-- b-ford@outlook.com

Liz McCullagh 7

-- liz.McCullagh@UHBW.nhs.uk

James McErlane 1

-- james.mcerlane1@gmail.com

Cian O'Donnell 8

-- cian.odonnell@bristol.ac.uk

Claire Durant 4

-- Claire.Durant@bristol.ac.uk

Ullrich Bartsch 5

-- Ullrich.Bartsch@bristol.ac.uk

Matt W Jones 5

-- Matt.Jones@bristol.ac.uk

Carlos Muñoz-Neira 1

-- carlos.munoz@bristol.ac.uk

Alfie R Wearn 1

-- alfie.wearn@bristol.ac.uk

John P Grogan 1,9

-- john.grogan@bristol.ac.uk

Elizabeth J Coulthard 1, 3 *

-- Elizabeth.Coulthard@bristol.ac.uk

*Corresponding authors

**Affiliations:**

1 Clinical Neurosciences, Translational Health Sciences, Bristol Medical School, University of

Bristol, Bristol

2 Digital Health, Faculty of Engineering, University of Bristol, Bristol

3 Southmead Hospital, North Bristol NHS Trust, Bristol

4 Experimental Psychology, University of Bristol, Bristol

5 School of Physiology, Pharmacology and Neuroscience, University of Bristol

6 Department of Neurology, Centre for Movement Disorders and Neuromodulation, Medical

Faculty, Heinrich-Heine-University, Düsseldorf.

7 Production Pharmacy, Bristol Royal Infirmary, University Hospitals Bristol and Weston NHS

Trust

8 School of Computer Science, Electrical and Electronic Engineering, and Engineering

Mathematics, University of Bristol, Bristol

9 Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford

**Abstract:**

The human brain selectively stores knowledge of the world to optimise future behaviour,

automatically rehearsing, contextualising or discarding information to create a robust record of

experiences. Storage or forgetting evolves over time, particularly during sleep. We have

previously shown that dopamine given in the form of L-DOPA tablets improves long-term

memory in Parkinson's disease, but only when given overnight. L-DOPA is already prescribed

widely with a good safety profile and could potentially be rapidly repurposed to improve

cognitive performance and improve quality of life in, for example, early Alzheimer's Disease, if

we understood the best time of day to prescribe. Therefore, we sought to test how dopamine

shaped long-term memory formation before and during sleep in a double-blind randomised

placebo-controlled cross-over trial of healthy older adults (n = 35). We administered L-DOPA

after word-list learning to be active during repeat exposure to a proportion of the words and

during subsequent nocturnal sleep. Nocturnal dopamine accelerated forgetting for words

presented once but it did not affect memory for words presented twice. During slow wave sleep,

L-DOPA also increased spindle amplitude around slow oscillation peaks. Larger dopamine-

induced difference in word memory was associated with a larger increase in spindle amplitude.

Dopamine-dependent memory processing may therefore modulate spindles dependent on slow-

oscillation phase. Further, overnight dopamine increased total slow wave sleep duration by

approximately 11%. This pharmaceutical modification of slow wave sleep may have potential

health-enhancing benefits in old age that could include cognitive enhancement and Alzheimer's

prevention.

## Introduction

The brain selectively extracts and stores important details of our daily lives, while demoting irrelevant information - you have probably forgotten where you parked your car while shopping last week, but you will remember your parking slot in an airport carpark after a week's holiday. When memories are first encoded they form traces known as engrams – or changes in neuronal and synaptic activity that represent a memory (1, 2). Depending on context and relevance, engrams can be integrated within memory networks for the long term or forgotten through a set of processes that start immediately and progress during wake and sleep (3-5).

The complex milieu that underpins memory formation depends on several neurotransmitters including dopamine. This transmitter is released from two midbrain areas - locus coeruleus and ventral tegmental area – which directly project to the hippocampus (6). Exogenous dopamine administration can modulate memory persistence, particularly after initial learning (7-10). In humans with dopamine depletion due to Parkinson's disease, memory consolidation improves with overnight administration of L-DOPA (Levodopa – which increases dopamine concentrations in the brain), but the timing of the dopamine manipulation relative to learning critically determines its effects on memory (8, 11).

During memory encoding and shortly after, engrams of important information can be prioritised for storage, based either on previous knowledge, repeated exposure, or other associations, such as financial or emotional value (12, 13). At a molecular level, synaptic, 'tagging' for information prioritised for storage, protein synthesis and synaptic modifications occur within hours of encountering information (14, 15). The dopaminergic connection between midbrain and hippocampus may selectively bias long-term memory storage by altering synaptic tagging or the protein synthesis involved in synaptic tagging (15-17).

As well as prospectively prioritising, or tagging, memories for later replay during sleep, dopamine

may directly act during sleep *per se (18)*. As engram storage evolves, newly acquired memories are

spontaneously repeated *(17)*; sleep affords an optimal neurophysiological state during which to

enact this process - although replay occurs during wake too *(19)*. Patterns of activation within

hippocampal neuronal assemblies are selectively replayed during sharp wave ripples which are, in

turn, temporally coupled to sleep spindles and slow oscillations, prominent during non-REM

sleep *(20-24)*.

Contextual information at a later time-point can also retroactively alter the likelihood that

previous memories are stored for longer term *(25-27)*. Dopaminergic modulation of memory

may underpin contextually-driven modification of engrams *(16, 28-30)*. Exogenous

administration of dopamine may therefore alter the likelihood of memory stabilisation when

given *after* initial learning, during re-exposure and consolidation.

It is important to point out that as well as actively prioritising relevant memories for storage,

several neurobiological substrates could promote forgetting *(31, 32)*. In drosophila, the

'forgetting cells' promote changes in cellular signalling that weaken the engram by releasing

dopamine. Dopamine enhances encoding of new memories at the cost of triggering forgetting of

competing information *(33, 34)*. This dopamine-induced strategic forgetting is selective to weakly

encoded memories – presumably, an automatic strategy for ensuring retention of more

behaviourally relevant information. In humans, retroactive retrieval of already learnt information

has been shown to simultaneously enhance memory for the retrieved information while inducing

forgetting of contextually related information *(27)*. This retrieval induced forgetting is caused by

an *active* inhibition of the competing memory traces during recall *(35, 36)*.

Here we propose a role for dopamine in *selecting* memories for long-term storage. Specifically, we

propose that, after initial learning, dopamine biases human memory in favour of strongly

encoded memory traces as opposed to weaker traces. We tested the hypothesis that L-DOPA

given after initial learning (to be active during re-exposure and subsequent sleep) will increase

memory retention compared to placebo. We also predicted that no such effect would be

observed for words that were not re-exposed. We predicted the primary effects of dopamine

during long-term memory evolution would be mediated through modulation of slow wave sleep

duration and sleep spindle characteristics.

In this double-blind randomised within-subjects placebo-controlled trial, we show that a single

dose of exogenous dopamine (L-DOPA) given after learning (to be active during re-exposure

and subsequent nocturnal sleep) unexpectedly accelerates forgetting of non-repeated information

effectively biasing memory selection away from weakly encoded items. Investigation of sleep

characteristics further showed that spindle amplitude during slow wave sleep was higher on L-

DOPA compared to placebo, and this effect was slow oscillation phase dependent. The

magnitude of this increase in spindle amplitude correlated with the behavioural effect of

dopamine on memory selection. Further, we show that nocturnal L-DOPA increases slow wave

sleep duration by ~11%.

We also report a secondary placebo-controlled randomised control study in which we found that

L-DOPA did not affect encoding or retrieval of learned words, further suggesting that the

effects of L-DOPA in the main experiment were enacted during repetition, consolidation and/or

sleep.

Results

To study the relationships between dopamine, sleep and forgetting, we carefully timed

administration of L-DOPA to increase dopamine concentration in healthy older adults across

two placebo-controlled double-blind randomised crossover experiments: the main experiment

and a secondary experiment. The overarching structure enabled targeting of L-DOPA to

different memory processes – in the main experiment (**Fig. 1a.**), we explored the effects of

dopamine on memory consolidation by administering long-acting L-DOPA after learning, to be

active after initial learning and during nocturnal sleep (37). In the secondary experiment, we used

short-acting L-DOPA to target memory retrieval (testing) or encoding (learning), but not sleep.

**A**

**b**

<div align="right"><em>Figures are not to scale.</em></div>

**Fig. 1. Experiment 1 Study procedure**

**a.** In this placebo-controlled randomised crossover trial, healthy elderly volunteers completed two overnight sleep polysomnography visits. In the evenings, they learnt a set of words (**Fig. 1b.**) 45min *before* receiving 200mg L-DOPA CR or placebo. 75min *after* dosing a portion of the words were re-exposed. The orange bracket denotes when L-DOPA was active. Therefore, L-DOPA was active during re-exposure and sleep but not during learning or memory tests on days 1, 3, or 5. Apart from treatment (L-DOPA or placebo) the nights were identical. Each volunteer completed both nights.

**b.** Participants were asked to memorise 80 words shown on a computer screen one at a time. The words were separated into four lists for testing (Lists *i, ii, iii, iv* – 20 words each) but during learning they were shown in a random, interleaved, order. 2h after learning,

participants were re-exposed to List *i* by a recognition test. The following morning, ~12 hours after initial learning, memory for Lists *i* and *ii* was tested (random, interleaved), while lists *iii* and *iv* were tested 3 and 5 days later over the phone. Each test was performed using a recognition test with a unique set of distractor words. The testing procedure was fully explained to participants before learning.

In the main experiment, 35 healthy elderly volunteers (age = 68.9 ± 3.5 years; 22 Female)

completed two overnight study visits (**Fig. 1a.**) which were identical except for treatment

allocation. On each visit, we administered controlled release L-DOPA (CR; co-beneldopa

200/50mg) or placebo *after* participants had learnt information (word Lists *i, ii, iii and iv,* **Fig.**

**1b.**). Participants were re-exposed to a quarter of the items (List *i*) shortly after L-DOPA (or

placebo) administration through a recognition memory test – this manipulation was performed

to strengthen the memory for each List *i* word. Memory for the re-exposed items (List *i*) was

tested the following day together with a matched number of items that had not been re-exposed

(List *ii* – weak memory). Memory for the remainder of the items was probed 3 or 5 days after

learning (Lists *iii* and i*v*). The participants knew some words would be tested both in the evening

and in the morning, and the remainder of the words would only be tested once.

Initial learning occurred before L-DOPA (/ placebo), whereas memory re-exposure and a full

night of sleep occurred after L-DOPA (/ placebo). Therefore, we were able to isolate the effects

of dopamine on re-exposure, consolidation and sleep-dependent processing from its effects on

initial encoding and retrieval. Items presented only once (Lists *ii, iii, iv*) were expected to have

induced weaker memory traces than the re-exposed items (List *i*).

We used d' (D-prime) as a measure of recognition memory accuracy for each list. D' is a

sensitivity index that takes into account both the accurately detected signal (hits) and inaccurately

identified noise (false alarms) *(38)*. In other words, d' captures not just correctly identified "old"

words during the recognition test, but it also accounts for incorrect judgements of "new" items

as "old". D' can be calculated as the difference between the Z-transformed rates of correct hit

responses and incorrect false alarms. A higher d' therefore indicates better ability at performing

the task, while 0 indicates performance at chance.

## L-DOPA accelerates forgetting during sleep

L-DOPA given after learning accelerated forgetting of items presented only once when memory was tested the next day (List $ii$) but not at greater delays (Lists $iii, iv$, **Fig. 2a.**). First, we performed pairwise comparisons between the L-DOPA and placebo conditions for each single-exposure list. Data was missing for one participant from Day 5 test following placebo, and therefore we analysed each list separately, as opposed to using an ANOVA which would require removing the participant's data from all analyses. These comparisons demonstrated that d' was reduced on L-DOPA (**d'** $_{\text{List } ii}$ = 1.249 ± 0.59) compared to placebo (**d'** $_{\text{List } ii}$ = 1.544, ± 0.65) at Day 1 (paired t(34) = -3.333, p = 0.002, $BF_{10}$ = 16.6, n = 35). By Day 3 there was no difference (**d'** $_{\text{List } iii}$: L-DOPA = 0.86 ± 0.46; placebo = 0.82 ± 0.63; Wilcoxon's Z = 338, p = 0.313, $BF_{01}$ = 5.2, n =35; **d'** $_{\text{List } iv}$: L-DOPA = 0.58 ± 0.58; placebo = 0.59 ± 0.55; t(33) = -0.02, p = 0.982, $BF_{01}$ = 5.4, n = 34). The reduction in List $ii$ accuracy remained after correcting for false discovery rate for the three tests (Benjamini-Hochberg corrected p = 0.006) (*39*). Together these findings show that L-DOPA accelerates the speed of forgetting for information over 1 night, but this information would be lost in the longer term even without L-DOPA (**Fig. 2a.**, *SM1*). This suggests that dopamine may play an important part in either selecting memories for storage or initiating forgetting.

We were also interested in any dose-dependent effects of L-DOPA on memory. Body weight is known to influence the cumulative dose and pharmacokinetic properties of L-DOPA (*40*), as well as L-DOPA's effect on memory in humans (*9*). We used a mixed linear model to investigate the effect of dose (based on body weight). A model with weight-adjusted dose (mg/kg), delay from learning (days) and the interaction term (delay * dose) as fixed effects and participants as random effects (including random intercepts and slopes of delay by participant) revealed a main effect of delay (t(33.7) = -9.142, p < 0.001), no overall effect of dose (t(20.3) = -1.36, p = 0.188)

and a delay * dose interaction (t(98.2) = 2.33, p = 0.022, n = 35). The two effects remained

following false discovery rate correction for the whole model (*SM 2*). Next, we performed a

series of post-hoc correlational analyses to determine which effects were driving this interaction.

The degree of forgetting correlated with L-DOPA dose (Spearman's $\varrho$ = -0.56, p < 0.001, n =

35, $p_{corrected}$ < 0.001 after correcting for all post hoc correlations) but not with placebo (**Fig. 2b.** –

Spearman's $\varrho$ = -0.23, p = 0.18, n = 35). The degree of forgetting did not correlate with L-

DOPA dose (p > 0.36) on days 3 or 5 in either condition. The lack of correlation in the placebo

arm suggests that these effects were not driven by bodyweight but were instead associated with

the treatment. The delay*dose interaction was therefore driven by L-DOPA affecting memory

for List *ii* on Day 1 but not at subsequent delays. This suggests that L-DOPA accelerates initial

forgetting in a dose-dependent manner, but it does not influence memory for more strongly

encoded items that would be retained 3 or 5 days later.

**a**

**b**

**c**

**Fig. 2. Nocturnal dopamine dose-dependently modulates memory**

**a.** Higher d' at Day 1 on placebo (green) compared with L-DOPA (red) shows that overnight L-DOPA increased forgetting when memory was tested next day (List *ii*) but not 3 or 5 days later (Lists *iii* and *iv* respectively) compared to placebo. Note that L-DOPA was no longer active during memory tests. Therefore, L-DOPA during sleep accelerates forgetting of weakly encoded information that is naturally forgotten by day 3. Boxplot shows quartiles with kernel densities plotted to the right.

**b.** Higher L-DOPA dose during consolidation was correlated with poorer Day 1 recall of List *ii* d' (Spearman's $\varrho$ = -.056, p < 0.001, red) but no such relationship was found on the placebo night ($\varrho$ = -0.23, p = 0.180, green). Notably, these two relationships were also different (Pearson's r-to-z transform z= -2.634, p = 0.008) suggesting the effect is driven by dose and not body weight alone. Lines of best fit are shown for illustration purposes.

**c.** L-DOPA increased the relative benefit of re-exposed compared to other items (List *i* d' minus List *ii* d'). This relative benefit was larger after L-DOPA (d' $_{List\ i\leftarrow ii}$= 0 .953 $\pm$ = 0.67) compared to placebo (d' $_{List\ i\leftarrow ii}$= 0.643 $\pm$ 0.56; t (34) = 2.48, p = 0.018, $BF_{10}$ = 2.6). This difference was driven both by an increase in List *i* d' and a decrease in List *ii* d' on L-DOPA, although the former was not significant (*SM 3*). Lines show maximum, median, and minimum values (horizontally) and kernel densities (vertically).

### L-DOPA accelerates forgetting of weak but not strong engrams

Next, we investigated whether dopamine modulates how re-exposure affects memory. As expected, strong memory traces (re-exposed items – List *i* ) were better retained (more 'hits') than others (List *ii*) both following L-DOPA (**Hits** $_{List\ i}$ = 18.1 $\pm$ 2.1; **Hits** $_{List\ ii}$ = 13.8 $\pm$ 3.3; t(34) = 8.49, p < 0.001) and following placebo (**Hits** $_{List\ i}$ = 18.0 $\pm$ 2.4; **Hits** $_{List\ ii}$ = 15.0 $\pm$ 3.0; t(34) = 7.18, p < 0.001).

While L-DOPA accelerated baseline forgetting for weaker items as shown above (d' $_{List\ ii}$ = 1.25 $\pm$ 0.59) compared to placebo (d' $_{List\ ii}$ = 1.54 $\pm$ 0.11, t(34) = -3.333, p = 0.002, $BF_{01}$ = 0.1, *SM 3*), re-exposed List *i* items were not affected by the treatment (**d'** $_{List\ i}$ = 2.20 $\pm$ 0.78; placebo **d'** $_{List\ i}$ = 2.19 $\pm$ 0.77; t(34) = 0.134, p = 0.894, $BF_{10}$ = 5.5, **Fig. 2c.**, *SM 4, 5*). Note that we expected that L-DOPA would have enhanced retention for strong engrams while leaving weaker memories unaffected. Instead, L-DOPA accelerated forgetting of weaker information leaving stronger memories largely unaffected.

To quantify this relative effect of dopamine on single-exposed compared to re-exposed items, we used the paired difference between the strongly and weakly encoded lists (i.e. d' for List $i$ minus d' for List $ii$) from the Day 1 recognition test. This relative benefit was larger following L-DOPA ($\mathbf{d'}_{\text{List } i\text{-}ii}$ = 0.953 ± 0.67) compared to placebo ($\mathbf{d'}_{\text{List } i\text{-}ii}$ = 0.643 ± 0.56, (t(34) = 2.48, p = 0.018, $BF_{10}$ = 2.6, **Fig. 2.c.**). Therefore, L-DOPA selectively biased memory retention away from non-repeated items resulting in a larger difference between the two lists on L-DOPA compared to placebo.

To reiterate, L-DOPA differentially modulated strong and weak memory traces, augmenting differences between them. Furthermore, we performed two post-hoc analyses that showed that the treatment had no effect on the false alarm rate (t(34) = 0.527, p = 0.601, $BF_{01}$ = 4.8). Rather, L-DOPA reduced the hit rate (List $ii$ – t(34) = -2.89, p = 0.007, $BF_{10}$ = 6.0) – the hits rather than the false alarms drive all the effects of L-DOPA on d' we identified. This implies that effects of dopamine are related to engram strength rather than modulation of noise that generates false responses.

Importantly, there was no difference in performance during the evening re-exposure tests between placebo and L-DOPA conditions (Day 0 List $i$ paired t(34) = .83, p = 0.412, $BF_{01}$ = 4.0, *SM 4*). The Bayes Factor ($BF_{01}$) suggested that these results were 4 times more likely to have been recorded under the null than the alternative distribution. Therefore, dopamine did not affect memory performance before sleep – the effects we report here only manifest after or during the re-exposure.

Together, these findings provide evidence that dopamine biases selection of memories for long

term storage by accelerating forgetting of weakly-encoded information, a process that could free

memory capacity for storage of more strongly encoded items. Dopamine may bias memory in

this way during sleep. Next, we explored polysomnography measures for potential

neurophysiological mechanisms underlying dopamine's effects on memory.

L-DOPA prolongs slow wave sleep

Nocturnal L-DOPA increased time spent in slow wave sleep (stage N3) by ~10.6% (**Fig. 3.a.**)

but did not markedly affect the time in other sleep stages or total sleep time (*SM 6*). As most

slow wave sleep occurs in the first 4 hours of sleep and the absorption profile of L-DOPA

controlled release strongly predicts that dopamine would be increased in the first half of the

night (*37*), we expected that dopamine would predominantly affect sleep during this time. As

predicted, the observed increase in slow wave sleep occurred only during the first half of the

night (as defined by the mid-point between lights-off and lights-on times) on L-DOPA (90.2 $\pm$

34.1 min) compared to placebo (76.8 $\pm$ 30.3 min, $(t(30) = -3.07, p = 0.005, BF_{10} = 8.7, n = 31,$

for missing data see *SM 7*). L-DOPA did not affect slow wave sleep duration during the second

half of the night $(t(30) = -0.387, p = 0.703, BF_{01} = 4.9)$.

Next, we explored the relationship between L-DOPA's effect on total slow wave sleep duration

with its effects on memory. Overall, longer slow wave sleep duration was associated with

enhanced accuracy for the repeated items (List *i*) on placebo (Spearman's $\varrho = 0.450, p = 0.009$).

This effect did not occur for List *ii* (non-repeated items), and it disappeared after participants

took L-DOPA (List *ii* Spearman's $\varrho = -0.043, p = 0.810$, **Fig. 3.b.,** *SM 8*). This suggests that

slow wave sleep duration is associated with consolidation of stronger memory traces, but it does

not suggest a direct relationship between slow wave sleep duration and acceleration of forgetting

of weaker memory traces on dopamine.

We next asked what neurophysiological processes underlie the quicker forgetting of weaker

compared to stronger memory traces on L-DOPA.

**Fig. 3. L-DOPA and slow wave sleep duration**

a. Paired differences in slow wave sleep duration show that most volunteers (dots above zero) had increased slow wave sleep on L-DOPA compared to placebo. The duration was increased by an average of ~10.6% on L-DOPA compared to placebo (t(31) = 2.702, p = 0.011, $BF_{10}$ = 4.0). This effect remained after false discovery rate correction accounting for each sleep stage ($p_{corrected}$ = 0.044).

b. Longer slow wave sleep duration was correlated with better memory for strongly encoded information on placebo (Spearman's ϱ = 0.45, p = 0.009, $p_{corrected}$ = 0.012, green), but after L-DOPA was given this effect disappeared (ϱ = 0.043, p = 0.810, red). The two relationships were different (Pearson's r-to-z = -1.99, p = 0.046). This strongly suggests that L-DOPA does not increase the relative effect of re-exposure by merely increasing slow wave sleep. Lines of best fit are presented for illustration.

L-DOPA increases spindle amplitude during slow wave sleep

Spindles are a prominent feature of NREM sleep and are associated with memory retention [41,

42]. L-DOPA increased slow wave sleep spindle amplitude, and while this increase was small on

average (m $_{placebo}$ = 28.3 ± 8.5 μV; m $_{L\text{-}DOPA}$ = 28.9 ± 83 μV; Wilcoxon's Z = 95.0, p$_{corrected}$ =

0.008, BF$_{10}$ = 3.6), this effect was manifest in 25 out of 31 participants with spindle data available

(**Fig. 4.a.,** *SM 9*). This change was not correlated with the weight adjusted dose (Pearson's r = -

0.139, p = 0.456), nor did we find any correlations between spindle amplitude and the d'

difference between Lists *i* and *ii* on either L-DOPA (Spearman's ϱ = 0.047, p = 0.801) or

Placebo (Spearman's ϱ = -0.040, p = 0.833, *SM 8*). However, the paired change of slow wave

spindle amplitude and the d' difference for strong and weak memories between the L-DOPA

and placebo nights was positively correlated (ϱ = 0.438, p = 0.015, n = 30, **Fig. 4.b.**).

In other words, the behavioural effect of L-DOPA on memory based on engram strength was

associated with the increase in spindle amplitude on L-DOPA. This effect was specific to the L-

DOPA-mediated *difference* in memory and spindle amplitude between strong and weak memory

traces, and it was not present for List *i* or *ii* alone (*SM 8*).

L-DOPA affects spindles most at slow oscillations peaks

Temporal coupling between slow oscillations and spindles have been shown to predict memory

performance, and this coupling is impaired by aging *(43)*. As we observed effects of L-DOPA on

both memory and spindle physiology, we next explored whether L-DOPA's effects on memory

performance could be due to an alteration in slow oscillation – spindle coupling.

Periods of time during which the maximum spindle amplitude occurred during a slow oscillation

events were identified, segmenting the slow oscillations into 16 phase bins (then grouped into 4

shaded and white areas in **Fig. 4c.**).L-DOPA had a slow oscillation phase dependent effect on

spindle amplitude, with a larger increase around the zero phase (**Fig. 4c.**). The peak change

occurred in the -π⁄4 to +π⁄4 grouping, the same that showed the highest mean spindle amplitude

for both L-DOPA and placebo conditions (**Fig. 4c., d.**).

To see whether this effect was specific to this spindle region-of-interest (ROI)we also

performed spectral composition analyses of each slow oscillation – spindle co-occurrence. A

spectral mean was calculated for each participant for L-DOPA and placebo conditions using

Morlet Wavelet convolution (**Fig. 4e**). Changes in power between L-DOPA and placebo were

then identified using a cluster-based permutation method. Cluster analysis of the *a priori* spindle

ROI revealed an increase in power on L-DOPA compared to placebo ($p = 0.002$). **Fig. 4e.**

demonstrates the primary cluster (threshold $\alpha = 0.01$ $p = 0.002$) when this same analysis was

carried out on the surrounding time-frequency space. It suggests that the power increase extends

beyond the spindle ROI, into theta and alpha bands following the slow oscillation peak.

L-DOPA therefore altered the neural dynamics that underlie the synchronised relationship

between slow oscillations, spindles, and potentially other frequencies. This may represent either a

phase-specific effect of dopamine during sleep, or a secondary effect on these dynamics caused

by a dopaminergic bias of early awake consolidation or re-exposure.

We found no associations between L-DOPA and other slow oscillation characteristics (all ps >

0.09, *SM 9*). Exploratory analyses revealed no differences between L-DOPA and placebo on

subjective sleep measures (St Mary's Hospital Sleep Questionnaire *(44)* or Leeds Sleep

Evaluation Questionnaire *(45)* (*SM 10).*

**a** L-DOPA minus placebo

**b** L-DOPA minus placebo

**e**

**Fig. 4. L-DOPA, memory, and spindle amplitude**

**a.** Nocturnal L-DOPA increased spindle amplitude (n = 31, Wilcoxon's z = 401, p = 0.002, p corrected = 0.008, $BF_{10}$ = 3.6) suggesting an effect of L-DOPA on regional coherence during slow wave spindles.

**b.** The L-DOPA mediated increase in spindle amplitude was associated with the L-DOPA mediated increase in the relative benefit of re-exposure on d' (**Fig. 2c.**) (Spearman's ϱ = 0.438, p = 0.015). Note that this relationship is non-linear, line is fitted in the figure for illustration.

**c.** The dopamine-induced spindle amplitude increase is slow oscillation phase-dependent. Mean spindle amplitude change (normalised to baseline amplitude ([placebo + L-DOPA]/2) is higher on L-DOPA around the zero phase of slow oscillations. We compared the effect of L-DOPA at the peak (zero phase) and trough ($\pi$ phase) of the slow oscillation. The L-DOPA mediated spindle amplitude increase was larger in the 4 zero-centric bins compared to the 4 $\pi$-centric bins (outermost on either side) – (paired t(30) = 2.12, p=0.043, $BF_{10}$ = 1.3). Yellow bars show the mean amplitude change with individual participants' spindle amplitude change overlaid. Spindle amplitude peaked in the -$\pi$/4 to -$\pi$/8 phase bin for both placebo and L-DOPA.

**d.** Peak-locked grand average mean slow oscillation events (grey) superimposed with the peak-locked average of all spindle events (blue) that occurred during slow oscillations – averaged across both L-DOPA and placebo nights.

**e.** Paired permutation cluster analysis in time-frequency space comparing L-DOPA and placebo conditions for all slow oscillation - spindle co-occurrence events, centered on the slow oscillation peak. All shown differences denote increased activity on L-DOPA (cluster threshold of $\alpha = 0.01$, time-frequency space outside significant clusters is greyed). Overall p = 0.002 for the largest cluster.

### L-DOPA does not modulate memory at encoding or retrieval – Secondary experiment

Whilst there were no differences in performance between L-DOPA and placebo during the re-

exposure test in the evening (*SM 4)*, it is possible that the observed effect of L-DOPA on

memory was driven by either L-DOPA enhancing encoding during the re-exposure or by

residual amounts of L-DOPA acting during retrieval. To investigate whether dopamine was

affecting these stages of memory, we ran a secondary placebo-controlled experiment

manipulating dopamine levels at encoding or retrieval.

In the secondary experiment, healthy elderly participants were given short-acting L-DOPA an

hour before encoding and, in a separate memory task, an hour before retrieval (*SM 11, 12)*. Note

that we used short-acting Levodopa here rather than controlled-release (as in the main

experiment) as the processes we were probing were discrete events rather than evolving

processes. We did not find an effect of L-DOPA on encoding (t(28) = - 0.352, p = 0.728, $BF_{01}$=

4.6, n = 32) or retrieval (t(27) = - 0.393, p = 0.698, $BF_{01}$= 4.6, n = 28) with a 24-hour delay

between learning and test (*SM 12, 14*, for missing data see *SM 7)*. Therefore, at the doses and

timings used here, dopamine appears to have a temporally and functionally specific effect biasing

memory towards important information after initial learning, during either re-exposure, sleep or

both.

Whilst the results from this control study support our initial interpretation that L-DOPA affects

memory *after* initial learning but *before* retrieval, it is important to note that due to practical

reasons direct statistical comparisons between the two studies cannot be made. First, due to the

different profiles of the treatments used (controlled release versus short acting), the L-DOPA

doses participants were exposed to were different; 2.1 mg/kg for the control study compared to

2.9 mg/kg for the main experiment. Second, whilst the memory tasks used were similar word-list

tasks, they were not identical.

Discussion

Dopamine accelerates forgetting for weakly encoded information during sleep – while more

strongly encoded information is relatively preserved – and increases duration of slow wave sleep

by 10.6%. The behavioural effect of dopamine on strongly versus weakly encoded information is

associated with a dopamine-driven increase in spindle amplitude during slow wave sleep. This

increase in spindle amplitude only occurs around the peak of slow oscillations.

Traditionally, forgetting is considered a passive process where information is "lost". However,

newer animal models support an active, more strategic, forgetting process mediated by dopamine

*(31, 33, 34, 46)*. We showed that dopamine increased forgetting for information at 1-day delay

but not at later timepoints. Therefore, dopamine may accelerate forgetting of low importance

information that would inevitably be lost over time. Our data suggest an *active* dopamine-

dependent forgetting mechanism in humans – which can be conceived as dopamine biasing

memory selection away from weakly encoded items. This may in turn allow prioritisation of

strongly encoded or salient items for consolidation.

Such prioritisation may be further explained – through analogy with drosophila experiments – by

a second dopaminergic system that protects important information from forgetting (46). Human

behavioural evidence supports preferential consolidation of salient or rewarded information

during sleep (19, 47, 48), and we tie this more closely to dopaminergic modulation. Contrary to

hypothesis,, we did not find evidence for a dopamine-driven direct enhancement in

consolidation of strongly encoded information here. The relative effect of dopamine on

forgetting of low versus high importance items  suggests a more nuanced dopaminergic effect -

biasing memory away from weaker memory traces Given L-DOPA is already widely used in

clinical practice for Parkinson's disease and could be quickly repurposed if this effect is beneficial

for memory overall, this certainly warrants attention and further clinical research.

There is clear evidence that memory processes before sleep can alter slow wave sleep

characteristics, particularly in the early part of the night (49). We administered dopamine while

participants were awake, 2h before their bedtime, thus it is possible that at least a portion of the

dopamine-driven increase in forgetting occurred during wake. Whilst we cannot rule this out, we

did observe that L-DOPA compared to placebo was associated with changes in scalp

electrophysiology during sleep, with some of these effects being associated with memory

performance. Therefore, we suggest that the dopamine-driven changes on memory were sleep-

dependent.

While changes in spindle characteristics are well known to be associated with memory and neurodegeneration (50), this study directly links dopamine with behavioural relevance of spindles. Spindle amplitude is shaped by the interplay between the thalamus and the cortex (51), and increased amplitude reflects a more coherent and wider topographical expression of spindle-related activity, i.e. better coordination between the brain regions (52, 53). Behaviourally, spindle amplitude has also been associated with enhanced memory retention during a motivated forgetting task (54) and during a tagging paradigm (55). This coordinated activity between the thalamus and cortex during sleep may thus be associated with selecting memories for later retention. Consistent with this, we showed that greater spindle amplitude was associated with a larger dopamine-induced difference between retention of strongly and weakly encoded information.

L-DOPA mainly increased spindle amplitude around the peak of slow oscillations, which occurred despite no change in slow-oscillation amplitude. Spindles peaked just before zero phase, consistent with previous findings in healthy elderly (43). Spindles, particularly when nested in slow oscillation peaks, are hallmarks of sleep-dependent memory consolidation (56). Age-related uncoupling of spindles from peak of slow oscillations increases overnight forgetting (43). We interpret dopaminergic increase in spindles synchronised to near zero phase of slow oscillation as enhancement of physiological spindle activity to modulate memory consolidation.

There are two possible explanations for our finding – (1) dopamine directly enhances spindle amplitude which in turn enhances the way in which memory is biased based on encoding strength (2) or dopamine during memory re-exposure before sleep results in stronger behavioural tags that in turn alter subsequent spindle amplitude to reflect the changes in the memory engram that took place during tagging. These effects are not mutually exclusive, and indeed could be

interacting. Future experiments separating the effects of sleep consolidation from re-exposure benefit are necessary to disentangle this.

We suggest that two simultaneous processes may be at play (**Fig. 5.**). First, during learning a portion of information is "tagged" as important (57), and dopamine enhances this process by creating a stronger tag (58, 59). Second, during subsequent sleep, dopamine increases forgetting for the less important, non-tagged items while the tag shields the important (or re-exposed) information from forgetting (60, 61). This theory has been proposed before, and the current study adds to it by implicating (dopamine-mediated) crosstalk between the thalamus and the cortex during spindles as a potential mechanism for the later effects.

Our observed findings may be specific to older people. Memory loss is a prominent problem in old age and our eventual goal is to improve quality of life through cognitive enhancement, justifying the use of a target population of interest to future trials. There is drop-out of dopaminergic neurons that comes with old age (62-64) (62-64) which has been shown to affect the impact of taking dopaminergic medications on cognition (65). Ageing decreases the duration of slow wave sleep, and the number and amplitude of spindles (66), with some reporting nearly a 50% reduction in spindle amplitude with advanced age (67). Models in non-aged animals suggest that D2 receptors promote wakefulness (68) and dopamine levels are generally higher during wake than sleep in animals (69). In young healthy adults, direct administration of a dopamine *antagonist* during slow wave sleep actually increases the duration of slow waves sleep (70). It has been noted before that the wake-promoting effects of dopamine in the young contradict the sleepiness that is a recognised side effect of L-DOPA in patients with Parkinson's disease (71). Therefore, age should be considered when interpreting the effects of dopamine on memory and sleep.

Furthermore, slow wave sleep may be affected early in Alzheimer's Disease (72). Interrupting

slow wave sleep is proposed to hinder clearance of amyloid from the brain and amyloid plaques

are one of the key pathological changes in Alzheimer's Disease (73, 74). L-DOPA is routinely

prescribed for Parkinson's disease with a good safety profile; however, the impacts of L-DOPA

on sleep have not been assessed in detail except in small studies of Parkinson's disease (75-77).

Our finding that L-DOPA may ameliorate age-dependent spindle loss with concomitant memory

benefits could be promising for treating age-related memory decline, or more severe memory

deficits found in Alzheimer's dementia. Perhaps more excitingly, our current findings may have

implications for prevention of Alzheimer's disease. Through increasing slow wave sleep duration

and spindle amplitude with nocturnal dopamine, we open up a new therapeutic avenue for

Alzheimer's disease prevention – repurposing L-DOPA to prevent Alzheimer's.

Together, our findings suggest that the repetition-benefit on memory is improved by dopamine

at the time of the repetition and during sleep-consolidation, which is mediated by increased slow

wave sleep duration and spindle amplitude. We propose that this dopamine-induced increase in

spindle amplitude reflects more synchronous cortical activity during spindles increasing

forgetting of weakly encoded items and with a net effect of augmenting the difference between

strongly and weakly encoded engrams (**Fig. 5.**). These findings have potential clinical impact in

enhancing sleep and memory selection in old age, and in mild amnesic disease.

**Fig. 5. Dopamine modulates memory after learning by enhancing memory prioritisation and subsequent sleep processes.**

A proportion of important information (yellow engram) is earmarked for retention by a neural "tag" during re-exposure (blue pin). Dopamine during re-exposure enhances this effect (red pins) at the expense of unpinned engrams (green). During sleep, weak engrams are preferentially forgotten whilst ear-marked information remains unaffected, leading to a more selective memory trace. Dopamine modulates these selective memory processes by enhancing synchronisation in cortical firing patterns during spindles, at the peak of slow oscillations. Together these two processes (enhanced prioritisation and synchronisation) bias subsequent memory.

Method

**Participants**

We recruited 70 elderly (65+ years) volunteers to complete the two studies reported here (n = 35

each study, see *SM 4, 15, 16*). All aspects of this research adhered with the Declaration of

Helsinki and we had relevant ethical and regulatory (UK) approvals in place (Study 1 ISRCTN:

90897064).

**Design**

<u>Study 1:</u> In the main placebo-controlled double-blind randomised study, volunteers were initially

screened over the phone for common exclusions, and then invited for three in-house visits. On

the first visit they were fully screened for eligibility, and they practiced the memory task. They

were asked about their usual sleeping pattern so that the second and third visits could be

designed to follow each participants' usual sleep routines as much as possible. On the second

visit, volunteers arrived on site in the evening where they were re-consented and screened for

continued eligibility. For an outline of the evening see **Fig. 1a.**, *SM 11.*

First, volunteers learnt a verbal memory task (**Fig. 1b.**). Thirty minutes after learning, they were

given 200mg L-DOPA or placebo. 75 min after dosing, a quarter of the items (List *i*) were re-

exposed by a recognition test where no feedback was given. The purpose of this test was to

create a stronger memory trace. 45 min after the re-exposure the volunteers went to bed. Each

evening was designed based on each participants' usual sleeping pattern (L-DOPA administered

2h prior to switching the lights off for the night at their usual bedtime).

Volunteers slept on-site for a full night, and they were woken up at their usual wake-up time.

Around 1.5h after waking up, approximately 12h after dosing, volunteers' verbal memory was

tested again (Lists *i* and *ii* ) before they left the study site. 2 and 4 days later (3 and 5 days after

learning) they were contacted over the phone for follow-up recognition memory tests (for Lists

*iii* and *iv*, respectively).

The second and third visits were identical except for treatment (L-DOPA / placebo) allocation.

This study obtained ethical approval from the South West Central Bristol NHS Research Ethics

Committee (REF: 16/SW0028) and clinical trial authorisation from the Medicines and

Healthcare products Regulatory Agency (IRAS ID:178711) .

Study 2: In the secondary placebo-controlled double-blind crossover experiment, volunteers

were first screened over the phone before inviting them on site for the test sets. Each test set

carried over for three days: On the Day -1 (relative to dosing) participants learnt word-list on

site, on Day 0 they were dosed with 150mg L-DOPA (or placebo, *SM 11, 12*) and then tested on

the previously learnt words (retrieval test). Immediately following this, they learnt another word-

list (encoding condition) for which their memory was tested the following day (Day 1) over the

phone. Therefore, participants learnt two word lists on each test There tests were timed so there

was approximately 24h in between learning and test.

This study obtained ethical approval from the University of Bristol Faculty of Medicine and

Dentistry Ethics Committee (REF: 12161).

**Treatment**

Study 1: In the main placebo-controlled randomised double-blind study, each participant was dosed with co-beneldopa controlled release (containing 200mg L-DOPA) was given in capsule form and placebo (encapsulated inert powder, matched for appearance). Blinding and randomisation were performed in blocks of 6 by author LM, Production Pharmacy, Bristol Royal Infirmary, University Hospitals Bristol and Weston NHS Trust. On the study nights, dose was given by an on-site medic who was blind to treatment condition and played no role in collecting data. The treatments were given at different visits. Both treatments were preceded by Domperidone 10mg (tablet) to alleviate possible nausea caused by L-DOPA.

Study 2: In the L-DOPA condition of the secondary study volunteers received 10mg of Domperidone (anti-emetic) 30 minutes before co-beneldopa (containing 150mg L-DOPA). Both medications were dispersible, and they mixed into cordial to hide taste and residue. In the placebo condition, volunteers received plain cordial in place of Domperidone and vitamin C mixed into cordial in place of L-DOPA. Blinding and randomisation were performed by members of the research group who had no other involvement in this study.

While the two experiments were designed to complement one another, for practical reasons there were several important differences in study designs. First, the L-DOPA given in the main study was long-acting and of higher dose (4-8 hours cf. 1-4 hours and 200mg cf. 150mg) to target consolidation during sleep which is a longer process than encoding or retrieval. Second, the controlled release L-DOPA in the main study was encapsulated, whilst in the secondary study we used dispersible L-DOPA. For this reason, the placebo used in the main experiment was encapsulated inert powder, whilst in the second study we used dispersible vitamin C. These

differences and individual differences in dopamine absorption and metabolism introduce

unmeasurable differences between the two experiments that need to be considered when

interpreting differences between them.

**Verbal memory test**

<u>Study 1:</u>  Volunteers learnt four lists (*i, ii, iii,* and *iv*) of 20 target words (total 80 targets)

presented on a computer screen one at a time, in a random, interleaved order (**Fig. 1b.,** *SM 12*).

Each word was presented once for 3.6s during which the volunteers were asked to determine if

the items were alive or not to assist learning. They were instructed to remember as many of the

words as they could.

During test phases, volunteers were presented with a list of 40 (days 0, 3, and 5) or 80 words (day

1), half of which were targets (present at learning) and half of which were distractors (not

presented previously). They were asked to judge whether words were targets or not. On days 0,

1, 3, and 5 memory was tested for Lists *i*, *i* and *ii*, *iii*, and *iv* respectively. Therefore, List *i* was

tested twice: First in the evening while L-DOPA (/ placebo) was active in the system and then

again in the morning together with List *ii.* The re-exposed and novel (List *i* and List *ii*,

respectively) targets tested on day 1 were assessed to study L-DOPA's effect on behavioural

tagging of 'important' information. The rationale was that when a word is presented a second

time (during re-exposure), it will be deemed more important and will be preferentially

remembered. The distractors were unique at each test.

<u>Study 2:</u> The purpose of this study was to test L-DOPA's effects on retrieval and encoding. Two

separate memory tests were conducted (*SM 13*).

*Retrieval*: During learning on D-1 (day before dosing) volunteers were presented with 48

complete nouns on a computer screen. They were instructed to read the words aloud and try to

memorise them for later. Each word was shown once for 5 seconds separated by a fixation cross

in the middle of the screen for 2 seconds and no responses to the words were made during

learning. There were no breaks in the learning block (total duration = 5mins 36secs). Memory

was tested using unique words 30 minutes (D-1, baseline) and 24 hours (D0) after learning. The

D0 test was given when L-DOPA was at its peak concentration (~ 1h following dosing). In the

test phases (D-1 and D0). This test took approximately 5 minutes to administer.

*Encoding*: D0 around 1.5 hours after dosing, after the test for the previous task had finished,

volunteers saw a list of 96 complete nouns presented on the computer screen. Each word was

displayed for 5 seconds, followed by a fixation cross for 2 seconds. The words were first

presented in a random order over two blocks, and then again in a different random order over

two more blocks, each word was presented twice (n blocks = 4, n words per block = 48, n

breaks = 3, block duration = 5 min 36 s). Memory was prompted immediately after learning

(D0), and 1, 3, and 5 days later. Each target was tested once with unique distractors (*SM 14*).

Across experiments, learning and tests were completed on a laptop on-site, or over the phone.

The experiments were programmed in the MATLAB environment (2015b or 2017a) using the

Psychophysics Toolbox V3 (78). The scripts and data are available from corresponding authors

upon request.

**Polysomnography**

In the main experiment, standard in-laboratory polysomnography, including video, was recorded

during both study nights using the Embla N9000 amplifier and Embla RemLogic software

(Natus Medical Inc., California) at CRIC Bristol, University of Bristol, Bristol, UK. We recorded

12 scalp EEG channels (F3, Fz, F4, C3, Cz, C4, M1, Pz, M2, O1, O2, and a ground electrode

approximately between Cz/P3 and C3/Pz) placed according to the 10-20-20 system. Eye

movements were detected by electro-oculogram recorded from E1 and E2 sites, and muscle tone

from electromyogram recorded below the chin. A 2-lead ECG was also recorded. All signals

were sampled at 500Hz. The recordings started 2.5h after dosing when lights were switched off

for the night and continued until the volunteer woke up.

**Analysis**

*EEG*

**Event scoring:** Sleep stages in 30s epochs were identified manually in accordance to standard

criteria *(79)* by two expert scorers, and a third scorer visually assessed a random 10% of ratings

for quality. Durations of N1, N2, N3 (i.e. slow wave sleep), REM, awake, asleep and total time in

bed were extracted in minutes. First and second halves of the nights were defined by the middle

time-point between switching lights ON and OFF. When there was an odd number of epochs,

they were rounded so that the first half of the night had the extra epoch.

**Spindle detection:** Spindle characteristics were then isolated with in-house written MATLAB

scripts using the EEGlab toolbox *(80)*. Electrodes were re-referenced to contralateral mastoid

and empty and high variance epochs were removed. Thereafter, only data from the Cz electrode

was used. First, the channel was visually inspected and epochs with high noise or clear artefacts

were removed manually. Data was then filtered (high pass 11Hz, low pass 17Hz) and rectified.

Next data was smoothed using a moving average window of 200ms before down-sampling to

100Hz (from 500Hz) for computational efficiency. An event was marked as a spindle if the

threshold exceeded the 90th percentile for that data set (i.e. sorting data into an ascending order

and including top 10%) for .5 – 3 seconds, with a minimum of 0.5s between spindle events.

**Slow oscillations:** The slow oscillation detection process followed the same re-referencing and

noise removal methods used for spindle detection, without smoothing. Data from the CZ

electrode was filtered between 0.16Hz and 1.25Hz and then z-scored. We applied a threshold of

75%; if the slow oscillation amplitude surpassed this threshold for 0.5 - 5 seconds (including

multiple events if separated by <0.25s), it was marked as a slow oscillation. The duration of the

event was determined by the closest oscillation maxima following the amplitude dropping below

a 60% threshold on each side.

**Spindles and slow oscillations:** We identified spindle-slow oscillation co-occurrences as cases

where the maximum amplitude of a spindle event coincided with a slow oscillation event, again

using the CZ electrode.  Using the time stamp of the spindle max amplitude as the centre point,

we calculated how spindle amplitude varied with slow oscillation phase over one cycle. First, we

divided the oscillation events into 16 bins, equally distributed in phase space around zero, to

calculate how the spindle amplitude varied with slow oscillation phase for each coinciding case

(for statistical analysis we grouped together 4 adjacent frequency data points (bins) to generate 4

bins as shown in **Fig. 4c**).

The spectral composition of each was done using a Morlet wavelet time-frequency method over

a 4s window centred on the slow oscillation peak. Morlet waves at 20 frequencies were used,

with 3 cycles for the lowest frequency (1Hz) and 6 cycles otherwise (2-20Hz). A spectral mean

was next calculated for each participant for L-DOPA and placebo conditions (**Fig. 4e.**). A

cluster-based permutation method (permutation n = 500, cluster threshold of $\alpha = 0.01$),

implemented with the Fieldtrip toolbox (*81*), was used to identify power differences. Based on

the finding that max spindle amplitude occurring near zero slow oscillation phase predicts to

memory performance in ageing {Helfrich, 2018 #3297}, an a-priori spindle region of interest of

11 - 16Hz, -0.5 - 0.5s was chosen for initial analysis. This same cluster method was then carried

out on the wider time-frequency space, $1 - 20Hz$, $-1 - 1s$, the primary cluster (p=0.002) of which

is shown in **Fig. 4e.**.

Behaviour

**Pairwise comparisons** (placebo versus L-DOPA) were calculated using either t-tests or

Wilcoxon's rank tests in R 3.5.3 using RStudio. We also employed a Bayesian paired t-tests in

JASP 0.9.2.0 (*82*) to obtain Bayes Factors (BF) – this allows more meaningful estimates of

confidence in both significantly different and null results than standard t-tests. BF gives the

probability of the data under either hypothesis. E.g. a $BF_{10}$ of 5 would denote that the data is 5

times more likely to have been sampled from the alternative compared to the null distribution,

while a $BF_{01}$ of 5 would denote that the data is 5 times less likely to have been sampled from the

alternative compared to the null distribution (i.e. 01 versus 10). We defined the prior (expected)

distribution as a Cauchy distribution with a mean of 0 and an interquartile range of .5 [$\delta \sim$

Cauchy (0, .5)]. In other words, we predicted that the $\delta$ lies between -.5 and .5 with a 50%

confidence. We selected this one as the $\delta$s in cognitive neurosciences typically are within those

bounds, and as we did not have an informed prediction for the effect sizes.

All mixed modelling was performed on R 3.5.3 using Rstudio, lme4 *(83)* and lmerTest *(84)*. We

included the participants as random effects and the dose (mg/kg) and the memory test delay

(Day 1, Day 3 and Day 5), or memory strength (re-enforced versus not), depending on the

analysis, as fixed effects. All fixed effects were mean-centred but not scaled. We selected the

model using the maximum feasible fit as this has previously been shown to be the best approach

for confirmatory hypothesis testing *(85)*.

**List of supplementary materials:**

SM 1 : Paired differences for delayed memory test

SM 2 : L-DOPA accelerates forgetting.

SM 3 : Re-activated items better retained in both conditions

SM 4 : Pairwise accuracy for List i across tests

SM 5: L-DOPA has disparate effects on forgetting rate depending on whether items were re-

exposed or not

SM 6 : Single dose of nocturnal L-DOPA increases time spent in slow wave sleep by 10.6%.

SM 7 : Missing data

SM 8 : Sleep and memory correlations on L-DOPA and placebo

SM 9 : L-DOPA increases spindle amplitude

SM 10 : Subjective sleep measures

SM 11 : Sleep visit timeline

SM 12 : Verbal Memory Task

SM 13 : Secondary study timeline

**References**

1.    S. Tonegawa, X. Liu, S. Ramirez, R. Redondo, Memory Engram Cells Have Come of Age.

*Neuron* **87**, 918-931 (2015).

2.    R. Semon, *Die nmemischen Empfindungen*.  (Willhelm Engelmann, Leipzig, 1909).

3.    H. Eichenbaum, P. Dudchenko, E. Wood, M. Shapiro, H. Tanila, The hippocampus, memory,

and place cells: is it spatial memory or a memory space? *Neuron* **23**, 209-226 (1999).

4.    J. L. McGaugh, Making lasting memories: remembering the significant. *Proc Natl Acad Sci U S*

*A* **110 Suppl 2**, 10402-10407 (2013).

5.    L. R. Squire, L. Genzel, J. T. Wixted, R. G. Morris, Memory Consolidation. *Csh Perspect Biol* **7**,

(2015).

6.    J. Lisman, A. A. Grace, E. Duzel, A neoHebbian framework for episodic memory; role of

dopamine-dependent late LTP. *Trends in Neurosciences* **34**, 536-547 (2011).

7.    G. B. Feld, L. Besedovsky, K. Kaida, T. F. Munte, J. Born, Dopamine D2-like Receptor

Activation Wipes Out Preferential Consolidation of High over Low Reward Memories during

Human Sleep. *J Cognitive Neurosci* **26**, 2310-2320 (2014).

8.    J. P. Grogan, R. Bogacz, D. Tsivos, A. Whone, E. J. Coulthard, Dopamine and Consolidation of

Episodic Memory: Timing is Everything. *J Cogn Neurosci* **27**, 2035-2050 (2015).

9.   R. Chowdhury, M. Guitart-Masip, N. Bunzeck, R. J. Dolan, E. Duzel, Dopamine Modulates

Episodic Memory Persistence in Old Age. *Journal of Neuroscience* **32**, 14193-14204 (2012).

10.  A. S. Franca *et al.*, D2 dopamine receptor regulation of learning, sleep and plasticity.

*European neuropsychopharmacology : the journal of the European College of*

*Neuropsychopharmacology* **25**, 493-504 (2015).

11.  E. J. Coulthard *et al.*, Distinct roles of dopamine and subthalamic nucleus in learning and

probabilistic decision making. *Brain* **135**, 3721-3734 (2012).

12.  U. Frey, R. G. M. Morris, Synaptic tagging and long-term potentiation. *Nature* **385**, 533-536

(1997).

13.  R. L. Redondo, R. G. Morris, Making memories last: the synaptic tagging and capture

hypothesis. *Nature reviews. Neuroscience* **12**, 17-30 (2011).

14.  S. H. Wang, R. L. Redondo, R. G. M. Morris, Relevance of synaptic tagging and capture to the

persistence of long-term potentiation and everyday spatial memory. *Proceedings of the*

*National Academy of Sciences of the United States of America* **107**, 19537-19542 (2010).

15.  A. J. Duszkiewicz, C. G. McNamara, T. Takeuchi, L. Genzel, Novelty and Dopaminergic

Modulation of Memory Persistence: A Tale of Two Systems. *Trends in Neurosciences* **42**, 102-

114 (2019).

16.  G. B. Feld, J. Born, Neurochemical mechanisms for memory processing during sleep: basic

findings in humans and neuropsychiatric implications. *Neuropsychopharmacol*,  (2019).

17.  F. Michon, J. J. Sun, C. Y. Kim, D. Ciliberti, F. Kloosterman, Post-learning Hippocampal Replay

Selectively Reinforces Spatial Memory for Highly Rewarded Locations. *Current biology : CB*

**29**, 1436-1444.e1435 (2019).

18.  D. Liu, Y. Dan, A Motor Theory of Sleep-Wake Control: Arousal-Action Circuit. *Annual review*

*of neuroscience* **42**, 27-46 (2019).

19.    D. Oudiette, J. W. Antony, J. D. Creery, K. A. Paller, The Role of Memory Reactivation during

Wakefulness and Sleep in Determining Which Memories Endure. *Journal of Neuroscience* **33**,

6672-U6758 (2013).

20.    R. Stickgold, Sleep-dependent memory consolidation. *Nature* **437**, 1272-1278 (2005).

21.    G. M. van de Ven, S. Trouche, C. G. McNamara, K. Allen, D. Dupret, Hippocampal Offline

Reactivation Consolidates Recently Formed Cell Assembly Patterns during Sharp Wave-

Ripples. *Neuron* **92**, 968-974 (2016).

22.    M. Molle, O. Yeshenko, L. Marshall, S. J. Sara, J. Born, Hippocampal sharp wave-ripples linked

to slow oscillations in rat slow-wave sleep. *J Neurophysiol* **96**, 62-70 (2006).

23.    M. Y. Yang, N. K. Logothetis, O. Eschenko, Occurrence of Hippocampal Ripples is Associated

with Activity Suppression in the Mediodorsal Thalamic Nucleus. *Journal of Neuroscience* **39**,

434-444 (2019).

24.    N. Maingret, G. Girardeau, R. Todorova, M. Goutierre, M. Zugaro, Hippocampo-cortical

coupling mediates memory consolidation during sleep. *Nat Neurosci* **19**, 959-964 (2016).

25.    A. Patil, V. P. Murty, J. E. Dunsmoor, E. A. Phelps, L. Davachi, Reward retroactively enhances

memory consolidation for related items. *Learning & Memory* **24**, 65-69 (2017).

26.    J. E. Dunsmoor, V. P. Murty, L. Davachi, E. A. Phelps, Emotional learning selectively and

retroactively strengthens memories for related events. *Nature* **520**, 345-348 (2015).

27.    M. C. Anderson, R. A. Bjork, E. L. Bjork, Remembering Can Cause Forgetting - Retrieval

Dynamics in Long-Term-Memory. *J Exp Psychol Learn* **20**, 1063-1087 (1994).

28.    M. Alizadeh Asfestani, V. Brechmann, J. Santiago, J. Born, G. B. Feld, Consolidation of reward

memory during sleep does not require dopaminergic activation. *BioRxiv*,  (2019, bioRxiv).

29.    C. G. McNamara, A. Tejero-Cantero, S. Trouche, N. Campo-Urriza, D. Dupret, Dopaminergic

neurons promote hippocampal reactivation and spatial memory persistence. *Nat Neurosci*

**17**, 1658-1660 (2014).

30.     S. N. Gomperts, F. Kloosterman, M. A. Wilson, VTA neurons coordinate with the

hippocampal reactivation of spatial experience. *Elife* **4**, (2015).

31.     R. L. Davis, Y. Zhong, The Biology of Forgetting-A Perspective. *Neuron* **95**, 490-503 (2017).

32.     J. T. Wixted, The psychology and neuroscience of forgetting. *Annu Rev Psychol* **55**, 235-269

(2004).

33.     J. A. Berry, I. Cervantes-Sandoval, E. P. Nicholas, R. L. Davis, Dopamine Is Required for

Learning and Forgetting in Drosophila. *Neuron* **74**, 530-542 (2012).

34.     J. A. Berry, A. Phan, R. L. Davis, Dopamine Neurons Mediate Learning and Forgetting through

Bidirectional Modulation of a Memory Trace. *Cell Rep* **25**, 651-+ (2018).

35.     M. C. Anderson, E. L. Bjork, R. A. Bjork, Retrieval-induced forgetting: Evidence for a recall-

specific mechanism. *Psychon B Rev* **7**, 522-530 (2000).

36.     M. Wimber, A. Alink, I. Charest, N. Kriegeskorte, M. C. Anderson, Retrieval induces adaptive

forgetting of competing memories via cortical pattern suppression. *Nat Neurosci* **18**, 582-+

(2015).

37.     A. Hsu, H. M. Yao, S. Gupta, N. B. Modi, Comparison of the pharmacokinetics of an oral

extended-release capsule formulation of carbidopa-levodopa (IPX066) with immediate-

release carbidopa-levodopa (Sinemet((R))), sustained-release carbidopa-levodopa

(Sinemet((R)) CR), and carbidopa-levodopa-entacapone (Stalevo((R))). *Journal of clinical*

*pharmacology* **55**, 995-1003 (2015).

38.     T. G. Birdsall, R. A. Roberts, On the Theory of Signal Detectability - an Optimum

Nonsequential Observation-Decision Procedure. *Ieee Transactions on Information Theory* **11**,

195-204 (1965).

39.     Y. Benjamini, Y. Hochberg, Controlling the False Discovery Rate - a Practical and Powerful

Approach to Multiple Testing. *J R Stat Soc B* **57**, 289-300 (1995).

40.     M. Zappia *et al.*, Body weight influences pharmacokinetics of levodopa in Parkinson's

disease. *Clin Neuropharmacol* **25**, 79-82 (2002).

41.    S. A. Cairney, A. A. V. Guttesen, N. El Marj, B. P. Staresina, Memory Consolidation Is Linked to Spindle-Mediated Information Processing during Sleep. *Curr Biol* **28**, 948-+ (2018).

42.    J. W. Antony *et al.*, Sleep Spindle Refractoriness Segregates Periods of Memory Reactivation. *Curr Biol* **28**, 1736-1743 e1734 (2018).

43.    R. F. Helfrich, B. A. Mander, W. J. Jagust, R. T. Knight, M. P. Walker, Old Brains Come Uncoupled in Sleep: Slow Wave-Spindle Synchrony, Brain Atrophy, and Forgetting. *Neuron* **97**, 221-230 e224 (2018).

44.    B. W. Ellis *et al.*, The St. Mary's Hospital Sleep Questionnaire: A Study of Reliability. *Sleep* **4**, 93-97 (1981).

45.    A. C. Parrott, I. Hindmarch, The Leeds Sleep Evaluation Questionnaire in psychopharmacological investigations - a review. *Psychopharmacology (Berl)* **71**, 173-179 (1980).

46.    J. A. Berry, I. Cervantes-Sandoval, M. Chakraborty, R. L. Davis, Sleep Facilitates Memory by Blocking Dopamine Neuron-Mediated Forgetting. *Cell* **161**, 1656-1667 (2015).

47.    I. Wilhelm *et al.*, Sleep Selectively Enhances Memory Expected to Be of Future Relevance. *Journal of Neuroscience* **31**, 1563-1569 (2011).

48.    E. van Rijn, C. Lucignoli, C. Izura, M. T. Blagrove, Sleep-dependent memory consolidation is related to perceived value of learned material. *J Sleep Res* **25**, 204-205 (2016).

49.    M. Mölle, O. Eschenko, S. Gais, S. J. Sara, J. Born, The influence of learning on sleep slow oscillations and associated spindles and ripples in humans and rats. **29**, 1071-1081 (2009).

50.    V. Latreille *et al.*, Sleep spindles in Parkinson's disease may predict the development of dementia. *Neurobiology of aging* **36**, 1083-1090 (2015).

51.    D. Contreras, A. Destexhe, T. J. Sejnowski, M. Steriade, Spatiotemporal patterns of spindle oscillations in cortex and thalamus. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **17**, 1179-1196 (1997).

52.    Y. Nir *et al.*, Regional slow waves and spindles in human sleep. *Neuron* **70**, 153-169 (2011).

53.  R. Cox, A. C. Schapiro, D. S. Manoach, R. Stickgold, Individual Differences in Frequency and

Topography of Slow and Fast Sleep Spindles. *Front Hum Neurosci* **11**, 433 (2017).

54.  B. Blaskovich, A. Szollosi, F. Gombos, M. Racsmany, P. Simor, The Benefit of Directed

Forgetting Persists After a Daytime Nap: The Role of Spindles and Rapid Eye Movement

Sleep in the Consolidation of Relevant Memories. *Sleep* **40**,  (2017).

55.  D. P. J. Heib *et al.*, Oscillatory Theta Activity during Memory Formation and Its Impact on

Overnight Consolidation: A Missing Link? *J Cognitive Neurosci* **27**, 1648-1658 (2015).

56.  B. P. Staresina *et al.*, Hierarchical nesting of slow oscillations, spindles and ripples in the

human hippocampus during sleep. *Nat Neurosci* **18**, 1679-1686 (2015).

57.  U. Frey, R. G. M. Morris, Weak before strong: Dissociating synaptic tagging and plasticity

factor accounts of late-LTP. *Eur J Neurosci* **10**, 20-20 (1998).

58.  C. Clopath, Synaptic consolidation: an approach to long-term learning. *Cogn Neurodynamics*

**6**, 251-257 (2012).

59.  S. Sajikumar, J. U. Frey, Late-associativity, synaptic tagging, and the role of dopamine during

LTP and LTD. *Neurobiol Learn Mem* **82**, 12-25 (2004).

60.  C. O'Donnell, T. J. Sejnowski, Selective Memory Generalization by Spatial Patterning of

Protein Synthesis. *Neuron* **82**, 398-412 (2014).

61.  G. Rauchs *et al.*, Sleep Contributes to the Strengthening of Some Memories Over Others,

Depending on Hippocampal Activity at Learning. *Journal of Neuroscience* **31**, 2563-2568

(2011).

62.  S. W. MacDonald, S. Karlsson, A. Rieckmann, L. Nyberg, L. Backman, Aging-related increases

in behavioral variability: relations to losses of dopamine D1 receptors. *The Journal of*

*neuroscience : the official journal of the Society for Neuroscience* **32**, 8186-8191 (2012).

63.  S. J. Kish, K. Shannak, A. Rajput, J. H. N. Deck, O. Hornykiewicz, Aging Produces a Specific

Pattern of Striatal Dopamine Loss - Implications for the Etiology of Idiopathic Parkinsons-

Disease. *J Neurochem* **58**, 642-648 (1992).

64.  A. Carlsson *et al.*, Biogenic amines in human brain in normal aging, senile dementia, and

chronic alcoholism. *Adv Biochem Psychopharmacol* **23**, 295-304 (1980).

65.  A. M. Morcom *et al.*, Memory Encoding and Dopamine in the Aging Brain: A

Psychopharmacological Neuroimaging Study. *Cerebral Cortex* **20**, 743-757 (2010).

66.  A. Nicolas, D. Petit, S. Rompre, J. Montplaisir, Sleep spindle characteristics in healthy

subjects of different age groups. *Clinical neurophysiology : official journal of the*

*International Federation of Clinical Neurophysiology* **112**, 521-527 (2001).

67.  K. Crowley, J. Trinder, Y. Kim, M. Carrington, I. M. Colrain, The effects of normal aging on

sleep spindle and K-complex production. *Clinical neurophysiology : official journal of the*

*International Federation of Clinical Neurophysiology* **113**, 1615-1622 (2002).

68.  W.-M. Qu *et al.*, Essential Role of Dopamine $D_2$ Receptor in the Maintenance of

Wakefulness, But Not in Homeostatic Regulation of Sleep, in Mice. **30**, 4382-4389 (2010).

69.  H. Dong *et al.*, Dorsal Striatum Dopamine Levels Fluctuate Across the Sleep–Wake Cycle and

Respond to Salient Stimuli in Mice. **13**,  (2019).

70.  D. N. Eder, M. Zdravkovic, G. Wildschiødtz, Selective alterations of the first NREM sleep cycle

in humans by a dopamine D1 receptor antagonist (NNC-687). *Journal of psychiatric research*

**37**, 305-312 (2003).

71.  C. R. Cantor, M. B. Stern, Dopamine agonists and sleep in Parkinson's disease. *Neurology* **26**,

71-78 (2002).

72.  M. P. Walker, The Role of Slow Wave Sleep in Memory Processing. *Journal of clinical sleep*

*medicine : JCSM : official publication of the American Academy of Sleep Medicine* **5**, S20-S26

(2009).

73.  L. Xie *et al.*, Sleep Drives Metabolite Clearance from the Adult Brain. *Science* **342**, 373-377

(2013).

74.  Y.-E. S. Ju *et al.*, Slow wave sleep disruption increases cerebrospinal fluid amyloid-β levels.

*Brain* **140**, 2104-2111 (2017).

75.    N. J. Diederich, V. Paolini, M. Vaillant, Slow wave sleep and dopaminergic treatment in

867           Parkinson's disease: a polysomnographic study. *Acta Neurol Scand* **120**, 308-313 (2009).

868    76.    S. P. Joy *et al.*, Serial macro-architectural alterations with levodopa in Parkinson's disease:

869           Polysomnography (PSG)-based analysis. *Annals of Indian Academy of Neurology* **18**, 309-313

870           (2015).

871    77.    S. Wailke, J. Herzog, K. Witt, G. Deuschl, J. Volkmann, Effect of controlled-release levodopa

872           on the microstructure of sleep in Parkinson's disease. *European journal of neurology* **18**, 590-

873           596 (2011).

874    78.    M. Kleiner, D. Brainard, D. Pelli, What's new in Psychtoolbox-3? *Perception* **36**, 14-14 (2007).

875    79.    R. B. Berry *et al.*, AASM Scoring Manual Updates for 2017 (Version 2.4). *J Clin Sleep Med* **13**,

876           665-666 (2017).

877    80.    A. Delorme, S. Makeig, EEGLAB: an open source toolbox for analysis of single-trial EEG

878           dynamics including independent component analysis. *J Neurosci Meth* **134**, 9-21 (2004).

879    81.    R. Oostenveld, P. Fires, E. Maris, J. Schoffelen, FieldTrip: Open Source Software for Advanced

880           Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Computational Intelligence*

881           *and Neuroscience* **2011**,  (2011).

882    82.    JASP. (2018).

883    83.    LME4-Authors. (2019).

884    84.    A. Kuznetsova, P. B. Brockhoff, R. H. B. Christensen. (2019).

885    85.    D. J. Barr, R. Levy, C. Scheepers, H. J. Tily, Random effects structure for confirmatory

886           hypothesis testing: Keep it maximal. *J Mem Lang* **68**, 255-278 (2013).

**Acknowledgements**

First, we would like to thank each of our participants for their time. We would also like to thank

doctors Nicholas O'Donnelly, Lisa Knight, Hilary Archer, Elizabeth Mallam, David Evans and

Tom Dalton for providing clinical cover for this study, and doctors Claire Rice and Catherine

Pennington for their time and guidance as the steering committee. We are grateful for Alex

Howat, Nerea Irigoras Izagirre, and Will Mears for their substantial help in collecting the data.

We would like to thank Phil Boreham, cardiologist, for advising us on the safe use of

domperidone. We would also like to thank Join Dementia Research, Bristol Brain Centre, North

Bristol NHS trust, and CRICBristol staff, and our funders; the Medical Research Council,

BRACE Bristol and the David Telling Trust.

**Author contributions**

HKI and EJC designed Study 1. MWJ, CD, CO, and LM contributed significantly to designing

Study 1. Randomisation and blinding were performed by LM for Study 1. HKI, JPG and EJC

designed Study 2. HKI and JPG developed the verbal memory tasks. HKI, GA, JPG, WJC and

UB wrote all analysis scripts. Sleep scoring was performed by WJC and OR and overseen by

HKI, spindle and slow oscillation analyses were carried out by HKI, WJC and UB. CO and UB

gave further statistical guidance. All data collection was overseen by HKI, JPG and EJC. Data

was collected by HKI, WJC, GA, OR, JS, RW, EF, JM, CD, ARW, CMN, and JPG. Further

clinical cover was provided by EJC, JM and JS. HKI, EJC, WJC, JPG, CO, MWJ and UB

interpreted the data. HKI and EJC wrote the manuscript, all authors contributed to the editing

of the manuscript and approved of the final version.

**Competing interests statement:**

913 The authors have no competing interests.

**Data availability:**

Contact the corresponding authors for copies of the MATLAB and R scripts used in analysis,

the experimental standard operating procedures, MATLAB scripts for the verbal memory tasks,

or word list. Data will be shared in line with sponsor's requirement for availability of

anonymised datasets from clinical trials. The data for the control study can also be shared upon

request.

**Code availability:**

Contact the corresponding authors for copies of the MATLAB and R scripts used in analysis.