Transmission and control strategies for porcine reproductive and respiratory

syndrome virus

Jason A. Galvis[1]; Joaquin M. Prada[2], Chris M. Jones[3], Gustavo Machado[1*]

[1] Department of Population Health and Pathobiology, College of Veterinary Medicine,

North Carolina State University, Raleigh, NC, USA·

[2] School of Veterinary Medicine, Faculty of Health and Medical Sciences, University of

Surrey, Guildford, UK

[3]Center for Geospatial Analytics, North Carolina State University, Raleigh, NC, USA,

Corresponding author: Dr. Gustavo Machado, gmachad@ncsu.edu

Summary

In the monitoring of porcine reproductive and respiratory syndrome virus (PRRSv),

knowledge about between-farm transmission dynamics is still lacking. Our objective was

to assess the relative contribution of between-farm PRRSv transmission routes through a

mechanistic epidemiological model calibrated with PRRSv occurrence, identify risk

areas, and estimate the impact of immunization strategies in the disease spread. We

developed a mathematical model of PRRSv transmission accounting for spatial

proximity, pig movements, and re-breaks in sow farms, parametrized on data collected

routinely by commercial pig farms. We then used the model to simulate the weekly

frequency of cases and built risk maps, and compared with the observed cases. We

simulated the implementation of two immunization strategies (preventive and reactive) to

mitigate the between-farm transmission. Our results indicated for sow and GDU farms'

local spread on average was above 60%, while for nurseries between-farm movements

represented 83% of transmissions and in finisher farms it was distributed almost 50%

local and 50% between-farm movement, the model allowed reproduce the weekly

frequency of observed cases and the risk maps built allowed the identification of

observed cases in the space. The increase in vaccine efficacy was the most important

parameter to mitigate between-farm transmission. Also, the implementation of

immunization by a preventive and reactive strategy combined had the better result to

mitigate between-farm transmission than implement these strategies individually. These

immunization strategies had a better performance with the use of rigorous protocols, such

as immunization as soon as the farm receives a shipment or increase the detection

coverage in all production types, as well as reduce time of disease detection. This model

can be used by the pork producers to improve surveillance activities, enhancing the

capacities to early detect circulation not only of endemic disease but also foreign animal

diseases (FADs).

KEYWORDS: swine disease spread, transmission dynamics, disease surveillance,

mechanistic modelling.

Introduction

Porcine reproductive and respiratory syndrome virus (PRRSv) is one of the most

relevant endemic swine diseases in North America causing major economic loss by

reproductive failures and reduction of average daily weight (Holtkamp et al., 2013). In

recent years the surveillance and control of PRRSv has increased in order to fight the

disease (Corzo et al., 2010; Perez et al., 2015, 2019; Emanuela Pileri & Mateu, 2016).

Despite the great advances, PRRSv has proven to be difficult to contain mainly because

of the continuous between-farms transmission events (Emanuela Pileri & Mateu, 2016).

As a consequence, new outbreaks have been reported each year, with epidemics peaks in

the colder seasons (Arruda, Vilalta, Puig, Perez, & Alba, 2018; Hermann et al., 2007;

Sanhueza, Vilalta, Corzo, & Arruda, 2019).

In the last two decades, several epidemiological studies have tried to explain the

between-farm transmission dynamics of PRRSv (Dee, Deen, Burns, Douthit, & Pijoan,

2004; Dee et al., 2003b; Rosendal et al., 2014; Thakur, Revie, Hurnik, Poljak, &

Sanchez, 2015). The contact among farms by pig movements and the proximity distance

between positive and negative farms have been described as the main route for PRRSv

spread (K. Lee et al., 2017; Silva, Machado, Baker, Holtkamp, & Linhares, 2019;

VanderWaal, Perez, Torremorrell, Morrison, & Craft, 2018; Velasova, Alarcon,

Williamson, & Wieland, 2012). Similarly, other studies pointed out that sow farms can

re-breaks due to PRRSv recirculation mainly by infected and latent sows, environmental

contamination and continuous birth of susceptible piglets (Arruda, Friendship, Carpenter,

Greer, & Poljak, 2016; Bierk et al., 2001). Despite these findings, the understanding and

quantification of PRRSv transmission continues to be a challenge, which has important

consequences in timely control of virus circulation and has limited our knowledge about

the impact of control measures over the yearly wave of PRRSv cases.

The construction of mechanistic disease spread models have provided a powerful

tool to understand disease transmission dynamics between populations (Dubé, Ribble,

Kelton, & McNab, 2009; Siettos & Russo, 2013). More recently studies used

mathematical models to investigate how COVID-19 spread between populations and had

shown to be useful to monitor the epidemic dynamics, evaluating number asymptomatic

populations, risk regions, temporal trend and impact of control strategies (Mandal et al.,

2020; Park, Cornforth, Dushoff, & Weitz, 2020; Shim, Tariq, Choi, Lee, & Chowell,

2020; Yang & Wang, 2020), information indispensable to organize effective and efficient

control plans and calculate the resources necessaries along the epidemic. Similarly, the

use of this mechanistic model in swine diseases can provide the same information, thus it

possible to increase our knowledge of the disease dynamic to face endemic disease, as

PRRSv, or be used to evaluate epidemic and control scenarios for the introduction of

exotic diseases, such a potential African Swine Fever introduction in the U.S. (Brown &

Bevins, 2018; Jurado et al., 2019). In this study, we show that a mechanistic spread

model can generate valuable information to understand PRRSv transmission dynamics

and evaluate control strategies to avoid the continuous between-farm spread. Our

objective was to assess the relative contribution of between-farm PRRSv transmission

routes through a mechanistic epidemiological model calibrated with PRRSv occurrence,

identify risk areas, and estimate the impact of immunization strategies in the disease

spread.

Methodology

We developed a discrete-time stochastic model to simulate between-farm PRRSv

transmission. The model included two main transmission routes, between-farm pig

movements and local transmission as a function of the distance between susceptible and

infected farms, named hereafter simply as local transmission. Additionally, we included

the probability of the virus to survive inside the farm causing re-break at sow farms

(Emanuela Pileri & Mateu, 2016). In the following sections we present i) a description of

the susceptible pig population, the between-farm pig movements and reported cases of

PRRSv, ii) a description of the mathematical model, projection of risk areas with future

outbreaks and how immunization strategies affect the between-farm transmission.

Data Sources

We collected PRRSv occurrences of 2,294 farms from three not commercially-

related pig production systems (here coded as A, B, and C), from a US region. Data of

each farm included national premises identification number, production types (sow,

nursery, finisher (which included wean-to-finish and finisher farms) and gilt development

unit (GDU), pig capacities, and geographic coordinates.

The record of pig movements between farms was used to reconstruct directed

weekly contact networks from June 01, 2019 to December 05, 2019. Each pig movement

included movement date, farm of origin and destination, number of pigs transported and

purpose of movement (e.g., weaning, market). Movement data with missing either farm

location, production type, farm of origin or destination were excluded prior to being

analyzed. Thus, the movement between 92% of the farms in the study region comprised a

total of 73,361 between-farm movements, in the past six months.

The weekly number of PRRSv cases from January 05, 2009 to December 05,

2019 were compiled from two sources i) for cases in sow farms we used the "PRRSv

hotlist" which is used and maintained by each production system to keep track of current

and new introductions, and ii) for downstream farms, nurseries, GDU, and finishers, we

used ORF-5 PRRSv sequences data, which is maintained by the swine veterinarians to

monitor PRRSv strain circulation. Briefly, a new sequence is often generated when

clinical signs are identified or there is a reason to request one (Jara, Rasmussen, Corzo, &

Machado, 2004). Thus, for sow farms we classified each PRRSv occurrence as "new" or

"recurrent case" based on the reported dates. Infection that occured within the past 41

weeks from a previous occurrence in the same sow farm was considered a recurrent case,

otherwise it was classified as a new case. We based this classification on the average

national time to breeding farms to reach PRRSv stable status, see (Sanhueza et al., 2019)

for more details. Similarly, for nurseries, GDUs and finishers farms, we used pig

production phase times to define what was a new PRRSv case, where a sequence of

PRRSv within 7, 22 and 25 weeks, respectively, was counted towards a recurrent case.

The rationale for the intervals used was based on the production characteristics of each

farm type, since these production sites are all-in and all-out continuous animal flow,

indeed all animals are introduced and removed from a farm within a short period of time.

Between-farm infection dynamics and transmission routes

The between-farm transmission was modeled through a discrete-time stochastic

model, using three farm infection status: susceptible (S), undetected infection (U) and

detected infection (D) at each time step (Figure 1A). The chosen one week time step

matches the frequency at which cases are reported. In this model, susceptible farms were

allowed to become infected, with the disease being initially undetected, through two main

between-farm transmission routes: infected incoming pigs from infected farms (both

undetected and detected) (VanderWaal et al., 2018), and local transmission from nearby

infected farms (Silva et al., 2019; Velasova et al., 2012). To model the local transmission

force of infection, we used a gravity model, while also accounting for the vegetation

surrounding each farm, as a possible barrier to transmission (Arruda, Vilalta, Perez, &

Morrison, 2017). Additionally, previous studies have shown that PRRSv can persist

undetected within recently infected farms, mainly in sow farms, contaminated

environments or fomites (Emanuela Pileri & Mateu, 2016). We thus included in the

model a probability of sow farms to re-break with PRRSv within two-year since their last

infection (Perez et al., 2019). The transmission rates from the between-farm pig

movements, local transmission and the re-break probability were represented by the beta

values βnet, $\beta local$ and $\beta rebreak$ and were fitted to the observed number of PRRSv

cases (see model fitting section) (Figure 1B).

Figure 1.

Between-farm pig movements

We used the weekly pig movement to identify contacts between susceptible farms

i and both infected detected and undetected farms j, which were used to estimate

infectious force by pig movements in the network (λnet):

$$\lambda net_{it} = \beta net * N_{it}; \ \forall = 1, \dots, n \ suceptible \ farms, \qquad (1)$$

163   N_i is the number of incoming contacts received from infected farms (undetected and

164   detected) at the time step t *(one week)*.

166   ***Local transmission***

167   The local transmission was modeled through a gravity model as follow:

$$a_{ij} = \frac{m_i m_j}{d_{ij}^2} \ , \qquad (2)$$

where the attraction force between the farms i and j was directly proportional to the

farms' pig capacity m_i and m_j divided by the squared Euclidean distance d_{ij}^2 between the

two farms. In addition, based on some preliminary simulations, we set the maximum

range of local transmission to 35km ($a_{ij} = 0$ when $d_{ij} > 35$km). Local transmission was

modulated by the enhanced vegetation index (EVI), which was used to quantify

vegetation greenness around each farm, here representing a potential barrier that reduced

local transmission of PRRSv (Arruda et al., 2017). Thus, susceptible farms at a high EVI

location were less likely to be infected via local transmission from nearby infected farms,

decreasing the attraction force between m_i and m_j such that,

$$g_{ij} = a_{ijt} * (1 - e_{it}) \; ; \; \forall_i = 1, \ldots, n \; suceptible \; farms, \tag{3}$$

177   where g_{ij} is the attraction force including the barrier effect, in which e_i is the barrier index

178   from farm i represented on a [0,1] scale (see SM for additional information). Thus, the

179   local infectious force ($\lambda local$) was defined as

$$\lambda local_{it} = \beta local * \sum_{j=1}^{n} g_{ijt} \; ; \; \forall_i = 1, \ldots, n \; suceptible \; farms \; , \tag{4}$$

$$; \forall_j = 1, \ldots, n \; infected \; farms$$

181   ***Sow farm re-break***

182   Exclusively for sow farms, a re-break probability which is computed with the

183   other two forces of infection was included as

$$\lambda rebreak_{it} = \beta rebreak \; ; \; \forall_i = 1, \ldots, n \; suceptible \; sow \tag{5}$$

$$farms \; with \; historical \; PRRSv \; records \; from \; June \; 2017 \; to \; June \; 2019,$$

185   **The transmission dynamics**

186   The total force of infection (λ) was calculated as the sum of all infectious forces

187   accounting for the monthly PRRSv seasonality,

$$\lambda_{it} = (\lambda net_{it} + \lambda local_{it} + \lambda rebreak_{it}) * T_{tm}, \tag{6}$$

where λ_{it} is the total infectious force from a susceptible farm i and T_{tm} is the seasonality

index from the time step t in month m. The monthly seasonality represents periods of

time where the PRRSv transmission is more or less frequent due to pigs' immunity and a

coordinated sequence of events related to the weather that may influence the total

transmission (Dee et al., 2003b, 2003a). Thus, the seasonality index values were

estimated from the moving averages of the monthly frequency of PRRSv cases between

2015 and 2019 (SM Figure 1), which were then scaled between 0.001 and 1. We also

include a term that accounts for the farms' ability to remain PRRSv negative (H), which

modulates the λ_{it}

$$\lambda_{it} = (\ (\lambda net_{it} + \ \lambda local_{it} + \ \lambda rebreak_{it}) * T_{tm} \ ) * (1 - H_{it} \ ), \qquad (7)$$

here H takes values between 0 and 1 and represents a reduction in the total force of

infection received by the susceptible farms with lower risk of infection ($H>0$). Thus, we

assigned H values greater than 0 only for sow farms without PRRSv cases from July 1,

2009 until June 1, 2019, which were denominated farms with low risk of infection as an

initial condition setting in the model. Furthermore, H value was fitted to the observed

number of PRRSv cases (see model fitting). Finally, we calculate the transmission

probability (Y_{it}):

$$Y_{it} = 1 - e^{-\lambda_{it}}, \qquad (8)$$

204   where the transition probability of a susceptible farm i to infected and undetected status at

205   time point t is given by Y_{it}.

207   The detection rate for an infected farm (transition from undetected to detected

208   infection status) is calculated by a detection probability derived from a logistic function

$$f(x) = \frac{L_p}{1 - e^{-k(x-x0)}} \tag{9}$$

where L is the maximum detection probability for PRRSv cases and its value was defined

accordingly to the farm production types p. Thus, for sow farms which undergo

periodically testing, we assumed L to be 0.95, while in nurseries, finishers and GDU

farm, PRRSv reported cases are frequently underestimated due to the lack of clinical

signs and samples are rarely collected, as a consequence the probability of detecting cases

in these farms is lower compared with sow farms. Therefore, to fit L in each production

type based on observed PRRSv cases and an empirical PRRSv prevalence (see model

fitting section for more details). Here, $X0$ is the sigmoid's midpoint to detect an infected

undetected farm (i.e. which was assumed to be the average time for PRRSv detection),

defined in 4 weeks, k is the logistic growth rate, which we assumed to be 0.5 to increase

the detection probability after each week and X were the weeks post infections. These

values were defined after consulting with veterinarians of the studied production systems,

which was also set to 4 weeks to detect an infected undetected farm. The assumption was

in agreement with the only publication to authors' knowledge, done in Chile where it was

found the virus could have been circulating for at least one month before it was detected

(Neira et al., 2017).

In the model we assumed the course of infection according to the farm production

type. Thus, an infected undetected sow farm stayed indefinitely infected until it was

detected, after detected the farm stayed infected by a predefined time calculated from a

poisson distribution using 41 weeks (average time to stability) as the mean of the

distribution (Sanhueza et al., 2019), then the farm transited to susceptible status after its

predefined infectious time elapsed in the simulation. On the other hand, due to the all-in

and all-out animal flow in nursery, GDU and finisher farms, these return to susceptible

status at the end of its production cycle, which means when all its pigs were moved out of

its premises.

Initial conditions settings

The initial farms assumed to be infected in the starting point of the simulation

where all farms previously infected within a number of days before June first 2019

accordingly farm types as follows: sow 41, nursery 7, GDU 22, finisher 25 weeks. These

time intervals correspond to time to stability in sow farms and cycle production time for

the other production types, respectively (Holtkamp et al., 2011).

Model fitting

We used an Approximate Bayesian Computation (ABC) method based on a

modified Sequential Monte Carlo (SMC) rejection algorithm, which allows us to recover

the approximate posterior distribution of the five unknown model parameters Bnet,

Blocal, Breinf, H, and L. A set of draws of these five parameters is denominated a

"particle" (Beaumont, 2010), and the value was fitted to the observed PRRSv cases for

each production type from June 1 until December 5, 2019. For nursery and finisher farms

an empirical expected weekly number of cases, derived from an annual prevalence of

30% (production system veterinarians, personal communications), was used. However,

considering the yearly PRRSv seasonality (Arruda et al., 2018; Jara et al., 2004), we

divided that empirical prevalence by week according to the seasonal index T calculated

previously and by the average number of weeks by month. For example, the expected

prevalence for the weeks in December was calculated as $(0.3\ /\ (T_{December}\ /$

$\sum_{m=1}^{n=12} T_m\ ))\ /\ 4.34\ ; \forall_m = January, \dots, December$.

The sequential algorithm had a two-step rejection procedure. In the first step, the

model was fitted to the weekly frequency of cases using eleven summary statistics, see

Table 1. In the second step, the sensitivity and specificity of the model to predict true risk

areas was evaluated through the comparison between simulated and observed infected

farm locations within cells of 10km by 10km (see SM for the sensitivity and specificity

calculation). The ABC rejection procedure was ran until the generation of 100 desired

particles (N), with a tolerance interval (ϵ) for all metrics as shown in Table 1. Finally, the

model was performed using the parameters listed in Table 2.

Model outputs

The model outputs generated were i) the weekly PRRSv number of infected

farms, both undetected and detected, ii) the estimated force of infection for each

transmission route and farm production type, iii) forecasting risk areas and comparison of

simulated and observed infected locations.

To assess the relative contribution of each transmission route, namely between-

farm, local and re-breaks in sow farms, we generated 1000 model runs with each

transmission route simulated one at a time. For example, between-farm transmission can

be calculated as the total number of farms infected in the runs with only between-farm

transmission divided by the total number of farms infected summed across the three

transmission routes.

To evaluate the efficacy of the disease spread model to predict the location of

future PRRSv cases, the 319 sow farms from our study region were grouped into 154

cells of 10km by 10km, in which the number of sow farms by cell was 1 in median, 3 in

the percentile 75% and a maximum of 12 farms. We calculated a risk of PRRSv detection

by cell into one week windows based on the output of the fitted model and the inclusion

of previous reported cases to avoid their influence in the forecasting. Thus, we calculated

the risk of PRRSv detection and new introductions for sow farms between October 30,

2019 and November 06, 2019 (week with the higher number of cases reported, 3 cases)

and identified the cell with a risk about the percentile 90 (arbitrarily chosen) as higher

risk.

Control strategies

We simulated two possible control strategies that could be implemented, both of

them relate to herd level immunization and were aimed at reducing between-farm

transmission (Moura et al., 2019). In the first control scenario simulated, referred to as

the preventive strategy, the nursery, GDU and finisher farms administer massive

vaccination the same week after receiving pig placement. In the second scenario

simulated, hereafter named reactive strategy, all farm types administer massive

vaccination the same week after detecting PRRSv. Finally, we considered a third scenario

combining the preventive and reactive strategy. Here we assumed twelve different

vaccine efficacies (V): 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 30%, 50%, 80% and 90%,

implemented as a reduction in the overall force of infection. Also, here we assumed V

followed a logistic growth curve and reached its maximum efficacy after 4 week of

implementation (Kick et al., 2019). Because information about the length of the herd

immunity levels post vaccination continues to be a gap in knowledge, in our formulated

scenarios vaccinated pigs maintain protection for 36 weeks, which was based on (Arruda

et al., 2016; Evans, Medley, Creasey, & Green, 2010) and an expert opinion study. Thus,

for nursery, GDU and finisher farms, herd immunity protected pigs throughout the whole

production phase. A brief explanation of the V to reduce the force of infection is

available in SM. Each intervention effectiveness was evaluated through the accumulated

number of infected farms by production type over the period simulated and compared

with the baseline model (V=0). In addition, we evaluated the influence of some

complementary protocols to reduce the average number of cases. Thus, for the preventive

strategy, we included a delay of 1, 2 and 3 weeks to vaccinate after receiving a pig

placement and for the reactive strategy, we included a delay of 1, 2 and 3 weeks to

vaccinate after detecting PRRSv, also a lower time for viral detection of 1, 2 and 3 week

and stronger detection surveillance of 0.5 and 0.95 probability in nursery, GDU and

finisher farms (i.e., reinforce sampling strategies). All these protocols were evaluated

through a multivariate linear regression for each control scenario tested.

Results

In table 3 we summarized the farms susceptible and the total number of infected

farms by production types. Sow farms showed more cases of PRRSv as expected, while

finisher and nursery farms together had around half of the sow cases. GDU farms were

less infected and no cases were reported in isolation and boar stud farms. These results

need to be interpreted with caution since the volume of cases are directly related to the

surveillance effort, in sow farms it often involves weekly testing at weaning or at

processing, in which viral detection may vary because of the test options available

(Henao-Diaz, Ji, Giménez-Lirola, Baum, & Zimmerman, 2020). To fit the model, 48

cases were identified from June 01, 2019 to December 05, 2019, being those reported in

sow, nursery and finisher farms.

The simulations indicated that on average 61% of the sow farms cases were due to

local transmission, while 33% and 6% were due to pig movements and re-break,

respectively (Figure 2). For nursery farms, 80% of cases were due to pig movements and

20% local transmission, while for finisher movements and local transmission had almost

the same relevance with 47% and 53% of cases, respectively. Regarding GDU farms,

local transmission was responsible for 99% of the cases.

Figure 2.

The estimated average number of new cases after 26 weeks thought the simulation

was 19 (IQR 16-21) for sow farms, 57 (IQR 52-63) for nursery farms, 9 (IQR 8-11) for

GDU farms and 100 (IQR 90-110) for finisher farms. The new detected cases estimated

by the model followed the same temporal pattern of the observed cases, being the most of

these observed cases equal to the median of the simulations or falling into the

interquartile range (IQR) calculated (Figure 3A). Regarding the new undetected cases

estimated by the model, nursery and finisher farms were the production types with a

higher weekly frequency (Figure 3B), principally in the last two months (November and

December), with a maximum of 13 and 28 cases per week, respectively. Contrary, sow

and GDU farms had a lower frequency of new undetected cases, with a maximum of 7

and 5 cases per week, respectively.

Figure 3.

In the spatial forecast of the observed cases in sow farms, the model had an

average maximum sensitivity of 62% (IQR 56%-83%) when a subset of cells with risk

values above percentile 40 was used as a threshold to identify infected observed cells

(SM Figure X). For example, the forecasted week had three sow farms with positive

PRRSv records located in three different cells (Figure 4A), our model calculated the risk

of disease detection for each cell (Figure 4B) and identified the 16 cells with risk above

percentile 90 (16 from 154 cells) (Figure 4C). As a result, two of the cells with observed

cases were also cells with risk about percentile 90%. Thus, the model was able to detect

two of the three reported cases. We used the model to calculate the risk for newly

infected sow farms for the simulated week (Figure 4D) and showed the cells with risk

above 90% (Figure 4E).

Figure 4.

Control measures

In general, the three scenarios evaluated had a significant lower average cases

starting from a vaccine efficacy (*ve*) of 1% (p > 0.05), except by sow and GDU farms in

the preventive scenario which needed a *ve* of 2% and 3%, respectively (Table 3 and SM

Table x and x). Independent of the statistical differences found, the combined scenario

had a lower average cases than using the preventive or reactive strategies individually,

that was observed along all the *ve* tested and for all the production types (Figure 5). For

example, the average total cases along the 26 weeks simulated without control was 185

new cases, the simulations in the combined scenario with *ve* = 5% had a total of 107 new

cases (42% less cases compared with no control scenario), while the same *ve* in the

preventive scenario had 136 new cases (26% less cases) and the reactive had 139 new

cases (25% less cases). The preventive and reactive scenario had similar results in

nursery farms along all *ve* tested, but had differences in finisher farms due to lower

average cases in the preventive scenario and in GDU and sow farms due to lower average

cases in the reactive scenario (Figure 5). Furthermore, it is important to note that the

combined and reactive scenario had a considerably lower average cases than the

preventive scenario in sow and GDU farms, that difference was easier to observe after a

$ve \geq 5\%$.

In table 3 was presented a multivariate linear regression from the combined

scenario in function of the *ve* and other complementary control protocols. As expected,

immunization was the most important variable to prevent cases as increased *ve* values

(Table 3). The protocol with higher importance was a shorter time to immunize the farm

after receiving a pig shipment, however this variable had a high influence only in finisher

farms with 2.2 cases prevented when the farm was immunized in the same week after

receiving a pig shipment, while in the other production types prevented lees than 1 case.

The remaining protocols for this scenario had a low influence to prevent cases in all

production types (<0.6 cases prevented). On the other hand, the contribution of these

protocols was different when the preventive and reactive strategies were evaluated

individually (SM Table 1 and 2). For both strategies the contribution of *ve* was lower

than the combined scenario, but still had the highest influence to prevent cases. However,

the influence of the protocols to prevent cases was higher than in the combined scenario,

being this difference more evident in finisher and nursery farms. For example, in the

preventive strategy, finisher and nursery farms had 5.4 and 1.6 cases less if the farms

were immunized in the same week after receiving a pig shipment. While for the reactive

strategy, increasing the probability detection of PRRSv to 0.95 in all farm types was the

most important protocol with 4.3 and 1.6 cases less in finisher and nursery farms,

respectively. The other protocols combined in the reactive strategy prevented 3.6 cases in

finisher farms and 1.6 cases less in nursery farms if the farms were detected and

immunized in the same week of virus introduction.

Figure 5.

Discussion

In this paper, we addressed two important points for the control of PRRSv in the

United States, i) we built a disease spread model able to reproduce and quantify between-

farm transmission dynamics. Our results indicated for sow and GDU farms' local spread

on average was above 60%, while for nurseries between-farm movements represented

83% of transmissions and finisher farms it was distributed almost 50% local and 50%

between-farm movement. ii) We evaluated the effect of PRRSv surveillance and control

strategies in reducing the number of new infected farms. The increase in vaccine efficacy

was the most important parameter to mitigate between-farm transmission. Also, the

implementation of immunization by a preventive and reactive strategy combined had the

better result to mitigate between-farm transmission than implement these strategies

individually. These immunization strategies had a better performance with the use of

rigorous protocols, such as immunize as soon as the farm receives a shipment or increase

the detection coverage in all production types, as well as reduce time of disease detection.

For this last protocol, our model is able to provide information about the probable

location of new cases by the construction of risk maps that could be used for early disease

detection.

PRRSv spread model

In our current model, the transmission, surveillance and diagnosis parameters

were fitted through the observed cases, previous scientific literature and the expert

opinion of local veterinarians about PRRSv, thus we tried to represent a close

approximation of what happens in field conditions. We highlight that transmission,

surveillance and diagnosis continue to be a stochastic process, while disease was not

randomly distributed in space. Our model differs from previous studies modeling PRRSv

due most of them focused on model within-farm transmission (Arruda et al., 2016;

Bitsouni, Lycett, Opriessnig, & Doeschl-Wilson, 2019; Evans et al., 2010; Phoo-ngurn,

Kiataramkul, & Chamchod, 2019), while other studies modeled between-farm

transmission, but without the inclusion of empirical farm' location or pig movements

between farms to spread the virus (H. S. Lee et al., 2019; Thakur, Sanchez, et al., 2015).

Although, there was a study which used empirical pig movements, this did not include

the actual spatial location of the farms (Thakur, Revie, et al., 2015). Also, in the

knowledge of the author at the moment to write this manuscript, there was no studies

with a model calibrated with historical records of PRRSv cases in pig farms. Thus, our

study may provide a more robust result about between-farm transmission of PRRSv.

Our findings indicating a high contribution of local transmission in sow (63%),

GDU (99%) and finisher (52%) farms was a controversial result because even proximity

distances between farms have been indicated as a risk factor associated with PRRSv

(Silva et al., 2019; Velasova et al., 2012), most of the previous studies indicated

movements as the main transmission route (Amirpour Haredasht et al., 2017; K. Lee et

al., 2017; VanderWaal, Paploski, Makau, & Corzo, 2020). Nevertheless, in our study pig

movements also played an important role to infect nursery (88%) farms and had

approximately half of the contribution to infect finishers farms (48%). Thus, the

transmission due pig movements was still relevant to spread the disease and both

transmission routes may be conditioned to the farms' production type. Despite these

results, it is necessary to consider that the model used in this study was calibrated trying

to identify the areas where PRRSv cases in sow farms were reported. Thus, the rate of the

movements, local transmission or re-break increased when any of these routes was able to

explain to a greater degree the spatial distribution of these observed cases. As a result, the

cases not explained by movements or re-break were explained by a greater contribution

of the local transmission. Also, the local transmission in our model may be filling the

lack of other potential transmission routes not included here, such contaminated vehicles,

vectors and visitors (Dee et al., 2003a; Pitkin, Deen, & Dee, 2009; Pitkin, Deen, Otake,

Moon, & Dee, 2009) and their future inclusion may change the contribution of the

transmission routes found in this study.

PRRSv surveillance and control

At the moment, studies evaluating farm level immunity against PRRSv are not

available, being the most of the studies focused on animal level immunity and making the

comparison with our results difficult. Thus, future studies evaluating farm level immunity

will bring valuable information to compare with the model result. The studies evaluating

the efficacy of immunization through adapted modified-live vaccine (MLV) at animal

level have indicated that due the nature of the virus, characterized by rapid mutation rates

and extensive genetic divergence (Cho & Dee, 2006; Kappes & Faaberg, 2015), the

current vaccines have not been sufficient to provide complete protection against the

emerging and reemerging viral strains (Hou et al., 2020; Lunney et al., 2016). Thus, it is

possible to assume that the current vaccine efficacy at farm level is low. Despite that, it

has been proved that the use of vaccines reduces PRRSv shedding spread (Jeong, Park,

Kang, Park, & Chae, 2017; E. Pileri et al., 2017; Emanuela Pileri et al., 2015), and

consequently provides a lower probability of transmission, whereby combined with an

effective implementation strategy could reach better results to contain the disease spread

between farms.

486   Our results showed that the combination of the preventive and reactive strategy

had the best performance to reduce the number of new cases than using each of these

separately, however this can be associated with a higher vaccination coverage in the

farms (Bitsouni et al., 2019; Gandon, Mackinnon, Nee, & Read, 2003). On the other

hand, the use of these strategies individually could be also useful, but it is necessary to

consider their strengths and weaknesses. For example, the preventive strategy had an

advantage against the reactive strategy by immunizing the farms after receiving a pig

shipment, whereby infected farms could implement immunization before the disease was

detected and reduce its infection force to other farms. Thus, the preventive strategy was

aimed to avoid cases mainly caused by animal movements, as a result the farms which

had a strong contribution of the movements (nursery and finisher farms) had a lower

incidence as ve increased than the farms which had stronger contribution of the local

transmission (sow and GDU farms). On the contrary, the reactive strategy here

simulated aimed to immunize detected farms, whereby the route of transmission of the

virus to the farm was not as relevant as in the preventive strategy, instead the virus

detection in the farms played a more important role. In this study we assumed that sow

farms had a high probability of virus detection (L = 0.95), consequently were more often

detected and immunized than the other production types which had a lower probability (L

= 0.09). Therefore, in the reactive strategy most of the infected sow farms were

vaccinated and its infection force to other farms was reduced, this was highly relevant for

the control of PRRSv because sow farms act as super-spreaders in the network due

they can reach a high number of farms by their trade contact chain (Passafaro,

Fernandes, Valente, Williams, & Rosa, 2020). Our result suggests that the reactive

strategy had equal or better performance to reduce cases in nursery, sow and GDU farms,

and slightly worse in finisher farms when compared with the preventive strategy.

Therefore, the reactive strategy seems to be a better option to implement, however

additional studies are necessary to consider implementing the reactive instead of the

preventive strategy. Thus, using both strategies combined seems to be the better option at

the moment to control PRRSv between-farm transmission.

516   The results of the built model can provide weekly reports (maps) of where

samples have elevated probabilities of return positive, thus providing information

necessary to implement PRRSv control based on target surveillance. In the surveillance

of PRRSv in the United States, we found two studies trying to predict spatial risk, one

was associated with the Dr. Morrison's Swine Health Monitoring Program (MSHMP),

which has contributed to collect data and evaluate the characteristics of this disease in the

country, one its products was the construction of spatial risk maps of PRRSv outbreaks

using machine learning algorithms with environmental and pig population variables

(Vilalta et al., 2017), but without the inclusion of between-farm transmission variables

that are crucial to understand the disease spread dynamic. The other study used pig trade

network matrices and geographical distance to predict the farm's PRRSv status through a

spatio-temporal analysis using Bayesian models with INLA (Amirpour Haredasht et al.,

2017); the model had a relatively high accuracy (87%) to predict PRRSv farm status,

however used 6-month windows to evaluate the probability of new infection and that is a

limiting to use into real-time target surveillance. Other studies outside US tried to predict

PRRSv recirculation within sow farms using conditional poisson models with farm

production index associated with PRRSv occurrence with (Fraile et al., 2020), although

this methodology did not considered between-farm transmission dynamic, it is an

interesting methodology to include in futures step in our model to calibrate the re-break

probability in the sow farms. Thus, considering the previous studies developed to predict

PRRSv, our model can provide valuable information for target surveillance in real time.

Limitations and final remarks

One limitation of this study is related to how sequences are collected in

downstream farms. Since each production system may follow a different protocol while

deciding in sending a sample to be sequenced, and this can also vary among the

veterinarians of the same company, said that until routine surveillance is not implemented

in those farm types, this data may be the best approximation of what has been circulating

in finisher and GDU and nursery farms. Another limitation of this study is the fact that

we do not consider the influence of viral genetics, we assume that all sequences from the

downstream farms are all the same and can be transmitted between-farms at the same

probability. It was not the goal of this study to model individual PRRSv strains, but this

should be considered in future studies. In addition, we calibrated our model using only

the spatial location of sow farms, whereby it may brought a bias in the transmission rates

calculated with the ABC, and consequently in the contribution of the transmission routes,

thus the future inclusion of the location of cases from nursery, GDU and finisher farms

will provide a more robust result, however it will only be possible with a stronger

surveillance to detect and report the presence of PRRS in these farms. Regarding the

control, in our model we assumed that vaccinated animals acquire life-long immunity, in

the future the heterologous response should be investigated in more detail in regard to its

effect in reducing between-farm spread, also due most of the literature about vaccines is

available at animal-level, the inclusion of compartmental transmission model at animal-

level in our model can allow to incorporate specific characteristics of vaccines.

In summary, our model reveals the transmission dynamic of PRRSv in our study

area, identifying the contribution of movements and local transmission to infect

individually sow, nursery, GDU and finisher farms. Also, increasing the vaccine efficacy

was crucial to avoid the continuous disease spread and its implementation through a

preventive and reactive strategy combined could prevent outbreaks in all production

types. Thus, this study provided producers and swine veterinarians a better understanding

of the disease spread process and which, when and where intervention(s) will better

impact the disease transmission chain. Ultimately, our model can be used to improve

tactical surveillance activities, enhancing the capacities to early detect circulation not

only of endemic disease but also foreign animal diseases (FADs).

Acknowledgements

This study was funded by the Swine Health Information Center (SHIC) Project #19-211

& thank the MSHMP participants for sharing information.

Conflict of interest

All authors confirm that there are no conflicts of interest to declare

Ethical statement

The authors confirm that the ethical policies of the journal, as noted on the journal's

author guidelines page. Since this work did not involve animal sampling neither

questioner data collection there was no need for ethics permits.

Data Availability Statement

The data that support the findings of this study are available from the regional veterinary

office. Restrictions apply to the availability of these data, which were used under

confidentiality agreements.

References

Amirpour Haredasht, S., Polson, D., Main, R., Lee, K., Holtkamp, D., & Martínez-López,

B. (2017). Modeling the spatio-temporal dynamics of porcine reproductive &

respiratory syndrome cases at farm level using geographical distance and pig trade

network matrices. *BMC Veterinary Research*, *13*(1), 163.

https://doi.org/10.1186/s12917-017-1076-6

Arruda, A. G., Friendship, R., Carpenter, J., Greer, A., & Poljak, Z. (2016). Evaluation of

Control Strategies for Porcine Reproductive and Respiratory Syndrome (PRRS) in

Swine Breeding Herds Using a Discrete Event Agent-Based Model. *PLOS ONE*,

11(11), e0166596. https://doi.org/10.1371/journal.pone.0166596

Arruda, A. G., Vilalta, C., Perez, A., & Morrison, R. (2017). Land altitude, slope, and

coverage as risk factors for Porcine Reproductive and Respiratory Syndrome

(PRRS) outbreaks in the United States. *PLOS ONE*, *12*(4), e0172638.

https://doi.org/10.1371/journal.pone.0172638

Arruda, A. G., Vilalta, C., Puig, P., Perez, A., & Alba, A. (2018). Time-series analysis for

porcine reproductive and respiratory syndrome in the United States. *PLOS ONE*,

13(4), e0195282. https://doi.org/10.1371/journal.pone.0195282

Beaumont, M. A. (2010). Approximate Bayesian Computation in Evolution and Ecology.

Annual Review of Ecology, Evolution, and Systematics, *41*(1), 379–406.

https://doi.org/10.1146/annurev-ecolsys-102209-144621

Bierk, M. D., Dee, S. A., Rossow, K. D., Otake, S., Collins, J. E., & Molitor, T. W.

(2001). Transmission of porcine reproductive and respiratory syndrome virus from

persistently infected sows to contact controls. *Canadian Journal of Veterinary*

Research, *65*(4), 261–266.

Bitsouni, V., Lycett, S., Opriessnig, T., & Doeschl-Wilson, A. (2019). Predicting vaccine

effectiveness in livestock populations: A theoretical framework applied to PRRS

virus infections in pigs. *PLOS ONE*, *14*(8), e0220738.

https://doi.org/10.1371/journal.pone.0220738

Brown, V. R., & Bevins, S. N. (2018). A Review of African Swine Fever and the

Potential for Introduction into the United States and the Possibility of Subsequent

Establishment in Feral Swine and Native Ticks. *Frontiers in Veterinary Science*, *5*.

https://doi.org/10.3389/fvets.2018.00011

Cho, J. G., & Dee, S. A. (2006). Porcine reproductive and respiratory syndrome virus.

Theriogenology, *66*(3), 655–662.

https://doi.org/10.1016/j.theriogenology.2006.04.024

Corzo, C. A., Mondaca, E., Wayne, S., Torremorell, M., Dee, S., Davies, P., & Morrison,

R. B. (2010). Control and elimination of porcine reproductive and respiratory

syndrome virus. *Virus Research*, *154*(1–2), 185–192.

https://doi.org/10.1016/j.virusres.2010.08.016

Dee, S., Deen, J., Burns, D., Douthit, G., & Pijoan, C. (2004). An assessment of

sanitation protocols for commercial transport vehicles contaminated with porcine

reproductive and respiratory syndrome virus. *Canadian Journal of Veterinary*

Research, *68*(3), 208–214.

Dee, S., Deen, J., Rossow, K., Weise, C., Eliason, R., Otake, S., … Pijoan, C. (2003a).

Mechanical transmission of porcine reproductive and respiratory syndrome virus

throughout a coordinated sequence of events during cold weather. *Canadian Journal*

of Veterinary Research, *66*(1), 232–239.

Dee, S., Deen, J., Rossow, K., Weise, C., Eliason, R., Otake, S., … Pijoan, C. (2003b).

Mechanical transmission of porcine reproductive and respiratory syndrome virus

throughout a coordinated sequence of events during warm weather. (612), 12–19.

Dubé, C., Ribble, C., Kelton, D., & McNab, B. (2009). A Review of Network Analysis

Terminology and its Application to Foot-and-Mouth Disease Modelling and Policy

Development. *Transboundary and Emerging Diseases*, *56*(3), 73–85.

https://doi.org/10.1111/j.1865-1682.2008.01064.x

Evans, C. M., Medley, G. F., Creasey, S. J., & Green, L. E. (2010). A stochastic

mathematical model of the within-herd transmission dynamics of porcine

reproductive and respiratory syndrome virus (PRRSV): Fade-out and persistence.

Preventive Veterinary Medicine, *93*(4), 248–257.

https://doi.org/10.1016/j.prevetmed.2009.11.001

Fraile, L., Fernández, N., Pena, R. N., Balasch, S., Castellà, G., Puig, P., … Valls, J.

(2020). A probabilistic Poisson-based model to detect PRRSV recirculation using

sow production records. *Preventive Veterinary Medicine*, *177*, 104948.

https://doi.org/10.1016/j.prevetmed.2020.104948

Gandon, S., Mackinnon, M., Nee, S., & Read, A. (2003). Imperfect vaccination: Some

epidemiological and evolutionary consequences. *Proceedings of the Royal Society*

B: Biological Sciences, *270*(1520), 1129–1136.

https://doi.org/10.1098/rspb.2003.2370

Henao-Diaz, A., Ji, J., Giménez-Lirola, L., Baum, D. H., & Zimmerman, J. (2020).

Understanding and interpreting PRRSV diagnostics in the context of "disease

transition stages." *Research in Veterinary Science*, *131*, 173–176.

https://doi.org/10.1016/j.rvsc.2020.04.023

Hermann, J., Hoff, S., Muñoz-Zanzi, C., Yoon, K.-J., Roof, M., Burkhardt, A., &

Zimmerman, J. (2007). Effect of temperature and relative humidity on the stability

of infectious porcine reproductive and respiratory syndrome virus in aerosols.

Veterinary Research, *38*(1), 81–93. https://doi.org/10.1051/vetres:2006044

Holtkamp, D. J., Kliebenstein, J. B., Neumann, E. J., Zimmerman, J. J., Rotto, H. F.,

Yoder, T. K., … Haley, C. A. (2013). Assessment of the economic impact of porcine

reproductive and respiratory syndrome virus on United States pork producers.

Journal of Swine Health and Production, *21*(2), 72–84.

Holtkamp, D. J., Morrison, B., Classen, D. M., Becton, L., Henry, S., Rodibaugh, M. T.,

… Zimmerman, J. (2011). Terminology for Classifying Swine Herds by PRRS

Status. *Journal Of Swine Health And Production*, *19*(February), 44–56.

Hou, F.-H., Lee, W.-C., Liao, J.-W., Chien, M.-S., Kuo, C.-J., Chung, H.-P., & Chia, M.-

Y. (2020). Evaluation of a type 2 modified live porcine reproductive and respiratory

syndrome vaccine against heterologous challenge of a lineage 3 highly virulent

isolate in pigs. *PeerJ*, *8*, e8840. https://doi.org/10.7717/peerj.8840

Jara, M., Rasmussen, D. A., Corzo, C. A., & Machado, G. (2004). On reproductive and

respiratory syndrome virus dissemination across pig production systems in the

United States. *MedRxiv*, *43*(2), 241–271, 372–373.

Jeong, J., Park, C., Kang, I., Park, S. J., & Chae, C. (2017). Concurrent vaccination of

boars with type 1 and type 2 porcine reproductive and respiratory syndrome virus

(Prrsv) reduces seminal shedding of type 1 and type 2 prrsv. *Canadian Journal of*

Veterinary Research, *81*(2), 108–117.

Jurado, C., Mur, L., Pérez Aguirreburualde, M. S., Cadenas-Fernández, E., Martínez-López, B., Sánchez-Vizcaíno, J. M., & Perez, A. (2019). Risk of African swine fever virus introduction into the United States through smuggling of pork in air passenger luggage. *Scientific Reports*, *9*(1), 14423. https://doi.org/10.1038/s41598-019-50403-w

Kappes, M. A., & Faaberg, K. S. (2015). PRRSV structure, replication and recombination: Origin of phenotype and genotype diversity. *Virology*, *479–480*, 475–486. https://doi.org/10.1016/j.virol.2015.02.012

Kick, A., Amaral, A., Cortes, L., Fogle, J., Crisci, E., Almond, G., & Käser, T. (2019). The T-Cell Response to Type 2 Porcine Reproductive and Respiratory Syndrome Virus (PRRSV). *Viruses*, *11*(9), 796. https://doi.org/10.3390/v11090796

Lee, H. S., Thakur, K. K., Bui, V. N., Bui, A. N., Dang, M. Van, & Wieland, B. (2019). Simulation of control scenarios of porcine reproductive and respiratory syndrome in Nghe An Province in Vietnam. *Transboundary and Emerging Diseases*, *66*(6), 2279–2287. https://doi.org/10.1111/tbed.13278

Lee, K., Polson, D., Lowe, E., Main, R., Holtkamp, D., & Martínez-López, B. (2017). Unraveling the contact patterns and network structure of pig shipments in the United States and its association with porcine reproductive and respiratory syndrome virus (PRRSV) outbreaks. *Preventive Veterinary Medicine*, *138*, 113–123. https://doi.org/10.1016/j.prevetmed.2017.02.001

Lunney, J. K., Fang, Y., Ladinig, A., Chen, N., Li, Y., Rowland, B., & Renukaradhya, G. J. (2016). Porcine Reproductive and Respiratory Syndrome Virus (PRRSV):

Pathogenesis and Interaction with the Immune System. *Annual Review of Animal*

Biosciences, *4*(1), 129–154. https://doi.org/10.1146/annurev-animal-022114-111025

Mandal, S., Bhatnagar, T., Arinaminpathy, N., Agarwal, A., Chowdhury, A., Murhekar,

M., … Sarkar, S. (2020). Prudent public health intervention strategies to control the

coronavirus disease 2019 transmission in India: A mathematical model-based

approach. *Indian Journal of Medical Research*.

https://doi.org/10.4103/ijmr.IJMR_504_20

Moura, C. A. A., Johnson, C., Baker, S. R., Holtkamp, D. J., Wang, C., & Linhares, D. C.

L. (2019). Assessment of immediate production impact following attenuated PRRS

type 2 virus vaccination in swine breeding herds. *Porcine Health Management*, *5*(1),

2–7. https://doi.org/10.1186/s40813-019-0120-2

Neira, V., Brito, B., Mena, J., Culhane, M., Apel, M. I., Max, V., … Ortega, R. (2017).

Epidemiological investigations of the introduction of porcine reproductive and

respiratory syndrome virus in Chile, 2013-2015. *PLOS ONE*, *12*(7), e0181569.

https://doi.org/10.1371/journal.pone.0181569

Park, S. W., Cornforth, D. M., Dushoff, J., & Weitz, J. S. (2020). The time scale of

asymptomatic transmission affects estimates of epidemic potential in the COVID-19

outbreak. *Epidemics*, 100392. https://doi.org/10.1016/j.epidem.2020.100392

Passafaro, T. L., Fernandes, A. F. A., Valente, B. D., Williams, N. H., & Rosa, G. J. M.

(2020). Network analysis of swine movements in a multi-site pig production system

in Iowa, USA. *Preventive Veterinary Medicine*, *174*(October 2019), 104856.

https://doi.org/10.1016/j.prevetmed.2019.104856

Perez, A. M., Davies, P. R., Goodell, C. K., Holtkamp, D. J., Mondaca-Fernández, E.,

Poljak, Z., … Morrison, R. B. (2015). Lessons learned and knowledge gaps about

the epidemiology and control of porcine reproductive and respiratory syndrome

virus in North America. *Journal of the American Veterinary Medical Association*,

246(12), 1304–1317. https://doi.org/10.2460/javma.246.12.1304

Perez, A. M., Linhares, D. C. L., Arruda, A. G., Van Der Waal, K., Machado, G., Vilalta,

C., … Corzo, C. A. (2019). Individual or common good? Voluntary data sharing to

inform disease surveillance systems in food animals. *Frontiers in Veterinary*

Science, *6*(JUN), 1–7. https://doi.org/10.3389/fvets.2019.00194

Phoo-ngurn, P., Kiataramkul, C., & Chamchod, F. (2019). Modeling the spread of

porcine reproductive and respiratory syndrome virus (PRRSV) in a swine

population: transmission dynamics, immunity information, and optimal control

strategies. *Advances in Difference Equations*, *2019*(1), 432.

https://doi.org/10.1186/s13662-019-2351-6

Pileri, E., Gibert, E., Martín-Valls, G. E., Nofrarias, M., López-Soria, S., Martín, M., …

Mateu, E. (2017). Transmission of Porcine reproductive and respiratory syndrome

virus 1 to and from vaccinated pigs in a one-to-one model. *Veterinary Microbiology*,

201, 18–25. https://doi.org/10.1016/j.vetmic.2016.12.012

Pileri, Emanuela, Gibert, E., Soldevila, F., García-Saenz, A., Pujols, J., Diaz, I., …

Mateu, E. (2015). Vaccination with a genotype 1 modified live vaccine against

porcine reproductive and respiratory syndrome virus significantly reduces viremia,

viral shedding and transmission of the virus in a quasi-natural experimental model.

Veterinary Microbiology, *175*(1), 7–16.

https://doi.org/10.1016/j.vetmic.2014.11.007

Pileri, Emanuela, & Mateu, E. (2016). Review on the transmission porcine reproductive

and respiratory syndrome virus between pigs and farms and impact on vaccination.

Veterinary Research, *47*(1), 108. https://doi.org/10.1186/s13567-016-0391-4

Pitkin, A., Deen, J., & Dee, S. (2009). Further assessment of fomites and personnel as

vehicles for the mechanical transport and transmission of porcine reproductive and

respiratory syndrome virus. *Canadian Journal of Veterinary Research*, *73*(4), 298–

302.

Pitkin, A., Deen, J., Otake, S., Moon, R., & Dee, S. (2009). Further assessment of

houseflies (Musca domestica) as vectors for the mechanical transport and

transmission of porcine reproductive and respiratory syndrome virus under field

conditions. *Canadian Journal of Veterinary Research*, *73*(2), 91–96.

Rosendal, T., Dewey, C., Friendship, R., Wootton, S., Young, B., & Poljak, Z. (2014).

Spatial and temporal patterns of porcine reproductive and respiratory syndrome

virus (PRRSV) genotypes in Ontario, Canada, 2004-2007. *BMC Veterinary*

Research, *10*. https://doi.org/10.1186/1746-6148-10-83

Sanhueza, J. M., Vilalta, C., Corzo, C., & Arruda, A. G. (2019). Factors affecting Porcine

Reproductive and Respiratory Syndrome virus time-to-stability in breeding herds in

the Midwestern United States. *Transboundary and Emerging Diseases*, *66*(2), 823–

830. https://doi.org/10.1111/tbed.13091

Shim, E., Tariq, A., Choi, W., Lee, Y., & Chowell, G. (2020). Transmission potential and

severity of COVID-19 in South Korea. *International Journal of Infectious Diseases*,

93, 339–344. https://doi.org/10.1016/j.ijid.2020.03.031

Siettos, C. I., & Russo, L. (2013). Mathematical modeling of infectious disease dynamics.

Virulence, *4*(4), 295–306. https://doi.org/10.4161/viru.24041

Silva, G. S., Machado, G., Baker, K. L., Holtkamp, D. J., & Linhares, D. C. L. (2019).

Machine-learning algorithms to identify key biosecurity practices and factors

associated with breeding herds reporting PRRS outbreak. *Preventive Veterinary*

Medicine, *171*, 104749. https://doi.org/10.1016/j.prevetmed.2019.104749

Thakur, K. K., Revie, C. W., Hurnik, D., Poljak, Z., & Sanchez, J. (2015). Simulation of

between-farm transmission of porcine reproductive and respiratory syndrome virus

in Ontario, Canada using the North American Animal Disease Spread Model.

Preventive Veterinary Medicine, *118*(4), 413–426.

https://doi.org/10.1016/j.prevetmed.2015.01.006

Thakur, K. K., Sanchez, J., Hurnik, D., Poljak, Z., Opps, S., & Revie, C. W. (2015).

Development of a network based model to simulate the between-farm transmission

of the porcine reproductive and respiratory syndrome virus. *Veterinary*

Microbiology, *180*(3–4), 212–222. https://doi.org/10.1016/j.vetmic.2015.09.010

VanderWaal, K., Paploski, I. A. D., Makau, D. N., & Corzo, C. A. (2020). Contrasting

animal movement and spatial connectivity networks in shaping transmission

pathways of a genetically diverse virus. *Preventive Veterinary Medicine*, *178*,

104977. https://doi.org/10.1016/j.prevetmed.2020.104977

VanderWaal, K., Perez, A., Torremorrell, M., Morrison, R. M., & Craft, M. (2018). Role

of animal movement and indirect contact among farms in transmission of porcine

epidemic diarrhea virus. *Epidemics*, *24*(January), 67–75.

https://doi.org/10.1016/j.epidem.2018.04.001

Velasova, M., Alarcon, P., Williamson, S., & Wieland, B. (2012). Risk factors for

porcine reproductive and respiratory syndrome virus infection and resulting

challenges for effective disease surveillance. *BMC Veterinary Research*, *8*(1), 1.

https://doi.org/10.1186/1746-6148-8-184

Vilalta, C., Arruda, A. G., Tousignant, S. J. P., Valdes-Donoso, P., Muellner, P.,

Muellner, U., … Perez, A. M. (2017). A Review of Quantitative Tools Used to

Assess the Epidemiology of Porcine Reproductive and Respiratory Syndrome in

U.S. Swine Farms Using Dr. Morrison's Swine Health Monitoring Program Data.

Frontiers in Veterinary Science, *4*. https://doi.org/10.3389/fvets.2017.00094

Yang, C., & Wang, J. (2020). A mathematical model for the novel coronavirus epidemic

in Wuhan, China. *Mathematical Biosciences and Engineering*, *17*(3), 2708–2724.

https://doi.org/10.3934/mbe.2020148

Table 1. Summary statistics used in the Approximate Bayesian Computation Sequential

Monte Carlo rejection algorithm.

Summary Statistic	Observed Values	Tolerance interval
Total number of new cases detected in sow farms	23	10
Av. weekly number of new cases detected in sow farms	0.85	0.5
Max. number of new cases detected in a week in sow farms	3	5
Total number of new cases detected in nursery farms	17	10
Av. weekly number of new cases detected in nursery farms	0.62	0.5
Max. number of new cases detected in nursery farms	3	5
Total number of new cases detected in finisher farms	8	10
Av. weekly number of new cases detected in finisher farms	0.3	0.5
Max. number of new cases detected in a week in finisher farms	3	5
Total number of expected cases in nursery and finisher farms	155	100
Av. weekly number of expected cases in nursery and finisher farms	5.7	5
Max. number of expected cases in nursery and finisher farms	20	10

810   **Table 2.** Transmission parameters used in the disease spread model to simulate PRRSv

811   infection dynamics.

Model parameters	Symbol	Values	Reference
Transmission rate by movements	βnet	0.24	Median values from ABC fitting
Transmission rate by local	$\beta local$	0.003	Median values from ABC fitting
Transmission rate by re-break	$\beta rebreak$	0.0019	Median values from ABC fitting
Farms' ability remain PRRSv stable	H	Sow = 0.53 GDU = 0.27	Median values from ABC fitting
Effective surveillance	L	Sow = 0.95 Nursery, GDU and Finisher = 0.094	Median values from ABC fitting
PRRSv cases seasonality	T	Weekly values calculated	Moving averages
Enhanced vegetation index	EVI	Values by farm each week	https://lpdaac.usgs.gov
Gravity with barrier effect	g	Calculated in each time step	Based on EVI (SM)
Av. time for PRRSv detection	X	4 weeks	Expert opinion
Av. infectious time sow farms	-	41 weeks	(Sanhueza et al., 2019)

813   **Table 3.** Number of PRRSv cases by production types, for a time series from 2009-2019

814   and a subset of cases used in the study

Production Type	Susceptible farms	Total new cases 2009-2019	Subset new cases Jun-Dec 2019
Sow	319 (14%)	1070	23
Gilt development unit (GDU)	187 (8.2%)	50	0
Nursery	468 (20.4%)	648	17
Finisher	1272 (55.4%)	537	8
Isolation	33 (1.4%)	0	0
Boar Stud	15 (0.6%)	0	0

815
816

817 **Table 4.** Multivariate linear regression from the combined control scenario.

		Sow 19 (16-21)	Nursery 57 (52-63)	GDU 9 (8-11)	Finisher 100 (90-110)
Cases without control					
Vaccine efficacy [c]	1%	-1.4 (-1.4,-1.5)	-4.3 (-4.1,-4.4)	-1 (-0.9,-1)	-14 (-13.7,-14.2)
	2%	-2.7 (-2.6,-2.8)	-8.1 (-7.9,-8.2)	-1.8 (-1.7,-1.8)	-24.8 (-24.5,-25)
	3%	-3.8 (-3.7,-3.9)	-11 (-10.9,-11.2)	-2.6 (-2.5,-2.6)	-33.9 (-33.7,-34.2)
	4%	-4.7 (-4.6,-4.8)	-13.6 (-13.5,-13.8)	-3.3 (-3.2,-3.3)	-41.7 (-41.4,-41.9)
	5%	-5.5 (-5.5,-5.6)	-15.9 (-15.7,-16.1)	-3.9 (-3.9,-4)	-47.7 (-47.4,-47.9)
	10%	-8.6 (-8.5,-8.6)	-23.3 (-23.1,-23.4)	-6.4 (-6.3,-6.4)	-66.1 (-65.8,-66.3)
	15%	-10.1 (-10,-10.2)	-27.7 (-27.6,-27.9)	-7.6 (-7.6,-7.6)	-74.3 (-74,-74.5)
	20%	-10.9 (-10.8,-11)	-31.2 (-31.1,-31.4)	-8.2 (-8.2,-8.3)	-78.8 (-78.5,-79.1)
	30%	-11.9 (-11.8,-12)	-36.7 (-36.6,-36.9)	-8.8 (-8.8,-8.9)	-84.7 (-84.5,-85)
	50%	-13.3 (-13.3,-13.4)	-45.9 (-45.8,-46.1)	-9.2 (-9.2,-9.2)	-91.6 (-91.4,-91.9)
	80%	-15.5 (-15.4,-15.6)	-54.7 (-54.5,-54.8)	-9.3 (-9.3,-9.4)	-96.8 (-96.6,-97.1)
	90%	-16.4 (-16.3,-16.4)	-56.2 (-56.1,-56.4)	-9.4 (-9.3,-9.4)	-97.9 (-97.6,-98.1)
Delay vaccine after shipment [a]	2 w	0 (0,-0.1)	-0.1 (0,-0.2)	0 (0,0)	-0.5 (-0.4,-0.7)
	1 w	0 (0,-0.1)	-0.3 (-0.2,-0.4)	0 (0,0)	-1.4 (-1.2,-1.5)
	0 w	-0.1 (-0.1,-0.1)	-0.6 (-0.5,-0.7)	0 (0,-0.1)	-2.2 (-2.1,-2.3)
Average time dx [a]	2 w	0 (0,-0.1)	0 (0.1,-0.1)	0 (0,-0.1)	-0.1 (0.1,-0.2)
	1 w	0 (0,-0.1)	0 (0.1,-0.1)	0 (0,-0.1)	-0.1 (0,-0.3)
	0 w	-0.1 (0,-0.1)	-0.1 (0,-0.2)	-0.1 (0,-0.1)	-0.2 (-0.1,-0.4)
Delay vaccine after dx [a]	2 w	-0.1 (-0.1,-0.2)	-0.1 (0,-0.2)	-0.1 (-0.1,-0.1)	-0.3 (-0.1,-0.4)
	1 w	-0.2 (-0.2,-0.3)	-0.3 (-0.2,-0.4)	-0.2 (-0.2,-0.2)	-0.6 (-0.4,-0.7)
	0 w	-0.2 (-0.2,-0.3)	-0.3 (-0.2,-0.4)	-0.2 (-0.2,-0.2)	-0.6 (-0.5,-0.7)
Probability. dx [b]	0.5	0 (0,-0.1)	-0.1 (0,-0.2)	0 (0,0)	-0.1 (0,-0.3)
	0.95	0 (0,-0.1)	-0.1 (0,-0.2)	0 (0,0)	-0.1 (0,-0.2)

818 *p > 0.05. [a] Base comparison category 3 weeks. [b] Base comparison category 0.09.
819 [c] Base comparison category 0%. (dx = detection, w = week(s))
820

Figure Legends

Figure 1. Between-farm PRRSv transmission modelling framework. A) Flow between

infection stages; and B) The modeled transmission routes.

Fig. 2. Infection force by each transmission route and production type. Y-axis

represents the proportion of transmission estimated by each transmission route, while x-

axis to which week the model was projected to make these inferences.

Figure 3. The weekly number of reported and simulated cases PRRSv for sow, nursery,

gilt development units (GDU) and finisher farms. **A-top**) Number of observed (in red)

and simulated detected PRRSv cases (in grey) and **A-bottom**) the grey line represent the

median of the model and shade area the first and third quartile; **B-top**) Number of

simulated undetected PRRSv cases and **B-bottom**) the grey line represent the median of

the model and shade area the first and third quartile.

Figure 4. Spatial forecasting of PRRSv in sow farms for a week between October 30,

2019, to November 06, 2019. A) cells with observed sow farms infected; B) risk of

PRRSv detection ; C) subset of cells with risk of PRRSv detection above percentile 90;

D) risk of new PRRSv introduction; and E) subset of cells with risk of new PRRSv

introduction above percentile 90. *(cells with risk = 0 have transparency).

Figure 5. Accumulated number of infected farms along 26 week after the implementation

of control strategies. In the graph, the vaccine efficacies are grouped by color, points are

the average cases and their segments represent one standard deviation. For the preventive

strategy, farms were vaccinated in the same week after receive a movement, for the

reactive strategy farms were vaccinated in the same week after detect positive animals,

also the average time of detection was fixed in 4 weeks and the effective surveillance was

0.95 for sow and 0.09 for the other production types (baseline model transmission

parameters), for the combined strategy the previous parameters were used.