Modeling the transmission and vaccination strategy for porcine reproductive and

respiratory syndrome virus

Running title: The dynamics of PRRS virus spread

Jason A. Galvis[1], Joaquin M. Prada[2] , Cesar A. Corzo[3], Gustavo Machado[1]*

[1] Department of Population Health and Pathobiology, College of Veterinary Medicine, Raleigh,

North Carolina.

[2] School of Veterinary Medicine, Faculty of Health and Medical Sciences, University of Surrey,

Guildford, UK

[3] Veterinary Population Medicine Department, College of Veterinary Medicine, University of

Minnesota, St Paul, MN, USA

***Corresponding author: gmachad@ncsu.edu**

Summary

Many aspects of the porcine reproductive and respiratory syndrome virus (PRRSV) between-farm

transmission dynamics have been investigated, but uncertainty remains about the significance of

farm type and different transmission routes on PRRSV spread. We developed a stochastic

epidemiological model calibrated on weekly PRRSV outbreaks accounting for the population

dynamics in different pig production phases, breeding herds, gilt development units, nurseries, and

finisher farms, of three hog producer companies. Our model accounted for indirect contacts by the

close distance between farms (local transmission), between-farm animal movements (pig flow),

and reinfection of sow farms (re-break). The fitted model was used to examine the effectiveness of

vaccination strategies and complementary interventions such as enhanced PRRSV detection and

vaccination delays and forecast the spatial distribution of PRRSV outbreak. The results of our

analysis indicated that for sow farms, 59% of the simulated infections were related to local

transmission (e.g. airborne, feed deliveries, shared equipment) whereas 36% and 5% were related

to animal movements and re-break, respectively. For nursery farms, 80% of infections were related

to animal movements and 20% to local transmission; while at finisher farms it was split between

local transmission and animal movements. Assuming that the current vaccines are 1% effective in

mitigating between-farm PRRSV transmission, weaned pigs vaccination would reduce the

incidence of PRRSV outbreaks by 2%, indeed under any scenario vaccination alone was

insufficient for completely controlling PRRSV spread. Our results also showed that intensifying

PRRSV detection and/or vaccination pigs at placement increased the effectiveness of all simulated

vaccination strategies. Our model reproduced the incidence and PRRSV spatial distribution;

therefore, this model could also be used to map current and future farms at-risk. Finally, this model

could be a useful tool for veterinarians, allowing them to identify the effect of transmission routes

and different vaccination interventions to control PRRSV spread.

KEYWORDS: PRRSV, swine, disease spread, transmission dynamics, disease surveillance,

mechanistic modeling

Introduction

Porcine reproductive and respiratory syndrome virus (PRRSV) is a widespread swine disease in

North America (Hermann et al., 2007; Arruda et al., 2018; Sanhueza et al., 2019). PRRSV is

seasonal and shown significant epidemic peak during cold months (Perez et al., 2019; Trevisan et

al., 2020), causing major economic losses (Holtkamp et al., 2013). PRRSV incidence has decreased

in the last couple of years, certainly as a result of biosecurity improvements (Donovan, 2019;

Miller, 2019; Thomas, 2019).

In the last two decades, epidemiological studies have described the between-farm PRRSV

transmission dynamics (Dee et al., 2003a, 2004; Rosendal et al., 2014; Thakur, Revie et al., 2015;

Jara et al., 2020; Andraud and Rose, 2020). Direct contact among farms through pig transportation

together with indirect contacts often defined as local spread or local transmission, (e.g. airborne,

feed deliveries, shared equipment) have been identified as key routes of swine bacteria and virus

dispersal (Velasova et al., 2012; Lee et al., 2017; VanderWaal et al., 2018; Silva et al., 2019;

Bastard et al., 2020; Salines et al., 2020). Previous experimental and observational studies have

reported frequent PRRSV reinfection events on stable breeding farms (i.e., farrow-to-wean), in this

paper we will referred to re-break events (Pileri and Mateu, 2016). Even though there is no clear

consensus in the definition of what constitutes a re-break, the rebound of PRRSV could be related

to the continued viral shedding within groups of pigs and/or reduced cleaning and disinfection

procedures among other unknown factors and dynamics (Bierk et al., 2001; Arruda et al., 2016).

Despite the initial findings, the quantification of PRRSV transmission among multiple farm types

(e.g., from finisher farms to farrow-to-wean), remains a major gap in PRRSV epidemiology (Jara

et al., 2020; Salines et al., 2020).

Reducing the incidence of PRRSV outbreaks is among long term goals of some North

American swine producing companies. Currently, the industry relies heavily on vaccination and

biosecurity practices as control and prevention strategies (Silva et al., 2019), with the vaccination

of weaned pigs and gilts with modified-live vaccines (MLV) being among the most common (Kroll

et al., 2018; Garcia-Morante et al., 2020; Rawal et al., 2020). Several studies have evaluated the

efficacy of vaccines in increasing the immunity level and reducing shedding magnitude and

duration (Pileri et al., 2015; Chase-Topping et al., 2020); however the absence of studies

quantifying the effect of vaccination programs on between-farm PRRSV transmission, has limited

our capacity to investigate the effectiveness of current vaccination strategies and the development

of new, or the enhancement of existing PRRSV vaccination protocols (Andraud and Rose, 2020).

Existing disease spread models have explored the dynamics of between-farm PRRSV

transmission (Thakur, Revie et al., 2015; Thakur, Sanchez et al., 2015); a limited number of these

studies have investigated the deployment of control strategies on PRRSV outbreak incidence

(Bitsouni et al., 2019). To our knowledge no modeling framework has accounted for the spatio-

temporal spread dynamics of PRRSV in an epidemiological model. We developed an original

mechanistic model that to disentangle the relevance of transmission routes and serve as a

framework to test control strategies *in silico* (Brown and Bevins, 2018; Jurado et al., 2019). The

current epidemiological model is built to track the between-farm PRRSV spread routes among farm

types, breeding herds, gilt development units and growing pigs. The model outputs recorded i) the

relative contribution of the between-farm direct contacts by the movement of animals, local

transmission by indirect contacts governed mainly by the proximity among farms and exclusively

for sow farm we also estimated the probability of re-breaks; ii) the weekly number of new PRRSV

outbreaks, iii) the effect of different vaccination strategies in combination with intensified PRRSV

detection and delay in vaccination, and iv) the use of the fitted baseline model to forecast and

visualize the spatial distribution of at-risk farm locations.

Material and Methods

We developed a discrete-time stochastic model to simulate between-farm PRRSV transmission.

The model included the between-farm pig movements and indirect contacts by local transmission

events as a function of the distance between susceptible and infected farms, named hereafter simply

as local transmission. The probability of virus to persist within the sow farm named as re-break

event was also considered. The survival probability was included to consider PRRSV accumulation

in the environment in cases where proper cleaning and drying management cannot be performed

(Vilalta et al., 2019) and/or driven by a group of breeding age females or piglets that continue to

shed the virus and consequently maintain the transmission chain (Pileri and Mateu, 2016). The

methods are divided into sections as follows: i) description of the utilized data, animal movements

and the PRRSV outbreak incidence and ii) the mathematical modelling infrastructure including the

effectiveness of simulated vaccination strategies on PRRSV transmission and mapping at-risk

farms.

Data sources

Data from a total of 2,294 farms from three U.S. non-related pig production systems (identified as

A, B, and C) were obtained from the Morrison Swine Health Monitoring Project (MSHMP) project

(Perez et al., 2019). Data from each farm included their national premises identification number,

production type (sow [which included farrow, farrow-to-wean and farrow-to-feeder farms]),

nursery, finisher [which included wean-to-feeder, wean-to-finish, feeder-to-finish], gilt

development unit (GDU), isolation and boar stud, pig spaces per farm, and geographic coordinates.

The between-farm pig flow were used to reconstruct directed weekly contact networks

between June 1st, 2019 and December 5th, 2019. Each pig movement included date, farm of origin

and destination, number of pigs transported, and purpose of movement (e.g., weaning). Multiple

movement between two farms in the same week were considered as one unique contact and the

number of animals moved was summed. Missing movement data (e.g., either farm location,

production type, number of animals moved, the farm of origin, or destination) were excluded prior

to the analysis. The model included movements between 92% of the farms in the studied region,

and comprise a total of 19,984 movements.

The weekly number of PRRSV infected farms between January 5th , 2009 and December

5th , 2019, was compiled from two sources: i) for the outbreaks in sow farms we used the farm list

that each company uses to track farms undergoing outbreaks, re-break or elimination, and ii) for

the downstream farms (e.g. growing pig farms) we utilized ORF-5 PRRSV sequence from MSHMP

to identify outbreaks, this record is used by the veterinarians to monitor PRRSV strain circulation

(Perez et al., 2019; Jara et al., 2020). Briefly, a new sequence is often generated when clinical signs

are identified or there is a reason to request a sequence (Jara et al, 2020 for more details about

sampling and PRRSV dissemination). For sow farms, we classified each PRRSV occurrence as

"new outbreak" or "persistence outbreak" based on the reported outbreak dates. PRRSV detection

that occurred within the past 41 weeks from a previously reported PRRSV outbreak in the same

sow farm was considered a persistence outbreak, otherwise, it was classified as a new outbreak.

We based this classification on the average time to sow farms to reach PRRSV stable status

(Sanhueza et al., 2019). Similarly, for nurseries, GDUs, and finishers farms, we used general pig

production phase duration to count new PRRSV outbreaks, thus a PRRSV sequence in nurseries,

GDUs, and finishers farms within 7, 22, and 25 weeks from a previous record, respectively, was

considered a persistent outbreak; otherwise it was counted as a new outbreak. These time windows

were based on the general production schedule for nurseries, GDUs, and finisher farms under all-

in and all-out system. The infected farms were divided in two groups, for the initial setting we

utilized historical cases of PRRSV from January 5[th], 2009 - June 1[st], 2019, and for model simulation

pig movement and PRRSV outbreaks from June 1st - December 5th, 2019.

Between-farm infection dynamics

In brief, it was a stochastic model in discrete-time with a weekly time-step. We chose a one-week

time-step to match the frequency at which infected farms are reported to the MSHMP project (Perez

et al., 2019). Three farm heath statuses susceptible to infection (S), undetected infection i.e.,

infected but not yet tested or absence of clinical signs (U), and detected infection (D) were

considered (Figure 1). In this study, we assumed that all sow farms have been vaccinated or exposed

to PRRSV previously, thus all sites have the same probability to report a PRRSV outbreak. The

latent period is not explicitly modeled, since at animal level we expect that nasal shedding would

be present at day 7 post-inoculation (Pileri and Mateu, 2016; Chase-Topping et al., 2020), thus

latent is embedded within the weekly time-step. To model the local transmission, we modified a

gravity model to account for the effect of vegetation coverage surrounding each farm, previous

showed to modulate between-farm transmission dynamics (Jara et al., 2020), see equations 2-4.

Because PRRSV can persist undetected for long periods of time, mainly in contaminated

environments and/or fomites (Pileri and Mateu, 2016), for sow farms we included a re-break

probability, in which any sow farm that had a PRRSV outbreak in the past two year could have a

re-break (Perez et al., 2019; Jara et al., 2020). Transmission rates of the between-farm pig

movements, local transmission, and the re-break probability are represented hereafter by the beta

values βnet, βlocal, and βrebreak, respectively, fitted to the observed number of PRRSV infected

farms (details in the model fitting section).

Between-farm pig movements

The weekly pig movement network was utilized to identify contacts between susceptible farms i

and infected undetected farms and detected farms j, used to estimate the force of infection driven

by the weekly between-farm pig network (λnet):

$$\lambda net_{it} = \beta net * N_{it}; \ \forall_i = 1, ..., n \ suceptible \ farms \qquad (1)$$

160   N_i is the number of all infected farms (undetected and detected) sending pigs to susceptible farms

161   i at the time step t (one week).

162   ***Local transmission***

163   The local transmission was modeled through a gravity model as follow:

$$a_{ij} = \frac{m_i m_j}{d_{ij}^2} , \qquad (2)$$

where the attraction force between farms i and j was directly proportional to the farms' pig capacity

of m_i and m_j divided by the squared Euclidean distance d_{ij}^2 between two farms. Based on a

descriptive analysis of the observed outbreaks, the maximum range allowed for the local

transmission was set at 35km ($a_{ij} = 0$ when $d_{ij} >$ 35 km), d_{ij} was selected based model

performance, it is worth noticing that short or long distances pathogen spread remains a challenge

in disease epidemiology in general (Anderson et al., 2017; Machado et al., 2019). In addition, the

local transmission was modulated by the enhanced vegetation index (EVI), which was used to

quantify vegetation greenness at each farm location, here representing a barrier that reduced local

transmission of PRRSV as described elsewhere (Jara et al., 2020). The attraction force, a_{ijt},

between m_i and m_j was modeled as:

$$g_{ijt} = a_{ijt} * (1 - e_{it}) ; \ \forall_i = 1, \dots, n \ suceptible \ farms, \tag{3}$$

174 where g_{ijt} is the attraction force including the barrier effect, in which e_i is the barrier index from

175 the recipient farm i, represented on a [0,1] scale (Supplementary section 2 for additional

176 information). Thus, the local infectious force ($\lambda local$) was defined as:

$$\lambda local_{it} = \beta local * \sum_{j=1}^{n} g_{ijt} ; \ \forall_i = 1, \dots, n \ suceptible \ farms , \tag{4}$$

$$; \forall_j = 1, \dots, n \ infected \ farms$$

177 ***The re-break in sow farms***

178 Exclusively for sow farms, a re-break probability was coupled with the other two forces of

179 infection and included as

$$\lambda rebreak_{it} = \beta rebreak ; \ \forall_i = 1, \dots, n \ suceptible \ sow \tag{5}$$

$$farms \ with \ PRRSV \ records \ from \ may \ 31st, 2017 \ to \ May \ 31st, 2019,$$

180 where sow farms that tested positive from May 2017 until May 2019 had equal probability to re-

181 break throughout the simulated period.

182 **The overall transmission dynamics**

183 The total force of infection (λ) was calculated as the sum of all force of infection accounting for

184 the yearly PRRSV seasonality (Jara et al., 2020; Sanhueza et al., 2020),

$$\lambda_{it} = ((\lambda net_{it} + \lambda local_{it} + \lambda rebreak_{it}) * T_t) * (1 - H_{it}), \tag{6}$$

where λ_{it} is the total force of infection of a susceptible farm i and T_t is the seasonality index of

time step t. The seasonality index values were estimated from the moving averages of the monthly

frequency of PRRSV outbreaks between 2015 and 2019, which were then scaled between 0.001

and 1 (Supplementary section 1). The seasonality captured the variability in PRRSV transmission

events due to assumed changes in pigs' immunity and other estimated weather and environmental

factors (Dee et al., 2003a, 2003b). Because on-farm biosecurity in sow and GDU farms is expected

to play an important role the introduction and spread of new pathogens, we included a biosecurity

term (H), which modulated λ_{it}. Thus, H_{it} is defined between 0 and 1, and represents a reduction

in the total force of infection. Since no farm information regarding biosecurity measures was

available at the moment, we hypothesized that sow and GDU farms without PRRSV historic

records were more likely to have enhanced biosecurity measures, thus H_{it} was applied only for

sow and GDU farms without PRRSV outbreaks since 2009 (Table 1), while nurseries and finishers

it was assumed to be zero. The H value for sow and GDU farms was fitted to the observed number

of PRRSV infected farms (details at model fitting section) (Table 1). Finally, we calculated the

transmission probability (Y_{it}):

$$Y_{it} = 1 - e^{-\lambda_{it}}, \tag{7}$$

200   where Y_{it} is the transition probability from susceptible farm i to infected (and undetected) status at

201   time t. The detection rate for infected farms (transition from undetected to detected infection status)

202   was assumed to be dependent on the elapsed time of the farms in the infected undetected status and

203   is calculated by a detection probability derived from a logistic function

$$f(x) = \frac{L_p}{1 - e^{-k(x-x0)}} \tag{8}$$

to increase the detection rate at each time step following a sigmoid curve, where L is the maximum

detection probability for PRRSV outbreaks and its values were defined according to the farm

production types p. Thus, for sow and GDU farms which undergo periodic testing, we assumed L

to be high (0.95), while at nurseries and finishers farms (downstream) PRRSV detection is expected

to be underestimated, mainly because sampling efforts are much lower than in sow farms.

Therefore, in downstream farms L was fitted on the PRRSV outbreaks and to the local empirical

PRRSV prevalence (details at model fitting section). Here, $X0$ is the sigmoid's midpoint to detect

an infected undetected farm (i.e. which was assumed to be the average time for PRRSV detection),

defined as four weeks (Neira et al., 2017); k is the logistic growth rate, which we assumed to be

0.5, based on the local veterinarian observations in the study region; X is the time post-infection (in

weeks).

The proportion of recovered sow farms is drawn from a Poisson distribution with a mean

of 41 weeks (average time to stability) (Sanhueza et al., 2019). Nursery, GDU, and finisher farms'

transition to susceptible status were driven by pig production scheduling, all-in all-out or by ingoing

or outgoing pig movement, whatever come first in the data. Briefly, nurseries, GDU and finisher

farms were assumed to become susceptible within 7, 22 and 25 weeks of pig placement,

respectively, or when a movement is recorded before reaching the scheduled production phase

timeline.

As an initial model setting, farms were considered to be infected if they presented a

previous outbreak within a determined number of weeks before June 1^{st}, 2019 (initial start point in

the simulations) according to their farm type as follows: sow 41, nursery 7, GDU 22, and finisher

25 weeks. All of the remaining farms were assumed to be susceptible.

Model fitting

We used an Approximate Bayesian Computation (ABC) method based on a modified Sequential

Monte Carlo (SMC) rejection algorithm, which allows us to recover the approximate posterior

distribution of six model parameters βnet, $\beta local$, $\beta rebreak$, H (sow and GDU fitted individually),

and L (nursery, GDU and finisher farms were fitted to one parameter). A set of draws of these six

parameters is denominated a "particle" (Beaumont, 2010), and their values were fitted to the

observed infected farms for each production type from June 1^{st}, 2019, until December 5^{th}, 2019.

Although outbreaks were reported at nursery and finisher farms, underdetection is a well-known

limitation of PRRSV surveillance in those downstream farms (Velasova et al., 2012). Thus, we

used an expected weekly number of nursery and finisher infected farms, derived from an annual

prevalence of 30% (personal communications from veterinarians from the production companies

we collected the data), and the outbreaks distributed throughout the year according to an estimated

seasonality term T (Supplementary section 3).

The modified SMC algorithm was a two-step rejection procedure. In the first step, the

model was fitted to the weekly number of infected farms by production type using eleven summary

statistics (Supplementary Table S1). In the second step, the sensitivity and specificity of the model

to predict true risk areas was evaluated through the comparison between simulated and observed

infected farm locations falling within square spatial areas of 10 km x 10 km, named hereafter as

cells (Supplementary section 3), these final fitting step considered PRRSV outbreaks in sow farms

(i.e. true outbreaks reported in our data). The ABC rejection procedure was run to generate 100

accepted particles (N), within the tolerance interval (ϵ) (Supplementary Table S1). We chose the

particle with the highest sensitivity in the second step of the fitting procedure for all the subsequent

simulations of the different control strategies. Table 1 summarizes the model parameters used.

The effect of vaccination strategies on PRRSV between-farm transmission

We simulated five vaccinations programs and investigated their impact at reducing the between-

farm transmission (Corzo et al., 2010; Azman and Lessler, 2015) (Table 2). We considered a range

of vaccine efficacies, here we assumed that mass vaccination reduced the force of infection (the

rate at which susceptible to infection farms acquire new infection, Figure 1) by 1%, 2%, 3%, 4%,

5%, 10%, 15%, 20%, 30%, 50%, 80% and 90%. We assumed that vaccine efficacy would have a

maximum effect in reducing force of PRRSV between-farm transmission four weeks after

vaccination, here it was modeled by a logistic growth curve described before (Kick et al., 2019)

(for more details about how ve was modeled refer to Supplementary material section 4). Because

information about the length of post-vaccination herd-level immunity was not available, we

assumed vaccinated pigs maintained protection for 36 weeks, based on expert opinion and previous

modelling studies (Evans et al., 2010; Arruda et al., 2016). Therefore, for nurseries, GDUs, and

finisher farms, one dose of vaccine protected pigs throughout the whole production phase. The

effectiveness of the vaccination strategies by production type was calculated by comparing the

reduction in the total number of simulated infected farms for 27 weeks to the counterfactual

scenario with no intervention.

Additionally, we also considered complementary interventions that match more closely

current field conditions and practical challenges in terms of vaccination delays and the effect of

testing efforts (i.e., decisions about testing finisher farms even if disease is clinical). We examined

the impact of i) delaying vaccination for one, two, and three weeks after pig placement, ii) delaying

vaccination for one, two, and three weeks post PRRSV detection, iii) enhanced detection scenarios

that assumed detection was able to detect 50% or 95% of PRRSV circulation in nurseries, GDUs,

and finisher farms, and iv) the average time to detect was set to be between one and three weeks of

the virus introduction. All control scenarios were run with 50 repeats to stabilize distributions of

the simulated incidence of infected farms. A multivariate linear regression analysis was used to

estimate the contribution of each vaccination strategy, individually for each farm type. The outcome

of interest was the proportion of prevented PRRSV outbreaks as a function of all complementary

interventions while accounting for the range of vaccine efficacies (Supplementary Tables S2 to S6).

Model outputs

The model outputs included i) the force of infection for each farm type and transmission route, ii)

the weekly number of infected undetected and detected farms, iii) the number of prevented PRRSV

infected farms for each vaccination strategy, and iv) maps of the location where PRRSV outbreaks

are more likely to occur and farms that have recently been infected (recent introductions). We

carried out 1000 runs to estimate the relative contribution of each transmission route. Here we

evaluated the number of infected farms resulting from each transmission route individually, which

were then divided by the number of simulated infected farms from the three combined routes.

To evaluate the model prediction performance of PRRSV infected farm locations, of all

sow farms (319 farms) were grouped into 154 cells (10 km x10 km), in which the number of sow

farms by cell had a median of 1 and a maximum of 12 farms (Figure 4). The at-risk PRRSV maps

were based on the average number of times farms in a given cell were estimated to be infected and

posteriorly detected by the model; these cell values were defined as a risk of PRRSV introduction

and detection, and were calculated through 100 simulations. Thus, the model spatial forecasting

produced two main outputs: a) the risk of PRRSV introduction, here representing areas with farms

recently infected; and b) the risk of PRRSV detection, here representing areas with infected but

undetected farms that are more likely to be detected. Therefore, we proposed a weekly mapping

forecast for both outputs calibrated on the transmission dynamics and observed infected farms from

prior weeks; details about predictive performance are described in supplementary material under

sensitivity analysis, and an illustrative example is shown in Figure 5.

3. Results

In Table 3 the number of susceptible and infected farms are summarized. The dataset used to

calibrate the model parameters included 48 infected farms reported between June 1[st] 2019, and

December 5[th] 2019, 27 weeks. Isolation farms and boar studs had no PRRSV detection.

Contribution of each transmission route to PRRSV spread

The results shown here are derived from the fitted particle with higher precision capturing the

spatial distribution of infected farms (Table 1). Among the three transmission routes, on average

59% of the simulated infections in sow farms were related to local transmission, and 36% and 5%

were related with animal movements and re-break, respectively (Figure 2). For nursery farms, 80%

of the simulated infections were related to animal movements and 20% to local transmission, while

for finisher, movements and local transmission were similarly important with 47% and 53% of the

transmissions, respectively. Finally, GDU, 99% of the infected farms were associated with local

transmission.

The estimated number of new PRRSV infected farms

The estimated average number of new infected farms in the simulated period was 189 (95% CI:

188-191), corresponding to 20 (95% CI: 19-20) newly infected sow farms, 63 (95% CI: 63-64)

nursery farms, 10 (95% CI 9-10) GDU farms and 96 (95% CI: 95-97) finisher farms (Figure 3A).

There was a good agreement between the weekly observed outbreaks and simulated detected farms

(Figure 3B). The model inferred that at the end of the 27 weeks on average 153 of the 189 infected

farms would not be detected, in which 81% of the infected sow farms were detected, while a much

lower proportion of detection was estimated for GDU 24%, nurseries 11% and for finisher 11%

(Figure 3).

The effectiveness of vaccination strategies

The total number of PRRSV infected farms declined as *vaccine efficacy* increased (Figure 5, and

Supplementary Tables S2-S6). A *ve* of 1% was sufficient to reduce on average at least 2% of the

total number of infected farms independently of farm types. The combined vaccination strategy at

vaccine efficacy of 1% reduced the average number of infected farms by 12%, likewise at *ve*=1%

prophylactic #2 and reactive #2 reduced infected farms by 7%, while prophylactic #1 and reactive

#1 prevented between 2% and 1%, respectively. Among the five vaccination strategies, the

combined strategy showed to be most effective (Figure 4-A), i.e., at *ve = 5%,* the average number

of infected farms would be reduced by 46%, which means four times more effective than the

prophylactic #1 and reactive #1, and almost two times more effective than prophylactic #2 and

reactive #2. The second most efficient strategies were the prophylactic #2 and reactive #2, which

showed similar results at *ve* < 20% (Figure 4-A), however reactive #2 performed better with *ve*

above 20%. Finally, the prophylactic #1 and reactive #1 strategies marginally reduced PRRSV

spread events (Figure 4-A, Supplementary Tables S2-S6).

In addition, in Figure 5-B the simulated vaccination strategies are showed by farm type.

Prophylactic #2 showed to work best for finisher farms, while sow, GDU, nursery farms were less

likely to be infected under reactive #2 (Figure 4-B). It is worth noting that the combined and

reactive #2 strategies rapidly reduced the number of new PRRSV outbreaks in sow and GDU farms,

reducing the number of infected farms by 25% with vaccine efficacies of 5% (Figure 4-B).

Supplementary Tables S2-S6 summarizes the results from the multivariate regression

analysis, which depicted the effect of delays in the implementation of vaccine interventions and

improvements in the PRRSV detections. The effect of complementary interventions ranged from

by 0.1% and 4.9%. However, the performance of these complementary interventions depended on

the vaccination strategies and the farm type (Supplementary table S2-S6). The most effective

interventions was to vaccinate pigs at the same week of placement and increasing the probability

of PRRSV detection in downstream farms (Supplementary Tables S2-S6). Overall most

complementary intervention showed to have a better impact on PRRSV spread when deployed to

finisher and nurseries farms.

The spatial-temporal risk of PRRSV spread

The calibrated model was used to map the PRRSV spatial risk patterns. The direct comparison of

predicted infected cells with the true infected farms weekly, was used to evaluate the model's

predictive ability. Under the an optimal threshold for disease detection, explained in details in

supplementary Figure S3, the model predicted correctly infected sow farms with a median

sensitivity of 62%. Figure 5 shows the main spatial features of the modelling forecast. Here we

defined high-risk cells were all values with risk above the 90[th] percentile; this percentile value was

chosen as a rigorous threshold in order to evaluate the model capabilities to identify locations with

high infection risks with the least number of cells (Supplementary Figure S4). In Figure 5-A, we

show a map with the true outbreaks reported in our data, Figure 5-B shows the predicted risk of

infection, and Figure 5-C shows the cells with risk values above the 90[th] percentile threshold. In

this example two of the three truly infected cells were correctly predicted (Supplementary Figure

S4). In addition, the model was also used to estimate the risk of a recent PRRSV introduction

(Figure 5-D and 5-E), thus the identified cells more likely to have at least one infected farms which

has not yet been identified as positive.

4. Discussion

We demonstrated that the dynamics of between-farm PRRSV transmission was strongly modulated

by farm types. Our results showed that local transmission generated the majority of the infections

in sow (that include farrow, farrow-to-wean and farrow-to-feeder farms) and GDU farms; however,

nurseries were more likely infected through animal movements as expected, while for finisher

farms (which included wean-to-feeder, wean-to-finish, feeder-to-finish farms) both routes were

equally important. Quantifying the routes of transmission can be used in the development of new

disease control strategies targeting the dominant transmission routes at each specific farm type

(Andraud et al., 2019; Machado et al., 2019). With regards to control strategies, we evaluated the

effect of vaccination, enhanced detection and diagnostic performance on the number of prevented

PRRSV infected farms. The vaccination of pigs at placement (prophylactic #2) combined with

mass-vaccination during PRRSV outbreak (reactive #2), showed to be the most efficient strategy.

It is worth noting that the performance of vaccination strategies with the same efficacy varied

among farm types. The best vaccination approach for sow and GDU farms was reactive #2. For

nursery, the vaccination of weaned pigs (prophylactic #1) was nearly as good as (re)vaccinating

finishers and gilts at placement (prophylactic #2). Similarly, prophylactic #2 was highly efficient

to prevent infection in finisher farms, and implement vaccination within one week of placement

was the best complementary strategy. We also showed that the control of PRRSV spread can be

optimized if diagnostic in downstream farms is enhanced and if vaccination is deployed early on

PRRSV infection (Henao-Diaz et al., 2020). Finally, based on our baseline model we present two

mapping results, a forecasting map showing the most likely infected farms and a map showing at-

risk locations in which the risk that PRRSV has been introduced within an average of four weeks

was estimated to be high. Ultimately, this can help the identification of farms where interventions

would be more effective at reducing the between-farm transmission (Stevenson et al., 1993;

Bilodeau et al., 1994; Hayama et al., 2013, 2020).

Contribution of transmission routes

The important contribution of the local transmission route in spreading PRRSV into sow, GDU and

finisher farms was consistent with previous studies, which demonstrated that farms located close

to infected sites were more likely to report PRRSV outbreaks (Velasova et al., 2012; Phoo-ngurn

et al., 2019; Silva et al., 2019; Jara et al., 2020), while others suggested that contact networks was

the preferred transmission route (Amirpour Haredasht et al., 2017; Lee et al., 2017; Bastard et al.,

2020; VanderWaal et al., 2020). Our approach to model the local PRRSV spread was unique in that

we considered the effect of vegetation around each farm location as a physical barrier against

mainly airborne transmission (Van Ryswyk et al., 2019; Jara et al., 2020). Consequently, in our

modelling framework, farms at a location with a high vegetation index were less likely to become

infected by local transmission route (Jara et al., 2020).

Interestingly, finisher farms transmission rates of finisher farms were less related to the in-

going movements when compared with nurseries. This could be attributed to several factors that

likely facilitated the between finisher farms local transmission, including high regional finisher

farm PRRSV prevalence (personal communication, local veterinarians) and the disproportionate

number of susceptible finisher farms. Descriptively, the median number of finisher farms in a 5 km

radius around each finisher farms (hereafter defined as a neighborhood) was nine, which was higher

than the median of six finishers around nursery farms (Supplementary Figure S5). It is worth

considering that the number of pigs flowing into the neighborhoods densely populated with finisher

farms is expected to be higher, which could also play an important role in sustaining spread at a

local scale due to the regular reintroduction of the virus (Otake et al., 2010; Machado et al., 2019;

Ezanno et al., 2020). Neighborhood contacts generally occur in a continuous manner over rather

long periods of time (Ezanno et al., 2020), which rise indirect contacts (e.g., airborne) which would

vary over time. Similarly, the intense local transmission into sow and GDU farms may be equally

associated with a high number of farms in those neighborhoods, especially in neighborhoods

heavily populated by farms with known restricted biosecurity, such as finisher farms

(Supplementary Figure S5). The results of our local transmission route could be also associated

with indirect contacts via vectors and fomites (e.g. farm visitors, other animals, insects, and

contaminated transport-tools) (Pitkin, Deen and Dee, 2009; Pitkin, Deen, Otake et al., 2009; Reiner

et al., 2009; Lowe et al., 2014). In future work it would be pivotal to include indirect contact

networks such as feed deliveries and mortality management through dead pickups (Porphyre et al.,

2020), which would allow PRRSV to spread throughout multilayer networks, for example. This

will likely shed light on other routes involved in PRRSV spread (Dee et al., 2020).

As described elsewhere, the number of in-going movements into sow and GDU farms have

not been associated with the introduction of pathogens (Lee et al., 2017; Bastard et al., 2020),

indeed, poorly connected sow and GDU farms are expected to be less infected through receiving

infected . In our study, this could be one of the reasons why pig movements were less relevant for

sow and GDU farms than the local transmissions; however, a thorough network analysis is still

necessary to fully evaluate this findings. Nursery and finisher farms on the other hand are more

prone to be infected by animal movements, previously associated with their network position

(Passafaro et al., 2020), as observed in other studies (Lee et al., 2017; Bastard et al., 2020), similarly

to the result described here.

This study uniquely considered PRRSV re-break events as a source of infection for sow

farms, which appears to be significantly lower in comparison with the local transmission and the

pig movement routes. On average, the re-breaks were associated with 5% of the outbreaks in sow

farms. Given the assumption and the lack of data to calibrate the re-breaks, it is possible that this

transmission route has been underestimated. Indeed, it does not mean that re-breaks are not relevant

in the epidemiology of PRRSV. Previous investigations described the relevance of sub-populations

which could maintain the virus population shedding for an extended period without being identified

(Bierk et al., 2001; Evans et al., 2010; Pileri and Mateu, 2016). Our limited understanding of

PRRSV infection dynamics in terms of herd immunity added to the complexity of the population

dynamics inside multi-barn pig farms, such as in sow farms that include farrowing and gestation

barns and the constant movement of animals, equipment and people across barns is a major gap

that needs further analysis before we can disentangle the role of re-breaks in the epidemiology of

PRRSV.

The effectiveness of control strategies

The vast majority of commercial pig producers in the US currently utilize commercial modified

live-virus (MLV) vaccines (Pileri and Mateu, 2016; USDA, 2016). The limited effectiveness of

vaccines in field conditions (Charerntantanakul, 2012; Roca et al., 2012), highlight the need to

estimate the level of protection required to successfully reduce PRRSV between-farm transmission.

As describe elsewhere (Linhares et al., 2012), suggested that vaccination can reduce PRRSV

shedding rates, a more recent experiment with a European type subtype 1 strain, indicated that the

current vaccines have a poor efficacy to reduce between animal transmission (Chase-Topping et

al., 2020). Therefore, remain under debate the impact MLV vaccines on the concentration of

PRRSV in the environment and the reduction of virus shedding. Thus *ve* for between-farm

transmission at 1% may be a fair assumption, however proper experimental studies are needed to

quantify it. According to our modeling, a vaccine efficiency of 80% would be necessary to

completely prevent new transmission events. Our simulations reinforce the need for improving the

current vaccine technologies (Bitsouni et al., 2019), meanwhile non-therapeutic interventions such

as enhancement of on-farm biosecurity will continue to be essential in reducing PRRSV spread

(Silva et al., 2019).

When comparing the effectiveness of prophylactic strategies #1 and #2 for finisher farms,

the former strategy was better. The extra vaccine boost likely reduced the force of infection of both

routes, animal movement and local transmission, either by increasing the protection of susceptible

farms or by decreasing the force of infection from infected and vaccinated farms. On the other

hand, both prophylactic strategies provided limited protection for sow and GDU farms (Figure 2

and 5). An added value of prophylactic #2 is that finisher farms may be the main source of virus to

other farm types within the neighborhood. In fact, studies showed that the vaccination of finisher

sites resulted in a reduction of the likelihood of airborne transmission and or transmission by

fomites due to the reduction of nasal shedding in vaccinated pigs (Linhares et al., 2012; Pileri et

al., 2015). It is important to remark that vaccination of either prophylactic program showed to work

best in nursery and finisher farms if deployed in the same week of placement when compared with

a later vaccination, for example 3 weeks after pigs arrive.

The vaccination of PRRSV positive farms was directly dependent on virus detection and

reporting (Perez et al., 2019). Therefore, if all farms with detected PRRSV infection were

vaccinated as proposed in the reactive #2 strategy, the number of prevented infected farms was

approximately the same as the preventive prophylactic #2 scheme, described above. The addition

of nursery, GDU and finisher farms in the vaccination of infected farms (reactive #2) was better

than reactive #1, even though the model assumed that PRRSV detection rates in downstream farms

was at 7% (Table 1). The main limitation of reactive #2 is related to PRRSV diagnostic in finisher

and nursery farms. The reactive #2 strategy could potentially be more effective than the

prophylactic #2 in terms the volume of vaccine doses. On the other hand, the prophylactic #2

strategy would be independent from diagnostics costs, and thus implementation could be

considered more feasible by the swine industry. Before recommendations can be made, future

longitudinal studies need to evaluate the economics of each vaccination strategy taking into

consideration the cost to stabilize an infected farm (Linhares et al., 2015; Nathues et al., 2017;

Thomann et al., 2020). Until diagnostic techniques become more accessible and easier to perform,

the effectiveness of strategies involving the identification of PRRSV in downstream pig operations

remains questionable.

The spatial-temporal risk of PRRSV

Unlike many other modelling frameworks, which calibrate models explicitly to fit on the number

of outbreaks over time (Brooks-Pollock et al., 2014; Funk et al., 2018), here we proposed the

calibration to both the space- and time- distribution of PRRSV outbreaks (Sumner et al., 2017;

VanderWaal et al., 2018). This approach allowed a closer approximation to the weekly incidence

and the true infected farms (Widgren et al., 2018), thus our model was able to correctly capture at-

risk areas with at least one infected farm (Supplementary Figure S3). It is important to note that the

accuracy of the forecast is expected to vary as PRRSV incidence changes along the season

(Trevisan et al., 2020); therefore, throughout our weekly simulations, the overall sensitivity ranged

from 50% to 73%, at the proposer optimal risk thresholds. Improving the spatial forecast sensitivity

to reproduce the PRRSV outbreaks spread an important challenge, this is because fitted models to

real data may not capture completely the disease transmission fluctuation, similarly to other studies

that have also attempted to model the distribution of *E. Coli* or Porcine Epidemic Diarrhea

(VanderWaal et al., 2018; Widgren et al., 2018; Machado et al., 2019). Our study was in agreement

with a previous study that also highlighted the challenges to predict PRRSV infection in sow farms,

when compared with nurseries, wean-to-finish and finisher (Amirpour Haredasht et al., 2017). The

challenges of model fitting could be attributed to the lack of information about other contact

networks that connect farms indirectly, such the routes taken by feed deliveries and dead picks,

which could explain in a greater detail the spatial dispersion of PRRSV. Recent work reinforced

the relevance of the indirect contacts among farms, in which sharing haulage vehicles has

significant potential for spreading infectious diseases within the pig sector in Great Britain

(Porphyre et al., 2020). Therefore, futures including other indirect contact networks into our model

framework, would also likely to improve our forecast accuracy.

The forecast sensitivity of 62% was obtained under the optimal risk threshold; however,

the number of cells identified to be at greater risk of PRRSV infection, were on average 55 cells.

In order to define a feasible threshold for the identification of at-risk areas, there is a need to not

only consider model precision, but also the economics of possible targeted deployment of control

strategies. In the example shown in this study (Figure 5), we proposed high risk cells to be those

with risk above the 90[th] percentile, thus on average 16 cells were identified to be at risk. As result

the model sensitivity was lower than the overall mean value (Supplementary Figure S3). On the

other hand, less being more extract to which cells are at risk, made this approach more useful, and

informative for the stakeholders' response and decision making, which could include the formation

of a forecast dashboard to expedite risk communication and guide mode precise interventions (e.g.,

vaccination). A more uneconomic threshold could be more effective in identifying outbreaks, but

it would be necessary to evaluate the capacity of the production companies to intensify testing

before these thresholds can be proposed. Our model also identified cells that did not present

outbreaks (false positives), which could be associated with the presence of infected not yet detected

farms. The swine industry could directly benefit from this approach by testing farms in which the

model predicted a recent PRRSV introduction or enhance control strategies i.e., vaccination,

enhancement of on-farm biosecurity to specific farms (Brooks-Pollock et al., 2014).

Limitations and further remarks

Our study has a number of simplifications and limitations starting with the study being conducted

in one region where three production companies are present which may not entirely represent what

occurs throughout the country. Also, the number of cases reported in downstream farms were

counted based in the number of ORF-5 PRRSV sequences, which was likely underestimated,

mainly because those sites do not undergo systematically testing. In addition, the fact that

individual veterinarians' together with the diagnostician decide which sample to be sequenced may

underestimate viral diversity. However, thus far we believe that the use of ORF-5 PRRSV

sequences to capture viral circulation in growing pig farms is still the best alternative to the current

PRRSV available databases. In addition, we did not consider the influence of individual viral

lineages while modeling PRRSV transmission forces, we assumed that all samples had the same

virulence, thus the probability of the between-farm transmission was assumed to be the same (Jara

et al., 2020). Modelling individual PRRSV strain dissemination was beyond the scope of this study,

but this is likely a potentially important future consideration as part of our active research focus

area. In the current modelling framework, we were interested in the between-farm transmission, in

which we did not include more detailed dynamics of individual pigs within farms, thus futures

studies including within-farm dynamics at animal level will be an important component of future

studies (Andraud and Rose, 2020). In addition, the model was calibrated on the incidence across

all farm types for the temporal model calibration, while we decided to fit the spatial component to

infected sow farms only, largely because PRRSV outbreaks records in sow farms were more

consistent allowing for a more confident assessment of model sensitivity. Transmission events not

explained by between-farm movements or re-breaks at sow farms were classified in the model as

local transmitted, thus future studies will include other movement flows such as feed truck

deliveries in order to explain in more details this route of transmission (Porphyre et al., 2020). For

the assessment of vaccination strategies, we assumed that vaccinated farms acquire life-long

immunity, future studies should also consider the heterologous host responses to vaccination

(Ezanno et al., 2020).

Conclusion

This study establishes a modeling framework to explore between-farm PRRSV transmission, which

was used to a) quantify the contribution of transmission routes for individual farm types, b) quantify

the number of infected farms prevented by vaccination strategies, and c) forecast current and future

PRRSV spatial distribution. Our findings reveal that local transmission is a key driver of PRRSV

spread into sow and GDU farms, whereas animal transportation into downstream farms causes the

majority of the infections in nurseries. Animal transportation and local transmission are equally

responsible for introducing infections into finisher farms. Vaccination alone was insufficient for

completely controlling PRRSV spread. Weaned pigs vaccination showed to be effective, however

when we assumed that vaccine efficacies were at ≥3% and ≥10%, the model estimated 10% and

25% reduction in PRRSV outbreaks, respectively. Our results also reiterated that the

implementation of additional strategies that can reduce PRRSV collateral spread, such as on-farm

biosecurity, will continue to play a key role in reducing collateral transmission, especially if those

strategies are also applied in finisher farms, from which PRRSV was easily spread via local

transmission route. The detailed outputs of our spatially explicit model closely reproduce the

empirical spatial distribution of PRRSV infection in sow farms. For this reason, the risk forecasted

by the fitted model has the potential to guide targeted control interventions. Ultimately, this study

offers producers and swine veterinarian's greater insight into mechanisms of PRRSV spread

between different farm types, and how vaccination programs could more effectively in reducing

disease transmission.

Acknowledgements

The primary funding support of this project is from the Swine Health Information Center under

project #19-211. This work was also supported by Critical Agricultural Research and Extension

2019-68008-29910 from the USDA National Institute of Food and Agriculture. This project also

fostered international collaboration between Dr. Machado and Dr. Prada laboratory from the travel

support provided by the University Global Partnership Network (UGPN) Research Collaboration

Fund. The Morrison Swine Health Monitoring Project is a Swine Health Information Center funded

project. Authors would like to acknowledge participating systems and veterinarians.

Authors' contributions

JAG and GM conceived the study. JAG, JP, and GM participated in the design of the study. CC

coordinated the PRRSV data collection via MSHMP. JAG conducted data processing, cleaning,

designed the model, simulation scenarios with the assistance of JP and GM. JAG designed the

computational analysis. JAG, JP, and GM wrote and edited the manuscript. All authors discussed

the results and critically reviewed the manuscript. GM recurred the USDA-NIFA funding.

Conflict of interest

All authors confirm that there are no conflicts of interest to declare

Ethical statement

The authors confirm the ethical policies of the journal, as noted on the journal's author guidelines

page. Since this work did not involve animal sampling nor questionnaire data collection by the

researchers there was no need for ethics permits.

Data Availability Statement

The data that support the findings of this study are not publicly available and are protected by

confidential agreements, therefore, are not available.

References

Amirpour Haredasht, S., D. Polson, R. Main, K. Lee, D. Holtkamp, and B. Martínez-López,

2017: Modeling the spatio-temporal dynamics of porcine reproductive & respiratory

syndrome cases at farm level using geographical distance and pig trade network matrices.

BMC Vet. Res. **13**, 163, DOI: 10.1186/s12917-017-1076-6.

Anderson, B.D., J.A. Lednicky, M. Torremorell, and G.C. Gray, 2017: The Use of Bioaerosol

Sampling for Airborne Virus Surveillance in Swine Production Facilities: A Mini Review.

Front. Vet. Sci. **4**, DOI: 10.3389/fvets.2017.00121.

Andraud, M., T. Halasa, A. Boklund, and N. Rose, 2019: Threat to the French Swine Industry of

African Swine Fever: Surveillance, Spread, and Control Perspectives. *Front. Vet. Sci.* **6**,

DOI: 10.3389/fvets.2019.00248.

Andraud, M., and N. Rose, 2020: Modelling infectious viral diseases in swine populations: a state

of the art. *Porc. Heal. Manag.* **6**, 22, DOI: 10.1186/s40813-020-00160-4.

Arruda, A.G., R. Friendship, J. Carpenter, A. Greer, and Z. Poljak, 2016: Evaluation of Control

Strategies for Porcine Reproductive and Respiratory Syndrome (PRRS) in Swine Breeding

Herds Using a Discrete Event Agent-Based Model. (Javier Sanchez, Ed.)*PLoS One* **11**,

e0166596, DOI: 10.1371/journal.pone.0166596.

Arruda, A.G., C. Vilalta, P. Puig, A. Perez, and A. Alba, 2018: Time-series analysis for porcine

reproductive and respiratory syndrome in the United States. (Yongchang Cao, Ed.)*PLoS

One* **13**, e0195282, DOI: 10.1371/journal.pone.0195282.

Azman, A.S., and J. Lessler, 2015: Reactive vaccination in the presence of disease hotspots. *Proc.

R. Soc. B Biol. Sci.* **282**, 20141341, DOI: 10.1098/rspb.2014.1341.

Bastard, J., M. Andraud, C. Chauvin, P. Glaser, L. Opatowski, and L. Temime, 2020: Dynamics

of livestock-associated methicillin resistant Staphylococcus aureus in pig movement

networks: Insight from mathematical modeling and French data. *Epidemics* **31**, 100389,

DOI: 10.1016/j.epidem.2020.100389.

Beaumont, M.A., 2010: Approximate Bayesian Computation in Evolution and Ecology. *Annu.*

Rev. Ecol. Evol. Syst. **41**, 379–406, DOI: 10.1146/annurev-ecolsys-102209-144621.

Bierk, M.D., S.A. Dee, K.D. Rossow, S. Otake, J.E. Collins, and T.W. Molitor, 2001:

Transmission of porcine reproductive and respiratory syndrome virus from persistently

infected sows to contact controls. *Can. J. Vet. Res.* **65**, 261–266.

Bilodeau, R., D. Archambault, S.A. Vézina, R. Sauvageau, M. Fournier, and S. Dea, 1994:

Persistence of porcine reproductive and respiratory syndrome virus infection in a swine

operation. *Can. J. Vet. Res.* **58**, 291–298.

Bitsouni, V., S. Lycett, T. Opriessnig, and A. Doeschl-Wilson, 2019: Predicting vaccine

effectiveness in livestock populations: A theoretical framework applied to PRRS virus

infections in pigs. (Yongchang Cao, Ed.)*PLoS One* **14**, e0220738, DOI:

10.1371/journal.pone.0220738.

Brooks-Pollock, E., G.O. Roberts, and M.J. Keeling, 2014: A dynamic model of bovine

tuberculosis spread and control in Great Britain. *Nature* **511**, 228–231, DOI:

10.1038/nature13529.

Brown, V.R., and S.N. Bevins, 2018: A Review of African Swine Fever and the Potential for

Introduction into the United States and the Possibility of Subsequent Establishment in Feral

Swine and Native Ticks. *Front. Vet. Sci.* **5**, DOI: 10.3389/fvets.2018.00011.

Charerntantanakul, W., 2012: Porcine reproductive and respiratory syndrome virus vaccines:

Immunogenicity, efficacy and safety aspects. *World J. Virol.* **1**, 23, DOI:

10.5501/wjv.v1.i1.23.

Chase-Topping, M., J. Xie, C. Pooley, I. Trus, C. Bonckaert, K. Rediger, R.I. Bailey, H. Brown,

V. Bitsouni, M.B. Barrio, S. Gueguen, H. Nauwynck, and A. Doeschl-Wilson, 2020: New

insights about vaccine effectiveness: Impact of attenuated PRRS-strain vaccination on

heterologous strain transmission. *Vaccine* **38**, 3050–3061, DOI:

10.1016/j.vaccine.2020.02.015.

Corzo, C.A., E. Mondaca, S. Wayne, M. Torremorell, S. Dee, P. Davies, and R.B. Morrison,

2010: Control and elimination of porcine reproductive and respiratory syndrome virus.

Virus Res. **154**, 185–192, DOI: 10.1016/j.virusres.2010.08.016.

Dee, S.A., J. Deen, S. Otake, and C. Pijoan, 2004: An experimental model to evaluate the role of

transport vehicles as a source of transmission of porcine reproductive and respiratory

syndrome virus to susceptible pigs. *Can. J. Vet. Res.* **68**, 128–133.

Dee, S.A., J. Deen, K. Rossow, C. Weise, R. Eliason, S. Otake, H.S. Joo, and C. Pijoan, 2003a:

Mechanical transmission of porcine reproductive and respiratory syndrome virus throughout

a coordinated sequence of events during cold weather. *Can. J. Vet. Res.* **66**, 232–239.

Dee, S.A., J. Deen, K. Rossow, C. Weise, R. Eliason, S. Otake, H.S. Joo, and C. Pijoan, 2003b:

Mechanical transmission of porcine reproductive and respiratory syndrome virus throughout

a coordinated sequence of events during warm weather. *Can. J. Vet. Res.* **67**, 12–19.

Dee, S.A., M.C. Niederwerder, G. Patterson, R. Cochrane, C. Jones, D. Diel, E. Brockhoff, E.

Nelson, G. Spronk, and P. Sundberg, 2020: The risk of viral transmission in feed: What do

we know, what do we do? *Transbound. Emerg. Dis.*tbed.13606, DOI: 10.1111/tbed.13606.

Donovan, T., 2019: The Hanor Company. p. 11. In: *2019 Allen D. Leman Swine Conf.* Saint Paul,

Minnesota.

Evans, C.M., G.F. Medley, S.J. Creasey, and L.E. Green, 2010: A stochastic mathematical model

of the within-herd transmission dynamics of porcine reproductive and respiratory syndrome

virus (PRRSV): Fade-out and persistence. *Prev. Vet. Med.* **93**, 248–257, DOI:

10.1016/j.prevetmed.2009.11.001.

Ezanno, P., M. Andraud, G. Beaunée, T. Hoch, S. Krebs, A. Rault, S. Touzeau, E. Vergu, and S.

Widgren, 2020: How mechanistic modelling supports decision making for the control of

enzootic infectious diseases. *Epidemics* **32**, 100398, DOI: 10.1016/j.epidem.2020.100398.

Funk, S., A. Camacho, A.J. Kucharski, R.M. Eggo, and W.J. Edmunds, 2018: Real-time

forecasting of infectious disease dynamics with a stochastic semi-mechanistic model.

Epidemics **22**, 56–61, DOI: 10.1016/j.epidem.2016.11.003.

Garcia-Morante, B., R. Friedrich, T. Kaiser, C. Kraft, P. Bridger, and M. Noguera, 2020: Gilt

Vaccination with a Mixed Administration of a PRRS MLV and a PPV1 Subunit Vaccine

Protects against Heterologous PRRSV1 Infection and Prevents Detrimental Effects on Piglet

Performance. *Viruses* **12**, 789, DOI: 10.3390/v12080789.

Hayama, Y., Y. Shimizu, Y. Murato, K. Sawai, and T. Yamamoto, 2020: Estimation of infection

risk on pig farms in infected wild boar areas—Epidemiological analysis for the reemergence

of classical swine fever in Japan in 2018. *Prev. Vet. Med.* **175**, 104873, DOI:

10.1016/j.prevetmed.2019.104873.

Hayama, Y., T. Yamamoto, S. Kobayashi, N. Muroga, and T. Tsutsui, 2013: Mathematical model

of the 2010 foot-and-mouth disease epidemic in Japan and evaluation of control measures.

Prev. Vet. Med. **112**, 183–193, DOI: 10.1016/j.prevetmed.2013.08.010.

Henao-Diaz, A., J. Ji, L. Giménez-Lirola, D.H. Baum, and J. Zimmerman, 2020: Understanding

and interpreting PRRSV diagnostics in the context of "disease transition stages." *Res. Vet.*

Sci. **131**, 173–176, DOI: 10.1016/j.rvsc.2020.04.023.

Hermann, J., S. Hoff, C. Muñoz-Zanzi, K.-J. Yoon, M. Roof, A. Burkhardt, and J. Zimmerman,

2007: Effect of temperature and relative humidity on the stability of infectious porcine

reproductive and respiratory syndrome virus in aerosols. *Vet. Res.* **38**, 81–93, DOI:

10.1051/vetres:2006044.

Holtkamp, D.J., J.B. Kliebenstein, E.J. Neumann, J.J. Zimmerman, H.F. Rotto, T.K. Yoder, C.

Wang, P.E. Yeske, C.L. Mowrer, and C.A. Haley, 2013: Assessment of the economic

impact of porcine reproductive and respiratory syndrome virus on United States pork

producers. *J. Swine Heal. Prod.* **21**, 72–84.

Holtkamp, D.J., B. Morrison, D.M. Classen, L. Becton, S. Henry, M.T. Rodibaugh, R.R.

Rowland, H. Snelson, B. Straw, P. Yeske, and J. Zimmerman, 2011: Terminology for

Classifying Swine Herds by PRRS Status. *J. Swine Heal. Prod.* **19**, 44–56.

Jara, M., D.A. Rasmussen, C.A. Corzo, and G. Machado, 2020: Porcine reproductive and

respiratory syndrome virus dissemination across pig production systems in the United

States. *Transbound. Emerg. Dis.* **43**, tbed.13728, DOI: 10.1111/tbed.13728.

Jurado, C., L. Mur, M.S. Pérez Aguirreburualde, E. Cadenas-Fernández, B. Martínez-López, J.M.

Sánchez-Vizcaíno, and A. Perez, 2019: Risk of African swine fever virus introduction into

the United States through smuggling of pork in air passenger luggage. *Sci. Rep.* **9**, 14423,

DOI: 10.1038/s41598-019-50403-w.

Kick, A., A. Amaral, L. Cortes, J. Fogle, E. Crisci, G. Almond, and T. Käser, 2019: The T-Cell

Response to Type 2 Porcine Reproductive and Respiratory Syndrome Virus (PRRSV).

Viruses **11**, 796, DOI: 10.3390/v11090796.

Kroll, J., M. Piontkowski, C. Kraft, T. Coll, and O. Gomez-Duran, 2018: Initial vaccination and

revaccination with Type I PRRS 94881 MLV reduces viral load and infection with porcine

reproductive and respiratory syndrome virus. *Porc. Heal. Manag.* **4**, 1–8, DOI:

10.1186/s40813-018-0096-3.

Lee, K., D. Polson, E. Lowe, R. Main, D. Holtkamp, and B. Martínez-López, 2017: Unraveling

the contact patterns and network structure of pig shipments in the United States and its

association with porcine reproductive and respiratory syndrome virus (PRRSV) outbreaks.

Prev. Vet. Med. **138**, 113–123, DOI: 10.1016/j.prevetmed.2017.02.001.

Linhares, D.C.L., J.P. Cano, T. Wetzell, J. Nerem, M. Torremorell, and S.A. Dee, 2012: Effect of

modified-live porcine reproductive and respiratory syndrome virus (PRRSv) vaccine on the

shedding of wild-type virus from an infected population of growing pigs. *Vaccine* **30**, 407–

413, DOI: 10.1016/j.vaccine.2011.10.075.

Linhares, D.C.L., C. Johnson, and R.B. Morrison, 2015: Economic Analysis of Vaccination

Strategies for PRRS Control. (Frederick C. Leung, Ed.)*PLoS One* **10**, e0144265, DOI:

10.1371/journal.pone.0144265.

Lowe, J., P. Gauger, K. Harmon, J. Zhang, J. Connor, P. Yeske, T. Loula, I. Levis, L. Dufresne,

and R. Main, 2014: Role of Transportation in Spread of Porcine Epidemic Diarrhea Virus

Infection, United States. *Emerg. Infect. Dis.* **20**, 872–874, DOI: 10.3201/eid2005.131628.

Machado, G., C. Vilalta, M. Recamonde-Mendoza, C. Corzo, M. Torremorell, A. Perez, and K.

VanderWaal, 2019: Identifying outbreaks of Porcine Epidemic Diarrhea virus through

animal movements and spatial neighborhoods. *Sci. Rep.* **9**, 1–12, DOI: 10.1038/s41598-018-

36934-8.

Miller, J., 2019: Decreasing PRRS incidence: A Production System View. p. 17. In: *2019 Allen*

D. Leman Swine Conf. Saint Paul, Minnesota.

Nathues, H., P. Alarcon, J. Rushton, R. Jolie, K. Fiebig, M. Jimenez, V. Geurts, and C. Nathues,

2017: Cost of porcine reproductive and respiratory syndrome virus at individual farm level –

An economic disease model. *Prev. Vet. Med.* **142**, 16–29, DOI:

10.1016/j.prevetmed.2017.04.006.

Neira, V., B. Brito, J. Mena, M. Culhane, M.I. Apel, V. Max, P. Perez, V. Moreno, C. Mathieu,

M. Johow, C. Badia, M. Torremorell, R. Medina, and R. Ortega, 2017: Epidemiological

investigations of the introduction of porcine reproductive and respiratory syndrome virus in

Chile, 2013-2015. (Sanjay B. Jadhao, Ed.)*PLoS One* **12**, e0181569, DOI:

10.1371/journal.pone.0181569.

Otake, S., S. Dee, C. Corzo, S. Oliveira, and J. Deen, 2010: Long-distance airborne transport of

infectious PRRSV and Mycoplasma hyopneumoniae from a swine population infected with

multiple viral variants. *Vet. Microbiol.* **145**, 198–208, DOI: 10.1016/j.vetmic.2010.03.028.

Passafaro, T.L., A.F.A. Fernandes, B.D. Valente, N.H. Williams, and G.J.M. Rosa, 2020:

Network analysis of swine movements in a multi-site pig production system in Iowa, USA.

Prev. Vet. Med. **174**, 104856, DOI: 10.1016/j.prevetmed.2019.104856.

Perez, A.M., D.C.L. Linhares, A.G. Arruda, K. Van Der Waal, G. Machado, C. Vilalta, J.M.

Sanhueza, J. Torrison, M. Torremorell, and C.A. Corzo, 2019: Individual or common good?

Voluntary data sharing to inform disease surveillance systems in food animals. *Front. Vet.*

Sci. **6**, 1–7, DOI: 10.3389/fvets.2019.00194.

Phoo-ngurn, P., C. Kiataramkul, and F. Chamchod, 2019: Modeling the spread of porcine

reproductive and respiratory syndrome virus (PRRSV) in a swine population: transmission

dynamics, immunity information, and optimal control strategies. *Adv. Differ. Equations*

2019, 432, DOI: 10.1186/s13662-019-2351-6.

Pileri, E., E. Gibert, F. Soldevila, A. García-Saenz, J. Pujols, I. Diaz, L. Darwich, J. Casal, M.

Martín, and E. Mateu, 2015: Vaccination with a genotype 1 modified live vaccine against

porcine reproductive and respiratory syndrome virus significantly reduces viremia, viral

shedding and transmission of the virus in a quasi-natural experimental model. *Vet.*

Microbiol. **175**, 7–16, DOI: 10.1016/j.vetmic.2014.11.007.

Pileri, E., and E. Mateu, 2016: Review on the transmission porcine reproductive and respiratory

syndrome virus between pigs and farms and impact on vaccination. *Vet. Res.* **47**, 108, DOI:

10.1186/s13567-016-0391-4.

Pitkin, A., J. Deen, and S. Dee, 2009: Further assessment of fomites and personnel as vehicles for

the mechanical transport and transmission of porcine reproductive and respiratory syndrome

virus. *Can. J. Vet. Res.* **73**, 298–302.

Pitkin, A., J. Deen, S. Otake, R. Moon, and S. Dee, 2009: Further assessment of houseflies

(Musca domestica) as vectors for the mechanical transport and transmission of porcine

reproductive and respiratory syndrome virus under field conditions. *Can. J. Vet. Res.* **73**,

91–96.

Porphyre, T., B.M. de C. Bronsvoort, G.J. Gunn, and C. Correia-Gomes, 2020: Multilayer

network analysis unravels haulage vehicles as a hidden threat to the British swine industry.

Transbound. Emerg. Dis. **67**, 1231–1246, DOI: 10.1111/tbed.13459.

Rawal, G., C. Rademacher, and D. Linhares, 2020: Comparison of productivity losses between

swine breeding herds adopting killed or attenuated PRRS virus vaccination protocols

following PRRS outbreaks. *Vet. Diagnostic Prod. Anim. Med. Publ.* **175**.

Reiner, G., C. Fresen, S. Bronnert, and H. Willems, 2009: Porcine Reproductive and Respiratory

Syndrome Virus (PRRSV) infection in wild boars. *Vet. Microbiol.* **136**, 250–258, DOI:

10.1016/j.vetmic.2008.11.023.

Roca, M., M. Gimeno, S. Bruguera, J. Segalés, I. Díaz, I.J. Galindo-Cardiel, E. Martínez, L.

Darwich, Y. Fang, J. Maldonado, R. March, and E. Mateu, 2012: Effects of challenge with a

virulent genotype II strain of porcine reproductive and respiratory syndrome virus on piglets

vaccinated with an attenuated genotype I strain vaccine. *Vet. J.* **193**, 92–96, DOI:

10.1016/j.tvjl.2011.11.019.

Rosendal, T., C. Dewey, R. Friendship, S. Wootton, B. Young, and Z. Poljak, 2014: Spatial and

temporal patterns of porcine reproductive and respiratory syndrome virus (PRRSV)

genotypes in Ontario, Canada, 2004-2007. *BMC Vet. Res.* **10**, DOI: 10.1186/1746-6148-10-

83.

Salines, M., M. Andraud, N. Rose, and S. Widgren, 2020: A between-herd data-driven stochastic

model to explore the spatio-temporal spread of hepatitis E virus in the French pig

production network. (Laurent Pujo-Menjouet, Ed.)*PLoS One* **15**, e0230257, DOI:

10.1371/journal.pone.0230257.

Sanhueza, J.M., M.A. Stevenson, C. Vilalta, M. Kikuti, and C.A. Corzo, 2020: Spatial relative

risk and factors associated with porcine reproductive and respiratory syndrome outbreaks in

United States breeding herds. *Prev. Vet. Med.*105128, DOI:

10.1016/j.prevetmed.2020.105128.

Sanhueza, J.M., C. Vilalta, C. Corzo, and A.G. Arruda, 2019: Factors affecting Porcine

Reproductive and Respiratory Syndrome virus time-to-stability in breeding herds in the

Midwestern United States. *Transbound. Emerg. Dis.* **66**, 823–830, DOI:

10.1111/tbed.13091.

Silva, G.S., G. Machado, K.L. Baker, D.J. Holtkamp, and D.C.L. Linhares, 2019: Machine-

learning algorithms to identify key biosecurity practices and factors associated with

breeding herds reporting PRRS outbreak. *Prev. Vet. Med.* **171**, 104749, DOI:

10.1016/j.prevetmed.2019.104749.

Stevenson, G.W., W.G. Van Alstine, C.L. Kanitz, and K.K. Keffaber, 1993: Endemic Porcine

Reproductive and Respiratory Syndrome Virus Infection of Nursery Pigs in Two Swine

Herds without Current Reproductive Failure. *J. Vet. Diagnostic Investig.* **5**, 432–434, DOI:

10.1177/104063879300500322.

Sumner, T., R.J. Orton, D.M. Green, R.R. Kao, and S. Gubbins, 2017: Quantifying the roles of

host movement and vector dispersal in the transmission of vector-borne diseases of

livestock. (James Lloyd-Smith, Ed.)*PLOS Comput. Biol.* **13**, e1005470, DOI:

10.1371/journal.pcbi.1005470.

Thakur, K.K., C.W. Revie, D. Hurnik, Z. Poljak, and J. Sanchez, 2015: Simulation of between-

farm transmission of porcine reproductive and respiratory syndrome virus in Ontario,

Canada using the North American Animal Disease Spread Model. *Prev. Vet. Med.* **118**,

413–426, DOI: 10.1016/j.prevetmed.2015.01.006.

Thakur, K.K., J. Sanchez, D. Hurnik, Z. Poljak, S. Opps, and C.W. Revie, 2015: Development of

a network based model to simulate the between-farm transmission of the porcine

reproductive and respiratory syndrome virus. *Vet. Microbiol.* **180**, 212–222, DOI:

10.1016/j.vetmic.2015.09.010.

Thomann, B., J. Rushton, G. Schuepbach-Regula, and H. Nathues, 2020: Modeling Economic

Effects of Vaccination Against Porcine Reproductive and Respiratory Syndrome: Impact of

Vaccination Effectiveness, Vaccine Price, and Vaccination Coverage. *Front. Vet. Sci.* **7**,

DOI: 10.3389/fvets.2020.00500.

Thomas, P., 2019: Iowa Select Farms – PRRS Incidence and Reduction. p. 11. In: *2019 Allen D.*

Leman Swine Conf. Saint Paul, Minnesota.

Trevisan, G., L.C.M. Linhares, B. Crim, P. Dubey, K.J. Schwartz, E.R. Burrough, C. Wang, R.G.

Main, P. Sundberg, M. Thurn, P.T.F. Lages, C.A. Corzo, J. Torrison, J. Henningson, E.

Herrman, G.A. Hanzlicek, R. Raghavan, D. Marthaler, J. Greseth, T. Clement, J.

Christopher-Hennings, D. Muscatello, and D.C.L. Linhares, 2020: Prediction of seasonal

patterns of porcine reproductive and respiratory syndrome virus RNA detection in the U.S.

swine industry. *J. Vet. Diagnostic Investig.* **32**, 394–400, DOI: 10.1177/1040638720912406.

USDA, 2016: Swine 2012 Part II: Reference of Swine Health and Health Management in the

United States, 2012 Vol. 2012*USDA*. Washington, D.C.

Van Ryswyk, K., N. Prince, M. Ahmed, E. Brisson, J.D. Miller, and P.J. Villeneuve, 2019: Does

urban vegetation reduce temperature and air pollution concentrations? Findings from an

environmental monitoring study of the Central Experimental Farm in Ottawa, Canada.

Atmos. Environ. **218**, 116886, DOI: 10.1016/j.atmosenv.2019.116886.

VanderWaal, K., I.A.D. Paploski, D.N. Makau, and C.A. Corzo, 2020: Contrasting animal

movement and spatial connectivity networks in shaping transmission pathways of a

genetically diverse virus. *Prev. Vet. Med.* **178**, 104977, DOI:

10.1016/j.prevetmed.2020.104977.

VanderWaal, K., A. Perez, M. Torremorell, R.M. Morrison, and M. Craft, 2018: Role of animal

movement and indirect contact among farms in transmission of porcine epidemic diarrhea

virus. *Epidemics* **24**, 67–75, DOI: 10.1016/j.epidem.2018.04.001.

Velasova, M., P. Alarcon, S. Williamson, and B. Wieland, 2012: Risk factors for porcine

reproductive and respiratory syndrome virus infection and resulting challenges for effective

disease surveillance. *BMC Vet. Res.* **8**, 1, DOI: 10.1186/1746-6148-8-184.

Vilalta, C., J. Sanhueza, J. Garrido, D. Murray, R. Morrison, C.A. Corzo, and M. Torremorell,

2019: Indirect assessment of porcine reproductive and respiratory syndrome virus status in

pigs prior to weaning by sampling sows and the environment. *Vet. Microbiol.* **237**, 108406,

DOI: 10.1016/j.vetmic.2019.108406.

Widgren, S., S. Engblom, U. Emanuelson, and A. Lindberg, 2018: Spatio-temporal modelling of

verotoxigenic Escherichia coli O157 in cattle in Sweden: exploring options for control. *Vet.*

Res. **49**, 78, DOI: 10.1186/s13567-018-0574-2.

Table list

Table 1. Transmission parameters used in simulations

Model parameters	Parameter	Value	Details & references
Between-farm pig movements (transmission rate)	βnet	0.26	ABC fitting
Local transmission (transmission rate)	$\beta local$	0.003	ABC fitting
Re-break for sow farms (transmission rate)	$\beta rebreak$	0.0019	ABC fitting
On-farm biosecurity levels	H (sow)	0.47	ABC fitting
	H (GDU)	0.38	ABC fitting
Detection effectiveness	L (nursery, GDU and finisher)	0.07	ABC fitting
	L (sow)	0.95	Expert opinion
PRRSV seasonality	T	Weekly values calculated	(supplementary material section 1)
Average time for PRRSV detection	X0	4 weeks	Expert opinion
Average infectious time for sow farms	-	41 weeks	(Holtkamp et al., 2011)

865 **Table 2. Vaccination strategies**

Vaccination strategy*	Vaccine deployment description
Prophylactic #1	Pigs sent from sow farms into nurseries and/or into wean-to-finisher farms, were vaccinated
Prophylactic #2	Pigs sent from either sow, nursery or finisher farms into nurseries, finisher, and replacement gilt units farms, were vaccinated (i.e. a finisher farm (re)vaccinate when receive pigs from a nursery)
Reactive #1	Whole-herd vaccination at infected and detected sow farms

| Reactive #2 | Whole-herd vaccination at any infected and detected farm, independent of the production type |
| Combined | Farms implement both prophylactic #2 and reactive #2 strategies |

866 *modeling assumed that all pig housed at vaccination were successfully vaccinated

867 **Table 3.** The summary of susceptible and infected farms by production type from 2009 to 2019.

Production type	Number of susceptible farms	Number of infected farms 2009-2019	Number of infected farms Jun-Dec 2019
Sow	319 (14%)	1070	23
Gilt development unit	187 (8.2%)	50	0
Nursery	468 (20.4%)	648	17
Finisher	1272 (55.4%)	537	8
Isolation	33 (1.4%)	0	0
Boar stud	15 (0.6%)	0	0

Figure Legends

Figure 1. The flow of infection stages in the farm's modeling framework. Each farm was classified

to either state: susceptible (S), infected undetected (U) or infected detected (D).

Figure. 2. Force of infection for each transmission route of each production type. The y-axis

represents the proportion of transmission by transmission routes for each of the simulated weeks

(x-axis). Weekly rate indices were calculated by dividing the number of simulated infected farms

for each transmission route by the number of simulated infected farms by the three routes

combined.

Figure 3. The simulated weekly number of A) infected farms and B) infected and detected

farms. The black line represents the median, the dark shade areas represent a 75% credible interval

and the light shade areas maximum and minimum generated by the model, and the red dots the

frequency of true outbreaks reported in our data. Uncertainty in the estimated model parameters is

reflected by 1000 repeated simulations.

Figure 4. Estimated reduction of PRRSV infected farm trajectories under five vaccine

efficacy scenarios. A) The total number of prevented infected farms and B) the number of

prevented infected farms depicted by production type. The y-axis represents the number of infected

farms within the 27 weeks and the x-axis the simulated vaccine efficacies. Among the prophylactics

strategies farms were vaccinated in the same week of receiving a pig movement and among the

reactive strategies vaccination occurred in the same week a farm reported an outbreak. The line

plot represents one upper and lower standard deviation and the median value represented as the

symbol in the center of the line, values calculated from 50 simulations by each vaccination strategy.

Figure 5. Predicted spatial distribution of PRRSV in sow farms for one target week

(November 03-09, 2019). A) cells with observed sow farms infected; B) estimated risk of PRRSV

detection over 100 model simulations; C) cells with risk for PRRSV detection above the 90th

percentile; D) estimated risk of PRRSV recent introduction; and E) cells with risk for PRRSV

introduction above the 90th percentile .

PRRSV detection forecast

A. Observed outbreaks

Status
- New case (red)
- Not reported (grey)

B. Risk of detection

Risk
- Maximum
- Minimum

C. Higher risk (risk > 90%)

Risk >90%
- High risk (red)
- Low risk (grey)

PRRSV introduction forecast

D. Risk of introduction

Risk
- Maximum
- Minimum

E. Higher risk (risk > 90%)

Risk >90%
- High risk (red)
- Low risk (grey)