Emergence and Propagation of Epistasis in Metabolic Networks

Sergey Kryazhimskiy[1*]

*For correspondence:
skryazhi@ucsd.edu (SK)

[1]Division of Biological Sciences, University of California San Diego, La Jolla, CA

Abstract   Epistasis is often used to probe functional relationships between genes, and it plays an important role in evolution. However, we lack theory to understand how functional relationships at the molecular level translate into epistasis at the level of whole-organism phenotypes, such as fitness. Here, I derive two rules for how epistasis between mutations with small effects propagates from lower- to higher-level phenotypes in a hierarchical metabolic network with first-order kinetics and how such epistasis depends on topology. Most importantly, weak epistasis at a lower level may be distorted as it propagates to higher levels. Computational analyses show that epistasis in more realistic models likely follows similar, albeit more complex, patterns. These results suggest that pairwise inter-gene epistasis should be common and it should generically depend on the genetic background and environment. Furthermore, the epistasis coefficients measured for high-level phenotypes may not be sufficient to fully infer the underlying functional relationships.

Introduction

Life emerges from an orchestrated performance of complex regulatory and metabolic networks within cells. The blueprint for these networks is encoded in the genome. Mutations alter the genome. Some of them, once decoded by the cell, perturb cellular networks and thereby change the phenotypes important for life. Understanding how mutations affect the function of cellular networks is key to solving many practical and fundamental problems, such as finding mechanistic causes of genetic disorders (*Hu et al., 2011*; *Fang et al., 2019*), deciphering the architecture of complex traits (*Zuk et al., 2012*; *Mackay, 2014*; *Wei et al., 2014*), building artificial cells (*Hutchison et al., 2016*), explaining past and predicting future evolution (*Blount et al., 2008*; *Wiser et al., 2013*; *De Visser and Krug, 2014*; *Harms and Thornton, 2014*; *Kryazhimskiy et al., 2014*; *Sailer and Harms, 2017a*; *Sohail et al., 2017*). Conversely, mutations can help us learn how cellular networks are organized (*Phillips, 2008*; *Van Opijnen and Camilli, 2013*).

To infer the wiring diagram of a cellular network that produces a certain phenotype, one approach in genetics is to measure the pairwise and higher-order genetic interactions (or "epistasis") between mutations that perturb it (*Phillips, 2008*). Much effort has been devoted in the past 20 years to a systematic collection of such genetic interaction data for several model organisms and cell lines (*Kelley and Ideker, 2005*; *Lehner et al., 2006*; *Jasnos and Korona, 2007*; *Collins et al., 2007*; *St Onge et al., 2007*; *Typas et al., 2008*; *Roguev et al., 2008*; *Costanzo et al., 2010*; *Szappanos et al., 2011*; *Huang et al., 2012*; *Roguev et al., 2013*; *Bassik et al., 2013*; *Babu et al., 2014*; *Costanzo et al., 2016*; *Van Leeuwen et al., 2016*; *Skwark et al., 2017*; *Du et al., 2017*; *Heigwer et al., 2018*; *Horlbeck et al., 2018*; *Norman et al., 2019*; *Liu et al., 2019*; *Kuzmin et al., 2018*; *New and Lehner, 2019*; *Celaj et al., 2020*). This approach is particulary powerful when the phenotypic effect of one mutation changes qualitatively depending on the presence or absence of a second mutation in another gene, for example when a mutation has no effect on the phenotype in the wildtype background, but abolishes the phenotype when introduced together with another mutation, such as synthetic

lethality (*Tong et al., 2001*). Such qualitative genetic interactions can often be directly interpreted in terms of a functional relationship between gene products (*Tong et al., 2001*; *Davierwala et al., 2005*; *Phillips, 2008*).

Most pairs of mutations do not exhibit qualitative genetic interactions. Instead, the phenotypic effect of a mutation may change measurably but not qualitatively depending on the presence or absence of other mutations in the genome (*Babu et al., 2014*; *Costanzo et al., 2016*). The genetic interactions can in this case be quantified with one of several metrics that are termed "epistasis coefficients" (*Wagner et al., 1998*; *Hansen and Wagner, 2001*; *Mani et al., 2008*; *Wagner, 2015*; *Poelwijk et al., 2016*). While some rules have been proposed for interpreting epistasis coefficients, in particular, their sign (*Dixon et al., 2009*; *Lehner, 2011*; *Baryshnikova et al., 2013*), the validity and robustness of these rules are unknown because there is no theory for how functional relationships translate into measurable epistasis coefficients in any system (*Lehner, 2011*; *Domingo et al., 2019*). To avoid this major difficulty, most large-scale empirical studies focus on correlations between epistasis coefficients rather than on their actual values (but see *Velenich and Gore, 2013*, for a notable exception). Genes with highly correlated epistasis profiles are then interpreted as being functionally related (*Segre et al., 2005*; *Bellay et al., 2011*; *Babu et al., 2014*; *Costanzo et al., 2016*; *Horlbeck et al., 2018*). Although this approach successfully groups genes into protein complexes and larger functional modules (*Michaut et al., 2011*; *Bellay et al., 2011*), it does not reveal the functional relationships themselves. As a result, many if not most, genetic interactions between genes and modules still await their biological interpretation (*Costanzo et al., 2016*; *Fang et al., 2019*).

While geneticists measure epistasis to learn the architecture of biological networks, evolutionary biologists face the reverse problem: they need to know how the genetic architecture constrains epistasis at the level of fitness. Epistasis determines the structure of fitness landscapes on which populations evolve (*Fragata et al., 2019*). Understanding it would bear on many important evolutionary questions, such as why so many organisms reproduce sexually (*Kondrashov, 2018*), how novel phenotypes evolve (*Blount et al., 2008*; *Bridgham et al., 2009*; *Natarajan et al., 2013*; *Harms and Thornton, 2014*), how predictable evolution is (*Weinreich et al., 2006*; *Tenaillon et al., 2012*; *Wiser et al., 2013*; *Kryazhimskiy et al., 2014*), etc. So far, evolutionary biologists have relied primarily on abstract models of fitness landscapes (see *Orr, 2005*, for a review), rather than those firmly grounded in organismal biochemistry and physiology (e.g., *Dykhuizen et al., 1987*; *Das et al., 2020*). For example, Fisher's geometric model—one of the most widely used fitness landscape models—is explicitly devoid of the physiological and biochemical details (*Fisher, 1930*; *Tenaillon, 2014*; *Martin, 2014*).

A theory of epistasis must address two challenges. First, it must specify how the architecture of a biological network constrains epistasis. Such knowledge is important not only for evolutionary questions, but also for the inference problem in genetics. Consider a biological network module that produces a phenotype of interest but whose internal structure is unknown. By genetically perturbing all genes within the module and measuring the phenotype in all single, double and possibly some higher-order mutants, we can obtain the matrix of epistasis coefficients. In principle, we can then fit a network topology and parameters to these data. However, without knowing what information about the network is contained in the matrix in the first place, we cannot be sure whether the inferred topology and parameters are close to their true values or represent one of many possible solutions consistent with the data.

The second challenge is that epistasis may arise at a different level of biological organization than where it is measured by the experimentalist or by natural selection. For example, geneticists are often interested in understanding the structures of specific regulatory or metabolic network modules (*Collins et al., 2007*; *Costanzo et al., 2010*). However, measuring the peformance of a module directly is often experimentally difficult or impossible. Then epistasis is measured for an experimentally accessible "high-level" phenotype, such as fitness, which depends on the performance of the focal "lower-level" module, but also on other unrelated modules. However, if

we do not know how epistasis that originally emerged in one module maps onto epistasis that is measured, it is unclear what we can infer about module's internal structure.

Evolutionary biologists encounter a related problem when they wish to learn the evolutionary history of a protein or a larger cellular module. To do so, they would in principle need to know how different mutations in this module affected fitness of the whole organism in its past environment. But such information is rarely available. Instead, it is sometimes possible to reconstruct past mutations and measure their biochemical effects in the lab (*Lunzer et al., 2005*; *Bridgham et al., 2009*; *Natarajan et al., 2013*; *Sarkisyan et al., 2016*). When interesting patterns of epistasis are identified at the biochemical level, it is usually assumed that the same patterns manifested themselves at the level of fitness and drove module's evolution. However, this is not obvious. If interactions with other modules distort epistasis as it propagates from the biochemical level to the level of fitness (*Snitkin and Segrè, 2011*), our ability to infer past evolutionary history from in vitro biochemical measurements could be diminished. Therefore, the second challenge that a theory of epistasis must address is how epistasis propagates from lower-level phenotypes to higher-level phenotypes.

There is a large body of theoretical and computational literature on epistasis. As early as 1934, Sewall Wright realized that epistasis naturally emerges in molecular networks (*Wright, 1934*). This was later explicitly demonstrated in many mathematical and computational models (e.g., *Kacser and Burns, 1981*; *Keightley, 1989*; *Szathmary, 1993*; *Gibson, 1996*; *Keightley, 1996*; *Wagner et al., 1998*; *Omholt et al., 2000*; *Peccoud et al., 2004*; *Gjuvsland et al., 2007*; *Gertz et al., 2010*; *Fiévet et al., 2010*; *Pumir and Shraiman, 2011*). Metabolic control analysis became one of the most successful and general frameworks for understanding epistasis between metabolic genes (*Kacser and Burns, 1973*). *Dean et al.* (*1986*), *Dykhuizen et al.* (*1987*), *Dean* (*1989*), *Lunzer et al.* (*2005*), *MacLean* (*2010*) used it to interpret the empirically measured fitness effects of mutations and their interactions in terms of the metabolic relationships between the products of mutated genes. *Kacser and Burns* (*1981*), *Hartl et al.* (*1985*), *Keightley* (*1989*), *Clark* (*1991*), *Keightley* (*1996*), *Bagheri-Chaichian et al.* (*2003*), *Bagheri and Wagner* (*2004*), *Fiévet et al.* (*2010*) explored the implications of epistasis in metabolism for genetic variation in populations, their response to selection, long-term evolutionary dynamics and outcomes, such as the evolution of dominance. However, most studies analyzed only the linear metabolic pathway (but see *Keightley, 1989*, *1996*) and assumed that fitness equals flux through the pathway (but see *Szathmary, 1993*), thereby bypassing the problem of epistasis propagation.

There have been few attempts to theoretically relate the molecular architecture of an organism to the types of epistasis that would arise for its high-level phenotypes, such as fitness. *Segre et al.* (*2005*) and *He et al.* (*2010*) used flux balance analysis (FBA, *Orth et al., 2010*) to compute genome-wide distributions of epistasis coefficients in metabolic models of *Escherichia coli* and *Saccharomyces cerevisiae* and arrived at starkly discordant conclusions. Recently, *Alzoubi et al.* (*2019*) showed that FBA is generally poor in predicting experimentally measured genetic interactions, suggesting that it might be difficult to understand the emergence and propagation of epistasis by relying exclusively on genome-scale computational models. *Sanjuán and Nebot* (*2008*) and *Macía et al.* (*2012*) modelled various abstract metabolic and regulatory networks and found a possible link between epistasis and network complexity. The work by *Chiu et al.* (*2012*) is a more systematic attempt to develop a general theory of epistasis. They established a fundamental connection between epistasis and the curvature of the function that maps lower-level phenotypes onto a higher-level phenotype. However, further progress has been so far hindered by uncertainty in what types of functions map phenotypes onto one another in real biological systems. Previous studies made various idiosyncratic choices with respect to this mapping, leaving us without a clear guidance as to the conditions or systems where they are expected to hold.

To overcome this problem, here I consider a whole class of hierarchical metabolic networks and obtain the family of all functions that determine how the effective activity of a larger metabolic module can depend on the activities of smaller constituent modules. There are several advantages to this approach. First, it leads to an intuitive understanding of how the structure of the network

influences epistasis emergence and propagation. Second, my approach is based on basic bio-chemical principles, so it should be relevant for many phenotypes. For example, epistasis is often measured at the level of growth rate (*Jasnos and Korona, 2007*; *St Onge et al., 2007*; *Babu et al., 2014*; *Costanzo et al., 2016*), and metabolism fuels growth. Moreover, metabolic genes occupy a large fraction of most genomes (*Orth et al., 2011*) and the general organization of metabolism is conserved throughout life (*Csete and Doyle, 2004*). Thus, by understanding genetic interactions between metabolic genes, we will gain an understanding of a large fraction of all genetic interactions.

In my model, I consider a hierarchical network with first-order kinetics but arbitrary topology, and ask two questions related to the two challenges mentioned above. (1) How does an epistasis coefficient that arose at some level of the metabolic hierarchy propagate to higher levels of the hierarchy? (2) How does the network topology constrain the value of an epistasis coefficient between two mutations that affect different enzymes in this network? I obtain answers to these questions analytically in the limiting case when the effects of mutations are vanishingly small. I then computationally probe the validity of the conclusions outside of the domain where they are expected to hold.

My model is not intended to generate predictions of epistasis for any specific organism. Instead, its main purpose is to provide a baseline expectation for how epistasis that emerges at lower-level phenotypes manifests itself at higher-level whole-organism phenotypes, such as fitness, and what kind of information may be gained from measurements of such higher-level epistasis. One possible outcome of this analysis is that there may be fundamental limitations to what an epistasis measurement at one level of biological organization can tell us about epistasis at another level. On the other hand, if it turns out that there there is a general correspondence between epistasis coefficients at different levels in this simple model, then it may be worth developing more sophisticated and general models on which inference from data can be based.

Model

Hierarchical metabolic network

Consider a set of metabolites $A = \{1, 2, \ldots, n\}$ with concentrations $S_1, \ldots, S_n$ which can be interconverted by reversible first-order biochemical reactions. The rate of the reaction converting metabolite i into metabolite j is $x_{ij} \left(S_i - S_j / K_{ij} \right)$ where K_{ij} is the equilibrium constant. The rate constants x_{ij}, which satisfy the Haldane relationships $x_{ji} = x_{ij} / K_{ij}$ (*Cornish-Bowden, 2013*), form the matrix $x = \left\| x_{ij} \right\|_{i,j=1}^{n}$. The metabolite set A and the rate matrix x define a biochemical network $\mathcal{N} = (A, x)$.

The first-order kinetics assumption makes the model analytically tractable, as discussed below; biologically, it is equivalent to assuming that all enzymes are far from saturation. The rate constants x_{ij} depend on the concentrations and the specific activities of enzymes and therefore can be altered by mutations. K_{ij} characterize the fundamental chemical nature of metabolites i and j and cannot be altered by mutations (*Savageau, 1976*).

The whole-cell metabolic network is large, and it is often useful to divide it into subnetworks that carry out certain functions important for the cell. I define subnetworks mathematically as follows. I say that two metabolites i and j are *adjacent* (in the graph-theoretic sense) if there exists an enzyme that catalyzes a biochemical reaction between them, i.e., if $x_{ij} > 0$. Now consider a subset of metabolites $A_\mu \subset A$. For this subset, let A_μ^{IO} be the set of all metabolites that do not belong to A_μ but are adjacent to at least one metabolite from A_μ. Let x_μ be the submatrix of x which corresponds to all reactions where both the product and the substrate belong to either A_μ or A_μ^{IO}. The metabolite subset A_μ and the rate matrix x_μ form a *subnetwork* $\mu = (A_\mu, x_\mu)$ of network $\mathcal{N}$. I refer to A_μ and A_μ^{IO} as the sets of internal and "input/output" ("I/O" for short) metabolites for subnetwork μ, respectively. Thus, all internal metabolites and all reactions that involve only internal and I/O metabolites are part of the subnetwork. Note that the I/O metabolites do not themselves belong to the subnetwork, but reactions between them, if they exist, are part of the subnetwork.

Figure 1. Illustration of a hierarchical metabolic network and its coarse-graining. (**A**) White rectangle represents the whole metabolic network $\mathcal{N}$. Example subnetworks μ and ν are represented by the dark and light gray rectangles. Only metabolites and reactions that belong to these subnetworks are shown; other metabolites and reactions in $\mathcal{N}$ are not shown. Metabolites 1 and 5 may be adjacent to other metabolites in $\mathcal{N}$; this fact is represented by short black lines that do not terminate in metabolites. Subnetworks μ and ν are both modules because there exists a simple path connecting their I/O metabolites that lies within μ and ν and contains all their internal metabolites (dashed blue line). (**B**) Network $\mathcal{N}$ can be coarse-grained by replacing module μ at steady state with an effective reaction between its I/O metabolites 1 and 2, with the rate constant is y_μ. (**C**) Network $\mathcal{N}$ can be coarse-grained by replacing module ν at steady state with an effective reaction between its I/O metabolites 1 and 5, with the rate constant is y_ν.

Metabolites that are neither internal nor I/O for μ are referred to as *external to subnetwork μ*. These
definitions are illustrated in *Figure 1*A.

The main objects in this work are biochemical modules, which are a special type of subnetworks.
To define modules, I introduce some auxiliary concepts. I say that two metabolites i and j are
connected if there exists a series of enzymes that interconvert i and j, possibly through a series
of intermediates. Mathematically, i and j are connected if there exists a simple (i.e., non-self-
intersecting) path between them. If all metabolites in this path are internal to the subnetwork μ
(possibly excluding the terminal metabolites i and j themselves) then i and j are *connected within*
the subnetwork μ, and such path is said to lie *within μ*. By this definition, metabolites i and j can be
connected within μ only if they are either internal or I/O metabolites for μ.

Definition 1. *A subnetwork μ is called a* module *if (a) it has two I/O metabolites, and (b) for every*
internal metabolite $i \in A_\mu$, there exists a simple path between the I/O metabolites that lies within μ and
contains i.

This definition is illustrated in *Figure 1*A. The assumption that modules only have two I/O
metabolites is not essential. However, mathematical calculations become unwieldy when the
number of I/O metabolites increases. Moreover, modules with just two I/O metabolites already
capture two most salient features of metabolism: its directionality, and its complex branched
topology (*Csete and Doyle, 2004*). Such modules are a natural generalization of the linear metabolic
pathway which has been extensively studied in the previous literature (*Kacser and Burns, 1973*;
Szathmary, 1993; *Bagheri-Chaichian et al., 2003*; *MacLean, 2010*).

Modules have two important properties. First, for any given concentrations of the two I/O
metabolites, all internal metabolites in the module can achieve a unique steady state which de-
pends only on concentrations of these I/O metabolites but not on the concentrations of any other
metabolites in the network (see *Proposition 1* in Materials and Methods). Now consider a module
μ whose I/O metabolites are (without loss of generality) labelled 1 and 2 (*Figure 1*A). The second
property is that, at steady state, the flux through this module is $J_\mu = y_\mu \left(S_1 - S_2/K_{12} \right)$, where

$$y_\mu = F\left(\boldsymbol{x}_\mu\right) \tag{1}$$

220 is the effective reaction rate constant of module μ (*Figure 1*B). Importantly, y_μ depends only on
221 the rate matrix $\boldsymbol{x}_\mu$, but not on any other rate constants (see *Corollary 2* in Materials and Methods),
222 and it can be recursively computed for any module, as described in Network coarse graining and

an algorithm for evaluating the effective rate constant for an arbitrary module in Materials and Methods. In other words, metabolic network $\mathcal{N}$ can be coarse-grained by replacing module μ at steady state with a single first-order biochemical reaction with rate y_μ. Importantly, such coarse-graining does not alter the dynamics of any metabolites outside of module μ (see **Proposition 1** in Materials and Methods). This statement is the biochemical analog of the star-mesh transformation (and its generalization, Kron reduction, **Rao et al., 2014**) well known in the theory of electric circuits (**Versfeld, 1970**). The biological interpretation of these properties is that a module is somewhat isolated from the rest of the metabolic network. And vice versa, the larger network (i.e., the cell) "cares" only about the total rate at which the I/O metabolites are interconverted by the module but "does not care" about the details of how this conversion is enzymatically implemented. In this sense, the effective rate y_μ quantifies the function of module μ (a macroscopic parameter) while the rates $\boldsymbol{x}_\mu$ describe the specific biochemical implementation of the module (microscopic parameters).

The effective rate constant y_μ of module μ depends on the entire rate matrix $\boldsymbol{x}_\mu$. In general, a single mutation may perturb several rate constants within a module, so that the entire shape of the function F may be important. Here, I focus on a special case when each mutation perturbs one reaction (real or effective) within a module, while all others remain constant. To examine epistasis between mutations, I will also consider two different mutations that perturb two separate reactions within a module. In these special cases, we do not need to know the entire function F. We only need to know how module's effective rate constant y_μ depends on the one or two rate constants of the perturbed reactions. When y_μ is considered as a function of the rate constant ξ of one reaction, I write

$$y_\mu = f_1(\xi), \tag{2}$$

and when y_μ is considered as a function of the rate constants ξ and η of two reactions, I write

$$y_\mu = f_2(\xi, \eta). \tag{3}$$

The rate constants of all other reactions within module μ play a role of parameters in functions f_1 and f_2.

Consider now a network $\mathcal{N}$ that has a hierarchical structure in the sense that there is a series of nested modules $\mu \subset \nu \subset \cdots$, in the sense that $A_\mu \subset A_\nu \subset \cdots$ (**Figure 1**A). Since any module at steady state can be replaced with an effective first-order biochemical reaction, there exists a hierarchy of quantitative metabolic phenotypes $y_\mu, y_\nu, \ldots$ (**Figure 1**B,C). These phenotypes are of course functionally related to each other. Specifically, because ν is a "higher-level" module (in the sense that it contains a "lower-level" module μ), the matrix $\boldsymbol{x}_\nu$ can be decomposed into two submatrices, $\boldsymbol{x}_\mu$ and $\boldsymbol{x}_{\nu \setminus \mu}$, where the latter is the matrix of rate constants of reactions that belong to module ν but not to module μ. Since replacing the lower-level module μ with an effective reaction with rate constant y_μ does not alter the dynamics of metabolites outside of μ, y_ν must depend on all elements of $\boldsymbol{x}_\mu$ only through y_μ, i.e.,

$$y_\nu = f_1\left(y_\mu\right), \tag{4}$$

where rates $\boldsymbol{x}_{\nu \setminus \mu}$ act as parameters of function f_1. Thus, in the hierarchy of metabolic phenotypes $y_\mu, y_\nu, \ldots$, a phenotype at each subsequent level depends on the phenotype at the preceding level according to **Equation 4**, and the lowest level phenotype y_μ depends on the actual rate constants accroding to **Equation 1**. This hierarchy of functionally nested phenotypes is conceptually similar to the hierarchical "ontotype" representation of genomic data proposed recently by **Yu et al.** (**2016**).

Quantification of epistasis

Consider a mutation A that perturbs only one rate constant x_{ij}, such that the wildtype value x_{ij}^0 changes to x_{ij}^A. This mutation can be quantified at the microscopic level by its relative effect $\delta^A x_{ij} = x_{ij}^A / x_{ij}^0 - 1$. If the reaction between metabolites i and j belongs to nested modules $\mu, \nu, \ldots$, then mutation A may impact the functions of these modules, which can be quantified by the relative effects $\delta^A y_\mu$, $\delta^A y_\nu$, etc. at each level of the hierarchy.

Consider now another mutation B that only perturbs the rate constant $x_{k\ell}$ of another reaction. Since mutations A and B perturb distinct enzymes, they by definition do not genetically interact at the microscopic level. However, if both perturbed reactions belong to the metabolic module μ (and, as a consequence, to all higher-level modules which contain μ), they may interact at the level of the function of this module, in the sense that the effect of mutation B on the effective rate y_μ may depend on whether mutation A is present or not. Such epistasis between mutations A and B can be quantified at the level μ of the metabolic hierarchy by a number of various epistasis coefficients (*Wagner et al., 1998*; *Hansen and Wagner, 2001*; *Mani et al., 2008*). I will quantify it with the epistasis coefficient

$$\varepsilon^{AB} y_\mu = \frac{\delta^{AB} y_\mu - \delta^A y_\mu - \delta^B y_\mu}{2\,\delta^A y_\mu\,\delta^B y_\mu}, \tag{5}$$

where $\delta^{AB} y_\mu$ denotes the effect of the combination of mutations A and B on phenotype y_μ relative to the wildtype. Since I only consider two mutations A and B, I will write εy_μ instead of $\varepsilon^{AB} y_\mu$ to simplify notations. Note that other epistasis coefficients can always be computed from εy_μ, $\delta^A y_\mu$ and $\delta^B y_\mu$, if necessary. Expressions for epistasis coefficients at other levels of the metabolic hierarchy are analogous.

Results

The central goal of this paper is to understand the patterns of epistasis between mutations that affect reaction rates in the hieararchical metabolic network described above. Specifically, I am interested in two questions. (1) Given that two mutations A and B have an epistasis coefficient εy_μ at a lower level μ of the metabolic hierarchy, what can we say about their epistasis coefficient εy_ν at a higher level ν of the hierarchy? In other words, how does epistasis propagate through the metabolic hierarchy? (2) If mutation A only perturbs the activity x_{ij} of one enzyme and mutation B only perturbs the activity $x_{k\ell}$ of another enzyme that belongs to the same module μ, then what values of εy_μ can we expect to observe based on the topological relationship between the two perturbed reactions within module μ? In other words, what kinds of epistasis emerge in a metabolic network?

Propagation of epistasis through the hierarchy of metabolic phenotypes

Assuming that the effects of both individual mutations and their combined effect at the lower-level μ are small, it follows from *Equation 4* and *Equation 5* that

$$\varepsilon y_\nu = \frac{\varepsilon y_\mu}{C} + \frac{H}{2\,C^2} + o(1), \tag{6}$$

$$\boxed{\begin{array}{ccc} 0 \le & C & \le 1 \\ 0 \le & \bar{\varepsilon} & \le 1 \end{array}}\quad \begin{array}{l}\text{If } \varepsilon y_\mu \le 0,\ \text{then } \varepsilon y_\nu \le 0 \\ \text{If } \varepsilon y_\mu \ge 1,\ \text{then } \varepsilon y_\nu \ge 1\end{array}$$

Figure 2. Propagation of epistasis. Properties of *Equation 6* that maps lower-level epistasis εy_μ onto higher-level epistasis εy_ν. Slope $1/C$ and fixed point $\bar{\varepsilon}$ depend on the topology and the rate constants of the higher-level module ν, but they are bounded, as shown. Thus, the fixed point $\bar{\varepsilon}$ of this map lies between 0 and 1 and is always unstable (open circle).

where $C = f_1' y_\mu / y_\nu$ and $H = f_1'' y_\mu^2 / y_\nu$ are the first- and second-order control coefficients of the lower-level module μ with respect to the flux through the higher-level module ν and $o(1)$ denotes

all terms that vanish as the effects of mutations tend to zero (see Derivation of **Equation 6** and **Equation 9** in Materials and Methods for details). Note that **Equation 6** is a special case of a more general **Equation 49** which describes the case when mutations affect multiple enzymes. **Equation 6** defines a linear map ϕ with slope $1/C$ and a fixed point $\bar{\varepsilon} = -H \left(2C \left(1 - C\right)\right)^{-1}$, which both depend on the topology of the higher-level module ν and the rate constants $\boldsymbol{x}_{\nu \backslash \mu}$.

To gain some intuition for how the map ϕ governs the propagation of epistasis from a lower level μ to a higher level ν, suppose that module ν is a linear metabolic pathway. In this case, it is intuitively clear that function f_1 is monotonically increasing (i.e., the higher y_μ, the more flux can pass through the linear pathway ν) and concave (i.e., as y_μ grows, other reactions in ν become increasingly more limiting, such that further gains in y_μ yield smaller gains in y_ν). Indeed, it is easy to show that $C = \left(1 + \alpha y_\mu\right)^{-1} > 0$ and $H = -2\alpha \, y_\mu \left(1 + \alpha y_\mu\right)^{-2} < 0$, where α is a positive constant that depends on other reactions in the pathway (see Calculation of epistasis in simple modules in Materials and Methods for details). It then immediately follows that any zero or negative epistasis εy_μ that already arose at the lower level would propagate to negative epistasis εy_ν at the level of the linear pathway ν. Moreover, since $C < 1$, the fixed point of the map in **Equation 6** is unstable. Therefore, if epistasis εy_μ was already sufficiently large at the lower level, it would induce even larger positive epistasis εy_ν at the level of the linear pathway ν. In fact, when module ν is a linear pathway, $\bar{\varepsilon} = 1$, so that $\varepsilon y_\nu > 1$ whenever $\varepsilon y_\mu > 1$.

The first result of this paper is the following theorem, which shows that the same rules of propagation of epistasis hold not only for a linear pathway but for any module (**Figure 2**).

Theorem 1. *For any module ν,*

$$0 \leq C \leq 1 \tag{7}$$

and

$$0 \leq \bar{\varepsilon} \leq 1. \tag{8}$$

The proof of **Theorem 1** is given in Proof of **Theorem 1** in Materials and Methods. Its main idea is the following. The functional form of f_1 in **Equation 4** depends on the topology of module ν. Since the number of topologies of ν is infinite, we might a priori expect that there is also an infinite number of functional forms of f_1. However, this is not the case. In fact, all higher-level modules that contain a lower-level module fall into three topological classes defined by the location of the lower-level module with respect to the I/O metabolites of the higher-level module (see **Proposition 2** and **Figure 7** in Materials and Methods). To each topological class corresponds a parametric family of the function f_1, so that there are only three such families. For each family, the values of C and H can be explicitly calculated, yielding the bounds in **Equation 7** and **Equation 8**.

Equation 6 together with **Equation 7** and **Equation 8** show that the linear map ϕ from epistasis at a lower-level to epistasis at the higher-level has an unstable fixed point between 0 and 1 (**Figure 2**). This implies that negative epistasis at a lower level of the metabolic hierarchy necessarily induces negative epistasis of larger magnitude at the next level of the hierarchy, i.e., $\varepsilon y_\nu \leq \varepsilon y_\mu < 0$. Therefore, once negative epistasis emerges somewhere along the hierarchy, it will induce negative epistasis at all higher levels of the hierarchy, irrespectively of the topology or the kinetic parameters of the network.

Similarly, if epistasis at the lower level of the metabolic hierarchy is positive and strong, $\varepsilon y_\mu > 1$, it will induce even stronger positive epistasis at the next level of the hierarchy, i.e., $\varepsilon y_\nu \geq \varepsilon y_\mu > 1$. Therefore, once strong positive epistasis emerges somewhere in the metabolic hierarchy, it will induce strong positive epistasis of larger magnitude at all higher levels of the hierarchy, irrespectively of the topology or the kinetic parameters of the network. If positive epistasis at a lower level of the hierarchy is weak, $0 < \varepsilon y_\mu < 1$, it could induce either negative, weak positive or strong positive epistasis at the higher level of the hierarchy, depending on the precise value of εy_μ, the topology of the higher-level module ν and the microscopic rate constants $\boldsymbol{x}_{\nu \backslash \mu}$.

In summary, there are three regimes of how epistasis propagates through a hierarchical metabolic network. Negative and strong positive epistasis propagate robustly irrespectively of the

Figure 3. Emergence of epistasis and its dependence on the topological relationship between the reactions affected by mutations. (**A**) An example of a simple module ν (same as in *Figure 1*A) where negative, weak positive and strong positive epistasis can emerge between two mutations A and B. (**B**) Epistasis between mutations A and B at the level of module ν depicted in (A) as a function of the rate constant z of a third reaction. The values of other parameters of the network are given in Calculation of epistasis in simple modules in Materials and Methods. (**C**) An example of a simple module where reactions affected by mutations are strictly parallel. In such cases, epistasis for the effective rate constant y_ν is non-positive. Dashed blue lines highlight paths that connect the I/O metabolites and each contain only one of the affected reactions. (**D**) An example of a simple module where reactions affected by mutations are strictly serial. In such cases, epistasis for the effective rate constant y_ν is equal to or greater than 1 (i.e., strongly positive). Dashed blue line highlights a path that connects the I/O metabolites and contains both affected reactions.

360 topology and kinetic parameters of the metabolic network, whereas the propagation of weakly
361 positive epistasis depends on these details. The strongest qualitative prediction that follows from
362 *Theorem 1* is that negative epistasis for a lower-level phenotype cannot turn into positive epistasis
363 for a higher-level phenotype, but the converse is possible.

Emergence of epistasis between mutations affecting different enzymes

Which of the three regimes described above can emerge in metabolic networks and under what
circumstances? In other words, if two mutations affect the same module, are there any constraints
on epistasis that might arise at the level of the effective rate constant of this module? To address
this question, I consider two mutations A and B that affect the rate constants of different single
reactions within a given module.

Consider a relatively simple module ν shown in *Figure 1*A and two mutations A and B that affect
the reactions, as shown in *Figure 3*A. I will now show that the epistasis coefficient ϵy_ν can take values
in all three domains described above, depending on the biochemical details of this module. Using
the recursive procedure for evaluating y_μ described in Network coarse graining and an algorithm
for evaluating the effective rate constant for an arbitrary module in Materials and Methods, it is
straightforward to obtain an analytical expression for y_ν as a function of the rate matrix x_ν, from
which ϵy_μ can also be obtained (see Calculation of epistasis in simple modules in Materials and
Methods for details). To demonstrate that ϵy_μ can take values below 0, between 0 and 1, and above
1, it is convenient to keep all of the rate constants fixed except for the rate constant $z \equiv x_{34}$ of a
reaction that is not affected by mutations A or B, as shown in *Figure 3*A. *Figure 3*B then shows
how the epistasis coefficient ϵy_μ varies as a function of z for one particular choice of all other rate
constants. When z is small, $\epsilon y_\mu < 0$. As z increases, it becomes weakly positive ($0 < \epsilon y_\mu < 1$) and
eventually strongly positive ($\epsilon y_\mu > 1$). Thus, in my model, there are no fundamental constraints on

the types of epistasis that can emerge between mutations.

384  This simple example also reveals that not only value but also sign of epistasis generically depends
385 on the rates of other reactions in the network, such that other mutations or physiological changes in
386 enzyme expression levels can modulate epistasis sign and strength. In other words, "higher-order"
387 and "environmental" epistasis are generic features of metabolic networks.

388  Upon closer examination, the toy example in *Figure 3* also suggests that the sign of εy_ν may
depend predictably on the topological relationship between the affected reactions. When $z = 0$,
the two reactions affected by mutations are parallel, and epistasis is negative. When z is very large,
most of the flux between the I/O metabolites passes through z such that the two reactions affected
by mutations become effectively serial, and epistasis is strongly positive. Other toy models show
consistent results: epistasis between mutations affecting different reactions in a linear pathway is
always positive and epistasis between mutations affecting serial reactions is always negative (see
Calculation of epistasis in simple modules in Materials and Methods for details). These observation
suggest an interesting conjecture. Do mutations affecting parallel reactions always exhibit negative
epistasis and do mutations affecting serial reactions always exhibit positive epistasis? In fact, such
relationship between sign of epistasis and topology has been previously suggested in the literature
(e.g., *Dixon et al., 2009*; *Lehner, 2011*).

400  To formalize and mathematically prove this hypothesis, I first define two reactions as *parallel*
401 within a given module if there exist at least two distinct simple (i.e., non-self-intersecting) paths that
402 connect the I/O metabolites, such that each path lies within the module and contains only one of
403 the two focal reactions. Analogously, two reactions are *serial* within a given module if there exists at
404 least one simple path that connects the I/O metabolites, lies within the module and contains both
405 focal reactions.

406  According to these definitions, two reactions can be simultaneously both parallel and serial, as,
407 for example, the reactions affected by mutations A and B in *Figure 3*A. I call such reaction pairs
408 *serial-parallel*. I define two reactions to be *strictly parallel* if they are parallel but not serial (*Figure 3*C)
409 and I define two reactions to be *strictly serial* if they are serial but not parallel (*Figure 3*D). Thus,
410 each pair of reactions within a module can be classified as either strictly parallel, strictly serial or
411 serial-parallel.

412  The second result of this paper is the following theorem.

413 **Theorem 2.** *Let ξ and η be the rate constants of two different reactions in module μ. Suppose that*
414 *mutation A perturbs only one of these reactions by $\delta^A \xi$ and mutation B perturbs only the other reaction*
415 *by $\delta^B \eta$. In the limit $\delta^A \xi \to 0$ and $\delta^B \eta \to 0$, the following statements are true. If the affected reactions are*
416 *strictly parallel then $\varepsilon y_\mu \leq 0$. If the affected reactions are strictly serial, then $\varepsilon y_\mu \geq 1$.*

417  The detailed proof of this theorem is given in Proof of *Theorem 2* in Materials and Methods. Its
418 key ideas and the logic are the following. It follows from *Equation 3* and *Equation 5* that

$$\varepsilon y_\mu = \frac{H_{\xi\eta}}{2\,C_\xi\,C_\eta} + o(1), \tag{9}$$

419 where $C_\xi = \frac{\partial f_2}{\partial \xi}\frac{\xi}{y_\mu}$, $C_\eta = \frac{\partial f_2}{\partial \eta}\frac{\eta}{y_\mu}$, $H_{\xi\eta} = \frac{\partial^2 f_2}{\partial \xi \partial \eta}\frac{\xi\eta}{y_\mu}$ are the first- and second-order control coefficients of the
420 affected reactions with respect to the flux through module μ and $o(1)$ denotes terms that vanish
421 when $\delta^A \xi$ and $\delta^B \eta$ approach zero (see Derivation of *Equation 6* and *Equation 9* in Materials and
422 Methods for details). Note that *Equation 9* was previously derived by *Chiu et al.* (*2012*).

423  To compute the epistasis coefficient εy_μ for an arbitrary module μ, we need to know the first and
second derivatives of function f_2. Analogous to function f_1, there is a finite number of parametric
families to which f_2 can belong. Specifically, all modules fall into nine topological classes with
respect to the locations of the affected reactions within the module (see *Figure 8*), and each of
these topologies defines a parametric family of function f_2 (see *Proposition 3* and its *Corollary 3*
in Materials and Methods). Most of these topological classes are broad and contain modules where
the affected reactions are strictly parallel, those where they are strictly serial as well as those where

Figure 4. Sensitivity of results of *Theorem 1* and *Theorem 2* with respect to the magnitude of mutational effects. (*A*) Distribution of the position of the fixed point $\bar{\varepsilon}$ of the function ϕ that maps lower-level epistasis εy_μ onto higher-level epistasis εy_ν in modules with random parameters and for mutations with positive effects on y_μ (see text and Materials and Methods for details). All cases are shown in *Figure 4–Figure Supplement 1* and *Figure 4–Figure Supplement 2*. The effect size of both mutations is indicated on each panel. "No f.p." indicates that no fixed point exists. (*B*) Fraction of sampled modules (averaged across generating topologies) where mutations affect strictly serial reactions but the epistasis coefficient is less than 1, contrary to the statement of *Theorem 2* (see text and Materials and Methods for details). All cases statified by generating topology are shown in *Figure 4–Figure Supplement 3*.

Figure 4–Figure supplement 1. Fixed-point position in random modules from the topological class $\mathcal{M}^{\mathrm{io}}$.

Figure 4–Figure supplement 2. Fixed-point position in random modules from the topological class $\mathcal{M}^{\mathrm{i}}$.

Figure 4–Figure supplement 3. Epistasis between strictly serial mutations in random modules, stratified by generating topology.

---

they are serial-parallel. And it is easy to show that not all members of each topological class have the
same sign of εy_μ. However, modules from the same topological class where the affected reactions
are strictly parallel or strictly serial fall into a finite number of topological sub-classes (see *Figure 10*
through *Figure 14*, *Table 2* and *Table 3*). Overall, there are only 17 distinct topologies where the
affected reactions are strictly parallel (*Table 2*), which define 17 parametric sub-families of function
f_2. For all members of these sub-families, *Equation 9* yields $\varepsilon y_\mu \leq 0$ (see *Proposition 7* in Materials
and Methods). Similarly, there are only 11 distinct topologies where the affected reactions are
strictly serial (*Table 3*), which define 11 parametric sub-families of function f_2. For all members of
these sub-families, *Equation 9* yields $\varepsilon y_\mu \geq 1$ (see *Proposition 8* in Materials and Methods).

The results of *Theorem 1* and *Theorem 2* together imply that the topological relationship at
the microscopic level between two reactions affected by mutations constrains the values of their
epistasis coefficient at all higher phenotypic levels. Specifically, if negative epistasis is detected
at any phenotypic level, the affected reactions cannot be strictly serial. And conversely, if strong
positive epistasis is detected at any phenotypic level, the affected reactions cannot be strictly
parallel. In this model, weak positive epistasis in the absence of any additional information does
not imply any specific topological relationship between the affected reactions.

Sensitivity of results with respect to the magnitude of mutational effects

Both *Theorem 1* and *Theorem 2* strictly hold only when the effects of both mutations are infinitesi-
mal. Next, I investigate how these results might change when the mutational effects are finite.

Propagation of epistasis between mutations with finite effect sizes

As mentioned above and discussed in detail in Materials and Methods, all higher-level modules
that contain a lower-level module fall into three topological classes, which I label $\mathcal{M}^{\mathrm{b}}$, $\mathcal{M}^{\mathrm{io}}$ and $\mathcal{M}^{\mathrm{i}}$,
depending on the location of the lower-level module within the higher-level module (see *Figure 7*).
The topological class specifies the parametric family of the function f_1 which maps the effective rate
constant y_μ onto the effective rate constant y_ν (see *Equation 4*). For all modules from the topological
class $\mathcal{M}^{\mathrm{b}}$, function f_1 is linear (see *Equation 29*), which implies that the results of *Theorem 1* hold

exactly even when the effects of mutations are finite. For modules from the topological classes $\mathcal{M}^{io}$ and $\mathcal{M}^{i}$, function f_1 is hyperbolic (see *Equation 30* and *Equation 31*), so that the results of *Theorem 1* may not hold when the effects of mutations are finite. To test the validity of *Theorem 1* in these cases, I calculated the non-linear function ϕ that maps the epistasis coefficient εy_μ onto the epistasis coefficient εy_ν for 1000 randomly generated modules from each of the two topological classes and for mutations that increase or decrease the effective rate constant of the lower-level module y_μ by up to 50% (see Materials and Methods for details).

The validity of *Theorem 1* depended on the sign of mutational effects. When at least one of the two mutations had a negative effect on y_μ, map ϕ had the same properties as described in *Theorem 1*, even for mutations with large effect, i.e., it had a fixed point $\bar{\varepsilon}$ in the interval $[0, 1]$ and this fixed point was unstable. When the effects of both mutations on y_μ were positive and small, these results also held in about 82% of sampled modules (see *Figure 4*A, *Figure 4–Figure Supplement 1*, *Figure 4–Figure Supplement 2*). In the remaining ~18% of sampled modules, the fixed point $\bar{\varepsilon}$ shifted slightly above 1. As the magnitude of mutational effects increased, the fraction of sampled modules with $\bar{\varepsilon} > 1$ grew, reaching 42% when both mutations increased y_μ by 50%. In most of these cases, $\bar{\varepsilon}$ remained below 2, and I found only one module with $\bar{\varepsilon} > 4$ (*Figure 4*A, *Figure 4–Figure Supplement 1*, *Figure 4–Figure Supplement 2*). Whenever the fixed point existed, it was unstable, with the exception of 12 modules for which ϕ was very close to the identity map. For 289 modules (14.5%), the fixed point disappeared when both mutations increased y_μ by 50%. In all these cases, $\varepsilon y_\nu < \varepsilon y_\mu$, indicating that even large positive epistasis may decline as it propagates through the metabolic hierarchy when the effects of mutations are finite.

Emergence of epistasis between mutations with finite effect sizes

As mentioned above and discussed in detail in Materials and Methods, modules where the reactions affected by mutations are strictly parallel fall into 17 topological classes (see *Table 2*) and modules where the reactions affected by mutations are strictly serial fall into 11 topological classes (see *Table 3*). The topological class specifies the parametric family of the function f_2 which maps the rate constants ξ and η of the affected reactions onto the effective rate constant y_μ. To test how well *Theorem 2* holds when the effects of mutations are finite, I calculated εy_μ for randomly generated modules from these topological classes and for mutations increasing or decreasing ξ and η by up to 50% (see Materials and Methods for details).

The validity of *Theorem 2* depended most strongly on the topological relationship between the reaction affected by mutations. Whenever the affected reactions were strictly parallel, the epistasis coefficient at the level of module μ was always less than or equal to zero, even when mutations perturbed the rate constants by as much as 50%, consistent with *Theorem 2*. This was also true for strictly serial reactions, as long as both mutations had positive effects. When the affected reaction were strictly serial and at least one of the mutations had a negative effect, the epistasis coefficient was always positive, but in some cases it was less than 1 (see *Figure 4*B, *Figure 4–Figure Supplement 3*), in disagreement with *Theorem 2*. This indicates that when the effects of mutations are not infinitesimal, even mutations that affect strictly serial reactions can potentially produce negative epistasis for higher-level phenotypes.

Taken together, these results suggest that both *Theorem 1* and *Theorem 2* extend reasonably well, but not perfectly, to mutations with finite effect sizes. The domains of validity of both theorems appear to depend on the sign of mutational effects. The way in which the theorems break down as their assumptions are violated appears to be stereotypical: when the mutational effects increase, more types of mutations produce weak epistasis, and the bias towards negative epistasis increases during propagation from lower to higher levels of the metabolic hierarchy.

Beyond first-order kinetics: epistasis in a kinetic model of glycolysis

The results of previous sections revealed a relationship between network topology and the ensuing epistasis coefficients in an analytically tractable model. However, the assumptions of this model are

Figure 5. Epistasis in a kinetic model of *Escherichia coli* glycolysis. (*A*) Simplified schematic of the model (see *Figure 5–Figure Supplement 1* for details). Different shades of gray in the background highlight four modules as indicated (see text). Light blue circles represent metabolites. Reactions are shown as lines with dark gray boxes. The enzymes catalyzing reactions whose control coefficients with respect to the flux through the module are positive are named; other enzyme names are ommitted for clarity (see *Table 5* and *Table 6* for abbreviations). Three reactions, catalyzed by PGI, PFK, PGDH, for which the epistasis coefficients are shown in panel B are highlighted in dark blue, red and orange, respectively. (*B*) Epistasis coefficients for flux through each module between mutations perturbing the respective reactions, computed at steady state (see text and Materials and Methods for details). Reactions catalyzed by PGI and PGDH are strictly parallel (path g6p-f6p-fdp-gap contains only PGI, path g6p-6pg-ribu5p-gap contains only PGDH and there is no simple path in UGPP between g6p and gap that contains both PGI and PGDH). Reactions catalyzed by PGI and PFK are serial-parallel (path g6p-f6p-fdp-gap contains both reactions, path g6p-f6p-gap contains only PGI, path g6p-6pg-ribu5p-f6p-fdp-gap contains only PFK). Reactions catalyzed by PFK and PGDH are also serial-parallel (path g6p-6pg-ribu5p-f6p-fdp-gap contains both reactions, path g6p-f6p-fdp-gap contains only PFK, path g6p-6pg-ribu5p-gap contains only PGDH).

Figure 5–Figure supplement 1. Detailed schematic of the model.

Figure 5–Figure supplement 2. Control coefficients for the output flux in the FULL module.

Figure 5–Figure supplement 3. Control and epistasis coefficients for the fluxes through multiple sub-modules within glycolysis.

most certainly violated in many realistic situations. It is therefore important to know whether the same or similar rules of epistasis emergence and propagation hold beyond the scope of this model. I address this question here by analyzing a computational kinetic model of glycolysis developed by *Chassagnole et al.* (*2002*). This model keeps track of the concentrations of 17 metabolites, reactions between which are catalyzed by 18 enzymes (*Figure 5*A and *Figure 5–Figure Supplement 1*; see Materials and Methods for details). This model falls far outside of the analytical framework introduced in this paper: some reactions are second-order, reaction kinetics are non-linear, and in several cases the reaction rates are modulated by other metabolites (*Chassagnole et al., 2002*).

Testing the predictions of the analytical theory in this computational model faces two complications. First, in a non-linear model, modules are no longer fully characterized by their effective rate constants, even at steady state. Instead, each module is described by the flux between its I/O metabolites which non-linearly depends on the concentrations of these metabolites. Consequently, the effects of mutations and epistasis coefficients also become functions of the I/O metabolite concentrations. An epistasis coefficient at the level of module ν can still be evaluated according to *Equation 5*, with y_ν now representing the flux through module ν evaluated at a particular concentration of the I/O metabolites. For simplicity, I computationally find the steady state of the full glycolysis network and evaluate the epistasis coefficients only at this steady state, i.e., for each module, I keep the concentrations of the I/O metabolites fixed at their steady-state values for the full network (see Materials and Methods for details).

The second complication is that some control coefficients are so small that they fall below the threshold of numerical precision. Perturbing such reactions has no detectable effect on flux (*Figure 5–Figure Supplement 2*). In the analysis that follows, I ignore such reactions because the epistasis coefficient defined by *Equation 5* can only be computed for mutations with non-zero effects on flux. In addition, the control coefficients of some reactions are negative, which implies that an increase in the rate of such reaction decreases the flux through the module (*Figure 5–Figure Supplement 2*). I also ignore such reactions because there is no analog for them in the analytical theory presented above. After excluding 7 reactions for these reasons, I examine epistasis in 55 pairs of mutations that affect the remaining 11 reactions.

The glycolysis network shown in *Figure 5*A (see also *Figure 5–Figure Supplement 1*) can be naturally partitioned into four modules which I name "LG" (lower glycolysis), "UGPP" (upper glycolysis and pentose phosphate), "GPP" (glycolysis and pentose phosophate) and "FULL". Modules LG and UGPP are non-overlappng and both of them are nested in module GPP which in turn is nested in the FULL module. Thus, at least for some reaction pairs it is possible to calculate epistasis coefficients at three levels of metabolic hierarchy. There are three such pairs, and the results for them are shown in *Figure 5*B. Epistasis for the remaining pairs of reactions can be evaluated only at one or two levels of the hierarchy because these reactions belong to different modules at the lowest levels or because their individual effects are too small. The results for all reaction pairs are shown in *Figure 5–Figure Supplement 3*.

The strongest qualitative prediction of the analytical theory described above is that negative epistasis for a lower-level phenotype cannot turn into positive epistasis for a higher-level phenotype, while the converse is possible. *Figure 5*B and *Figure 5–Figure Supplement 3* show that the data are consistent with this prediction. Another prediction is that epistasis between strictly parallel reactions should be negative. There is only one pair of reactions that are strictly parallel, those catalyzed by glucose-6-phosphate isomerase (PGI) and 6-phosphogluconate dehydrogenase (PGDH), and indeed the epistasis coefficients between mutations affecting these reactions are negative at all levels of the hierarchy (*Figure 5*B). Finally, the analytical theory predicts that mutations affecting strictly serial reactions should exhibit strong positive epistasis. There are 36 reaction pairs that are strictly serial. Epistasis is positive between mutations in 33 of them, and it is strongly positive in 17 of them (*Figure 5–Figure Supplement 3*). Three pairs of strictly serial reactions (those where one reaction is catalyzed by PK and the other is catalyzed by PGI, PGDH or PFK) exhibit negative epistasis (*Figure 5–Figure Supplement 3*). These results suggest that, although one may not be

556 able to naively extrapolate the rules of emergence and propagation of epistasis derived in the
557 simple analytical model to more complex networks, some generalized versions of these rules may
558 nevertheless hold more broadly.

Discussion

Genetic interactions are a powerful tool in genetics, and they play an important role in evolution.
Yet, how epistasis emerges from the molecular architecture of the cell and how it propagates to
higher-level phenotypes, such as fitness, remains largely unknown. Several recent studies made a
statistical argument that the structure of the fitness landscape (and, as a consequence, the epistatic
interactions between mutations at the level of fitness) may be largely independent of the underlying
molecular architecture of the organism (*Martin, 2014*; *Lyons et al., 2020*; *Reddy and Desai, 2020*).
If mutations are typically highly pleiotropic (i.e., affect many independent phenotypes relevant for
fitness) or are engaged in a large number of idiosyncratic epistatic interactions with other mutations
in the genome, the resulting fitness landscapes converge to certain limiting shapes, such as the
Fisher's geometric model (*Martin, 2014*; *Tenaillon, 2014*). To what extent these arguments indeed
apply in practice is unclear. But if they do, most genetic interactions detected at the fitness level
may be uninformative about the architecture of the underlying biological networks.

In this paper, I took a "mechanistic" approach, which is in a sense orthogonal to the statistical
one. In my model of a hierarchical metabolic network, mutations are highly pleiotropic (a mutation
in any enzyme affects all the fluxes in the module) and highly epistatic (a mutation in any enzyme
interacts with mutations in any other enzyme). Yet, these pleiotropic and epistatic effects appear to
be sufficiently structured that some information about the topology of the network is preserved
through all levels of the hierarchy. Indeed, the emergence and propagation of epistasis follow
two simple rules in my model. First, once epistasis emerges at some level of the hierarchy, its
propagation through the higher levels of the hierarchy depends weakly on the details of the network.
Specifically, negative epistasis at a lower level induces negative epistasis at all higher levels and
strong positive epistasis induces strong positive epistasis at all higher levels, irrespectively of the
topology or the kinetic parameters of the network. Second, what type of epistasis emerges in the
first place depends on the topological relationship between the reactions affected by mutations.
In particular, negative epistasis emerges between mutations that affect strictly parallel reactions
and positive epistasis emerges between mutations that affect strictly serial reactions. Insofar as my
model is relevant to nature, the key conclusion from it is that epistasis at high-level phenotypes
carries some, albeit incomplete, information about the underlying topological relationship between
the affected reactions.

These results have implications for the interpretation of empirically measured epistasis coef-
ficients. It is often assumed that a positive epistasis coefficient between mutations that affect
distinct genes signals that their gene products act in some sense serially, whereas a negative
epistasis coefficient is a signal of genetic redundancy, i.e., a parallel relationship between gene
products (*Dixon et al., 2009*). My results suggest that this reasoning is generally correct, but that the
relationship between epistasis and topology is more nuanced. In particular, the sign of the epistasis
coefficient in my model constrains but does not uniquely specify the topological relationship, such
that a negative epistasis coefficient implies that the affected reactions are not strictly serial (but
may or may not be strictly parallel) and an epistasis coefficient exceeding unity excludes a strictly
parallel relationship (but does not necessarily imply a strictly serial relationship). My model suggests
that one should also be careful with inferences going in the other direction, i.e., extrapolating the
patterns of epistasis measured at the biochemical level to those at the level of fitness. For example,
if one wishes to infer the past evolutionary trajectory of an enzyme and finds two amino acid
changes that exhibit a positive interaction at the level of enzymatic activity, it does not automatically
imply that these mutations will exhibit a positive interaction at the level of fitness, even if this
enzyme is essential.

The strongest results presented here rely on several assumptions. I proved *Theorem 1* and *Theorem 2* in the limit of vanishingly small mutational effects. Some results of the metabolic control analysis, notably the summation theorem, are sensitive to this assumption (*Bagheri-Chaichian et al., 2003*; *Bagheri and Wagner, 2004*). To test the sensitivity of my analytical results with respect to this assumption I used numerical simulations of networks with randomly sampled kinetic parameters and found that the results hold reasonably well when the effects of mutations are not infinitesimal.

The most restrictive assumption in the present work is that of first-order kinetics. Networks with only first-order kinetics clearly fail to capture some biologically important phenomena, such as sign epistasis (*Weinreich et al., 2005*; *Chou et al., 2014*; *Ewald et al., 2017*; *Kemble et al., 2020*). I discuss possible ways to relax this assumption below. But at present, a major question remains whether the rules of epistasis and propagation described here hold for realistic biological networks and whether they can be directly used to interpret empirical epistasis coefficients. My analysis of a fairly realistic computational model of glycolysis cautions against overinterpreting empirical epistasis coefficients using the rules derived here. But it also suggests that more general rules of propagation and emergence of epistasis may be found for more realistic networks. Thus, the simple rules derived here should probably be thought of as null expectations.

Relaxing the first-order kinetics assumption is analytically challenging because it is critical for replacing a module with a single effective reaction without altering the dynamics of the rest of the network. Although such lossless replacement is almost certainly not possible in networks with more complex kinetics, advanced network coarse-graining techniques may offer a promising way forward (*Rao et al., 2014*). Flux balance analysis (FBA) is an alternative approach (*Orth et al., 2010*). FBA is appealing because it entirely removes the dependence of the model on reaction kinetics. However, this comes at a substantial cost. In FBA models, fitness and other high-level phenotypes become independent of the internal kinetic parameters, which is clearly unrealistic. Nevertheless, FBA is often very good at capturing the effects of mutations that change the topology of metabolic networks, such as reaction additions and deletions (reviewed in *Gu et al., 2019*). At the same time, there is no natural way within FBA to model mutations that perturb reaction kinetics (*He et al., 2010*; *Alzoubi et al., 2019*). In short, FBA and my approach are complementary (see *Appendix 5* for a more detailed discussion).

Generic properties of epistasis in biological systems

Simple models help us identify generic phenomena—those that are shared by a large class of systems—which should inform our "null" expectations in empirical studies. Deviations from such null in a given system under examination inform us about potentially interesting peculiarities of this system. The model presented here suggests several generic features of epistasis between genome-wide mutations.

Epistasis has two contributions

My analysis shows that the value of an epistasis coefficient measured for a higher level phenotype is a result of two contributions (*Domingo et al., 2019*), propagation and emergence, which correspond to two terms in *Equation 6* (or the more general *Equation 49*). The first term, propagation, shows that if two mutations exhibit epistasis for a lower-level phenotype they also generally exhibit epistasis for a higher-level phenotype. The second contribution comes from the fact that lower-level phenotypes map onto higher-level phenotypes via non-linear functions. This is true even in a simple model with linear kinetics considered here. As a result, even if two mutations exhibit no epistasis at the lower-level phenotype, epistasis must emerge for the higher-level phenotype, as previously pointed out by multiple authors (e.g. *Kacser and Burns, 1981*; *DePristo et al., 2005*; *Martin et al., 2007*; *Chiu et al., 2012*; *Otwinowski et al., 2018*; *Domingo et al., 2019*; *Husain and Murugan, 2020*).

Epistasis depends on the genetic background and environment

My analysis shows that the value of an epistasis coefficient for a particular pair of mutations is in large part determined by the topological relationship between reactions affected by them. Since the topology of the metabolic network itself depends on the genotype (which genes are present in the genome) and on the environment (which enzymes are active or not), the topological relationship between two specific reactions might change if, for example, a third mutation knocks out another enzyme or if an enzyme is up- or down-regulated due to an environmental change (see *Figure 3*). Thus, we should generically expect epistasis between mutations to depend on the environment and on the presence or absence of other mutations in the genome. In other words, $G \times G \times G$ interactions (higher-oder epistasis) and $G \times G \times E$ interactions (environmental epistasis) should be common (*Snitkin and Segrè, 2011*; *Flynn et al., 2013*; *Lindsey et al., 2013*; *Taylor and Ehrenreich, 2015*; *Sailer and Harms, 2017a*). This fact certainly complicates the interpretation of inter-gene epistasis since mutations in the same pair of genes can exhibit qualitatively different genetic interactions in different strains, organisms and environments, as has been observed (*St Onge et al., 2007*; *Musso et al., 2008*; *Tischler et al., 2008*; *Dowell et al., 2010*; *Heigwer et al., 2018*; *Li et al., 2019*). However, the situation may not be hopeless because the topological relationship between two reactions cannot change arbitrarily after addition or removal of a single reaction. For example, if two reactions are strictly parallel, removing a third reaction does not alter their relationship (see *Proposition 5*). Thus, comparing matrices of epistasis coefficients measured in different environments or genetic backgrounds could inform us about how the organism rewires its metabolic network in response to these perturbations (*St Onge et al., 2007*; *Musso et al., 2008*; *Heigwer et al., 2018*; *Li et al., 2019*).

Skew in the distribution of epistasis coefficients

Studies that measure epistasis for fitness-related phenotypes among genome-wide mutations usually find both positive and negative epistasis, but the preponderance of positive and negative epistasis varies. Some authors reported a skew towards positive interactions among deleterious mutations (*Jasnos and Korona, 2007*; *He et al., 2010*; *Johnson et al., 2019*), while others reported a skew towards negative interactions (*Szappanos et al., 2011*; *Costanzo et al., 2016*). Beneficial mutations appear to predominantly exhibit negative epistasis, also known as "diminishing returns" epistasis (e.g., *Martin et al., 2007*; *Khan et al., 2011*; *Chou et al., 2011*; *Kryazhimskiy et al., 2014*; *Schoustra et al., 2016*). The reasons for these patterns are currently unclear. Several recent theoretical papers offer possible statistical explanations for them (*Martin, 2014*; *Lyons et al., 2020*; *Reddy and Desai, 2020*). On the other hand, mechanistic predictions for the distribution of epistasis coefficients are not yet available (but see *Sanjuán and Nebot, 2008*; *Macía et al., 2012*; *Chiu et al., 2012*). The present work does not directly address this problem either, but it provides some additional clues.

First, my model shows that the sign of epistasis at least to some extent reflects the topology of the network. Thus, the distribution of epistasis coefficients at high-level phenotypes in real organisms should ultimately depend on the preponderance of different topological relationships between the edges in biological networks. It then seems *a priori* unlikely that positive and negative interactions would be exactly balanced. Thus, we should expect the distribution of epistasis coefficients to be skewed in one or another direction.

The second observation is that in the metabolic model considered here a positive epistasis coefficient at one level of the hierarchy can turn into a negative one at a higher level, but the reverse is not possible. This bias towards negative epistasis at higher-level phenotypes appears to be even stronger in networks with saturating kinetics (*Figure 5* and *Figure 5–Figure Supplement 3*).

The third observation is that epistasis among beneficial and deleterious mutations affecting metabolic genes should be identical at the level where they arise, provided that beneficial and deleterious mutations are identically distributed among metabolic reactions. Thus, a stronger skew towards negative epistasis among beneficial mutations at the level of fitness could arise in my

model for two mutually non-exclusive reasons. One possibility is that beneficial mutations tend to affect certain special subsets of genes, those that predominantly give rise to positive epistasis. For example, beneficial mutations may for some reason predominantly arise in enzymes that catalyze strictly parallel reactions. Another possibility is that when epistasis between beneficial mutations propagates through the metabolic hierarchy it tends to exhibit a stronger negative bias compared to epistasis between deleterious mutations. Indeed, this phenomenon arises in my model among mutations with large effect (see *Figure 4*A, *Figure 4–Figure Supplement 1* and *Figure 4–Figure Supplement 2*).

Inter-gene epistasis is generic

Perhaps the most important—and also the most intuitive—conclusion of this work is that we should expect epistasis for high-level phenotypes, such as fitness, to be extremely common. Consider first a unicellular organism growing exponentially. Its fitness is fully determined by its growth rate, which can be thought of as the rate constant of an effective biochemical reaction that converts external nutrients into cells (see *Appendix 6* for a simple mathematical model of this statement). In other words, growth rate is the most coarse-grained description of a metabolic network and, as such, it depends on the rate constants of all underlying biochemical reactions. Many previous studies have shown that within-protein epistasis is extremely common (e.g. *Lunzer et al., 2005*; *DePristo et al., 2005*; *Sailer and Harms, 2017b*; *Husain and Murugan, 2020*). Present work shows that, once epistasis arises at the level of protein activity, it will propagate all the way up the metabolic hierarchy and will manifest itself as epistasis for growth rate. It also suggests that growth rate is a generically non-linear function of the rate constants of the underlying biochemical reactions, such that all mutations that affect growth rate individually would also exhibit pairwise epistasis for growth rate with each other (*Kacser and Burns, 1981*; *DePristo et al., 2005*; *Martin et al., 2007*; *Chiu et al., 2012*; *Otwinowski et al., 2018*; *Husain and Murugan, 2020*).

In more complex organisms and/or in certain variable environments, it may be possible to decompose fitness into multiplicative or additive components, e.g., plant's fitness may be equal to the product of the number of seeds it produces and their germination probability, as pointed out by *Khan et al.* (*2011*). Then, mutations that affect different components of fitness would exhibit no epistasis. However, such situations should be considered exceptional, as they require fitness to be decomposable and mutations to be non-pleiotropic.

If epistasis is in fact generic for high-level phenotypes, why do we not observe it more frequently? For example, a recent study that tested almost all pairs of gene knock-out mutations in yeast found genetic interactions for fitness for only about 4% of them (*Costanzo et al., 2016*). One possibility is that many pairs of mutations exhibit epistasis that is simply too small to detect with current methods. As the precision of fitness measurements improves, we would then expect the fraction of interacting gene pairs to grow. Another possibility is that systematic shifts in the distribution of estimated epistasis coefficients away from zero are taken by researchers as systematic errors rather than real phenomena, and are normalized out. Thus, some epistasis that would otherwise be detectable may be lost during data processing.

If epistasis is indeed as ubiquituous as the present analysis suggests, it would call into question how observations of inter-gene epistasis are interpreted. In particular, contrary to a common belief, a non-zero epistasis coefficient does not necessarily imply any specific functional relationship between the components affected by mutations beyond the fact that both components somehow contribute to the measured phenotype (*Boyle et al., 2017*). The focus of future research should then be not merely on documenting epistasis but on developing theory and methods for a robust inference of biological relationships from measured epistasis coefficients.

Materials and Methods

Key ideas and logic of proofs of Theorems 1 and 2

Before proceeding to the detailed proofs of *Theorem 1* and *Theorem 2*, I informally outline some key ideas and the basic logic.

The central object of the theory is a metabolic module. Modules have two key properties. First, a module is somewhat isolated from the rest of the metabolic network, in the sense that all metabolites inside it interact with only two metabolites outside, the I/O metabolites. The second property is that the metabolites within the module are sufficiently connected that each of them individually as well as any subset of them collectively can achieve a quasi-steady state (QSS), given the concentrations of the remaining metabolites. This property is proven in *Proposition 1*.

When some metabolites are at QSS, they can be effectively removed from the network and replaced with effective reactions among the remaining metabolites. In other words, one can "coarse-grain" the network by removing metabolites. This approach is a standard biochemical-network reduction technique (*Segel, 1988*); for example, the Briggs-Haldane derivation of the Michaelis-Menten formula is based on this idea. In general, the resulting effective reactions have more complex (non-linear) kinetics than the original reactions. However, when the original reactions are first-order, the effective reactions are also first-order, i.e., there is no increase in complexity. In Network coarse graining and an algorithm for evaluating the effective rate constant for an arbitrary module, I formally define the coarse-graining procedure (CGP) that eliminates one or multiple metabolites and replaces them with effective reactions.

CGP is an essential concept in my theory. I use it to compute the QSS concentrations for internal metabolites within a module (*Corollary 1*) and thereby prove *Proposition 1*, mentioned above. Since any module μ can achieve a QSS at any concentrations of its I/O metabolites and since any module has only two I/O metabolites, it can be replaced with a single effective reaction (*Corollary 2*). CGP provides a way to calculate the rate constant y_μ of this reaction. In other words, the CGP is an algorithm for evaluating function $F(\boldsymbol{x}_\mu)$ in *Equation 1* for any module μ.

CGP has an important property: its result does not depend on the order in which metabolites are eliminated. Therefore, in computing the effective rate constant of a module, we can choose any convinient way to eliminate its metabolites. Suppose that one module μ is nested within another module ν as in *Figure 1*A. A convenient way to compute the effective rate y_ν of the larger module is to first coarse-grain the smaller module μ, replacing it with an effective rate y_μ, and then eliminate all the remaining metabolites in ν. Since effective rates after coarse-graining do not depend on the order of metabolite elimination, y_ν must depend on the rate constants $\boldsymbol{x}_\mu$ only indirectly, through y_μ. In other words, all the information about the smaller module μ that is relevant for the performance of the larger module ν is contained in y_μ. Therefore, if a mutation or mutations perturb only reactions inside of the smaller module μ, we only need to know their effects on the effective rate constant y_μ to completely understand how they will perturb the performance of the larger module ν. Specifically, if we have two such mutations A and B, all the information about them is contained in three numbers, $\delta^A y_\mu$, $\delta^B y_\mu$ and εy_μ. *Theorem 1* then describes how epistasis at the level of module μ propagates to epistasis at the level of module ν.

The proof of *Theorem 1* proceeds as follows. Let a be the effective reaction with rate constant y_μ that represents module μ within the larger module ν, and consider y_ν as a function of y_μ, as in *Equation 4*. To obtain $f_1(y_\mu)$, it is convenient to first eliminate all metabolites that do not participate in reaction a. No matter what the initial structure of module ν is, such coarse-graining will produce only one of three topologically distinct "minimal" modules, which differ by the location of reaction a with respect to the I/O metabolites of module ν (*Figure 7*). This implies that the function f_1 can belong to three parameteric families, where the parameters are the effective rate constants of reactions other than a in each of the minimal modules. This is the essence of *Proposition 2*. Then *Theorem 1* can be easily proven by explicitly evaluating the first- and second-order control coefficients for each of the three functions and showing that the statements of the theorem hold

for all of them, irrespectively of the function's parameters.

Now consider two reactions a and b with rate constants ξ and η, and imagine the two mutations A and B that affect these reactions. To understand what value of εy_μ will occur, we need to obtain y_μ as a function of ξ and η (*Equation 3*). To do so, it is convenient to first eliminate all metabolites that do not participate in reactions a or b. No matter what the initial structure of module μ is, such coarse-graining will produce only one of nine topologically distinct minimal modules, which differ by the location of reactions a and b with respect to the I/O metabolites of module μ and each other (*Figure 8*). This implies that the function f_2 can belong to nine parameteric families. This is the essence of *Proposition 3* and *Corollary 3*.

How does the topological relationship between reactions a and b translate into epistasis? First, there are only three types of relationships between any pair of reactions in a module: strictly serial, strictly parallel or serial-parallel (see *Figure 3*). Second, *Proposition 4* and *Corollary 4* show that coarse-graining does not alter the strict relationships, i.e., if reactions a and b are strictly serial or strictly parallel before coarse-graining they will remain so after coarse-graining. This is important because it implies that to prove *Theorem 2* we do not need to consider an infinitely large space of all modules but only a much smaller space of all minimal modules, i.e., those that have only those metabolites that participate in the affected reactions a and b. Although the space of all minimal modules is still infinite, the space of their topologies is finite (see *Figure 8*). For some minimal topologies, the connection between the strictly serial or strictly parallel relationship and epistasis can be established very easily. For example, if reaction a and reaction b both share an I/O metabolite as a substrate (see topological class $\mathcal{M}^{\text{io,io,IO}}$ in *Figure 8*), then they are always strictly parallel, no matter what the rest of the module looks like. Evaluating the first- and second-order control coefficients for the function f_2 that corresponds to this topological class reveals that $\varepsilon y_\mu \leq 0$ for any parameter values of f_2.

Unfortunately, most topological classes are too broad and include modules where reactions a and b are strictly serial as well as modules where they are strictly parallel or serial-parallel (for example, class $\mathcal{M}^{\text{io,io,}\emptyset}$). Consequently, the sign of εy_μ for such modules can change depending on the values of the rate constants. However, since the number of distinct minimal topologies is finite, it is possible to identify all minimal topologies where the reactions a and b are strictly serial or strictly parallel. These topological sub-classes define parametric sub-families of function f_2, and we can explicitly calculate εy_μ for all such functions. However, such brute-force approach is extremely cumbersome because the number of distinct minimal topologies is very large.

Fortunately, the following simple and intuitive fact greatly simplifies this problem. If two reactions are strictly serial or strictly parallel, this relationship does not change if a third reaction is removed from the module. This statement is the essence of *Proposition 5*. However, if the two reactions are serial-parallel, removal of a third reaction can change the relationship to a strictly serial or a strictly parallel one. As a consequence, there exist certain most connected "generating" topologies where the relationship between the focal reactions is strictly parallel or strictly serial, and any other strictly serial minimal topology can be produced from at least one of the generating topologies by removal of reactions. This is the essence of *Proposition 6*. All generating topologies can be discovered by a simple algorithm provided in *Appendix 3*. They are listed in *Table 2* and *Table 3* and shown in *Figure 10* throgh *Figure 14*. Each generating topology defines a parametric sub-family of function f_2, and I explicitly evaluate the first- and second-order control coefficients for all these sub-families (see *Proposition 7* and *Proposition 8*) which essentially completes the proof of *Theorem 2*.

Network coarse-graining

Notations and definitions

Here, I give a more precise definition of the model and introduce additional notations and definitions. As mentioned above, all reactions are first order and reversible. Thus, each reaction $i \leftrightarrow j$ has one substrate $i \in A$ and one product $j \in A$, and it is fully described by its rate constant x_{ij}. By definition, $x_{ii} = 0$. I denote the set of all reactions by $R = \{i \leftrightarrow j \,:\, i, j \in A, \ x_{ij} > 0\}$. The dynamics of metabolite

concentrations $S_1, \dots, S_n$ in the network $\mathcal{N}$ are governed by equations

$$\dot{S}_i = \sum_{j=1}^{n} x_{ji} \, S_j - D_i \, S_i, \quad i \in A \tag{10}$$

where

$$D_i = \sum_{j=1}^{n} x_{ij}, \quad i \in A. \tag{11}$$

In what follows, it will be important to distinguish three types of reactions within a module, based on their topological relationship to the I/O metabolites of that module. The topology of the module μ is determined by its set of reactions $R_\mu = \left\{ i \leftrightarrow j \in R : i, j \in A_\mu \cup A_\mu^{\mathrm{IO}} \right\}$. I call all reactions where both the substrate and the product are internal to module μ as *reactions internal to μ*, and I denote the set of all such reactions by $R_\mu^{\mathrm{i}} \subset R_\mu$. For example, module μ in **Figure 1**A has one internal reaction $3 \leftrightarrow 4$. I call all reactions where one of the metabolites is internal to μ and the other is an I/O metabolite as the *i/o reactions for μ*, and I denote the set of all such reactions by $R_\mu^{\mathrm{io}} \subset R_\mu$. (I reserve upper-case "I/O" for metabolites and use lower-case "i/o" for reactions.) For example, module μ in **Figure 1**A has three i/o reactions $1 \leftrightarrow 3$, $1 \leftrightarrow 4$ and $2 \leftrightarrow 4$. Finally, I refer to reactions between any two I/O metabolites for module μ as *bypass reactions for module μ*, and I denote the set of all such reactions by $R_\mu^{\mathrm{b}} \subset R_\mu$. For example, module μ in **Figure 1**A has no bypass reactions but reaction $1 \leftrightarrow 5$ is a bypass reaction for module ν. By definition, all these three sets of reactions R_μ^{i}, R_μ^{io} and R_μ^{b} are non-overlapping, and $R_\mu = R_\mu^{\mathrm{i}} \cup R_\mu^{\mathrm{io}} \cup R_\mu^{\mathrm{b}}$.

Another important concept are the simple paths that lie within a module (see Model). For any two metabolites $i, j \in A \cup A_\mu^{\mathrm{IO}}$, I denote a simple path between them that lies within μ as p_{ij}^μ or, equivalently as $i \leftrightarrow k_1 \leftrightarrow \dots \leftrightarrow k_m \leftrightarrow j$ (where all $k_\ell \in A_\mu$). I say that each of the metabolites k_ℓ *belongs to* (or *is contained in*) path p_{ij}^μ (denoted by $k_\ell \in p_{ij}^\mu$). Similarly, I say that each of the reactions $k_\ell \leftrightarrow k_{\ell+1}$ *belong to* (or *are contained in*) path p_{ij}^μ (denoted by $k_\ell \leftrightarrow k_{\ell+1} \in p_{ij}^\mu$). I will drop superindex μ from p_{ij}^μ if it is clear what module is being referred to.

Network coarse graining and an algorithm for evaluating the effective rate constant for an arbitrary module

In this section, I formally introduce and characterize the coarse-graining procedure (CGP). In Elimination of a single metabolite, I introduce the main idea, which is to eliminate a metabolite that is at QSS and to replace it with a set of new reactions between metabolites adjacent to the eliminated one. This is exactly analogous to the star-mesh transformation in the theory of electric circuits (**Versfeld, 1970**). The resulting network is a coarse-grained version of the original network in the sense that it has one less metabolite. Next, in The coarse-graining procedure (CGP), I define the CGP, which is simply multiple metabolite eliminations applied successively. I prove **Proposition 1**, which justifies applying the CGP to a whole module and replacing it with a single effective reaction (**Corollary 2**). Finally, in Computation of effective rate constants for simple modules, I show how to apply the CGP in practice to compute function F from **Equation 1** for modules with some simple topologies.

Elimination of a single metabolite

I begin by outlining the main idea behind the CGP, which is to replace one metabolite internal to a module with a series of effective reactions between metabolites adjacent to it. If the effective rate constants are defined appropriately, the dynamics of all metabolites in the resulting coarse-grained network are the same as in the original network, provided that the eliminated metabolite is at QSS in the original network.

To formalize this idea, suppose that module $\mu = \left(A_\mu, \mathbf{x}_\mu \right)$ contains m internal metabolites. Let $k \in A_\mu$ be the internal metabolites that will be eliminated. Let $A^{\{k\}} = A \setminus \{k\}$ be the reduced

890 metabolite set and let $\boldsymbol{x}^{\{k\}}$ be the reduced $(n-1) \times (n-1)$ matrix of rate constants defined by

$$x_{ij}^{\{k\}} \;=\; x_{ij} + \frac{x_{ik}\, x_{kj}}{D_k}, \quad i, j \in A^{\{k\}}, \quad i \neq j, \tag{12}$$

$$x_{ii}^{\{k\}} \;=\; 0, \quad i \in A^{\{k\}}, \tag{13}$$

where D_k is given by *Equation 11*.

Such metabolite elimination has three properties that follow immediately from *Equation 12*
and *Equation 13*. First, the rate constants of reactions do not change as long as the eliminated
metabolite does not participate in them. Mathematically, $x_{ij}^{\{k\}} = x_{ij}$ for all metabolites i and j
that are not adjacent to the eliminated metabolite k. In particular, this is true for all metabolites
external to module μ. Second, because equilibrium constants have the property $K_{ij} = K_{i\ell} K_{\ell j}$ for
any metabolites i, j, ℓ, the rate constants $x_{ij}^{\{k\}}$ obey Haldane's relationships. Therefore, the reduced
metabolite set $A^{\{k\}}$ and the reduced rate matrix $\boldsymbol{x}^{\{k\}}$ define a new "coarse-grained" metabolic
network $\mathcal{N}^{\{k\}} = \left(A^{\{k\}}, \boldsymbol{x}^{\{k\}} \right)$. It is easy to show that subnetwork μ after the elimination of metabolite
k is still a module. Third, the reaction set of module μ (i.e., its topology) in the coarse-grained
network $\mathcal{N}^{\{k\}}$ depends only on the reaction set of this module in the original network $\mathcal{N}$, but not
on the particular values in the rate matrix $\boldsymbol{x}_\mu$.

Next, I will show that the dynamics of metabolites in the coarse-grained network $\mathcal{N}^{\{k\}}$ are
identical to the dynamics of metabolites in the original network $\mathcal{N}$ where metabolite k is at QSS.
Note that if metabolite k is at QSS in the network $\mathcal{N}$, its concentration is given by

$$S_k = \sum_{j \in A_\mu^{\mathrm{IO}} \cup A_\mu \setminus \{k\}} \frac{x_{jk}\, S_j}{D_k}, \tag{14}$$

906 which follows from *Equation 10*. Now, the dynamics of metabolites in the network $\mathcal{N}^{\{k\}}$ are
907 governed by equations

$$\dot{S}_i = \sum_{j \in A^{\{k\}}} x_{ji}^{\{k\}}\, S_j - D_i^{\{k\}}\, S_i, \quad \text{for } i \in A^{\{k\}}, \tag{15}$$

where $D_i^{\{k\}} = \sum_{j \in A^{\{k\}}} x_{ij}^{\{k\}}$. As mentioned above, $x_{ij}^{\{k\}} = x_{ij}$ for all pairs of metabolites where at least
one metabolite is external to module μ. Therefore, *Equation 15* for the external metabolites are
identical to *Equation 10* that govern the dynamics of these metabolites in the original network $\mathcal{N}$.
Next, consider the dynamics of the I/O and internal metabolites for module μ in the coarse-grained
network $\mathcal{N}^{\{k\}}$, i.e., those in the set $A_\mu^{\mathrm{IO}} \cup A_\mu \setminus \{k\}$. For any such metabolite i, the sum in the righthand
side of *Equation 15* can be re-written as

$$\sum_{j \in A_\mu^{\mathrm{IO}} \cup A_\mu \setminus \{k\}} x_{ji}^{\{k\}}\, S_j = \sum_{j \in A_\mu^{\mathrm{IO}} \cup A_\mu \setminus \{k\}} \left(x_{ji} + \frac{x_{jk}\, x_{ki}}{D_k} \right) S_j - \frac{x_{ik}\, x_{ki}}{D_k}\, S_i$$

$$= \sum_{j \in A_\mu^{\mathrm{IO}} \cup A_\mu \setminus \{k\}} x_{ji}\, S_j + x_{ki} \sum_{j \in A_\mu^{\mathrm{IO}} \cup A_\mu \setminus \{k\}} \frac{x_{jk}\, S_j}{D_k} - \frac{x_{ik}\, x_{ki}}{D_k}\, S_i. \tag{16}$$

914 According to *Equation 14*, the second term in *Equation 16* equals $x_{ki} S_k$, so that *Equation 16* becomes
915

$$\sum_{j \in A_\mu^{\mathrm{IO}} \cup A_\mu \setminus \{k\}} x_{ji}^{\{k\}}\, S_j = \sum_{j \in A_\mu^{\mathrm{IO}} \cup A_\mu} x_{ji} S_j - \frac{x_{ik}\, x_{ki}}{D_k}\, S_i. \tag{17}$$

916 For any metabolite $i \in A_\mu^{\mathrm{IO}} \cup A_\mu \setminus \{k\}$, the second term in the righthand side of *Equation 15* can be
917 re-written as

$$D_i^{\{k\}} = \sum_{j \in A_\mu^{\mathrm{IO}} \cup A_\mu \setminus \{k\}} \left(x_{ij} + \frac{x_{ik}\, x_{kj}}{D_k} \right) - \frac{x_{ik}\, x_{ki}}{D_k} = D_i - \frac{x_{ik}\, x_{ki}}{D_k}. \tag{18}$$

918 Substituting *Equation 17* and *Equation 18* into *Equation 15*, we see that *Equation 15* is in fact
919 equivalent to *Equation 10* for all $i \in A \setminus \{k\}$ with S_k given by *Equation 14*.

The coarse-graining procedure (CGP)

Here, I define the CGP for an arbitrary set of internal metabolites by applying metabolite elimination recursively.

Let $E \subseteq A_\mu$ be an arbitrary subset of metabolites internal to module μ in the metabolic network $\mathcal{N}$ and let n_E be the number of elements in E. Let $A^E = A \setminus E$ be the reduced metabolite set after the metabolites have been eliminated. I define the reduced $(n - n_E) \times (n - n_E)$ matrix of rate constants $\boldsymbol{x}^E$ as follows. If $n_E = 1$, the matrix $\boldsymbol{x}^E$ is defined by *Equation 12* and *Equation 13*. If $n_E > 1$, then I define it recursively. Suppose that all metabolites in E other than some metabolite $k \in E$ have been previously eliminated, such that the coarse-grained network $\mathcal{N}^{E'} = \left(A^{E'}, \boldsymbol{x}^{E'}\right)$ is defined, with the set of eliminated metabolites $E' = E \setminus \{k\}$, $A^{E'} = A \setminus E'$, and the known matrix $\boldsymbol{x}^{E'}$. Then, I define the matrix $\boldsymbol{x}^E$ through the elimination of metabolite k from $\mathcal{N}^{E'}$, i.e.,

$$x_{ij}^E \;=\; x_{ij}^{E'} + \frac{x_{ik}^{E'} x_{kj}^{E'}}{D_k^{E'}}, \quad i, j \in A^E,\ i \neq j, \tag{19}$$

$$x_{ii}^E \;=\; 0, \quad i \in A^E, \tag{20}$$

with

$$D_k^{E'} = \sum_{j \in A^{E'}} x_{kj}^{E'}. \tag{21}$$

I define the *the coarse-graining procedure that eliminates the metabolite set E* as a map

$$\mathrm{CG}^E \;:\; \mathcal{N} \mapsto \mathcal{N}^E = \left(A^E, \boldsymbol{x}^E\right).$$

As with the elimination of a single metabolite, it is straightforward to show that the rate constants x_{ij}^E obey Haldane's relationships, so that $\mathcal{N}^E$ is indeed a metabolic network. CG^E maps module μ within the metabolic network $\mathcal{N}$ onto a subnetwork μ' within the metabolic network $\mathcal{N}^E$. It is straightforward to show that μ' is a module. Whenever there is no ambiguity, I will label both the original and the coarse-grained versions of the module by μ. To simplify notations, if the CGP eliminates the entire module μ (i.e., if $E = A_\mu$), I label it CG^μ. I label the coarse-grained network that restults from the application of CG^μ by $\mathcal{N}^\mu$ and I label the effective rate of the reaction substituting module μ in network $\mathcal{N}^\mu$ as y_μ.

Intuitively, the result of coarse-graining should not depend on the order in which the metabolites are eliminated. To see this, let us obtain explicit (i.e., not recursive) expressions for x_{ij}^E. First, by applying the recursion *Equation 19*, it is easy to show that elimination of two metabolites $E = \{k, \ell\}$ yields effective rate constants

$$
\begin{aligned}
x_{ij}^{\{k,\ell\}} \;&=\; x_{ij}^{\{k\}} + \frac{x_{i\ell}^{\{k\}} x_{\ell j}^{\{k\}}}{D_\ell^{\{k\}}} \\[2mm]
&=\; x_{ij} + \frac{D_k\, x_{i\ell}\, x_{\ell j} + D_\ell\, x_{ik}\, x_{kj} + x_{ik}\, x_{k\ell}\, x_{\ell j} + x_{i\ell}\, x_{\ell k}\, x_{kj}}{D_k\, D_\ell - x_{k\ell}\, x_{\ell k}} \tag{22} \\[2mm]
&=\; x_{ij} + \frac{x_{i\ell}\, x_{\ell j}}{D_\ell^{\{k\}}} + \frac{x_{ik}\, x_{kj}}{D_k^{\{\ell\}}} + \frac{x_{ik}\, x_{k\ell}\, x_{\ell j}}{D_k^{\{\ell\}}\, D_\ell} + \frac{x_{i\ell}\, x_{\ell k}\, x_{kj}}{D_\ell^{\{k\}}\, D_k}, \tag{23} \\[2mm]
& \quad i, j \in A \setminus \{k, \ell\},\ i \neq j.
\end{aligned}
$$

As expected, *Equation 22* and *Equation 23* are symmetric with respect to the eliminated metabolites k and ℓ. Extrapolating from *Equation 23*, it is possible to show that for an arbitrary metabolite subset $E \subseteq A_\mu$ that contains n_E metabolites,

$$x_{ij}^E = x_{ij} + \sum_{L=1}^{n_E} \sum_{(k_1, \ldots, k_L)} \frac{x_{ik_1}}{1} \frac{x_{k_1 k_2}}{D_{k_1}^{E \setminus \{k_1\}}} \cdots \frac{x_{k_{L-1} j}}{D_{k_L}^{E \setminus \{k_1, k_2, \ldots, k_L\}}}, \quad i, j \in A \setminus E,\ i \neq j. \tag{24}$$

Here, the second sum is taken over all $n_E! / (n_E - L)!$ ordered lists of metabolites $(k_1, \ldots, k_L)$ from E. Each list can be thought of as a simple path within E that connects metabolites i and j. The

proof of *Equation 24* can be found in *Appendix 1*. As expected, *Equation 24* shows that the effective reation rate x_{ij}^E does not depend on the order in which metabolites are eliminated. This and other properties of the CGP are listed in *Box 1*.

One of key building blocks of the proofs of *Theorem 1* and *Theorem 2* is the fact that modules can be classified into a finite number of topological classes (see Classification of modules with respect to "marked" reactions, and the parametric families of functions f_1 and f_2 below). To arrive at this classification, it will be convenient to define a composition of coarse-graining procedures, as follows. Suppose that CG^{E_1} and CG^{E_2} are two coarse-graining procedures of network $\mathcal{N}$ for two subsets of metabolites $E_1 \subset A_\mu$ and $E_2 \subset A_\mu$. If the sets E_1 and E_2 are non-overlapping, CG^{E_2} is also defined for the coarse-grained network $\mathcal{N}^{E_1}$ which is the result of applying CG^{E_1} to the original network $\mathcal{N}$. The result of applying CG^{E_2} to the $\mathcal{N}^{E_1}$ is called *the composition of coarse-graining procedures* CG^{E_1} and CG^{E_2} of the original network $\mathcal{N}$ and is denoted as $\mathrm{CG}^{E_1} \circ \mathrm{CG}^{E_2}$.

As defined above, coarse-graining is a formal procedure, and there is no a priori guarantee that (a) it can in fact be carried out for every set of metabolites and (for example, because a metabolite set does not have a steady-state solution); and (b) it will not distort the dynamics in the rest of the network. The following proposition alleviates both of these concerns and thereby justifies the use of the CGP for any subset of internal metabolites within a module (including the entire module μ). It is straightforward to prove it by induction, using the same logic as in the Elimination of a single metabolite.

Proposition 1. *Let E be any subset of metabolites internal to module μ. Then*

1. *There exists a joint QSS solution $\overline{S}_i$ for all metabolites $i \in E$, given the concentrations of the remaining internal and I/O metabolites for module μ.*
2. *The dynamics of all remaining metabolites in $A \setminus E$ in the coarse-grained metabolic network $\mathcal{N}^E$ are the same as in $\mathcal{N}$ where all metabolites in E are at QSS.*

Corollary 1. *Without loss of generality, suppose that the I/O metabolites for module μ are labelled 1 and 2 and its internal metabolites are labelled $A_\mu = \{3, 4, \dots, m\}$. There exists a unique QSS $\overline{S}_i$ for all $i \in A_\mu$. The QSS concentrations can be obtained by recursively applying equation*

$$\overline{S}_k = \frac{1}{D_k^{\{k+1,\dots,m\}}} \left(x_{1k}^{\{k+1,\dots,m\}} S_1 + x_{2k}^{\{k+1,\dots,m\}} S_2 + \sum_{j \in \{3,\dots,k-1\}} x_{jk}^{\{k+1,\dots,m\}} \overline{S}_j \right) \tag{25}$$

for $k = 3, 4, \dots, m$.

Equation 25 follows from *Equation 10* for the coarse-grained network obtained by eliminating metabolites $k + 1, \dots, m$.

Corollary 2. *Without loss of generality, suppose that the I/O metabolites for module μ are labelled 1 and 2. Module μ can be replaced with a single effective reaction between its I/O metabolites, whose rate constant y_μ can be calculated using* **Equation 19** *and* **Equation 20** *or* **Equation 24**. *The dynamics of all metabolites in the resulting coarse-grained metabolic network are identical to their dynamics in the original network $\mathcal{N}$ where all metabolites internal to module μ are at the QSS determined by* **Equation 25**.

Computation of effective rate constants for simple modules

Corollary 2 provides a method for replacing any module μ at QSS with an effective rate $y_\mu = F(\boldsymbol{x}_\mu)$, which can be calculated using *Equation 19* and *Equation 20* or *Equation 24*. Here, I show how to implement this calculation for three simple metabolic modules.

Linear pathway

Consider a linear pathway with I/O metabolites 1 and m and internal metabolites $2, 3, \dots, m-1$ (*Figure 6*A). This labelling of metabolites is more convenient for the linear pathway. To calculate y_μ, I will apply recursion *Equation 19* and *Equation 20*. I start by eliminating metabolite 2. After

Box 1. Properties of the coarse-graining procedure

The coarse-graining procedure CG^E eliminating metabolites in set $E \subseteq A_\mu$ has the following useful properties which follow from *Equation 24*.

1. The effective rate constants x_{ij}^E do not depend on the order in which metabolites are eliminated. Therefore, the composition of coarse-graining procedures is commutative, i.e., if $E = E_1 \cup E_2$, where E_1 and E_2 are two non-overlapping subsets of A_μ, then

$$\mathrm{CG}^{E_1} \circ \mathrm{CG}^{E_2} = \mathrm{CG}^{E_2} \circ \mathrm{CG}^{E_1} = \mathrm{CG}^E.$$

2. If at least one of the metabolites i or j is not adjacent to any of the eliminated metabolites, then $x_{ij}^E = x_{ij}$. In particular, $x_{ij}^E = x_{ij}$ if either i and/or j are external to μ.
3. The topology of module μ after the application of CG^E depends only on its original topology but not on the values of its rate constants $\boldsymbol{x}_\mu$.
4. If both metabolites i and j are adjacent to at least one eliminated metabolite, then

$$x_{ij}^E = x_{ij} + \alpha,$$

   where $\alpha \geq 0$ depends only on the rate constants of reactions that involve at least one eliminated metabolite. In particular, if both k and ℓ are from $A \setminus E$, then x_{ij}^E is independent of $x_{k\ell}$.
5. If $E = A_\mu$, then the effective rate constant y_μ of module μ depends on the rate matrix $\boldsymbol{x}_\mu$ but does not depend on any other reaction rates, i.e., *Equation 1* holds. Furthermore, the functional form of $F(\boldsymbol{x}_\mu)$ depends only on the topology of module μ, i.e., all modules with the same topology are mapped onto y_μ by the same function F.
6. Suppose that module μ is nested in a larger module $v = (A_v, \boldsymbol{x}_v)$ (see *Figure 1*A). It follows from Property #1 that $\mathrm{CG}^v = \mathrm{CG}^\mu \circ \mathrm{CG}^{A_v \setminus A_\mu}$, i.e., y_v can be obtained by carrying out the CGP in two stages, by first eliminating module μ and replacing it with the effective reaction with the rate constant y_μ and then eliminating the remaining metabolites in A_v. In the network $\mathcal{N}^\mu$ after the first stage of coarse-graining, all rate constants $\boldsymbol{x}_{v \setminus \mu}$ are identical to those in the original network, i.e., they are independent of $\boldsymbol{x}_\mu$, by virtue of Property #2. Therefore y_v depends on the rate constants $\boldsymbol{x}_\mu$ of reactions within module μ only through the effective rate constant y_μ of module μ. In other words, *Equation 4* holds.
7. If, in the original network $\mathcal{N}$, metabolites i and j are adjacent or connected by a simple path that contains only the eliminated metabolites, then metabolites i and j are adjacent in the coarse-grained network $\mathcal{N}^E$.
8. If, in the metabolic network $\mathcal{N}$, metabolites i and j are not adjacent (i.e., $x_{ij} = 0$) and no simple path exists within the set E (i.e., such that all non-terminal metabolites in this path are from E) that connects metabolites i and j, then metabolites i and j are also not adjacent in the coarse-grained network $\mathcal{N}^E$ (i.e., $x_{ij}^E = 0$).
9. It follows from properties #7 and #8 that for a simple path $p_{\ell_1 \ell_m} = \ell_1 \leftrightarrow \ell_2 \leftrightarrow \cdots \leftrightarrow \ell_m$ to exist in module μ after the application of CG^μ, it is neccessary and sufficient that for each $i = 1, \ldots, m-1$, either ℓ_i and ℓ_{i+1} are adjacent in the original module μ or there exists a simple path $\ell_i \leftrightarrow \cdots \leftrightarrow \ell_{i+1}$ within the original module μ all of whose non-terminal metabolites are from E.

Figure 6. Simple modules. (**A**) Linear pathway. (**B**) Two parallel pathways.

this initial coarse-graining step, the resulting module is still a linear pathway, where two reactions $1 \leftrightarrow 2 \leftrightarrow 3$ were replaced with a single reaction $1 \leftrightarrow 3$ with the effective rate constant

$$x_{13}^{\{2\}} = \frac{x_{12}\,x_{23}}{x_{21}+x_{23}} = \left(\frac{1}{K_{12}\,x_{23}} + \frac{1}{x_{12}}\right)^{-1}.$$

All other rate constants remain unchanged. Next, I eliminate metabolite 3. The resulting module is still a linear pathway, where now three reactions $1 \leftrightarrow 2 \leftrightarrow 3 \leftrightarrow 4$ were replaced with a single reaction $1 \leftrightarrow 4$ with the effective rate constant

$$x_{14}^{\{2,3\}} = \frac{x_{13}^{\{2\}}\,x_{34}}{x_{31}^{\{2\}}+x_{34}} = \left(\frac{1}{K_{13}\,x_{34}} + \frac{1}{K_{12}\,x_{23}} + \frac{1}{x_{12}}\right)^{-1}.$$

All other rate constants remain unchanged. Continuing this process until all internal metabolites are eliminated, I obtain

$$y_\mu = \left(\sum_{i=1}^{m-1} \frac{1}{K_{1i}\,x_{i\,i+1}}\right)^{-1}, \tag{26}$$

which is identical to the expression originally obtained by **Kacser and Burns** (**1973**).

Two parallel pathways

Consider two parallel pathways with I/O metabolites 1 and 2 and internal metabolites 3 and 4 (**Figure 6**B). I obtain the effective rate constant using **Equation 22** with $i = 1$, $j = 2$, $k = 3$, $\ell = 4$. Since $x_{12} = x_{34} = 0$, **Equation 22** simplifies to

$$y_\mu = \frac{D_3\,x_{14}\,x_{42} + D_4\,x_{13}\,x_{32}}{D_3\,D_4} = \frac{x_{14}\,x_{42}}{x_{41}+x_{42}} + \frac{x_{13}\,x_{32}}{x_{31}+x_{32}}. \tag{27}$$

Thus, the contributions of parallel pathways are simply added.

Module μ in **Figure 1**

To obtain the effective rate constant for module μ shown in **Figure 1**, I again use **Equation 22** with $i = 1$, $j = 2$, $k = 3$, $\ell = 4$.

$$y_\mu = \frac{D_3\,x_{14}\,x_{42} + D_4\,x_{13}\,x_{32} + x_{13}\,x_{34}\,x_{42} + x_{14}\,x_{43}\,x_{32}}{D_3\,D_4 - x_{34}\,x_{43}}, \tag{28}$$

with $D_3 = x_{31} + x_{32} + x_{34}$ and $D_4 = x_{41} + x_{42} + x_{43}$.

Classification of modules with respect to "marked" reactions, and the parametric families of functions f_1 and f_2.

The CGP described above allows us to calculate the function F that maps the rate matrix $\boldsymbol{x}_\mu$ for an arbitrary module μ onto the module's effective rate constant y_μ. F is a multivariate function of the entire matrix $\boldsymbol{x}_\mu$. However, in many applications, only one or two reactions are varied at a time while all others remain constant, and we want to know how module's effective parameter y_μ depends on these one or two perturbed reactions. I refer to such singled-out reactions as "marked". When y_μ is considered as a function of the rate constant ξ of one marked reaction, I write $y_\mu = f_1(\xi)$, as in **Equation 2**. When y_μ is considered as a function of the rate constants ξ and η of two marked reactions, I write $y_\mu = f_2(\xi, \eta)$ as in **Equation 3**.

Figure 7. Classification of single-marked modules. Left column shows a general module from each topological class. The right column shows a minimal fully connected module in each topological class (see text for details). Circles represent metabolites and lines represent reactions. Only the I/O metabolites and the metabolites that participate in the marked reaction are shown, all other metabolites are suppressed. Short lines that have only one terminal metabolite represent all remaining reactions in which this metabolite participates, reactions between all other metabolites are suppressed. Metabolites are labelled according to the conventions listed in the text. The marked reaction is colored orange and labelled a. The module is represented by a gray rectangle, and the rest of the network is not shown.

The functional form of F and, as a consequence, the functional forms of f_1 and f_2 depend only on the topology of module μ (see Property #5 of the CGP in *Box 1*). In other words, modules with identical topologies have the same functional forms of f_1 and f_2, such that each topology of module μ defines a parametric family of functions f_1 and f_2, where all rate constants within module μ other than ξ, or ξ and η, play a role of parameters.

Since the number of possible topologies is infinite, there is an infinite number of functional forms of F. However, the number of parameteric families of functions f_1 and f_2 is finite, and it turns out to be small. To see this, notice that for any module with a single marked reaction, the CGP can be carried out in two stages. In the first stage, we can eliminate all metabolites that do not participate in the marked reaction. The resulting coarse-grained module is minimal in the sense that it can have at most two internal metabolites. Such minimal modules (and, as a consequence, all modules with one marked reaction) fall into three distinct topological classes, which are specified by the location of the marked reaction with respect to the I/O metabolites, as shown in *Figure 7*. This implies that there are only three parameteric families of the function f_1. The topologies of the three minimal modules are sufficiently simple that the three corresponding parametric functional forms of f_1 can be easily computed by applying the coarse-graining *Equation 19* or *Equation 22*. This result is formulated in *Proposition 2*.

The same logic applies to modules with two marked reactions. CGP that eliminates all metabolites that do not participate in the marked reactions maps all such modules onto respective minimal modules, which can have at most four internal metabolites (see *Figure 8*). This result is formulated in *Proposition 3*. Minimal modules (and, as a consequence, all modules with two marked reactions) fall into nine distinct topological classes, which are specified by the locations of the marked reactions. All modules from the same topological class are described by functions f_2 from the same parametric family. These families are characterized in *Corollary 3*.

These topological classifications are extremely useful for the following reason. If we can show

that all functions from the same parameteric family (corresponding to a given topological class) have some common property irrespectively of the values of the parameters, it would imply that this property holds for all modules from the corresponding topological class. This logic is a key part of the proofs of both *Theorem 1* and *Theorem 2*.

To formalize this reasoning, consider module $\mu = (A_\mu, x_\mu)$ and let $a = i_a \leftrightarrow j_a$ and $b = i_b \leftrightarrow j_b$ be two reactions from its set of reactions R_μ. I call a pair (μ, a) *a single-marked module* and I call a triplet (μ, a, b) *a double-marked module*, and I refer to reactions a and b as *marked* within module μ. The topology of a single-marked module (μ, a) is determined not only by the reaction matrix R_μ, but also by the position of the marked reaction, so I refer to the pair (R_μ, a) as the *topology of the single-marked module* (μ, a). Similarly, I refer to the triplet (R_μ, a, b) as the *topology of the double-marked module* (μ, a, b). I denote by $x_{\mu \backslash a}$ the matrix of rate constants of all reactions in module μ other than reaction a and I denote by $x_{\mu \backslash \{a,b\}}$ the matrix of all rate constants in module μ other than reactions a and b. I denote the sets of all single- and double-marked modules by $\mathcal{M}_1$ and $\mathcal{M}_2$, respectively. To avoid metabolite labelling ambiguities, I adopt the following conventions:

(i) The I/O metabolites are labelled $1, 2$ and the internal metabolites are labelled $3, 4, \ldots$;

(ii) $i_a, j_a \in \{1, 2, 3, 4\}$; $i_b, j_b \in \{1, 2, 3, 4, 5, 6\}$;

(iii) $i_a < j_a$, $i_b < j_b$, $i_a \le i_b$, $j_a \le j_b$.

It is easy to see that the set of all single-marked modules $\mathcal{M}_1$ can be partitioned into three non-overlapping topological classes depending on the type of the marked reaction a. I denote the classes of all single-marked modules where the marked reaction is bypass, i/o or internal (see Notations and definitions) by $\mathcal{M}^b$, $\mathcal{M}^{io}$ and $\mathcal{M}^i$, respectively (*Figure 7*). Similarly, the set $\mathcal{M}_2$ can be partitioned into 9 non-overlapping topological classes according to the types of marked reactions and the type of metabolite that is shared by both of these reactions (I/O, internal, or none). These classes are listed in *Table 1* and illustrated in *Figure 8*.

Each topological class contains infinitely many modules, with various numbers of metabolites and various topologies. However, for each topological class $\mathcal{M}$, there is a minimum number of internal metabolites $m_{\mathcal{M}}$, such that all modules within $\mathcal{M}$ must have at least $m_{\mathcal{M}}$ internal metabolites. I denote the set of metabolites minimal in the topological class $\mathcal{M}$ by $A_{\mathcal{M}}$. It is clear that for the single-marked module classes $\mathcal{M}^b$ and for $\mathcal{M}^{io}$, $m_{\mathcal{M}^b} = m_{\mathcal{M}^{io}} = 1$ and $A_{\mathcal{M}^b} = A_{\mathcal{M}^{io}} = \{3\}$, and for $\mathcal{M}^i$, $m_{\mathcal{M}^i} = 2$ and $A_{\mathcal{M}^i} = \{3, 4\}$ (see second column in *Figure 7*). For the double-marked modules, $m_{\mathcal{M}}$ and $A_{\mathcal{M}}$ are given in *Table 1* and illustrated in *Figure 8*.

If a single-marked module from the topological class $\mathcal{M}$ has the minimum number of metabolites $n_{\mathcal{M}}$ in that class, I call such module and its topology *minimal* in $\mathcal{M}$. There may be several minimal topologies in a topological class, but there is only one minimal topology that is fully connected. A topology (R_μ, a) is called *fully connected* if the reaction set R_μ is *complete* in the sense that it contains reactions between all pairs of metabolites in the minimal metabolite set $A_{\mathcal{M}}$. I denote such complete reaction set for the class $\mathcal{M}$ by $R_{\mathcal{M}}$, and I denote the respective fully connected topology by $(R_{\mathcal{M}}, a)$. I employ the same terminology and analogous notations for double-marked modules. The minimal fully connected topologies are shown in the second column in *Figure 7* and *Figure 8*.

Next, I prove *Proposition 2*, which is the key step towards the proof of *Theorem 1*. This proposition states that there are only three functional forms for the function f_1 and characterizes them. The idea of the proof is the following. According to Property 5 (*Box 1*), all single-marked modules that are mapped by the CGP onto a minimal module with the same topology (R_μ, a) have the same functional form of f_1. In other words, each minimal topology (R_μ, a) specifies a parameteric family of the function f_1. Since the number of possible minimal topologies is finite, the claim of *Theorem 1* can be tested for each corresponding functional form of f_1. However, the number of minimal topologies is rather large. Fortunately, another simplification is possible. Since the reaction set R_μ of any minimal single-marked module is a subset of the complete reaction set $R_{\mathcal{M}}$, the fully connected topology $(R_{\mathcal{M}}, a)$ specifies the largest parametric family of the function f_1 for the class $\mathcal{M}$, such that all other families can be obtained from it by setting some parameters to zero, which is equivalent to

removing reactions from the fully connected topology. In other words, *all* single-marked modules that belong to the topological class $\mathcal{M}$ are described by functions f_1 that belong to one parametric family corresponding to the fully connected topology minimal in $\mathcal{M}$. The three parametric families of f_1 are characterized by *Proposition 2*.

One important consequence of *Proposition 2* is that it is not necessary to test the claim of *Theorem 1* for each family of f_1 that corresponds to each minimal topology. Instead, it is sufficient to test it for the three families that correspond to the fully connected minimal topologies in each class.

Proposition 2. *Let (μ, a) be a single-marked module, and let ξ be the rate constant of reaction a. Then $y_\mu = f_1(u)$, where $u = \xi + \alpha$ for some $\alpha \geq 0$, and the function f_1 is given by one of the following expressions.*

$$f_1(u) = u, \quad if\ (\mu, a) \in \mathcal{M}^{\mathrm{b}}, \tag{29}$$

$$f_1(u) = w_{12} + \frac{u\, w_{32}}{u/K_{13} + w_{32}}, \quad if\ (\mu, a) \in \mathcal{M}^{\mathrm{io}}, \tag{30}$$

$$f_1(u) = w_{12} + \frac{D_3\, w_{14}\, w_{42} + D_4\, w_{13}\, w_{32} + w_{13}\, w_{42}\, u + w_{14}\, w_{32}\, u/K_{34}}{D_3\, D_4 - u^2/K_{34}}, \quad if\ (\mu, a) \in \mathcal{M}^{\mathrm{i}}. \tag{31}$$

Here $D_3 = w_{31} + w_{32} + u$, $D_4 = w_{41} + w_{42} + u/K_{34}$, and all $w_{ij} \geq 0$ are independent of ξ.

Proof. Since any single-marked module (μ, a) belongs to exactly one of three classes $\mathcal{M}^{\mathrm{b}}$, $\mathcal{M}^{\mathrm{io}}$ and $\mathcal{M}^{\mathrm{io}}$, I consider these three cases one by one.

Case $(\mu, a) \in \mathcal{M}^{\mathrm{b}}$**.** According to the labelling conventions outlined above, $a = 1 \leftrightarrow 2$ (see *Figure 7*). *Equation 29* follows directly from Property #4 of the CGP (*Box 1*).

Case $(\mu, a) \in \mathcal{M}^{\mathrm{io}}$**.** According to the labelling conventions, $a = 1 \leftrightarrow 3$ (see *Figure 7*). According to Property #1 of the CGP, module μ can be coarse-grained in two stages, by first applying $\mathrm{CG}^{A_\mu \backslash \{3\}}$ which eliminates metabolites $4, \dots, m$ (those that do not participate in the marked reaction) and then applying $\mathrm{CG}^{\{3\}}$ which eliminates the remaining metabolite 3. Mathematically, $\mathrm{CG}^\mu = \mathrm{CG}^{A_\mu \backslash \{3\}} \circ \mathrm{CG}^{\{3\}}$. After applying $\mathrm{CG}^{A_\mu \backslash \{3\}}$, the resulting coarse-grained module μ' has a single internal metabolite 3 and at most three effective reactions $1 \leftrightarrow 2$, $1 \leftrightarrow 3$ and $2 \leftrightarrow 3$ (*Figure 7*), i.e., it is minimal in $\mathcal{M}^{\mathrm{io}}$. By virtue of Properties #2 and #4 of the CGP, the effective rate constants w_{12}, w_{23} are independent of ξ and $u \equiv w_{13} = \xi + \alpha$. Note that w_{12} may equal zero, but $w_{23} \neq 0$ because μ' is a module. Regardless, the reaction set $R_{\mu'}$ of module μ' is always a subset of the complete reaction set $R_{\mathcal{M}^{\mathrm{io}}}$. Thus, to obtain the effective rate constant y_μ, I consider the most general case when μ' is fully connected and eliminate the remaining internal metabolite 3, which leads to *Equation 30*.

Case $(\mu, a) \in \mathcal{M}^{\mathrm{io}}$**.** According to the labelling conventions, $a = 3 \leftrightarrow 4$ (see *Figure 7*). Otherwise, the logic of the proof is exactly the same as for the case $(\mu, a) \in \mathcal{M}^{\mathrm{io}}$. $\square$

Next I prove *Proposition 3* which states that, for any double-marked module that belongs to a given topological class, there exists a double-marked module that is minimal in the same class, such that both modules have the same function f_2. The corresponding minimal module is obtained from the original module by applying the CGP. This proposition is important because it implies that all functions f_2 can be completely characterized by only examing minimal modules. Then, analogously to single-marked modules, *Corollary 3* states that function f_2 can belong to one of nine parametric families which are defined by the fully connected minimal topologies in each topological class.

Proposition 3. *Let (μ, a, b) be a double-marked module that belongs to the topological class $\mathcal{M}$, and let ξ and η be the rate constants of reactions a and b, respectively. Then there exist non-negative constants α and β and a double-marked module (μ', a, b) minimal in $\mathcal{M}$ such that $y_\mu = y_{\mu'} = f_2(u, v)$, where*

$$u = \xi + \alpha, \tag{32}$$

$$v = \eta + \beta \tag{33}$$

are the rate constants of the marked reactions a and b in μ', respectively, and all other rate constants in μ' are independent of ξ or η. Module μ' is obtained from μ by the coarse-graining procedure $\mathrm{CG}^{\mu \backslash \{a, b\}}$ that eliminates all metabolites internal to module μ that do not participate in reactions a or b.

Table 1. Classification of double-marked modules. Metabolites are labelled according to conventions described in the text. $m_{\mathcal{M}}$ is the minimum number of internal metabolites in a module from class $\mathcal{M}$. $A_{\mathcal{M}}$ is the set of internal and I/O metabolites in all minimal modules in class $\mathcal{M}$.

Class	a	b	Shared metab.	Verbal description	$m_{\mathcal{M}}$	$A_{\mathcal{M}}$	Equation for f_2
$\mathcal{M}^{\mathrm{b,io,IO}}$	$(1,2)$	$(1,3)$	1	bypass and i/o reactions, shared I/O metabolite	1	$\{1,2,3\}$	(34)
$\mathcal{M}^{\mathrm{b,i,\emptyset}}$	$(1,2)$	$(3,4)$	–	bypass and internal reactions, no shared metabolies	2	$\{1,2,3,4\}$	(35)
$\mathcal{M}^{\mathrm{io,io,I}}$	$(1,3)$	$(2,3)$	3	i/o reactions, shared internal metabolite	1	$\{1,2,3\}$	(36)
$\mathcal{M}^{\mathrm{io,io,IO}}$	$(1,3)$	$(1,4)$	1	i/o reactions, shared I/O metabolite	2	$\{1,2,3,4\}$	(37)
$\mathcal{M}^{\mathrm{io,io,\emptyset}}$	$(1,3)$	$(2,4)$	–	i/o reactions, no shared metabolites	2	$\{1,2,3,4\}$	(38)
$\mathcal{M}^{\mathrm{io,i,I}}$	$(1,3)$	$(3,4)$	3	i/o and internal reactions, shared internal metabolite	2	$\{1,2,3,4\}$	(39)
$\mathcal{M}^{\mathrm{io,i,\emptyset}}$	$(1,3)$	$(4,5)$	–	i/o and internal reactions, no shared metabolites	3	$\{1,2,3,4,5\}$	(40)
$\mathcal{M}^{\mathrm{i,i,I}}$	$(3,4)$	$(3,5)$	3	internal reactions, share internal metabolite	3	$\{1,2,3,4,5\}$	(41)
$\mathcal{M}^{\mathrm{i,i,\emptyset}}$	$(3,4)$	$(5,6)$	–	internal reactions, no shared metabolites	4	$\{1,2,3,4,5,6\}$	(42)

Figure 8. Classification of double-marked modules. Notations as in *Figure 7*.

Proof. To prove this proposition, I will construct the double-marked module (μ', a, b) minimal in
$\mathcal{M}$ by applying $\mathrm{CG}^{\mu \backslash \{a,b\}}$. Let $m_{\mathcal{M}}$ be the mimimal number of internal metabolites in class $\mathcal{M}$ (see
Table 1). According to the metabolite labelling conventions, metabolites $n_{\mathcal{M}} + 3, n_{\mathcal{M}} + 4, \dots$ are
neither I/O nor do they participate in the marked reactions. $\mathrm{CG}^{\mu \backslash \{a,b\}}$ eliminates all these metabolites
and maps module μ onto module μ', all of whose internal metabolites participate in reactions
a and/or b. Therefore, (μ', a, b) is minimal in class $\mathcal{M}$ (*Figure 8*). According to Properties #2 and
#4 of the CGP (*Box 1*), the effective rate constants u and v of reactions a and b in module μ' are
given by linear relationships in *Equation 32* and *Equation 33*, and the remaining effective rate
constants are independent of ξ and η. The fact that $y_{\mu} = y_{\mu'}$ follows from Property #1 of the CGP,
$\mathrm{CG}^{\mu} = \mathrm{CG}^{\mu \backslash \{a,b\}} \circ \mathrm{CG}^{A_{\mu'}}$. $\square$

Corollary 3. *Let (μ, a, b) be a double-marked module, and let ξ and η be the rate constants of reactions*
a and b, respectively. The function f_2 that maps ξ and η onto module's effective rate constant y_{μ} belongs
to one of nine parametric families. If $(\mu, a, b) \in \mathcal{M}^{\mathrm{b,io,IO}}$, then

$$f_2(u, v) \quad = \quad u + \frac{v\, w_{32}}{v/K_{13} + w_{32}}. \tag{34}$$

1195 *If $(\mu, a, b) \in \mathcal{M}^{\mathrm{b,i,\emptyset}}$, then*

$$f_2(u, v) \quad = \quad u + \frac{D_3\, w_{14}\, w_{42} + D_4\, w_{13}\, w_{32} + w_{13}\, w_{42}\, v + w_{14}\, w_{32}\, v/K_{34}}{D_3\, D_4 - v^2/K_{34}}, \tag{35}$$

$$D_3 \quad = \quad w_{31} + w_{32} + v,$$

$$D_4 \quad = \quad w_{41} + w_{42} + v/K_{34}.$$

1196 *If $(\mu, a, b) \in \mathcal{M}^{\mathrm{io,io,I}}$, then*

$$f_2(u, v) \quad = \quad w_{12} + \frac{u\, v}{u/K_{13} + v}, \tag{36}$$

1197 *If $(\mu, a, b) \in \mathcal{M}^{\mathrm{io,io,IO}}$, then*

$$f_2(u, v) \quad = \quad w_{12} + \frac{D_3\, v\, w_{42} + D_4\, u\, w_{32} + u\, w_{34}\, w_{42} + v\, w_{43}\, w_{32}}{D_3\, D_4 - w_{34}\, w_{43}}, \tag{37}$$

$$D_3 \quad = \quad u/K_{13} + w_{32} + w_{34},$$

$$D_4 \quad = \quad v/K_{14} + w_{42} + w_{43}.$$

1198 *If $(\mu, a, b) \in \mathcal{M}^{\mathrm{io,io,\emptyset}}$, then*

$$f_2(u, v) \quad = \quad w_{12} + \frac{D_3\, w_{14}\, v/K_{24} + D_4\, u\, w_{32} + u\, w_{34}\, v/K_{24} + w_{14}\, w_{43}\, w_{32}}{D_3\, D_4 - w_{34}\, w_{43}}, \tag{38}$$

$$D_3 \quad = \quad u/K_{13} + w_{32} + w_{34},$$

$$D_4 \quad = \quad w_{41} + v/K_{24} + w_{43}.$$

1199 *If $(\mu, a, b) \in \mathcal{M}^{\mathrm{io,i,I}}$, then*

$$f_2(u, v) \quad = \quad w_{12} + \frac{D_3\, w_{14}\, w_{42} + D_4\, u\, w_{32} + u\, v\, w_{42} + w_{14}\, w_{32}\, v/K_{34}}{D_3\, D_4 - v^2/K_{34}}, \tag{39}$$

$$D_3 \quad = \quad u/K_{13} + w_{32} + v,$$

$$D_4 \quad = \quad w_{41} + w_{42} + v/K_{43}.$$

1200 *If $(\mu, a, b) \in \mathcal{M}^{\mathrm{io,i,\emptyset}}$, then*

$$f_2(u, v) \quad = \quad W_{12} + \frac{W_{13}\, W_{32}}{W_{31} + W_{32}}, \tag{40}$$

$$W_{ij} \quad = \quad w_{ij} + \frac{D_4\, w_{i5}\, w_{5j} + D_5\, w_{i4}\, w_{4j} + w_{i4}\, v\, w_{5j} + w_{i5}\, w_{4j}\, v/K_{45}}{D_4\, D_5 - v^2/K_{45}},$$

$$D_4 \quad = \quad w_{41} + w_{42} + w_{43} + v,$$

$$D_5 \quad = \quad w_{51} + w_{52} + w_{53} + v/K_{45},$$

$$w_{13} \quad \equiv u.$$

1201 *If $(\mu, a, b) \in \mathcal{M}^{\text{i,i,I}}$, then*

$$f_2(u, v) = W_{12} + \frac{D_3 \, W_{14} \, W_{42} + D_4 \, W_{13} \, W_{32} + W_{13} \, W_{34} \, W_{42} + W_{14} \, W_{43} \, W_{32}}{D_3 \, D_4 - W_{34} \, W_{43}}, \tag{41}$$

$$W_{ij} = w_{ij} + \frac{w_{i5} \, w_{5j}}{D_5},$$

$$D_3 = W_{31} + W_{32} + W_{34},$$

$$D_4 = W_{41} + W_{42} + W_{43},$$

$$D_5 = w_{51} + w_{52} + w_{53} + w_{54},$$

$$w_{34} \equiv u,$$

$$w_{35} \equiv v.$$

1202 *If $(\mu, a, b) \in \mathcal{M}^{\text{i,i,}\emptyset}$, then*

$$f_2(u, v) = W_{12} + \frac{D_3 \, W_{14} \, W_{42} + D_4 \, W_{13} \, W_{32} + W_{13} \, W_{34} \, W_{42} + W_{14} \, W_{43} \, W_{32}}{D_3 \, D_4 - W_{34} \, W_{43}}, \tag{42}$$

$$W_{ij} = w_{ij} + \frac{D_5 \, w_{i6} \, w_{6j} + D_6 \, w_{i5} \, w_{5j} + w_{i5} \, w_{6j} \, v + w_{i6} \, w_{5j} \, v / K_{56}}{D_5 \, D_6 - v^2 / K_{56}}, \tag{43}$$

$$D_3 = W_{31} + W_{32} + W_{34},$$

$$D_4 = W_{41} + W_{42} + W_{43},$$

$$D_5 = w_{51} + w_{52} + w_{53} + w_{54} + v,$$

$$D_6 = w_{61} + w_{62} + w_{63} + w_{64} + v / K_{56},$$

$$w_{34} \equiv u.$$

*In **Equation 34** through **Equation 42**, u and v are given by **Equation 32** and **Equation 33**. All effective*
activities $w_{ij} \geq 0$ are constants (other than cases where they stand for u or v) that depend on the topology
of module μ and on the parameters $\boldsymbol{x}_{\mu \setminus \{a,b\}}$ but do not depend on ξ and η.

Proof. This statement and *Equation 34* through *Equation 42* follow directly from *Proposition 3* and
the fact that the reaction set of any double-marked module in any given topological class is a subset
of the complete reaction set in that topological class. □

Derivation of *Equation 6* and *Equation 9*
Consider a higher-level phenotype y, such as the effective activity of a module, which is function
of a multivariate lower-level phenotype $\boldsymbol{x} = (x_1, x_2, \ldots, x_n)$, such as the rates of individual reactions
within the module, $y = F(\boldsymbol{x})$. Denote the wildtype values of the phenotypes as $\boldsymbol{x}^0 = (x_1^0, x_2^0, \ldots, x_n^0)$
and $y^0 = F(\boldsymbol{x}^0)$. Consider a mutation that perturbes these values, so that the mutant has lower-
level phenotypic values $\boldsymbol{x}' = (x_1', x_2', \ldots, x_n')$. The relative effect of the mutation on phenotype x_i
is $\delta x_i = x_i'/x_i^0 - 1$. If all $\|\delta x_i\| \ll 1$ where $\|\boldsymbol{x}\|$ denotes the length of vector $\boldsymbol{x}$, then the value of the
higher-level phenotype y' in the mutant is given by

$$y' = y^0 \left(1 + \sum_{i=1}^{n} C_i \, \delta x_i + \frac{1}{2} \sum_{i,j=1}^{n} H_{ij} \, \delta x_i \, \delta x_j \right) + o\left(\|\delta \boldsymbol{x}\|^2 \right). \tag{44}$$

1217 where

$$C_i = \left. \frac{x_i^0}{y^0} \frac{\partial F}{\partial x_i} \right|_{\boldsymbol{x}=\boldsymbol{x}^0}, \quad i = 1, \ldots, n, \tag{45}$$

$$H_{ij} = \left. \frac{x_i^0 x_j^0}{y^0} \frac{\partial^2 F}{\partial x_i \, \partial x_j} \right|_{\boldsymbol{x}=\boldsymbol{x}^0}, \quad i, j = 1, \ldots, n, \tag{46}$$

1218 which I refer to as *first-* and *second-order control coefficients* of the lower-level phenotypes x_i and x_j
1219 with respect to the higher-level phenotype y.

Now consider two single mutants, A and B, and the double-mutant AB. Each mutation A and B and their combination may perturb all x_i phenotypes such that $x_i^A = x_i^0 \left(1 + \delta^A x_i\right)$, $x_i^B = x_i^0 \left(1 + \delta^B x_i\right)$, and

$$x_i^{AB} = x_i^0 \left(1 + \delta^{AB} x_i\right) = x_i^0 \left(1 + \delta^A x_i + \delta^B x_i + 2\,\delta^A x_i\, \delta^B x_i\, \varepsilon x_i\right).$$

Assuming that $\|\delta^A \boldsymbol{x}\| \ll 1$, $\|\delta^B \boldsymbol{x}\| \ll 1$ and $\|\delta^{AB} \boldsymbol{x}\| \ll 1$, using the approximation in *Equation 44* and the definition of εx_i (analogous to *Equation 5*), I obtain

$$\delta^A y = \sum_{i=1}^{n} C_i\, \delta^A x_i + o\left(\|\delta^A \boldsymbol{x}\|\right), \tag{47}$$

$$\delta^B y = \sum_{i=1}^{n} C_i\, \delta^B x_i + o\left(\|\delta^B \boldsymbol{x}\|\right), \tag{48}$$

$$\varepsilon y = \frac{\sum_{i=1}^{n} C_i\, \varepsilon x_i\, \delta^A x_i\, \delta^B x_i + \frac{1}{2} \sum_{i,j=1}^{n} H_{ij}\, \delta^A x_i\, \delta^B x_j}{\sum_{i=1}^{n} \sum_{j=1}^{n} C_i\, C_j\, \delta^A x_i\, \delta^B x_j} + o(1), \tag{49}$$

where $o(1)$ refers to terms that are vanishingly small as all $\delta^A x_i \to 0$, $\delta^B x_i \to 0$, $i = 1, \dots n$.

I examine two special cases of *Equation 49*. The first special case is when both mutations affect a single phenotype x_k, i.e., when all $\delta^A x_i = 0$ and all $\delta^B x_i = 0$ except for $i = k$. Then *Equation 47*, *Equation 48*, *Equation 49* simplify to

$$\delta^A y = C_k\, \delta^A x_k + o\left(|\delta^A x_k|\right), \tag{50}$$

$$\delta^B y = C_k\, \delta^B x_k + o\left(|\delta^B x_k|\right), \tag{51}$$

$$\varepsilon y = \frac{\varepsilon x_k}{C_k} + \frac{H_{kk}}{2\, C_k^2} + o(1). \tag{52}$$

Equation 52 is equivalent to *Equation 6*.

The second special case is when mutation A affects a single phenotypes x_k and mutation B affects a single phenotype x_ℓ ($k \neq \ell$), i.e., all $\delta^A x_i = 0$ except for $i = k$, all $\delta^B x_i = 0$ except for $i = \ell$, and all $\varepsilon x_i = 0$. Then *Equation 47*, *Equation 48*, *Equation 49* simplify to

$$\delta^A y = C_k\, \delta^A x_k + o\left(|\delta^A x_k|\right), \tag{53}$$

$$\delta^B y = C_\ell\, \delta^B x_\ell + o\left(|\delta^B x_\ell|\right), \tag{54}$$

$$\varepsilon y = \frac{H_{k\ell}}{2\, C_k\, C_\ell} + o(1). \tag{55}$$

Equation 55 is equivalent to *Equation 9*.

Calculation of epistasis in simple modules

Equation 52 and *Equation 55* allow me to compute how epistasis propagates and emerges in arbitrary metabolic networks. In this section, I show how to implement these calculations for three simple metabolic modules considered above (see Computation of effective rate constants for simple modules) and in module ν shown in *Figure 3*.

Linear pathway

First, consider how epistasis propagates through a linear pathway (*Figure 6*A). For simplicity, assume that both mutations A and B affect the same reaction $1 \leftrightarrow 2$. It follows from *Equation 26* that

$$y_\mu = f_1(\xi) = \left(\frac{1}{\xi} + \alpha\right)^{-1},$$

where α is a positive constant. Therefore, the first- and second-order control coefficients of reaction $1 \leftrightarrow 2$ with respect to the flux through the linear pathway μ are given by

$$C = \frac{y_\mu}{\xi} = \frac{1}{1 + \alpha\, \xi},$$

$$H = -\frac{2\alpha\, \xi}{(1 + \alpha\, \xi)^2}.$$

Substituting these expressions into the expression for the fixed point $\bar{\varepsilon} = -H\left(2C\left(1-C\right)\right)^{-1}$, I find that $\bar{\varepsilon} = 1$, irrespective of the rates of other reactions in the linear pathway. This implies that epistasis $\varepsilon\xi < 1$ at the level of reaction $1 \leftrightarrow 2$ would induce a lower value of epistasis $\varepsilon y_\mu < \varepsilon\xi < 1$ at the level of the entire linear pathway, any value $\varepsilon\xi > 1$ would induce a higher value of epistasis $\varepsilon y_\mu > \varepsilon\xi > 1$, and $\varepsilon\xi = 1$ would induce $\varepsilon y_\mu = 1$.

Now consider emergence of epistasis in a linear pathway. Suppose that mutation A affects reaction $1 \leftrightarrow 2$ and mutation B affects reaction $2 \leftrightarrow 3$. Denote the rate constant of reactions $1 \leftrightarrow 2$ and $2 \leftrightarrow 3$ by $\xi \equiv x_{12}$ and $\eta \equiv x_{23}$, respectively. It follows from *Equation 26* that

$$y_\mu = f_2(\xi, \eta) = \left(\frac{1}{\xi} + \frac{1}{K_{12}\eta} + \beta\right)^{-1},$$

where β is a positive constant. Therefore,

$$C_\xi = \frac{y_\mu}{\xi},$$

$$C_\eta = \frac{y_\mu}{K_{12}\,\eta},$$

$$H_{\xi\eta} = \frac{2\,y_\mu^2}{K_{12}\,\xi\,\eta},$$

which, after substituting into *Equation 9*, yield $\varepsilon y_\mu = 1$. Together with the fact that $\bar{\varepsilon} = 1$ (see above), this result implies that epistasis coefficient between any two mutations that affect different reactions in a linear pathway equals 1.

Two parallel pathways

Suppose that mutation A affects reaction $1 \leftrightarrow 3$ and mutation B affects reaction $1 \leftrightarrow 4$ in the linear metabolic pathway shown in *Figure 6*B. Denote the rate constants of reaction $1 \leftrightarrow 3$ and $1 \leftrightarrow 4$ by $\xi \equiv x_{13}$ and $\eta \equiv x_{14}$. It follows from *Equation 27* that

$$y_\mu = f_2(\xi, \eta) = \left(\frac{1}{\xi} + \alpha\right)^{-1} + \left(\frac{1}{\eta} + \beta\right)^{-1},$$

where $\alpha = 1/(K_{13}\,x_{32})$ and $\beta = 1/(K_{14}\,x_{42})$. Thus, we have $H_{\xi\eta} = 0$, and there is no epistasis between such mutations.

Module ν in *Figure 3*A

I denote the rate of the reactions affected by mutations A and B by $\xi = x_{13}$ and $\eta = x_{42}$, and I also denote $z = x_{34}$. I will calculate the epistasis coefficient εy_ν in two stages, by first calculating the epistasis coefficient εy_μ and then propagating it to εy_ν using *Equation 6*. Here I am specifically interested in how εy_ν depends on the rate constant z.

To compute epistasis between mutations A and B at the level of module μ, I rewrite *Equation 28* as

$$y_\mu = \frac{a\,\xi\,\eta + b_\xi\,\xi + b_\eta\,\eta + c}{d\,\xi\,\eta + e_\xi\,\xi + e_\eta\,\eta + f},$$

where $a = x_{14}/K_{13} + x_{32} + z$, $b_\xi = x_{32}\left(x_{41} + z/K_{34}\right)$, $b_\eta = x_{14}\left(x_{32} + z\right)$, $c = x_{14}\,x_{32}\,z/K_{34}$, $d = 1/K_{13}$, $e_\xi = \left(x_{41} + z/K_{34}\right)/K_{13}$, $e_\eta = x_{32} + z$, $f = x_{32}\,z/K_{34} + x_{41}\,z + x_{32}\,x_{41}$. I obtain the following expressions for the first- and second-order control coefficients.

$$C_\xi = \frac{\xi}{y_\mu}\left(\frac{\tilde{c}_1\,z + \tilde{d}_1}{D}\right)^2, \tag{56}$$

$$C_\eta = \frac{\eta}{y_\mu}\frac{1}{K_{14}}\left(\frac{\tilde{c}_2\,z + \tilde{d}_2}{D}\right)^2, \tag{57}$$

$$H_{\xi\eta} = \frac{\xi\,\eta}{y_\mu}\frac{2\,z}{K_{14}}\frac{\left(\tilde{c}_1\,z + \tilde{d}_1\right)\left(\tilde{c}_2\,z + \tilde{d}_2\right)}{D^3}, \tag{58}$$

where $D = d\,\xi\,\eta + e_\xi\,\xi + e_\eta\,\eta + f$, $\tilde{c}_1 = x_{23}/K_{24} + \eta$, $\tilde{d}_1 = x_{32}\,(x_{41} + \eta)$, $\tilde{c}_2 = \xi + x_{14}$, $\tilde{d}_2 = x_{14}\,(\xi/K_{13} + x_{32})$.

Substituting *Equation 56* through *Equation 58* into *Equation 53* through *Equation 55*, I obtain

$$\delta^A y_\mu \;=\; \frac{\xi}{y_\mu}\left(\frac{\tilde{c}_1\,z + \tilde{d}_1}{D}\right)^2 \delta^A \xi, \tag{59}$$

$$\delta^B y_\mu \;=\; \frac{\eta}{y_\mu}\frac{1}{K_{14}}\left(\frac{\tilde{c}_2\,z + \tilde{d}_2}{D}\right)^2 \delta^B \eta. \tag{60}$$

$$\varepsilon\,y_\mu \;=\; \frac{z\,(\tilde{a}\,z + \tilde{b})}{(\tilde{c}_1\,z + \tilde{d}_1)\,(\tilde{c}_2\,z + \tilde{d}_2)}, \tag{61}$$

where $\tilde{a} = \tilde{c}_1\,\tilde{c}_2$ and $\tilde{b} = (\xi/K_{13} + x_{32})\,x_{14}\,\eta + (x_{41} + \eta)\,x_{32}\,\xi$.

To obtain the expression for $\varepsilon\,y_\nu$, I coarse-grain module ν by eliminating the only remaining internal metabolite 2 and obtain

$$y_\nu = x_{15} + \frac{y_\mu\,x_{25}}{y_\mu/K_{12} + x_{25}}.$$

I then apply equation *Equation 6* with

$$C \;=\; \frac{y_\mu}{y_\nu}\frac{x_{25}^2}{\left(y_\mu/K_{12} + x_{25}\right)^2}, \tag{62}$$

$$H \;=\; -\frac{2\,y_\mu^2}{y_\nu\,K_{12}}\frac{x_{25}^2}{\left(u/K_{12} + x_{25}\right)^3}. \tag{63}$$

Figure *Figure 3*B illustrates how $\varepsilon\,y_\nu$ changes as a function of z. It was generated using the following matrix of rate constants:

$$x = \begin{pmatrix} 0 & 0.378 & 0.514 & 0.237 & 0 \\ 1.810 & 0 & 0 & 0 & 1.001 \\ 42.232 & 0 & 0 & z & 2.446 \\ 7.957 & 0 & z/2.44 & 0 & 0.259 \\ 0 & 6.982 & 0.994 & 0.257 & 0 \end{pmatrix}.$$

The Matlab code is available at https://github.com/skryazhi/epistasis_theory.

Next, I consider thress special cases of the toy network depicted in *Figure 3*A that relate this network to those in *Figure 3*C and D.

1. $z = 0$. According to *Equation 61*, $\varepsilon\,y_\mu = 0$ and hence $\varepsilon\,y_\nu = H/2C^2 \le 0$, with C and H given by *Equation 62* and *Equation 63*. It is easy to see that in this case the reactions $1 \leftrightarrow 3$ and $2 \leftrightarrow 4$ that are affected by mutations are strictly parallel because there is a simple path $1 \leftrightarrow 3 \leftrightarrow 2 \leftrightarrow 5$ that contains only reaction $1 \leftrightarrow 3$ and there is a simple path $1 \leftrightarrow 4 \leftrightarrow 2 \leftrightarrow 5$ that contains only reaction $2 \leftrightarrow 4$ (*Figure 3*C).

2. $x_{15} = x_{32} = x_{14} = 0$. In this case, module ν is a linear pathway. Therefore, $\varepsilon\,y_\nu = 1$ independently of z, as shown above. This fact can also be obtained directly from *Equation 61*, *Equation 62*, *Equation 63* and *Equation 6*.

3. $z \to \infty$. In this case, module μ becomes an effectively linear pathway because most of the metabolic flux between the I/O metabolites 1 and 2 passes through reaction $3 \leftrightarrow 4$. Thus, it follows from *Equation 61* that $\varepsilon\,y_\mu \to \tilde{a}/(\tilde{c}_1\,\tilde{c}_2) = 1$, as expected. Then, according to *Theorem 1*, $\varepsilon\,y_\nu \ge 1$.

Proof of *Theorem 1*

As discussed above, the key step towards the proof is *Proposition 2*, which states that the function f_1 belongs to one of three parameteric families, given by *Equation 29*, *Equation 30*, *Equation 31*. To complete the proof, I now explicitly evaluate the control coefficient C and the H in *Equation 6* for each of these functions and show that the inequalities in *Equation 7* and *Equation 8* hold for all parameter values.

Proof of **Theorem 1.** Let a be the effective reaction within higher-level module ν that represents
the lower-level module μ. To simplify notations, I denote $y_\mu \equiv \xi$. According to **Proposition 2**, the
functionional from of f_1 depends only on the topological class of the single-marked module (ν, a).
So, I consider the three classes one by one.

Case $(\nu, a) \in \mathcal{M}^{\mathrm{b}}$**.** From **Equation 29**, $C = \xi/y_\nu$ and $H = 0$. Therefore, inequalities in **Equation 7**
and **Equation 8** hold.

Case $(\nu, a) \in \mathcal{M}^{\mathrm{io}}$**.** From **Equation 30**,

$$C \;=\; \frac{\xi}{y_\nu}\left(\frac{w_{32}}{D}\right)^2, \tag{64}$$

$$H \;=\; -2\frac{\xi^2}{y_\nu}\frac{w_{32}^2}{D^3}\frac{1}{K_{13}} = -2\,C\,\frac{\xi/K_{13}}{D}, \tag{65}$$

1308 where $D = (\xi + \alpha)/K_{13} + w_{32}$. From **Equation 64**, it is clear that $C \geq 0$. Re-writing **Equation 64** as

$$C = \left(\frac{\xi\,w_{32}/D}{y_\nu}\right)\left(\frac{w_{32}}{D}\right)$$

1309 it is also clear that $C \leq 1$ since both ratios in this expression do not exceed 1. From **Equation 65**
1310 and the fact that $0 \leq C \leq 1$, it follows that $\bar{\varepsilon} \geq 0$. To show that $\bar{\varepsilon} \leq 1$, note that

$$D\,(1 - C) = \frac{\xi + \alpha}{K_{13}} + w_{32}\left(1 - \frac{\xi\,w_{32}/D}{y_\nu}\right) \geq \frac{\xi}{K_{13}}.$$

1311 Therefore,

$$\bar{\varepsilon} = \frac{\xi/K_{13}}{D\,(1 - C)} \leq 1.$$

1312 Therefore, inequalities in **Equation 7** and **Equation 8** hold.

1313   **Case** $(\nu, a) \in \mathcal{M}^{\mathrm{i}}$**.** I re-write **Equation 31** as

$$y_\nu = w_{12} + \frac{\tilde{A}\,u + \tilde{B}}{D},$$

1314 with $u = \xi + \alpha$, $D = \tilde{C}\,u + \tilde{D}$, $\tilde{A} = (w_{13} + w_{14})(w_{42} + w_{32}/K_{34})$, $\tilde{B} = (w_{31} + w_{32})\,w_{14}\,w_{42} + (w_{41} + w_{42})\,w_{13}\,w_{32}$,
1315 $\tilde{C} = (w_{31} + w_{32})/K_{34} + (w_{41} + w_{42})$, $\tilde{D} = (w_{31} + w_{32})(w_{41} + w_{42})$, which yields

$$C \;=\; \frac{\xi}{y_\nu}\frac{\tilde{A}\tilde{D} - \tilde{B}\tilde{C}}{D^2}, \tag{66}$$

$$H \;=\; -2\frac{\xi^2}{y_\nu}\frac{(\tilde{A}\tilde{D} - \tilde{B}\tilde{C})\,\tilde{C}}{D^3} = -2\,C\,\frac{\tilde{C}\,\xi}{D}. \tag{67}$$

1316 Next, it is straightforward to show that $\tilde{A}\tilde{D} - \tilde{B}\tilde{C} = (w_{41}\,w_{32} - w_{31}\,w_{42})^2/K_{31} \geq 0$, which implies that
1317 $C \geq 0$. To show that $C \leq 1$, I expand the denominator in **Equation 66** and obtain

$$y_\nu\,D^2 \geq (\tilde{A}\,u + \tilde{B})(\tilde{C}\,u + \tilde{D}) \geq u\,(\tilde{A}\,\tilde{D} + \tilde{B}\,\tilde{C}) \geq \xi\,(\tilde{A}\,\tilde{D} - \tilde{B}\,\tilde{C}).$$

1318 Therefore, numerator in **Equation 66** cannot exceed the denominator. The fact that $\bar{\varepsilon} \geq 0$ follows
1319 directly from **Equation 67** together with $C \leq 1$. To show that $\bar{\varepsilon} \leq 1$, first note that

$$y_\nu = w_{12} + \frac{\tilde{A}\,\xi}{D} + \frac{\tilde{A}\,\alpha + \tilde{B}}{D} \geq \frac{\tilde{A}\,\xi}{D}.$$

1320 Therefore,

$$D\,(1 - C) = \tilde{C}\,\xi + \tilde{C}\,\alpha + \tilde{D}\left(1 - \frac{\tilde{A}\,\xi/D}{y_\nu}\right) + \frac{\xi}{D}\frac{\tilde{B}\,\tilde{C}}{y_\nu} \geq \tilde{C}\,\xi.$$

1321 Hence,

$$\bar{\varepsilon} = \frac{\tilde{C}\,\xi}{D(1 - C)} \leq 1.$$

1322 Therefore, inequalities in **Equation 7** and **Equation 8** hold in this case as well, which completes the
1323 proof.   □

Proof of *Theorem 2*

Proving *Theorem 2* involves several auxiliary steps. First, I note that any two reactions a and b within any module μ can be either strictly serial, strictly parallel or serial-parallel. Then, *Proposition 4* and its *Corollary 4* establish that strictly parallel (serial) reactions in (μ, a, b) are also strictly parallel (serial) in a minimal module (μ', a, b), which is obtained from μ by eliminating all metabolites that do not participate in the marked reactions. Next, recall that in both modules (μ, a, b) and (μ', a, b) the same function f_2 maps the rate constants of two marked reactions onto module's effective rate constant (*Proposition 3*). Since the epistasis coefficient depends only on the shape of this function, we only need to consider all minimal modules in order to understand what kinds of epistasis may arise between mutations affecting strictly serial and strictly parallel reactions in any module. This is a big simplification because the number of different minimal topologies is finite and the parameteric families of function f_2 are known for all of them (see *Corollary 3*). As a consequence, to prove *Theorem 2*, we could in principle list all of the minimal topologies, identify those where the marked reactions are strictly serial or strictly parallel and evaluate the epistasis coefficient using *Equation 9* for every respective function f_2.

Unfortunately, the number of minimal topologies is very large, so that such brute-force approach would be quite cumbersome. In Generating topologies, I take a less cumbersome approach which is based on the realization that a strictly serial or strictly parallel relationship between two reactions cannot be altered by removing a third reaction from the module (*Proposition 5*). This implies that every minimal topology where the two reactions are strictly serial can be produced from another, more connected, "generating" topology by removing some reactions; and similarly for minimal modules where the reactions are strictly parallel (*Proposition 6*). All generating topologies can be identified by a simple algorithm given in *Appendix 3*.

Finally, in Topological relationship between reactions and epistasis, I prove *Theorem 2* in three steps. First, *Proposition 7* shows that $\varepsilon y_\mu \leq 0$ for any minimal module μ with any strictly parallel generating topology. Second, *Proposition 8* shows that that $\varepsilon y_\mu \geq 1$ for any minimal module μ with any strictly serial generating topology. Third, the proof of *Theorem 2* formally extends this argument to all modules with strictly serial and strictly parallel reactions.

Topological relationships between reactions within a module

Consider module μ with the I/O metabolites 1 and 2. As described in Notations and definitions, a simple path connecting two metabolites i and j within module μ is denoted by $p^\mu_{ij} = i \leftrightarrow k \leftrightarrow \ldots \leftrightarrow \ell \leftrightarrow j$. If such path contains reactions $a_1, a_2, \ldots$ and does not contain reactions $b_1, b_2, \ldots$, I denote it as $p^\mu_{ij}(a_1, a_2, \ldots, \bar{b}_1, \bar{b}_2, \ldots)$. I denote the set of all paths $p^\mu_{ij}(a_1, a_2, \ldots, \bar{b}_1, \bar{b}_2, \ldots)$ by $\mathcal{P}^\mu_{ij}(a_1, a_2, \ldots, \bar{b}_1, \bar{b}_2, \ldots)$.

According to *Lemma 1* proven in *Appendix 2*, any reaction in the module belongs to at least one simple path within module μ that connects the I/O metabolites. Mathematically, $\mathcal{P}^\mu_{12}(a) \neq \emptyset$ for any reaction $a \in R_\mu$. Thus, we can define different topological relationships between any two reactions within a module based on the existence of various paths that do or do not contain them. Consequently, we can classify any double-marked module (μ, a, b) based on the toplogocial relationship between its marked reactions. This classification is orthogonal to that given in *Table 1*.

Two reactions $a \in R_\mu$ and $b \in R_\mu$ are called *parallel* within module μ if there exists a simple path $p^\mu_{12}(a, \bar{b})$ and a simple path $p^\mu_{12}(b, \bar{a})$. Two reactions $a \in R_\mu$ and $b \in R_\mu$ are called *serial* within module μ if there exist at least one simple path $p^\mu_{12}(a, b)$. Two reactions are called *strictly parallel* within module μ if they are parallel but not serial, they are called *strictly serial* within module μ if they are serial but not parallel, and they are called *serial-parallel* within module μ if they are both serial and parallel. It is straightforward to show that there are no other logical possibilities for any two reactions to be anything other than strictly serial, strictly parallel or serial-parallel. This implies that two reactions are strictly parallel if they are not serial, and they are strictly serial if they are not parallel. If reactions a and b are serial, parallel, strictly serial, strictly parallel or serial-parallel within module μ, I also refer to the double-marked module (μ, a, b) as serial, parallel, etc. Since the relationship between reactions depends only on the topology of the module, but not on its rate

Figure 9. A counter example illustrating that the converse to claim 2 in *Proposition 4* may not be true.
Reactions a and b are parallel in (μ, a, b). CGP maps the double-marked module (μ, a, b) onto the minimal double-marked module (μ', a, b) where reactions a and b are not parallel.

constants, I also refer to the topology (R_μ, a, b) as serial, parallel, etc.

Recall that coarse-graining procedure $\mathrm{CG}^{\mu \setminus \{a,b\}}$ eliminates all metabolites internal to module μ

other than those participating in reactions a and b. If the double-marked module (μ, a, b) belongs

to the topological class $\mathcal{M}$, then, according to ***Proposition 3***, $\mathrm{CG}^{\mu \setminus \{a,b\}}$ maps (μ, a, b) onto a minimal

double-marked module (μ', a, b) from the same class $\mathcal{M}$. The following proposition, which is easy to

prove using Property #9 of the CGP (see ***Box 1***), establishes how this procedure alters the topological

relationship between reactions a and b.

Proposition 4. *Let (μ, a, b) be a double-marked module from the topological class $\mathcal{M}$ (**Table 1**) and let*

(μ', a, b) be the minimal double-marked module in $\mathcal{M}$ onto which (μ, a, b) is mapped by $\mathrm{CG}^{\mu \setminus \{a,b\}}$.

1. Reactions a and b are serial in (μ', a, b) if and only if they are serial in (μ, a, b).

2. If reactions a and b are parallel in (μ', a, b), then they are also parallel in (μ, a, b).

Note that the converse of the second claim in ***Proposition 4*** is not true. In other words, if

two reactions a and b are parallel in (μ, a, b), they may not be parallel in (μ', a, b). ***Figure 9*** shows a

counter-example illustrating this.

Corollary 4.   *1. If reactions a and b are strictly serial in (μ, a, b), they are also strictly serial in (μ', a, b).*

2. If reactions a and b are strictly parallel in (μ, a, b), they are also strictly parallel in (μ', a, b).

3. If reactions a and b are serial-parallel in (μ, a, b), they are either strictly serial or serial-parallel in

(μ', a, b).

Corollary 4 is an important step towards proving ***Theorem 2***. According to ***Proposition 3***, the

function that maps the rate constants ξ and η of the reactions a and b in module μ onto the effective

rate constant y_μ is the same function that maps the rate constants u and v of these reactions

in the minimal module μ' onto the effective rate constant $y_{\mu'}$. It then immediately follows from

Equation 9 that the epistasis coefficient εy_μ between mutations affecting reactions a and b in the

original module μ is the same as the epistasis coefficient $\varepsilon y_{\mu'}$ in the minimal module μ'. Now, if the

reactions a and b are strictly parallel in (μ, a, b), then, according to ***Corollary 4***, these reactions are

also strictly parallel in (μ', a, b). Hence, to demonstrate that $\varepsilon y_\mu \leq 0$ for any such double-marked

module (μ, a, b), it is sufficient to show that $\varepsilon y_{\mu'} \leq 0$ for all double-marked modules (μ', a, b) that are

minimal in $\mathcal{M}$ and where the reactions a and b are strictly parallel. And similarly for the strictly serial

reactions.

According to this logic, ***Theorem 2*** can be proven by identifying all double-marked modules that

are minimal in each of the topological classes listed in ***Table 1*** and where the marked reactions

are strictly serial or strictly parallel, evaluating epistasis for all of them, and showing that it is

non-positive, irrespectively of the rate constants of any reactions.

Generating topologies

Since the number of topologically distinct minimal double-marked modules is finite, the approach

outlined above is in principle feasible. Unfortunately, the number of topologies to be considered is

very large, so in practice it is very cumbersome. To avoid this complication, I take an alternative

approach that is based on the same key idea as the proof of ***Theorem 1***. Rather than considering

one by one each minimal topology where the marked reactions are strictly serial or strictly parallel
(and the corresponding parametric families of f_2), the idea is to identify the most connected minimal
topologies (and the corresponding largest parametric families of f_2) such that all the other minimal
topologies with the strictly serial or strictly parallel reactions (and the corresponding parametric
families) can obtained from them by removing reactions (i.e., setting some parameters to zero).

This idea can be implemented using the following observations. If the two marked reactions are
strictly parallel or strictly serial in a minimal module, then removing a third reaction from it does not
change this relationship. This statement is proven in *Proposition 5*. As a consequence, all minimal
modules in the topological classes $\mathcal{M}^{\text{b,io,IO}}$, $\mathcal{M}^{\text{b,i,}\emptyset}$ and $\mathcal{M}^{\text{io,io,IO}}$ must be strictly parallel because the
fully connected minimal topologies are strictly parallel (*Figure 8*). Similarly, all minimal modules in
the topological class $\mathcal{M}^{\text{io,io,I}}$ must be strictly serial because the fully connected minimal topology is
strictly serial (*Figure 8*). The fully connected minimal topologies in all other topological classes are
serial-parallel. If the two reactions are serial-parallel, removing a third reaction can change their
relationship into a strictly serial or strictly parallel one, depending on which reaction is removed, as
shown for example in *Figure 3*A, C and D. In fact, by removing reactions from the fully connected
minimal modules shown in *Figure 8*, it is easy to show that the topological classes $\mathcal{M}^{\text{io,io,}\emptyset}$, $\mathcal{M}^{\text{io,i,I}}$,
$\mathcal{M}^{\text{io,i,}\emptyset}$, $\mathcal{M}^{\text{i,i,I}}$, $\mathcal{M}^{\text{i,i,}\emptyset}$ contain both strictly serial and strictly parallel modules.

These observations suggests that adding reactions to a minimal module where the marked
reactions are strictly serial or strictly parallel will either change their relationship into serial-parallel
or will preserve their relationship until the minimal module is fully connected. Therefore, among
all minimal modules in a topological class, there must exist the most connected modules where
the marked reactions are strictly parallel or strictly serial, such that all other less connected strictly
serial or strictly parallel modules can be produced from the most connected ones by removal of
reactions. This statement is proven in *Proposition 6*. Such most connected strictly parallel and
strictly serial minimal topologies, which I refer to as "generating", are listed in *Table 2* and *Table 3*.
They define the largest parameteric families of functions f_2 which I then examine for the value of
εy_μ.

Proposition 5. *Let (μ, a, b) and (μ', a, b) be two minimal double-marked modules from the same topo-*
logical class whose sets of reactions are R_μ and $R_{\mu'}$, respectively, and $R_{\mu'} = R_\mu \setminus \{c\}$ where $c \in R_\mu \setminus \{a, b\}$.

1. *If reactions a and b are strictly parallel in (μ, a, b), they are also strictly parallel in (μ', a, b).*
2. *If reactions a and b are strictly serial in (μ, a, b), they are also strictly serial in (μ', a, b).*

Proof. Denote the I/O metabolites in both modules μ and μ' by 1 and 2. Since μ' and μ are topo-
logically identical except for μ' lacking one reaction c, it must be true that $\mathcal{P}_{12}^{\mu'}(a_1, a_2, \ldots, \bar{b}_1, \bar{b}_2, \ldots) \subseteq$
$\mathcal{P}_{12}^{\mu}(a_1, a_2, \ldots, \bar{b}_1, \bar{b}_2, \ldots)$ for any reactions $a_1, a_2, \ldots, b_1, b_2, \ldots$ from $R_{\mu'}$. In other words, there could only
be fewer paths connecting the I/O metabolites within module μ' compared to module μ. The rest of
the proof follows immediately from this fact and the definitions of strictly serial and strictly parallel
relatioships. $\square$

Next, I define a minimal topology as generating either if it is a fully connected topology (as in
topological classes $\mathcal{M}^{\text{b,io,IO}}$, $\mathcal{M}^{\text{b,i,}\emptyset}$ and $\mathcal{M}^{\text{io,io,IO}}$, $\mathcal{M}^{\text{io,io,I}}$) or if adding any reaction to it would make the
marked reactions serial-parallel.
Denote the sets of all double-marked topologies minimal in class $\mathcal{M}$ where the marked reactions
are strictly serial, strictly parallel and serial-parallel by by $\mathcal{R}_{\mathcal{M}}^{\text{ser}}$, $\mathcal{R}_{\mathcal{M}}^{\text{par}}$ and $\mathcal{R}_{\mathcal{M}}^{\text{sp}}$, respectively.

Definition 2. *Topology (R, a, b) minimal in $\mathcal{M}$ is called a* strictly serial generating topology *in $\mathcal{M}$ if it is*
strictly serial (i.e., $(R, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{ser}}$) and either it is fully connected (i.e., $R = R_{\mathcal{M}}$) or $(R \cup \{c\}, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{sp}}$ for
any reaction $c \in R_{\mathcal{M}} \setminus R$.

Definition 3. *Topology (R, a, b) minimal in $\mathcal{M}$ is called a* strictly parallel generating topology *in $\mathcal{M}$ if it*
is strictly parallel (i.e., $(R, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{par}}$) and either it is fully connected (i.e., $R = R_{\mathcal{M}}$) or $(R \cup \{c\}, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{sp}}$
for any reaction $c \in R_{\mathcal{M}} \setminus R$.

Clearly, a topological class $\mathcal{M}$ may have multiple generating topologies, and it is easy to show that every topological class has at least one generating topology. I denote the set of all strictly serial generating topologies for the class $\mathcal{M}$ by $\mathcal{G}_{\mathcal{M}}^{\text{ser}}$ and I denote the set of all strictly parallel generating topologies for class $\mathcal{M}$ by $\mathcal{G}_{\mathcal{M}}^{\text{par}}$. The following proposition justifies the name "generating topology". It states that any strictly serial minimal topology can be produced from some strictly serial generating topology by removing one or multiple reactions, and similarly for any strictly parallel minimal topology.

Proposition 6. *If (R, a, b) is a strictly parallel topology minimal in the topological class $\mathcal{M}$, then there exists a strictly parallel generating topology (R_g, a, b) in $\mathcal{M}$, such that $R \subseteq R_g$. If (R, a, b) is a strictly serial topology minimal in the topological class $\mathcal{M}$, then there exists a strictly serial generating topology (R_g, a, b) in $\mathcal{M}$, such that $R \subseteq R_g$.*

Proof. Suppose that $\mathcal{M}$ is one of the topological classes $\mathcal{M}^{\text{b,io,IO}}$, $\mathcal{M}^{\text{b,i,\emptyset}}$, or $\mathcal{M}^{\text{io,io,IO}}$. Since the fully connected minimal topology $(R_{\mathcal{M}}, a, b)$ is strictly parallel, it is a generating topology in $\mathcal{M}$. Then, according to *Proposition 5*, any topology (R, a, b) minimal in $\mathcal{M}$ is strictly parallel, and *Proposition 6* holds. By the same logic, *Proposition 6* holds for the topological class $\mathcal{M}^{\text{io,io,I}}$.

Suppose that $\mathcal{M}$ is one of the remaining topological classes $\mathcal{M}^{\text{io,io,\emptyset}}$, $\mathcal{M}^{\text{io,i,I}}$, $\mathcal{M}^{\text{io,i,\emptyset}}$, $\mathcal{M}^{\text{i,i,I}}$ or $\mathcal{M}^{\text{i,i,\emptyset}}$. Then the fully connected minimal topology $(R_{\mathcal{M}}, a, b)$ is serial-parallel. Suppose that (R, a, b) is strictly parallel. Then R must be a strict subset of $R_{\mathcal{M}}$, so that the set $C = R_{\mathcal{M}} \setminus R$ is not empty. Then, either $(R, a, b) \in \mathcal{G}_{\mathcal{M}}^{\text{par}}$ or $(R, a, b) \notin \mathcal{G}_{\mathcal{M}}^{\text{par}}$. If $(R, a, b) \in \mathcal{G}_{\mathcal{M}}^{\text{par}}$, the *Proposition 6* holds. Suppose that $(R, a, b) \notin \mathcal{G}_{\mathcal{M}}^{\text{par}}$. This implies that there exists a reaction $c_1 \in C$, such that $R_1 = R \cup \{c_1\}$ and $(R_1, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{par}}$ ((R_1, a, b) cannot be in $\mathcal{R}_{\mathcal{M}}^{\text{ser}}$ due to *Proposition 5*). There are three possibilities.

(a) $R_1 = R_{\mathcal{M}}$.
(b) $R_1 \subset R_{\mathcal{M}}$ and $(R_1, a, b) \in \mathcal{G}_{\mathcal{M}}^{\text{par}}$.
(c) $R_1 \subset R_{\mathcal{M}}$ and $(R_1, a, b) \notin \mathcal{G}_{\mathcal{M}}^{\text{par}}$.

Option (a) is not possible since $(R_1, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{par}}$ while $(R_{\mathcal{M}}, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{sp}}$. Option (b) implies that the *Proposition 6* holds. Option (c) implies that there exists a reaction $c_2 \in C \setminus \{c_1\}$, such that $R_2 = R_1 \cup \{c_2\}$ and $(R_2, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{par}}$, and we have the same three possibilities for R_2 as above. Thus, by induction, *Proposition 6* must hold. The proof is analogous if (R, a, b) is strictly serial. $\square$

Discovering all strictly serial and strictly parallel generating topologies in any given topological class $\mathcal{M}$ is straightforward because all minimal topologies within $\mathcal{M}$ can be produced by removing reactions from the unique fully connected topology minimal in $\mathcal{M}$ shown in *Figure 8*. In *Appendix 3*, I provide an algorithm that discovers all strictly serial and strictly parallel generating topologies by sequentially removing reactions from the fully connected minimal topology in each topological class. The code implementing this algorithm in Matlab is available at https://github.com/skryazhi/epistasis_theory. All strictly parallel generating topologies are listed in *Table 2* and all strictly serial generating topologies are listed in *Table 3*. They are also illustrated in *Figure 8* and *Figure 10* through *Figure 14*. I label each generating topology by a four-letter combination (see column 4 in *Table 2* and *Table 3*): the first three letters denote the topological class and the last letter (F, P or S) denotes the specific generating topology within that class. Letter "F" (stands for "full") denotes the fact that the reaction set in the generating topology is complete. Letters "P" (for "parallel") and "S" (stands for "serial") denote strictly parallel and strictly serial generating topologies, respectively; if there are a multiple generating topologies within the same class, they are distinguished by subindices, e.g., $\text{io}, \text{i}, \emptyset, \text{P}_1$; $\text{io}, \text{i}, \emptyset, \text{P}_2$, etc.

Topological relationship between reactions and epistasis

Each strictly serial and strictly parallel generating topology in a given class $\mathcal{M}$ (listed in *Table 2* and *Table 3*) is produced by removing reactions from the fully connected topology minimal in $\mathcal{M}$ (shown in *Figure 8*). This implies that the parametric family of function f_2 that corresponds to any

Table 2. Strictly parallel generating topologies.

Class	Marked reactions		Generating topology		Fig.
	a	b	ID	Excluded reactions	
$\mathcal{M}^{\mathrm{b,io,IO}}$	$1 \leftrightarrow 2$	$1 \leftrightarrow 3$	b, io, IO, F	$\emptyset$	8
$\mathcal{M}^{\mathrm{b,i,\emptyset}}$	$1 \leftrightarrow 2$	$3 \leftrightarrow 4$	b, i, $\emptyset$, F	$\emptyset$	8
$\mathcal{M}^{\mathrm{io,io,IO}}$	$1 \leftrightarrow 3$	$1 \leftrightarrow 4$	io, io, IO, F	$\emptyset$	8
$\mathcal{M}^{\mathrm{io,io,\emptyset}}$	$1 \leftrightarrow 3$	$2 \leftrightarrow 4$	io, io, $\emptyset$, P	$\{3 \leftrightarrow 4\}$	10
$\mathcal{M}^{\mathrm{io,i,I}}$	$1 \leftrightarrow 3$	$3 \leftrightarrow 4$	io, i, I, P	$\{2 \leftrightarrow 4\}$	11
$\mathcal{M}^{\mathrm{io,i,\emptyset}}$	$1 \leftrightarrow 3$	$4 \leftrightarrow 5$	io, i, $\emptyset$, P_1	$\{3 \leftrightarrow 4, 3 \leftrightarrow 5\}$	12
			io, i, $\emptyset$, P_2	$\{2 \leftrightarrow 5, 3 \leftrightarrow 5\}$	
			io, i, $\emptyset$, P_3	$\{2 \leftrightarrow 4, 2 \leftrightarrow 5\}$	
$\mathcal{M}^{\mathrm{i,i,I}}$	$3 \leftrightarrow 4$	$3 \leftrightarrow 5$	i, i, I, P_1	$\{2 \leftrightarrow 4, 2 \leftrightarrow 5\}$	13
			i, i, I, P_2	$\{1 \leftrightarrow 5, 2 \leftrightarrow 5\}$	
$\mathcal{M}^{\mathrm{i,i,\emptyset}}$	$3 \leftrightarrow 4$	$5 \leftrightarrow 6$	i, i, $\emptyset$, P_1	$\{3 \leftrightarrow 5, 3 \leftrightarrow 6, 4 \leftrightarrow 5, 4 \leftrightarrow 6\}$	14
			i, i, $\emptyset$, P_2	$\{1 \leftrightarrow 5, 1 \leftrightarrow 6, 2 \leftrightarrow 5, 2 \leftrightarrow 6\}$	
			i, i, $\emptyset$, P_3	$\{2 \leftrightarrow 4, 2 \leftrightarrow 6, 3 \leftrightarrow 6, 4 \leftrightarrow 5, 4 \leftrightarrow 6\}$	
			i, i, $\emptyset$, P_4	$\{2 \leftrightarrow 4, 2 \leftrightarrow 5, 2 \leftrightarrow 6, 4 \leftrightarrow 5, 4 \leftrightarrow 6\}$	
			i, i, $\emptyset$, P_5	$\{1 \leftrightarrow 6, 2 \leftrightarrow 4, 2 \leftrightarrow 5, 2 \leftrightarrow 6, 4 \leftrightarrow 6\}$	
			i, i, $\emptyset$, P_6	$\{1 \leftrightarrow 4, 1 \leftrightarrow 6, 2 \leftrightarrow 4, 2 \leftrightarrow 6, 4 \leftrightarrow 6\}$	
			i, i, $\emptyset$, P_7	$\{1 \leftrightarrow 4, 1 \leftrightarrow 5, 2 \leftrightarrow 3, 2 \leftrightarrow 6, 3 \leftrightarrow 5, 4 \leftrightarrow 6\}$	

Table 3. Strictly serial generating topologies.

Class	Marked reactions		Generating topologies		Fig.
	a	b	ID	Excluded reactions	
$\mathcal{M}^{\mathrm{io,io,I}}$	$1 \leftrightarrow 3$	$2 \leftrightarrow 3$	io, io, I, F	$\emptyset$	8
$\mathcal{M}^{\mathrm{io,io,\emptyset}}$	$1 \leftrightarrow 3$	$2 \leftrightarrow 4$	io, io, $\emptyset$, S	$\{2 \leftrightarrow 3\}$	10
$\mathcal{M}^{\mathrm{io,i,I}}$	$1 \leftrightarrow 3$	$3 \leftrightarrow 4$	io, i, I, S_1	$\{2 \leftrightarrow 3\}$	11
			io, i, I, S_2	$\{1 \leftrightarrow 4\}$	
$\mathcal{M}^{\mathrm{io,i,\emptyset}}$	$1 \leftrightarrow 3$	$4 \leftrightarrow 5$	io, i, $\emptyset$, S_1	$\{1 \leftrightarrow 4, 1 \leftrightarrow 5\}$	12
			io, i, $\emptyset$, S_2	$\{2 \leftrightarrow 3, 2 \leftrightarrow 4, 3 \leftrightarrow 5\}$	
			io, i, $\emptyset$, S_3	$\{1 \leftrightarrow 5, 2 \leftrightarrow 3, 2 \leftrightarrow 5\}$	
$\mathcal{M}^{\mathrm{i,i,I}}$	$3 \leftrightarrow 4$	$3 \leftrightarrow 5$	i, i, I, S_1	$\{1 \leftrightarrow 3, 2 \leftrightarrow 3\}$	13
			i, i, I, S_2	$\{2 \leftrightarrow 3, 2 \leftrightarrow 5, 4 \leftrightarrow 5\}$	
$\mathcal{M}^{\mathrm{i,i,\emptyset}}$	$3 \leftrightarrow 4$	$5 \leftrightarrow 6$	i, i, $\emptyset$, S_1	$\{2 \leftrightarrow 3, 2 \leftrightarrow 5, 2 \leftrightarrow 6, 4 \leftrightarrow 5, 4 \leftrightarrow 6\}$	14
			i, i, $\emptyset$, S_2	$\{1 \leftrightarrow 3, 1 \leftrightarrow 6, 2 \leftrightarrow 3, 2 \leftrightarrow 6, 4 \leftrightarrow 6\}$	

generating topology is obtained from *Equation 34* through *Equation 42* by setting some parameters w_{ij} to zero. In other words, these parametric families are known. Next, I prove *Proposition 7* that shows that $\varepsilon y_\mu \leq 0$ for every member of every parameteric family of f_2 that corresponds to a strictly parallel generating topology and the analogous *Proposition 8* for strictly serial topologies.

Now, any minimal strictly parallel topology can in turn be produced by removing reactions from some strictly parallel generating topology, and any minimal strictly serial topology can be produced by removing reactions from some strictly serial generating topology. This implies that any function f_2 that corresponds to any strictly parallel minimal topology belongs to the parametric family that corresponds to some strictly parallel generating topology; and any function f_2 that corresponds to any strictly serial minimal topology belongs to the parametric family that corresponds to some strictly serial generating topology. Therefore, Propositions 7 and 8 imply that $\varepsilon y_\mu \leq 0$ for *any* minimal strictly parallel topology and that $\varepsilon y_\mu \geq 1$ for *any* minimal strictly serial topology. The proof of *Theorem 2* is then concluded by recalling that every strictly parallel module is mapped onto its effective rate constant via function f_2 that corresponds to some minimal strictly parallel module, and similarly for strictly serial modules.

Proposition 7. *Let (μ, a, b) be a minimal double-marked module in the topological class $\mathcal{M}$, with u and v being the rates of reactions a and b, respectively, and let y be the effective rate constant of this module. Suppose that mutation A perturbs only reaction a by $\delta^A u$, and mutation B perturbs only reaction b by $\delta^B v$, such that $|\delta^A u| \ll 1$, $|\delta^B v| \ll 1$. If reactions a and b are strictly parallel in (μ, a, b), then $\varepsilon y \leq 0$.*

Proof. According to *Proposition 6*, the topology of module (μ, a, b) can be produced by removing reactions from some strictly parallel generating topology (R_g, a, b). Therefore, the function f_2 that maps u and v in this module onto its effective rate constant y belongs to the parametric family that corresponds to (R_g, a, b). According to *Equation 55*,

$$\varepsilon y = \frac{H_{uv}}{2\, C_u\, C_v},\tag{68}$$

where

$$C_u = \frac{u}{y}\frac{\partial f_2}{\partial u},\tag{69}$$

$$C_v = \frac{v}{y}\frac{\partial f_2}{\partial v},\tag{70}$$

$$H_{uv} = \frac{u\,v}{y}\frac{\partial^2 f_2}{\partial u\,\partial v}.\tag{71}$$

According to *Theorem 1*, $0 \leq C_u \leq 1$ and $0 \leq C_v \leq 1$. And since all $y > 0$, $u > 0$, $v > 0$, to prove *Proposition 7*, it is sufficient to show that $\frac{\partial^2 f_2}{\partial u\,\partial v} \leq 0$ for any member of any parametric family that corresponds to generating topologies listed in *Table 2*.

Generating topologies $\mathrm{b, io, IO, F}$ **and** $\mathrm{b, i, \emptyset, F}$ (*Figure 8*). According to *Equation 34* and *Equation 35*, $y = f_2(u, v) = u + \phi(v)$, which implies that $\frac{\partial^2 f_2}{\partial u\,\partial v} = 0$.

Generating topology $\mathrm{io, io, IO, F}$ (*Figure 8*). According to *Equation 37*,

$$y = f_2(u, v) = w_{12} + \frac{A\,u\,v + B\,u + B\,v}{D},$$

where $D = E\,u\,v + F\,u + G\,v + B$, $A = w_{42}/K_{13} + w_{32}/K_{14}$, $B = w_{32}\,w_{42} + w_{32}\,w_{43} + w_{34}\,w_{42}$, $E = 1/\left(K_{13}\,K_{14}\right)$, $F = \left(w_{42} + w_{43}\right)/K_{13}$, $G = \left(w_{32} + w_{34}\right)/K_{14}$. Therefore,

$$\frac{\partial^2 f_2}{\partial u\,\partial v} = -2\,\frac{w_{34}}{K_{14}}\,\frac{\left(w_{32}\,v/K_{14} + B\right)\left(w_{42}\,u/K_{13} + B\right)}{D^3} \leq 0.$$

Generating topology $\mathrm{io, io, \emptyset, P}$ (*Figure 10*). According to *Equation 38*, $y = f_2(u, v) = w_{12} + \phi(u) + \psi(v)$, which implies $\frac{\partial^2 f_2}{\partial u\,\partial v} = 0$.

Generating topology $\mathrm{io, i, I, P}$ (*Figure 11*). Notice that metabolite 4 together with reactions $\mathrm{I} \leftrightarrow 4$, a and b form a double-marked module (μ', a, b) whose I/O metabolites are 1 and 3 and which is

Figure 10. Graphical representation of strictly serial and strictly parallel generating topologies in the class $\mathcal{M}^{io,io,\emptyset}$. Fully connected topology $io, io, \emptyset, F$ is shown for reference (same as in *Figure 8*).

Figure 11. Graphical representation of strictly serial and strictly parallel generating topologies in class $\mathcal{M}^{io,i,I}$. Fully connected topology io, i, I, F is shown for reference (same as in *Figure 8*).

Figure 12. Graphical representation of strictly serial and strictly parallel generating topologies in class $\mathcal{M}^{io,i,\emptyset}$. Fully connected topology $io, i, \emptyset, F$ is shown for reference (same as in *Figure 8*).

Figure 13. Graphical representation of strictly serial and strictly parallel generating topologies in class $\mathcal{M}^{\mathrm{i,i,I}}$. Fully connected topology i, i, I, F is shown for reference (same as in *Figure 8*).

minimal in the topological calss $\mathcal{M}^{\mathrm{b,io,IO}}$. Denote the effective reaction rate of module μ' by y'. Therefore, $\varepsilon y' = 0$, as shown above. Since module μ' is contained in μ, by *Theorem 1*, $\varepsilon y \leq 0$.

Generating topology io, i, $\emptyset$, P_1 (*Figure 12*). According to Property 1 of the CGP (*Box 1*), module μ can be coarse-grained by first eliminating metabolite 3. In the resulting module μ', mutation A perturbs only the rate constant u' of the effective reaction $a' \equiv 1 \leftrightarrow 2$ (by Properties 2 and 4 of the CGP). Then, according to *Equation 50* and *Theorem 1*, $|\delta^A u'| \ll 1$. The double-marked module (μ', a', b) is minimal in the topological class $\mathcal{M}^{\mathrm{b,i,\emptyset}}$ which implies that $\varepsilon y = 0$, as shown above.

Generating topology io, i, $\emptyset$, P_2 (*Figure 12*). Module μ can be coarse-grained by first eliminating metabolite 5, which will result in a double-marked module (μ', a, b') that is minimal in the topological class $\mathcal{M}^{\mathrm{io,io,IO}}$. The rest of the proof for this topology is analogous to that for the topology io, i, $\emptyset$, P_1.

Generating topology io, i, $\emptyset$, P_3 (*Figure 12*). Notice that metabolites 4 and 5 together with reactions a, b, $1 \leftrightarrow 4$, $1 \leftrightarrow 5$, $3 \leftrightarrow 4$ and $3 \leftrightarrow 5$ form a double-marked module (μ', a, b) whose I/O metabolites are 1 and 3 and which is minimal in the topological calss $\mathcal{M}^{\mathrm{b,i,\emptyset}}$. The rest of the proof for this topology is analogous to that for the topology io, i, I, P.

Generating topology i, i, I, P_1 (*Figure 13*). Notice that metabolites 4 and 5 together with reactions a, b, $1 \leftrightarrow 3$, $1 \leftrightarrow 4$, $1 \leftrightarrow 5$ and $4 \leftrightarrow 5$ form a double-marked module (μ', a, b) whose I/O metabolites are 1 and 3 and which is minimal in the topological calss $\mathcal{M}^{\mathrm{io,io,IO}}$. The rest of the proof for this topology is analogous to that for the topology io, i, I, P.

Generating topology i, i, I, P_2 (*Figure 13*). Notice that metabolite 5 together with reactions a, b, and $4 \leftrightarrow 5$ form a double-marked module (μ', a, b) whose I/O metabolites are 3 and 4 and which is minimal in the topological calss $\mathcal{M}^{\mathrm{b,io,IO}}$. The rest of the proof for this topology is analogous to that for the topology io, i, I, P.

Generating topology i, i, $\emptyset$, P_1 (*Figure 14*). According to *Equation 42*, $y = f_2(u, v) = x_{12} + \phi(u) + \psi(v)$, which implies $\frac{\partial^2 f_2}{\partial u \, \partial v} = 0$.

Generating topology i, i, $\emptyset$, P_2 (*Figure 14*). Notice that metabolites 5 and 6 together with reactions a, b, $3 \leftrightarrow 5$, $3 \leftrightarrow 6$, $4 \leftrightarrow 5$ and $5 \leftrightarrow 6$ form a double-marked module (μ', a, b) whose I/O metabolites are 3 and 4 and which is minimal in the topological calss $\mathcal{M}^{\mathrm{b,i,\emptyset}}$. The rest of the proof for this topology is analogous to that for the topology io, i, I, P.

Generating topology i, i, $\emptyset$, P_3 (*Figure 14*). Module μ can be coarse-grained by first eliminating metabolites 4 and 6, which will result in a double-marked module (μ', a', b') that is minimal in the topological class $\mathcal{M}^{\mathrm{io,io,IO}}$. The rest of the proof for this topology is analogous to that for the topology io, i, $\emptyset$, P_1.

Generating topology i, i, $\emptyset$, P_4 (*Figure 14*). Module μ can be coarse-grained by first eliminating

Figure 14. Graphical representation of strictly serial and strictly parallel generating topologies in class $\mathcal{M}^{i,i,\emptyset}$. Fully connected topology $i, i, \emptyset, F$ is shown for reference (same as in *Figure 8*).

---

metabolite 4, which will result in a double-marked module (μ', a', b) that is minimal in the topological class $\mathcal{M}^{io,i,\emptyset}$ with a strictly parallel generating topology $io, i, \emptyset, P_3$. The rest of the proof for this topology is analogous to that for the topology $io, i, \emptyset, P_1$.

    Generating topology $i, i, \emptyset, P_5$ (*Figure 14*). Module μ can be coarse-grained by first eliminating metabolite 6, which will result in a double-marked module (μ', a, b') that is minimal in the topological class $\mathcal{M}^{i,i,I}$ with a strictly parallel generating topology i, i, I, P_1. The rest of the proof for this topology is analogous to that for the topology $io, i, \emptyset, P_1$.

    Generating topology $i, i, \emptyset, P_6$ (*Figure 14*). Notice that metabolites 4 and 6 together with reactions a, b, $3 \leftrightarrow 5$, $3 \leftrightarrow 6$, $4 \leftrightarrow 5$ form a double-marked module (μ', a, b) whose I/O metabolites are 3 and 5 and which is minimal in the topological calss $\mathcal{M}^{io,io,\emptyset}$. The rest of the proof for this topology is analogous to that for the topology io, i, I, P.

    Generating topology $i, i, \emptyset, P_7$ (*Figure 14*). Using *Equation 42*, I show in *Appendix 4* that

$$\frac{\partial^2 f_2}{\partial u \, \partial v} = \frac{2\,\beta}{K_{31}} \, \frac{\left(A_u\,v + B_u\right)\left(A_v\,u + B_v\right)}{\left(E_u\,v + F_u\right)^3}, \tag{72}$$

where A_v, B_v, E_u, F_u are all non-negative constants and $\beta \leq 0$. $\qquad\square$

Proposition 8. *Let (μ, a, b) be a minimal double-marked module in the topological class $\mathcal{M}$, with u and v being the rates of reactions a and b, respectively, and let y be the effective rate constant of this module. Suppose that mutation A perturbs only reaction a by $\delta^A u$, and mutation B perturbs only reaction b by $\delta^B v$, such that $|\delta^A u| \ll 1$, $|\delta^B v| \ll 1$. If reactions a and b are strictly serial in (μ, a, b), then $\varepsilon y \geq 1$.*

Proof. The logic of the proof is the same as for *Proposition 7*, i.e., it is sufficient to show that $\varepsilon y \geq 1$ for any double-marked module (μ, a, b) with any strictly serial generating topology listed in *Table 3*.

    Generating topology io, io, I, F (*Figure 8*). According to *Equation 36*,

$$y = f_2(u, v) = w_{12} + \frac{u\,v}{D} \tag{73}$$

where $D = u/K_{13} + v$. Therefore,

$$
\begin{aligned}
C_u &= \left(\frac{v}{D}\right)^2 \frac{u}{y}, \\
C_v &= \frac{1}{K_{12}} \left(\frac{u}{D}\right)^2 \frac{v}{y}, \\
H_{uv} &= \frac{2}{K_{12}} \frac{1}{y\,D} \left(\frac{u\,v}{D}\right)^2.
\end{aligned}
$$

Substituting these expressions into *Equation 68*, I obtain

$$
\varepsilon y = \frac{y}{u\,v/D} \geq 1
$$

because $y \geq u\,v/D$ according to *Equation 73*.

    Generating topology $\mathrm{io, io, \emptyset, S}$ (*Figure 10*). According to Property 1 of the CGP (*Box 1*), module μ can be coarse-grained by first eliminating metabolite 3. In the resulting module μ', mutation A perturbs only the rate constant u' of the effective reaction $a' \equiv 1 \leftrightarrow 4$ (by Properties 2 and 4 of the CGP). Then, according to *Equation 50* and *Theorem 1*, $|\delta^A u'| \ll 1$. The double-marked module (μ', a', b) is minimal in the topological class $\mathcal{M}^{\mathrm{io,io,I}}$ which implies that $\varepsilon y \geq 1$, as shown above.

    Generating topology $\mathrm{io, i, I, S_1}$ (*Figure 11*). Notice that metabolite 3 together with reactions a, b, and $1 \leftrightarrow 4$ form a double-marked module (μ', a, b) whose I/O metabolites are 1 and 4 and which is minimal in the topological calss $\mathcal{M}^{\mathrm{io,io,I}}$. Therefore, if the effective reaction rate of module μ' is y', $\varepsilon y' \geq 1$, as shown above. According to *Equation 50*, *Equation 51* and *Theorem 1*, $|\delta^A y'| \ll 1$, $|\delta^B y'| \ll 1$. Since module μ' is contained in μ, by *Theorem 1*, $\varepsilon y \geq 1$.

    Generating topology $\mathrm{io, i, I, S_2}$ (*Figure 11*). Module μ can be coarse-grained by first eliminating metabolite 4, which results in a double-marked module (μ', a, b') that is minimal in the topological class $\mathcal{M}^{\mathrm{io,io,I}}$. The rest of the proof for this topology is analogous to that for the topology $\mathrm{io, io, \emptyset, S}$.

    Generating topology $\mathrm{io, i, \emptyset, S_1}$ (*Figure 12*). Module μ can be coarse-grained by first eliminating metabolites 4 and 5, which results in a double-marked module (μ', a, b') that is minimal in the topological class $\mathcal{M}^{\mathrm{io,io,I}}$. The rest of the proof for this topology is analogous to that for the topology $\mathrm{io, io, \emptyset, S}$.

    Generating topology $\mathrm{io, i, \emptyset, S_2}$ (*Figure 12*). Notice that metabolites 3 and 4 together with reactions a, b, $1 \leftrightarrow 4$, $1 \leftrightarrow 5$ and $3 \leftrightarrow 4$ form a double-marked module (μ', a, b) whose I/O metabolites are 1 and 5 and which is minimal in the topological calss $\mathcal{M}^{\mathrm{io,io,\emptyset}}$ and whose reaction set is the strictly serial generating set $\mathrm{io, io, \emptyset, S}$. The rest of the proof for this topology is analogous to that for the topology $\mathrm{io, i, I, S_1}$.

    Generating topology $\mathrm{io, i, \emptyset, S_3}$ (*Figure 12*). Notice that metabolites 3 and 5 together with reactions a, b, $1 \leftrightarrow 4$, $3 \leftrightarrow 4$ and $3 \leftrightarrow 5$ form a double-marked module (μ', a, b) whose I/O metabolites are 1 and 4 and which is minimal in the topological calss $\mathcal{M}^{\mathrm{io,io,\emptyset}}$ and whose reaction set is the strictly serial generating set $\mathrm{io, io, \emptyset, S}$. The rest of the proof for this topology is analogous to that for the topology $\mathrm{io, i, I, S_1}$.

    Generating topology $\mathrm{i, i, I, S_1}$ (*Figure 13*). Notice that metabolite 3 together with reactions a, b, and $4 \leftrightarrow 5$ form a double-marked module (μ', a, b) whose I/O metabolites are 4 and 5 and which is minimal in the topological calss $\mathcal{M}^{\mathrm{io,io,I}}$. The rest of the proof for this topology is analogous to that for the topology $\mathrm{io, i, I, S_1}$.

    Generating topology $\mathrm{i, i, I, S_2}$ (*Figure 13*). Notice that metabolites 3 and 5 together with reactions a, b, $1 \leftrightarrow 3$, $1 \leftrightarrow 4$ and $1 \leftrightarrow 5$ form a double-marked module (μ', a, b) whose I/O metabolites are 1 and 4 and which is minimal in the topological calss $\mathcal{M}^{\mathrm{io,i,I}}$ and whose reaction set is the strictly serial generating set $\mathrm{io, i, I, S_2}$. The rest of the proof for this topology is analogous to that for the topology $\mathrm{io, i, I, S_1}$.

    Generating topology $\mathrm{i, i, \emptyset, S_1}$ (*Figure 14*). Module μ can be coarse-grained by first eliminating metabolites 5 and 6, which results in a double-marked module (μ', a, b') that is minimal in the

topological class $\mathcal{M}^{\mathrm{io,i,I}}$ with the strictly serial generating topology $\mathrm{io, i, I, S_1}$. The rest of the proof for this topology is analogous to that for the topology $\mathrm{io, io, \emptyset, S}$.

Generating topology $\mathrm{i, i, \emptyset, S_2}$ (*Figure 14*). Module μ can be coarse-grained by first eliminating metabolite 6, which results in a double-marked module (μ', a, b') that is minimal in the topological class $\mathcal{M}^{\mathrm{i,i,I}}$ with the strictly serial generating topology $\mathrm{i, i, I, S_1}$. The rest of the proof for this topology is analogous to that for the topology $\mathrm{io, io, \emptyset, S}$. □

Proof of **Theorem 2.** According to *Proposition 3*, the coarse-graining procedure $\mathrm{CG}^{\mu \backslash \{a,b\}}$ maps the double-marked module (μ, a, b) onto a double-marked module (μ', a, b) that is minimal in the same topological class as (μ, a, b), and the rates u, v of reactions a, b in μ' are given by linear relations in *Equation 32* and *Equation 33*. Clearly, $|\delta^A u| \ll 1$ and $|\delta^B v| \ll 1$. Furthermore, none of the other reaction rates w_{ij} in μ' depend on ξ or η, so that $\delta^A w_{ij} = 0$ and $\delta^B w_{ij} = 0$ for all w_{ij} other than u and v, and $\varepsilon w_{ij} = 0$ for all w_{ij} including u and v. It then follows from *Proposition 3* that $\varepsilon y_\mu = \varepsilon y_{\mu'}$.

Now, according to *Corollary 4*, if reactions a and b are strictly parallel in (μ, a, b), they are also strictly parallel in (μ', a, b). Therefore, by *Proposition 7*, $\varepsilon y_{\mu'} \leq 0$. Analogously, if reactions a and b are strictly serial in (μ, a, b), they are also strictly serial in (μ', a, b). Therefore, by *Proposition 8*, $\varepsilon y_{\mu'} \geq 1$. □

Sensitivity of *Theorem 1* and *Theorem 2* with respect to the magnitude of mutational effects

According to *Proposition 2*, function f_1 for any module belongs to one of three parametric families, which correspond to the three minimal fully connected modules shown in *Figure 7*. As mentioned in Results, for modules in the class $\mathcal{M}^{\mathrm{b}}$, function f_1 is linear, so that the claims of *Theorem 1* continue to hold for mutations with finite effects. To evaluate the sensitivity of *Theorem 1* with respect to the effect sizes of mutations for the topological classes $\mathcal{M}^{\mathrm{io}}$ and $\mathcal{M}^{\mathrm{i}}$, I generated 1000 minimal single-marked modules ν from each of these topological classes with random parameters. Evaluating only minimal modules is sufficient because for any module from a given topological class there exists a minimal module from the same class, such that both of them map the lower level phenotype y_μ onto the higher-level phenotype y_ν via the same function f_1 (see *Proposition 2*).

To this end, I drew each x_{ij} ($i < j$) from a mixture of a point measure at 0 (with weight 0.25) and an exponential distribution with mean 1 (with weight 0.75). The point measure at 0 ensures that minimal modules that are not fully connected are represented in the sample. I drew each K_{ij} ($i < j$) as a ratio of two random numbers from an exponential distribution with mean 1. As a result, the distribution of non-zero x_{ij} values had the interdecile range of $(5.7 \times 10^{-2}, 3.91)$ with median 0.65.

I denote the effective rate constant of the reaction that represents the lower-level module μ by $\xi \equiv y_\mu$. In modules from the topological class $\mathcal{M}^{\mathrm{io}}$, it is reaction $1 \leftrightarrow 3$ and in modules from the topological class $\mathcal{M}^{\mathrm{i}}$, it is the reaction $3 \leftrightarrow 4$. I perturbed ξ by two mutations A and B with relative effects $\delta^A \xi$ and $\delta^B \xi$ and epistasis $\varepsilon \xi$. I chose nine different pairs of mutational effects $(\delta^A \xi, \delta^B \xi)$: $(-0.01, -0.01)$, $(-0.1, -0.1)$, $(-0.5, -0.5)$, $(0.01, 0.01)$, $(0.1, 0.1)$, $(0.5, 0.5)$, $(-0.01, 0.01)$, $(-0.1, 0.1)$, $(-0.5, 0.5)$, and 16 different values of $\varepsilon \xi$ ranging from -1 to 2 with an increment of 0.2. Since the rate constant ξ^{AB} of the double mutant cannot be negative, I skipped those combinations of perturbations and epistasis values for which $\delta^A \xi + \delta^B \xi + 2 \left(\varepsilon \xi \right) \left(\delta^A \xi \right) \left(\delta^B \xi \right) < -1$. I then computed the resulting values $\delta^A y_\nu$, $\delta^B y_\nu$ and εy_ν at the level of the effective rate constant y_ν of the higher-level module ν.

Using these data, I inferred the function ϕ that maps lower-level epistasis $\varepsilon \xi$ onto higher-level epistasis εy_ν, as follows. For any minimal single-marked module from the topological classes $\mathcal{M}^{\mathrm{io}}$ or $\mathcal{M}^{\mathrm{i}}$, the effective rate constant y_ν can be written as

$$y_\nu = x_{12} + \frac{\tilde{A} \xi + \tilde{B}}{D},$$

where $D = \tilde{C} \xi + \tilde{D}$ and $\tilde{A} = x_{32}$, $\tilde{B} = 0$, $\tilde{C} = 1/K_{13}$, $\tilde{D} = x_{32}$ for modules from the topological class $\mathcal{M}^{\mathrm{io}}$ (see *Equation 30*), and $\tilde{A} = \left(x_{13} + x_{14} \right) \left(x_{42} + x_{32}/K_{34} \right)$, $\tilde{B} = \left(x_{31} + x_{32} \right) x_{14} x_{42} + \left(x_{41} + x_{42} \right) x_{13} w_{32}$,

1683 $\tilde{C} = (x_{31} + x_{32})/K_{34} + (x_{41} + x_{42})$, $\tilde{D} = (x_{31} + x_{32})(x_{41} + x_{42})$ for modules from the topological class $\mathcal{M}^i$

1684 (see *Equation 31*). Therefore, for any perturbation $\delta\xi$, we have

$$\delta y_v = \frac{\tilde{A}\,\tilde{D} - \tilde{B}\,\tilde{C}}{D^2} \frac{\xi}{y_v} \frac{\delta\xi}{1 + (\tilde{C}\,\xi/D)\,\delta\xi}.$$

1685 Since $\delta^{AB}\xi$ is a linear function of $\varepsilon\xi$, $\delta^{AB}y_v$ is a hyperbolic function of $\varepsilon\xi$. Therefore, εy_v is also a

1686 hyperbolic function of $\varepsilon\xi$,

$$\varepsilon y_v = \phi(\varepsilon\xi) = a - \frac{b}{\varepsilon\xi + c}, \tag{74}$$

where constants a, b and c depend on the parameters of module v and on the mutational effect

sizes $\delta^A\xi$ and $\delta^B\xi$. I numerically calculated these parameters for each sampled module and each

pair of mutational effects.

The main results of *Theorem 1* are that, when the effects of mutations are infinitesimal, the map

ϕ has a fixed point $\bar{\varepsilon}$, this fixed point is located between 0 and 1, and it is unstable. I use equation

Equation 74 to test whether these statements also hold when the effects of mutations are finite.

Specifically, it is easy to see that the map ϕ has a fixed point $\bar{\varepsilon}$ if the discriminant $d = (a-c)^2 - 4(b-ac)$

is positive. In this case, I designate $\bar{\varepsilon}$ as the one of two roots $1/2\left(a - c \pm \sqrt{d}\right)$ that is closer to zero.

I then check whether this fixed point is located between 0 and 1. I check whether it is unstable by

comparing the derivative of ϕ at $\bar{\varepsilon}$ with 1.

According to *Proposition 6*, function f_2 for any module where the reactions affected by mutations

are strictly parallel belongs to one of 17 parameteric families, which correspond to the strictly

parallel generating topologies listed in *Table 2*. And similarly, function f_2 for any module where

the reactions affected by mutations are strictly serial belongs to one of 11 parameteric families,

which correspond to the strictly serial generating topologies listed in *Table 3*. Therefore, to evaluate

the sensitivity of *Theorem 2* with respect to the effect sizes of mutations I generated 10^4 double-

marked modules (μ, a, b) with each of the strictly serial and strictly parallel topologies with random

parameters. I drew x_{ij} and K_{ij} as described above. I chose the same nine pairs of mutational

effects $(\delta^A\xi, \delta^B\eta)$ as above, where ξ and η are the rate constants of reactions affected by mutations

A and B: $(-0.01, -0.01)$, $(-0.1, -0.1)$, $(-0.5, -0.5)$, $(0.01, 0.01)$, $(0.1, 0.1)$, $(0.5, 0.5)$, $(-0.01, 0.01)$, $(-0.1, 0.1)$,

$(-0.5, 0.5)$.

I found that, for some modules, individual mutational perturbations $\delta^A y_\mu$ and/or $\delta^B y_\mu$ at the

level of the whole module were too small, which resulted in numerical instabilities. To avoid them, I

calculated epistasis εy_μ only for cases where the effects of both mutations $\delta^A y_\mu$ and $\delta^B y_\mu$ exceeded

the precision threshold of 10^{-5}. As a result, I evaluated epistasis in less than 10^4 modules per

generating topology and pair of mutational effects, but this number never fell below 1000. When

comparing the values of epistasis with 0 and 1, I used the same precision threshold of 10^{-5} to

avoid numerical problems. In addition, I found that for mutations affecting strictly serial reactions

there is a substantial fraction of modules where εy_μ falls between 0.99 and 1 (see *Figure 4–Figure*

Supplement 3). This is not a numerical artifact, but probably reflects real clustering of epistasis

coefficients around 1, which is expected for the linear pathway irrespective of its parameters (see

Calculation of epistasis in simple modules).

The Matlab code for this analysis is available at https://github.com/skryazhi/epistasis_theory.

Kinetic model of glycolysis

I downloaded the kinetic metabolic model of *E. coli* glycolysis by *Chassagnole et al.* (*2002*) from the

BioModels database (*Malik-Sheriff et al., 2019*) on September 15, 2015 (model ID BIOMD0000000051).

I used the Matlab SimBiology toolbox to interpret the model. To validate the model, I simulated it

for 40 seconds and reproduced Figures 4 and 5 from (*Chassagnole et al., 2002*). The Matlab code is

available at https://github.com/skryazhi/epistasis_theory.

Modifications to the original model

I simplified and modified the model by (a) fixing the concentrations of ATP, ADP, AMP, NADPH,
NADP, NADH, NAD at their steady-state values given in Table V of (*Chassagnole et al., 2002*) and (b)
removing dilution by growth. I then created four models of sub-modules of glycolysis by retaining
the subsets of metabolites and enzymes shown in *Figure 5–Figure Supplement 1* and *Table 4*
and removing other metabolites and enzymes. Each sub-module has one input and one output
metabolite. Note that, since some reactions are irreversible, it is important to distinguish the input
metabolite from the output metabolite. The concentrations of the input and the output metabolites
in each model are held constant at their steady-state values given in *Table 4*. I defined the flux
through the sub-module as the flux towards the output metabolite contributed by the sub-module
(*Table 4*). This flux is the equivalent of the quantitative phenotype y_μ of a module in the analytical
model. In addition, I made the following modifications specific to individual sub-modules.

1. In the FULL model, the stoichiometry of the PTS reaction was changed to

$$[\text{Ext glu}] + [\text{pep}] \leftrightarrow [\text{g6p}] + [\text{pyr}]$$

and the value of the constant $K_{\text{PTS,a1}}$ was set to 0.02 mM, based on the values found in the
literature (*Stock et al., 1982*; *Natarajan and Srienc, 1999*).
2. In all models other than FULL, the extracellular compartment was deleted.
3. In all models, the concentrations of the I/O metabolites were set to values shown in *Table 4*,
which are the steady-state concentrations achieved in the FULL model with the concentration
of extracellular glucose being 2 µM and pyruvate concentration being 10 µM.

Table 4. Definition of modules in the glycolysis network shown in *Figure 5–Figure Supplement 1*. Enzyme abbreviations are listed in *Table 6*. Metabolite abbreviations are listed in *Table 5*.

Model	Internal metabolites	Concentrations of I/O metabolites	Reactions	Output flux
UGPP	6pg, dhap, e4p, f6p, fdp, rib5p, ribu5p, sed7p, xyl5p	[g6p] = 3.82 mM, [gap] = 0.44 mM	ALDO, G6PDH, PFK, PGDH, PGI, Ru5P, R5PI, TA, TIS, TKa, TKb	$J_{\text{ALDO}} + J_{\text{TIS}} + J_{\text{TKb}} + J_{\text{TKa}} - J_{\text{TA}}$
LG	2pg, 3pg, pgp	[gap] = 0.44 mM, [pep] = 0.08 mM	ENO, GAPDH, PGK, PGM	J_{ENO}
GPP	all in UGPP and in LG, gap	[g6p] = 3.82 mM, [pep] = 0.08 mM	all in UGPP and in LG	J_{ENO}
FULL	all in GPP, g6p, pep	[Ext glu] = 2 µM, [pyr] = 10 µM	all in GPP, PTS, PK, PEPCxyl	$J_{\text{PK}} + J_{\text{PTS}}$

Calculation of flux control coefficients and epistasis coefficients

I calculate the first- and second-order flux control coefficients (FCC) C_i and H_{ij} for flux J with respect
to reactions i and j as follows (see *Equation 45* and *Equation 46*). I perturb the $r_{\text{max},i}$ of reaction i by
factor between 0.75 and 1.25 (10 values in a uniformly-spaced grid), such that $\delta r_{\text{max},i} \in [-0.25, 0.25]$.
Then, I obtain the steady-state flux J' in each perturbed model and calculate the flux perturbations
$\delta J = J'/J^0 - 1$, where J^0 is the corresponding flux in the unperturbed model. Then, to obtain C_i
and H_{ii}, I fit the linear model

$$\delta J \sim C_i \left(\delta r_{\text{max},i} \right) + \frac{H_{ii}}{2} \left(\delta r_{\text{max},i} \right)^2$$

1752 by least squares. If the estimated value of C_i was below 10^{-4} for a given sub-module, I set C_i to zero
1753 and exclude this reaction from further consideration in that sub-module because it does not affect

Table 5. Names of metabolites used in the kinetic model of glycolysis.

2pg	2-phosphoglycerate
3pg	3-phosphoglycerate
6pg	6-phosphogluconate
dhap	dihydroxyacetonephosphate
e4p	erythrose-4-phosphate
f6p	fructose-6-phosphate
fdp	fructose-1,6-bisphosphate
g6p	glucose-6-phosphate
gap	glyceraldehyde-3-phosphate
glu	glucose
pep	phosphoenolpyruvate
pgp	1,3-diphosphoglycerate
pyr	pyruvate
rib5p	ribose-5-phosphate
ribu5p	ribulose-5-phosphate
sed7p	sedoheptulose-7-phosphate
xyl5p	xylulose-5-phosphate

Table 6. Names of enzymes used in the kinetic model of glycolysis.

ALDO	aldolase
ENO	enolase
G6PDH	glucose-6-phosphate dehydrogenase
GAPDH	glyceraldehyde-3-phosphate dehydrogenase
PFK	phosphofructokinase
PGDH	6-phosphogluconate dehydrogenase
PGI	glucose-6-phosphateisomerase
PGK	phosphoglycerate kinase
PGM	phosphoglycerate mutase
PEPCxyl	PEP carboxylase
PK	pyruvate kinase
PTS	phosphotransferase system
R5PI	ribose-phosphateisomerase
Ru5P	ribulose-phosphate epimerase
TA	transaldolase
TIS	triosephosphate isomerase
TKa	transketolase, reaction a
TKb	transketolase, reaction b

1754 flux to the degree that is accurately measurable. If the estimated value of H_{ii} is below 10^{-4}, I set H_{ii}
1755 to zero.

1756 To calculate the non-diagonal second-order control coefficients H_{ij}, I create a 4×4 grid of
1757 perturbations of $\delta r_{\max,i}$ and $\delta r_{\max,j}$ and calculate the resulting flux perturbations δJ (16 perturbations
1758 total). Since C_i, C_j, H_{ii} and H_{jj} are known, I obtain H_{ij}, by regressing

$$\delta J - \left(C_i \left(\delta r_{\max,i} \right) + \frac{H_{ii}}{2} \left(\delta r_{\max,i} \right)^2 \right) - \left(C_j \left(\delta r_{\max,j} \right) + \frac{H_{jj}}{2} \left(\delta r_{\max,j} \right)^2 \right)$$

1759 against

$$\left(\delta r_{\max,i} \right) \left(\delta r_{\max,j} \right).$$

1760 If the estimated value of H_{ij} is below 10^{-4}, I set H_{ij} to zero. I estimate the epistasis coefficient εJ
1761 between mutations affecting reactions i and j as

$$\varepsilon J = \frac{H_{ij}}{2 \, C_i \, C_j}.$$

Establishing the topological relationships between pairs of reactions
To establish the topological relationship (strictly serial, strictly parallel or serial-parallel) between two
reactions, I consider the smallest module (LG, UGPP, GPPP or FULL) which contains both reactions. I
then manually identify whether there exists a simple path connecting the input metabolite with
the output metabolite for that module that passes through both reactions. (Note that, since some
reactions are irreversible in this model, it is important to distinguish the input metabolite from the
output metabolite). If such path does not exist, I classify the topological relationship between the
two reactions as strictly parallel. If such path exists, I check if there are two paths connecting the
input to the output metabolites such that each path contains only one of the two focal reactions. If
such paths do not exist, I classify the topological relationship between the two reactions as strictly
serial. Otherwise, I classify it as serial-parallel.

Acknowledgements
I thank Chris Marx for discussions that stimulated this work, David McCandlish for making me
aware of the Kron reduction, Daniel P. Rice, the Kryazhimskiy lab, and the reviewers for feedback on
the manuscript. This work was supported by the BWF Career Award at Scientific Interface (Grant
1010719.01), the Alfred P. Sloan Foundation (Grant FG-2017-9227), the Hellman Foundation and
NIH (Grant 1R01GM137112).

References
Alzoubi D, Desouki AA, Lercher MJ. Flux balance analysis with or without molecular crowding fails to predict
two thirds of experimentally observed epistasis in yeast. Scientific reports. 2019; 9(1):1–9.

Babu M, Arnold R, Bundalovic-Torma C, Gagarinova A, Wong KS, Kumar A, Stewart G, Samanfar B, Aoki H, Wagih
O, et al. Quantitative genome-wide genetic interaction screens reveal global epistatic relationships of protein
complexes in *Escherichia coli*. PLoS Genet. 2014; 10(2).

Bagheri HC, Wagner GP. Evolution of dominance in metabolic pathways. Genetics. 2004; 168(3):1713–1735.

Bagheri-Chaichian H, Hermisson J, Vaisnys JR, Wagner GP. Effects of epistasis on phenotypic robustness in
metabolic pathways. Math Biosci. 2003; 184(1):27–51.

Baryshnikova A, Costanzo M, Myers CL, Andrews B, Boone C. Genetic interaction networks: toward an
understanding of heritability. Annu Rev of Genomics Hum Genet. 2013; 14:111–133.

Bassik MC, Kampmann M, Lebbink RJ, Wang S, Hein MY, Poser I, Weibezahn J, Horlbeck MA, Chen S, Mann M,
et al. A systematic mammalian genetic interaction map reveals pathways underlying ricin susceptibility. Cell.
2013; 152(4):909–922.

Bellay J, Atluri G, Sing TL, Toufighi K, Costanzo M, Ribeiro PSM, Pandey G, Baller J, VanderSluis B, Michaut M, et al. Putting genetic interactions in context through a global modular decomposition. Genome Res. 2011; 21(8):1375–1387.

Blount ZD, Borland CZ, Lenski RE. Historical contingency and the evolution of a key innovation in an experimental population of Escherichia coli. Proc Natl Acad Sci USA. 2008; 105(23):7899–7906.

Boyle EA, Li YI, Pritchard JK. An expanded view of complex traits: from polygenic to omnigenic. Cell. 2017; 169(7):1177–1186.

Bridgham JT, Ortlund EA, Thornton JW. An epistatic ratchet constrains the direction of glucocorticoid receptor evolution. Nature. 2009; 461(7263):515–519.

Celaj A, Gebbia M, Musa L, Cote AG, Snider J, Wong V, Ko M, Fong T, Bansal P, Mellor JC, Nguyen GSM, Zhou S, Wang L, Stagljar NKI, Suzuki Y, Yachie N, Roth FP. Highly Combinatorial Genetic Interaction Analysis Reveals a Multi-Drug Transporter Influence Network. Cell Syst. 2020; 10(1):25–38.

Chassagnole C, Noisommit-Rizzi N, Schmid JW, Mauch K, Reuss M. Dynamic modeling of the central carbon metabolism of *Escherichia coli*. Biotechnol Bioeng. 2002; 79(1):53–73.

Chiu HC, Marx CJ, Segrè D. Epistasis from functional dependence of fitness on underlying traits. Proc R Soc B. 2012; 279(1745):4156–4164.

Chou HH, Chiu HC, Delaney NF, Segrè D, Marx CJ. Diminishing returns epistasis among beneficial mutations decelerates adaptation. Science. 2011; 332(6034):1190–1192.

Chou HH, Delaney NF, Draghi JA, Marx CJ. Mapping the fitness landscape of gene expression uncovers the cause of antagonism and sign epistasis between adaptive mutations. PLoS Genet. 2014; 10(2).

Clark A. Mutation-selection balance and metabolic control theory. Genetics. 1991; 129(3):909–923.

Collins SR, Miller KM, Maas NL, Roguev A, Fillingham J, Chu CS, Schuldiner M, Gebbia M, Recht J, Shales M, et al. Functional dissection of protein complexes involved in yeast chromosome biology using a genetic interaction map. Nature. 2007; 446(7137):806–810.

Cornish-Bowden A. Fundamentals of enzyme kinetics. John Wiley & Sons; 2013.

Costanzo M, Baryshnikova A, Bellay J, Kim Y, Spear ED, Sevier CS, Ding H, Koh JL, Toufighi K, Mostafavi S, et al. The genetic landscape of a cell. Science. 2010; 327(5964):425–431.

Costanzo M, VanderSluis B, Koch EN, Baryshnikova A, Pons C, Tan G, Wang W, Usaj M, Hanchard J, Lee SD, et al. A global genetic interaction network maps a wiring diagram of cellular function. Science. 2016; 353(6306):aaf1420.

Csete M, Doyle J. Bow ties, metabolism and disease. Tr Biotech. 2004; 22(9):446–450.

Das SG, Direito SO, Waclaw B, Allen RJ, Krug J. Predictable properties of fitness landscapes induced by adaptational tradeoffs. Elife. 2020; 9:e55155.

Davierwala AP, Haynes J, Li Z, Brost RL, Robinson MD, Yu L, Mnaimneh S, Ding H, Zhu H, Chen Y, et al. The synthetic genetic interaction spectrum of essential genes. Nat Genet. 2005; 37(10):1147–1152.

De Visser JAG, Krug J. Empirical fitness landscapes and the predictability of evolution. Nat Rev Genet. 2014; 15(7):480–490.

Dean AM. Selection and neutrality in lactose operons of *Escherichia coli*. Genetics. 1989; 123(3):441–454.

Dean AM, Dykhuizen DE, Hartl DL. Fitness as a function of β-galactosidase activity in *Escherichia coli*. Genetics Research. 1986; 48(1):1–8.

DePristo MA, Weinreich DM, Hartl DL. Missense meanderings in sequence space: a biophysical view of protein evolution. Nature Reviews Genetics. 2005; 6(9):678–687.

Dixon SJ, Costanzo M, Baryshnikova A, Andrews B, Boone C. Systematic mapping of genetic interaction networks. Ann Rev Genet. 2009; 43:601–625.

Domingo J, Baeza-Centurion P, Lehner B. The causes and consequences of genetic interactions (epistasis). Annu Rev of Genomics Hum Genet. 2019; 20:433–460.

Dowell RD, Ryan O, Jansen A, Cheung D, Agarwala S, Danford T, Bernstein DA, Rolfe PA, Heisler LE, Chin B, et al. Genotype to phenotype: a complex problem. Science. 2010; 328(5977):469–469.

Du D, Roguev A, Gordon DE, Chen M, Chen SH, Shales M, Shen JP, Ideker T, Mali P, Qi LS, et al. Genetic interaction mapping in mammalian cells using CRISPR interference. Nat Methods. 2017; 14(6):577.

Dykhuizen DE, Dean AM, Hartl DL. Metabolic flux and fitness. Genetics. 1987; 115(1):25–31.

Ewald J, Bartl M, Dandekar T, Kaleta C. Optimality principles reveal a complex interplay of intermediate toxicity and kinetic efficiency in the regulation of prokaryotic metabolism. PLoS Comput Biol. 2017; 13(2).

Fang G, Wang W, Paunic V, Heydari H, Costanzo M, Liu X, Liu X, VanderSluis B, Oately B, Steinbach M, et al. Discovering genetic interactions bridging pathways in genome-wide association studies. Nat Commun. 2019; 10(1):1–18.

Feist AM, Palsson BO. The biomass objective function. Current opinion in microbiology. 2010; 13(3):344–349.

Fiévet JB, Dillmann C, de Vienne D. Systemic properties of metabolic networks lead to an epistasis-based model for heterosis. Theoretical and Applied Genetics. 2010; 120(2):463.

Fisher RA. The Genetical Theory of Natural Selection. Oxford: The Clarendon Press; 1930.

Flynn KM, Cooper TF, Moore FB, Cooper VS. The environment affects epistatic interactions to alter the topology of an empirical fitness landscape. PLoS Genet. 2013; 9(4):e1003426.

Fragata I, Blanckaert A, Louro MAD, Liberles DA, Bank C. Evolution in the light of fitness landscape theory. Trends in ecology & evolution. 2019; 34(1):69–82.

Gertz J, Gerke JP, Cohen BA. Epistasis in a quantitative trait captured by a molecular model of transcription factor interactions. Theoretical population biology. 2010; 77(1):1–5.

Gibson G. Epistasis and pleiotropy as natural properties of transcriptional regulation. Theoretical population biology. 1996; 49(1):58–89.

Gjuvsland AB, Hayes BJ, Omholt SW, Carlborg Ö. Statistical epistasis is a generic feature of gene regulatory networks. Genetics. 2007; 175(1):411–420.

Gu C, Kim GB, Kim WJ, Kim HU, Lee SY. Current status and applications of genome-scale metabolic models. Genome biology. 2019; 20(1):121.

Hansen TF, Wagner GP. Modeling genetic architecture: a multilinear theory of gene interaction. Theoretical population biology. 2001; 59(1):61–86.

Harms MJ, Thornton JW. Historical contingency and its biophysical basis in glucocorticoid receptor evolution. Nature. 2014; 512(7513):203–207.

Hartl DL, Dykhuizen DE, Dean AM. Limits of adaptation: the evolution of selective neutrality. Genetics. 1985; 111(3):655–674.

He X, Qian W, Wang Z, Li Y, Zhang J. Prevalent positive epistasis in *Escherichia coli* and *Saccharomyces cerevisiae* metabolic networks. Nature genetics. 2010; 42(3):272–276.

Heigwer F, Scheeder C, Miersch T, Schmitt B, Blass C, Jamnani MVP, Boutros M. Time-resolved mapping of genetic interactions to model rewiring of signaling pathways. eLife. 2018; 7:e40174.

Horlbeck MA, Xu A, Wang M, Bennett NK, Park CY, Bogdanoff D, Adamson B, Chow ED, Kampmann M, Peterson TR, et al. Mapping the genetic landscape of human cells. Cell. 2018; 174(4):953–967.

Hu T, Sinnott-Armstrong NA, Kiralis JW, Andrew AS, Karagas MR, Moore JH. Characterizing genetic interactions in human disease association studies using statistical epistasis networks. BMC bioinformatics. 2011; 12(1):364.

Huang W, Richards S, Carbone MA, Zhu D, Anholt RR, Ayroles JF, Duncan L, Jordan KW, Lawrence F, Magwire MM, et al. Epistasis dominates the genetic architecture of Drosophila quantitative traits. Proc Natl Acad Sci USA. 2012; 109(39):15553–15559.

Husain K, Murugan A. Physical constraints on epistasis. Molecular Biology and Evolution. 2020; 37(10):2865–2874.

Hutchison CA, Chuang RY, Noskov VN, Assad-Garcia N, Deerinck TJ, Ellisman MH, Gill J, Kannan K, Karas BJ, Ma L, et al. Design and synthesis of a minimal bacterial genome. Science. 2016; 351(6280).

Jasnos L, Korona R. Epistatic buffering of fitness loss in yeast double deletion strains. Nat Genet. 2007; 39(4):550–554.

Johnson MS, Martsul A, Kryazhimskiy S, Desai MM. Higher-fitness yeast genotypes are less robust to deleterious mutations. Science. 2019; 366(6464):490–493.

Kacser H, Burns JA. The control of flux. Symp Soc Exp Biol. 1973; 32:65–104.

Kacser H, Burns JA. The molecular basis of dominance. Genetics. 1981; 97(3-4):639–666.

Keightley PD. Models of quantitative variation of flux in metabolic pathways. Genetics. 1989; 121(4):869–876.

Keightley PD. Metabolic models of selection response. Journal of Theoretical Biology. 1996; 182(3):311–316.

Kelley R, Ideker T. Systematic interpretation of genetic interactions using protein networks. Nat Biotech. 2005; 23(5):561–566.

Kemble H, Eisenhauer C, Couce A, Chapron A, Magnan M, Gautier G, Le Nagard H, Nghe P, Tenaillon O. Flux, toxicity, and expression costs generate complex genetic interactions in a metabolic pathway. Science Advances. 2020; 6(23):eabb2236.

Khan AI, Dinh DM, Schneider D, Lenski RE, Cooper TF. Negative epistasis between beneficial mutations in an evolving bacterial population. Science. 2011; 332(6034):1193–1196.

Kondrashov AS. Through sex, nature is telling us something important. Trends in Genetics. 2018; 34(5):352–361.

Kryazhimskiy S, Rice DP, Jerison ER, Desai MM. Global epistasis makes adaptation predictable despite sequence-level stochasticity. Science. 2014; 344(6191):1519–1522.

Kuzmin E, VanderSluis B, Wang W, Tan G, Deshpande R, Chen Y, Usaj M, Balint A, Usaj MM, Van Leeuwen J, Koch EN, Pons C, Dagilis AJ, Pryszlak M, Wang JZY, Hanchard J, Riggi M, Xu K, Heydari H, Luis BJS, et al. Systematic analysis of complex genetic interactions. Science. 2018; 360(6386):eaao1729.

Lehner B. Molecular mechanisms of epistasis within and between genes. Tr Genet. 2011; 27(8):323–331.

Lehner B, Crombie C, Tischler J, Fortunato A, Fraser AG. Systematic mapping of genetic interactions in *Caenorhabditis elegans* identifies common modifiers of diverse signaling pathways. Nat Genet. 2006; 38(8):896–903.

Li X, Lalić J, Baeza-Centurion P, Dhar R, Lehner B. Changes in gene expression predictably shift and switch genetic interactions. Nature communications. 2019; 10(1):1–15.

Lindsey HA, Gallie J, Taylor S, Kerr B. Evolutionary rescue from extinction is contingent on a lower rate of environmental change. Nature. 2013; 494(7438):463–467.

Liu X, Liu Z, Dziulko AK, Li F, Miller D, Morabito RD, Francois D, Levy SF. iSeq 2.0: A Modular and Interchangeable Toolkit for Interaction Screening in Yeast. Cell Systems. 2019; 8(4):338–344.

Lunzer M, Miller SP, Felsheim R, Dean AM. The biochemical architecture of an ancient adaptive landscape. Science. 2005; 310(5747):499–501.

Lyons DM, Zou Z, Xu H, Zhang J. Idiosyncratic epistasis creates universals in mutational effects and evolutionary trajectories. Nature Ecology & Evolution. 2020; 4(12):1685–1693.

Macía J, Solé RV, Elena SF. The causes of epistasis in genetic networks. Evolution. 2012; 66(2):586–596.

Mackay TF. Epistasis and quantitative traits: using model organisms to study gene–gene interactions. Nat Rev Genet. 2014; 15(1):22–33.

MacLean R. Predicting epistasis: an experimental test of metabolic control theory with bacterial transcription and translation. J Evol Biol. 2010; 23(3):488–493.

Malik-Sheriff RS, Glont M, Nguyen TVN, Tiwari K, Roberts MG, Xavier A, Vu MT, Men J, Maire M, Kananathan S, Fairbanks EL, Meyer JP, Arankalle C, Varusai TM, Knight-Schrijver V, Li L, Dueñas-Roca C, Dass G, Keating SM, Park YM, et al. BioModels—15 years of sharing computational models in life science. Nucleic Acids Research. 2019 11; https://doi.org/10.1093/nar/gkz1055, doi: 10.1093/nar/gkz1055, gkz1055.

Mani R, Onge RPS, Hartman JL, Giaever G, Roth FP. Defining genetic interaction. Proc Natl Acad Sci USA. 2008;
105(9):3461–3466.

Martin G. Fisher's geometrical model emerges as a property of complex integrated phenotypic networks.
Genetics. 2014; 197(1):237–255.

Martin G, Elena SF, Lenormand T. Distributions of epistasis in microbes fit predictions from a fitness landscape
model. Nature genetics. 2007; 39(4):555–560.

Michaut M, Baryshnikova A, Costanzo M, Myers CL, Andrews BJ, Boone C, Bader GD. Protein complexes are
central in the yeast genetic landscape. PLoS Comput Biol. 2011; 7(2).

Musso G, Costanzo M, Huangfu M, Smith AM, Paw J, San Luis BJ, Boone C, Giaever G, Nislow C, Emili A, et al. The
extensive and condition-dependent nature of epistasis among whole-genome duplicates in yeast. Genome
Res. 2008; 18(7):1092–1099.

Natarajan A, Srienc F. Dynamics of glucose uptake by single *Escherichia coli* cells. Metabolic engineering. 1999;
1(4):320–333.

Natarajan C, Inoguchi N, Weber RE, Fago A, Moriyama H, Storz JF. Epistasis among adaptive mutations in deer
mouse hemoglobin. Science. 2013; 340(6138):1324–1327.

New AM, Lehner B. Harmonious genetic combinations rewire regulatory networks and flip gene essentiality.
Nat Commun. 2019; 10(1):1–12.

Norman TM, Horlbeck MA, Replogle JM, Alex YG, Xu A, Jost M, Gilbert LA, Weissman JS. Exploring genetic
interaction manifolds constructed from rich single-cell phenotypes. Science. 2019; 365(6455):786–793.

Omholt SW, Plahte E, Øyehaug L, Xiang K. Gene regulatory networks generating the phenomena of additivity,
dominance and epistasis. Genetics. 2000; 155(2):969–980.

Orr HA. The genetic theory of adaptation: a brief history. Nature Reviews Genetics. 2005; 6(2):119–127.

Orth JD, Conrad TM, Na J, Lerman JA, Nam H, Feist AM, Palsson BØ. A comprehensive genome-scale reconstruc-
tion of *Escherichia coli* metabolism–2011. Mol Syst Biol. 2011; 7(1).

Orth JD, Thiele I, Palsson BØ. What is flux balance analysis? Nature biotechnology. 2010; 28(3):245–248.

Otwinowski J, McCandlish DM, Plotkin JB. Inferring the shape of global epistasis. Proceedings of the National
Academy of Sciences. 2018; 115(32):E7550–E7558.

Peccoud J, Vander Velden K, Podlich D, Winkler C, Arthur L, Cooper M. The selective values of alleles in a
molecular network model are context dependent. Genetics. 2004; 166(4):1715–1725.

Phillips PC. Epistasis—the essential role of gene interactions in the structure and evolution of genetic systems.
Nature Reviews Genetics. 2008; 9(11):855–867.

Poelwijk FJ, Krishna V, Ranganathan R. The context-dependence of mutations: a linkage of formalisms. PLoS
computational biology. 2016; 12(6).

Pumir A, Shraiman B. Epistasis in a model of molecular signal transduction. PLoS computational biology. 2011;
7(5).

Rao S, Van der Schaft A, Van Eunen K, Bakker BM, Jayawardhana B. A model reduction method for biochemical
reaction networks. BMC systems biology. 2014; 8(1):52.

Reddy G, Desai MM. Global epistasis emerges from a generic model of a complex trait. bioRxiv. 2020; doi:
https://doi.org/10.1101/2020.06.14.150946.

Roguev A, Bandyopadhyay S, Zofall M, Zhang K, Fischer T, Collins SR, Qu H, Shales M, Park HO, Hayles J, et al.
Conservation and rewiring of functional modules revealed by an epistasis map in fission yeast. Science. 2008;
322(5900):405–410.

Roguev A, Talbot D, Negri GL, Shales M, Cagney G, Bandyopadhyay S, Panning B, Krogan NJ. Quantitative
genetic-interaction mapping in mammalian cells. Nat Methods. 2013; 10(5):432.

Sailer ZR, Harms MJ. High-order epistasis shapes evolutionary trajectories. PLoS Comput Biol. 2017;
13(5):e1005541.

Sailer ZR, Harms MJ. Molecular ensembles make evolution unpredictable. Proceedings of the National Academy of Sciences. 2017; 114(45):11938–11943.

Sanjuán R, Nebot MR. A network model for the correlation between epistasis and genomic complexity. PLoS One. 2008; 3(7).

Sarkisyan KS, Bolotin DA, Meer MV, Usmanova DR, Mishin AS, Sharonov GV, Ivankov DN, Bozhanova NG, Baranov MS, Soylemez O, et al. Local fitness landscape of the green fluorescent protein. Nature. 2016; 533(7603):397–401.

Savageau MA. Biochemical systems analysis. A study of function and design in molecular biology. Reading, MA: Addison-Wesley Publishing Company, Inc.; 1976.

Schoustra S, Hwang S, Krug J, de Visser JAG. Diminishing-returns epistasis among random beneficial mutations in a multicellular fungus. Proceedings of the Royal Society B: Biological Sciences. 2016; 283(1837):20161376.

Segel LA. On the validity of the steady state assumption of enzyme kinetics. Bulletin of mathematical biology. 1988; 50(6):579–593.

Segre D, DeLuna A, Church GM, Kishony R. Modular epistasis in yeast metabolism. Nat Genet. 2005; 37(1):77–83.

Skwark MJ, Croucher NJ, Puranen S, Chewapreecha C, Pesonen M, Xu YY, Turner P, Harris SR, Beres SB, Musser JM, et al. Interacting networks of resistance, virulence and core machinery genes identified by genome-wide epistasis analysis. PLoS Genet. 2017; 13(2).

Snitkin ES, Segrè D. Epistatic interaction maps relative to multiple metabolic phenotypes. PLoS Genet. 2011; 7(2):e1001294.

Sohail M, Vakhrusheva OA, Sul JH, Pulit SL, Francioli LC, van den Berg LH, Veldink JH, de Bakker PI, Bazykin GA, Kondrashov AS, et al. Negative selection in humans and fruit flies involves synergistic epistasis. Science. 2017; 356(6337):539–542.

St Onge RP, Mani R, Oh J, Proctor M, Fung E, Davis RW, Nislow C, Roth FP, Giaever G. Systematic pathway analysis using high-resolution fitness profiling of combinatorial gene deletions. Nat Genet. 2007; 39(2):199–206.

Stock JB, Waygood EB, Meadow N, Postma P, Roseman S. Sugar transport by the bacterial phosphotransferase system. The glucose receptors of the *Salmonella typhimurium* phosphotransferase system. J Biol Chem. 1982; 257(23):14543–14552.

Szappanos B, Kovács K, Szamecz B, Honti F, Costanzo M, Baryshnikova A, Gelius-Dietrich G, Lercher MJ, Jelasity M, Myers CL, et al. An integrated approach to characterize genetic interaction networks in yeast metabolism. Nat Genet. 2011; 43(7):656–662.

Szathmary E. Do deleterious mutations act synergistically? Metabolic control theory provides a partial answer. Genetics. 1993; 133(1):127–132.

Taylor MB, Ehrenreich IM. Higher-order genetic interactions and their contribution to complex traits. Trends in genetics. 2015; 31(1):34–40.

Tenaillon O. The utility of Fisher's geometric model in evolutionary genetics. Annual review of ecology, evolution, and systematics. 2014; 45:179–201.

Tenaillon O, Rodríguez-Verdugo A, Gaut RL, McDonald P, Bennett AF, Long AD, Gaut BS. The molecular diversity of adaptive convergence. Science. 2012; 335(6067):457–461.

Tischler J, Lehner B, Fraser AG. Evolutionary plasticity of genetic interaction networks. Nat Genet. 2008; 40(4):390–391.

Tong AHY, Evangelista M, Parsons AB, Xu H, Bader GD, Pagé N, Robinson M, Raghibizadeh S, Hogue CW, Bussey H, et al. Systematic genetic analysis with ordered arrays of yeast deletion mutants. Science. 2001; 294(5550):2364–2368.

Typas A, Nichols RJ, Siegele DA, Shales M, Collins SR, Lim B, Braberg H, Yamamoto N, Takeuchi R, Wanner BL, et al. High-throughput, quantitative analyses of genetic interactions in *E. coli*. Nat Methods. 2008; 5(9):781.

Van Leeuwen J, Pons C, Mellor JC, Yamaguchi TN, Friesen H, Koschwanez J, Ušaj MM, Pechlaner M, Takar M, Ušaj M, et al. Exploring genetic suppression interactions on a global scale. Science. 2016; 354(6312):aag0839.

Van Opijnen T, Camilli A. Transposon insertion sequencing: a new tool for systems-level analysis of microorganisms. Nature Reviews Microbiology. 2013; 11(7):435–442.

Velenich A, Gore J. The strength of genetic interactions scales weakly with mutational effects. Genome biology. 2013; 14(7):R76.

Versfeld L. Remarks on star-mesh transformation of electrical networks. Electron Lett. 1970; 6(19):597–599.

Wagner GP. Two rules for the detection and quantification of epistasis and other interaction effects. In: *Epistasis* Springer; 2015.p. 145–157.

Wagner GP, Laubichler MD, Bagheri-Chaichian H. Genetic measurement theory of epistatic effects. In: *Mutation and Evolution* Springer; 1998.p. 569–580.

Wei WH, Hemani G, Haley CS. Detecting epistasis in human complex traits. Nature Reviews Genetics. 2014; 15(11):722–733.

Weinreich DM, Delaney NF, DePristo MA, Hartl DL. Darwinian evolution can follow only very few mutational paths to fitter proteins. science. 2006; 312(5770):111–114.

Weinreich DM, Watson RA, Chao L. Sign epistasis and genetic costraint on evolutionary trajectories. Evolution. 2005; 59(6):1165–1174.

Wiser MJ, Ribeck N, Lenski RE. Long-term dynamics of adaptation in asexual populations. Science. 2013; 342(6164):1364–1367.

Wright S. Physiological and evolutionary theories of dominance. The American Naturalist. 1934; 68(714):24–53.

Yu MK, Kramer M, Dutkowski J, Srivas R, Licon K, Kreisberg JF, Ng CT, Krogan N, Sharan R, Ideker T. Translation of genotype to phenotype by a hierarchy of cell subsystems. Cell Syst. 2016; 2(2):77–88.

Zuk O, Hechter E, Sunyaev SR, Lander ES. The mystery of missing heritability: Genetic interactions create phantom heritability. Proc Natl Acad Sci USA. 2012; 109(4):1193–1198.

 Appendix 1

Proof of *Equation 24*

First, the terms in *Equation 24* can be re-arranged as follows

$$
\begin{aligned}
x_{ij}^E &= x_{ij} + \frac{x_{ik_1} x_{k_1 j}}{D_{k_1}^{E\backslash\{k_1\}}} + \frac{x_{ik_2} x_{k_2 j}}{D_{k_2}^{E\backslash\{k_2\}}} + \cdots \\
&+ \frac{x_{ik_1} x_{k_1 k_2} x_{k_2 j}}{D_{k_1}^{E\backslash\{k_1\}} D_{k_2}^{E\backslash\{k_1,k_2\}}} + \cdots + \frac{x_{ik_2} x_{k_2 k_1} x_{k_1 j}}{D_{k_2}^{E\backslash\{k_2\}} D_{k_1}^{E\backslash\{k_1,k_2\}}} + \cdots \\
&= x_{ij} + \frac{x_{ik_1}}{D_{k_1}^{E\backslash\{k_1\}}} \underbrace{\left(x_{k_1 j} + \frac{x_{k_1 k_2} x_{k_2 j}}{D_{k_2}^{E\backslash\{k_1,k_2\}}} + \cdots \right)}_{= x_{k_1 j}^{E\backslash\{k_1\}}} \\
&+ \frac{x_{ik_2}}{D_{k_2}^{E\backslash\{k_2\}}} \underbrace{\left(x_{k_2 j} + \frac{x_{k_2 k_1} x_{k_1 j}}{D_{k_1}^{E\backslash\{k_1,k_2\}}} + \cdots \right)}_{= x_{k_2 j}^{E\backslash\{k_2\}}} + \cdots \\
&= x_{ij} + \sum_{\ell \in E} \frac{x_{i\ell} x_{\ell j}^{E\backslash\{\ell\}}}{D_\ell^{E\backslash\{\ell\}}}.
\end{aligned}
\tag{75}
$$

Next, I will demonstrate the validity of *Equation 75* by induction. It is clear that for $E = \{k\} \subset A_\mu$, *Equation 24* reduces to *Equation 12*. Now suppose that $n_E > 1$ and that there exists a metabolite $k \in E$, such that *Equation 75* holds for the subset $E' = E \setminus \{k\}$, i.e.,

$$
x_{ij}^{E'} = x_{ij} + \sum_{\ell \in E'} \frac{x_{i\ell} x_{\ell j}^{E'\backslash\{\ell\}}}{D_\ell^{E'\backslash\{\ell\}}}.
\tag{76}
$$

I will now show that then *Equation 75* also holds for E. To do so, I use the definition of x_{ij}^E (*Equation 19*) and apply *Equation 76* to expand terms $x_{ij}^{E'}$ and $x_{ik}^{E'}$ as follows.

$$
\begin{aligned}
x_{ij}^E &= x_{ij}^{E'} + \frac{x_{ik}^{E'} x_{kj}^{E'}}{D_k^{E'}} \\
&= x_{ij} + \sum_{\ell \in E'} \frac{x_{i\ell} x_{\ell j}^{E'\backslash\{\ell\}}}{D_\ell^{E'\backslash\{\ell\}}} + \frac{x_{ik} x_{kj}^{E'}}{D_k^{E'}} + \sum_{\ell \in E'} \frac{x_{i\ell} x_{\ell k}^{E'\backslash\{\ell\}} x_{kj}^{E'}}{D_\ell^{E'\backslash\{\ell\}} D_k^{E'}} \\
&= x_{ij} + \sum_{\ell \in E'} \frac{x_{i\ell}}{D_\ell^{E'\backslash\{\ell\}}} \left(x_{\ell j}^{E'\backslash\{\ell\}} + \frac{x_{\ell k}^{E'\backslash\{\ell\}} x_{kj}^{E'}}{D_k^{E'}} \right) + \frac{x_{ik} x_{kj}^{E'}}{D_k^{E'}}.
\end{aligned}
\tag{77}
$$

Recalling that $E = E' \setminus \{k\}$, I re-write *Equation 75* as

$$
x_{ij}^E = x_{ij} + \sum_{\ell \in E'} \frac{x_{i\ell} x_{\ell j}^{E\backslash\{\ell\}}}{D_\ell^{E\backslash\{\ell\}}} + \frac{x_{ik} x_{kj}^{E'}}{D_k^{E'}}.
\tag{78}
$$

Since the first and third terms of *Equation 77* and *Equation 78* are identical, to complete the proof, it is sufficient to show that for any $\ell \in E'$,

$$
\frac{1}{D_\ell^{E'\backslash\{\ell\}}} \left(x_{\ell j}^{E'\backslash\{\ell\}} + \frac{x_{\ell k}^{E'\backslash\{\ell\}} x_{kj}^{E'}}{D_k^{E'}} \right) \equiv \frac{1}{D_\ell^{E\backslash\{k,\ell\}}} \left(x_{\ell j}^{E\backslash\{k,\ell\}} + \frac{x_{\ell k}^{E\backslash\{k,\ell\}} x_{kj}^{E\backslash\{k\}}}{D_k^{E\backslash\{k\}}} \right) = \frac{x_{\ell j}^{E\backslash\{\ell\}}}{D_\ell^{E\backslash\{\ell\}}}.
\tag{79}
$$

To show that **Equation 79** holds, I first use **Equation 19** and **Equation 21** to express $x_{kj}^{E\setminus\{k\}}$ and $D_k^{E\setminus\{k\}}$ in terms of effective reaction rates after the elimination of the metabolite set $E \setminus \{k, \ell\}$,

$$
\begin{aligned}
x_{kj}^{E\setminus\{k\}} &= x_{kj}^{E\setminus\{k,\ell\}} + \frac{x_{k\ell}^{E\setminus\{k,\ell\}} \, x_{\ell j}^{E\setminus\{k,\ell\}}}{D_\ell^{E\setminus\{k,\ell\}}} \\[2mm]
D_k^{E\setminus\{k\}} &= D_k^{E\setminus\{k,\ell\}} - \frac{x_{k\ell}^{E\setminus\{k,\ell\}} \, x_{\ell k}^{E\setminus\{k,\ell\}}}{D_\ell^{E\setminus\{k,\ell\}}},
\end{aligned}
\tag{80}
$$

which imply that

$$
\begin{aligned}
&\frac{1}{D_\ell^{E\setminus\{k,\ell\}}}\left(x_{\ell j}^{E\setminus\{k,\ell\}} + \frac{x_{\ell k}^{E\setminus\{k,\ell\}} \, x_{kj}^{E\setminus\{k\}}}{D_k^{E\setminus\{k\}}}\right) = \frac{D_k^{E\setminus\{k,\ell\}}}{D_\ell^{E\setminus\{k,\ell\}} \, D_k^{E\setminus\{k\}}}\left(x_{\ell j}^{E\setminus\{k,\ell\}} + \frac{x_{\ell k}^{E\setminus\{k,\ell\}} \, x_{kj}^{E\setminus\{k,\ell\}}}{D_k^{E\setminus\{k,\ell\}}}\right) \\[2mm]
&= \frac{D_k^{E\setminus\{k,\ell\}} \, x_{\ell j}^{E\setminus\{\ell\}}}{D_\ell^{E\setminus\{k,\ell\}} \, D_k^{E\setminus\{k\}}}.
\end{aligned}
$$

Finally, it follows from **Equation 80** that $D_\ell^{E\setminus\{k,\ell\}} \, D_k^{E\setminus\{k\}} = D_k^{E\setminus\{k,\ell\}} \, D_\ell^{E\setminus\{\ell\}}$, which completes the proof of **Equation 79**. Thus, **Equation 75** and equivalently **Equation 24** hold for any metabolite subset $E \subseteq A_\mu$.

Appendix 2

Existence of a simple path that contains a given reaction

Lemma 1. *Let μ be a module with the reaction set R_μ. Then, for any reaction $a \in R_\mu$, there*
exists a simple path $p_{12}(a)$ within μ that connects the I/O metabolites and contains reaction a, i.e.,
$\mathcal{P}_{12}^\mu(a) \neq \emptyset$.

Proof. Reaction a is either a bypass, i/o, or internal reaction for module μ. If a is a bypass
reaction, then the statement is trivially true. If a is an i/o reaction, then, without loss of
generality, let $a = 1 \leftrightarrow j$. Since μ is a module, there exists a simple path $j \leftrightarrow j_1 \leftrightarrow \cdots \leftrightarrow 2$ that
connects the internal metabolite j to the I/O metabolite 2. Therefore, the path $1 \leftrightarrow j \leftrightarrow j_1 \leftrightarrow$
$\cdots \leftrightarrow 2$ connects the I/O metabolites and contains reaction a.

Suppose that $a = i \leftrightarrow j$ is an internal reaction. To prove the statement, it is sufficient to
show that there exists a pair of non-intersecting paths p'_{1i} and p'_{2i}, such that one of them
contains a and the other does not. Since μ is a module, there exists a pair of non-intersecting
paths p_{1i} and p_{2i} and a pair of non-intersecting paths p_{1j} and p_{2j} within module μ (I omitted
super-index μ to simplify notations). There are two mutually exclusive possibilities. (i)
Metabolite j is contained in either of the paths p_{1i} or p_{2i} and/or metabolite i is contained in
either of the paths p_{1j} or p_{2j}. (ii) Metabolite j is not contained in either of the paths p_{1i} or
p_{2i} and metabolite i is not contained in either of the paths p_{1j} or p_{2j}, i.e., $j \notin p_{ui}$ and $i \notin p_{uj}$,
$u = 1, 2$. It is trivial to construct the neccessary paths p'_{1i} and p'_{2i} in case (i).

2102
2104 **Appendix 2 Figure 1. Illustration for the proof of *Lemma 1*.**

Consider case (ii). If paths p_{2j} and p_{1i} do not intersect, then let $p'_{1i} = p_{1i}$ and

$$p'_{2i} = \underbrace{i \leftrightarrow}_{=a} \overbrace{j \leftrightarrow \cdots \leftrightarrow 2}^{=p_{2j}},$$

and the statement is true. Suppose paths p_{2j} and p_{1i} intersect. Then, among all metabolites
that belong to both p_{1i} and p_{2j}, let metabolite k be the one closest to j along the path p_{2j} (see
Figure). Then the segment p_{kj} of path p_{2j} and the path p_{1i} do not intersect. Let

$$p''_{1i} = \overbrace{1 \leftrightarrow \cdots \leftrightarrow k}^{\text{along } p_{1i}} \underbrace{\leftrightarrow \cdots \leftrightarrow j}_{\text{along } p_{2j}} \overbrace{\leftrightarrow i}^{=a},$$

If p''_{1i} and p_{2i} do not intersect, then the lemma is true. If p''_{1i} and p_{2i} do intersect, this intersetion
can only occur within the segment p_{kj} of path p''_{1i}, excluding metabolites k and j (see Figure).
This is because the remaining segment p_{1k} of path p''_{1i} is also a segment of p_{1i}, which, by
assumption, does not intersect p_{2i}. Suppose that among all metabolites that belong to both

the segment p_{kj} of path p''_{1i} and the path p_{2i} metabolite ℓ is the one closest to j along the path p''_{1i}. Then let

$$
\begin{aligned}
p'_{1i} &= p_{1i}, \\
p'_{2i} &= \underbrace{2 \leftrightarrow \cdots \leftrightarrow \overset{\text{along } p_{2i}}{\overbrace{\ell}} \leftrightarrow \cdots \leftrightarrow \overset{=a}{\overbrace{j}} \leftrightarrow i}_{\text{along } p_{2j}} .
\end{aligned}
$$

The path p'_{2i} does not intersect the path p_{1i} because its first segment $p_{2\ell}$ belongs to path p_{2i} and its second segment $p_{\ell j}$ belongs to the segment p_{kj} of path p''_{1i} (and, as mentioned above, segment p_{kj} does not intersect p_{1i}). Thus, the statement holds for case (ii) as well. $\qquad \square$

Appendix 3

An algorithm for discovering all strictly serial and strictly parallel gen-
erating topologies

Suppose that $(R, a, b) \in \mathcal{G}_{\mathcal{M}}^{\text{ser}}$, i.e., (R, a, b) is a strictly serial generating topology in the topologi-
cal class $\mathcal{M}$. Since $R \subseteq R_{\mathcal{M}}$ where $R_{\mathcal{M}}$ is the complete reaction set $R_{\mathcal{M}}$ for class $\mathcal{M}$, R can be
discovered by sequentially removing reactions from $R_{\mathcal{M}}$. The same logic holds for strictly
parallel generating topologies. The following algorithm implements this idea.

1. **Define function `generate_topology_list`.** This function takes a topology $(R, a, b) \in$
$\mathcal{R}_{\mathcal{M}}^{\text{sp}}$ as input and returns a new list of topologies L as output, which is produced as
follows. Initialize $L = \emptyset$. For every reaction $c_i \in R \setminus \{a, b\}$, construct the reaction subset
$R_i = R \setminus \{c_i\}$ and use **Definition 1** to test whether R_i corresponds to a valid module. If
R_i corresponds to a module, add (R_i, a, b) to list L; otherwiese, discard.[a] Return list L.

2. **Initialization.**

(a) Pick a topological class $\mathcal{M}$.
(b) Test whether $(R_{\mathcal{M}}, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{ser}}$. If so, $\mathcal{G}_{\mathcal{M}}^{\text{ser}} = \{(R_{\mathcal{M}}, a, b)\}$ and $\mathcal{G}_{\mathcal{M}}^{\text{par}} = \emptyset$. Return $\mathcal{G}_{\mathcal{M}}^{\text{ser}}$, $\mathcal{G}_{\mathcal{M}}^{\text{par}}$.
(c) Test whether $(R_{\mathcal{M}}, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{par}}$. If so, $\mathcal{G}_{\mathcal{M}}^{\text{par}} = \{(R_{\mathcal{M}}, a, b)\}$ and $\mathcal{G}_{\mathcal{M}}^{\text{ser}} = \emptyset$. Return $\mathcal{G}_{\mathcal{M}}^{\text{ser}}$,
$\mathcal{G}_{\mathcal{M}}^{\text{par}}$.
(d) Set $\mathcal{G}_{\mathcal{M}}^{\text{ser}} = \emptyset$, $\mathcal{G}_{\mathcal{M}}^{\text{par}} = \emptyset$. Use function `generate_topology_list` with $(R_{\mathcal{M}}, a, b)$ as input
and obtain the list of reaction sets L. Proceed to Step 3 with list L.

3. Take list $L = \big((R_1, a, b), (R_2, a, b), \dots, (R_k, a, b)\big)$ as input. Set $L' = \emptyset$. Proceed to Step 3a
with $i = 1$.

(a) Test whether (R_i, a, b) belongs to $\mathcal{R}_{\mathcal{M}}^{\text{par}}$, $\mathcal{R}_{\mathcal{M}}^{\text{ser}}$ or $\mathcal{R}_{\mathcal{M}}^{\text{sp}}$. Choose one of the alternatives
3b, 3c, or 3d.
(b) $(R_i, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{par}}$. If (R_i, a, b) is not in the set $\mathcal{G}_{\mathcal{M}}^{\text{par}}$ and if $(R_i \cup \{c\}, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{sp}}$ for all
$c \in R_{\mathcal{M}} \setminus R_i$, then add (R_i, a, b) to $\mathcal{G}_{\mathcal{M}}^{\text{par}}$. Proceed to Step 3e.
(c) $(R_i, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{ser}}$. If (R_i, a, b) is not in the set $\mathcal{G}_{\mathcal{M}}^{\text{ser}}$ and if $(R_i \cup \{c\}, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{sp}}$ for all
$c \in R_{\mathcal{M}} \setminus R_i$, then add (R_i, a, b) to $\mathcal{G}_{\mathcal{M}}^{\text{ser}}$. Proceed to Step 3e.
(d) $(R_i, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{sp}}$. Use function `generate_topology_list` with (R_i, a, b) as input and
obtain the list of reaction sets L_i. Replace L' with $L' \cup L_i$. Proceed to Step 3e.
(e) If $i = k$, proceed to Step 3f. Otherwise, proceed to Step 3a with $i + 1$.
(f) If $L' \neq \emptyset$, then proceed to Step 3 with L' as input. Otherwise, return $\mathcal{G}_{\mathcal{M}}^{\text{ser}}$, $\mathcal{G}_{\mathcal{M}}^{\text{par}}$.

---

[a] It can be proven that, as long as $(R, a, b) \in \mathcal{R}_{\mathcal{M}}^{\text{sp}}$, there exists at least one $c_i \in R$, such that R_i corresponds to a module, i.e., $L \neq \emptyset$.

2158 **Appendix 4**

2159 **Derivation of *Equation 72***

In this case, equations (43) simplify to

$$W_{12}(v) = w_{12} + \frac{w_{16}\, w_{52}\, v/K_{56}}{D_{56}(v)},$$

$$W_{13}(v) = w_{13} + \frac{w_{16}\, w_{63}\, D_5(v)}{D_{56}(v)},$$

$$W_{14}(v) = \frac{w_{16}\, w_{54}\, v/K_{56}}{D_{56}(v)},$$

$$W_{23}(v) = \frac{w_{25}\, w_{63}\, v}{D_{56}(v)},$$

$$W_{24}(v) = w_{24} + \frac{w_{25}\, w_{54}\, D_6(v)}{D_{56}(v)},$$

$$W_{34}(u,v) = u + \frac{w_{36}\, w_{54}\, v/K_{56}}{D_{56}(v)},$$

where

$$D_{56}(v) = D_5(v)\, D_6(v) - \frac{v^2}{K_{56}},$$

2160 $$D_5(v) = w_{52} + w_{54} + v,$$

2161 $$D_6(v) = w_{61} + w_{63} + v/K_{56}.$$

2162
2163 Notice that the only effective rate constant that depends on u is W_{34}, and $\frac{\partial W_{34}}{\partial u} = 1$. Thus, it is
2164 easy to differentiate y, given by *Equation 42*, with respect to u, if we isolate the term W_{34} in
2165 both the numerator and the denomintor,

2166
2167 $$y = W_{12} + \frac{A_{56,34}\, W_{34} + B_{56,34}}{D_{56,34}}, \tag{81}$$

2168
2169 where

2170 $$D_{56,34} = E_{56,34}\, W_{34} + F_{56,34},$$

2171
2172 $$A_{56,34} = W_{14}\, W_{42} + \frac{W_{13}\, W_{32}}{K_{34}} + W_{13}\, W_{42} + \frac{W_{14}\, W_{32}}{K_{34}} \left(W_{13} + W_{14}\right) \left(\frac{W_{32}}{K_{34}} + W_{42}\right),$$

2173 $$B_{56,34} = W_{14}\, W_{42} \left(W_{31} + W_{32}\right) + W_{13}\, W_{32} \left(W_{41} + W_{42}\right),$$

2174
2175 $$E_{56,34} = \frac{W_{31} + W_{32}}{K_{34}} + \left(W_{41} + W_{42}\right),$$

2176 $$F_{56,34} = \left(W_{31} + W_{32}\right) \left(W_{41} + W_{42}\right).$$

2177

It is also useful to obtain another expression for y, which is easier to differentiate with respect to v. To do that, we can first eliminate metabolites 3 and 4 to obtain effective reaction

rates

$$V_{12}(u) = w_{12} + \frac{w_{13} \, w_{42} \, u}{D_{34}(u)},$$

$$V_{15}(u) = \frac{w_{13} \, w_{45} \, u}{D_{34}(u)},$$

$$V_{16}(u) = w_{16} + \frac{w_{13} \, w_{36} \, D_4(u)}{D_{34}(u)},$$

$$V_{25}(u) = w_{25} + \frac{w_{24} \, w_{45} \, D_3(u)}{D_{34}(u)},$$

$$V_{26}(u) = \frac{w_{24} \, w_{36} \, u/K_{34}}{D_{34}(u)},$$

$$V_{56}(u, v) = v + \frac{w_{36} \, w_{54} \, u/K_{34}}{D_{34}(u)},$$

where

$$D_{34}(u) = D_3(u) \, D_4(u) - \frac{u^2}{K_{34}},$$

$$D_3(u) = w_{31} + u + w_{36},$$

$$D_4(u) = w_{42} + u/K_{34} + w_{45}.$$

The only effective activity that depends on v is V_{56}, and $\frac{\partial V_{56}}{\partial u} = 1$. Thus, isolating the term V_{56} (and recalling that $V_{65} = V_{56}/K_{56}$), we obtain the following expression for y, which is easy to differentiate with respect to v.

$$y = V_{12} + \frac{A_{34,56} \, V_{65} + B_{34,56}}{D_{34,56}}, \tag{82}$$

where

$$D_{34,56} = E_{34,56} \, V_{65} + F_{34,56},$$

$$A_{34,56} = \left(V_{16} + V_{15} \right) \left(\frac{V_{62}}{K_{65}} + V_{52} \right),$$

$$B_{34,56} = V_{15} \, V_{52} \left(V_{61} + V_{62} \right) + V_{16} \, V_{62} \left(V_{51} + V_{52} \right),$$

$$E_{34,56} = \frac{V_{61} + V_{62}}{K_{65}} + \left(V_{51} + V_{52} \right),$$

$$F_{34,56} = \left(V_{61} + V_{62} \right) \left(V_{51} + V_{52} \right).$$

Using symbolic computation it is possible to show that (see Mathematica notebook (Supplementary file 1) and Mathematica notebook pdf (Supplementary file 2), p. 2)

$$D_{56,34} \, D_{56} = D_{34,56} \, D_{34}. \tag{83}$$

Also notice that, any module with the reaction set $i, i, \emptyset, P_7$ is symmetric with respect to swapping metabolite labels 1 with 2, 3 with 5, and 4 with 6. It is easy to check that equations (81), (82) respect this symmetry.

Differentiating **Equation 81** with respect to u, after some algebra I obtain

$$\frac{\partial y}{\partial u} = \frac{\partial y}{\partial W_{34}} = \frac{1}{K_{31}} \left(\frac{W_{31} \, W_{42} - W_{32} \, W_{41}}{D_{56,34}} \right)^2. \tag{84}$$

Analogously, differentiating **Equation 82** with respect to v, I obtain

$$\frac{\partial y}{\partial v} = \frac{1}{K_{56}} \frac{\partial y}{\partial V_{65}} = \frac{1}{K_{51}} \left(\frac{V_{52} \, V_{61} - V_{51} \, V_{62}}{D_{34,56}} \right)^2. \tag{85}$$

Notice that **Equation 85** can also be obtained from **Equation 84** by symmetry with respect to the aforementioned metabolite relabelling.

Next, using symbolic computation (see Mathematica notebook (Supplementary file 1) and Mathematica notebook pdf (Supplementary file 2), p. 3), it is possible to show that

$$\frac{(W_{31}\,W_{42} - W_{32}\,W_{41})\,D_{56}}{D_{56,34}\,D_{56}} = \frac{A_u\,v + B_u}{E_u\,v + F_u}, \tag{86}$$

where all coefficients

$$A_u = \frac{w_{31}}{K_{56}}\,\psi + w_{42}\,\phi,$$

$$B_u = \psi\,\phi$$

$$E_u = u\left[\frac{w_{31}\,w_{52} + w_{31}\,w_{54} + w_{36}\,w_{52}}{K_{34}\,K_{56}} + (w_{42}\,w_{61} + w_{42}\,w_{63} + w_{45}\,w_{61})\right] + D_4(u)\,\phi + \frac{D_3(u)}{K_{56}}\,\psi$$

$$F_u = u\left(\frac{w_{52} + w_{54}}{K_{34}}\,\phi + (w_{61} + w_{63})\,\psi\right) + \phi\,\psi.$$

and

$$\phi = w_{31}\,w_{61} + w_{31}\,w_{63} + w_{36}\,w_{61},$$

$$\psi = w_{42}\,w_{52} + w_{42}\,w_{54} + w_{45}\,w_{52}$$

are independent of v and are non-negative. Similarly (see Mathematica notebook (Supplementary file 1) and Mathematica notebook pdf (Supplementary file 2), p. 4),

$$\frac{(V_{52}\,V_{61} - V_{51}\,V_{62})\,D_{34}}{D_{34,56}\,D_{34}} = \frac{A_v\,u + B_v}{E_v\,u + F_v}, \tag{87}$$

where

$$A_v = w_{61}\,\psi + \frac{w_{52}}{K_{34}}\,\phi,$$

$$B_v = \psi\,\phi,$$

$$E_v = v\left[(w_{42}\,w_{61} + w_{42}\,w_{63} + w_{45}\,w_{61}) + \frac{w_{31}\,w_{52} + w_{31}\,w_{54} + w_{36}\,w_{52}}{K_{34}\,K_{56}}\right] + \frac{D_5(v)}{K_{34}}\,\phi + D_6(v)\,\psi$$

$$F_v = v\left((w_{42} + w_{45})\,\phi + \frac{w_{31} + w_{36}}{K_{56}}\,\psi\right) + \phi\,\psi.$$

We can now obtain the second derivative $\frac{\partial^2 y}{\partial u\,\partial v}$, taking into account **Equation 86**. Alternatively, we can obtain $\frac{\partial^2 y}{\partial u\,\partial v}$ by differentiating $\frac{\partial y}{\partial v}$ with respect to u, taking into account **Equation 87**. The denominators in both expressions would be identical due to **Equation 83**. Therefore the expression for the second derivative must have the form given by **Equation 72**, i.e.,

$$\frac{\partial^2 y}{\partial u\,\partial v} = \frac{2\,\beta}{K_{31}}\,\frac{(A_u\,v + B_u)\,(A_v\,u + B_v)}{(E_u\,v + F_u)^3},$$

where β is independent of u and v. Thus, according to **Equation 84**, **Equation 86**,

$$\beta = \frac{A_u\,F_u - B_u\,E_u}{A_v\,u + B_v} = -(w_{36}\,\psi/K_{56} + w_{45}\,\phi),$$

which is verified in Mathematica notebook (Supplementary file 1) and Mathematica notebook pdf (Supplementary file 2), p. 4. Similarly, according to *Equation 85*, *Equation 87*,

$$\frac{\beta}{K_{31}} = \frac{1}{K_{51}} \frac{A_v F_v - B_v E_v}{A_u v + B_v} = -\frac{w_{63} \psi + w_{54}/K_{34} \phi}{K_{51}}$$

which is verified in Mathematica notebook (Supplementary file 1) and Mathematica notebook pdf (Supplementary file 2), p. 5.

2256 **Appendix 5**

Relationship to the Flux Balance Analysis

2257
Here, I discuss the relationship of the model presented in this paper to the flux balance
analysis (FBA), a widely used approach to modeling whole-cell metabolism (***Orth et al., 2010***).
FBA and my model are designed to address different questions. My model was designed to
explore how the flux depends on the kinetic parameters of a module. FBA was designed to
avoid this dependence. Nevertheless, the two models are conceptually and mathematically
related and in fact are in some sense equivalent, as discussed below. The most important
similarity is that both models rely on flux balance—the assumption that internal metabolites
are at steady state—as a key simplification.

The major difference is that my model assumes that all reaction kinetics are first order,
which allows for analytical tractability, while FBA is agnostic with respect to reaction kinetics
and takes into account only their stoichiometries and possibly some additional "capacity"
constraints on the reaction rates (***Orth et al., 2010***). This difference deserves some additional
discussion. In real cells, the concentrations of internal metabolites and the resulting fluxes
(which are functions of these concentrations) are determined by the stoichiometries of
reactions, the activities of enzymes and the concentrations of external nutrients. A general
approach to modelling such systems, adopted by the metabolic control analysis and related
theories, is to explicitly specify the dynamic equations for the metabolites, such as ***Equa-***
tion 10, and solve them to find steady-state concentrations of internal metabolites (see e.g.,
Savageau, 1976). The steady-state fluxes are then determined automatically. In my model,
the internal steady state exists and is unique for any module (see ***Corollary 1***). In contrast,
there are no dynamic equations within FBA. Instead, the steady-state fluxes are subject only
to the mass-balance equations and the capacity conditions, which typically form a severely
underdetermined system. Therefore, arriving at a unique flux distribution in FBA requires
additional assumptions. The typical approach is to first fix at least some nutrient uptake
rates and then maximize an objective function, such as the growth rate or biomass yield
(***Feist and Palsson, 2010***). In other words, FBA trades-off the ability to handle reactions with
arbitrary kinetics and unknown kinetic parameters against the necessity to impose auxiliary
conditions to find the right solution among many plausible ones.

As a consequence of these model design choices, mutations that add or remove reactions
can be naturally studied within the FBA framework, but there is no natural way to incorporate
mutations that perturb the kinetic parameters of reactions (***He et al., 2010***; ***Alzoubi et al.,***
2019). In contrast, mutations of small or large effect, including reaction additions and
deletions, can be in principle naturally studied within my model.

Due to differences in the assumptions, my model and FBA describe different sets of
biological systems. However, there are special cases which can be described by both FBA
and my model, and in these special cases the two models are mathematically equivalent.

From the perspective of FBA, modules with two I/O metabolites and first-order kinetics
described by my model are a special case. So, let us apply FBA to a metabolic module
$\mu = (A_\mu, \mathbf{x}_\mu)$ with two I/O metabolites 1 and 2. For the purposes of FBA, let the I/O metabolite
1 be the "external nutrient" and let the I/O metabolite 2 be the "biomass". Suppose that
metabolites in the set $A_{\mathrm{in}} \subset A_\mu$ are adjacent to the external nutrient and metabolites in
the set $A_{\mathrm{out}} \subset A_\mu$ are adjacent to the biomass. Then reactions $1 \leftrightarrow i$ for $i \in A_{\mathrm{in}}$, i.e., those
that convert the nutrient into intermediate metabolites, are the "uptake reactions". And the
reactions $i \leftrightarrow 2$ for $i \in A_{\mathrm{out}}$, i.e., those that convert intermediate metabolites into biomass, are
the "biomass reactions". Denote the rate of reaction consuming metabolite i and producing
metabolite j by v_{ij} and let $J_{\mathrm{in}} = \sum_{i \in A_{\mathrm{in}}} v_{1i}$ and $J_{\mathrm{out}} = \sum_{i \in A_{\mathrm{out}}} v_{i2}$ be the input and output fluxes,

respectively. To study epistasis within the FBA framework, we would need to obtain J_{out}.

The assumption of my model that all reaction kinetics are first-order translates into the FBA formalism as the fact that the elements of the stoichiometry matrix S take values $-1, +1$ or 0 and there are no capacity constraints on the rates of reactions. Then, the mass-balance equation $S\,v = 0$ leads to the equality $J_{\text{in}} = J_{\text{out}}$, so that the flux through the module does not depend on module's internal structure. Instead, it must be given as an auxiliary condition.

In my model, the steady-state flux J through the module depends in general on the internal structure of the module (through the effective rate constant y_μ) and is given by $J = y_\mu \left(S_1 - S_2/K_{12} \right)$ for any concentrations S_1 and S_2 of the external nutrient and the biomass. However, in the degenerate case where all uptake reactions are irreversible (i.e., $K_{1i} = \infty$ for all $i \in A_{\text{in}}$), $J = J_{\text{in}} = \sum_{i \in A_{\text{in}}} x_{1i} S_1$, such that it is independent of the internal structure of the module. In this sense, this special case of my model is equivalent to FBA.[a]

---

[a]However, my model and FBA are not equivalent in terms of the distribution of internal fluxes. My model still produces a unique steady sate and the corresponding flux distribution, whereas FBA does not specify a unique flux distribution in this case without additional auxiliary conditions.

Appendix 6

Connection between metabolism and growth

Here I describe a simple "bioreactor" model for connecting a hierarchical metabolic network
described in this paper with cellular growth. Suppose that module μ with I/O metabolites
1 and n is at the top level of the metabolic hierarchy and describes the whole cell. The
I/O metabolite 1 can be thought of as nutrients, and the I/O metabolite n can be thought
of as proteins, so that cellular metabolism converts nutrients into biomass. Denote the
concentrations of metabolites and proteins within cells by S_i and p. Suppose that cells have
a fixed volume v and the number of cells is N. Then, the absolute abundance of proteins
and metabolites across all N cells is $P = p\,v\,N$ and $A_i = S_i\,v\,N$, $i = 1, \ldots, n-1$.

The proteins produced by the cell are the enzymes that catalyze all the reactions inside
module μ. Let the relative expression level of the enzyme catalyzing reaction $i \leftrightarrow j$, $i < j$ be
ϕ_{ij} (with $\sum_{i=1}^{n-1} \sum_{j>i} \phi_{ij} = 1$), so that its concentration inside cells is $\phi_{ij}\,p$. If the specific forward
and reverse activities of this enzyme are a_{ij} and a_{ji} (so that a_{ij} obey the Haldane equalities
and $a_{ii} = 0$), then the total forward and reverse activities are $x_{ij} = a_{ij}\,\phi_{ij}\,p$ and $x_{ji} = a_{ji}\,\phi_{ij}\,p$.

Finally, I assume that all reactions converting internal metabolites into biomass are
irreversible, and I assume that the cell density in the bioreactor is small enough that the
nutrient concentration S_1 stays constant. In other words, I model early exponential growth.
2331 Then the dynamics of metabolite and protein abundances are governed by equations

2332

2333
$$\dot{A}_i \quad = \quad \sum_{i=1}^{n-1} x_{ji}\,A_j - D_i\,A_i, \quad i = 2, \ldots n-1. \tag{88}$$
2334

2335
2336
$$\dot{P} \quad = \quad \sum_{i=2}^{n-1} x_{in}\,A_i, \tag{89}$$
2337

2338 where $D_i = \sum_{i=1}^{n} x_{ij}$.

I assume that all cells are identical and at steady state, such that protein and metabolite
concetrations inside cells are constants and the production of proteins and metabolites
2339 manifests itself in the multiplication of cells. Then *Equation 88* and *Equation 89* become
2340

2341
$$\lambda\,S_i \quad = \quad \sum_{i=1}^{n-1} x_{ji}\,S_j - D_i\,S_i, \quad i = 2, \ldots n-1. \tag{90}$$
2342

2343
$$\dot{N} \quad = \quad \lambda\,N, \tag{91}$$
2344

where $\lambda = \sum_{i=2}^{n-1} x_{in} S_i$ is the steady-state flux into biomass which defines the exponential
growth rate of the system.

Finally, assuming that the rates of consumption and production of all metabolites are
much greater than their dilution by growth, I set $\lambda\,S_i \approx 0$. Then *Equation 90* defines the same
steady state for module μ as described in Network coarse-graining. Therefore, module μ can
be replaced with the effective activity x_μ between nutrients and biomass, yielding $\lambda = x_\mu$.

Additional files

- **Supplementary file 1.** Mathematica notebook "Case i,i,emptyset,P7.nb" for evaluating epistasis for the generating topology $i, i, \emptyset, P_7$.
- **Supplementary file 2.** PDF version of the Mathematica notebook "Case i,i,emptyset,P7.nb".

2355

Figure 4–Figure supplement 1. Distribution of the position of the fixed point $\bar{\varepsilon}$ in 1000 modules from the topological class $\mathcal{M}^{\text{io}}$ with random parameters. Notations are as in **Figure 4**.

2356

Figure 4–Figure supplement 2. Distribution of the position of the fixed point $\bar{\varepsilon}$ in 1000 modules from the topological class $\mathcal{M}^i$ with random parameters. Notations are as in **Figure 4**.

2357

Figure 4–Figure supplement 3. Fraction of sampled modules with different strictly serial generating topologies where the epistasis coefficient falls between γ **and** $1 - \gamma$**.** Dark gray bars show cases with epistasis substantially below 1 ($\gamma = 0.01$). The same data is shown in **Figure 4**B, averaged over 11 generating topologies. Light gray bars show additional cases where epistasis is clustered around 1 (see Materials and Methods for details). The effects of both mutations are indicated on each panel.

2358

Figure 5–Figure supplement 1. Detailed schematic of the kinetic model of glycolysis. Blue circles indicate metabolites (see **Table 5** for abbreviations). Orange rectangles indicate enzymes (see **Table 6** for abbreviations). Double-arrows indicate reversible reactions. Arrows with a fletch indicate irreversible reactions. Black circles indicate reactions with multiple substrates or products. Different shades of gray indicate the FULL, GPP, UGPP and LG models defined in **Table 4**.

2359

Figure 5–Figure supplement 2. Control coefficients for the output flux in the FULL module.
The first-order flux control coefficients (FCCs) of all 18 reactions with respect to the output flux in the unpperturbed FULL model are shown. 11 out of 18 reactions have positive FCCs, five (TA, TKa, TKb, R5PI, Ru5P) have zero FCCs and two (G6PDH and PEPCxyl) have negative FCCs. The 11 reactions with positive FCCs were retained for further analysis, the others were excluded. For these 11 reactions, the FCCs and epistasis coefficients with respect to fluxes through all modules (FULL, GPP, UGPP, LG) are shown in *Figure 5–Figure Supplement 3*.

2360

Figure 5–Figure supplement 3. Control and epistasis coefficients for the fluxes through multiple sub-modules within glycolysis. The top row shows the FCCs of 11 reactions whose FCCs in the FULL model are positive (see *Figure 5–Figure Supplement 2*). The matrix below shows the epistasis coefficients for each pair of these reactions. The topological relationship between reactions is indicated in the lower left corner of each panel ("P", strictly parallel; "S", strictly serial; "SP" serial-parallel). Points are colored orange (green) if the epistasis coefficient is less than zero (greater than one). Backgrounds of different shades of gray indicate the sub-modules, as in *Figure 5–Figure Supplement 1*.