Social structure defines spatial transmission of African swine fever in wild boar

Kim M. Pepin[1,*], Andrew Golnar[1], and Tomasz Podgórski[2,3]

1.   National Wildlife Research Center, USDA, APHIS, Wildlife Services, 4101 Laporte

Ave., Fort Collins, CO, 80526.

2.   Mammal Research Institute, Polish Academy of Sciences, Stoczek 1, 17-230 Białowieża,

Poland

3.   Department of Game Management and Wildlife Biology, Faculty of Forestry and Wood

Sciences, Czech University of Life Sciences, Kamýcká 129, 165 00 Praha 6, Czech Republic

*Corresponding author: Kim Pepin, National Wildlife Research Center, United States

Department of Agriculture, 4101 Laporte Ave., Fort Collins, CO, 80526;

kim.m.pepin@aphis.usda.gov

RH: Pepin et al. ˙Spatial spread of African swine fever in wild boar

Author E-mails:

Kim Pepin: kim.m.pepin@aphis.usda.gov

Tomasz Podgórski:t_podgorski@ibs.bialowieza.pl

Abstract

1. African swine fever virus (ASFv) is endemic in wild boar in Eastern Europe, challenging

elimination in domestic swine. Estimates of the distances between transmission events

are crucial for predicting rates of disease spread to guide allocation of surveillance and

control resources. Transmission distances are mainly defined by spatial and social

processes in hosts, but effects of these processes on spread are poorly understood, and

inferences often include only one process.

2. To understand effects of spatial and social processes on disease dynamics we developed

spatially-explicit transmission models with different assumptions about social and/or

spatial contact processes. We fit the models to ASFv surveillance data from Eastern

Poland from 2014-2015 and evaluated how inclusion of social structure affected

inference.

3. The model that accounted for social along with spatial processes provided better

inference of spatial spread and predicted that ~80% of transmission events are within the

same family group.

4. The models predicted dramatically different effective reproductive numbers, both in

magnitude and variation.

5. Specifying contact structure with spatial but not social processes can lead to very

different disease dynamics and inference of epidemiological parameters. Uncertainty in

these processes should be accounted for in predicting spatial spread in social species.

KEYWORDS: African swine fever, Effective reproduction number, Spatial transmission kernel,

Surveillance, Wild boar,

INTRODUCTION

Spatial transmission kernels (STKs) are probability distribution functions of transmission

distance between sequential cases of infection. They describe the variation and limits of spatial

disease spread per transmission event and inform how surveillance, containment, or mitigation

strategies should be deployed (Salje et al. 2016). For example, information on where cases may

arise can inform what spatial radius should be used for ring culling, ring vaccination, spatial

quarantine or intensive surveillance (Riley 2007). Without detailed genetic data or contact

tracing data to reconstruct transmission history, STKs are predominantly estimated indirectly by

fitting disease transmission models to available case data (Riley 2007, Salje et al. 2016, Pepin et

al. 2017) providing valuable insight for developing intervention strategies (Ferguson et al. 2001a,

b, Keeling et al. 2001, Woolhouse 2011). However, models often make simplifying assumptions

based on the available information that could have negative impacts on policy decisions if the

models are not robust to violation of these assumptions. Common assumptions in models for

estimating STKs include assuming a single introduction event and assuming observation of all

transmission events (Chipperfield et al. 2011). Methods that account for these processes are

important for providing more realistic predictions of spatial spread in systems where these

assumptions are violated.

Another common issue is that the potential scope of contact heterogeneities is often

simplified or lacking, such that uncertainty in model specification cannot be considered despite

its potential importance (Shea et al. 2014). In non-vector-borne disease systems, key drivers of

contact heterogeneities include social (Sah et al. 2018) and spatial processes (Riley 2007), yet

our understanding of the relative role of these processes in driving spatial disease dynamics is

weak in most systems (Emch et al. 2012, Dougherty et al. 2018) and analyses that consider

contact heterogeneities tend to focus on one or the other. Understanding the potential effects of

these different elements of contact heterogeneity on inference of disease dynamics will provide

insight on the appropriate scope of model uncertainty that should be considered in practical

applications.

African swine fever virus (ASFv), a virulent virus of swine, emerged in domestic swine

in Georgia in 2007 following a single introduction event from Africa (Chapman et al. 2011).

After its initial emergence, the virus spread quickly to Eastern Europe becoming endemic in wild

boar, which has challenged elimination. With no effective vaccine or treatment options, control

strategies are focused on reducing swine movement, decontamination, and culling of infected

animals (Authority et al. 2020). Effectiveness of these strategies depends on being able to rapidly

find new cases and target high-risk areas, thus models that can predict spatial spread are crucial

tools. However, it has become clear that modeling efforts need to include realistic ecological

details to improve predictions of spatial spread and our understanding of factors that drive it

(Arias et al. 2018). Wild boar have limited spatial movement and cluster into family groups

suggesting that both social and spatial processes likely need to be considered for estimating

STKs and thus spatial spread (Pepin et al. 2016, Podgórski et al. 2018, Pepin et al. 2020). Also

surveillance is passive with only a small proportion of cases likely being observed, and genetic

(Mazur-Panasiuk and Woźniakowski 2019) and other analyses (Pepin et al. 2020) suggest that

re-introductions are common. Thus estimates of STKs that account for these realities are likely

important for providing more accurate guidance for disease control policies.

In previous work we used ASFv surveillance data from wild boar in Poland during 2014-

2015 to estimate the frequency of carcass-based transmission and re-introduction in outbreak

dynamics while accounting for partially observed data (Pepin et al. 2020). Here, we extended the

modeling approach to three different assumptions about the form of spatial transmission

processes: 1) neighborhood (local transmission only), 2) exponential decay (distance distribution

that includes long-distance processes), and 3) distance distribution with social structure (Fig. 1).

We then predicted the STKs under each set of assumptions using the fitted models and evaluated

the effects of spatial and social processes on STKs and a key epidemiological parameter – the

effective reproduction number. Our results show striking differences in disease dynamics when

social structure modifies spatially-determined contact heterogeneities, showing that both

processes need to be considered for accurate predictions of spatial spread in socially-structured

species.

METHODS

Surveillance data and study site

The index case of ASFv in wild boar was detected in February 2014 in the north-eastern

part of the country (53°19'33"N, 23°45'31"E), less than 1 km from the border with Belarus.

Subsequent cases occurred close to the Belarusian border (Śmietanka et al. 2016, Woźniakowski

et al. 2016). By the end of 2015, 139 wild boar tested positive for ASFv in the area, with

maximum distance of 27.4km west of the border and a 100km range along the border. The

affected area is dominated by a mosaic of woodlands and agricultural land (crop fields, pastures,

meadows) with several large (several hundred square kilometers), continuous forests. On

average, forest covers 53% of the area. In 2014, average wild boar densities were estimated at

1.5 - 2.5 boar/km^2, locally ranging from 0.5-1 boar /km^2 to 3-5 boar/km^2 (Regional Directorate

of State Forests, Białystok, Poland). ASFv surveillance used a combination of active and passive

mechanisms, with samples being obtained through hunter recovery (active) or reports of road

kills or carcasses found on the landscape opportunistically (passive). A total of 4625 samples

were from hunters, while 271 were from road kill or non-road kill carcasses. Samples, collected

by veterinary services and hunters, were submitted to the National Reference Laboratory for

ASFv at the National Veterinary Research Institute in Puławy, Poland. Detailed description of

laboratory procedures and tests can be found in Woźniakowski et al. 2015 and Śmietanka et al.

2016. We used surveillance data from the area of 8 administrative districts where ASFv occurred

during 2014-2015 ('infected zone', 2224 samples tested) to fit the model and define spatio-

temporal intensity of sampling in our model.

Process model

We used a spatially-explicit, individual-based modeling framework fitted to the ASFv

surveillance data using Approximate Bayesian Computation (ABC) as in Pepin et al. 2020. We

evaluated three models that differed by social and spatial transmission process assumptions

(described below) in host populations that were structured by family groups and dispersal as

defined by field data (Fig. 1). We estimated transmission parameters and some other

epidemiological and demographic parameters as described below. With parameters from the

fitted models we then predicted cases over time, spatial spread over time, spatial transmission

kernels, effective reproductive numbers over time, and age- and sex-structure of infected

individuals. All analyses were implemented in Matlab (Version R2016b, The MathWorks, Inc.,

Natick, Massachusetts, United States). A full description of the individual-based model is given

in Pepin et al. (2020). Below is an overview of the approach with emphasis on differences from

our previous work.

We used a 5 x 5 km (25 km^2) gridded landscape to map spatial movement. The total

landscape size was 120 x 50 km (6000 km^2), similar to the 'infected' zone. Grid cells each had a

carrying capacity of 0.5 or 2 boars/km^2, which controlled heterogeneity in population density

across the landscape through density-dependent reproduction. In previous work we found that

this level of heterogeneous boar density fit the surveillance data better than homogenous

densities of 1, 2 or 4 wild boar / km^2 (Pepin et al. 2020).

Individual-boar attributes were monitored and updated at a daily time step. These

included age, unique group identification, X and Y coordinates of the home range centroid, grid-

cell ID; and status of life, reproduction, disease. Thus, the distribution of wild boar locations was

continuous but density was controlled at the grid cell level. The variable attributes changed based

on time, age, group size, grid-cell density, natal dispersal timing, and the disease transmission

process. Attributes that were fixed at birth included sex, dispersal distance, dispersal age, and age

at natural death. Thus, natal dispersal age and distance, and natural death, occurred at pre-set

ages and distance (for dispersal).

Individual-boar status was updated by the following order of processes: daily movement

(defined by the contact processes described below) and disease transmission, natural mortality

(occurring according to the pre-set age), natal dispersal (occurring according to the pre-set age),

dispersal due to other factors (i.e., family groups becoming too large, single females searching

for groups; occurring based on current family group size), surveillance sampling (permanent

removal of individuals from the landscape), conception (rates dependent on current grid cell

density), and new births (occurring with gestating females reach the end of their gestation

period). Fixed parameters included longevity (a data-based distribution), litter size (6), age at

reproductive maturity (180 days), minimum time between conception and farrowing (90 days),

gestation time (115 days), age of natal dispersal (~Poisson(13 months) truncated between 10-24

months), dispersal distance (~Weibull(2.5,0.5)), maximum size of family groups (10), incubation

period for ASFv (~Poisson(4 days) truncated at 1), infectious period for ASFv (~Poisson(5 days)

truncated at 1), and disease-induced mortality (100%). There were also fixed seasonal trends that

varied monthly for conception probability and carcass persistence that were based on data

(Bieber and Ruf 2005, Selva et al. 2005, Ježek et al. 2011, Rosell et al. 2012). Rationale and

sources for these parameters are derived from ecological studies of wild boar and are described

in Pepin et al. 2020.

Based on the attributes and processes described above, epidemiological states for

individual boar included: susceptible, exposed, infectious, infectious carcass, uninfectious

carcass, and removed from the landscape. Mortality only occurred from the disease (leading to

an infectious carcess) or reaching the age of longevity (leading to an uninfectious carcass). The

hunting process of alive individuals caused direct removal from the landscape (no carcass). Our

model also included multiple spatio-temporal scales of spatial processes because the dispersal

process (~Weibull(2.5,0.5) allowed for longer-distance movements and occurred less frequently

relative to the contact process that occurred daily and mostly at shorter distances.

We compared three different forms of contact structure: 1) neighborhood (local

transmission only), 2) exponential decay (distance distribution that includes long-distance

processes), and 3) distance distribution with social structure (Fig. 1). Forms 1 and 2 are common

ways of considering contact in space at population-level scales (Keeling and Rohani 2008),

whereas 3 incorporates heterogeneity due to social groups. For 1, infectious individuals could

transmit to all susceptible individuals within a fixed radius with equal probability. The radius of

the local neighborhood was constant across individuals and time – thus similar to a queen's

neighbor effect (Eq. 1). For 2, infectious individuals could transmit to all susceptible individuals

on the landscape, but the probability of transmission decayed with distance (Eq. 2). Model

structure 3 was the same as 2, except that transmission rates varied due to both group

membership and space - individuals in the same family group had higher transmission rates with

each other relative to those among family groups (Eq. 3). In general the daily force of infection

(λ) for each contact structure was defined as follows:

$$187 \quad \lambda = \sum_{k=1}^{K}\sum_{j=1}^{J} I_{k,d} \begin{cases} S_j\beta_d, & x_{k,j} < \zeta \\ 0, & otherwise \end{cases} + \sum_{k=1}^{K}\sum_{j=1}^{J} I_{k,c} \begin{cases} S_j\beta_c, & x_{k,j} < \zeta \\ 0, & otherwise \end{cases} + \beta_{0,\{j\}} \qquad (1)$$

$$189 \quad \lambda = \sum_{k=1}^{K}\sum_{j=1}^{J} I_{k,d} S_j\beta_d e^{-\alpha x_{k,j}} + \sum_{k=1}^{K}\sum_{j=1}^{J} I_{k,c} S_j\beta_c e^{-\alpha x_{k,j}} + \beta_{0,\{j\}} \qquad (2)$$

$$188 \quad \lambda = \sum_{k=1}^{K}\sum_{j=1}^{J} I_{k,d}(S_j\beta_d\, e^{-\alpha x_{k,j}} + S_{j,w}\beta_{w,d}) + \sum_{k=1}^{K}\sum_{j=1}^{J} I_{k,c}(S_j\beta_c e^{-\alpha x_{k,j}} + S_{j,w}\beta_{w,c}) + \beta_{0,\{Sj\}} \qquad (3)$$
190

Where ζ is a fixed local neighborhood (Eq. 1) delimiting the contact radius, $x_{i,j}$ is the distance

between infectious individual k (I_k) and susceptible individual j (S_j), α is the rate at which

transmission decays with distance (Eq. 2 and 3), d denotes alive individuals or direct

transmission, c denotes infectious carcasses or carcass-based transmission, β is the transmission

rate that is specific to the transmission mechanism (d or c), $\beta_{0,\{j\}}$ is the baseline rate at which re-

introduction occurs to susceptible individuals near the Eastern border ($\{Sj\}$), and family group

structure (w denotes contacts within the same family group, absence of w denotes among family

groups; Eq. 3).

Observation model

Because surveillance sampling was small compared to the full population it was

important to calibrate the process model with an observation model. Thus, we sampled the true

disease dynamics according to the surveillance process that was used in Poland, i.e., alive

individuals were available to be harvested by hunters, and carcasses were available to be found

for carcass sampling. As negative samples could not be georeferenced to the grid cell level (they

were only available at the district level), we were not able to account for the spatial distribution

of sampling accurately. However, there were strong temporal trends in the number of samples

collected thus we used those trends to describe sampling heterogeneity. First, we calculated the

relative number of boar sampled by hunters and carcass-sampling from the data (number

sampled on day t/maximum ever sampled separately for each method) to produce seasonal trends

in the proportion of the population sampled. Then we multiplied the seasonal trend data for each

method by the scaling factors (ρ_h and ρ_c) to determine the daily proportion of boar that would be

sampled (detection probability) by hunter harvesting or dead carcasses across the landscape at

random. We assumed that boar < 6 months of age would not be hunted (typically not targeted by

hunters) and that boar < 3 months of age would not be sampled by the dead carcass method

(because they are unlikely to be found).

Model fitting and evaluation

Unknown parameters were estimated based on Approximate Bayesian Computation

(ABC) with rejection sampling as described in Pepin et al. 2020. For all models, estimated

parameters included: frequency of introduction at the eastern border ($\beta_{0,\{j\}}$), β_d, β_c, scaling

parameters on seasonal trends of hunted hosts (ρ_h) and carcass sampling (ρ_c), a scaling parameter

on seasonal trends in the length of carcass persistence on the landscape (π), and a scaling

parameter on seasonal patterns of host birth probabilities (θ). In addition, we estimated spatial

parameters that describe three different contact structures: 1) ξ (nearest-neighbor), 2) α (the

decay of contact probability with distance, and 3) $\beta_{w,d}$ and $\beta_{w,c}$ (direct and carcass-based

transmission rates for within-group contacts). Prior distributions are listed in Table S1 (with

restrictions: $\beta_d > \beta_c$, $\beta_{w,d} > \beta_d$, $\beta_{w,c} > \beta_c$) and were informed by movement and contact data

(Podgórski et al. 2013, Pepin et al. 2016, Kay et al. 2017, Podgórski et al. 2018).

To sample across parameter space efficiently we used a Latin hypercube algorithm to

generate 979,592 parameter sets and then ran the model twice on each parameter set (for a total

of 1,959,184 iterations; or 2 chains of 979,592). β_d, β_c, and ρ_c were sampled on a log scale. A

two-tiered approach was applied for evaluating parameter sets to improve efficiency. Simulations

were terminated early if they were unrealistic, specifically: 1) when landscape-wide host density

< 20% of the initial density, 2) > 150 new cases per day; 3) no new cases sampled for 6 months,

or 4) > 300 total cases (more than double the actual number). We then only considered parameter

sets for which the simulation reached the end of the two-year time frame. The posterior

distributions consisted of all unique parameter sets (considering both chains) that were within the

absolute distance of three metrics: the sum of absolute differences between observed and

simulated surveillance data for monthly cases from live and dead animals (considered

separately), and the maximum monthly Euclidian distance of cases from the eastern border.

Distance metric tolerance values were 48 for monthly cases from carcasses, 24 for monthly cases

from hunter-harvest samples, and 120 for maximum distance from the border. This allowed

average error rates of 2 (carcass) and 1 (hunter harvest) cases, and 5 km from the border per

month on average. These error rates represent levels of uncertainty that we expected from the

data sources in our system, sensitivity analyses revealed that less stringent error rates would

affect the posterior distribution estimates (data not shown), and more stringent error rates would

require restrictively large computational resources unless prior distributions are more informed.

Average distance metrics for parameter sets from the posterior distribution were used to

evaluate goodness of fit along with R^2 values (squared correlation of observed and predicted case

and spatial distance trajectories, Table S1) and mean absolute error (MAE, Fig. 2). For each

fitted model we predicted outbreak dynamics using 1000 random samples from the posterior

distribution. The average of the 1000 predictions was used to calculate R^2 and MAE. We also

tested the ability of our models to forecast ASF dynamics by using the parameters estimated

from fits to the 2014-2015 data to predict the first 7 months of 2016 (Jan.-Jul.). We predicted

underlying spatial transmission kernels, effective reproductive number over time, and age-sex

structure of cases by simulating from the fitted models.

RESULTS

Parameter inference and model fit

The model with both social and spatial processes (Model 3) qualitatively captured spatial

spread better than Models 1 (Local neighborhood only) and 2 (Distance distribution only), and

Model 1 largely overestimated cases during the largest peak (Fig. 2). Also, the posterior

distribution of transmission probabilities were much lower and more realistic for Models 2 and 3

relative to Model 1 (see β_d and β_c in Table S1). The inferred STK for each model revealed two

distinct peaks symbolic of within- and between-group transmission (Fig. 3), but predicted

different amounts of within group transmission when within and between-group transmission

probabilities were allowed to vary (Model 3). Model 3 predicted the highest proportion of within

group transmission (0.8), followed by model 2 (0.6) and model 1 (0.3) (Fig. 3). For both Models

2 and 3, between-group transmission peaked between 0.5 km and 1 km, with a peak proportion

of transmission events reaching 0.05 and 0.02 for Model 2 and Model 3, respectively (Fig. 3).

For Model 1, between-group transmission events plateaued between 1-1.5 km at a frequency of

0.2 before dropping rapidly to a frequency of 0 around 1.5km (Fig. 3). The realized STKs for

Models 2 and 3 had long tails that indicated a low frequency of long-distance pathogen dispersal

(Fig. 3).

Impacts of model structure on epidemiological processes

The specification of spatial and social transmission processes in the model structure

resulted in dramatically different inference of R_e. Model 1 predicted much higher R_e over time,

followed by Model 3, and then Model 2 (Fig. 4). However, predictions from Model 2 suggest R_e

is relatively homogenous over time, while Models 1 and 3 predicted much more variability, with

R_e values reaching above a value of 4 on multiple occasions, and above a value of 8 at least once

(Fig. 4). Model 3 predicted higher R_e during annual birth pulses (Fig. 4). Models 1 and 2

predicted lower contributions of carcass-based transmission in overall R_e whereas Model 3

predicted more similar levels of each transmission mechanism (with carcass-based transmission

being slightly lower on average). All models predicted that the infected class is predominantly

composed by juveniles (<6 months of age; Fig. S1-S3), reflecting the age-structure in the

population. However, Models 1 and 2 predicted a slight male-bias in infected individuals while

Model 3 predicted a slight female bias (Fig. S1-S3).

DISCUSSION

Understanding how infectious diseases spread in space and time is key for developing

effective surveillance and intervention strategies (Keeling et al. 2001). Statistical inference of

wildlife disease systems is often challenged by partially observed data, multiple pathogen

introductions, and a limited understanding of host contact processes, all of which create

uncertainty in our ability to characterize patterns of infectious disease spread. Here we

demonstrate an approach for estimating patterns of ASFv spread that accounts for these

complexities while evaluating how different assumptions about contact structure influence the

inference of spatial disease dynamics. Results show that both spatial and social sources of

contact heterogeneity are important for capturing the spatial dynamics of ASFv in wild boar of

Eastern Poland, and that not accounting for both processes can lead to very different inferences

of a key epidemiological quantity, R_e.

Our results highlight the importance of considering both spatial and social processes in

estimating STKs. There are numerous approaches for modeling contact heterogeneities due to

spatial (Riley 2007, Keeling and Rohani 2008, White et al. 2018b, a) and social processes (Sah et

al. 2018). Network models are a useful approach for accounting for contact heterogeneity but

these are often focused only on heterogeneities due to social structure (Sah et al. 2018), while

neglecting spatial processes. Including both of these processes in a single network makes it

difficult to disentangle the role of each process, which can be important for designing optimal

control strategies. On the flip side, the field of movement ecology has developed new strategies

for accounting for heterogeneities due to animal movement (Dougherty et al. 2018), but these

methods remain underdeveloped for application in disease ecology, and also have not been used

to disentangle spatial and social processes. In order to provide practical inference for control of

important animal diseases it will be important to develop methods that can account for

heterogeneities due to both social and spatial processes, in a manner that the role of each process

can be inferred. This will allow control to be targeted to the appropriate process while improving

predictions of spatial spread for optimizing risk-based surveillance and control.

Our models suggested that between 30-80% of transmission events were within the same

family group and almost all transmission events were within 1.5 km, with some rare events at

longer distances. STK estimates can be used to establish control and surveillance zones. For

example, depopulation efforts could be intensified within 1.5 km where most transmission is

occurring, with surveillance intensified out to further distances (i.e., the tail of the STK) to

capture the rare long-distance spreading events. A useful approach could be to employ an

adaptive radius that focuses intervention efforts within 99% (or more – this should be validated

with modeling) of the STK, but adapts surveillance based on real-time surveillance. However,

the precise recommendations will depend on how soon a detection is made relative to where the

infection front is currently. With ASFv travelling at 1-2 km per month (Podgórski et al. 2018),

the radii for high-intensity culling and surveillance would need to be increased by 1-2 km for

each month that detection has lagged behind the infection front, highlighting the importance of

accurate predictions of spatial spread. A longer lag time for detection will also amplify

challenges that arise from long-distance jumps highlighting that this process is especially

important to understand.

Long-distance jumps were observed on multiple occasions in Poland after 2015 (Mazur-

Panasiuk and Woźniakowski 2019) and are thought to be due to human-mediated activities,

(Authority et al. 2017, Chenais et al. 2019). Thus, developing estimates of STKs that account for

mechanisms of long-distance dispersal will be important for appropriately targeting disease

control going forward. Our approach allowed for some long-distance events but we assumed a

monotonic functional form for contact distances, such that we did not infer the effects of spatial

contact processes occurring on multiple spatial scales. In order to infer spatial spread with later

surveillance data (i.e., 2016-present when longer-distance events occurred multiple times), it will

be important to incorporate other spatial mechanisms in the inference of the STK, perhaps using

covariate data that can inform these long-distance processes. Such an approach would provide

refined recommendations for surveillance and control targets at longer distances.

Our estimates of R_e (Model 1 (1.8-3.2), Model 2 (1.0-1.3) and Model 3 (1.1-2.0)) were

similar to an estimate of R_0 of ASFv in wild boar (1.13-3.77) in Russia (Iglesias et al. 2016).

However, although our estimates were similar in magnitude to Iglesias et al. (2016) on average,

they revealed substantial temporal variation in R_e that could impact policy decisions. For

example, all models estimated higher R_e during seasonal birth pulses indicating that at this time

of year we may expect to see higher rates of spatial spread. Additionally, estimates of the R_e over

time suggested a different role for carcass-based transmission in driving ASF transmission, with

Models 1 and 2 suggesting that carcass-based transmission is lower than direct transmission

while Model 3 suggested that the two types of transmission occurred at similar frequencies.

In addition to differences in R_e, the different contact structures predict differences in the role

of sex in transmission over time. Model 3 predicted a slight female bias because this model

predicted that 80% of transmission events occurred within family groups, which are female-

biased. In contract, Models 1 and 2 predicted a slight male bias because they predicted more

between-group transmission and with a 50:50 sex ratio there are more independent males relative

to family groups. Although it is known that males will travel longer distances than females

{Podgorski, 2014 #49}, especially during mating season to seek out females, we did not account

for this temporal heterogeneity in dispersal. Considering these types of movement

heterogeneities in future work could be important for improving our understanding of which sex

might present a higher risk of ASFv transmission and persistence.

Considered separately, spatial and social processes can have similar impacts on disease

dynamics. For example, social aggregation and spatial structuring can both reduce epidemic

potential by fragmented populations or by restricting the spread of pathogens (Keeling 1999, Sah

et al. 2018). However, our results highlight that spatial and social processes can also have quite

different impacts on epidemiological quantities, especially estimates of STKs, R_e, the frequency

of different transmission mechanisms, and potential risk factors such as sex. Being able to

appropriately infer the role of these quantities is crucial for optimizing disease control strategies.

When these contact heterogeneities are inappropriately accounted for it can bias inference (e.g.,

(Ames et al. 2011) and potentially misguide policy decisions (Shea et al. 2014). Moving forward

the field of disease ecology needs to develop mainstream methods that account for multiple

source of contact heterogeneities in a manner that their relative roles can be inferred so that

uncertainties in contact processes can be appropriately evaluated and incorporated in predictions

of spatial spread (Shea et al. 2014) and control can be targeted to the most important risk factors.

Acknowledgements

KMP was supported by the United States Department of Agriculture, Animal and Plant Health

Inspection Service's National Feral Swine Damage Management Program. AG was supported by

the United States Department of Agriculture, Animal and Plant Health Inspection Service's

APHIS Science Fellowship. TP was supported by the National Science Centre, Poland (grant

number 2014/15/B/NZ9/01933). We thank M. Łyjak, A. Kowalczyk, K. Śmietanka, and G.

Woźniakowski from Department of Swine Diseases, National Veterinary Research Institute in

Pulawy, Poland, for surveillance data. N. Selva provided valuable information on carcass

persistence time.

Figures

Fig 1. Schematic of methods and contact structures.

Fig. 2. Model fits to the observed surveillance data (red). Lines are the mean predictions from

1000 simulations with each fitted model (see legend in B for color code). Shading around the

lines areas are 95% prediction intervals of the means. Shading on the right side of plots indicates

the time frame for out-of-sample predictions.

407 **Fig. 3.** Realized spatial transmission kernels for each model (see legend). The X-axis is the

408 distance between home range centroids of infectious and susceptible individuals for which

409 transmission occurred. Y-axis is the proportion of all transmission events. Lines are the means of

410 100 simulations using random samples from the posterior distributions of the fitted models.

Fig. 5. Effective reproduction number (R_e) over time for each model. Effective reproduction

number at a given time point was calculated as the average number of transmissions made

throughout the infectious period for individuals that initially became infectious on day *t* (where *t*

is a day on the X-axis). Dark lines are the means of 100 simulations using random samples from

the posterior distributions of the fitted models; shading indicates 95% prediction intervals of the

means. Overall means with 95% prediction intervals for each model and each transmission

mechanism (black: direct, red: carcass-based) were: A) 1.9 [1.4, 2.3], 0.6 [0.4, 0.9], B) 0.9 [0.8

1.0], 0.2 [0.2, 0.3], C) 0.9 [0.7, 1.1], 0.6 [0.4, 0.9].

References

Ames, G. M., D. B. George, C. P. Hampson, A. R. Kanarek, C. D. McBee, D. R. Lockwood, J. D. Achter,

and C. T. Webb. 2011. Using network properties to predict disease dynamics on human contact

networks. Proceedings of the Royal Society B-Biological Sciences **278**:3544-3550.

Arias, M., C. Jurado, C. Gallardo, J. Fernández-Pinero, and J. Sánchez-Vizcaíno. 2018. Gaps in African

swine fever: analysis and priorities. Transboundary and emerging diseases **65**:235-247.

Authority, E. F. S., K. Depner, C. Gortazar, V. Guberti, M. Masiulis, S. More, E. Oļševskis, H. Thulke,

A. Viltrop, and G. Woźniakowski. 2017. Epidemiological analyses of African swine fever in the

Baltic States and Poland: (Update September 2016–September 2017). EFSA Journal **15**:e05068.

Authority, E. F. S., A. Miteva, A. Papanikolaou, A. Gogin, A. Boklund, A. Bøtner, A. Linden, A. Viltrop,

C. G. Schmidt, and C. Ivanciu. 2020. Epidemiological analyses of African swine fever in the

European Union (November 2018 to October 2019). EFSA Journal **18**:e05996.

Bansal, S., B. T. Grenfell, and L. A. Meyers. 2007. When individual behaviour matters: homogeneous

and network models in epidemiology. Journal of the Royal Society Interface **4**:879-891.

Bieber, C., and T. Ruf. 2005. Population dynamics in wild boar Sus scrofa: ecology, elasticity of growth

rate and implications for the management of pulsed resource consumers. Journal of applied

ecology **42**:1203-1213.

Chapman, D. A., A. C. Darby, M. Da Silva, C. Upton, A. D. Radford, and L. K. Dixon. 2011. Genomic

analysis of highly virulent Georgia 2007/1 isolate of African swine fever virus. Emerging

infectious diseases **17**:599.

Chenais, E., K. Depner, V. Guberti, K. Dietze, A. Viltrop, and K. Ståhl. 2019. Epidemiological

considerations on African swine fever in Europe 2014–2018. Porcine health management **5**:1-6.

Chipperfield, J. D., E. P. Holland, C. Dytham, C. D. Thomas, and T. Hovestadt. 2011. On the

approximation of continuous dispersal kernels in discrete-space models. Methods in Ecology and

Evolution **2**:668-681.

Dougherty, E. R., D. P. Seidel, C. J. Carlson, O. Spiegel, and W. M. Getz. 2018. Going through the

449   motions: incorporating movement analyses into disease research. Ecology letters **21**:588-604.

450   Emch, M., E. D. Root, S. Giebultowicz, M. Ali, C. Perez-Heydrich, and M. Yunus. 2012. Integration of

451   spatial and social network analysis in disease transmission studies. Annals of the Association of

452   American Geographers **102**:1004-1015.

453   Ferguson, N. M., C. A. Donnelly, and R. M. Anderson. 2001a. The foot-and-mouth epidemic in Great

454   Britain: pattern of spread and impact of interventions. Science **292**:1155-1160.

455   Ferguson, N. M., C. A. Donnelly, and R. M. Anderson. 2001b. Transmission intensity and impact of

456   control policies on the foot and mouth epidemic in Great Britain. Nature **413**:542-548.

457   Iglesias, I., M. Muñoz, F. Montes, A. Perez, A. Gogin, D. Kolbasov, and A. De la Torre. 2016.

458   Reproductive ratio for the local spread of African swine fever in wild boars in the Russian

459   Federation. Transboundary and emerging diseases **63**:e237-e245.

460   Ježek, M., K. Štípek, T. Kušta, J. Červený, and J. Vícha. 2011. Reproductive and morphometric

461   characteristics of wild boar (Sus scrofa) in the Czech Republic. Journal of forest science **57**:285-

462   292.

463   Kay, S. L., J. W. Fischer, A. J. Monaghan, J. C. Beasley, R. Boughton, T. A. Campbell, S. M. Cooper, S.

464   S. Ditchkoff, S. B. Hartley, and J. C. Kilgo. 2017. Quantifying drivers of wild pig movement

465   across multiple spatial and temporal scales. Movement Ecology **5**:14.

466   Keeling, M. J. 1999. The effects of local spatial structure on epidemiological invasions. Proceedings of

467   the Royal Society of London. Series B: Biological Sciences **266**:859-867.

468   Keeling, M. J., and P. Rohani. 2008. Modeling infectious diseases in humans and animals. Princeton

469   University Press, NJ, USA.

470   Keeling, M. J., M. E. Woolhouse, D. J. Shaw, L. Matthews, M. Chase-Topping, D. T. Haydon, S. J.

471   Cornell, J. Kappey, J. Wilesmith, and B. T. Grenfell. 2001. Dynamics of the 2001 UK foot and

472   mouth epidemic: stochastic dispersal in a heterogeneous landscape. Science **294**:813-817.

Mazur-Panasiuk, N., and G. Woźniakowski. 2019. The unique genetic variation within the O174L gene of

Polish strains of African swine fever virus facilitates tracking virus origin. Archives of virology

164:1667-1672.

Nurmoja, I., K. Mõtus, M. Kristian, T. Niine, K. Schulz, K. Depner, and A. Viltrop. 2018.

Epidemiological analysis of the 2015–2017 African swine fever outbreaks in Estonia. Preventive

veterinary medicine.

Pepin, K. M., A. J. Davis, J. Beasley, R. Boughton, T. Campbell, S. M. Cooper, W. Gaston, S. Hartley, J.

C. Kilgo, and S. M. Wisely. 2016. Contact heterogeneities in feral swine: implications for disease

management and future research. Ecosphere **7**:e01230.

Pepin, K. M., A. J. Davis, D. G. Streicker, J. W. Fischer, K. C. VerCauteren, and A. T. Gilbert. 2017.

Predicting spatial spread of rabies in skunk populations using surveillance data reported by the

public. PLoS neglected tropical diseases **11**:e0005822.

Pepin, K. M., A. Golnar, Z. Abdo, and T. Podgorski. 2020. Ecological drivers of African swine fever

virus persistence in wild boar populations: insight for control. Ecol Evolv. **10**:2846–2859.

Podgórski, T., M. Scandura, and B. Jedrzejewska. 2014. Next of kin next door - philopatry and

socio-genetic population structure in wild boar. The Journal of Zoology **294**:190-197.

Podgórski, T., M. Apollonio, and O. Keuling. 2018. Contact rates in wild boar populations: Implications

for disease transmission. The Journal of Wildlife Management **82**:1210-1218.

Podgórski, T., G. Baś, B. Jędrzejewska, L. Sönnichsen, S. Śnieżko, W. Jędrzejewski, and H. Okarma.

2013. Spatiotemporal behavioral plasticity of wild boar (Sus scrofa) under contrasting conditions

of human pressure: primeval forest and metropolitan area. Journal of Mammalogy **94**:109-119.

Riley, S. 2007. Large-scale spatial-transmission models of infectious disease. Science **316**:1298-1301.

Rosell, C., F. Navas, and S. Romero. 2012. Reproduction of wild boar in a cropland and coastal wetland

area: implications for management. Animal Biodiversity and Conservation **35**:209-217.

Sah, P., J. Mann, and S. Bansal. 2018. Disease implications of animal social network structure: a

synthesis across social systems. Journal of Animal Ecology **87**:546-558.

Salje, H., D. A. Cummings, and J. Lessler. 2016. Estimating infectious disease transmission distances

using the overall distribution of cases. Epidemics **17**:10-18.

Scherer, C., V. Radchuk, M. Franz, H. H. Thulke, M. Lange, V. Grimm, and S. Kramer–Schadt. 2020.

Moving infections: individual movement decisions drive disease persistence in spatially

structured landscapes. Oikos **129**:651-667.

Selva, N., B. Jędrzejewska, W. Jędrzejewski, and A. Wajrak. 2005. Factors affecting carcass use by a

guild of scavengers in European temperate woodland. Canadian Journal of Zoology **83**:1590-

1601.

Shea, K., M. J. Tildesley, M. C. Runge, C. J. Fonnesbeck, and M. J. Ferrari. 2014. Adaptive management

and the value of information: learning via intervention in epidemiology. PLoS biology

12:e1001970.

Śmietanka, K., G. Woźniakowski, E. Kozak, K. Niemczuk, M. Frączyk, Ł. Bocian, A. Kowalczyk, and Z.

Pejsak. 2016. African swine fever epidemic, Poland, 2014–2015. Emerging infectious diseases

22:1201.

White, L. A., J. D. Forester, and M. E. Craft. 2018a. Disease outbreak thresholds emerge from

interactions between movement behavior, landscape structure, and epidemiology. Proceedings of

the National Academy of Sciences **115**:7374-7379.

White, L. A., J. D. Forester, and M. E. Craft. 2018b. Dynamic, spatial models of parasite transmission in

wildlife: Their structure, applications and remaining challenges. Journal of Animal Ecology

87:559-580.

Woolhouse, M. 2011. How to make predictions about future infectious disease risks. Philosophical

Transactions of the Royal Society B: Biological Sciences **366**:2045-2054.

Woźniakowski, G., E. Kozak, A. Kowalczyk, M. Łyjak, M. Pomorska-Mól, K. Niemczuk, and Z. Pejsak.

2016. Current status of African swine fever virus in a population of wild boar in eastern Poland

(2014-2015). Archives of virology **161**:189-195.

Supplemental Material

Social structure defines spatial transmission of African swine fever in wild boar

Kim M. Pepin[1,*], Andrew Golnar[1], and Tomasz Podgórski[2,3]

4.   National Wildlife Research Center, USDA, APHIS, Wildlife Services, 4101 Laporte Ave., Fort Collins, CO, 80526.
5.   Mammal Research Institute, Polish Academy of Sciences, Stoczek 1, 17-230 Białowieża, Poland
6.   Department of Game Management and Wildlife Biology, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, Kamýcká 129, 165 00 Praha 6, Czech Republic

*Corresponding author: Kim Pepin, National Wildlife Research Center, United States Department of Agriculture, 4101 Laporte Ave., Fort Collins, CO, 80526;
kim.m.pepin@aphis.usda.gov

Contents

Table S1	Contact structure:	Neighborhood	Exp. Decay (ED)	ED & social structure
	R^2			
Monthly cases	in sample	0.57 ± 0.0059	0.51 ± 0.0054	0.55 ± 0.0060
(Median of the R^2's for 1000 individual time series ± 95% confidence interval)	all	0.48 ± 0.0054	0.43 ± 0.0046	0.47 ± 0.0049
Monthly distance from border	in sample	0.28 ± 0.018	0.29 ± 0.017	0.31 ± 0.018
(Median of the R^2's for 1000 individual time series ± 95% confidence interval)	all	0.19 ± 0.016	0.14 ± 0.011	0.28 ± 0.011
Monthly cases	in sample	0.65	0.57	0.63
(R^2 of median values from 1000 points at each month)	all	0.55	0.48	0.53
Monthly distance from border	in sample	0.53	0.49	0.53
(R^2 of median values from 1000 points at each month)	all	0.49	0.38	0.55
	Distance metrics[a]			
Median absolute error in monthly live cases ± 95% confidence interval	in sample	20 ± 0.8	20 ± 0.9	20 ± 0.5
Median absolute error in monthly carcass cases ± 95% confidence interval	in sample	44 ± 1.4	43 ± 2.8	44 ± 0.8
Median absolute error in monthly distance from border ± 95% confidence interval	in sample	109.5 ± 5.0	111.7 ± 5.7	106.6 ± 3.0
Number of values in posterior distribution		16 / 1,959,184 = 0.00082%	7 / 1,959,184 = 0.00036%	53 / 1,959,184 = 0.0027%
	Uniform Priors	95% credible intervals of posterior distributions		
Probability of direct transmission given proximity (β_d)	$[0.0001,1]$[b]	[0.0046, 1.0]	[0.039, 0.15]	[0.0003, 0.18]
Probability of carcass-based transmission given proximity (β_c)	$[0.0001,1]$[b]	[0.0002, 0.53]	[0.0038, 0.029]	[0.0001, 0.029]
Annual frequency of spillover from neighboring country ($\beta_{0,\{\varphi\}}$)	$[0,60]$[c]	[13, 59]	[9, 60]	[6, 59]
Detection probability in hunted boar samples (ρ_h)	$[0.0005,0.1]$[b]	[0.0006, 0.012]	[0.0007, 0.021]	[0.0007, 0.017]
Detection probability in carcass samples (ρ_c)	$[0.0005,0.8]$[b]	[0.012, 0.080]	[0.017, 0.045]	[0.014, 0.30]
Scaling parameter on seasonal trends in carcass persistence (π)	$[0.1,1.5]$	[0.58, 1.47]	[0.77, 1.41]	[0.17, 1.47]
Scaling parameter on seasonal trends in conception probability (θ)	$[0.5,6]$	[0.93, 5.73]	[1.31, 5.87]	[0.67, 5.88]
Constant contact radius (ζ)	$[0.5,5]$	[1.09, 3.35]	NA	NA
Rate parameter for decay of contact probability with distance (α)	$[0.1,2.5]$	NA	[1.62, 2.49]	[0.05, 2.49]
Probability of direct transmission given contact is in the same group ($\beta_{w,d}$)	$[0.01,1]$	NA	NA	[0.077, 0.98]
Probability of carcass-based transmission given contact is in the same group ($\beta_{w,c}$)	$[0.001,1]$	NA	NA	[0.14, 0.97]

[a]Median distance metrics ± 95% confidence intervals for 1000 simulations from the posterior distribution.

[b]These prior distributions were sampled on a natural log scale.

[c]0 indicates one introduction ever whereas values > 0 indicate the number of introductions / year.

Figure S1. The demographic dynamics of African swine fever virus based on Model 1 output (Neighborhood). Top plots show the frequency of infection in wild boar of different age classes (left) and sexes (right). Each line is a prediction from a separate sample of the posterior distribution of the fitted exponential decay & social structure model (100 trajectories in total). Bottom plots show the corresponding age class and sex distributions of infection over all time.

**Figure S2. The demographic dynamics of African swine fever virus based on Model 2
output (exponential decay).** Top plots show the frequency of infection in wild boar of different
age classes (left) and sexes (right). Each line is a prediction from a separate sample of the
posterior distribution of the fitted exponential decay & social structure model (100 trajectories in
total). Bottom plots show the corresponding age class and sex distributions of infection over all
time.

**Figure S3. The demographic dynamics of African swine fever virus based on Model 3
output (social and exponential).** Top plots show the frequency of infection in wild boar of
different age classes (left) and sexes (right). Each line is a prediction from a separate sample of
the posterior distribution of the fitted exponential decay & social structure model (100
trajectories in total). Bottom plots show the corresponding age class and sex distributions of
infection over all time.