TITLE

DNA Binding Induces a *cis* to *trans* Switch in Cre

Recombinase to Enable Intasome Assembly

AUTHORS

Aparna Unnikrishnan[1], Carlos D. Amero[2], Deepak Kumar Yadav[1], Kye Stachowski[1], Devante

Potter[1] and Mark P. Foster[1*]

[1]Department of Chemistry and Biochemistry, The Ohio State University, Columbus, Ohio,

USA 43210. [2]Laboratorio de Bioquímica y Resonancia Magnética Nuclear, Centro de

Investigaciones Químicas, Instituto de Investigación en Ciencias Básicas y Aplicadas,

Universidad Autónoma del Estado de Morelos, Avenida Universidad 1001, Colonia Chamilpa,

Cuernavaca, Morelos, 62209, Mexico.

*Corresponding Author: Mark P. Foster (email: foster.281@osu.edu; 614-292-1377).

Department of Chemistry and Biochemistry, Columbus, OH 43210.

ABSTRACT

Mechanistic understanding of DNA recombination in the Cre-*loxP* system has largely been guided by crystallographic structures of tetrameric synaptic complexes. Those studies have suggested a role for protein conformational dynamics that has not been well characterized at the atomic level. We used solution NMR to discover the link between intrinsic flexibility and enzyme function in Cre recombinase. Remarkably, in the absence of DNA the C-terminal helix αN, implicated in assembly of synaptic complexes and regulation of DNA cleavage activity via *trans* protein-protein interactions, is found to adopt an apparent auto-inhibitory *cis* conformation. Binding to *loxP* DNA dislodges the C-terminus from this *cis* conformation, thereby enabling the *trans* protein-protein interactions required for assembly of recombinogenic Cre intasomes. These findings necessitate a re-examination of the mechanisms by which this widely utilized gene-editing tool selects target sites, avoids spurious DNA cleavage activity, and controls DNA recombination efficiency.

INTRODUCTION

Site specific DNA recombinases represent an attractive option for genome engineering – the insertion or exchange of genes into precise locations in chromosomes[1–4]. The tyrosine recombinase family of phage-derived enzymes (e.g., λ-integrase, Cre and Flp recombinases), which evolved to facilitate viral infection, gene transposition and bacterial pathogenesis, have proven useful for applications that include DNA subcloning without restriction enzymes, and conditional expression of target genes[5–7]. These proteins bind specifically to pairs of inverted short palindromic DNA sequences (*recombinase binding elements*; RBEs) and mediate recombination by assembly of tetrameric *intasomes* (Fig. 1a, b) which perform concerted DNA strand cleavage, exchange and ligation reactions[5,8,9]. Compared to other genome engineering approaches[10,11], site specific DNA recombinases have the advantage that they allow a high degree of specificity, don't require involvement of additional host-encoded factors, and are capable of generating cleanly integrated double-stranded DNA products[3,12–14]. These advantages motivate efforts to understand and manipulate the mechanisms of this family of enzymes.

Cre (Causes Recombination) is the best studied member of the tyrosine recombinase family, yet fundamental gaps exist in our understanding of its mechanism of DNA site selection, intasome assembly, and allosteric control of DNA cleavage. The overall mechanism of Cre-mediated DNA recombination has been mapped out via many biochemical and crystallographic studies[5,15–19]. The enzyme catalyzes recombination between a pair of homologous 34 bp *loxP* sites (Fig. 1c) via a highly orchestrated series of events involving recognition and binding of a pair of Cre molecules to its semi-palindromic recognition site containing two RBEs (*loxP* half-sites), "synapsis" of an antiparallel pair of Cre dimers bound to *loxP* to form a stable tetrameric structure, coordinated cleavage of two opposing DNA

strands to produce an intermediate with two of the Cre protomers covalently attached to a 3'-

phosphate via a tyrosine linkage, followed by strand transfer and re-ligation to form a four-

way DNA Holliday junction intermediate[5]. Two-fold asymmetry in the synaptic structures (Fig.

1a) is implicated in regulation of DNA strand cleavage activity, and thereby influence the order

of strand cleavage and direction by which the Holliday junction is resolved[17,20,21].

Although much is known about the overall mechanism of the reactions catalyzed by

Cre and related site-specific recombinases, the nature of the conformational changes in the

protein and protein-DNA complexes that facilitate the various steps in the pathway remain

poorly understood[16,17,19,22–25]. Cre binds its target sequences by forming a C-shaped clamp

with a C-terminal catalytic domain (Cre^{Cat}) possessing the namesake tyrosine residue on one

side of the DNA, and on the other an N-terminal domain (Cre^{NTD}), connected by an extended

8 amino acid linker (Fig. 1). A series of sequential inter-protomer protein-protein interactions

across the tetrameric synapse distinguish the structures of the active and inactive protomers.

Within the catalytic domain, the tyrosine residue that serves as the nucleophile to catalyze

phosphoryl transfer and formation of the covalent intermediate is located on the penultimate

helix αM, while the αM-N loop and C-terminal helix αN from each protomer makes a *trans*

contact (in a "clockwise" manner as viewed in Fig. 1a) to the neighboring protomer in the

tetrameric complex. The β2-3 loop of each protomer abuts its own helix αM, and in the

tetramer packs proximally to helix αM of the neighboring protomer, in the opposite direction

compared to helix αN. Structural asymmetry in tetrameric structures is localized to these two

regions of the protein (Fig. 1a, b). The asymmetry in the C-terminal and inter-protomer

interfaces observed in the synaptic complexes of Cre and other tyrosine recombinase family

members suggest that the intrinsic dynamic behavior of the enzymes is important in

controlling its function, both in mediating tetramer assembly, and for regulating protomer

activity[26–32]. Despite the importance of understanding the basis for conformational differences

in Cre, structural insights have been largely limited to crystal structures of isosteric tetrameric

synaptic complexes[5,16,17,19].

We have used solution nuclear magnetic resonance (NMR) spectroscopy to explore

the link between the intrinsic dynamic behavior of Cre and its function. First, NMR spectra of

full-length Cre and of the isolated catalytic domain (Cre^{Cat}) support the premise that the N-

and C-terminal domains of Cre are structurally uncoupled. Nuclear spin relaxation

experiments reveal flexibility in regions of Cre^{Cat} that are associated with both protein-protein

and protein-DNA interactions. Unexpectedly, we found that the region of the protein

comprising C-terminal helix αN is not highly dynamic, contrary to the expectation from crystal

structures that it would be extended in solution[5,16,17,19]. Instead, NMR chemical shift

perturbations and paramagnetic relaxation enhancement (PRE) experiments show the C-

terminus of unbound Cre to be located in the DNA binding active site, in an apparent auto-

inhibitory conformation. Upon binding to *loxP* DNA, the data show that the C-terminus is

displaced, after which it is able to participate in fledgling intermolecular protein-protein

interactions with other Cre molecules. These findings shed light on the previous paradoxical

reports of *trans* interactions regulating Cre activity[33], and represent a transformative advance

in our understanding of the role of protein dynamics in regulating conservative site-specific

DNA recombination.

RESULTS

NMR spin relaxation reveals unexpected rigidity in the Cre C-terminal helix αN

To explore the role of protein dynamics in enabling the interconversion of Cre reaction

intermediates in solution, we expressed and purified full-length Cre (38.5 kDa, residues 1-

343) and a C-terminal fragment encoding the interdomain linker along with the C-terminal

domain (23.8 kDa, residues 127-343). This domain, herein termed Cre^{Cat} (Fig. 1b, d) includes

all seven active site residues (R173, E176, K201, H289, R292, W315 and Y324), the Cre-

DNA binding specificity determinants (R259 and E262) as well as the C-terminal region

implicated in *loxP* DNA binding cooperativity and synapsis (Fig. 1a)[17,33–37]. Both full-length

Cre and Cre^{Cat} are predominantly monomeric in the absence of DNA, as indicated by their

elution times in size exclusion chromatography (Supplementary Fig. 1a). Comparison of 2D

^{1}H-^{15}N TROSY-HSQC correlation spectra of Cre^{Cat} and full-length Cre showed that the well-

dispersed signals overlay well (Supplementary Fig. 1b). This indicates that Cre^{Cat} folds

independently in solution and that the N-terminal domain of Cre does not interact with the

catalytic domain in solution, thereby justifying structural and dynamics studies of the isolated

domain.

For backbone resonance assignments of Cre^{Cat} we expressed, purified the [U-^{15}N] and

[U-^{15}N,^{13}C] labeled protein and recorded HSQC- and TROSY-based double- and triple-

resonance NMR spectra. Homogeneity and integrity of protein constructs were verified by

SDS-PAGE and mass spectrometry (data not shown). We obtained backbone resonance

assignments for 195 of 214 non-proline amides (> 91%) (Fig. 1d, Supplementary Fig. 2 and

3). Unassigned amide resonances include the two N-terminal residues (A127 and G128), and

a few residues in the loop regions of Cre^{Cat}, due to line broadening from intermediate

conformational exchange.

To probe the fast timescale backbone dynamics in Cre[Cat], we measured ^{15}N NMR (R_1,

R_2) and {^1H}-^{15}N heteronuclear NOE (HetNOE) relaxation data (Fig. 2). The data show that

Cre[Cat] backbone amides largely exhibits uniform relaxation rates, with an overall rotational

correlation time (τ_C) of 13.2 ± 0.9 ns, computed from the trimmed-mean R_2/R_1 ratios[38].

Distinctly lower R_2/R_1 ratios and decreased {^1H}-^{15}N hetNOE values indicating fast (ps-ns)

internal motions, were observed for the linker residues through A134 preceding αF, the β2-3

loop, αJ-K loop, αM-N loop and a few residues after the C-terminal helix αN. The β2-3 loop

forms inter-protomer contacts in Cre-DNA synaptic tetramers and adopts distinct

conformations between the catalytically "active" and "inactive" forms of Cre (Fig. 1a, b)[5]. The

data show the αJ-K loop to be flexible in free Cre[Cat] though it does not differ structurally

between the two Cre isomers, consistent with a role in DNA binding, not in regulating

recombination[5]. Surprisingly, although residues bracketing the C-terminal helix αN exhibit

evidence of fast internal motions, the relaxation rates for the helix αN (A334- L339) are close

to the average values, indicating that it tumbles with the catalytic core of the protein. This

suggests that instead of being extended in solution as might be expected from the *trans*

conformations observed in tetrameric crystal structures, αN might adopt a distinct

conformation in the free protein.

Truncation of C-terminal helix αN results in widespread CSPs in the core of Cre[Cat]

In tetrameric DNA-bound crystal structures of Cre, the C-terminal helix αN of each protomer

packs in *trans* in a cyclic (non-reciprocal) manner into a surface cavity on a neighboring

protomer composed of residues from αF, αG, αH, β2-3 loop, αI, αK and αL regions (Fig. 1a).

Since Cre is monomeric in the absence of DNA, the Van Duyne group had previously used

crosslinking experiments to test the hypothesis that in free Cre the C-terminal region folds

back to dock over its own *trans* cavity in *cis*; those experiments failed to detect intramolecular

crosslinks[39]. Nevertheless, the ^{15}N relaxation data (Fig. 2) would be consistent with a *cis*

docking model in which rotational diffusion of αN is coincident with overall tumbling of the

protein.

To test whether the C-terminal region of Cre^{Cat} might be docked in *cis*, instead of being

extended for *trans* docking, we examined the effect of deleting the C-terminal residues on the

NMR spectra. We constructed a C-terminal deletion lacking thirteen C-terminal residues

including helix αN: Cre^{Cat}ΔC (residues 127- 330), obtained backbone resonance assignments

from triple resonance NMR data, and compared the ^{1}H-^{15}N TROSY-HSQC spectra of WT

Cre^{Cat} and Cre^{Cat}ΔC (Fig. 3a, Supplementary Fig. 4). Large chemical shift perturbations

(CSPs) were observed for residues in the interdomain linker region (Q133- A136), helices

αH, αK, αM and the αJ-K loop (Fig. 3a). Each of the regions in the protein core that show

large CSPs is in excess of 20 Å from the center of helix αN in the synaptic structures. These

findings are consistent with a *cis* docking arrangement of the C-terminus, and with the

restricted internal motions inferred from ^{15}N relaxation studies.

Remarkably, when the CSPs for Cre^{Cat}ΔC are mapped to the crystal structure of the

protein (Fig. 3b), the CSPs do not map to the synaptic *trans* docking surface of Cre. The

region of the interdomain linker that shows large CSPs (Q133- A136) is adjacent to the DNA

binding surface and was found to be largely rigid on the ps-ns timescale (Fig. 2). Helix αH

contains active site residues R173 and E176, αJ-K loop and helix αK contain the active site

residues H289 and R292, while helix αM contains the active site residue Y324.

These CSPs show that deletion of the C-terminal region alters the environment of

residues surrounding the active site of Cre and part of its DNA binding surface. To ensure

that the CSPs observed in these experiments are not an artifact of working with the catalytic

domain Cre^{Cat}, we compared spectra between full-length Cre and the same C-terminal

deletion: CreΔC (residues 1-330). An overlay of ^{1}H-^{15}N TROSY-HSQC spectrum of full-length

Cre on that of CreΔC showed the same CSPs for residues from the catalytic domain, and

minimal perturbation of signals attributed to Cre^{NTD} (Supplementary Fig. 5). Although CSPs

can be due to either direct or indirect interactions[40,41], the Cre^{Cat}ΔC CSP data suggest that

the C-terminal residues in Cre^{Cat} pack in *cis* over the active site and DNA binding surface of

Cre^{Cat}, not over the *trans* docking surface.

PRE-NMR data support *cis* docking of C-terminal helix αN of Cre

To clarify whether the CSPs in the core of Cre^{Cat} domain were induced by direct interactions

or via indirect allosteric effects, we attached a nitroxide spin probe at the extreme C-terminus

of Cre^{Cat} and performed paramagnetic relaxation enhancement (PRE)-NMR experiments[42–

44]. We constructed a variant of Cre^{Cat} (C155A/C240A/D343C) that enabled attachment of the

spin probe to the C-terminal cysteine residue and used ^{1}H-^{15}N TROSY-HSQC spectra to

verify that the mutations did not significantly perturb the structure of the protein (data not

shown). The single cysteine mutant was then conjugated with a S-(1-oxyl-2,2,5,5-tetramethyl-

2,5-dihydro-1H-pyrrol-3-yl) methyl methanesulfonothioate (MTSL) paramagnetic spin probe.

Near 100% tagging efficiency was verified using MALDI mass spectrometry (data not shown).

^{1}H-^{15}N TROSY-HSQC spectra of the MTSL tagged-protein under oxidized (paramagnetic)

and reduced (diamagnetic) conditions were obtained and a comparison of the peak intensities

was used to determine per-residue PRE effects in free Cre^{Cat}. In such a PRE-NMR

experiment, the lone-pair electron on a paramagnetic spin probe is expected to induce

distance-dependent line broadening for protons up to 25 Å away[45].

A strong correspondence between the C-terminus-linked PRE effects and previously

observed CSPs provide evidence for *cis* packing of the C-terminus over the active site and

DNA binding surface of Cre. In addition to residues sequentially neighboring C343, we

observed strong PRE effects from the C-terminal C343-MTSL spin probe to a part of the

interdomain linker (E129- A136), helix αH (R173- A178), helix αJ (S257- I264), αJ-K loop

(L284- R292) and helix αM (G314- Y324) (Fig. 4a, Supplementary Fig. 6). To interpret the

PRE effects in a structural context, we built a model of the MTSL tagged-protein using

PyMOL[46] with PDB ID 2HOI, chain B as the template, manually added the two missing C-

terminal residues (G342, C343), D343 substituted with cysteine, and attaching an MTSL

moiety (coordinates from PDB ID 2XIU)[47] to the C343 sidechain, and mapped the PRE data

onto this Cre^{Cat} structural model (Fig. 4c: left). Each of the regions showing strong PRE effects

in free Cre^{Cat} are at distances in excess of 25 Å as measured from the C343-MTSL-oxygen

to backbone amide nitrogen atoms: further than ~ 42 Å, ~ 47 Å, ~ 51 Å, ~ 42 Å, and ~ 37 Å,

respectively (Supplementary Fig. 6a).

To rule out the possibility that the observed PRE effects were due to intermolecular

interactions, we performed an intermolecular PRE experiment by recording ^{1}H-^{15}N TROSY-

HSQC spectra of a 1:1 mixture of [U-^{15}N]-Cre^{Cat} (non-tagged) with MTSL tagged- unlabeled

Cre^{Cat} C155A/C240A/D343C, under paramagnetic and diamagnetic conditions. Because the

^{15}N labels and MTSL tags are on different molecules any PRE effects must arise from

intermolecular interactions (Supplementary Fig. 7a)[48]. Only a few sparse residues located in

the *trans* docking surface of Cre showed intermolecular PRE effects, suggestive of very weak

transient intermolecular interaction (Supplementary Fig. 7b, c). Thus, the strong

intramolecular PRE effects in the ^{1}H-^{15}N TROSY-HSQC amide spectrum of free Cre^{Cat}

217 strongly support a model in which the C-terminal region makes a *cis* interaction with its own

218 DNA binding surface (Fig. 4c: left and right).

220 **PRE-restraints yield a *cis* docked model of the C-terminus of Cre, consistent with CSPs**

221 To determine whether the covalent structure of Cre^{Cat} would allow *cis* docking of helix αN region

222 without distorting other structural elements, we performed structure refinement using PRE-

223 derived distances as restraints. Analysis of the free Cre^{Cat} PRE I_{ox}/I_{red} values and ^{15}N R$_2$ relaxation

224 rates yielded 124 restraints (Supplementary Table 1). To account for a distribution of spin probe

225 positions, lower bounds were set to 5 Å below calculated distances while the upper bounds were

226 set to 10 Å higher than the calculated distances. ROSETTA energy minimization was performed

227 using upper and lower distance bounds, assuming a flexible C-terminus, β2-3 loop and αJ-K loop

228 regions; crystallographically observed secondary structure elements were enforced. Members of

229 the PRE-derived Cre^{Cat} ensemble (Fig. 5a) adopt stereochemically robust *cis* docked C-terminal

230 conformations that differ significantly from the conformation observed in synaptic complexes (Fig.

231 5b). These structures are consistent with CSP data, even though those were not used during

232 refinement (Fig. 5c). Difference Cα-Cα contact map between the top ten conformers and the

233 synaptic crystal structure (PDB ID 2HOI, chain A) show that the average changes in C-terminal

234 inter-residue distances when *cis* docked, ranged in values as large as 25 Å (Fig. 5d).

236 **DNA binding perturbs the C-terminal region in Cre^{Cat}**

237 Coincidence of the DNA binding site and *cis* docking surface of the Cre^{Cat} C-terminal region

238 led us to examine the effects of binding to a *loxP* DNA half-site substrate (Fig. 1c), on the

239 protein C-terminus. The ^{1}H-^{15}N TROSY-HSQC spectrum of the Cre^{Cat}-DNA half-site complex

240 showed CSPs in residues corresponding to the Cre-DNA interaction surface, comprising of

structural elements involved in DNA binding and the active site regions. Moreover, large

CSPs (and broadening of resonances) were also observed in the C-terminal residues E331-

D343, including helix αN (Fig. 6a, b). The CSPs mapped on the crystal structure (Fig. 6c)

show that *loxP* DNA half-site binds to Cre^{Cat} in the manner predicted from available tetrameric

structure models, but DNA binding also induces a change in the environment of the C-terminal

residues, resulting in the observed CSPs.

PRE-NMR data reveal a DNA binding-induced conformational change in the Cre C-

terminus

Like for the free protein, we used PRE-NMR to determine the conformational space

populated by the C-terminus of Cre upon binding to *loxP* DNA half-site. PRE effects were

measured using MTSL tagged-Cre^{Cat} C155A/C240A/D343C bound to the *loxP* half-site

hairpin DNA. PRE effects were remarkably different in the DNA complexes in comparison to

free Cre^{Cat} (Fig. 4b, Supplementary Fig. 8). Low I_{ox}/I_{red} ratios were observed for helix αF

(E138- M149), helix αG (G165- E176), β2 and β2-3 loop (G191- I195, G198 and V204), β3-

αI region (A212- R223), and helices αK- αL region (A291- G314). These regions together

form the *trans* docking cavity that accommodates the C-terminus of neighboring Cre

protomers in synaptic complexes (Fig. 1a, 4d: left and right). This change in PRE pattern (Fig.

4a versus 4b) clearly identifies a DNA-induced conformational change that involves

displacing the C-terminus from the DNA binding surface, to enable new protein-protein

interactions; however, they do not establish whether those new interactions occur in *cis*

(intramolecular), or in *trans* (intermolecular).

Intermolecular PRE-NMR studies with mixed isotopic labeling were then used to clarify

whether the PRE effects observed in the presence of DNA arise from intra or intermolecular

interactions. We mixed DNA-bound non-tagged [U-^{15}N]-Cre^{Cat} and MTSL-tagged unlabeled

Cre^{Cat} C155A/C240A/D343C at a 1:1 molar ratio, to study possible intermolecular interactions

in Cre^{Cat} bound to DNA. Because these mixed samples have the potential to interact in four

different relative orientations (Supplementary Fig. 9a), only one of which would generate

measurable PRE effects (i.e., MTSL tagged- unlabeled Cre^{Cat} C155A/C240A/D343C with C-

terminus docking onto a [U-^{15}N]-Cre^{Cat}), the experiment would be predicted to yield weaker

PRE effects in comparison to the experiments with uniformly labeled and MTSL tagged-

Cre^{Cat}-DNA complex (Fig. 4b); although observation of such effects in the same regions of

the protein would support *trans* interactions. Despite the resonances being broader and data

quality being generally lower, we indeed observed strong PRE effects consistent with docking

of DNA-bound Cre C-terminal region over the *trans* docking surface (Supplementary Fig. 9b,

c); these effects *must* arise from intermolecular interactions because of the labeling pattern

used. Yet, a comparison to the similarly strong PRE effects in uniformly tagged and labelled

Cre^{Cat}-DNA complex studies (Supplementary Fig. 9c versus Fig. 4b) illustrates the possibility

of a population of DNA-bound Cre C-terminus docking in *cis* over its own *trans* docking cavity,

while another population extends out to dock in *trans* into the same cavity on another DNA-

bound Cre molecule.

DISCUSSION

Cre recombinase has emerged as an important tool in molecular and cellular biology and has

several features that make it an attractive reagent for gene editing[7,49,50]. Reaching that

potential requires thorough characterization of its mechanism for DNA site selection, and

control of its DNA cleavage and recombination activity. High-resolution structural studies of

Cre, largely limited to crystal structures of DNA bound tetrameric synaptic complexes, have

suggested a role for conformational plasticity in facilitating the steps within the reaction

mechanism. We set out to characterize the solution behavior of Cre in its free and DNA bound

states in order to understand the link between protein dynamics and regulation of Cre

assembly and activity. While solution NMR studies yielded some results consistent with

previous understanding of this prototypical member of the tyrosine recombinase family, other

surprising results call for a revised view of the reaction mechanism.

Solution NMR data showed Cre^{Cat} to be independent of Cre^{NTD}. The well-dispersed

signals in the ^1H-^{15}N TROSY-HSQC NMR spectrum of Cre^{Cat} are nearly superimposable on

that of full-length Cre, except for the signals attributable to N-terminal domain Cre^{NTD}

(Supplementary Fig. 1b). The domains' structural independence could imply variability in

inter-domain orientations that would be expected to prevent formation of suitable crystal

lattices for x-ray diffraction. It however facilitated detailed NMR studies with the Cre^{Cat} domain,

wherein reside the active site residues and sequence specificity determinants, focused on

testing the role of protein dynamics in regulation of Cre function[33–36].

Our findings of the protein dynamics in specific regions within Cre^{Cat} revealed unique

spectral signatures linked to their functional roles. Underlying flexibility in Cre^{Cat} revealed by

nuclear spin relaxation measurements (^{15}N R_1, R_2 and {^1H}-^{15}N HetNOE), were consistent

with the monomeric 217-residue protein construct (τ_C, 13.2 ns). A portion of the linker, the

β2-3 loop, the αJ-K loop and the αM-N loop each exhibit higher R_1, lower R_2, and reduced

HetNOE values, indicative of fast internal motions. Within the Cre_4-$loxP_2$ tetrameric synaptic

complexes, conformational differences in the β2-3 loop and the C-terminal regions (including

helix αN) that form structural bridges that link adjacent protomers, and in helix αM bearing

the catalytic tyrosine, distinguish the catalytically "active" and "inactive" Cre protomers[5,16,22].

Flexibility of the β2-3 loop observed in free Cre^{Cat} could be expected to prevent stabilization

of the active site geometry in uncoordinated Cre protomers. The alternating conformations of

the loop in tetrameric synaptic complexes indicates that this flexibility will be still retained,

thereby enabling its regulatory role over adjacent active site structures. The protein dynamics

observed in the αJ-K loop in the free protein may serve to facilitate DNA binding and could

thereafter be predicted to undergo quenching since this region does not show structural

plasticity between the various Cre reaction intermediates. The surprise within the protein

dynamics analysis lies in the observation that while residues flanking C-terminal helix αN

exhibit flexibility on the ps-ns timescale, αN itself does not, and instead exhibits relaxation

rates consistent with overall tumbling of the protein, counter to the expectation that the Cre

C-terminus would flexibly adopt a range of extended conformations to facilitate capturing an

adjacent protomer via *trans* docking contacts.

An unexpected C-terminal *cis* docking interaction in the proximity of the free Cre active

site was revealed from our highly correlated CSP and PRE-NMR experiments. Given the

known *trans* docking site, and prior incongruities regarding *cis* or *trans* cleavage by Cre and

other recombinases[51], a logical presumption was that in free Cre, helix αN might dock in *cis*

in the same site it occupies when assembled in *trans* in synaptic complexes; indeed, such a

premise was previously tested experimentally using disulfide crosslinking strategies[39]. Were

that the case, one would predict that deletion of helix αN would result in CSPs for residues

flanking that *trans* docking surface cavity. Instead, we observed the largest CSPs for residues

on the other side of the protein – namely, the DNA binding and active site surfaces (Fig. 3).

Since CSPs can arise from both proximity and induced conformational changes, we employed

distance-dependent effects in PRE-NMR experiments to conclusively demonstrate the

proximity of the protein C-terminus to the active site and DNA binding regions (Fig. 4a).

Consistent with size exclusion chromatography (Supplementary Fig. 1a), NMR relaxation

(Fig. 2) and previous sedimentation velocity experiments[17] that show unbound Cre to be

monomeric in solution, the absence of strong intermolecular PREs (Supplementary Fig. 7)

indicates that docking of the C-terminus of free Cre onto the DNA binding surface indeed

occurs *intra*-molecularly, in *cis*. *Cis*-docked models generated using distance bounds derived

from the PRE data satisfy the restraints, covalent structure, and are in excellent agreement

with the CSPs obtained upon deletion of the C-terminal region (Fig. 5). A comparison between

the free Cre^{Cat} ensemble model and the Cre^{Cat} structure from crystallographic DNA bound

synaptic complex protomers shows changes in inter-residue contacts as large as 20- 25 Å

(Fig. 5d). Thus, these data provide strong evidence that in the absence of DNA the C-terminal

residues dock in *cis* over the DNA binding and active site surfaces, in an apparent auto-

inhibitory state.

A Cre structure in which the C terminus is docked in *cis* over the active site would

seem incompatible with a DNA-bound state – so, how is this conundrum resolved? PRE-NMR

data on the Cre^{Cat}-DNA complex show that the C-terminus of Cre^{Cat} is displaced from the *cis*

docking surface when DNA binds there, since PRE effects are instead observed at the *trans*

docking cavity (Fig. 4b). This *loxP* binding-induced pattern of PREs at the *trans* docking site

clearly indicates relocation of C-terminal helix αN, but does not clarify whether docking occurs

in *cis*, or in *trans* to a nearby Cre molecule. The use of a DNA *loxP* half-site substrate would

be expected to prevent assembly of pre-synaptic complexes with two Cre molecules bound

to DNA; however, PRE effects can be observed even when interactions are transient[52].

Although partially attenuated, intermolecular PRE effects were indeed observed at the same

trans docking cavity in experiments where the NMR signals from DNA-bound Cre^{Cat} arise

from protein molecules not tagged with MTSL (Supplementary Fig. 9). This attenuation could

result from the fact that only one-fourth of protein-protein contacts within DNA-bound

complexes can produce *intermolecular* PRE effects (i.e., when C-terminus from MTSL-tagged

un-labeled Cre[Cat] docks in *trans* over non-tagged [U-[15]N]-Cre[Cat] when bound to DNA;

Supplementary Fig. 9a), or may indicate that in the presence of DNA, *intramolecular cis*

docking over the protomer's own *trans* docking cavity is also possible. However, molecular

modeling calculations starting from the DNA-bound crystal structure (data not shown)

indicated that the C-terminal helix αN region could not pack over its own *trans* docking cavity

without significant conformational strain, involving additional remodeling that would be

inconsistent with the NMR CSP data (Fig. 5). We conclude that in the absence of DNA, the

C-terminal residues spanning helix αN occupy the DNA binding surface of Cre, and that

binding to DNA produces a large conformational change that enables productive *trans*

protein-protein interactions with adjacent Cre molecules.

The rigidity and apparent auto-inhibitory conformation observed in C-terminal residues

in free Cre could be expected to disfavor oligomerization of the protein prior to *loxP* DNA

binding and play a role in selectively inhibiting spurious uncoordinated DNA cleavage and

recombination activity. Although it is unclear whether the *cis* docking conformation of C-

terminal region in free Cre serves additional roles, for example by obstructing inadvertent

binding to non-cognate DNA substrates, our data suggests that the conformational

rearrangements following the displacement of *cis* docking C-terminal upon *loxP* DNA half-site

binding, comprise the mechanistic step that could trigger Cre protomer stepwise assembly on

loxP through protein-protein interactions via the now exposed C-terminal residues.

These studies expand our understanding of the regulatory role played by C-terminal

residues in Cre [5,16,17]. DNA-binding associated conformational changes mediated by C-

terminal regions of Cre and related recombinases have been implicated in catalytic activation

and initiation of oligomerization[2,5,28]. However, the structural basis for Cre to undergo

stepwise assembly on the DNA recognition sites have remained poorly understood due to

lack of structural information on pre-synaptic reaction intermediates. The features of Cre and

Cre-*loxP* complexes presented here provide insight into both the unliganded state of Cre, as

well as the mechanism of initiation of protein-protein interactions that lead to stepwise

assembly of recombinogenic intasomes. Knowledge gaps remain, including understanding of

precise structural rearrangements that accompany the *cis* to *trans* switch in pre-synaptic Cre

recombinase. Nevertheless, these investigations provide a new NMR perspective on Cre

recombinase and represent a transformative step towards understanding the structure,

function and mechanism of this important and widely used gene editing tool.

METHODS

Protein expression, purification, and site-directed mutagenesis

The deletion construct containing the catalytic domain and the interdomain linker (Cre^{Cat},

residues 127- 343) was prepared using QuikChange Site-Directed Mutagenesis Kit (Agilent)

from a pET21A vector (Novagen) encoding WT Cre Recombinase (provided by Dr. Gregory

Van Duyne, U. Penn). Further truncation and point mutants of Cre^{Cat} construct were prepared

using the Q5 Site-Directed Mutagenesis Kit (NEB). Protein expression from *E. coli* BL21(DE3)

cells was carried out by growing freshly transformed cells (using Ampicillin antibiotic, 100

mg/L) in LB media (natural abundance Cre) or M9 minimal medium (isotopically labeled [U-

^{15}N] and [U-^{13}C,^{15}N] Cre) supplemented with 1 g/L [^{15}N]-ammonium chloride (Cambridge

Isotopes) as the sole nitrogen source or 2 g/L [^{13}C]-glucose (Cambridge Isotopes) as the sole

carbon source in addition to [^{15}N]-ammonium chloride. The cells were grown (shaking at 220

rpm, 37 °C) to an OD$_{600}$ of 0.6- 0.8 and induced with 0.75 mM isopropyl β-D-1-

thiogalactopyranoside (Gold Biotechnology) for 12 h. After harvesting the cells (4000 g, 15

min, 4 °C) and sonication, the Cre constructs in 40 mM tris, 100 mM NaCl, pH 7.0 buffer

containing 1 protease inhibitor cocktail tablet (Roche), were purified using 5 ml SPFF cation

exchange column (GE Healthcare) with a 100 ml gradient of 0.1 to 1 M NaCl. The protein

fractions (eluted at ~ 0.4 M NaCl) were combined and diluted four times with low salt buffer

40 mM tris, 100 mM NaCl pH 7.0 and further purified on a 5 ml Heparin affinity column (GE

Healthcare) with a 100 ml gradient of 0.1 to 1.5 M NaCl. Protein fractions (eluted at ~ 1 M

NaCl) were then purified using a HiLoad 16/60 Superdex 75 (GE Healthcare) column (SEC)

in 40 mM tris, 500 mM NaCl pH 7.0 buffer. Purified protein concentrations were determined

using UV-Vis spectroscopy (ε for Cre^{Cat} = 23,950 M^{-1} cm^{-1} at 280 nm). Typical yields obtained

were ~20-25 mgs protein per L of culture.

Cre^{Cat}-DNA complex NMR sample preparation

loxP half-site hairpin was formed from single-stranded DNA oligos (IDT) containing

recombinase binding elements along with additional GC bps at ends, symmetrized spacer

bps and a GAA hairpin[53]:

5'-GCATAACTTCGTATAGCATATGCGAAGCATATGCTATACGAAGTTATGC-3'.

The lyophilized DNA oligo was resuspended in H_2O and heated at 95 °C for 15 mins and

immediately cooling on ice. To avoid precipitation, Cre^{Cat} and *loxP* DNA hairpin at low

concentrations (< 50 µM) in high salt buffer (10 mM Tris, > 500 mM NaCl, pH 7.0) were mixed

by stepwise titration of the protein into DNA. The solution was then dialyzed into low salt NMR

buffer (10 mM tris, 15 mM NaCl, 0.02% NaN_3, pH 7.0) at 4 °C. The protein-DNA complex

sample was then concentrated using 500 µL, 1 kDa centrifugal filters (VWR).

NMR data collection

Purified [U-^{15}N]- or [U-^{15}N,^{13}C]- Cre^{Cat} samples were concentrated to 0.5- 1.0 mM after dialysis into buffer containing 10 mM Tris, 100 mM NaCl, 0.02 % NaN$_3$, pH 7.0. When required, the samples were exchanged into 10 mM d$_{11}$-Tris (Sigma), 100 mM NaCl, 0.02 % NaN$_3$, pH 7.0 buffer, using Sephadex PD10 columns (GE Healthcare). 5-10 % (v/v) D$_2$O (99 %) and 0.66 mM DSS were added to the NMR samples. NMR data were recorded on Bruker Avance III HD spectrometers operating at 850, 800 or 600 MHz equipped with a 5 mm triple resonance cryoprobes and z axis gradients, at 25 °C. Data were processed with NMRPipe[54] or NMRFx[55] and visualized using NMRView[56]. Typical ^{1}H-^{15}N TROSY-HSQC 2D correlation spectra were recorded using 16- 24 scans and 2048 x 128 data points.

NMR chemical shift assignment and perturbation mapping

For backbone chemical shift assignments, TROSY triple-resonance spectra HNCO, HNCA, HN(CO)CA, HN(CA)CO, and non-TROSY CBCA(CO)NH and HA(CO)NH were recorded on purified [U-^{13}C,^{15}N] Cre^{Cat} sample. Backbone assignment was achieved using NMRView aided by PINE[57] and CARA[58].

CSPs were calculated from differences in peak positions using:

$$\Delta\delta_{NH} = \sqrt{\Delta\delta_H{}^2 + \frac{(\Delta\delta_N)^2}{25}}$$

where $\Delta\delta_H$ is the chemical shift difference in the ^{1}H dimension and $\Delta\delta_N$ is the chemical shift change in the ^{15}N dimension. Corrected standard deviations were determined by calculating

the SD of CSPs of all assigned residues, removing CSP data with > 3 SD and recalculating

the SD.

^{15}N relaxation measurements

Backbone amide relaxation measurements were performed at 800 MHz proton Larmor

frequency using [U-^{15}N] Cre^{Cat} at a concentration of 800 µM in 10 mM Tris, 100 mM NaCl,

0.02 % NaN$_3$, pH 7.0 buffer (5 % D$_2$0 v/v). TROSY versions of R$_1$ and R$_2$ experiment data

were acquired using recycle delay of 2 seconds between experiments, and the following

relaxation delays for R$_1$: 0, 560, 1120 and 1680 ms with repeats and R$_2$: 2, 24.2, 48.2, and

72.2 ms. The R$_1$ and R$_2$ values were obtained by fitting the intensity of peaks to exponential

decay function and errors were determined by relaxation curve fitting. 80-90 % of the amides

were analyzed, after excluding those with significant resonance overlap. {^{1}H}-^{15}N HetNOE

values were obtained by recording spectra with and without a ^{1}H pre-saturation period (8 s),

in which ^{1}H signals were saturated using a train of 90° pulses, applied before the start of

experiment. Non-overlapping assigned peaks were analyzed and the intensity ratio were

determined using the HetNOE analysis tool within NMRViewJ; uncertainties were obtained

from standard deviation of noise in the spectra.

Paramagnetic Relaxation Enhancement (PRE) - NMR

We constructed a single-cysteine Cre^{Cat} variant C155A/C240A/D343C to avoid unintended

paramagnetic tagging of the native cysteines and thereby enable probing just the C-terminus.

S-(1-oxyl-2,2,5,5-tetramethyl- 2,5-dihydro-1H-pyrrol-3-yl) methyl methanesulfonothioate

(MTSL) tagging of Cre^{Cat} C155A/C240A/D343C was achieved by following a published

protocol[59]. Briefly, a 200 mM MTSL (Toronto Research Chemicals) stock was made by adding

189 µL of acetonitrile to 10 mg of MTSL (stored at -20 °C, protected from light). DTT was

added to purified protein in high salt conditions (10 mM Tris, 500 mM NaCl, 5 mM DTT, pH

7.0) to reduce possible disulfide bonds. The reducing agent was then removed by rapid buffer

exchange into 10 mM Tris, 500 mM NaCl, pH 7.0 buffer using a PD10 desalting column (GE

Healthcare). The sample was collected into exchange buffer containing ten-fold molar excess

of MTSL and stirred at room temperature overnight. Excess MTSL was removed after reaction

by buffer exchange with a second PD10 column, followed by dialysis (MWCO 3 kDa) into

NMR buffer. Near 100 % MTSL tagging of the protein was verified using MALDI MS

measurements as indicated by an increase in mass by 186 Da (weight of attached probe).

Final sample concentrations ranged between 100-200 µM at a volume of ~ 500 µL.

For PRE-NMR measurements, ^{1}H-^{15}N TROSY-HSQC spectra were recorded on the same

MTSL tagged-Cre^{Cat} C155A/C240A/D343C sample before (paramagnetic) and after

(diamagnetic) reduction of the spin probe by treatment with five-fold molar excess of sodium

ascorbate (250 mM stock used for minimal sample dilution) for three hours at room

temperature. Peak intensities were measured and normalized using intensities of residues

(D153, E222, V230) > 45 Å away from C-terminus in PDB ID 2HOI, chain B. For MTSL

probes, I_{ox}/I_{red} ratios between 0 and 1 indicate that the distance of probe to proton is within

13- 25 Å. Uncertainties in I_{ox}/I_{red} values were propagated from signal-to-noise ratio of each

resonance in the two spectra using:

$$\sigma I_{ox}/I_{red} = I_{ox}/I_{red} \sqrt{(\frac{\sigma I_{ox}}{I_{ox}})^2 + (\frac{\sigma I_{red}}{I_{red}})^2}$$

where $\sigma I_{ox}/I_{red}$ is the calculated error in I_{ox}/I_{red} ratio, σI_{ox} is the standard deviation of the noise

in the MTSL oxidized spectrum, and σI_{red} is the standard deviation in the reduced spectrum.

For the intermolecular PRE studies, [U-^{15}N]-Cre^{Cat} (non-MTSL-tagged) was mixed with MTSL

tagged-natural abundance Cre^{Cat} C155A/C240A/D343C, at a 1:1 molar ratio in NMR buffer

(10 mM Tris, 100 mM NaCl, pH 7.0) at 25 °C. For similar studies with the protein-DNA

complex, the complex samples with/without MTSL tags were prepared separately and mixed

at a molar ratio of 1:1.

PRE-restrained modeling using ROSETTA

PRE derived-distance constraints for use in structural modeling were generated using the

approach of Battiste and Wagner[42]. Briefly, per residue R_2^{sp} values (paramagnetic relaxation

rate due to the spin tag) were obtained from:

$$I_{ox}/I_{red} = \frac{R_2 e^{-R_2^{sp}t}}{R_2 + R_2^{sp}}$$

where R_2 is the transverse relaxation rate of each amide and t is the acquisition time in the

proton dimension. R_2^{sp} values were then used to obtain the approximate distance, r (Å)

between the spin tag to the amide proton of each residue:

$$r = \left[\frac{K}{R_2^{sp}}\left(4\tau_C + \frac{3\tau_C}{1 + \omega_h^2\tau_c^2}\right)\right]^{\frac{1}{6}}$$

where τ_C (s) is per-residue correlation time of amide protons determined from R_2/R_1 relaxation

ratios, ω_h (Hz) is the Larmor frequency of each amide proton, and K is a constant (1.23 x 10^{-}

^{32} cm^6 sec^{-2}) that encapsulates the spin properties of the MTSL tag[42].

Distance restraints were used in a bounded form (typical for NOE constraints). Due to the

relative imprecision of this method of constraint distance calculation[44,60], residues with an

I_{ox}/I_{red} value less than 0.8 were assigned a lower bound of calculated r - 5 Å and an upper

bound was set to $r + 10$ Å. Residues with an I_{ox}/I_{red} value greater than 0.8 were assigned a

lower bound of 20 Å and no upper bound. Constraints for 124 residues in Cre^{Cat} were obtained

wherein data (I_{ox}, $I_{red,}$ R$_2$, ω_h, and τ_C) for each amide proton was available.

ROSETTA energy minimization was performed using the relax application[61–65]. Chain

A of PDB ID 2HOI (residues 127- 341, and G342 and D343C modeled) was used as the input

structure with the D343C mutation. A move map file retained secondary structure of Cre^{Cat}

along with the following impositions: the β2-3 and αJ-K loops and interdomain linker residues

(A127-A134) were allowed to have backbone φ, ψ, and sidechain χ$_1$ torsional freedom based

on flexibility observed via ^{15}N R$_2$/R$_1$ data. C-terminal E331- C343 were also allowed to sample

backbone and χ$_1$ angles allowing movement of helix αN through space. Other residues were

only allowed to undergo sidechain minimization. Constraints were used to generate 100 initial

structures subsequently scored using the ref2015_cst weights function[66]. The ten structures

with the lowest *atom_pair_constraint* energy were selected as the input structures for a

second round of minimization wherein ten structures were generated for each input structure.

The resulting 100 structures constitute the PRE-NMR constrained ensemble model of free

Cre^{Cat}. Coarse grained contact maps were generated using MATLAB by calculating backbone

inter-residue Cα-Cα distances from PDB files (2HOI, chain A residues 127- 341 and G342

and D343C modeled, and top ten lowest energy members of free Cre^{Cat} PRE-derived

ensemble). Absolute values of change in inter-residue distances between the contact maps

were determined by subtracting one distance matrix from the other.

DATA AVAILABILITY

The Cre^{Cat} (residues 127- 343) backbone chemical shift assignments have been deposited

in the Biological Magnetic Resonance Data Bank, www.bmrb.wisc.edu (accession no. 50270).

ACKNOWLEDGEMENTS

We thank Dr. Chunhua Yuan, Dr. Alex Hansen and Dr. Arpad Somogyi of the Ohio State University Campus Chemical Instrumentation Center for assistance with NMR experimental setup and mass spectrometry analysis and Dr. Eric Danhart for guidance with NMR data processing. We also thank Dr. Gregory Van Duyne (U. Penn) for providing the plasmids encoding Cre. This study was supported by NIH grant R01 GM122432 (to M.P.F), NSF MCB grant 0092962 (to M.P.F), and NIH/NIGMS grant 1P41GM111135 (to NMRBox).

AUTHOR CONTRIBUTIONS

Author contributions: A.U., C.A. and D.K.Y. and M.P.F. designed experiments; A.U., C.A., D.K.Y and D.P. performed research; A.U., D.K.Y., C.A., K.S. and M.P.F. analyzed and interpreted data; and A.U. and M.P.F. wrote the manuscript.

COMPETING INTERESTS

The authors declare no competing financial interests.

REFERENCES

1.   Bucholtz, F. Principles of site-specific recombinase (SSR) technology. *J. Vis. Exp.* (2008) doi:10.3791/718.

2.   Nunes-Düby, S. E., Kwon, H. J., Tirumalai, R. S., Ellenberger, T. & Landy, A. Similarities and differences among 105 members of the Int family of site-specific recombinases. *Nucleic Acids Res.* **26**, 391–406 (1998).

3.   Wirth, D. *et al.* Road to precision: recombinase-based targeting technologies for
genome engineering. *Current Opinion in Biotechnology* (2007)
doi:10.1016/j.copbio.2007.07.013.

4.   Fogg, P. C. M., Colloms, S., Rosser, S., Stark, M. & Smith, M. C. M. New applications
for phage integrases. *Journal of Molecular Biology* (2014)
doi:10.1016/j.jmb.2014.05.014.

5.   Van Duyne, G. D. Cre Rcombinase. *Microbiol. Spectr.* (2015).

6.   Meinke, G., Bohm, A., Hauber, J., Pisabarro, M. T. & Buchholz, F. Cre Recombinase
and Other Tyrosine Recombinases. *Chem. Rev.* (2016)
doi:10.1021/acs.chemrev.6b00077.

7.   Lanza, A. M., Dyess, T. J. & Alper, H. S. Using the Cre/lox system for targeted
integration into the human genome: LoxFAS-loxP pairing and delayed introduction of
Cre DNA improve gene swapping efficiency. *Biotechnol. J.* (2012)
doi:10.1002/biot.201200034.

8.   Grindley, N. D. F., Whiteson, K. L. & Rice, P. A. Mechanisms of Site-Specific
Recombination. *Annu. Rev. Biochem.* (2006)
doi:10.1146/annurev.biochem.73.011303.073908.

9.   Landy, A. The λ Integrase Site-specific Recombination Pathway. *Microbiol. Spectr.*
(2015) doi:10.1128/microbiolspec.mdna3-0051-2014.

10.   Gaj, T., Gersbach, C. A. & Barbas, C. F. ZFN, TALEN, and CRISPR/Cas-based
methods for genome engineering. *Trends in Biotechnology* (2013)
doi:10.1016/j.tibtech.2013.04.004.

11.   Cox, D. B. T., Platt, R. J. & Zhang, F. Therapeutic genome editing: Prospects and
challenges. *Nature Medicine* (2015) doi:10.1038/nm.3793.

12.   Groth, A. C. & Calos, M. P. Phage integrases: Biology and applications. *Journal of*
Molecular Biology (2004) doi:10.1016/j.jmb.2003.09.082.

13.   Craig, N. L. Conservative site-specific recombination. 77–105 (1988).

14.   Birling, M. C., Gofflot, F. & Warot, X. Site-specific recombinases for manipulation of the
mouse genome. *Methods Mol. Biol.* (2009) doi:10.1007/978-1-60327-019-9_16.

15.   Sternberg, N. Bacteriophage P1 site-specific recombination. *J. Mol. Biol.* **150**, 603–608
597   (1981).

598   16.   Guo, F., Gopaul, D. N. & Van Duyne, G. D. Structure of Cre recombinase complexed
599   with DNA in a site-specific recombination synapse. *Nature* (1997) doi:10.1038/37925.

600   17.   Ghosh, K., Guo, F. & Van Duyne, G. D. Synapsis of loxP sites by cre recombinase. *J.*
601   *Biol. Chem.* (2007) doi:10.1074/jbc.M703283200.

602   18.   Ghosh, K. & Van Duyne, G. D. Cre-loxp biochemistry. *Methods* (2002)
603   doi:10.1016/S1046-2023(02)00244-X.

604   19.   Gopaul, D. N., Guo, F. & Van Duyne, G. D. Structure of the Holliday junction
605   intermediate in Cre-loxP site-specific recombination. *EMBO J.* **17**, 4175–4187 (1998).

606   20.   Martin, S. S., Pulido, E., Chu, V. C., Lechner, T. S. & Baldwin, E. P. The order of strand
607   exchanges in Cre-LoxP recombination and its basis suggested by the crystal structure
608   of a Cre-LoxP holliday junction complex. *J. Mol. Biol.* (2002) doi:10.1016/S0022-
609   2836(02)00246-2.

610   21.   Ghosh, K., Lau, C. K., Gupta, K. & Van Duyne, G. D. Preferential Synapsis of Loxp
611   Sites Drives Ordered Strand Exchange in Cre-Loxp Site-Specific Recombination. *Nat.*
612   *Chem. Biol.* (2005) doi:10.1038/nchembio733.

613   22.   Van Duyne, G. D. A structural view of Cre-loxP site-specific recombination. *Annual*
614   *Review   of   Biophysics   and   Biomolecular   Structure*   (2001)
615   doi:10.1146/annurev.biophys.30.1.87.

616   23.   Rajeev, L., Malanowska, K. & Gardner, J. F. Challenging a Paradigm: the Role of DNA
617   Homology in Tyrosine Recombinase Reactions. *Microbiol. Mol. Biol. Rev.* (2009)
618   doi:10.1128/mmbr.00038-08.

619   24.   Seah, N. E. *et al.* Nucleoprotein architectures regulating the directionality of viral
620   integration   and   excision.   *Proc.   Natl.   Acad.   Sci.   U.   S.   A.*   (2014)
621   doi:10.1073/pnas.1413019111.

622   25.   Baldwin, E. P. *et al.* A specificity switch in selected Cre recombinase variants is
623   mediated   by   macromolecular   plasticity   and   water.   *Chem.   Biol.*   (2003)
624   doi:10.1016/j.chembiol.2003.10.015.

625   26.   Duyne, G. D. Van. A STRUCTURAL VIEW OF Cre-loxP SITE -SPECIFIC
626   RECOMBINATION. *Annu. Rev. Biophys. Biomol. Struct.* (2001).

627   27.   Zhang, C. *et al.* Redesign of the monomer-monomer interface of Cre recombinase
628   yields an obligate heterotetrameric complex. *Nucleic Acids Res.* (2015)
629   doi:10.1093/nar/gkv901.

630   28.   Aihara, H., Kwon, H. J., Nunes-Düby, S. E., Landy, A. & Ellenberger, T. A
631   conformational switch controls the DNA cleavage activity of λ integrase. *Mol. Cell*
632   (2003) doi:10.1016/S1097-2765(03)00268-5.

633   29.   Subramaniam, S., Tewari, A. K., Nunes-Duby, S. E. & Foster, M. P. Dynamics and
634   DNA substrate recognition by the catalytic domain of lambda integrase. *J. Mol. Biol.*
635   **329**, 423–439 (2003).

636   30.   Serre, M. C. *et al.* The Carboxy-Terminal αN Helix of the Archaeal XerA Tyrosine
637   Recombinase Is a Molecular Switch to Control Site-Specific Recombination. *PLoS One*
638   (2013) doi:10.1371/journal.pone.0063010.

639   31.   Hickman, A. B., Waninger, S., Scocca, J. J. & Dyda, F. Molecular organization in site-
640   specific recombination: The catalytic domain of bacteriophage HP1 integrase at 2.7 Å
641   resolution. *Cell* (1997) doi:10.1016/S0092-8674(00)80202-0.

642   32.   Kazmierczak, R. A., Swalla, B. M., Burgin, A. B., Gumport, R. I. & Gardner, J. E.
643   Regulation of site-specific recombination by the C-terminus of λ integrase. *Nucleic*
644   *Acids Res.* **30**, 5193–5204 (2002).

645   33.   Gibb, B. *et al.* Requirements for catalysis in the Cre recombinase active site. *Nucleic*
646   *Acids Res.* **38**, 5817–5832 (2010).

647   34.   Hoess, R., Abremski, K., Irwin, S., Kendall, M. & Mack, A. DNA specificity of the cre
648   recombinase resides in the 25 kDa carboxyl domain of the protein. *J. Mol. Biol.* **216**,
649   873–882 (1990).

650   35.   Rufer, A. W. Non-contact positions impose site selectivity on Cre recombinase. *Nucleic*
651   *Acids Res.* **30**, 2764–2771 (2002).

652   36.   Kim, S. T., Kim, G. W., Lee, Y. S. & Park, J. S. Characterization of Cre-loxP interaction
653   in the major groove: Hint for structural distortion of mutant Cre and possible strategy

for HIV-1 therapy. *J. Cell. Biochem.* **80**, 321–327 (2000).

37.   Lee, L. & Sadowski, P. D. Identification of Cre residues involved in synapsis,
isomerization, and catalysis. *J. Biol. Chem.* **278**, 36905–36915 (2003).

38.   Kay, L. E., Torchia, D. A. & Bax, A. Backbone Dynamics of Proteins As Studied by 15N
Inverse Detected Heteronuclear NMR Spectroscopy: Application to Staphylococcal
Nuclease. *Biochemistry* (1989) doi:10.1021/bi00449a003.

39.   Yuan, P. Structural and biochemical studies of site-specific recombinases. *Diss.*
available from ProQuest 1–182 (2008).

40.   Foster, M. P. *et al.* Chemical shift as a probe of molecular interfaces: NMR studies of
DNA binding by the three amino-terminal zinc finger domains from transcription factor
IIIA. *J. Biomol. NMR* (1998) doi:10.1023/A:1008290631575.

41.   Williamson, M. P. Using chemical shift perturbation to characterise ligand binding.
Progress in Nuclear Magnetic Resonance Spectroscopy (2013)
doi:10.1016/j.pnmrs.2013.02.001.

42.   Battiste, J. L. & Wagner, G. Utilization of site-directed spin labeling and high-resolution
heteronuclear nuclear magnetic resonance for global fold determination of large
proteins with limited nuclear overhauser effect data. *Biochemistry* (2000)
doi:10.1021/bi000060h.

43.   Marius Clore, G. & Iwahara, J. Theory, practice, and applications of paramagnetic
relaxation enhancement for the characterization of transient low-population states of
biological macromolecules and their complexes. *Chem. Rev.* (2009)
doi:10.1021/cr900033p.

44.   Clore, G. M. Practical Aspects of Paramagnetic Relaxation Enhancement in Biological
Macromolecules. in *Methods in Enzymology* (2015). doi:10.1016/bs.mie.2015.06.032.

45.   Altenbach, C., Oh, K. J., Trabanino, R. J., Hideg, K. & Hubbell, W. L. Estimation of
inter-residue distances in spin labeled proteins at physiological temperatures:
Experimental strategies and practical limitations. *Biochemistry* (2001)
doi:10.1021/bi011544w.

46.   DeLano, W. L. Pymol: An open-source molecular graphics tool. *CCP4 Newsl. Protein*

Crystallogr. (2002).

47.   Gruene, T. *et al.* Integrated analysis of the conformation of a protein-linked spin label
by crystallography, EPR and NMR spectroscopy. *J. Biomol. NMR* (2011)
doi:10.1007/s10858-011-9471-y.

48.   Janowska, M. K. & Baum, J. Intermolecular paramagnetic relaxation enhancement
(PRE) studies of transient complexes in intrinsically disordered proteins. in *Methods in*
Molecular Biology (2015). doi:10.1007/978-1-4939-2978-8_3.

49.   Brault, V., Besson, V., Magnol, L., Duchon, A. & Hérault, Y. Cre/loxP-mediated
chromosome engineering of the mouse genome. *Handbook of Experimental*
Pharmacology (2007) doi:10.1007/978-3-540-35109-2-2.

50.   Sengupta, R. *et al.* Viral Cre-LoxP tools aid genome engineering in mammalian cells.
J. Biol. Eng. (2017) doi:10.1186/s13036-017-0087-y.

51.   Esposito, D. & Scocca, J. J. The integrase family of tyrosine recombinases: Evolution
of a conserved active site domain. *Nucleic Acids Res.* (1997)
doi:10.1093/nar/25.18.3605.

52.   Clore, G. M. Seeing the invisible by paramagnetic and diamagnetic NMR. *Biochem.*
Soc. Trans. (2013) doi:10.1042/BST20130232.

53.   Yoshizawa, S., Kawai, G., Watanabe, K., Miura, K. I. & Hirao, I. GNA trinucleotide loop
sequences producing extraordinarily stable DNA minihairpins. *Biochemistry* (1997)
doi:10.1021/bi961738p.

54.   Delaglio, F. *et al.* NMRPipe: A multidimensional spectral processing system based on
UNIX pipes. *J. Biomol. NMR* (1995) doi:10.1007/BF00197809.

55.   Norris, M., Fetler, B., Marchant, J. & Johnson, B. A. NMRFx Processor: a cross-
platform NMR data processing program. *J. Biomol. NMR* (2016) doi:10.1007/s10858-
016-0049-6.

56.   Johnson, B. A. Using NMRView to visualize and analyze the NMR spectra of
macromolecules. *Methods Mol. Biol.* (2004) doi:10.1385/1-59259-809-9:313.

57.   Bahrami, A., Assadi, A. H., Markley, J. L. & Eghbalnia, H. R. Probabilistic interaction
network of evidence algorithm and its application to complete labeling of peak lists from

protein NMR spectroscopy. *PLoS Comput. Biol.* (2009)
doi:10.1371/journal.pcbi.1000307.

58. Keller, R. *The computer aided resonance assignment tutorial. Goldau, Switzerland:*
Cantina Verlag (2004).

59. Sjodt, M. & Clubb, R. Nitroxide Labeling of Proteins and the Determination of
Paramagnetic Relaxation Derived Distance Restraints for NMR Studies. *BIO-*
PROTOCOL (2017) doi:10.21769/bioprotoc.2207.

60. Iwahara, J., Schwieters, C. D. & Clore, G. M. Ensemble Approach for NMR Structure
Refinement against 1H Paramagnetic Relaxation Enhancement Data Arising from a
Flexible Paramagnetic Group Attached to a Macromolecule. *J. Am. Chem. Soc.* (2004)
doi:10.1021/ja031580d.

61. Conway, P., Tyka, M. D., DiMaio, F., Konerding, D. E. & Baker, D. Relaxation of
backbone bond geometry improves protein energy landscape modeling. *Protein Sci.*
(2014) doi:10.1002/pro.2389.

62. Khatib, F. *et al.* Algorithm discovery by protein folding game players. *Proc. Natl. Acad.*
Sci. U. S. A. (2011) doi:10.1073/pnas.1115898108.

63. Nivón, L. G., Moretti, R. & Baker, D. A Pareto-Optimal Refinement Method for Protein
Design Scaffolds. *PLoS One* (2013) doi:10.1371/journal.pone.0059004.

64. Tyka, M. D. *et al.* Alternate states of proteins revealed by detailed energy landscape
mapping. *J. Mol. Biol.* (2011) doi:10.1016/j.jmb.2010.11.008.

65. Leaver-Fay, A. *et al.* Rosetta3: An object-oriented software suite for the simulation and
design of macromolecules. in *Methods in Enzymology* (2011). doi:10.1016/B978-0-12-
381270-4.00019-6.

66. Park, H. *et al.* Simultaneous Optimization of Biomolecular Energy Functions on
Features from Small Molecules and Macromolecules. *J. Chem. Theory Comput.* (2016)
doi:10.1021/acs.jctc.6b00819.

Fig. 1 Conformational changes underlie coordinated DNA recombination by Cre recombinase.

a Tetrameric synaptic complex of four Cre protomers bound to two *loxP* DNA duplexes (PDB ID 2HOI), viewed from the catalytic domain, Cre^Cat. C-terminal helix αN of each protomer packs into a *trans* surface cavity (labeled in blue) in the adjacent protomer.

b Superposition of Cre^Cat domains from "inactive" (red) and "active" (gray) Cre protomers (PDB ID 2HOI: chains A and B, respectively) highlight differences in the β2-3 loop, helix αM, αM-N loop and C-terminal helix αN.

J, exhibit fast timescale dynamics.

relaxation rate constant (s^{-1}), ^{15}N

onding R$_2$/R$_1$ ratios for Cre^Cat; 800

values calculated excluding regions

ements of Cre^Cat synaptic complex

e the plot for comparison.

769 **b** {1H}-15N HetNOE values for Cre^{Cat} mapped onto the crystal structure (PDB ID 2HOI, chain

770 B) as a linear gradient from red (= 0.3) to white ($\geq$ 0.7). Residues with no data are shown in

771 gray.

Fig. 3 C-terminal deletion results in CSPs in the core of Cre^{Cat}.

a Per-residue amide CSPs [$\Delta\delta_{NH} = (\Delta\delta_H^2 + \Delta\delta_N^2/25)^{1/2}$] of Cre^{Cat} upon truncation of last 13

amino acid from the C terminus Δ(E331- D343) (red bars). The secondary structure elements

of Cre synaptic complex crystal structure (PDB ID 2HOI, chain B) are shown above the plot.

Asterisks indicate regions of high CSP containing active site residues R173, E176, H289,

R292, W315 and Y324. Residues that showed largest CSPs or peak broadening beyond

detection (assigned in Cre^{Cat} but unassigned in Cre$^{Cat}\Delta$C spectrum) are shown in maroon.

Unassigned residues are indicated by small negative black bars.

b Amide CSPs mapped onto the crystal structure (PDB ID 2HOI, chain B) as a linear gradient

from white (= 0.0 ppm) to red ($\geq$ 0.15 ppm). Sidechains of active site residues R173, E176,

K201 (modeled using PyMOL), H289, R292, W315 and Y324 are shown in blue. Truncated

C-terminal residues Δ(E331- D343) and unassigned residues are indicated in gray.

Fig. 4 PRE-NMR data reveal a *cis* to *trans* docking conformational switch in the Cre[Cat] C-terminus upon DNA binding.

a Per-residue amide PRE-NMR normalized intensity ratios (I_{ox}/I_{red}) of free Cre[Cat] C155A/C240A/D343C generated using an MTSL paramagnetic probe at C-terminal residue C343 (green arrow); strong PRE effects ($I_{ox}/I_{red} < 0.65$) are shown in red. Secondary structure elements of Cre synaptic complex crystal structure (PDB ID 2HOI, chain B) are shown. Uncertainties are propagated from the signal-to-noise ratio of individual resonances.

b Per-residue amide PRE-NMR normalized intensity ratios (I_{ox}/I_{red}) for Cre^{Cat}
C155A/C240A/D343C bound to *loxP* half-site DNA generated using MTSL paramagnetic
probe at C-terminal residue C343 (green arrow); strong PRE effects (I_{ox}/I_{red} < 0.65) are shown
in blue. Uncertainties are propagated from the signal-to-noise ratio of individual resonances.

c I_{ox}/I_{red} PRE values from free Cre^{Cat} mapped to the structure of a protomer in the crystal
structure (PDB ID 2HOI, chain B; G342, C343-MTSL modeled) as a gradient from red (I_{ox}/I_{red}=
0) to white (I_{ox}/I_{red}= 0.65). Residues with no PRE data are shown in gray. Juxtaposed
equivalent surface representation (same orientation) along with a portion of the *loxP* DNA, as
seen in the crystal (PDB ID 2HOI, chains C and D) is shown in orange to illustrate the PRE
effects mapping to the DNA binding surface.

d I_{ox}/I_{red} PRE values from the Cre^{Cat}-DNA complex mapped to the structure of a protomer in
the crystal structure (PDB ID 2HOI, chain B; G342, C343 modeled) as a gradient from blue
(I_{ox}/I_{red}= 0) to white (I_{ox}/I_{red}= 0.65). Bound DNA (PDB ID 2HOI, chains C and D) is shown in
pale orange. Residues with no PRE data are shown in gray. Juxtaposed equivalent surface
representation (same orientation) along with C-terminal helix αN region from an adjacent
MTSL-tagged protomer (PDB ID 2HOI, chain G: E331-D341 and G342, C343-MTSL
modeled) is shown in yellow to illustrate the PRE effects mapping to protein-protein *trans*
docking surface cavity. Orientation of Cre^{Cat} is rotated approximately 180° between Fig. 4**c**
and **d.**

Fig. 5 PRE-derived ensemble models of free Cre[Cat] shows *cis* docking of C-terminus over the Cre[Cat] active site.

a Top fifty PRE-derived ROSETTA models of free Cre[Cat] showing *cis* docking of C-terminal region (E331- D343; in magenta).

b Overlay of lowest ROSETTA energy member of the PRE-derived ensemble model (magenta) and the crystal structure protomer (PDB ID 2HOI, chain A) (gray).

c CSPs in free Cre[Cat] upon C-terminal truncation Δ(E331- D343) (magenta) mapped onto the lowest energy member of the PRE-derived ensemble model as a linear gradient from white (0.0 ppm) to red (≥ 0.15 ppm). Unassigned residues are shown in gray (as in Fig. 3**b**).

d Difference in the inter-residue Cα-Cα contact map between the top ten PRE-derived models and the DNA bound tetrameric crystal structure (PDB ID 2HOI, chain A) (absolute values of changes in inter-residue distances |Δdistance$_{C\alpha-C\alpha}$| (Å) are shown by a three-color heatmap from blue to yellow). The secondary structure elements of Cre synaptic complex crystal structure (PDB ID 2HOI, chain B) are shown along the axes. Diagonal is shown as dashed line (orange).

Fig. 6. C-terminal helix αN region of Cre[Cat] is perturbed upon binding to *loxP* DNA

a Per-residue amide CSPs [$\Delta\delta_{NH} = (\Delta\delta_H^2 + \Delta\delta_N^2/25)^{1/2}$] of Cre[Cat] upon binding to *loxP* half-site hairpin DNA with a linear color ramp from gray (= 0 ppm) to red ($\geq$ twice the $\Delta\delta$ SD (0.047 ppm)). Largest CSPs, or those that resulted in peak-broadening beyond detectability (assigned in free Cre[Cat] but unassigned in the Cre[Cat]- DNA complex) are shown in maroon. Unassigned residues are indicated by small negative black bars. Secondary structure elements of Cre synaptic complex crystal structure (PDB ID 2HOI, chain B) are shown above the plot.

b Overlay of ^{1}H-^{15}N TROSY-HSQC spectra of free Cre[Cat] (black) and bound to a DNA half-site (red) with select residue assignments indicated (> 3 SD ppm).

845　**c** CSPs mapped onto the crystal structure (PDB ID 2HOI, chain B) as a linear gradient from

846　white ($\leq$ 1 SD ppm) to red ($\geq$ 2 SD ppm). Unassigned residues are shown in gray.